

Pancharatnam-Zak phase

Vivek M. Vyas, Dibyendu Roy, and Joseph Samuel
Raman Research Institute,
C. V. Raman Avenue, Sadashivnagar,
Bangalore 560080, India

The geometric phase acquired by an electron in a one-dimensional periodic lattice due to weak electric perturbation is found and referred to as the Pancharatnam-Zak phase. The underlying mathematical structure responsible for this phase is unveiled. As opposed to the well-known Zak phase, the Pancharatnam-Zak phase is a gauge invariant observable phase, and correctly characterizes the energy bands of the lattice. We demonstrate the gauge invariance of the Pancharatnam-Zak phase in two celebrated models displaying topological phases. A filled band generalization of this geometric phase is constructed and is observed to be sensitive to the Fermi-Dirac statistics of the band electrons. The measurement of the single-particle Pancharatnam-Zak phase in individual topological phases, as well as the statistical contribution in its many-particle generalization, should be accessible in various controlled quantum experiments.

I. INTRODUCTION

The concept of geometric phase [1] appears surprisingly late in the development of quantum mechanics. The work of Pancharatnam [2] on the interference of polarized light is now recognized as an early precursor of this phase [3]. Other anticipations of this effect are found in the works of Mead and Truhlar [4], Herzberg and Longuet-Higgins [5] and Aharonov and Bohm [6]. The work of Berry [7] on the adiabatic theorem of quantum mechanics drew wide attention to the scope and generality of the geometric phase [3] which manifests in a variety of phenomena across physics [8].

The notion of geometric phase in a condensed matter system perhaps makes its first appearance in the work of Thouless [9, 10] in the guise of a topological invariant, known as the Chern number, characterizing different quantum Hall states. This topological invariant was in fact, the integral of a geometric quantity (the Berry curvature) over a closed manifold [11], rather like the Gauss-Bonnet theorem in classical geometry. Since then, such topological invariants have been employed to understand and classify the phases of several condensed matter systems [8, 12, 13]. The central theme in the current understanding of electromagnetic properties of insulators and superconductors is the usage of the geometric phase as an invariant, describing various quantum states in such systems [14–18].

The purpose of this paper is to study the origin and structure of the geometric phase in a one-dimensional (1D) periodic lattice under the influence of a weak electric field. The earliest attempt in evaluating the geometric phase in such a system was due to Zak [19], who is claimed to have derived an expression for the same, which is often called the *Zak phase*. It was later understood that the Zak phase is closely related to the Chern number characterizing the quantized particle transport in a 1D insulator [14, 20]. These works subsequently lead to the modern understanding of the electric polarization in dielectric materials purely in terms of the geometrical

phase [17, 20–23]. The emergence of the Zak phase has been studied in strongly coupled LC circuits [24] and photonic waveguide lattices [25]. It has also been employed to classify the edge states in planar honeycomb lattice systems [26, 27] and to understand the topological states of a spin chain with long-range interactions [28]. The Zak phase is also used for characterizing topological edge states of 1D photonic lattice for the experimental demonstration of lasing in topological edge states [29] and for isolating topological defects from trivial defects [30].

The geometric phase has been observed in several experiments in diverse areas of physics [8, 31]. The value of the geometric phase, a physically measurable quantity, can not be altered by spatially translating the system. Nevertheless, it was found that the value of the Zak phase itself was changed as one translated the system spatially or employed a different gauge convention for the Bloch states [17, 19, 22, 23, 32–34]. This problem leads to a general belief that the Zak phase in such systems itself is *not* a physically measurable quantity [32]. It essentially means that the Zak phase can not be a genuine geometric phase, as claimed in the literature, which by definition is invariant under gauge transformation and spatial translation.

In this light, one naturally wonders about the correct notion of the geometric phase in a 1D periodic lattice. By careful consideration of the concept of the geometric phase in its generality, here we find the geometric phase gained by (a) a single electron, and (b) by electrons of a filled band of a lattice when influenced by a weak electric field. The geometric phase in the single electron case, referred to as the *Pancharatnam-Zak phase*, possesses the essential invariances under gauge transformation and spatial translation. The underlying geometrical and topological properties of the system are uncovered in the course of its derivation. In particular, it is shown that this system displays cyclicity under time evolution in a generalized sense, rather than the existing notion of cyclicity based on returning of the density matrix in the course of evolution. The Pancharatnam-Zak phase is found to be quantized for systems with inversion symme-

try, in the sense that it either equal to 0 or π . The geometrical phase for the filled band case is properly formulated, and surprisingly, it is found to receive a contribution owing to the Fermi-Dirac statistics of the electrons. An explicit calculation of the Pancharatnam-Zak phase is also presented for the Su-Schrieffer-Heeger (SSH) model and the Kitaev model of 1D p -wave superconductors. The theoretical value for the difference of (single-particle) Pancharatnam-Zak phase in the SSH model across the topological phase transition is observed to agree with the experimental measurement [33] by Atala *et al.* in an optical lattice setup.

The paper is organized as follows. In section (II), the concept of the geometric phase, as defined in its generality, is briefly reviewed. Subsequently, the problem of a charged particle in a 1D periodic lattice, subjected to a weak electric field, is formulated and its kinematic aspects are studied in section (III), bringing out the underlying mathematical structure. In section (IV), the adiabatic quantum dynamics of such a motion discussed in (a) single-particle case, and (b) in many-particle filled band case; and the manifestation of the geometric Pancharatnam-Zak phase in both the cases is found. An explicit calculation of the single-particle Pancharatnam-Zak phase for SSH and Kitaev model is provided in section (V), followed by the conclusion.

II. GEOMETRIC PHASE

Consider a quantum mechanical system being described by some Hermitian Hamiltonian $\hat{H}(t)$ which generates time evolution of the (normalized) state $|\psi(t)\rangle$ as per the Schrödinger equation:

$$\hat{H}(t)|\psi(t)\rangle = i\hbar\frac{\partial}{\partial t}|\psi(t)\rangle. \quad (1)$$

In the course of time evolution, the state $|\psi(t_f)\rangle$ differs from $|\psi(t_i)\rangle$, and one can ask about the net phase acquired in the process. The acquired phase is given by $\gamma_T = \text{Arg} \langle \psi(t_i) | \psi(t_f) \rangle$, and is often referred to as the *Pancharatnam total phase* [35, 36][37]. At any given time t , the state $|\psi(t)\rangle$ and $e^{i\Lambda(t)}|\psi(t)\rangle$ ($\Lambda(t)$ is some real function of t) represent the same physical state, since the corresponding density matrices are identical. Nevertheless, the total phase γ_T acquired is different for these states $|\psi(t)\rangle$ and $e^{i\Lambda(t)}|\psi(t)\rangle$, showing that the total phase γ_T is not a physical observable.

Motivated by the requirement that a physically observable phase must be the same for whether the system is described by states $|\psi(t)\rangle$ or $e^{i\Lambda(t)}|\psi(t)\rangle$, it turns out that a unique phase γ_g can be extracted from γ_T by subtracting what is called the dynamical phase γ_{dyn} [35, 36]:

$$\gamma_g = \gamma_T - \gamma_{dyn} \quad (2)$$

$$= \text{Arg} \langle \psi(t_i) | \psi(t_f) \rangle + i \int_{t_i}^{t_f} dt \langle \psi(t) | \frac{\partial}{\partial t} |\psi(t)\rangle. \quad (3)$$

It can be readily checked that this phase is *invariant* under the *local gauge transformation* $|\psi(t)\rangle \rightarrow e^{i\Lambda(t)}|\psi(t)\rangle$.

Interestingly, the phase γ_g also possesses another property called *reparameterisation invariance*. Let a real parameter $r(t)$, an increasing function of time t , be used to relabel the states $|\psi(t)\rangle$. The parameter r takes the values between $r_i = r(t_i)$ and $r_f = r(t_f)$, so that one can now express states $|\psi(t)\rangle$ as parameterised by $r(t)$: $|\tilde{\psi}(r)\rangle = |\psi(t)\rangle$. Upon changing the variable t with r in (3), it is immediately clear that:

$$\begin{aligned} \text{Arg} \langle \psi(t_i) | \psi(t_f) \rangle + i \int_{t_i}^{t_f} dt \langle \psi(t) | \frac{\partial}{\partial t} |\psi(t)\rangle \\ = \text{Arg} \langle \tilde{\psi}(r_i) | \tilde{\psi}(r_f) \rangle + i \int_{r_i}^{r_f} dr \langle \tilde{\psi}(r) | \frac{\partial}{\partial r} |\tilde{\psi}(r)\rangle. \end{aligned}$$

This property of form invariance of γ_g under reparameterisation, along with the local gauge invariance, is a clear evidence of the geometric origin of this phase. Owing to this fact, γ_g , so defined as (3), is called the *geometric phase*. The above expression for (3) and its proper generalization encompassing the case of non-unitary evolution were originally obtained long back using a manifestly geometric route [35]. The existence of such a phase is well established through several experiments [3, 8, 31, 38, 39].

The geometric phase γ_g possesses yet another invariance under the *static unitary operations*. Let the state $|\bar{\psi}\rangle$ obtained from $|\psi\rangle$ as: $|\bar{\psi}(t)\rangle = \hat{T}|\psi(t)\rangle$, where \hat{T} is time independent unitary operator. Then it is clear that $\langle \bar{\psi}(t_i) | \bar{\psi}(t_f) \rangle = \langle \psi(t_i) | \psi(t_f) \rangle$, and $\langle \bar{\psi}(t) | \frac{\partial}{\partial t} |\bar{\psi}(t)\rangle = \langle \psi(t) | \frac{\partial}{\partial t} |\psi(t)\rangle$, which implies the invariance of γ_g under action of \hat{T} . Noting the that the coordinate translation operator $\hat{T}_x(\epsilon) = e^{-\frac{i\epsilon}{\hbar}\hat{p}}$ and momentum translation operator $\hat{T}_p(\epsilon) = e^{-\frac{i\epsilon}{\hbar}\hat{x}}$ are both time independent unitary operations, it thus follows that the geometric phase γ_g is *invariant* under coordinate and momentum translations.

It is known [35] that the expression (3) can be cast into a familiar form by using the natural connection $\mathcal{A}(t) = i\langle \psi(t) | \frac{\partial}{\partial t} |\psi(t)\rangle$ and by noting that the Pancharatnam total phase $\text{Arg} \langle \psi(t_i) | \psi(t_f) \rangle$ can be expressed as a line integral over the shortest geodesic connecting $|\psi(t_f)\rangle$ and $|\psi(t_i)\rangle$:

$$\text{Arg} \langle \psi(t_i) | \psi(t_f) \rangle = \int_{t_f}^{t_i} ds \mathcal{A}(s) \Big|_{\text{geodesic}}. \quad (4)$$

Here the states $|\psi(s)\rangle$ used to define $\mathcal{A}(s)$, do not evolve by the Schrödinger equation but rather by the geodesic equation:

$$D_s^2 |\psi(s)\rangle = 0, \quad (5)$$

where the covariant derivative $D_s = \frac{\partial}{\partial s} - i\mathcal{A}(s)$. With

this identification the expression for γ_g now reads:

$$\gamma_g = \oint_C dl \mathcal{A}(l), \quad (6)$$

where the integration is to be done along the closed curve C , which consists of (i) the curve traced by $|\psi(t)\rangle$ in course of its Schrödinger evolution from t_i to t_f , and (ii) the shortest geodesic curve connecting $|\psi(t_f)\rangle$ to $|\psi(t_i)\rangle$. This expression manifests the local gauge invariance of the geometric phase.

In the special case, when the final state $|\psi(t_f)\rangle$ turns out to be *colinear* to the initial state $|\psi(t_i)\rangle$, which means:

$$|\psi(t_f)\rangle = e^{i\theta} |\psi(t_i)\rangle, \quad (7)$$

then the corresponding geometric phase is often referred to as *cyclic*. The reason behind this nomenclature is the fact that even though the initial and final states are not identical, the corresponding density matrices are identical, indicating that the system returns to its initial state in the course of time evolution.

The notion of geometric phase in the systems, whose time evolution is both adiabatic and cyclic, is well studied by Berry [7]. Usually it is assumed that the time dependence in the Hamiltonian only appears via a parameter $\lambda(t)$, which is cyclic: $\lambda(t_i) = \lambda(t_f)$. Owing to the adiabaticity condition, the state of the system is given by: $|\psi(t)\rangle \simeq e^{i\phi(t)} |\bar{n}(\lambda)\rangle$ ($\phi(t)$ is a real phase factor), where the vector $|\bar{n}(\lambda)\rangle$ is the instantaneous eigenket of the Hamiltonian. Generally, it is assumed that $|\bar{n}(\lambda)\rangle$ is a smooth function of λ as also $|\bar{n}(\lambda_i)\rangle = |\bar{n}(\lambda_f)\rangle$, in which case the expression for the geometric phase, as obtained by Berry, appears without the Pancharatnam total phase contribution (since it vanishes).

However, if the time evolution in a given system is adiabatic but not strictly cyclic, then the notion of geometric phase of Berry does not hold, and one needs to work with its general definition, as given by (3).

III. PERIODIC POTENTIAL PROBLEM

Consider a (spinless) charged particle of mass μ and charge e defined on a line, under the influence of a 1D periodic potential $V(x)$ with lattice constant a . Owing to the periodicity of potential $V(x) = V(x+a)$, it follows that:

$$[\hat{H}(0), \hat{T}_x(a)] = 0, \quad (8)$$

where $\hat{T}_x(a)$ is a spatial translation operator over a unit cell, and

$$\hat{H}(0) = \frac{\hat{p}^2}{2\mu} + V(\hat{x}). \quad (9)$$

The simultaneous eigenstates $\Psi_{nk_m}(x) \equiv \langle x | \Psi_{nk_m} \rangle$, which diagonalise both these operators solve:

$$\hat{H}(0)\Psi_{nk_m}(x) = E_{nk_m}\Psi_{nk_m}(x), \quad (10)$$

$$\hat{T}_x(a)\Psi_{nk_m}(x) = e^{ik_m a}\Psi_{nk_m}(x). \quad (11)$$

These eigenvalue problems are defined with the *periodic boundary condition* (PBC):

$$\Psi_{nk_m}(x+L) = \Psi_{nk_m}(x), \quad (12)$$

where $L = Na$, and N represents the number of unit cells in the system. Here, the wave vector $k_m = \frac{2\pi}{L}m$ (m is an integer) takes values within the first Brillouin zone $[0, \frac{2\pi}{a}]$, whereas n stands for the band index. By construction the eigenstates $\Psi_{nk_m}(x)$ possess the Bloch property: $\Psi_{nk_m}(x+a) = e^{ik_m a}\Psi_{nk_m}(x)$. As a result, one finds that:

$$\Psi_{nk_{m+N}}(x) = \Psi_{nk_m}(x)e^{i\chi}, \quad (13)$$

where χ is some arbitrary real number. The equation (13) shows that both the states $\Psi_{nk_{m+N}}(x)$ and $\Psi_{nk_m}(x)$ correspond to the same physical state, since the corresponding density matrices are identical. It is often assumed that $\chi = 0$, and is referred to as the *periodic gauge condition* [22, 32]. Since there is no physical reason to work with such a gauge restriction, in what follows no condition is imposed on χ , and we shall demand that the physical observables must be insensitive to the choice of χ .

The cell periodic Bloch states $u_{nq}(x)$ are given by the application of the momentum translation operator [40] $\hat{T}_p(k) = e^{-ik\hat{x}}$ on $\Psi_{nq}(x)$:

$$u_{nq}(x) = e^{-iqx}\Psi_{nq}(x), \quad (14)$$

so that they solve the eigenvalue problem:

$$\begin{aligned} \hat{H}_q|u_{nq}\rangle &= E_{nq}|u_{nq}\rangle, \quad \text{where} \\ \hat{H}_q &= \frac{1}{2\mu}(\hat{p} + \hbar q)^2 + V(\hat{x}). \end{aligned} \quad (15)$$

Consider that this system is further influenced by an external weak DC electric field, so that the Hamiltonian now reads:

$$\hat{H}(t) = \frac{1}{2\mu}(\hat{p} - eA(t))^2 + V(\hat{x}). \quad (16)$$

Since the time dependence in the above Hamiltonian appears only from the vector potential $A(t) = -Et$, it is useful to define a real parameter: $\alpha(t) = -eA(t)/\hbar$, which allows the Hamiltonian to be written as:

$$\hat{H}(t) \equiv \frac{1}{2\mu}(\hat{p} + \hbar\alpha(t))^2 + V(\hat{x}) \equiv \hat{H}_{\alpha(t)}. \quad (17)$$

It may be noted that the parameter $\alpha(t)$ is a monotonically increasing function of t , and is not cyclic. Owing to

the periodicity of potential $V(x) = V(x + a)$ and therefore of \hat{H}_α , one has $[\hat{H}_\alpha, \hat{T}_x(a)] = 0$, which allows for their simultaneous instantaneous eigenstate, denoted by $\Psi_{nk_m\alpha}$, which solves the eigenvalue problems:

$$\hat{H}_\alpha \Psi_{nk_m\alpha}(x) = E_{nk_m\alpha} \Psi_{nk_m\alpha}(x), \quad (18)$$

$$\hat{T}_x(a) \Psi_{nk_m\alpha}(x) = e^{ik_m a} \Psi_{nk_m\alpha}(x), \quad (19)$$

where the wavefunction $\Psi_{nk_m\alpha}(x)$ obeys the PBC.

The fact that $[\hat{H}_\alpha, \hat{T}_x(a)] = 0$ gives rise to important consequences on the time evolution of the system. Firstly, $\hat{T}_x(a)$ is conserved under time evolution. Furthermore if the system is prepared initially in an eigenstate $|\Psi_{nk_l\alpha(0)}\rangle$ of $\hat{T}_x(a)$, then it immediately follows that under time evolution the system is forbidden to evolve into any other eigenstate $|\Psi_{n'k_l\alpha(t)}\rangle$ ($l \neq l'$) in any band n' at any time t . This clearly shows that the wave vector k_l is a conserved quantum number under time evolution. However, it must be noted that this conservation law does not prohibit the system into evolving to the state $|\Psi_{n'k_l\alpha(t)}\rangle$, with the same wave vector k_l albeit in a different band n' .

The Hamiltonian (17) has very interesting property under time evolution. The vector potential at any time t can be written as a gauge transformation:

$$A(t) = A(0) + \frac{i\hbar}{e} U^\dagger(x, t) \partial_x U(x, t), \quad (20)$$

where

$$U(x, t) = \exp\left(\frac{i}{\hbar} e E t x\right), \quad (21)$$

and $A(0) = 0$ by virtue of its definition. Under such a transformation, the momentum operator transforms as: $\hat{p} - eA(t) = U^\dagger(x, t) \hat{p} U(x, t)$, which allows the Hamiltonian at some time t and at $t = 0$ to be written as:

$$\hat{H}(t) = U^\dagger(x, t) \hat{H}(0) U(x, t). \quad (22)$$

The gauge transformation $U(x, t)$, which is a function of x , must respect the PBC, $U(x, t) = U(x + L, t)$ in order to be a well defined operator. It immediately follows that only for time $t = j\tau$ (j is an integer), the boundary condition is respected, where

$$\tau = \frac{2\pi\hbar}{eEL}. \quad (23)$$

This shows that the Hamiltonian $\hat{H}(j\tau)$ (for different j s) are physically the same (they are *gauge equivalent*), their spectra are identical, and further their instantaneous eigenstates are related to each other by the gauge transformation $U(x, j\tau)$. It is straightforward to see that the instantaneous eigenstates $\Psi_{nk_m\alpha(j\delta)}(x)$ are expressible in terms of $\Psi_{nk_m\alpha(0)}(x)$ as:

$$\Psi_{nk_m\alpha(j\delta)}(x) = U^\dagger(x, j\tau) \Psi_{nk_{m+j}\alpha(0)}(x) \quad (24)$$

$$= \exp\left(-i\frac{2\pi x j}{L}\right) \Psi_{nk_{m+j}\alpha(0)}(x), \quad (25)$$

and the energies are related by: $E_{nk_m\alpha(j\delta)} = E_{nk_{m+j}\alpha(0)}$.

The gauge transformation factor $U(x, j\tau) = e^{i\frac{2\pi x j}{L}}$ has a very interesting topological property. It is a function of x , albeit with the PBC, implying that the points $x = 0$ and $x = L$ are identified since $U(0, j\tau) = U(L, j\tau)$. This shows that it lives on a circle with circumference L . Now, $U(x, j\tau)$ by definition is a phase and takes values only on the unit circle in the complex plane. Thus, $U(x, j\tau)$ is a map from one circle (with circumference L) to the unit circle. Such maps are classified in terms of homotopy classes [41], with each of them characterized by an integer called the *winding number*, which measures the number of times one circle is wound on another. In the case of $U(x, j\tau)$, it is easy to see that the integer j is actually the winding number; under one rotation in x space, the factor $e^{i\frac{2\pi x j}{L}}$ completes j rotations of the unit circle. As a result, it is not possible to continuously deform $U(x, j\tau)$ to some $U(x, j'\tau)$ for $j \neq j'$. The class of such gauge transformations, which can not be continuously deformed into the identity (gauge) transformation, is often referred to as *large gauge transformations*.

This shows that the system owing to the PBC only allows for the large gauge transformations $U(x, j\tau)$, which are topologically non-trivial in the sense that each $U(x, j\tau)$ belongs to different homotopy class with the winding number j .

IV. GEOMETRIC PHASE IN THE PERIODIC POTENTIAL PROBLEM

As seen in section (II), in order to determine the geometric phase acquired during the evolution, the knowledge of the state $|\Phi(t)\rangle$ of the system during the course of time evolution is essential. As shown earlier for the periodic potential problem described by (16), if the system is prepared to be in the instantaneous eigenstate $|\Phi(0)\rangle = |\Psi_{nk_l\alpha(0)}\rangle$ at time $t = 0$, then the system is constrained to evolve with the same quantum number k_l at any other time t . This dictates that the system can only evolve into a linear combination of states $|\Psi_{mk_l\alpha(t)}\rangle$ with different values of m , which all have the same quantum number k_l , albeit different energies $E_{mk_l\alpha}$. Assuming that the external electric field is sufficiently weak such that the evolution to other band states $|\Psi_{mk_l\alpha(t)}\rangle$ ($m \neq n$) is energetically suppressed, such transition can be ignored in the leading order. Thus, one assumes that the time evolution is *adiabatic* [42, 43] in this approximation. So, the system only evolves into the instantaneous eigenstate $|\Psi_{nk_l\alpha(t)}\rangle$ along with an overall phase:

$$|\Phi(t)\rangle = e^{i\phi(t)} |\Psi_{nk_l\alpha(t)}\rangle, \quad (26)$$

where $\phi(t)$ is given by [7, 42]:

$$\phi(t) = \phi_g(t) - \frac{1}{\hbar} \int_0^t ds E_{nk_l\alpha(s)}, \quad (27)$$

with $\phi_g(t) = i \int_0^t ds \langle \Psi_{nk_l\alpha(s)} | \frac{\partial}{\partial s} | \Psi_{nk_l\alpha(s)} \rangle$. This can be expressed in the light of (24) as:

$$|\Phi(j\tau)\rangle = e^{i\phi(j\tau)} \hat{U}^\dagger(x, j\tau) |\Psi_{nk_{l+j}\alpha(0)}\rangle, \quad (28)$$

which shows that the system, which was initially in the eigenstate with a wave vector k_l , in the course of adiabatic evolution goes into the eigenstate with a wave vector k_{l+j} (in the same band) after time $t = j\tau$, modulo a large gauge transformation with the winding number $-j$. This remarkable result leads to the evaluation of the geometric phase acquired by the system in two distinct cases: the single-particle case and the filled band many-particle case.

A. Single-particle case

From the above relation (28), it immediately follows that after the time evolution for time $N\tau$, the state of the system is given by:

$$|\Phi(N\tau)\rangle = e^{i\chi} e^{i\phi(N\tau)} \hat{U}^\dagger(x, N\tau) |\Psi_{nk_l\alpha(0)}\rangle, \quad (29)$$

indicating that the system returns to the initial state with a large gauge transformation. It may be noted that in general,

$$|\langle \Psi_{nk_l\alpha(0)} | \hat{U}^\dagger(x, N\tau) | \Psi_{nk_l\alpha(0)} \rangle| \neq 1. \quad (30)$$

This shows that neither the initial and final states are colinear:

$$|\Phi(N\tau)\rangle \neq e^{i\theta} |\Phi(0)\rangle, \quad (31)$$

and nor the corresponding density matrices are identical. Thus, strictly speaking the system does not return to its initial state after the time evolution for time $N\tau$. However, owing the gauge transformation factor $U^\dagger(x, N\tau)$ in (29), it is straightforward to see that the average of any gauge covariant observable $\hat{F}(\hat{x}, \hat{p} - eA(t))$ returns after time $N\tau$:

$$\begin{aligned} & \langle \Psi_{nk_l\alpha(0)} | \hat{F}(\hat{x}, \hat{p} - eA(0)) | \Psi_{nk_l\alpha(0)} \rangle \\ &= \langle \Psi_{nk_l\alpha(N\tau)} | \hat{F}(\hat{x}, \hat{p} - eA(N\tau)) | \Psi_{nk_l\alpha(N\tau)} \rangle. \end{aligned} \quad (32)$$

So the states $|\Phi(0)\rangle$ and $|\Phi(N\tau)\rangle$ are though not colinear, nevertheless they represent the same physical state of the system, albeit expressed in different gauges. Thus, the time evolution of the system in this case must be understood to be of *adiabatic* and *cyclic* kind. It must be mentioned that this notion of cyclicity generalizes the existing notion in the literature [35, 36] based on the requirement of returning of the density matrix.

The geometric phase gained by the system after such a cyclic adiabatic evolution then follows from the substitution of (26) in (3), and it reads:

$$\begin{aligned} \gamma_g(n) &= \text{Arg} \langle \Psi_{nk_l\alpha(0)} | \Psi_{nk_l\alpha(N\tau)} \rangle \\ &+ i \int_0^{N\tau} dt \langle \Psi_{nk_l\alpha(t)} | \frac{\partial}{\partial t} | \Psi_{nk_l\alpha(t)} \rangle. \end{aligned} \quad (33)$$

Employing the reparameterisation invariance of the geometric phase, enables us to rewrite $\gamma_g(n)$ while treating α as a parameter, as:

$$\begin{aligned} \gamma_g(n) &= \text{Arg} \langle \Psi_{nk_l\alpha=0} | \Psi_{nk_l\alpha=\frac{2\pi}{a}} \rangle \\ &+ i \int_0^{\frac{2\pi}{a}} d\alpha \langle \Psi_{nk_l\alpha} | \frac{\partial}{\partial \alpha} | \Psi_{nk_l\alpha} \rangle. \end{aligned} \quad (34)$$

The above equation can be further simplified by working with the cell periodic Bloch states $|u(k_l + \alpha(t))\rangle$ which are defined as:

$$|\Psi_{nk_l\alpha}\rangle = e^{ik_l\hat{x}} |u_n(k_l + \alpha)\rangle, \quad (35)$$

which solve the eigenvalue problem for $\hat{H}_{k_l+\alpha}$:

$$\hat{H}_{k_l+\alpha} |u_n(k_l + \alpha)\rangle = E_{nk_l\alpha} |u_n(k_l + \alpha)\rangle, \quad (36)$$

where

$$\hat{H}_{k_l+\alpha} = \frac{1}{2\mu} (\hat{p} + \hbar\alpha + \hbar k_l)^2 + V(\hat{x}). \quad (37)$$

These states $|u_n(k_l + \alpha)\rangle$ are generalization of states $|u_n(q)\rangle$ encountered earlier in (14), albeit the parameter q is now understood to be continuous with the identification $q = k_l + \alpha$. Setting $k_l = 0$ without loss of generality, one gets a very important relation for the geometric phase $\gamma_g(n)$ expressed in terms of $|u_n(\alpha)\rangle$:

$$\gamma_g(n) = \text{Arg} \langle u_n(0) | u_n(2\pi/a) \rangle + i \int_0^{\frac{2\pi}{a}} d\alpha \langle u_n(\alpha) | \frac{\partial}{\partial \alpha} | u_n(\alpha) \rangle. \quad (38)$$

The above expression for $\gamma_g(n)$, which is referred to as *Pancharatnam-Zak phase*, is one of the main results of this paper. Comprising of the contributions from all the states in the band n , this geometric phase properly characterizes the band. This expression can also be derived using a direct axiomatic route as shown in the appendix.

The idea of evaluating the geometric phase in such a 1D periodic lattice potential influenced by a weak electric field has a long history starting from the work of Zak [19]. The expression thus obtained, and often claimed to be the geometric phase, is popularly referred to as *Zak phase* [17, 22, 23, 32, 34]. It is given by:

$$\gamma_Z(n) = i \int_0^{\frac{2\pi}{a}} dk \langle u_n(k) | \frac{\partial}{\partial k} | u_n(k) \rangle, \quad (39)$$

wherein the states $|u_n(k)\rangle$ are required to obey the periodic gauge condition [17, 19, 22, 23]:

$$|u_n(k + 2\pi/a)\rangle = e^{-i\frac{2\pi}{a}\hat{x}} |u_n(k)\rangle.$$

In derivation of (39), the wave vector is treated as a continuous variable k parametrising the states, and the limit $N \rightarrow \infty$ is assumed. Comparing (38) and (39), it is evident that the second term in the (38) can be identified with the Zak phase [44]. This clearly shows that the Zak phase is only a partial contribution to the net geometric phase $\gamma_g(n)$, and it is devoid of the non-vanishing contribution of the Pancharatnam total phase $\text{Arg}\langle u_n(0)|u_n(2\pi/a)\rangle$ [45]. A clear testimony of this fact is that the value of the Zak phase is known to suffer from the well acknowledged flaws of gauge-dependence and non-invariance under spatial translation [22, 23, 32–34].

Often in the literature it is inaccurately claimed [17, 19, 22, 34] that the Zak phase is gauge invariant, by showing that $\gamma_Z(n) \rightarrow \gamma_Z(n) - 2\pi m$ under a restricted class of gauge transformations, for which $\Lambda(2\pi/a) = \Lambda(0) + 2\pi m$ (m being an integer). As argued earlier, the physical requirement of invariance under *any* general gauge transformation stems from the fact that the state $|u_n(q)\rangle$ and $e^{i\Lambda(q)}|u_n(q)\rangle$ ($\Lambda(q)$ is some arbitrary function of q), represent the same physical state of the system, since the corresponding density matrices are identical. As a result one demands that a physically observable quantity must remain invariant under a general gauge transformation for any choice of $\Lambda(q)$. This is clearly not the case with the Zak phase. (The gauge dependence of the Zak phase in an alternate way is also shown in the appendix.)

It follows from section (II) and (III) that the Pancharatnam total phase contribution can be expressed as a line integral:

$$\text{Arg}\langle u_n(0)|u_n(2\pi/a)\rangle = i \int_{2\pi/a}^0 ds \langle u_n(s) | \frac{\partial}{\partial s} | u_n(s) \rangle \Big|_{\text{geodesic}}, \quad (40)$$

which is to be evaluated along the shortest geodesic connecting $|u_n(2\pi/a)\rangle$ to $|u_n(0)\rangle$. With the identification of natural connection $\mathcal{A}_n(l) = i\langle u_n(l) | \frac{\partial}{\partial l} | u_n(l) \rangle$, the Pancharatnam-Zak phase thus takes a manifestly gauge invariant form [46]:

$$\gamma_g(n) = \oint_C dl \mathcal{A}_n(l). \quad (41)$$

The local gauge invariance of $\gamma_g(n)$ can be explicitly checked by considering a local gauge transformation of the form:

$$|u_n(q)\rangle \rightarrow e^{i\Lambda(q)}|u_n(q)\rangle. \quad (42)$$

It is immediately clear that the contributions arising from the first and the second term in (38) due to such a gauge transformation exactly cancel each other.

Apart from the local gauge invariance, the Pancharatnam-Zak phase $\gamma_g(n)$ is also invariant under spatial translation. Consider two observers whose coordinate origins are separated by a distance d . If one

observer finds that the system is in the state $|\Psi_{nq}\rangle$, then the other observer finds that it is described by $|\tilde{\Psi}_{nq}\rangle = e^{i\frac{d}{\hbar}\hat{p}}|\Psi_{nq}\rangle$. From (14), it follows that the corresponding Bloch states $|u_n(q)\rangle$ and $|\tilde{u}_n(q)\rangle$ are related by a local gauge transformation and translation:

$$|\tilde{u}_n(q)\rangle = e^{iqd} e^{i\frac{d}{\hbar}\hat{p}}|u_n(q)\rangle. \quad (43)$$

We have already seen earlier in section (II), and it can also be checked explicitly using the relation (43) in (38) that the geometric phase $\gamma_g(n)$ remains invariant under the above operation. So, the two observers located at different points in space indeed agree upon the Pancharatnam-Zak phase $\gamma_g(n)$ of the system, which means that the value of the Pancharatnam-Zak phase does not depend upon the choice of the origin of the unit cell. Here, one again notes that neither the argument term nor the Zak phase term in (38) are individually invariant under translation operation (43) [47].

It is worth noting that the Pancharatnam-Zak phase so obtained above is independent of the total number of cells N in the system, and the continuum limit $N \rightarrow \infty$ has not been assumed in deriving it.

The spatial inversion (unitary) operator $\hat{\Pi}$ is defined such that $-\hat{x} = \hat{\Pi} \hat{x} \hat{\Pi}^\dagger$ and $-\hat{p} = \hat{\Pi} \hat{p} \hat{\Pi}^\dagger$. So for the lattices which are inversion symmetric, that is $V(-\hat{x}) = \hat{\Pi} V(\hat{x}) \hat{\Pi}^\dagger = V(\hat{x})$, one finds that $|u_n(-\kappa)\rangle = \hat{\Pi}|u_n(\kappa)\rangle$ from (35). Using this in (38) along with the reparameterisation invariance of $\gamma_g(n)$, one immediately finds that the Pancharatnam-Zak phase for such a system is quantized:

$$\gamma_g(n) = 0 \text{ or } \pi. \quad (44)$$

In the subsequent section, the Pancharatnam-Zak phase γ_g has been explicitly calculated for the SSH model and the Kitaev chain to show the validity of the above assertions.

B. Many-particle case

From equation (28), it is clear that if the system was prepared in the initial state $|\Psi_{nk_l\alpha(0)}\rangle$, then after time τ it adiabatically evolves as:

$$|\Phi(\tau)\rangle = e^{i\phi(\tau)} \hat{U}^\dagger(x, \tau) |\Psi_{nk_{l+1}\alpha(0)}\rangle, \quad (45)$$

and the Hamiltonian returns modulo a large gauge transformation:

$$\hat{H}(\tau) = \hat{U}^\dagger(x, \tau) \hat{H}(0) \hat{U}(x, \tau), \quad (46)$$

indicating that the spectra at time $t = 0$ and $t = \tau$ are identical.

This observation motivates one to consider the N -particle generalization of this problem, the case wherein the band n is completely filled by N non-interacting spinless fermions. The many-particle wavefunction $\tilde{\Psi}$ representing such a filled band at any time t in the adiabatic approximation is given by the Slater determinant:

$$\bar{\Psi}_n(x_1, x_2, \dots, x_N; \alpha(t)) = \frac{1}{\sqrt{N!}} \begin{vmatrix} \Phi_{nk_0\alpha}(x_1) & \Phi_{nk_1\alpha}(x_1) & \cdots & \Phi_{nk_{N-1}\alpha}(x_1) \\ \Phi_{nk_0\alpha}(x_2) & \Phi_{nk_1\alpha}(x_2) & \cdots & \Phi_{nk_{N-1}\alpha}(x_2) \\ \vdots & \vdots & & \vdots \\ \Phi_{nk_0\alpha}(x_N) & \Phi_{nk_1\alpha}(x_N) & \cdots & \Phi_{nk_{N-1}\alpha}(x_N) \end{vmatrix}. \quad (47)$$

Here, $\Phi_{nk_l\alpha(t)}(x_j) \equiv \langle x_j | \Phi(t) \rangle$ represents the j^{th} particle wavefunction whose adiabatic evolution can be expressed using equations (26,27) as:

$$\Phi_{nk_l\alpha(t)}(x_j) = e^{-\int_0^t ds \langle \Psi_{nk_l\alpha(s)} | \partial_s | \Psi_{nk_l\alpha(s)} \rangle} \times e^{-\frac{i}{\hbar} \int_0^t ds E_{nk_l\alpha(s)} \Psi_{nk_l\alpha(s)}(x_j)}.$$

The wavefunction $\Psi_{nk_l\alpha(t)}(x_j)$ is the instantaneous eigenstate of the Hamiltonian $\hat{H}_{\alpha(t)}^j = \frac{1}{2m} (\hat{p}_j + \hbar\alpha(t))^2 + V(x_j)$. Employing equation (45), we can write the many-particle wavefunction at time τ as:

$$\bar{\Psi}_n(x_1, x_2, \dots, x_N; \alpha(\tau)) = \frac{e^{i\Gamma(\tau)}}{\sqrt{N}} G(x_1, x_2, \dots, x_N; \tau) \begin{vmatrix} \Psi_{nk_1\alpha(0)}(x_1) & \Psi_{nk_2\alpha(0)}(x_1) & \cdots & \Psi_{nk_N\alpha(0)}(x_1) \\ \Psi_{nk_1\alpha(0)}(x_2) & \Psi_{nk_2\alpha(0)}(x_2) & \cdots & \Psi_{nk_N\alpha(0)}(x_2) \\ \vdots & \vdots & & \vdots \\ \Psi_{nk_1\alpha(0)}(x_N) & \Psi_{nk_2\alpha(0)}(x_N) & \cdots & \Psi_{nk_N\alpha(0)}(x_N) \end{vmatrix}.$$

Here, the N -particle large gauge transformation G is given by the product:

$$G(x_1, x_2, \dots, x_N; \tau) = \prod_{j=1}^N U^\dagger(x_j, \tau), \quad (48)$$

whereas the phase factor $\Gamma(\tau)$ reads:

$$\Gamma(\tau) = i \int_0^{\frac{2\pi}{\alpha}} d\alpha \langle u_n(\alpha) | \frac{\partial}{\partial \alpha} | u_n(\alpha) \rangle - \frac{1}{\hbar} \sum_{j=0}^{N-1} \int_0^\tau dt E_{nk_j\alpha(t)}. \quad (49)$$

Note that $\Psi_{nk_N\alpha}(x_j) = e^{i\chi} \Psi_{nk_0\alpha}(x_j)$ due to the Bloch property. We further use the anti-symmetric nature of the Slater determinant to rewrite the many-particle wavefunction at time τ :

$$\bar{\Psi}_n(x_1, x_2, \dots, x_N; \alpha(\tau)) = (-1)^{N-1} e^{i\chi} e^{i\Gamma(\tau)} \times G(x_1, x_2, \dots, x_N; \tau) \bar{\Psi}_n(x_1, x_2, \dots, x_N; \alpha(0)). \quad (50)$$

This shows that the wavefunction $\bar{\Psi}$ at times $t = 0$ and $t = \tau$ represents the same physical state with identical spectrum, which is the filled band n . So the evolution of the system in such a case should be understood as cyclic, albeit in the generalized sense as the single-particle case; since the system returns to its initial state after time τ , modulo a large gauge transformation. Recapitulating that the only gauge-invariant contribution to the total

phase gained by the system during time evolution is the geometric phase, allows one to straight away read off the geometric phase Γ_g acquired by the filled band n from (50) using (38), and it reads:

$$\Gamma_g = \gamma_g(n) + ((N-1) \bmod 2)\pi. \quad (51)$$

This remarkable expression tells us that the geometric phase acquired by the band n over the cyclic evolution after time τ is not only comprises of the Pancharatnam-Zak phase $\gamma_g(n)$, but also a contribution coming from the Fermi-Dirac statistics of the N -particle state, which is sensitive to odd/even nature of the number of states in the band. The geometric phase for filled bands has been studied for some time now [21, 22, 48]. However, such a contribution of Fermi-Dirac statistics to the geometric phase has not been reported. This is an example of the many-particle geometric phase, which has been studied earlier theoretically [49] in the context of intensity interferometry, followed by an experimental observation in the quantum optical regime [39] and as a classical light analog of the multi-particle Aharonov-Bohm effect [50]. While the theoretical predictions and the experiments of this effect were not in dispute; the interpretation was criticized [51] perhaps because it was formulated in the language of Feynman path integrals, rather than the quantum Hilbert space description.

V. EXPLICIT EXAMPLES

In this section, the calculation of the Pancharatnam-Zak phase for two popular soluble models, e.g., the SSH model and the Kitaev model of 1D p -wave superconductors, is presented. The aim is to show explicitly the validity of the conclusions drawn about the Pancharatnam-Zak phase in the previous sections.

A. SSH model

The SSH model is a 1D lattice of atoms with an unit cell consisting of two atoms. This model is formulated within the tight-binding approximation with nearest-neighbour couplings between the atoms [16, 32]. In the recent years, there are many experimental realizations of this model in photonics [29, 30, 33]. The Hamiltonian describing the model reads as:

$$\hat{H}_{SSH} = \sum_{m=-N/2}^{N/2} (-v|ma + r_\alpha\rangle\langle ma + r_\beta| - w|(m+1)a + r_\alpha\rangle\langle ma + r_\beta| + \text{h.c.}). \quad (52)$$

Here, $r_{\alpha,\beta}$ represent the coordinates of the two atoms respectively within the unit cell, whereas a is the lattice constant. The spatially localized electron state on atom in m^{th} unit cell at site $r_{\alpha,\beta}$ is described by $|ma + r_{\alpha,\beta}\rangle$, whereas the distance between the two atoms in the unit cell is $b = r_\beta - r_\alpha$. The parameter v is the intracell electron hopping amplitude, whereas w is the intercell hopping amplitude. The system consists of N cells with PBC. Going over to the momentum space allows one to define the free particle states $|k\rangle_{\alpha,\beta}$ as:

$$|k\rangle_{\alpha,\beta} = \frac{1}{\sqrt{N}} \sum_{m=-N/2}^{N/2} e^{ik(ma+r_{\alpha,\beta})} |ma + r_{\alpha,\beta}\rangle, \quad (53)$$

so that the above Hamiltonian reads:

$$\hat{H}_{SSH} = \sum_{k \in \text{FBZ}} (|k\rangle_\alpha, |k\rangle_\beta) [\mathcal{H}(k)] \begin{pmatrix} \langle \alpha | k \rangle \\ \langle \beta | k \rangle \end{pmatrix}. \quad (54)$$

Here, the sum is over all the allowed values of k in the first Brillouin zone (FBZ). The 2×2 matrix $[\mathcal{H}(k)]$ has only two non-vanishing off diagonal elements $\mathcal{H}_{\alpha\beta} = \mathcal{H}_{\beta\alpha}^* = x(k) + iy(k) = (-ve^{ikb} - we^{ik(b-a)})$. This can be diagonalized to find two eigenvalues $E_\pm(k) = \pm E(k)$, where $E(k) = \sqrt{x^2 + y^2} = \sqrt{v^2 + w^2 + 2vw \cos(ka)}$. The corresponding eigenvectors $[u_\pm(k)]$ are given by:

$$[u_\pm(k)] = \begin{pmatrix} u_{1,\pm}(k) \\ u_{2,\pm}(k) \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \pm \frac{x(k) + iy(k)}{E(k)} \\ 1 \end{pmatrix}. \quad (55)$$

Note that, there is an ambiguity (upto a local gauge transformation) in defining these eigenvectors, since $[u_\pm(k)]$ and $e^{i\theta(k)}[u_\pm(k)]$ (where $\theta(k)$ is any general function of k) both solve the eigenvalue problem for $[\mathcal{H}(k)]$ for the same eigenvalues. It follows that the Hamiltonian (52) diagonalises in terms of $|\Psi_\pm(k)\rangle$:

$$\hat{H}_{SSH} = \sum_{k \in \text{FBZ}} E_\pm(k) |\Psi_\pm(k)\rangle \langle \Psi_\pm(k)|, \quad (56)$$

which are defined as:

$$|\Psi_\pm(k)\rangle = u_{1,\pm}(k)|k\rangle_\alpha + u_{2,\pm}(k)|k\rangle_\beta.$$

One can now invoke relation (35) to find the Bloch states [52] $|u_\pm(\kappa)\rangle$ as:

$$|u_\pm(\kappa)\rangle = u_{1,\pm}(\kappa)|0\rangle_\alpha + u_{2,\pm}(\kappa)|0\rangle_\beta, \quad (57)$$

so that the Pancharatnam-Zak phase $\gamma_g(\pm)$ from (38) is given by:

$$\gamma_g(\pm) = \text{Arg} (u_{1,\pm}^*(0)u_{1,\pm}(2\pi/a) + u_{2,\pm}^*(0)u_{2,\pm}(2\pi/a)) + i \int_0^{2\pi/a} d\kappa (u_{1,\pm}^* \partial_\kappa u_{1,\pm} + u_{2,\pm}^* \partial_\kappa u_{2,\pm}). \quad (58)$$

Evidently, the ambiguity of local phase factor $e^{i\theta(\kappa)}$ in the definition of $|u_\pm(\kappa)\rangle$ does not affect $\gamma_g(\pm)$. It can be readily checked that the contributions from the first and second terms in the expression (58) arising from such a phase factor get exactly cancelled, displaying yet again the gauge invariance of the Pancharatnam-Zak phase. It can be immediately seen from (43) and from the above expression for $|u_\pm(\kappa)\rangle$, that the $\gamma_g(\pm)$ is also invariant under spatial translation operation.

This is to be contrasted with the range of values of the Zak phase $\gamma_Z(\pm)$ reported in literature [16, 23, 32, 33] which is arising due to the different choices in defining $|u_\pm(\kappa)\rangle$ and the choice of the origin of the unit cell.

Employing relation (58), we find that $\gamma_g(\pm)$ in this model takes two values: (a) it is equal to π when $v/w < 1$, and (b) it is equal to 0 when $v/w > 1$; where we have assumed that $b < a/2$. Interestingly, it turns out that when $b > a/2$, the system resembles itself with $b < a/2$

case, albeit with the roles of v and w now interchanged. Thus, one finds that (a) $\gamma_g(\pm) = \pi$ when $v/w > 1$, and (b) $\gamma_g(\pm) = 0$ when $v/w < 1$ for $b > a/2$.

The SSH model is thus found to exist in two discrete phases: the non-topological phase when $\gamma_g(\pm) = 0$, and the topological phase when $\gamma_g(\pm) = \pi$. This sharp change in the Pancharatnam-Zak phase shows the topological phase transition of this model.

In the above paragraphs, the spectrum as well as the geometric phase of the SSH model are derived using the PBC. However, it is the open boundary condition that is obeyed by the system in the physical realization of this model. Upon diagonalizing the Hamiltonian (52) numerically [16] with the open boundary condition, one finds that the system admits a degenerate pair of zero energy modes within the bulk-gap and localized at the edges when $v/w < 1$, and no zero mode for $v/w > 1$, irrespective of the value of b/a . As mentioned earlier, the SSH model, employing the PBC, is often only studied in the region $b \leq a/2$, in which case the topological phase of the model corresponds to the parameter regime $v/w < 1$. The fact that the SSH model admits zero modes (with open boundary condition) and non-trivial Pancharatnam-Zak phase ($= \pi$) (with PBC), leads one to think that this is an example of the bulk-boundary correspondence, in the sense, that the finite/zero value of Pancharatnam-Zak phase corresponds to the presence/absence of the zero energy edge modes. However, it is evident from the above discussion that the existence of such a correspondence in this model needs to be relooked, since the value of the Pancharatnam-Zak phase changes as a function of b/a , whereas the presence/absence of zero energy modes is indifferent to the changes in b/a .

It is worthwhile to consider the work of Atala *et al.*, wherein the experimental observation of the single-particle Zak phase in the SSH model was reported [33]. The SSH model in their experimental setup was realized in an optical lattice setup, and the difference in the geometric phase between the topological and non-topological phase of the system was observed to be equal to π . Such a phase difference is also in agreement with the difference of the calculated single-particle Pancharatnam-Zak phase across the topological transition point. Nevertheless, it would be interesting to experimentally measure the abso-

lute value of the geometric phase separately in the topological and non-topological phase of the model so as to validate the assertions made about the Pancharatnam-Zak phase.

B. Kitaev model of 1D p -wave superconductors

The Kitaev model is a mean-field Hamiltonian of 1D p -wave superconductors with triplet (p -wave) superconducting pairing between electrons with the same spin orientation. It is well known that this model, like SSH model, admits two distinct phases respectively having trivial and non-trivial topological property. This model in the topological phase hosts two zero energy Majorana bound states at the two ends of the system when defined with open boundary condition.

The Hamiltonian of the Kitaev model [15], with the PBC, in terms of the creation and destruction operators of electrons reads:

$$\hat{H}_K = \sum_{j=-N/2}^{N/2} (-J\hat{c}_j^\dagger\hat{c}_{j+1} + \Delta\hat{c}_j\hat{c}_{j+1} + \text{h.c.}) + \varepsilon(\hat{c}_j^\dagger\hat{c}_j - \frac{1}{2}). \quad (59)$$

Here, ε represents the on-site energy, J is the nearest-neighbour hopping amplitude, and Δ stands for the superconducting pairing amplitude; and all them are assumed to be real. The electron creation and destruction (Fermi) operators at site j , obey the anticommutation relations:

$$\{\hat{c}_j, \hat{c}_{j'}^\dagger\} = \delta_{j,j'}, \quad \text{and} \quad \{\hat{c}_j, \hat{c}_{j'}\} = 0. \quad (60)$$

These creation and annihilation operators can be decomposed in terms of Hermitian Majorana operators, $\hat{\eta}_{j\alpha} (= \hat{\eta}_{j\alpha}^\dagger)$ and $\hat{\eta}_{j\beta} (= \hat{\eta}_{j\beta}^\dagger)$ as:

$$\hat{c}_j = \frac{1}{2}(\hat{\eta}_{j\alpha} + i\hat{\eta}_{j\beta}), \quad (61)$$

for which the relation $\{\hat{\eta}_{j\alpha}, \hat{\eta}_{j'\beta}\} = 2\delta_{j,j'}\delta_{\alpha,\beta}$ holds. The Hamiltonian \hat{H}_K can now be expressed as:

$$\hat{H}_K = \sum_{j=-N/2}^{N/2} \frac{i}{4} \left(\varepsilon(\hat{\eta}_{j\alpha}\hat{\eta}_{j\beta} - \hat{\eta}_{j\beta}\hat{\eta}_{j\alpha}) - (J + \Delta)(\hat{\eta}_{j+1\alpha}\hat{\eta}_{j\beta} - \hat{\eta}_{j\beta}\hat{\eta}_{j+1\alpha}) + (\Delta - J)(\hat{\eta}_{j\alpha}\hat{\eta}_{j+1\beta} - \hat{\eta}_{j+1\beta}\hat{\eta}_{j\alpha}) \right). \quad (62)$$

The Majorana operators can be written in terms of their

k -space counterparts as:

$$\hat{\eta}_{j\alpha} = \frac{1}{\sqrt{N}} \sum_k e^{-ikaj} \hat{\eta}_{k\alpha}, \quad (63)$$

$$\hat{\eta}_{j\beta} = \frac{1}{\sqrt{N}} \sum_k e^{-ikaj} \hat{\eta}_{k\beta}, \quad (64)$$

which are such that, $\hat{\eta}_{-k\alpha} = \hat{\eta}_{k\alpha}^\dagger$, $\hat{\eta}_{-k\beta} = \hat{\eta}_{k\beta}^\dagger$, and

$$\{\hat{\eta}_{k\alpha}, \hat{\eta}_{-k'\beta}\} = 2\delta_{k,k'}\delta_{\alpha,\beta}. \quad (65)$$

The operators $\hat{\eta}_{-k\alpha}$ and $\hat{\eta}_{-k\beta}$ respectively create single (Majorana) particle state $|k\rangle_{\alpha,\beta} = \hat{\eta}_{-k\alpha,\beta}|0\rangle_{\alpha,\beta}$, with wave vector k . The Hamiltonian \hat{H}_K can now be expressed in terms of these k -space operators as:

$$\hat{H}_K = \sum_{k \in FBZ} \begin{pmatrix} \hat{\eta}_{k\alpha}^\dagger & \hat{\eta}_{k\beta}^\dagger \end{pmatrix} [\mathcal{H}(k)] \begin{pmatrix} \hat{\eta}_{k\alpha} \\ \hat{\eta}_{k\beta} \end{pmatrix}. \quad (66)$$

The 2×2 matrix $[\mathcal{H}(k)]$ has only non-zero elements $\mathcal{H}_{\alpha\beta} = \mathcal{H}_{\beta\alpha}^* = X(k) + iY(k)$, where $X(k) = \frac{\Delta}{2} \sin(ka)$, and $Y(k) = \frac{\varepsilon}{4} - \frac{J}{2} \cos(ka)$. The matrix $[\mathcal{H}(k)]$ when diagonalized yields two bands of eigenvalues: $E_\pm(k) = \pm E(k)$, where $E(k) = \sqrt{X^2(k) + Y^2(k)}$. The corresponding eigenvectors are given by $[u_\pm(k)]$:

$$[u_\pm(k)] = \begin{pmatrix} u_{1,\pm}(k) \\ u_{2,\pm}(k) \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \pm \frac{X(k) + iY(k)}{E(k)} \\ 1 \end{pmatrix}. \quad (67)$$

As was noted in the earlier case of the SSH model, these eigenvectors are ambiguous upto a local phase factor $e^{i\theta(k)}$. The Hamiltonian \hat{H}_K in the diagonal form reads:

$$\hat{H}_K = \sum_{k \in FBZ} \left(E_\pm(k) \hat{\Psi}_\pm^\dagger(k) \hat{\Psi}_\pm(k) \right), \quad (68)$$

where the field operator $\hat{\Psi}_\pm^\dagger(k) = u_{1,\pm}(k) \hat{\eta}_{-k\alpha} + u_{2,\pm}(k) \hat{\eta}_{-k\beta}$. From here, it follows that the energy eigenstates are:

$$|\Psi_\pm(k)\rangle = u_{1,\pm}(k)|k\rangle_\alpha + u_{2,\pm}(k)|k\rangle_\beta, \quad (69)$$

and the corresponding cell periodic Bloch states $|u_\pm(\kappa)\rangle$ are given by:

$$|u_\pm(\kappa)\rangle = u_{1,\pm}(\kappa)|0\rangle_\alpha + u_{2,\pm}(\kappa)|0\rangle_\beta. \quad (70)$$

It is easy to see that the Pancharatnam-Zak phase γ_g in this model can be evaluated exactly as the earlier model, using (58), with the expression of $u_{1,2}(\kappa)$ now given by (67).

It is found that the Pancharatnam-Zak phase in this model takes two values: (a) $\gamma_g(\pm) = 0$ when $|\varepsilon| > 2J$, and (b) $\gamma_g(\pm) = \pi$ when $|\varepsilon| < 2J$.

VI. CONCLUSION

In this paper, we have attempted to provide a correct and consistent understanding of the notion of the geometric phase in 1D periodic lattice systems weakly perturbed by an electric field. The expression of the Pancharatnam-Zak phase, which is the geometric phase acquired by a single electron in such a system, is found, and the underlying mathematical structure is unveiled. It is shown

that the Pancharatnam-Zak phase is an example of cyclic adiabatic geometric phase, albeit the concept of cyclicity of time evolution in this system is a generalization of the existing notion.

For a given system, if the density matrix over time evolution returns to its initial state, then the time evolution is considered to be cyclic. However, in the case of systems that were considered in this work, the density matrix in the course of evolution never actually returns to its initial state but instead returns with a topologically non-trivial large gauge transformation. So the time evolution is not strictly cyclic; nevertheless it turns out that, owing to the nature of the gauge transformation, all the physical observables, like expectation values, return to their respective initial values in the course of evolution. It shows that evolution should be considered cyclic, albeit in a generalized sense. In these cases, however, the concept of the geometric phase, as pioneered by Berry, can not be applied, since it is only defined for the systems displaying strict cyclicity [7]. Thus one is naturally drawn to work with the notion of geometric phase defined in its generality, which is meaningful even while the system does not display strict cyclicity.

The Pancharatnam-Zak phase, so constructed keeping in mind the above consideration, is gauge invariant and spatial translation invariant. In the case of systems with inversion symmetry, it is found that Pancharatnam-Zak phase can either be equal to 0 or π . An explicit calculation of this geometric phase is demonstrated for the SSH model and Kitaev model. It would be exciting to experimentally confirm the calculated values of the single-particle Pancharatnam-Zak phase separately in the topological and trivial phases of the SSH and Kitaev model by generalizing current experimental schemes [33] which have already probed the difference in geometric phases.

A many-particle generalization of this cyclic adiabatic geometric phase for a filled band case is obtained, and remarkably it is found to receive a contribution owing to the Fermi-Dirac statistics of the band electrons. This statistical contribution to the geometric phase is completely missed in the previous treatments dealing with filled band systems which are common in condensed matter physics [15, 16]. It would also be interesting to observe such statistical contribution to the geometric phase in the experimental set-up of Atala *et al.* [33] with multiple fermionic cold atoms in optical lattices.

ACKNOWLEDGMENTS

We thank Dr. Nikolai Sinitsyn for a critical reading of our manuscript. DR gratefully acknowledges the funding from the Department of Science and Technology, India via the Ramanujan Fellowship.

ALTERNATE DERIVATION OF PANCHARATNAM-ZAK PHASE

It has been known since long that the notion of geometric phase γ_g is essentially a kinematic one, and can be defined for any given set of states $|a_i\rangle$ (where $i = 0, 1, 2, \dots, M$), as the argument of the cyclic object D_M [35, 36] called the Bargmann invariant:

$$D_M = \langle a_0 | a_M \rangle \langle a_M | a_{M-1} \rangle \cdots \langle a_2 | a_1 \rangle \langle a_1 | a_0 \rangle.$$

The expression for the Pancharatnam-Zak phase can be straight away obtained using this route. Let the Bargmann invariant be defined as:

$$\Delta_M = \langle u_{n,0} | u_{n,M} \rangle \langle u_{n,M} | u_{n,M-1} \rangle \cdots \langle u_{n,2} | u_{n,1} \rangle \langle u_{n,1} | u_{n,0} \rangle,$$

where $|u_{n,i}\rangle \equiv |u_n(\frac{2\pi i}{Ma})\rangle$. The expression (38) for Pancharatnam-Zak phase can be seen to be obtained as $\gamma_g(n) = \text{Arg } \Delta_M$, in the limit $M \rightarrow \infty$. The cyclic nature of Δ_M ensures that it is invariant under local gauge

transformations $|u_{n,j}\rangle \rightarrow e^{i\Lambda_j} |u_{n,j}\rangle$ (here Λ_j are some arbitrary real numbers), and under unitary operation both of which preserve the value of the amplitudes in Δ_M .

It is argued in the literature [17, 22] that the Zak phase (39) can be obtained from the argument of the non-cyclic object as:

$$\gamma_Z(n) = \text{Arg} \langle u_{n,N} | u_{n,N-1} \rangle \cdots \langle u_{n,2} | u_{n,1} \rangle \langle u_{n,1} | u_{n,0} \rangle.$$

It follows from (13) that:

$$|u_{n,N}\rangle = e^{i\chi} e^{-i\frac{2\pi}{a}\hat{x}} |u_{n,0}\rangle,$$

which allows $\gamma_Z(n)$ to be written as:

$$\gamma_Z(n) = \text{Arg} \langle u_{n,0} | e^{-i\chi} e^{i\frac{2\pi}{a}\hat{x}} | u_{n,N-1} \rangle \cdots \langle u_{n,2} | u_{n,1} \rangle \langle u_{n,1} | u_{n,0} \rangle.$$

The explicit occurrence of phase χ clearly shows that the absolute value of the Zak phase is arbitrary, since χ can be fixed to any desired value by a gauge convention. This evidently shows that the Zak phase can not be an observable phase, since the value of the physical observables are required to be independent of the choice of convention.

-
- [1] J. Anandan, The geometric phase, *Nature* **360**, 307 (1992).
- [2] S. Pancharatnam, Generalized theory of interference and its applications, *Proc. Indian Acad. of Sci. A* **44**, 398 (1956).
- [3] M. V. Berry, The adiabatic phase and Pancharatnam's phase for polarized light, *J. Mod. Opt.* **34**, 1401 (1987).
- [4] C. A. Mead and D. G. Truhlar, On the determination of Born-Oppenheimer nuclear motion wave functions including complications due to conical intersections and identical nuclei, *J. Chem. Phys.* **70**, 2284 (1979).
- [5] G. Herzberg and H. Longuet-Higgins, Intersection of potential energy surfaces in polyatomic molecules, *Discuss. Faraday Soc.* **35**, 77 (1963).
- [6] Y. Aharonov and D. Bohm, Significance of electromagnetic potentials in the quantum theory, *Phys. Rev.* **115**, 485 (1959).
- [7] M. V. Berry, Quantal phase factors accompanying adiabatic changes, *Proc. Roy. Soc. Lond. A* **392**, 45 (1984).
- [8] A. Shapere and F. Wilczek, *Geometric phases in physics* (World Scientific, 1989).
- [9] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, Quantized Hall conductance in a two-dimensional periodic potential, *Phys. Rev. Lett.* **49**, 405 (1982).
- [10] Q. Niu, D. J. Thouless, and Y.-S. Wu, Quantized Hall conductance as a topological invariant, *Phys. Rev. B* **31**, 3372 (1985).
- [11] B. Simon, Holonomy, the quantum adiabatic theorem, and Berry's phase, *Phys. Rev. Lett.* **51**, 2167 (1983).
- [12] F. D. M. Haldane, Model for a quantum Hall effect without Landau levels: Condensed-matter realization of the "parity anomaly", *Phys. Rev. Lett.* **61**, 2015 (1988).
- [13] F. D. M. Haldane, Berry curvature on the fermi surface: Anomalous Hall effect as a topological Fermi-liquid property, *Phys. Rev. Lett.* **93**, 206602 (2004).
- [14] D. J. Thouless, Quantization of particle transport, *Phys. Rev. B* **27**, 6083 (1983).
- [15] B. A. Bernevig and T. L. Hughes, *Topological insulators and topological superconductors* (Princeton University, 2013).
- [16] J. K. Asbóth, L. Oroszlány, and A. Pályi, *A short course on topological insulators*, Vol. 919 (Springer, 2016).
- [17] D. Vanderbilt, *Berry Phases in Electronic Structure Theory: Electric Polarization, Orbital Magnetization and Topological Insulators* (Cambridge University Press, 2018).
- [18] M. Z. Hasan and C. L. Kane, Colloquium: Topological insulators, *Rev. Mod. Phys.* **82**, 3045 (2010).
- [19] J. Zak, Berry's phase for energy bands in solids, *Phys. Rev. Lett.* **62**, 2747 (1989).
- [20] R. D. King-Smith and D. Vanderbilt, Theory of polarization of crystalline solids, *Phys. Rev. B* **47**, 1651 (1993).
- [21] R. Resta, Macroscopic polarization in crystalline dielectrics: the geometric phase approach, *Rev. Mod. Phys.* **66**, 899 (1994).
- [22] R. Resta, Manifestations of Berry's phase in molecules and condensed matter, *J. Phys. C* **12**, R107 (2000).
- [23] D. Xiao, M.-C. Chang, and Q. Niu, Berry phase effects on electronic properties, *Rev. Mod. Phys.* **82**, 1959 (2010).
- [24] T. Goren, K. Plekhanov, F. Appas, and K. Le Hur, Topological Zak phase in strongly coupled LC circuits, *Phys. Rev. B* **97**, 041106(R) (2018).
- [25] S. Longhi, Zak phase of photons in optical waveguide lattices, *Opt. Lett.* **38**, 3716 (2013).
- [26] P. Delplace, D. Ullmo, and G. Montambaux, Zak phase and the existence of edge states in graphene, *Phys. Rev. B* **84**, 195452 (2011).
- [27] T. Kameda, F. Liu, S. Dutta, and K. Wakabayashi, Topological edge states induced by the Zak phase in A_3B

- monolayers, *Phys. Rev. B* **99**, 075426 (2019).
- [28] S. Sarkar, Quantization of geometric phase with integer and fractional topological characterization in a quantum Ising chain with long-range interaction, *Sci. Rep.* **8**, 5864 (2018).
- [29] P. St-Jean, V. Goblot, E. Galopin, A. Lemaître, T. Ozawa, L. Le Gratiet, I. Sagnes, J. Bloch, and A. Amo, Lasing in topological edge states of a one-dimensional lattice, *Nature Photon.* **11**, 651 (2017).
- [30] A. Blanco-Redondo, I. Andonegui, M. J. Collins, G. Harari, Y. Lumer, M. C. Rechtsman, B. J. Eggleton, and M. Segev, Topological optical waveguiding in Silicon and the transition between topological and trivial defect states, *Phys. Rev. Lett.* **116**, 163901 (2016).
- [31] R. Bhandari and J. Samuel, Observation of topological phase by use of a laser interferometer, *Phys. Rev. Lett.* **60**, 1211 (1988).
- [32] N. R. Cooper, J. Dalibard, and I. B. Spielman, Topological bands for ultracold atoms, *Rev. Mod. Phys.* **91**, 015005 (2019).
- [33] M. Atala, M. Aidelsburger, J. T. Barreiro, D. Abanin, T. Kitagawa, E. Demler, and I. Bloch, Direct measurement of the Zak phase in topological Bloch bands, *Nat. Phys.* **9**, 795 (2013).
- [34] J. E. Moore, An introduction to topological phases of electrons, in *Lecture Notes of the Les Houches Summer School, August 2014*, Vol. 103, edited by C. Chamon, M. O. Goerbig, R. Moessner, and L. F. Cugliandolo (Oxford University Press, 2017) pp. 3–61.
- [35] J. Samuel and R. Bhandari, General setting for Berry's phase, *Phys. Rev. Lett.* **60**, 2339 (1988).
- [36] N. Mukunda and R. Simon, Quantum kinematic approach to the geometric phase. I. General formalism, *Ann. Phys. (N.Y.)* **228**, 205 (1993).
- [37] It must be noted that the total phase can only be defined if the initial and final states are non-orthogonal.
- [38] T. Chyba, L. Wang, L. Mandel, and R. Simon, Measurement of the Pancharatnam phase for a light beam, *Opt. Lett.* **13**, 562 (1988).
- [39] A. Martin, O. Alibart, J.-C. Flesch, J. Samuel, S. Sinha, S. Tanzilli, and A. Kastberg, Non-local geometric phase in two-photon interferometry, *Europhys. Lett.* **97**, 10003 (2012).
- [40] The momentum translation operator translates momentum eigenstates $|q - k\rangle = \hat{T}_p(k)|q\rangle$, where $p|q\rangle = \hbar q|q\rangle$.
- [41] M. Nakahara, *Geometry, topology and physics* (CRC Press, 2003).
- [42] A. Messiah, *Quantum Mechanics, Two Volumes* (Dover Publications, 1999).
- [43] D. M. Tong, K. Singh, L. C. Kwek, and C. H. Oh, Sufficiency criterion for the validity of the adiabatic approximation, *Phys. Rev. Lett.* **98**, 150402 (2007).
- [44] This identification is valid in the $N \rightarrow \infty$ limit only, so that the parameter α can be identified with k in (39).
- [45] This justifies the nomenclature for $\gamma_g(n)$ as Pancharatnam-Zak phase.
- [46] In the literature often the Zak phase $\gamma_Z(n)$ is also expressed in the form a closed line integral, which is incorrect as is evident from here.
- [47] In fact $\gamma_Z(n) \rightarrow \gamma_Z(n) + \frac{2\pi}{\alpha}d$ under such a spatial translation.
- [48] E. Yaschenko, L. Fu, L. Resca, and R. Resta, Macroscopic polarization as a discrete Berry phase of the Hartree-Fock wave function: The single-point limit, *Phys. Rev. B* **58**, 1222 (1998).
- [49] P. Mehta, J. Samuel, and S. Sinha, Nonlocal Pancharatnam phase in two-photon interferometry, *Phys. Rev. A* **82**, 034102 (2010).
- [50] N. Satapathy, D. Pandey, P. Mehta, S. Sinha, J. Samuel, and H. Ramachandran, Classical light analogue of the non-local Aharonov-Bohm effect, *Europhys. Lett.* **97**, 50011 (2012).
- [51] Arvind, S. Chaturvedi, and N. Mukunda, A classical optical approach to the non-local Pancharatnam-like phases in Hanbury-Brown-Twiss correlations, *Phys. Lett. A* **381**, 1272 (2017).
- [52] The parameter κ is a continuous parameter, and is not to be confused as the wave vector.