

## TOTAL POSITIVITY IN SPRINGER FIBRES

G. LUSZTIG

## CONTENTS

1. The totally positive part of  $\mathcal{B}_u$ .
2. Examples.
3. Partial flag manifolds.
4. A conjecture and its consequences.
5. The map  $g \mapsto P_g$  from  $G_{\geq 0}$  to  $\mathcal{P}_{\geq 0}$ .

1. THE TOTALLY POSITIVE PART OF  $\mathcal{B}_u$ 

**1.1.** Let  $G$  be a reductive connected algebraic group over  $\mathbf{C}$  and let  $\mathcal{B}$  be the variety of Borel subgroups of  $G$ . For  $g \in G$  the Springer fibre at  $g$  is the subvariety  $\mathcal{B}_g = \{B \in \mathcal{B}; g \in B\}$  of  $\mathcal{B}$ . The Springer fibres can be quite complicated; they play an important role in representation theory (for example in character formulas for finite reductive groups over a finite field). In this paper we assume that a pinning of  $G$  is given so that the totally positive part  $G_{\geq 0}$  of  $G$  and the totally positive part  $\mathcal{B}_{\geq 0}$  of  $\mathcal{B}$  are defined (see [L94, 2.2, 8.1]). Recall that  $G_{\geq 0}$  is a submonoid and a closed subset of  $G$  and  $\mathcal{B}_{\geq 0}$  is a closed subset of  $\mathcal{B}$  on which  $G_{\geq 0}$  acts. We are interested in the interaction of the theory of total positivity with that of Springer fibres. More precisely, for any  $g \in G_{\geq 0}$  we consider the closed subset  $\mathcal{B}_{g, \geq 0} = \mathcal{B}_g \cap \mathcal{B}_{\geq 0}$  of  $\mathcal{B}_g$ , which we call the totally positive part of the Springer fibre  $\mathcal{B}_g$ . Let  $\mathcal{U}$  be the variety of unipotent elements of  $G$  and let  $\mathcal{U}_{\geq 0} = \mathcal{U} \cap G_{\geq 0}$ . Our main result is that if  $g \in \mathcal{U}_{\geq 0}$ ,  $\mathcal{B}_{g, \geq 0}$  has a surprisingly simple structure, much simpler than that of  $\mathcal{B}_g$  itself (see Cor. 1.16); namely it has a canonical cell decomposition which is part of the canonical cell decomposition of  $\mathcal{B}_{\geq 0}$ . By a similar argument we show that an analogous result holds for the fixed point set of  $g$  on a partial flag manifold intersected with the totally positive part [L98] of that partial flag manifold (see 3.9). In §4 we state a conjecture (see 4.4) on the compatibility of two ways to define a positive structure on the cells of  $\mathcal{B}_{\geq 0}$  and give some of its consequences; this conjecture extends a result of [L97, §3] which concerns the open dense cell  $\mathcal{B}_{>0}$  of  $\mathcal{B}_{\geq 0}$  defined in [L94, 8.8]. Let  $G_{>0}$  the open dense sub-semigroup of  $G_{\geq 0}$  defined in [L94]. In [L94, 8.9(c)], a map  $g \mapsto B_g$  from

---

Supported by NSF grant DMS-1855773.

$G_{>0}$  to  $\mathcal{B}_{>0}$  was defined. In §5 we give a definition of this map which is simpler than that in [L94] and we define an extension of this map to a map from  $G_{\geq 0}$  to the set of parabolic subgroups of  $G$ . As a result we obtain a new definition of  $\mathcal{B}_{>0}$ .

We shall always assume that  $G$  is simply laced; the non-simply laced case can be reduced to the simply laced case by descent.

**1.2.** Recall that we have fixed a pinning of  $G$ . Thus we are given a maximal torus  $T$  of  $G$  and a pair  $B^+, B^-$  of opposed Borel subgroups of  $G$  containing  $T$  with unipotent radicals  $U^+, U^-$ . The pinning also includes root homomorphisms  $x_i : \mathbf{C} \rightarrow U^+$ ,  $y_i : \mathbf{C} \rightarrow U^-$  indexed by a finite set  $I$  (corresponding to simple roots). Let  $NT$  be the normalizer of  $T$  in  $G$  and let  $W$  be the Weyl group. For  $i \in I$  we set  $\dot{s}_i = y_i(1)x_i(-1)y_i(1) \in NT$ ; let  $s_i$  be the image of  $\dot{s}_i$  in  $W$ . Now  $W$  is a Coxeter group with simple reflections  $\{s_i; i \in I\}$ ; let  $w \mapsto |w|$  be the standard length function. Let  $w_I$  be the unique element of maximal length of  $W$ . More generally, for any  $H \subset I$  we denote by  $w_H$  the longest element in the subgroup  $W_H$  of  $W$  generated by  $\{s_i; i \in H\}$ .

Let  $\leq$  be the standard partial order on  $W$ . For  $w \in W$  let  $\mathcal{I}_w$  be the set of sequences  $(i_1, i_2, \dots, i_m)$  in  $I$  such that  $w = s_{i_1}s_{i_2}\dots s_{i_m}$ ,  $m = |w|$ ; let  $\text{supp}(w)$  be the set of all  $i \in I$  which appear in some (or equivalently any)  $(i_1, i_2, \dots, i_m) \in \mathcal{I}_w$ . For  $w \in W$  we set  $\dot{w} = \dot{s}_{i_1}\dot{s}_{i_2}\dots\dot{s}_{i_m} \in NT$  where  $(i_1, i_2, \dots, i_m) \in \mathcal{I}_w$ ; this is known to be well defined.

For  $B, B'$  in  $\mathcal{B}$  there is a unique  $w \in W$  denoted by  $\text{pos}(B, B')$  such that for some  $g \in G$  we have  $gBg^{-1} = B^+$ ,  $gB'g^{-1} = \dot{w}B^+\dot{w}^{-1}$ . There is a unique isomorphism  $\phi : G \xrightarrow{\sim} G$  such that  $\phi(x_i(a)) = y_i(a)$ ,  $\phi(y_i(a)) = x_i(a)$  for all  $i \in I, a \in \mathbf{C}$  and  $\phi(t) = t^{-1}$  for all  $t \in T$ . This carries Borel subgroups to Borel subgroups hence induces an isomorphism  $\phi : \mathcal{B} \rightarrow \mathcal{B}$  such that  $\phi(B^+) = B^-$ ,  $\phi(B^-) = B^+$ . For  $i \in I$  we have  $\phi(\dot{s}_i) = x_i(1)y_i(-1)x_i(1) = \dot{s}_i^{-1}$ . Hence  $\phi$  induces the identity map on  $W$ . We show:

(a) *Let  $B, B'$  in  $\mathcal{B}$  be such that  $\text{pos}(B, B') = w$ . Then  $\text{pos}(\phi(B), \phi(B')) = w_I w w_I$ .*

Assume first that for some  $i \in I$  we have  $B = B^+, B' = \dot{s}_i B^+ \dot{s}_i^{-1}$  so that  $\text{pos}(B, B') = s_i$ . We have

$$\begin{aligned} \text{pos}(\phi(B), \phi(B')) &= \text{pos}(B^-, \dot{s}_i^{-1} B^+ \dot{s}_i) = \text{pos}(\dot{w}_I B^+ \dot{w}_I^{-1}, \dot{s}_i^{-1} \dot{w}_I B^+ \dot{w}_I^{-1} \dot{s}_i) \\ &= \text{pos}(B^+, \dot{w}_I^{-1} \dot{s}_i^{-1} \dot{w}_I B^+ \dot{w}_I^{-1} \dot{s}_i \dot{w}_I) = w_I s_i w_I. \end{aligned}$$

It follows that (a) holds when  $w = s_i$  for some  $i \in I$ . The general case can be easily reduced to this special case. This proves (a).

**1.3.** Let  $V$  be an irreducible rational  $G$ -module over  $\mathbf{C}$  with a given highest weight vector  $\eta$ . Let  $\beta$  be the basis of  $V$  (containing  $\eta$ ) obtained by specializing at  $v = 1$  the canonical basis [L90] of the corresponding module over the corresponding quantized enveloping algebra. For any  $B \in \mathcal{B}$  let  $L_B$  be the unique  $B$ -stable line in  $V$ . We can assume that  $V$  is such that  $B \mapsto L_B$  is a bijection

(a)  $\mathcal{B} \xrightarrow{\sim} \mathcal{X}$

onto a subset  $\mathcal{X}$  of the set of lines in  $V$ . Let

(b)  $V_+ = \sum_{b \in \beta} \mathbf{R}_{\geq 0} b \subset V$ .

If  $\xi_1 \in V_+, \xi_2 \in V_+$  then  $\xi_1 + \xi_2 \in V_+$ ; moreover,

(c) if  $\xi_1 + \xi_2 = 0$ , then  $\xi_1 = \xi_2 = 0$ .

**1.4.** Following [KL], for any  $v, w$  in  $W$  we set  $\mathcal{B}_{v,w} = \{B \in \mathcal{B}; \text{pos}(B^+, B) = w, \text{pos}(B^-, B) = w_I v\}$ . It is known that  $\mathcal{B}_{v,w} \neq \emptyset$  if and only if  $v \leq w$ . The subvarieties  $\mathcal{B}_{v,w}$  with  $v \leq w$  form a partition of  $\mathcal{B}$ . For  $v \leq w$  in  $W$  we have  $ww_I \leq vw_I$ ; moreover,

(a)  $\phi$  defines an isomorphism  $\mathcal{B}_{ww_I, vw_I} \xrightarrow{\sim} \mathcal{B}_{v,w}$ .

We use that for  $B \in \mathcal{B}$  we have:

$$\text{pos}(B^+, B) = vw_I \implies \text{pos}(B^-, \phi(B)) = w_I v; \text{pos}(B^-, B) = w_I ww_I \implies \text{pos}(B^+, \phi(B)) = w.$$

**1.5.** As in [L94] let  $G_{\geq 0}$  be the submonoid of  $G$  generated by  $\{x_i(a); i \in I, a \in \mathbf{R}_{\geq 0}\}$ ,  $\{y_i(a); i \in I, a \in \mathbf{R}_{\geq 0}\}$  and by  $\chi(a)$  for various algebraic group homomorphisms  $\chi : \mathbf{C}^* \rightarrow T$  and various  $a \in \mathbf{R}_{> 0}$ ; this is a closed subset of  $G$ . We have

(a)  $\phi(G_{\geq 0}) = G_{\geq 0}$ .

By [L94, 3.2] we have in the setup of 1.3:

(b)  $G_{\geq 0} V_+ \subset V_+$ .

From [L94, 8.7] it follows that

(c)  $\phi(\mathcal{B}_{\geq 0}) = \mathcal{B}_{\geq 0}$ .

Let  $\mathcal{X}_{\geq 0}$  be the set of lines in  $\mathcal{X}$  which meet  $V_+ - \{0\}$ . From [L94, 8.17] we have that:

(d) under the bijection 1.3(a),  $\mathcal{B}_{\geq 0}$  corresponds to  $\mathcal{X}_{\geq 0}$ .

Using (b), (d) we deduce

(e)  $g \in G_{\geq 0}, B \in \mathcal{B}_{\geq 0} \implies gBg^{-1} \in \mathcal{B}_{\geq 0}$ .

(This also follows from [L94, 8.12].)

**1.6.** Recall that  $\mathcal{U}_{\geq 0} = \mathcal{U} \cap G_{\geq 0}$  and that for  $u \in \mathcal{U}_{\geq 0}$  we have  $\mathcal{B}_{u, \geq 0} = \mathcal{B}_u \cap \mathcal{B}_{\geq 0}$ . Let  $V, V_+$  be as in 1.3. For  $u \in \mathcal{U}_{\geq 0}$  let  $V_+^u = \{\xi \in V_+; u\xi = \xi\}$ . In particular  $V_+^{x_i(a)}, V_+^{y_i(a)}$  are defined for  $i \in I, a \in \mathbf{R}_{> 0}$ . We show:

(a) Let  $u, u', u''$  in  $\mathcal{U}_{\geq 0}$  be such that  $u = u'u''$ . We have  $V_+^u = V_+^{u'} \cap V_+^{u''}$ .

Let  $\xi \in V_+^{u'} \cap V_+^{u''}$ . Then  $u\xi = u'(u''\xi) = u'\xi = \xi$  hence  $\xi \in V_+^u$ . Conversely, let  $\xi \in V_+^u$ . By [L94, 6.2, 3.2] for any  $b \in \beta$  we have  $u'b - b \in V_+, u''b - b \in V_+$ . Hence for any  $\xi', \xi''$  in  $V_+$  we have  $u'\xi' - \xi' \in V_+, u''\xi'' - \xi'' \in V_+$ . Thus  $u''\xi = \xi + \xi'$  where  $\xi' \in V_+$  so that  $\xi = u\xi = u'(u''\xi) = u'(\xi + \xi')$ . We have  $u'\xi = \xi + \xi'_1, u'\xi' = \xi' + \xi'_2$  where  $\xi'_1 \in V_+, \xi'_2 \in V_+$ . Thus  $\xi = \xi + \xi'_1 + \xi' + \xi'_2$  so that  $\xi'_1 + \xi' + \xi'_2 = 0$ ; using 1.3(c) we see that  $\xi'_1 = \xi' = \xi'_2 = 0$ . Thus we have  $u''\xi = \xi, u'\xi = \xi$  and  $\xi \in V_+^{u'} \cap V_+^{u''}$ . This proves (a).

We show:

(b) Let  $i \in I, a \in \mathbf{R}_{>0}$ . We have  $V_+^{x_i(a)} = V_+^{x_i(1)}$ ,  $V_+^{y_i(a)} = V_+^{y_i(1)}$ .

Note that  $x_i(a) = x_i(1)^a : V \rightarrow V$  and  $x_i(1) = x_i(a)^{1/a} : V \rightarrow V$ . Hence we have  $\xi \in V_+^{x_i(a)}$  if and only if  $\xi \in V_+^{x_i(1)}$ . This proves the first equality in (b). The proof of the second equality is entirely similar.

Let  $u \in \mathcal{U}_{\geq 0}$ . From 1.5(d) we see that under the bijection 1.3(a),  $\mathcal{B}_{u,\geq 0}$  corresponds to the set of lines  $L$  in  $\mathcal{X}_{\geq 0}$  which are  $u$ -stable. Since  $u : V \rightarrow V$  is unipotent, for any such  $L$ ,  $u$  automatically acts on  $L$  as identity. Thus we have:

(c) Under the bijection 1.3(a),  $\mathcal{B}_{u,\geq 0}$  corresponds to the set of lines in  $\mathcal{X}$  which meet  $V_+^u - \{0\}$ .

**1.7.** Following [L94, 8.15] we define a partition  $\mathcal{B}_{\geq 0} = \sqcup_{(v,w) \in W \times W; v \leq w} \mathcal{B}_{\geq 0,v,w}$  by  $\mathcal{B}_{\geq 0,v,w} = \mathcal{B}_{v,w} \cap \mathcal{B}_{\geq 0}$ . In [L94, 8.15] it was conjectured that for  $v \leq w$ ,

(a)  $\mathcal{B}_{\geq 0,v,w}$  is homeomorphic to  $\mathbf{R}_{>0}^{|w|-|v|}$ .

This conjecture was proved by Rietsch [R98], [R99] and in a more explicit form by Marsh and Rietsch [MR].

For  $v \leq w$  let  $[v, w]$  be the subset of  $\mathcal{X}_{\geq 0}$  corresponding to  $\mathcal{B}_{\geq 0,v,w}$  under the bijection  $\mathcal{B}_{\geq 0} \leftrightarrow \mathcal{X}_{\geq 0}$  in 1.5(d). We have  $\mathcal{X}_{\geq 0} = \sqcup_{(v,w) \in W \times W; v \leq w} [v, w]$ .

**1.8.** Let  $v \leq w$  in  $W$  and let  $\mathbf{i} = (i_1, i_2, \dots, i_m) \in \mathcal{I}_w$ . According to Marsh and Rietsch [MR], there is a unique sequence  $t_1, t_2, \dots, t_m$  with  $t_k \in \{s_{i_k}, 1\}$  for  $k \in [1, m]$ ,  $t_1 t_2 \dots t_m = v$  and such that  $t_1 \leq t_1 t_2 \leq \dots \leq t_1 t_2 \dots t_m$  and  $t_1 \leq t_1 s_{i_2}, t_1 t_2 \leq t_1 t_2 s_{i_3}, \dots, t_1 t_2 \dots t_{m-1} \leq t_1 t_2 \dots t_{m-1} s_{i_m}$ . Following [MR] we define a subset  $\mathcal{Y}_{v,w,\mathbf{i}}$  of  $G$  to be the set of products  $g_1 g_2 \dots g_m$  in  $G$  where  $g_k = s_{i_k}$  if  $t_k = s_{i_k}$  and  $g_k = y_{i_k}(a_k)$  with  $a_k \in \mathbf{R}_{>0}$  if  $t_k = 1$ ; according to [MR], the map  $\mathcal{Y}_{v,w,\mathbf{i}} \rightarrow \mathcal{B}$ ,  $g \mapsto g B^+ g^{-1}$  is a homeomorphism

(a)  $\mathcal{Y}_{v,w,\mathbf{i}} \xrightarrow{\sim} \mathcal{B}_{\geq 0,v,w}$ .

Moreover we have a homeomorphism

(b)  $\mathcal{Y}_{v,w,\mathbf{i}} \xrightarrow{\sim} \mathbf{R}_{>0}^{|w|-|v|}$

(to  $g_1 g_2 \dots g_m$  we associate the sequence consisting of the  $a_k$  with  $k$  such that  $t_k = 1$ .) In particular, the factors  $g_1, g_2, \dots, g_m$  are uniquely determined by their product  $g_1 g_2 \dots g_m$ . The composition of the inverse of (b) with (a) is a homeomorphism

(c)  $\tau_{\mathbf{i}} : \mathbf{R}_{>0}^{|w|-|v|} \xrightarrow{\sim} \mathcal{B}_{\geq 0,v,w}$ .

**1.9.** We preserve the setup of 1.8. Assume that  $s_{i_1} w \leq w$ . We show:

(a) We have  $t_1 = 1$  if and only if  $v \leq s_{i_1} w$ .

If  $t_1 = 1$  then from  $v = t_1 \dots t_m$  we deduce  $v = t_2 t_3 \dots t_m$ ; thus  $v$  is equal to a product of a subsequence of  $s_{i_2}, s_{i_3}, \dots, s_{i_m}$  so that  $v \leq s_{i_1} w$ . Conversely, if  $v \leq s_{i_1} w$  then by the results in 1.8 applied to  $v, s_{i_1} w$  instead of  $v, w$ , we can find a sequence  $t'_2, \dots, t'_m$  with  $t'_k \in \{s_{i_k}, 1\}$  for  $k \in [2, m]$ ,  $t'_2 \dots t'_m = v$  and such that  $t'_2 \leq t'_2 t'_3 \leq \dots \leq t'_2 \dots t'_m$  and  $t'_2 \leq t'_2 s_{i_3}, \dots, t'_2 \dots t'_{m-1} \leq t'_2 \dots t'_{m-1} s_{i_m}$ . Taking  $t'_1 = 1$  we have  $t'_1 t'_2 \dots t'_m = v$ ,  $t'_1 \leq t'_1 t'_2 \leq \dots \leq t'_1 t'_2 \dots t'_m$  and  $t'_1 \leq t'_1 s_{i_2}, t'_1 t'_2 \leq t'_1 t'_2 s_{i_3}, \dots, t'_1 t'_2 \dots t'_{m-1} \leq t'_1 t'_2 \dots t'_{m-1} s_{i_m}$ . By uniqueness we have  $(t'_1, t'_2, \dots, t'_m) = (t_1, t_2, \dots, t_m)$ . Thus  $t_1 = 1$  and (a) is proved.

**1.10.** For any  $h \in G$  let  $[h] : \mathcal{B} \rightarrow \mathcal{B}$  be the map  $B \mapsto hBh^{-1}$ .

Let  $i \in I, a \in \mathbf{R}_{>0}$  and let  $v, w$  in  $W$  be such that  $v \leq w$ . We show:

(a) *If  $v \leq s_i w \leq w$ , then  $[y_i(a)] : \mathcal{B} \rightarrow \mathcal{B}$  restricts to a map  $\mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v, w}$  which is fixed point free.*

(b) *If  $s_i w \leq w$ ,  $v \not\leq s_i w$ , then  $[y_i(a)] : \mathcal{B} \rightarrow \mathcal{B}$  restricts to the identity map  $\mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v, w}$ .*

(c) *If  $w \leq s_i w$  then  $[y_i(a)] : \mathcal{B} \rightarrow \mathcal{B}$  restricts to a map  $\mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v, s_i w}$ .*

Assume first that  $s_i w \leq w$ . We can find  $\mathbf{i} = (i_1, i_2, \dots, i_m) \in \mathcal{I}_w$  such that  $i_1 = i$ . We use the notation of 1.8 relative to  $v, w, \mathbf{i}$ .

In case (a) we have  $t_1 = 1$ , see 1.9(a). In this case, left multiplication by  $y_i(a)$  restricts to a map  $\mathcal{Y}_{v, w, \mathbf{i}} \rightarrow \mathcal{Y}_{v, w, \mathbf{i}}$  given by  $g_1 g_2 \dots g_m \mapsto g'_1 g'_2 \dots g'_m$  where  $g_1 = y_1(a_1), g'_1 = y_1(a_1 + a), g'_k = g_k$  for  $k > 1$ . To prove (a), it remains to use that  $a_1 \mapsto a_1 + a$  from  $\mathbf{R}_{>0}$  to  $\mathbf{R}_{>0}$  is fixed point free.

In case (b) we have  $t_1 = s_i$ , see 1.9(a). In this case, for  $g_1 g_2 \dots g_m \in \mathcal{Y}_{v, w, \mathbf{i}}$  we have  $g_1 = \dot{s}_i$  and for some  $b \in B^+$  we have  $y_i(a) g_1 g_2 \dots g_m = y_i(a) \dot{s}_i g_2 \dots g_m = \dot{s}_i x_i(-a) g_2 \dots g_m = \dot{s}_i g_2 \dots g_m b$  (the last equality follows by an argument in [MR, 11.9]). Hence (b) holds.

Assume that we are in case (c). Let  $B \in \mathcal{B}_{\geq 0, v, w}$ ,  $B' = y_i(a) B y_i(a)^{-1}$ . We have  $\text{pos}(B^+, B') = \text{pos}(y_i(a)^{-1} B^+ y_i(a), B)$ . This equals  $s_i w$  since

$$\text{pos}(y_i(a)^{-1} B^+ y_i(a), B^+) = s_i, \text{pos}(B^+, B) = w$$

and  $s_i w \geq w$ . We have

$$\text{pos}(B^-, B') = \text{pos}(y_i(a)^{-1} B^- y_i(a), B) = \text{pos}(B^-, B) = v$$

since  $y_i(a) \in B^-$ . Thus  $B' \in \mathcal{B}_{v, s_i w}$ . By 1.5(e) we have  $B' \in \mathcal{B}_{\geq 0}$  hence  $B' \in \mathcal{B}_{\geq 0, v, s_i w}$ . Hence (c) holds.

**1.11.** Let  $i \in I, a \in \mathbf{R}_{>0}$  and let  $v, w$  in  $W$  be such that  $v \leq w$ . We show:

(a) *If  $v \leq s_i v \leq w$ , then  $[x_i(a)] : \mathcal{B} \rightarrow \mathcal{B}$  restricts to a map  $\mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v, w}$  which is fixed point free.*

(b) *If  $v \leq s_i v$ ,  $s_i v \not\leq w$ , then  $[x_i(a)] : \mathcal{B} \rightarrow \mathcal{B}$  restricts to the identity map  $\mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v, w}$ .*

(c) *If  $s_i v \leq v$ , then  $[x_i(a)] : \mathcal{B} \rightarrow \mathcal{B}$  restricts to a map  $\mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, s_i v, w}$ .*

We apply 1.10(a)-(c) to  $ww_I, vw_I$  instead of  $v, w$  (note that  $ww_I \leq vw_I$ ) and we apply the automorphism  $\phi$ . We obtain the following statements.

(1) *If  $ww_I \leq s_i vw_I \leq vw_I$ , then  $[x_i(a)] : \mathcal{B} \rightarrow \mathcal{B}$  restricts to a map*

$$\phi(\mathcal{B}_{\geq 0, ww_I, vw_I}) \rightarrow \phi(\mathcal{B}_{\geq 0, ww_I, vw_I})$$

*which is fixed point free.*

(2) *If  $s_i vw_I \leq vw_I$ ,  $ww_I \not\leq s_i vw_I$  then  $[x_i(a)] : \mathcal{B} \rightarrow \mathcal{B}$  restricts to the identity map  $\phi(\mathcal{B}_{\geq 0, ww_I, vw_I}) \rightarrow \phi(\mathcal{B}_{\geq 0, ww_I, vw_I})$ .*

(3) If  $vw_I \leq s_i vw_I$ , then  $[x_i(a)] : \mathcal{B} \rightarrow \mathcal{B}$  restricts to a map  $\phi(\mathcal{B}_{\geq 0, ww_I, vw_I}) \rightarrow \phi(\mathcal{B}_{\geq 0, ww_I, s_i vw_I})$ .

It remains to note that

(d)  $\phi(\mathcal{B}_{\geq 0, ww_I, vw_I}) = \mathcal{B}_{\geq 0, v, w}$   
for any  $v \leq w$  in  $W$  (see 1.4(a) and 1.5(c)).

**1.12.** Let  $i \in I, a \in \mathbf{R}_{>0}$ . From 1.10 we deduce:

$$(a) \quad \{B \in \mathcal{B}_{\geq 0}; y_i(a) \in B\} = \sqcup_{(v,w) \in W \times W; v \leq w, s_i w \leq w, v \not\leq s_i w} \mathcal{B}_{\geq 0, v, w}.$$

From 1.11 we deduce:

$$(b) \quad \{B \in \mathcal{B}_{\geq 0}; x_i(a) \in B\} = \sqcup_{(v,w) \in W \times W; v \leq w, v \leq s_i v, s_i v \not\leq w} \mathcal{B}_{\geq 0, v, w}.$$

**1.13.** Let  $(W \times W)_{disj} = \{(z, z') \in W \times W; \text{supp}(z) \cap \text{supp}(z') = \emptyset\}$ . For  $(z, z') \in (W \times W)_{disj}$  with  $m = |z|, m' = |z'|$ , let  $\mathcal{U}_{\geq 0, z, z'}$  be the image of the (injective) map  $\mathbf{R}_{>0}^m \times \mathbf{R}_{>0}^{m'} \rightarrow G_{\geq 0}$  given by

$$\begin{aligned} & ((a_1, a_2, \dots, a_m), (a'_1, a'_2, \dots, a'_{m'})) \\ & \mapsto y_{i_1}(a_1)y_{i_2}(a_2) \dots y_{i_m}(a_m)x_{i'_1}(a'_1)x_{i'_2}(a'_2) \dots x_{i'_{m'}}(a'_{m'}) \\ & = x_{i'_1}(a'_1)x_{i'_2}(a'_2) \dots x_{i'_{m'}}(a'_{m'})y_{i_1}(a_1)y_{i_2}(a_2) \dots y_{i_m}(a_m) \end{aligned}$$

where  $\mathbf{i} = (i_1, \dots, i_m) \in \mathcal{I}_w, \mathbf{i}' = (i'_1, \dots, i'_{m'}) \in \mathcal{I}_{w'}$ . This map is injective and its image is independent of the choice of  $\mathbf{i}, \mathbf{i}'$  (see [L94, 2.7, 2.9]). Note that  $\mathcal{U}_{\geq 0, z, z'} \subset \mathcal{U}_{\geq 0}$ . More precisely, we have (see [L94, 6.6]):

$$\mathcal{U}_{\geq 0} = \sqcup_{(z, z') \in (W \times W)_{disj}} \mathcal{U}_{\geq 0, z, z'}.$$

We state the following result.

**Theorem 1.14.** Let  $(z, z') \in (W \times W)_{disj}$  and let  $u \in \mathcal{U}_{\geq 0, z, z'}$ ,  $J = \text{supp}(z)$ ,  $J' = \text{supp}(z')$ . We have

$$\mathcal{B}_{u, \geq 0} = \cap_{i \in J} \mathcal{B}_{y_i(1), \geq 0} \cap \cap_{j \in J'} \mathcal{B}_{x_j(1), \geq 0}.$$

Using 1.6(c) we see that the following statement implies the theorem.

$$(a) \quad V_+^u = \cap_{i \in J} V_+^{y_i(1)} \cap \cap_{j \in J'} V_+^{x_j(1)}.$$

Let  $m, m', \mathbf{i}, \mathbf{i}'$  be as in 1.13 and let  $((a_1, a_2, \dots, a_m), (a'_1, a'_2, \dots, a'_{m'}))$  be corresponding to  $u$  as in 1.13. We argue by induction on  $m + m'$ . If  $m + m' = 0$  the result is obvious. Assume now that  $m + m' \geq 1$ . If  $m \geq 1$  let  $z_1 = s_{i_1} z$  and let  $u_1 = y_{i_1}(a_1)^{-1}u$ ; we have  $|z_1| = m - 1$  and  $u_1 \in \mathcal{U}_{\geq 0, z_1, z'}, u = y_{i_1}(a_1)u_1$ . If  $m' \geq 1$  let  $z'_1 = s_{i'_1} z'$  and let  $u'_1 = x_{i'_1}(a'_1)^{-1}u$ ; we have  $|z'_1| = m' - 1$  and  $u'_1 \in \mathcal{U}_{\geq 0, z, z'_1}, u = x_{i'_1}(a'_1)u'_1$ . From 1.6(a),(b), we have  $V_+^u = V_+^{y_{i_1}(1)} \cap V_+^{u_1}$  if  $m \geq 1$ ,  $V_+^u = V_+^{x_{i'_1}(1)} \cap V_+^{u'_1}$  if  $m' \geq 1$ . Since the induction hypothesis is applicable to  $V_+^{u_1}$  (if  $m \geq 1$ ) and to  $V_+^{u'_1}$  (if  $m' \geq 1$ ) we see that (a) is proved. This proves the theorem.

**1.15.** Let  $J \subset I, J' \subset I$  be such that  $J \cap J' = \emptyset$ . We define

$$\begin{aligned} Z_{J,J'} &= \{(v, w) \in W \times W; v \leq w; s_i w \leq w, v \not\leq s_i w \quad \forall i \in J; \\ &\quad v \leq s_j v, s_j v \not\leq w \quad \forall j \in J'\}. \end{aligned}$$

Combining 1.14 with 1.12(a),(b) we obtain our main result.

**Corollary 1.16.** *Let  $(z, z') \in (W \times W)_{disj}$  and let  $u \in \mathcal{U}_{\geq 0, z, z'}$ ,  $J = \text{supp}(z)$ ,  $J' = \text{supp}(z')$ . We have*

$$\mathcal{B}_{u, \geq 0} = \cup_{(v, w) \in Z_{J, J'}} \mathcal{B}_{\geq 0, v, w}.$$

Thus  $\mathcal{B}_{u, \geq 0}$  admits a canonical, explicit, cell decomposition in which each cell is part of the canonical cell decomposition of  $\mathcal{B}_{\geq 0}$ . The zero dimensional cells of  $\mathcal{B}_{u, \geq 0}$  are  $\mathcal{B}_{\geq 0, w, w}$  where  $w \in W$  is such that the set  $\{i \in I; s_i w \leq w\}$  contains  $J$  and is contained in  $I - J'$ . (For example  $w_J$  is such a  $w$  since  $J \subset I - J'$ .) In particular,  $Z_{J, J'} \neq \emptyset$ , so that  $\mathcal{B}_{u, \geq 0} \neq \emptyset$ . (This also follows from [L94, 8.11].)

**1.17.** In [L94, 2.11],  $G_{\geq 0}$  is partitioned into pieces indexed by  $W \times W$ ; by results in [L94] each piece is a cell. In [L19] the set of pieces of  $G_{\geq 0}$  is interpreted as a monoid  $G(\{1\})$ , the value of  $G$  at the semifield  $\{1\}$  with 1 element, so that pieces appear precisely as the fibres of a natural surjective map  $G_{\geq 0} \rightarrow G(\{1\}) = W \times W$  compatible with the monoid structures (the monoid structure on  $W \times W$  thus obtained is not the usual group structure). In particular, the product of two pieces of  $G_{\geq 0}$  is contained in a single piece of  $G_{\geq 0}$ . Similarly, we can view the set of cells  $\mathcal{B}_{\geq 0, v, w}$  of  $\mathcal{B}_{\geq 0}$  (see 1.7) as  $\mathcal{B}(\{1\})$  or the value of  $\mathcal{B}$  at the semifield  $\{1\}$ . From 1.10 and 1.11 we see that in the action of  $G_{\geq 0}$  on  $\mathcal{B}_{\geq 0}$ , the result of applying a piece of  $G_{\geq 0}$  to a cell of  $\mathcal{B}_{\geq 0}$  is contained in a single cell of  $\mathcal{B}_{\geq 0}$ . It follows that the action of  $G_{\geq 0}$  on  $\mathcal{B}_{\geq 0}$  induces an action of  $G(\{1\}) = W \times W$  on  $\mathcal{B}(\{1\})$ . We can identify  $\mathcal{B}(\{1\})$  with  $\{(v, w) \in W \times W; v \leq w\}$ . Then the action of  $W \times W$  becomes:

$$(s_i, 1) : (v, w) \mapsto (v, s_i * w), (1, s_i) : (v, w) \mapsto (s_i \circ v, w)$$

where for  $i \in I, v \in W, w \in W$  we define

$$\begin{aligned} s_i * w &= w \text{ if } s_i w \leq w, \quad s_i * w = s_i w \text{ if } w \leq s_i w, \\ s_i \circ v &= v \text{ if } v \leq s_i v, \quad s_i \circ v = s_i v \text{ if } s_i v \leq v. \end{aligned}$$

Now let  $u \in \mathcal{U}_{\geq 0}$  and let  $z, z', J, J'$  be associated to  $u$  as in 1.16. Let  $\mathcal{B}_u(\{1\})$  be the set of cells of  $\mathcal{B}_{u, \geq 0}$  described in 1.16; this set is in bijection with  $Z_{J, J'}$  (see 1.16) and can be viewed as a subset of  $\mathcal{B}(\{1\})$ . Let  $\mathfrak{Z}_u(\{1\})$  be the set of all  $(w, w') \in W \times W = G(\{1\})$  such that in the  $W \times W$ -action on  $\mathcal{B}(\{1\})$  (as above), we have  $(w, w')\mathcal{B}_u(\{1\}) \subset \mathcal{B}_u(\{1\})$ . Clearly,  $\mathfrak{Z}_u(\{1\})$  is a submonoid of  $W \times W = G(\{1\})$ .

Let  $\mathfrak{Z}(u)$  be the inverse image of  $\mathfrak{Z}_u(\{1\})$  under the canonical monoid homomorphism  $G_{\geq 0} \rightarrow G(\{1\})$  (as above). Note that  $\mathfrak{Z}(u)$  is a submonoid of  $G_{\geq 0}$  which is related to the centralizer of  $u$  in  $G$  (although it is not in general contained in it). Clearly the  $G_{\geq 0}$ -action of  $\mathcal{B}_{\geq 0}$  restricts to a  $\mathfrak{Z}(u)$ -action on  $\mathcal{B}_{u, \geq 0}$ .

**1.18.** Let  $\tilde{\mathcal{B}} = \{(u, B) \in \mathcal{U} \times \mathcal{B}; u \in B\}$ . Let  $\tilde{\mathcal{B}}_{\geq 0} = \{(u, B) \in \mathcal{U}_{\geq 0} \times \mathcal{B}_{\geq 0}; u \in B\}$ . From 1.16 we can deduce that  $\tilde{\mathcal{B}}_{\geq 0}$  is the disjoint union of the sets

$$(a) \quad \tilde{\mathcal{B}}_{\geq 0, z, z', v, w} = \mathcal{U}_{\geq 0, z, z'} \times \mathcal{B}_{\geq 0, v, w}$$

where  $(z, z')$  runs through  $(W \times W)_{disj}$  and  $(v, w)$  runs through  $Z_{J, J'}$  where  $J = \text{supp}(z)$ ,  $J' = \text{supp}(z')$ . The subset (a) is a cell of dimension  $|z| + |z'| + |w| - |v|$ . Note that the first projection  $\tilde{\mathcal{B}}_{\geq 0} \rightarrow \mathcal{U}_{\geq 0}$  is a trivial fibration over each  $\mathcal{U}_{\geq 0, z, z'}$ . Let  $\tilde{\mathcal{B}}(\{1\})$  be the indexing set for the set of cells of  $\tilde{\mathcal{B}}_{\geq 0}$ . This is the set of all  $(z, z', v, w) \in W^4$  such that  $J = \text{supp}(z)$ ,  $J' = \text{supp}(z')$  are disjoint and  $(v, w) \in Z_{J, J'}$ .

**1.19.** Let  $B \in \mathcal{B}_{\geq 0}$  and let  $\underline{B} = B \cap \mathcal{U}_{\geq 0}$ . This is a closed subset of  $B$  closed under multiplication. (If  $u \in \underline{B}$ ,  $u' \in \underline{B}$  then  $u, u'$  are contained in the unipotent radical of  $B$  hence  $uu'$  is also contained in that unipotent radical, so that  $uu'$  is unipotent. Since  $u \in G_{\geq 0}$ ,  $u' \in G_{\geq 0}$  we have also  $uu' \in G_{\geq 0}$ . Thus  $uu' \in \underline{B}$ .) We have  $B \in \mathcal{B}_{\geq 0, v, w}$  for well defined  $v \leq w$  in  $W$ . We set

$$\begin{aligned} \Xi_{v, w} &= \{(z, z') \in (W \times W)_{disj}; s_i w \leq w, v \not\leq s_i w \quad \forall i \in \text{supp}(z); \\ v &\leq s_j v, s_j v \not\leq w \quad \forall j \in \text{supp}(z')\}. \end{aligned}$$

We show:

$$(a) \quad \underline{B} = \sqcup_{(z, z') \in \Xi_{v, w}} \mathcal{U}_{\geq 0, z, z'}.$$

Let  $E$  be the right hand side of (a). Let  $u \in \underline{B}$ . Let  $(z, z') \in (W \times W)_{disj}$  be such that  $u \in \mathcal{U}_{\geq 0, z, z'}$ . Let  $J = \text{supp}(z)$ ,  $J' = \text{supp}(z')$ . We have  $B \in \mathcal{B}_{u, \geq 0}$  hence by 1.16 we have  $(v, w) \in Z_{J, J'}$  hence  $(z, z') \in \Xi_{v, w}$ . Thus,  $u \in E$ . We see that  $\underline{B} \subset E$ . Conversely, assume that  $u \in E$ . We can find  $(z, z') \in \Xi_{v, w}$  such that  $u \in \mathcal{U}_{\geq 0, z, z'}$ . Let  $J = \text{supp}(z)$ ,  $J' = \text{supp}(z')$ . We have  $(v, w) \in Z_{J, J'}$ . Since  $B \in \mathcal{B}_{\geq 0, v, w}$  from 1.16 we see that  $B \in \mathcal{B}_{u, \geq 0}$  so that  $u \in B$  and  $u \in \underline{B}$ . We see that  $E \subset \underline{B}$ . This proves (a).

$$\text{Let } H = \{i \in I; s_i w \leq w, v \not\leq s_i w\}, H' = \{j \in I; v \leq s_j v, s_j v \not\leq w\}.$$

If  $i \in H$  then  $v \leq w, s_i w \leq w$  and this is known to imply  $s_i v \leq w$ . Thus  $i \in H \implies i \notin H'$  so that

$$(b) \quad H \cap H' = \emptyset.$$

We have  $\Xi_{v, w} = \{(z, z') \in (W \times W)_{disj}; \text{supp}(z) \subset H, \text{supp}(z') \subset H'\}$ . Using (b) we see that

$$(c) \quad \Xi_{v, w} = \{(z, z') \in W \times W; \text{supp}(z) \subset H, \text{supp}(z') \subset H'\}.$$

Hence (a) becomes:

$$(d) \quad \underline{B} = \sqcup_{(z, z') \in W \times W; \text{supp}(z) \subset H, \text{supp}(z') \subset H'} \mathcal{U}_{\geq 0, z, z'}.$$

From (d) we see that  $\underline{B}$  has a canonical cell decomposition with cells indexed by (c). One of these cells is  $\mathcal{U}_{\geq 0, w_H, w_{H'}}$  (of dimension  $|w_H| + |w_{H'}|$ ); all other cells



have dimension strictly less than  $|w_H| + |w_{H'}|$  and are contained in the closure of  $\mathcal{U}_{\geq 0, w_H, w_{H'}}$ . We see that

(e)  $\underline{B}$  is connected of dimension  $|w_H| + |w_{H'}|$ .

If  $(v, w) = (1, w_I)$  then  $H = H' = \emptyset$  and from (d) we see that  $\underline{B} = \mathcal{U}_{\geq 0, 1, 1} = \{1\}$ . Recall that  $\mathcal{B}_{\geq 0, 1, w_I} = \mathcal{B}_{> 0}$ . We see that:

(f) If  $u \in \mathcal{U}_{\geq 0}$  and  $B \in \mathcal{B}_u$  satisfies  $B \in \mathcal{B}_{> 0}$  then  $u = 1$ .

It is likely that, more generally, for any  $B \in \mathcal{B}_{> 0}$ ,  $B \cap G_{\geq 0}$  consists of semisimple elements.

**1.20.** Let  $g \in G_{\geq 0}$  and let  $[g] : \mathcal{B} \rightarrow \mathcal{B}$  be as in 1.10. Let  $\mathcal{B}_{g, \geq 0}$  be as in 1.1. If  $v \leq w$  in  $W$  then, by 1.17, there are three possibilities:

(i)  $[g]\mathcal{B}_{\geq 0, v, w} = \mathcal{B}_{\geq 0, v', w'}$  where  $v' \leq w'$ ,  $(v', w') \neq (v, w)$ ;

(ii)  $[g]\mathcal{B}_{\geq 0, v, w} = \mathcal{B}_{\geq 0, v, w}$  and  $B \mapsto [g]B$ ,  $\mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v, w}$  has empty fixed point set;

(iii)  $[g]\mathcal{B}_{\geq 0, v, w} = \mathcal{B}_{\geq 0, v, w}$  and  $B \mapsto [g]B$ ,  $\mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v, w}$  has non-empty fixed point set (denoted by  $\mathcal{B}_{g, \geq 0, v, w}$ ).

It follows that  $\mathcal{B}_{g, \geq 0} = \sqcup_{v, w} \mathcal{B}_{g, \geq 0, v, w}$  where  $v, w$  is as in (iii). We conjecture that if  $v, w$  is as in (iii), then  $\mathcal{B}_{g, \geq 0, v, w}$  is a cell. (When  $g$  is unipotent this holds by 1.16.)

## 2. EXAMPLES

**2.1.** Assume that  $G = SL_{n+1}(\mathbf{C})$  and  $I = \{1, 2, \dots, n\}$  with  $n \geq 2$  and with  $s_1 s_2, s_2 s_3, \dots, s_{n-1} s_n$  of order 3. Let  $J = \{1, 2, \dots, n-1\}$ . Let  $-$  be the unit element of  $W$ . We have  $(w_J, -) \in (W \times W)_{disj}$  and  $Z_{J, \emptyset}$  consists of

$$(a) \quad (w_J, w_J), (w_J s_n, w_J s_n), (w_J, w_J s_n), (w_J s_n s_{n-1}, w_J s_n s_{n-1}), (w_J s_n, w_J s_n s_{n-1}), \\ \dots, (w_J s_n s_{n-1} \dots s_2 s_1, w_J s_n s_{n-1} \dots s_2 s_1), (w_J s_n s_{n-1} \dots s_2, w_J s_n s_{n-1} \dots s_2 s_1).$$

Thus if  $u \in \mathcal{U}_{\geq 0, w_J, -}$  then  $\mathcal{B}_{u, \geq 0}$  is a union of  $n+1$  cells of dimension 0 and  $n$  cells of dimension 1.

We have also  $(w_J, s_n) \in (W \times W)_{disj}$  and  $Z_{J, \{n\}}$  consists of the pairs in (a) other than the last two. Thus, if  $u \in \mathcal{U}_{\geq 0, z, s_n}$  then  $\mathcal{B}_{u, \geq 0}$  is a union of  $n$  cells of dimension 0 and  $n-1$  cells of dimension 1.

Assume now that  $n = 3$ . We have  $(s_1 s_3, -) \in (W \times W)_{disj}$ . Let  $J = \{1, 3\}$ . Then  $Z_{J, \emptyset}$  consists of

$$(b) \quad (132, 132), (13, 132), (1321, 1321), (132, 1321), (1323, 1323), (132, 1323), \\ (13231, 13231), \\ (1321, 13213), (1323, 13213), (132, 13213), (132312, 132312), (13231, 132132).$$

(We write  $i_1 i_2 \dots$  instead of  $s_{i_1} s_{i_2} \dots$ .) Thus, if  $u \in \mathcal{U}_{\geq 0, s_1 s_3, -}$  then  $\mathcal{B}_{u, \geq 0}$  is a union of 6 cells of dimension 0, 6 cells of dimension 1 and one cell of dimension 2.

We have also  $(s_1 s_3, s_2) \in (W \times W)_{disj}$  and  $Z_{J, \{2\}}$  consists of the pairs in (b) other than the last two. Thus, if  $u \in \mathcal{U}_{\geq 0, s_1 s_3, s_2}$  then  $\mathcal{B}_{u, \geq 0}$  is a union of 5 cells of dimension 0, 5 cells of dimension 1 and one cell of dimension 2.

In these examples  $\mathcal{B}_{u, \geq 0}$  is contractible; but in the last two examples,  $\mathcal{B}_{u, \geq 0}$  is not of pure dimension, unlike  $\mathcal{B}_u$ .

We have  $(s_1, -) \in (W \times W)_{disj}$ . Then  $Z_{\{1\}, \emptyset}$  consists of

$(2, 2), (21, 21), (2, 21), (23, 23), (2, 23), (212, 212), (21, 212),$   
 $(232, 232), (23, 232), (213, 213), (21, 213), (23, 213), (2, 213), (2123, 2123),$   
 $(212, 2123), (213, 2123), (21, 2123), (2321, 2321), (232, 2321), (231, 2321),$   
 $(23, 2321), (2132, 2132), (213, 2132), (232, 2132), (212, 2132), (21, 2132), (23, 2132),$   
 $(21232, 21232), (2123, 21232), (2132, 21232), (212, 21232), (21, 21232),$   
 $(23212, 23212), (2321, 23212), (2312, 23212), (232, 23212), (231, 23212),$   
 $(23, 23212), (213213, 213213), (32132, 213213), (12312, 213213), (2132, 213213).$

(We write  $i_1 i_2 \dots$  instead of  $s_{i_1} s_{i_2} \dots$ .) Thus if  $u \in \mathcal{U}_{\geq 0, s_1, -}$  then  $\mathcal{B}_{u, \geq 0}$  is a union of cells of dimension  $\leq 3$ , two of which have dimension 3.

In each of the examples above, for any cell  $\mathcal{B}_{\geq 0, v, w}$  of maximal dimension of  $\mathcal{B}_{u, \geq 0}$ , we have  $w = vw_H$  where  $H \subset I$  is such that  $u$  is conjugate in  $G$  to a regular unipotent element in the subgroup of  $G$  generated by  $\{y_i(a), x_i(a); i \in I - H, a \in \mathbf{C}\}$  and by  $T$ . This is likely to be a general phenomenon.

For such  $u$  one can show that  $|w_H| \leq \dim \mathcal{B}_u$  (complex dimension); this is compatible with  $|w_H| = \dim \mathcal{B}_{u, \geq 0}$  (real dimension).

**2.2.** In this subsection we assume that  $G, I$  are as in 2.1 and  $n = 2$ . In this case we can take  $V$  in 1.3 to be the adjoint representation of  $G$ . The canonical basis  $\beta$  of  $V$  can be denoted by  $X_{-12}, X_{-1}, X_{-2}, t_1, t_2, X_1, X_2, X_{12}$  and the action of  $x_i(a), y_i(a), i \in I, a \in \mathbf{C}$  is as follows:

$$\begin{aligned} x_i(a)X_{12} &= X_{12} \\ x_i(a)X_j &= X_j + aX_{12} \text{ if } i \neq j \in I \\ x_i(a)X_j &= X_j \text{ if } i = j \\ x_i(a)X_{-j} &= X_{-j} \text{ if } i \neq j \in I \\ x_i(a)X_{-j} &= X_{-j} + at_j + a^2X_j \text{ if } i = j \\ x_i(a)X_{-12} &= X_{-12} + aX_{-j} \text{ if } i \neq j \in I \\ x_i(a)t_j &= t_j + aX_i \text{ if } i \neq j \in I \\ x_i(a)t_j &= t_j + 2aX_i \text{ if } i = j \\ y_i(a)X_{12} &= X_{12} + aX_j \text{ if } i \neq j \in I \\ y_i(a)X_j &= X_j \text{ if } i \neq j \in I \\ y_i(a)X_j &= X_j + at_j + a^2X_{-j} \text{ if } i = j \\ y_i(a)X_{-j} &= X_{-j} + aX_{-12} \text{ if } i \neq j \in I \\ y_i(a)X_{-j} &= X_{-j} \text{ if } i = j \\ y_i(a)X_{-12} &= X_{-12} \end{aligned}$$

$$y_i(a)t_j = t_j + aX_{-i} \text{ if } i \neq j \in I$$

$$y_i(a)t_j = t_j + 2aX_{-i} \text{ if } i = j.$$

The set  $\mathcal{X}_{\geq 0}$  consists of all

$$a_{-12}X_{-12} + a_{-1}X_{-1} + a_{-2}X_{-2} + c_1t_1 + c_2t_2 + a_1X_1 + a_2X_2 + a_{12}X_{12}$$

where  $a_{-12}, a_{-1}, a_{-2}, c_1, c_2, a_1, a_2, a_{12}$  are in  $\mathbf{R}_{\geq 0}$  (not all 0) such that

$$\begin{aligned} & a_2a_{-12} = c_2a_{-1}, a_1a_{-12} = c_1a_{-2}, a_{-1}a_{12} = c_1a_2, \\ & a_{-2}a_{12} = c_2a_1, a_{12}(c_1 + c_2) = a_1a_2, a_{-12}(c_1 + c_2) = a_{-1}a_{-2}, \\ (a) \quad & c_1c_2 = a_{12}a_{-12}, c_1(c_1 + c_2) = a_1a_{-1}, c_2(c_1 + c_2) = a_2a_{-2} \end{aligned}$$

modulo the homothety action of  $\mathbf{R}_{>0}$ .

The subsets  $[v, w]$  of  $\mathcal{X}_{\geq 0}$  can be described as follows (the coefficients

$$a_{-12}, a_{-1}, a_{-2}, c_1, c_2, a_1, a_2, a_{12}$$

are required to be in  $\mathbf{R}_{>0}$  and are taken up to simultaneous multiplication by an element in  $\mathbf{R}_{>0}$ ):

$$\begin{aligned} [121, 121]: & \{a_{-12}X_{-12}\}, \\ [12, 12]: & \{a_{-1}X_{-1}\}, \\ [21, 21]: & \{a_{-2}X_{-2}\}, \\ [2, 2]: & \{a_1X_1\}, \\ [1, 1]: & \{a_2X_2\}, \\ [-, -]: & \{a_{12}X_{12}\}, \\ [21, 121]: & \{a_{-12}X_{-12} + a_{-2}X_{-2}\}, \\ [12, 121]: & \{a_{-12}X_{-12} + a_{-1}X_{-1}\}, \\ [1, 12]: & \{a_{-1}X_{-1} + a_2X_2\}, \\ [2, 21]: & \{a_{-2}X_{-2} + a_1X_1\}, \\ [-, 2]: & \{a_1X_1 + a_{12}X_{12}\}, \\ [-, 1]: & \{a_2X_2 + a_{12}X_{12}\}, \\ [2, 12]: & \{a_{-1}X_{-1} + c_1t_1 + a_1X_1; a_{-1}a_1 = c_1^2\}, \\ [1, 21]: & \{a_{-2}X_{-2} + c_2t_2 + a_2X_2; a_{-2}a_2 = c_2^2\}, \end{aligned}$$

$$\begin{aligned} [2, 121]: & \{a_{-12}X_{-12} + a_{-1}X_{-1} + a_{-2}X_{-2} + c_1t_1 + a_1X_1, a_{-1}a_1 = c_1^2, \\ & c_1a_{-12} = a_{-1}a_{-2}, a_1a_{-12} = c_1a_{-2}\}, \end{aligned}$$

$$\begin{aligned} [1, 121]: & \{a_{-12}X_{-12} + a_{-1}X_{-1} + a_{-2}X_{-2} + c_2t_2 + a_2X_2; a_{-2}a_2 = c_2^2, \\ & c_2a_{-12} = a_{-1}a_{-2}, a_2a_{-12} = c_2a_{-1}\}, \end{aligned}$$

$$\begin{aligned} [-, 12]: & \{a_{-1}X_{-1} + c_1t_1 + a_1X_1 + a_2X_2 + a_{12}X_{12}; a_{-1}a_1 = c_1^2, \\ & c_1a_{12} = a_1a_2, a_{-1}a_{12} = c_1a_2\}, \end{aligned}$$

$$: [-, 21] : \{a_{-2}X_{-2} + c_2t_2 + a_2X_2 + a_1X_1 + a_{12}X_{12}; a_{-2}a_2 = c_2^2, \\ c_2a_{12} = a_1a_2, a_{-2}a_{12} = c_2a_1\},$$

$$[-, 121] :$$

$$\{a_{-12}X_{-12} + a_{-1}X_{-1} + a_{-2}X_{-2} + c_1t_1 + c_2t_2 + a_1X_1 + a_2X_2 + a_{12}X_{12} \\ \text{such that (a) holds}\}.$$

**2.3.** We preserve the setup of 2.2.

(a) *For any  $v \leq w$  in  $W$  there is a well defined subset  $[[v, w]]$  of  $\beta$  such that  $[v, w]$  consists of all lines in  $\mathcal{X}_{\geq 0}$  which contain some vector spanned by an  $\mathbf{R}_{>0}$ -linear combination of vectors in  $[[v, w]]$ .*

We have

$$\begin{aligned} [[121, 121]] &= \{X_{-12}\}, \\ [[12, 12]] &= \{X_{-1}\}, \\ [[21, 21]] &= \{X_{-2}\}, \\ [[2, 2]] &= \{X_1\}, \\ [[1, 1]] &= \{X_2\}, \\ [[-, -]] &= \{X_{12}\}, \\ [[21, 121]] &= \{X_{-12}, X_{-2}\}, \\ [[12, 121]] &= \{X_{-12}, X_{-1}\}, \\ [[1, 12]] &= \{X_{-1}, X_2\}, \\ [[2, 21]] &= \{X_{-2}, X_1\}, \\ [[-, 2]] &= \{X_1, X_{12}\}, \\ [[-, 1]] &= \{X_2, X_{12}\}, \\ [[2, 12]] &= \{X_{-1}, t_1, X_1\}, \\ [[1, 21]] &= \{X_{-2}, t_2, X_2\}, \\ [[2, 121]] &= \{X_{-12}, X_{-1}, X_{-2}, t_1, X_1\}, \\ [[1, 121]] &= \{X_{-12}, X_{-1}, X_{-2}, t_2, X_2\}, \\ [[-, 12]] &= \{X_{-1}, t_1, X_1, X_2, X_{12}\}, \\ [[-, 21]] &= \{X_{-2}, t_2, X_2, X_1, X_{12}\}, \\ [[-, 121]] &= \{X_{-12}, X_{-1}, X_{-2}, t_1, t_2, X_1, X_2, X_{12}\}. \end{aligned}$$

We have

(b)  $[[v, w]] = [[v, 121]] \cap [[-, w]]$  for any  $v \leq w$  in  $W$ .

(c) *There is a well defined partition  $\beta = \sqcup_{z \in W} \beta_z^-$  such that*

$$[[-, w]] = \sqcup_{z \in W; z \leq w} \beta_z^-$$

*for any  $w \in W$ . There is a well defined partition  $\beta = \sqcup_{z \in W} \beta_z^+$  such that*

$$[[v, 121]] = \sqcup_{z \in W; v \leq z} \beta_z^+.$$

We have

$$\begin{aligned} \beta_-^- &= \{X_{12}\}, \beta_1^- = \{X_2\}, \beta_2^- = \{X_1\}, \beta_{12}^- = \{X_{-1}, t_1\}, \beta_{21}^- = \{X_{-2}, t_2\}, \\ \beta_{121}^- &= \{X_{-12}\}. \\ \beta_-^+ &= \{X_{12}\}, \beta_1^+ = \{t_2, X_2\}, \beta_2^+ = \{t_1, X_1\}, \beta_{21}^+ = \{X_{-2}\}, \beta_{12}^+ = \{X_{-1}\}, \\ \beta_{121}^+ &= \{X_{-12}\}. \end{aligned}$$

**2.4.** We return to the general case. Now 2.3(a),(b),(c) make sense in the general case (in (b),(c) we replace 121 by  $w_I$  and  $-$  by the unit element of  $W$ ); we expect that these statements hold in the general case. In particular  $\beta_z^- \subset \beta$  and  $\beta_z^+ \subset \beta$  are defined for  $z \in W$ .

Let  $\eta^-$  be the unique vector in  $\beta$  such that the stabilizer of  $\mathbf{C}\eta^-$  in  $G$  is  $B^-$ . We can regard  $V$  naturally as a module over the universal enveloping algebra of the Lie algebra of  $G$  hence as a module over the universal enveloping algebra  $\mathfrak{U}^+$  of the Lie algebra of  $U^+$  and as a module over the universal enveloping algebra  $\mathfrak{U}^-$  of the Lie algebra of  $U^-$ . Now  $\mathfrak{U}^+$  (resp.  $\mathfrak{U}^-$ ) has a canonical basis  $\tilde{\beta}^+$  (resp.  $\tilde{\beta}^-$ ), see [L90]) and the map  $c^+ : \tilde{b} \mapsto \tilde{b}\eta^-$  (resp.  $c^- : \tilde{b} \mapsto \tilde{b}\eta$ ) from  $\tilde{\beta}^+$  (resp.  $\tilde{\beta}^-$ ) to  $V$  has image  $\beta \cup \{0\}$ . From [L19, 10.2] we have a partition  $\tilde{\beta}^+ = \sqcup_{w \in W} \tilde{\beta}_w^+$ ; similarly we have a partition  $\tilde{\beta}^- = \sqcup_{w \in W} \tilde{\beta}_w^-$ . We expect that for  $z \in W$ ,  $\beta_z^-$  is equal to  $c^-(\tilde{\beta}_z^-)$  with 0 removed and that  $\beta_z^+$  is equal to  $c^+(\tilde{\beta}_{zw_I}^+)$  with 0 removed.

### 3. PARTIAL FLAG MANIFOLDS

**3.1.** We fix  $H \subset I$ . Let  $W^H$  be the set of all  $w \in W$  such that  $w$  has minimal length in  $wW_H$ . Let  $P_H$  be the subgroup of  $G$  generated by  $\{x_i(a); i \in I, a \in \mathbf{C}\}$ ,  $\{y_i(a); i \in H, a \in \mathbf{C}\}$  and by  $T$  (a parabolic subgroup containing  $B^+$ ). Let  $\mathcal{P}_H$  be the variety whose points are the subgroups of  $G$  conjugate to  $P_H$ . (We have  $P_\emptyset = B^+$ ,  $\mathcal{P}_\emptyset = \mathcal{B}$ .) Define  $\pi_H : \mathcal{B} \rightarrow \mathcal{P}_H$  by  $B \mapsto P$  where  $P \in \mathcal{P}_H$  contains  $B$ . As observed in [L98] to any  $P \in \mathcal{P}_H$  we can attach two Borel subgroups  $B', B''$  of  $P$  such that  $\text{pos}(B^+, B') = b \in W^H$ ,  $\text{pos}(B^-, B'') = a' \in W^H$ ; moreover  $B', B''$  (hence  $a', b$ ) are uniquely determined by  $P$ . Let  $c = \text{pos}(B', B'') \in W_H$  and let  $a = w_I a'$ . We have  $a \leq bc$  (since  $B'' \in \mathcal{B}_{a,bc}$ ) and  $ac^{-1} \leq b$  (since  $B' \in \mathcal{B}_{ac^{-1}, b}$ ). Conversely, if

(a)  $(a, b, c) \in (w_I W^H) \times W^H \times W_H$  satisfy  $a \leq bc$  or equivalently  $ac^{-1} \leq b$ , then the set of all  $P \in \mathcal{P}_H$  which give rise as above to  $a, b, c$  is non-empty; we denote this set by  $\mathcal{P}_{H,a,b,c}$ . The subsets  $\mathcal{P}_{H,a,b,c}$  form a partition of  $\mathcal{P}_H$  indexed by the set  $\mathcal{P}_H(\{1\})$  of triples  $(a, b, c)$  as in (a).

Following [L98] we set  $\mathcal{P}_{H,\geq 0} = \pi_H(\mathcal{B}_{\geq 0})$ ,  $\mathcal{P}_{H,> 0} = \pi_H(\mathcal{B}_{> 0})$ . For  $(a, b, c) \in \mathcal{P}_H(\{1\})$  we set

$$\mathcal{P}_{H,\geq 0,a,b,c} := \mathcal{P}_{H,a,b,c} \cap \mathcal{P}_{H,\geq 0}$$

so that we have

$$\mathcal{P}_{H,\geq 0} = \sqcup_{(a,b,c) \in \mathcal{P}_H(\{1\})} \mathcal{P}_{H,\geq 0,a,b,c}.$$

Let

$$\mathcal{Z} = \{(r, t) \in (w_I W^H) \times W; r \leq t\}, \mathcal{Z}' = \{(r', t') \in W \times W^H; r' \leq t'\}.$$

We have bijections

$$\alpha : \mathcal{P}_H(\{1\}) \xrightarrow{\sim} \mathcal{Z}, (a, b, c) \mapsto (a, bc), \alpha' : \mathcal{P}_H(\{1\}) \xrightarrow{\sim} \mathcal{Z}', (a, b, c) \mapsto (ac^{-1}, b).$$

We shall write  ${}_{r,t}\mathcal{P}_{H,\geq 0}$  instead of  $\mathcal{P}_{H,\geq 0,a,b,c}$  where  $(r, t) \in \mathcal{Z}$ ,  $(a, b, c) = \alpha^{-1}(r, t)$  and  ${}^{r',t'}\mathcal{P}_{H,\geq 0}$  instead of  $\mathcal{P}_{H,\geq 0,a,b,c}$  where  $(r', t') \in \mathcal{Z}'$ ,  $(a, b, c) = \alpha'^{-1}(r', t')$ . Thus we have

$\mathcal{P}_{H,\geq 0} = \sqcup_{(r,t) \in \mathcal{Z}} \mathcal{P}_{H,\geq 0} = \sqcup_{(r',t') \in \mathcal{Z}'} \mathcal{P}_{H,\geq 0}$   
and

$${}_{r,t}\mathcal{P}_{H,\geq 0} = {}_{r',t'}\mathcal{P}_{H,\geq 0} \text{ if } (r',t') = \alpha'\alpha^{-1}(r,t).$$

In [R98, p.50,51] it is shown that for  $(r,t) \in \mathcal{Z}$ ,  $\pi_H$  restricts to a bijection

$$(b) \tilde{\alpha} : \mathcal{B}_{\geq 0,r,t} \xrightarrow{\sim} {}_{r,t}\mathcal{P}_{H,\geq 0}$$

and that for  $(r',t') \in \mathcal{Z}'$ ,  $\pi_H$  restricts to a bijection

$$(c) \tilde{\alpha}' : \mathcal{B}_{\geq 0,r',t'} \xrightarrow{\sim} {}_{r',t'}\mathcal{P}_{H,\geq 0}.$$

This implies (by 1.7(a)) that  $\mathcal{P}_{H,\geq 0,a,b,c}$  is a cell for any  $(a,b,c) \in \mathcal{P}_H(\{1\})$ , so that the various  $\mathcal{P}_{H,\geq 0,a,b,c}$  form a cell decomposition of  $\mathcal{P}_{H,\geq 0}$ ; this justifies the notation  $\mathcal{P}_H(\{1\})$ .

From 1.5(d) we deduce:

$$(d) g \in G_{\geq 0}, P \in \mathcal{P}_{H,\geq 0} \implies gPg^{-1} \in \mathcal{P}_{H,\geq 0}.$$

Let  $\mathcal{P} = \sqcup_{H \subset I} \mathcal{P}_H$ ,  $\mathcal{P}_{\geq 0} = \sqcup_{H \subset I} \mathcal{P}_{H,\geq 0}$ ,  $\mathcal{P}_{> 0} = \sqcup_{H \subset I} \mathcal{P}_{H,> 0}$ .

**3.2.** Let  $H$  be as in 3.1. Let  $V_H$  be an irreducible rational  $G$ -module over  $\mathbf{C}$  such that  $\{g \in G; g\mathcal{L} = \mathcal{L}\} = P_H$  for some (necessarily unique) line  $\mathcal{L}$  in  $V$ ; let  $\eta \in \mathcal{L} - \{0\}$ . Let  $\beta$  be the basis of  $V_H$  (containing  $\eta$ ) obtained by specializing at  $v = 1$  the canonical basis [L90] of the corresponding module over the corresponding quantized enveloping algebra. Let  $\mathcal{X}_H$  be the  $G$ -orbit of  $\mathcal{L}$  in the set of lines in  $V_H$ . We have a bijection

$$(a) \mathcal{P}_H \xrightarrow{\sim} \mathcal{X}_H$$

given by  $P \mapsto L_P$  where  $L_P$  is the unique  $P$ -stable line in  $V_H$ . Let

$$(b) V_{H+} = \sum_{b \in \beta} \mathbf{R}_{\geq 0} b \subset V_H.$$

From [L98] we see that  $V_H$  above can be chosen so that:

(c) *under the bijection (a),  $\mathcal{P}_{H,\geq 0}$  corresponds to the set of lines in  $\mathcal{X}_H$  which meet  $V_{H+} - \{0\}$ .*

In the sequel we assume that  $V_H$  has been chosen so that (c) holds.

**3.3.** For  $u \in \mathcal{U}_{\geq 0}$  let  $\mathcal{P}_{H,u,\geq 0} = \{P \in \mathcal{P}_{H,\geq 0}; u \in P\}$ ,  $V_{H+}^u = \{\xi \in V_{H+}; u\xi = \xi\}$ . The proof of (a),(b),(c) below is identical to that of 1.6(a),(b),(c).

(a) *Let  $u, u', u''$  in  $\mathcal{U}_{\geq 0}$  be such that  $u = u'u''$ . We have  $V_{H+}^u = V_{H+}^{u'} \cap V_{H+}^{u''}$ .*

(b) *Let  $i \in I, a \in \mathbf{R}_{> 0}$ . We have  $V_{H+}^{x_i(a)} = V_{H+}^{x_i(1)}$ ,  $V_{H+}^{y_i(a)} = V_{H+}^{y_i(1)}$ .*

(c) *Under the bijection 3.2(c),  $\mathcal{P}_{H,u,\geq 0}$  corresponds to the set of lines in  $\mathcal{X}_H$  which meet  $V_{H+}^u - \{0\}$ .*

**3.4.** For any  $h \in G$  let  $[h]_H : \mathcal{P}_H \rightarrow \mathcal{P}_H$  be the map  $P \mapsto hPh^{-1}$ .

Let  $i \in I, a \in \mathbf{R}_{> 0}$ . Let  $(r,t) \in \mathcal{Z}$ . Now (a),(b),(c) below follow immediately from 1.10(a),(b),(c) using 3.1(b).

(a) *If  $r \leq s_i t \leq t$ , then  $[y_i(a)]_H : \mathcal{P}_H \rightarrow \mathcal{P}_H$  restricts to a map  ${}_{r,t}\mathcal{P}_{H,\geq 0} \rightarrow {}_{r,t}\mathcal{P}_{H,\geq 0}$  which is fixed point free.*

(b) *If  $s_i t \leq t$ ,  $r \not\leq s_i t$ , then  $[y_i(a)]_H : \mathcal{P}_H \rightarrow \mathcal{P}_H$  restricts to the identity map  ${}_{r,t}\mathcal{P}_{H,\geq 0} \rightarrow {}_{r,t}\mathcal{P}_{H,\geq 0}$ .*

(c) *If  $t \leq s_i t$  then  $[y_i(a)]_H : \mathcal{P}_H \rightarrow \mathcal{P}_H$  restricts to a map  ${}_{r,t}\mathcal{P}_{H,\geq 0} \rightarrow {}_{r,s_i t}\mathcal{P}_{H,\geq 0}$ ; note that  $(r, s_i t) \in \mathcal{Z}$ .*

**3.5.** Let  $i \in I, a \in \mathbf{R}_{>0}$ . Let  $(r', t') \in \mathcal{Z}'$ . Now (a),(b),(c) below follow immediately from 1.11(a),(b),(c) using 3.1(c).

(a) If  $r' \leq s_i r' \leq t'$ , then  $[x_i(a)]_H : \mathcal{P}_H \rightarrow \mathcal{P}_H$  restricts to a map  ${}^{r',t'}\mathcal{P}_{H,\geq 0} \rightarrow {}^{r',t'}\mathcal{P}_{H,\geq 0}$  which is fixed point free.

(b) If  $r' \leq s_i r', s_i r' \not\leq t'$ , then  $[x_i(a)]_H : \mathcal{P}_H \rightarrow \mathcal{P}_H$  restricts to the identity map  ${}^{r',t'}\mathcal{P}_{H,\geq 0} \rightarrow {}^{r',t'}\mathcal{P}_{H,\geq 0}$ .

(c) If  $s_i r' \leq r'$  then  $[x_i(a)]_H : \mathcal{P}_H \rightarrow \mathcal{P}_H$  restricts to a map  ${}^{r',t'}\mathcal{P}_{H,\geq 0} \rightarrow {}^{s_i r',t'}\mathcal{P}_{H,\geq 0}$ ; note that  $(s_i r', t') \in \mathcal{Z}'$ .

**3.6.** Let  $i \in I, a \in \mathbf{R}_{>0}$ . From 3.4 we deduce:

$$(a) \quad \{P \in \mathcal{P}_{H,\geq 0}; y_i(a) \in P\} = \sqcup_{(r,t) \in \mathcal{Z}; s_i t \leq t, r \not\leq s_i t} ({}_{r,t}\mathcal{P}_{H,\geq 0}).$$

From 3.5 we deduce:

$$(b) \quad \{P \in \mathcal{P}_{H,\geq 0}; x_i(a) \in P\} = \sqcup_{(r',t') \in \mathcal{Z}'; r' \leq s_i r', s_i r' \not\leq t'} {}^{r',t'}\mathcal{P}_{H,\geq 0}.$$

The proof of the following result is entirely similar to that of 1.14, using 3.3(a),(b) instead of 1.6(a),(b).

**Theorem 3.7.** Let  $(z, z') \in (W \times W)_{disj}$  and let  $u \in \mathcal{U}_{\geq 0, z, z'}, J = \text{supp}(z), J' = \text{supp}(z')$ . We have

$$\mathcal{P}_{H,u,\geq 0} = \cap_{i \in J} \mathcal{P}_{H,y_i(1),\geq 0} \cap \cap_{j \in J'} \mathcal{P}_{H,x_j(1),\geq 0}.$$

**3.8.** Let  $J \subset I, J' \subset I$  be such that  $J \cap J' = \emptyset$ . We set

$$\begin{aligned} Z_{H;J,J'} &= \{((r, t), (r', t')) \in \mathcal{Z} \times \mathcal{Z}'; \\ (r', t') &= \alpha' \alpha^{-1}(r, t), s_i t \leq t, r \not\leq s_i t \quad \forall i \in J; r' \leq s_j r', s_j r' \not\leq t' \quad \forall j \in J'\}. \end{aligned}$$

Combining 3.7 with 3.6(a),(b) we obtain the following result.

**Corollary 3.9.** Let  $(z, z') \in (W \times W)_{disj}$  and let  $u \in \mathcal{U}_{\geq 0, z, z'}, J = \text{supp}(z), J' = \text{supp}(z')$ . We have

$$\mathcal{P}_{H,u,\geq 0} = \cup_{((r,t),(r',t')) \in Z_{H;J,J'}} ({}_{r,t}\mathcal{P}_{H,\geq 0}).$$

Thus  $\mathcal{P}_{H,u,\geq 0}$  admits a canonical, explicit, cell decomposition in which each cell is part of the canonical cell decomposition of  $\mathcal{P}_{H,\geq 0}$ . The zero dimensional cells of  $\mathcal{P}_{H,u,\geq 0}$  are indexed by

$$\{((r, r), (r', r')); r' \in W^H, r = r' w_H, s_i r \leq r \quad \forall i \in J, r' \leq s_j r' \quad \forall j \in J'\}.$$

The last set contains for example  $((r, r), (r', r'))$  where

$$r' = w_J w_{J \cap H}, r = w_J w_{J \cap H} w_H.$$

**3.10.** From 3.4 and 3.5 we see that in the action of  $G_{\geq 0}$  on  $\mathcal{P}_{H,\geq 0}$ , the result of applying a piece of  $G_{\geq 0}$  to a cell of  $\mathcal{P}_{H,\geq 0}$  is contained in a single cell of  $\mathcal{P}_{H,\geq 0}$ . It follows that the action of  $G_{\geq 0}$  on  $\mathcal{P}_{H,\geq 0}$  induces an action of  $G(\{1\}) = W \times W$  on  $\mathcal{P}_H(\{1\})$ . We can identify  $\mathcal{P}_H(\{1\})$  with

$$\{((r, t), (r', t')) \in \mathcal{Z} \times \mathcal{Z}'; (r', t') = \alpha' \alpha^{-1}(r, t)\}.$$

Then the action of  $W \times W$  becomes:

$$(s_i, 1) : ((r, t), (r', t')) \mapsto ((r, s_i * t), \alpha' \alpha^{-1}(r, s_i * t)),$$

$$(1, s_i) : ((r, t), (r', t')) \mapsto (\alpha \alpha'^{-1}(s_i \circ r', t'), (s_i \circ r', t')),$$

(notation of 1.17).

**3.11.** Let  $P \in \mathcal{P}_{H,\geq 0}$  and let  $\underline{P} = P \cap \mathcal{U}_{\geq 0}$ . This is a closed subset of  $P$ . We have  $P \in {}_{r,t}\mathcal{P}_{H,\geq 0} = {}^{r',t'}\mathcal{P}_{H,\geq 0}$  for a well defined  $((r, t), (r', t')) \in \mathcal{Z} \times \mathcal{Z}'$  such that  $(r', t') = \alpha' \alpha^{-1}(r, t)$ . We set

$$\begin{aligned} \Xi_{H,(r,t),(r',t')} &= \{(z, z') \in (W \times W)_{disj}; s_i t \leq t, r \not\leq s_i t \quad \forall i \in \text{supp}(z); \\ &r' \leq s_j r', s_j r' \not\leq t' \quad \forall j \in \text{supp}(z')\}. \end{aligned}$$

We show:

$$(a) \quad \underline{P} = \sqcup_{(z,z') \in \Xi_{H,(r,t),(r',t')}} \mathcal{U}_{\geq 0,z,z'}.$$

Let  $E$  be the right hand side of (a). Let  $u \in \underline{P}$ . Let  $(z, z') \in (W \times W)_{disj}$  be such that  $u \in \mathcal{U}_{\geq 0,z,z'}$ . Let  $J = \text{supp}(z)$ ,  $J' = \text{supp}(z')$ . We have  $P \in \mathcal{P}_{H,u,\geq 0}$  hence by 3.9 we have  $((r, t), (r', t')) \in Z_{H;J,J'}$  hence  $(z, z') \in \Xi_{H,(r,t),(r',t')}$ . Thus,  $u \in E$ . We see that  $\underline{P} \subset E$ . Conversely, assume that  $u \in E$ . We can find  $(z, z') \in \Xi_{H,(r,t),(r',t')}$  such that  $u \in \mathcal{U}_{\geq 0,z,z'}$ . Let  $J = \text{supp}(z)$ ,  $J' = \text{supp}(z')$ . We have  $((r, t), (r', t')) \in Z_{H;J,J'}$ . Since  $P \in {}_{r,t}\mathcal{P}_{H,\geq 0}$ , from 3.9 we see that  $P \in \mathcal{P}_{H,u,\geq 0}$  so that  $u \in P$  and  $u \in \underline{P}$ . We see that  $E \subset \underline{P}$ . This proves (a).

From (a) we see that  $\underline{P}$  has a canonical cell decomposition with cells indexed by  $\Xi_{H,(r,t),(r',t')}$ .

#### 4. A CONJECTURE AND ITS CONSEQUENCES

**4.1.** An  $\mathbf{R}_{>0}$ -positive structure on a set  $\mathfrak{X}$  is a finite collection  $f_e : \mathbf{R}_{>0}^m \xrightarrow{\sim} \mathfrak{X}$  of bijections ( $e \in E$ ) with  $m \geq 0$  fixed, such that for any  $e, e'$  in  $E$ ,  $f_e^{-1} f_{e'} : \mathbf{R}_{>0}^m \rightarrow \mathbf{R}_{>0}^m$  is admissible in the sense of [L19, 1.2]. If  $(\mathfrak{X}; f_e : \mathbf{R}_{>0}^m \rightarrow \mathfrak{X}, e \in E)$ ,  $(\tilde{\mathfrak{X}}; \tilde{f}_{\tilde{e}} : \mathbf{R}_{>0}^{\tilde{m}} \rightarrow \tilde{\mathfrak{X}}, \tilde{e} \in \tilde{E})$  are two sets with  $\mathbf{R}_{>0}$ -positive structure, a map  $\xi : \mathfrak{X} \rightarrow \tilde{\mathfrak{X}}$  is said to be a morphism if for some (or equivalently any)  $e \in E, \tilde{e} \in \tilde{E}$ , the map  $\tilde{f}_{\tilde{e}}^{-1} \xi f_e : \mathbf{R}_{>0}^m \rightarrow \mathbf{R}_{>0}^{\tilde{m}}$  is admissible in the sense of [L19, 1.2]. We say that  $\xi$  is an isomorphism if it is a bijective morphism and  $\xi^{-1}$  is a morphism.

If  $\mathfrak{X}, \mathfrak{X}'$  have positive  $\mathbf{R}_{>0}$ -structures, then  $\mathfrak{X} \times \mathfrak{X}'$  has a natural  $\mathbf{R}_{>0}$ -positive structure.



**4.2.** Let  $T_{>0} = T \cap G_{\geq 0}$ . For any basis  $\chi_* = (\chi_1, \chi_2, \dots, \chi_n)$  of  $\text{Hom}(\mathbf{C}^*, T)$  we have a bijection  $\mathbf{R}_{>0}^n \xrightarrow{\sim} T_{>0}$  given by  $(a_1, \dots, a_n) \mapsto \chi_1(a_1) \dots \chi_n(a_n)$ . These bijections (for various  $\chi_*$ ) define an  $\mathbf{R}_{>0}$ -positive structure on  $T_{>0}$ .

Let  $v \leq w$  in  $W$ . From [R08] we see that the bijections  $\tau_{\mathbf{i}} : \mathbf{R}_{>0}^{|w|-|v|} \xrightarrow{\sim} \mathcal{B}_{\geq 0, v, w}$  (see 1.8(c)) with  $\mathbf{i} \in \mathcal{I}_w$  form an  $\mathbf{R}_{>0}$ -positive structure on  $\mathcal{B}_{\geq 0, v, w}$ .

From 1.10 we see that for  $i \in I$ ,

(a) *the map  $\mathbf{R}_{>0} \times \mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v, s_i * w}$  given by  $(a, B) \mapsto y_i(a) B y_i(a)^{-1}$  is a well defined morphism of sets with  $\mathbf{R}_{>0}$ -positive structure.*

From the definitions, for any  $\chi \in \text{Hom}(\mathbf{C}^*, T)$ ,

(b) *the map  $\mathbf{R}_{>0} \times \mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v, w}$  given by  $(a, B) \mapsto \chi(a) B \chi(a)^{-1}$  is a well defined morphism of sets with  $\mathbf{R}_{>0}$ -positive structure.*

**4.3.** Replacing in 4.2  $v, w$  by  $w_I w, w_I v$ , we deduce that the bijections

$$\tau_{\mathbf{i}'} : \mathbf{R}_{>0}^{|w|-|v|} = \mathbf{R}_{>0}^{|w_I v|-|w_I w|} \xrightarrow{\sim} \mathcal{B}_{\geq 0, w_I w, w_I v}$$

with  $\mathbf{i}' \in \mathcal{I}_{w_I v}$  form an  $\mathbf{R}_{>0}$ -positive structure on  $\mathcal{B}_{\geq 0, w_I w, w_I v}$  and that for  $i \in I$ , the map

$$\mathbf{R}_{>0} \times \mathcal{B}_{\geq 0, w_I w, w_I v} \rightarrow \mathcal{B}_{\geq 0, w_I w, s_i * (w_I v)}$$

given by  $(a, B) \mapsto y_i(a) B y_i(a)^{-1}$  is a well defined morphism of sets with  $\mathbf{R}_{>0}$ -positive structure. We state:

**Conjecture 4.4.** *The bijection  $\mathcal{B}_{\geq 0, w w_I, v w_I} \xrightarrow{\sim} \mathcal{B}_{\geq 0, v, w}$  defined by  $\phi$  (see 1.11(d)) is an isomorphism of  $\mathbf{R}_{>0}$ -positive structures.*

When  $v = 1, w = w_I$  this can be deduced from [L97, §3].

*In the remainder of this section we assume that this conjecture holds.*

**4.5.** Let  $v, w$  be as in 4.2. Using 4.4 we can reformulate the last statement in 4.3 as follows. For  $i \in I$ ,

(a) *the map  $\mathbf{R}_{>0} \times \mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, s_i \circ v, w}$  given by  $(a, B) \mapsto x_i(a) B x_i(a)^{-1}$  is a well defined morphism of sets with  $\mathbf{R}_{>0}$ -positive structure.*

**4.6.** We consider a sequence of admissible maps (see [L19, 1.2]):

$$\begin{aligned} \Phi_1 : \mathbf{R}_{>0} \times \mathbf{R}_{>0}^{n_0} &\rightarrow \mathbf{R}_{>0}^{n_1}, \Phi_2 : \mathbf{R}_{>0} \times \mathbf{R}_{>0}^{n_1} \rightarrow \mathbf{R}_{>0}^{n_2}, \\ \dots, \Phi_\sigma : \mathbf{R}_{>0} \times \mathbf{R}_{>0}^{n_{\sigma-1}} &\rightarrow \mathbf{R}_{>0}^{n_\sigma} \end{aligned}$$

where  $n_0, n_1, \dots, n_\sigma$  are in  $\mathbf{N}$ . We define  $\Phi : \mathbf{R}_{>0}^\sigma \times \mathbf{R}_{>0}^{n_0} \rightarrow \mathbf{R}_{>0}^{n_\sigma}$  by

$$\Phi((a_1, a_2, \dots, a_\sigma), b) = \dots \Phi_3(a_3, \Phi_2(a_2, \Phi_1(a_1, b))) \dots$$

where  $b \in \mathbf{R}_{>0}^{n_0}$ . Clearly,  $\Phi$  is admissible.

**4.7.** Consider the inverse image  $G_{r,-s}$  of  $(r, s) \in W \times W$  under the map  $G_{\geq 0} \rightarrow W \times W$  in 1.17. Now  $G_{r,-s}$  has a natural  $\mathbf{R}_{>0}$ -positive structure (see [L19]). Let  $(i_1, i_2, \dots, i_m) \in \mathcal{I}_r, (j_1, j_2, \dots, j_l) \in \mathcal{I}_s$ . Let  $v \leq w$  be elements of  $W$ . Define  $v' \leq w'$  by  $v' = s_{i_1} \circ (\dots s_{i_{m-1}} \circ (s_{i_m} \circ v)) \dots$ ,  $w' = s_{j_1} * (\dots s_{j_{l-1}} * (s_{j_l} \circ w)) \dots$  (notation of 1.17). The  $G_{\geq 0}$ -action on  $\mathcal{B}_{\geq 0}$  restricts to a map  $G_{r,-s} \times \mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v', w'}$ .

(a) *This is a morphism of sets with  $\mathbf{R}_{>0}$ -positive structure.*

Indeed, our map can be identified with a  $\Phi$  in 4.6 where each of  $\Phi_1, \dots, \Phi_\sigma$  is a map as in 4.2(a),(b) or 4.5(a) hence is admissible.

**4.8.** We now fix a semifield  $K$ . Let  $(\mathfrak{X}; f_e : \mathbf{R}_{>0}^m \rightarrow \mathfrak{X}, e \in E)$  be a set with an  $\mathbf{R}_{>0}$ -positive structure. For any  $e, e'$  in  $E$  the admissible bijection  $f_e^{-1} f_{e'} : \mathbf{R}_{>0}^m \xrightarrow{\sim} \mathbf{R}_{>0}^m$  induces a bijection  $\underline{f}_{e, e'} : K^m \xrightarrow{\sim} K^m$ . (This is obtained by replacing the indeterminates which appear in the formula for  $f_e^{-1} f_{e'}$  by elements of  $K$  instead of elements of  $\mathbf{R}_{>0}$ .) There is a well defined set  $\mathfrak{X}(K)$  with bijections  $\underline{f}_e : K^m \xrightarrow{\sim} \mathfrak{X}(K)$  such that  $\underline{f}_{e, e'} = \underline{f}_e^{-1} \underline{f}_{e'}$  for any  $e, e'$  in  $E$ . If  $\xi : \mathfrak{X} \rightarrow \mathfrak{X}'$  is a morphism of sets with  $\mathbf{R}_{>0}$ -positive structure then  $\xi$  induces a map of sets  $\xi(K) : \mathfrak{X}(K) \rightarrow \mathfrak{X}'(K)$ .

**4.9.** In the setup of 4.7, the sets  $G_{r,-s}(K), \mathcal{B}_{\geq 0, v, w}(K), \mathcal{B}_{\geq 0, v', w'}(K)$  are defined as in 4.8 and by 4.8(a), the map  $G_{r,-s} \times \mathcal{B}_{\geq 0, v, w} \rightarrow \mathcal{B}_{\geq 0, v', w'}$  induces a map

(a)  $G_{r,-s}(K) \times \mathcal{B}_{\geq 0, v, w}(K) \rightarrow \mathcal{B}_{\geq 0, v', w'}(K)$ .

We set  $G(K) = \sqcup_{(r,s) \in W \times W} G_{r,-s}(K)$ ,  $\mathcal{B}(K) = \sqcup_{(v,w) \in W \times W; v \leq w} \mathcal{B}_{\geq 0, v, w}(K)$ . The maps (a) define a map  $G(K) \times \mathcal{B}(K) \rightarrow \mathcal{B}(K)$ . This is an action of  $G(K)$  (with the monoid structure induced from that of  $G_{\geq 0}$ ) on the set  $\mathcal{B}(K)$ .

**4.10.** Let  $H \subset I$ . For any  $(a, b, c) \in \mathcal{P}_H(\{1\})$ , the cell  $\mathcal{P}_{H, \geq 0, a, b, c}$  in  $\mathcal{P}_{H, \geq 0}$  has a  $\mathbf{R}_{>0}$ -positive structure via a bijection as in 3.1(a) or as in 3.1(b) (these two bijections define the same  $\mathbf{R}_{>0}$ -positive structure, as we see easily from the definitions). Thus the set  $\mathcal{P}_{H, \geq 0, a, b, c}(K)$  is defined. We set

$$\mathcal{P}_H(K) = \sqcup_{(a,b,c) \in \mathcal{P}_H(\{1\})} \mathcal{P}_{H, \geq 0, a, b, c}(K).$$

An argument similar to that in 4.9 shows that the  $G_{\geq 0}$ -action on  $\mathcal{P}_{H, \geq 0}$  (see 3.1(d)) induces an action of the monoid  $G(K)$  on  $\mathcal{P}_H(K)$ .

**4.11.** Let  $u \in \mathcal{U}_{\geq 0}$ . Let  $z, z', J, J'$  be as in 1.16. We set

$$\mathcal{B}_u(K) = \cup_{(v,w) \in Z_{J, J'}} \mathcal{B}_{\geq 0, v, w}(K),$$

$$\mathfrak{Z}(u)(K) = \cup_{(r,s) \in \mathfrak{Z}_u(\{1\})} G_{r,-s}(K).$$

(notation of 1.15, 1.17). Then the action of  $G(K)$  on  $\mathcal{B}(K)$  restricts to an action of  $\mathfrak{Z}(u)(K)$  (with the monoid structure induced from that of  $G(K)$ ) on the set  $\mathcal{B}_u(K)$ .

5. THE MAP  $g \mapsto P_g$  FROM  $G_{\geq 0}$  TO  $\mathcal{P}_{\geq 0}$ 

**5.1.** In this section we give a new, simpler, definition of the map  $g \mapsto B_g$  from  $G_{>0}$  to  $\mathcal{B}_{>0}$  in [L94, 8.9(c)] and we extend it to a map  $g \mapsto P_g$  from  $G_{\geq 0}$  to  $\mathcal{P}_{\geq 0}$ .

**5.2.** For a closed subgroup  $G'$  of  $G$  we denote by  $\mathfrak{L}G'$  the Lie algebra of  $G'$ . Let  $\mathfrak{g} = \mathfrak{L}G$ . Let  $g \in G_{\geq 0}$ . We associate to  $g$  a parabolic subgroup  $P = P_g$  of  $G$  containing  $g$  as follows. By [L19, 9.1(a)], we have  $\mathfrak{g} = \bigoplus_{a \in \mathbf{R}_{>0}} \mathfrak{g}_a$  where  $\mathfrak{g}_a$  is the generalized  $a$ -eigenspace of  $\text{Ad}(g) : \mathfrak{g} \rightarrow \mathfrak{g}$ . It follows that we have  $\mathfrak{g} = \mathfrak{g}_{<1} \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_{>1}$  where  $\mathfrak{g}_{<1} = \bigoplus_{a:0 < a < 1} \mathfrak{g}_a$ ,  $\mathfrak{g}_{>1} = \bigoplus_{a:a > 1} \mathfrak{g}_a$  and  $\mathfrak{g}_{\leq 1} = \mathfrak{g}_{<1} \oplus \mathfrak{g}_1$ ,  $\mathfrak{g}_{\geq 1} = \mathfrak{g}_1 \oplus \mathfrak{g}_{>1}$  are opposed parabolic subalgebras of  $\mathfrak{g}$  with common Levi subalgebra  $\mathfrak{g}_1$ . Let  $P = P_g$  be the parabolic subgroup of  $G$  with  $\mathfrak{L}P = \mathfrak{g}_{\geq 1}$ . Let  $L = L_g$  be the Levi subgroup of  $P$  with  $\mathfrak{L}L = \mathfrak{g}_1$ . Note that  $L$  is the centralizer in  $G$  of the semisimple part  $g_s$  of  $g$ . In particular we have  $g \in L$ ; more precisely,  $g$  is a central element of  $L$  times a unipotent element of  $L$ . For example, if  $g \in G_{\geq 0}$  is unipotent then  $P_g = G$ .

**5.3.** We now assume that  $g \in G_{>0}$ . We show:

(a)  $B_g = P_g$ .

Let  $U_{>0}^+ \subset U^+$ ,  $U_{>0}^- \subset U^-$  be as in [L94, 2.12]. Let  $T_{>0} \subset T$  be as in 4.7. By [L94, 8.10] we can find  $u \in U_{>0}^+$ ,  $u' \in U_{>0}^-$ ,  $t \in T_{>0}$  such that  $g = u'tuu'^{-1}$  and all eigenvalues of  $\text{Ad}(t)$  on  $\mathfrak{L}U^+$  are  $> 1$ . Then all eigenvalues of  $\text{Ad}(t)$  on  $\mathfrak{L}B^+$  are  $\geq 1$ . Now  $tu$  is  $U^+$ -conjugate to  $t$  hence all eigenvalues of  $\text{Ad}(tu)$  on  $\mathfrak{L}B^+$  are  $\geq 1$ . It follows that all eigenvalues of  $\text{Ad}(g)$  on  $\mathfrak{L}(u'B^+u'^{-1})$  are  $\geq 1$ , so that  $\mathfrak{L}(u'B^+u'^{-1}) \subset \mathfrak{L}P_g$ . Since  $t$  (and  $tu$ ) is regular semisimple,  $L_g$  is a maximal torus of  $G$  so that  $P_g \in \mathcal{B}$ ; this implies that  $\mathfrak{L}(u'B^+u'^{-1}) = \mathfrak{L}P_g$ . From the definition of  $\mathcal{B}_{>0}$  we have  $u'B^+u'^{-1} \in \mathcal{B}_{>0}$ . Since  $g \in u'B^+u'^{-1}$ , we have  $u'B^+u'^{-1} = B_g$ . This proves (a).

**5.4.** Let  $g \in G_{\geq 0}$ . We show:

(a)  $P_g \in \mathcal{P}_{\geq 0}$ .

By [L94, 4.4] we can find a sequence  $g_1, g_2, g_3, \dots$  in  $G_{>0}$  such that  $g = \lim_{n \rightarrow \infty} g_n$  in  $G$ . Since  $\mathcal{B}$  is compact, some subsequence of the sequence of Borel subgroups  $P_{g_n}$  converges in  $\mathcal{B}$  to a Borel subgroup  $B \in \mathcal{B}$ . We can assume that  $\lim_{n \rightarrow \infty} P_{g_n} = B$  in  $\mathcal{B}$ . Since  $P_{g_n} \in \mathcal{B}_{>0} \subset \mathcal{B}_{\geq 0}$  and  $\mathcal{B}_{\geq 0}$  is closed in  $\mathcal{B}$ , we have  $B \in \mathcal{B}_{\geq 0}$ . Let  $\mathfrak{p}_n = \mathfrak{L}P_{g_n}$ ,  $n = 1, 2, \dots$ ,  $\mathfrak{p} = \mathfrak{L}P_g$ , and let  $\mathfrak{b} = \mathfrak{L}B$ . We show:

(b)  $\mathfrak{b} \subset \mathfrak{p}$ .

We have  $\lim_{n \rightarrow \infty} \mathfrak{p}_n = \mathfrak{b}$  in the Grassmannian of  $\dim \mathfrak{b}$ -dimensional subspaces of  $\mathfrak{g}$ . Since  $\mathfrak{p}_n$  is stable under  $\text{Ad}(g_n)$  for  $n = 1, 2, \dots$ , we see that  $\mathfrak{b}$  is stable under  $\text{Ad}(g)$ . Moreover since all eigenvalues of  $\text{Ad}(g_n) : \mathfrak{p}_n \rightarrow \mathfrak{p}_n$  are  $\geq 1$ , we see that all eigenvalues of  $\text{Ad}(g) : \mathfrak{b} \rightarrow \mathfrak{b}$  are  $\geq 1$ . Hence (b) holds.

From (b) we deduce that  $B \subset P_g$ . Since  $B \in \mathcal{B}_{\geq 0}$  we see that  $P_g \in \mathcal{P}_{\geq 0}$ ; this proves (a) and completes the verification of the statements in 5.1.

**5.5.** Let  $B \in \mathcal{B}_{>0}$ . We show:

(a)  $B \cap G_{>0}$  is non-empty.

We have  $B = uB^+u^{-1}$  for some  $u \in U_{>0}^-$ . Let  $\tilde{u} \in U_{>0}^+$ . By [L94, 7.2] we can

find  $t \in T_{>0}$  such that  $u\tilde{u}tu^{-1} \in G_{>0}$ . We have  $u\tilde{u}tu^{-1} \in uB^+u^{-1} = B$ . Thus  $u\tilde{u}tu^{-1} \in B \cap G_{>0}$  and (a) follows.

We conjecture that  $B \cap G_{>0}$  is homeomorphic to  $\mathbf{R}_{>0}^{\dim B}$ . Consider for example the case where  $G = SL_2(\mathbf{C})$ . In this case there is a unique  $z \in \mathbf{R}_{>0}$  such that  $B$  equals  $\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix}; a, b, c, d \text{ in } \mathbf{C}, ad = bc + 1, c + dz = z(a + bz) \right\}$ . Then  $B \cap G_{>0}$  can be identified with the set  $\{(a, b, c, d) \in \mathbf{R}_{>0}^4; ad = bc + 1, c + dz = z(a + bz)\}$  or (via the substitution  $d = (bc + 1)/a$ ), with the set  $X_z = \{(a, b, c) \in \mathbf{R}_{>0}^3; c + z\frac{bc+1}{a} = z(a + bz)\} = \{(a, b, c) \in \mathbf{R}_{>0}^3; (az - c)(bz + a) = z\}$ . Setting  $\epsilon = a - cz^{-1}$  we can identify  $X_z$  with  $X'_z = \{(b, c, \epsilon) \in \mathbf{R}_{>0}^3; \epsilon^2 + \epsilon(bz + cz^{-1}) - 1 = 0\}$ . The map  $X'_z \rightarrow \mathbf{R}_{>0}^2, (b, c, \epsilon) \mapsto (b, c)$  is a homeomorphism: for any  $(b, c) \in \mathbf{R}_{>0}^2$  there is a unique  $\epsilon \in \mathbf{R}_{>0}$  such that  $\epsilon^2 + \epsilon(bz + cz^{-1}) - 1 = 0$ . This proves the conjecture in our case.

**5.6.** We show:

(a) *If  $g \in G_{>0}$  then  $P_g \in \mathcal{B}$ ; the image of the map  $g \mapsto P_g$  from  $G_{>0}$  to  $\mathcal{B}$  is exactly  $\mathcal{B}_{>0}$ .*

The fact that  $P_g \in \mathcal{B}_{>0}$  for  $g \in G_{>0}$  follows from 5.3(a). Let  $B \in \mathcal{B}_{>0}$ . Let  $g \in B \cap G_{>0}$  (see 5.5(a)). From [L94, 8.9(a)] we see that  $B = B_g$  hence by 5.3(a) we have  $B = P_g$ . This proves (a).

Now (a) provides a new definition of  $\mathcal{B}_{>0}$ ; it is the image of the map  $g \mapsto P_g$  from  $G_{>0}$  to  $\mathcal{B}$ .

Let  $G_{reg}$  be the set of regular elements in  $G$  and let  $G_{reg, \geq 0} = G_{reg} \cap G_{\geq 0}$ . We conjecture that the map  $g \mapsto P_g$  from  $G_{\geq 0}$  to  $\mathcal{P}_{\geq 0}$  is surjective and that, moreover, its restriction  $G_{reg, \geq 0} \rightarrow \mathcal{P}_{\geq 0}$  is surjective.

## REFERENCES

- [KL] D. Kazhdan and G. Lusztig, *Representations of Coxeter groups and Hecke algebras*, Invent.Math. **53** (1979), 165-184.
- [L90] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), 447-498.
- [L94] G. Lusztig, *Total positivity in reductive groups*, Lie theory and geometry, Progr.in Math. 123, Birkhäuser Boston, 1994, pp. 531-568.
- [L97] G. Lusztig, *Total positivity and canonical bases*, Algebraic groups and Lie groups, ed. G.I.Lehrer, Cambridge U.Press, 1997, pp. 281-295.
- [L98] G. Lusztig, *Total positivity in partial flag manifolds*, Repres.Th. **2** (1998), 70-78.
- [L19] G. Lusztig, *Total positivity in reductive groups, II*, arxiv:1904.07198 (to appear Bull. Inst. Math. Acad. Sinica).
- [MR] R. J. Marsh and K. Rietsch, *Parametrizations of flag varieties*, Repres.Th. **8** (2004), 212-242.
- [R98] K. Rietsch, *Total positivity and real flag varieties*, MIT Ph.D. thesis (1998).
- [R99] K. Rietsch, *An algebraic cell decomposition of the nonnegative part of a flag manifold*, J.Algebra **213** (1999), 144-154.
- [R08] K. Rietsch, *A mirror symmetric construction of  $qH(T)^*(G/P)((q))$* , Adv.Math. **217** (2008), 2401-2442.