ASSOCIATED VARIETIES AND MINIMAL HIGHEST WEIGHT MODULES

ZHANQIANG BAI, JIA-JUN MA, WEI XIAO AND XUN XIE

ABSTRACT. For a complex simple Lie algebra \mathfrak{g} , it is well-known that the associated variety of a highest weight module $L(\lambda)$ of g is the union of some orbital varieties associated with the nilpotent orbit $\mathcal{O}_{\operatorname{Ann} L(\lambda)}$ whose closure is the associated variety of Ann $L(\lambda)$. But in general, it is a very difficult problem to determine the associated variety of a highest weight module. The highest weight modules with minimal Gelfand-Kirillov dimension (or $\mathcal{O}_{\operatorname{Ann} L(\lambda)} = \mathcal{O}_{\min}$ is the unique minimal nilpotent orbit of \mathfrak{g}) play important roles in the study of representations of Lie groups and Lie algebras. Joseph found some necessary and sufficient condition for weak quantization (if there is a $U(\mathfrak{g})$ -module whose associated variety is the given orbital variety) of these minimal orbital varieties, and he also classified those minimal highest weight modules whose annihilator ideals are completely prime. In this paper, we will give a classification of minimal highest weight modules with their annihilator ideals being not necessarily completely prime for all complex simple Lie algebras. We also describe the associated varieties of these modules. By comparing with Joseph's work, we have found all minimal highest weight modules which can weakly quantize a given minimal orbital variety (in the sense of Joseph).

Key Words: Highest weight module, associated variety, orbital variety, Kazhdan-Lusztig cell, Gelfand-Kirillov dimension.

1. INTRODUCTION

Let \mathfrak{g} be a complex simple Lie algebra. For every finitely generated $U(\mathfrak{g})$ -module M, Bernstein [Ber71] constructed a variety V(M) in \mathfrak{g}^* , which is called the associated variety of M and whose dimension is equal to the Gelfand-Kirillov dimension of M. Let $L(\lambda)$ be the simple highest weight \mathfrak{g} -module with highest weight $\lambda - \rho$, where ρ is half the sum of all the positive roots. For an element w of the Weyl group W, we put $L_w := L(-w\rho)$. Tanisaki [Tan88] showed that there exist examples with reducible associated varieties in type B and C. For a long time, people believed that $V(L_w)$ is irreducible for type A, see [BB85, Mel93]. However, counterexamples of type A were found by Williamson [Will5] in 2014. The structure of $V(L_w)$ or $V(L(\lambda))$ is still mysterious.

For any simple complex Lie algebra \mathfrak{g} , there is a unique non-zero nilpotent orbit \mathcal{O}_{min} of minimal dimension. From Wang [Wan99], we know that dim $\mathcal{O}_{min} = 2(\rho, \beta^{\vee})$, where β is the highest root. From [Jos85] and [BB85], the associated variety of the annihilator ideal of any highest weight module with minimal Gelfand-Kirillov dimension (ρ, β^{\vee}) will equal to the closure of the minimal orbit \mathcal{O}_{min} . In this paper, we simply call these modules minimal highest weight modules. Similarly the Harish-Chandra modules with minimal GK dimension will be called minimal

²⁰¹⁰ Mathematics Subject Classification. Primary 22E47; Secondary 17B08.

Harish-Chandra modules. They play an important role in the study of representations of Lie groups and Lie algebras. For example, Joseph [Jos98] used some minimal highest weight modules (the annihilator ideals of these module are completely prime) to study quantization problems of minimal orbital varieties. Mathieu [Mat00] used some minimal highest weight modules in the classification of weight modules for type A and C. Meng [Men11] found that some unitary minimal highest weight modules satisfy a family of quadratic relations when he reconstructed the various Kepler-type problems in the unified language of Euclidean Jordan algebras. Hilgert-Kobayashi-Möllers [HKM14] constructed some unitary minimal highest weight modules whose annihilator ideals are completely prime. Tamori [Tam] classified irreducible minimal Harish-Chandra modules whose annihilator ideals are completely prime for type A. Some more examples can be found in [BBL97, GS05, Li00, Sun08]. These previous studies on minimal highest weight modules focused on modules whose annihilator ideals are completely prime. In this article, we will give a complete classification of minimal highest weight modules for all complex simple Lie algebras. From Bernstein-Gelfand [BG80], problems about Harish-Chandra modules can often be transformed into problems about highest weight modules. So the classification of minimal Harish-Chandra modules will be given in our next paper. As Kobayashi [Kob11] said that small representations of a group=large symmetries in a representation space, we think that these minimal highest weight modules will continue to play important role in representation theory of Lie groups and Lie algebras.

Let \mathfrak{g} be a complex simple Lie algebra with adjoint group G. Let $B \subseteq G$ be a Borel subgroup and let $\mathfrak{n} \subseteq \mathfrak{g}$ be the nilpotent radical of the Lie algebra of B. Fix a triangular decomposition $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{h} \oplus \overline{\mathfrak{n}}$ such that \mathfrak{h} is a Cartan subalgebra. Let Φ be the root system of $(\mathfrak{g}, \mathfrak{h})$ with the positive system Φ^+ and simple system Δ . Denote by W the Weyl group of Φ . Let w_0 be the longest element of W. For a nilpotent orbit \mathcal{O} , each irreducible component of $\overline{\mathcal{O}} \cap \mathfrak{n}$ is called an *orbital variety* by Joseph [Jos98]. When $\mathcal{O} = \mathcal{O}_{min}$, each irreducible component of $\overline{\mathcal{O}} \cap \mathfrak{n}$ is called a *minimal orbital variety*. An orbital variety is called *weakly quantizable* by Joseph if it is the associated variety of a highest weight module. Joseph [Jos98] classified those minimal highest weight modules whose annihilator ideals are completely prime. Such ideals are called *Joseph ideals*, which are unique when \mathfrak{g} is not of type A. For type A, there is a one parameter family of such ideals. See [Jos76] for more details. Savin noticed that there is a gap in the proof of in [Jos76, Lemma 8.8]. Then Gan and Savin [GS04] provided a simple proof of the uniqueness of the Joseph ideal.

The first purpose of this paper is to determine the highest weights of minimal highest weight modules for all complex simple Lie algebras, with their annihilator ideals being not necessarily completely prime. Our method is based on the algorithms of Gelfand-Kirillov dimensions for highest weight modules found by Bai-Xie [BX19] and Bai-Xiao-Xie [BXX]. Our results in type A and C imply that the minimal highest weight modules appeared in Mathieu [Mat00] exhaust all minimal highest weight modules. The modules in Sun [Sun08] also exhaust all minimal highest weight modules in type C, and all of them are Harish-Chandra modules. We also note that Tamori [Tam] classified the minimal highest weight modules whose annihilator ideals are completely prime. So there is some overlap with our results in type A.

The second purpose of this paper is to determine the associated varieties of these minimal highest weight modules. In our paper, we use \sim_L , \sim_R , \sim_{LR} to denote the equivalence relations of Kazhdan-Lusztig cells, see [KL79] or §2.3. We denote $I_w = \operatorname{Ann}(L_w)$ and from [BB82] and [Jos85], we have $V(I_w) := V(U(\mathfrak{g}/I_w) = \overline{\mathcal{O}}_w)$, where \mathcal{O}_w is a special nilpotent orbit (in the sense of Lusztig [Lus79]). There is a one-to-one correspondence between two-sided cells and special nilpotent orbits. We know that two primitive ideals $I_x = I_y$ if and only if $x \sim_L y$, and $V(I_x) = V(I_y)$ if and only if $x \sim_{LR} y$. So in general there are many left cells in a given two-sided cell. Joseph ideal is unique when \mathfrak{g} is not of type A. It only corresponds to one left cell in the two-sided cell corresponding to the minimal nilpotent orbit (when it is special). In this paper, our minimal highest weight modules may have annihilator ideals equal to other primitive ideals corresponding to the other left cells.

Let us introduce more notation. For $\lambda \in \mathfrak{h}^*$, define

$$\Phi_{[\lambda]} := \{ \alpha \in \Phi \mid \langle \lambda, \alpha^{\vee} \rangle \in \mathbb{Z} \}$$
$$W_{[\lambda]} := \{ w \in W \mid w\lambda - \lambda \in \mathbb{Z}\Phi \}$$

Then $\Phi_{[\lambda]}$ is a root system with Weyl group $W_{[\lambda]}$. For any $\lambda \in \mathfrak{h}^*$, there exists a unique shortest element $w_{\lambda} \in W_{[\lambda]}$ such that $\mu = w_{\lambda}^{-1}\lambda$ is anti-dominant. Thus w_{λ} is a minimal length coset representative of $W_{[\mu]}/W_J$, where $J = \{\alpha \in \Delta_{[\mu]} \mid \langle \mu, \alpha^{\vee} \rangle = 0\}$.

For $\lambda \in \mathfrak{h}^*$, put

(1.1)
$$I_{\lambda} = \{ \alpha \in \Delta \mid \langle \lambda, \alpha^{\vee} \rangle \in \mathbb{Z}_{>0} \}.$$

Then we have the following.

Theorem 1.1 (Theorem 4.2). Let $\lambda \in \mathfrak{h}^*$. If $L(\lambda)$ is a minimal highest weight module, then

$$V(L(\lambda)) = \bigcup_{\alpha \in \Delta \setminus I_{\lambda}} \overline{Be_{\alpha}}.$$

Moreover, I_{λ} contains all the simple short roots.

By using Tanisaki's result [Tan88, Conjecture 3.4] on highest weight modules, we will have the following.

Proposition 1.2. Suppose \mathfrak{g} is of classical type. Let $w, y \in W$, then $w \sim_R y$ if and only if $V(L_w) = V(L_y)$.

For different type Lie algebra \mathfrak{g} , a minimal highest weight module $L(\lambda)$ may be integral or nonintegral.

Theorem 1.3 (Theorem 5.1). Let $\lambda \in \mathfrak{h}^*$ be integral. Then $L(\lambda)$ is minimal if and only if there exist $\gamma_1, \gamma_2 \cdots, \gamma_m \in \Delta$ $(m \ge 1)$ such that

- (1) λ is regular or semiregular (i.e., there is the unique positive root $\gamma \in \Phi^+$ such that $\langle \lambda, \gamma^{\vee} \rangle = 0$).
- (2) $s_{\gamma_m} \cdots s_{\gamma_2} s_{\gamma_1} \lambda$ is dominant.
- (3) For any $k \leq m$, γ_k is the unique simple root with $\langle s_{\gamma_{k-1}} \cdots s_{\gamma_1} \lambda, \gamma_k^{\vee} \rangle \leq 0$. If λ is semiregular, the equality holds if and only if k = m.

Moreover, an integral module $L(\lambda)$ can be minimal when Φ is of type A, D or E.

Corollary 1.4 (Corollary 5.5). Let Φ be irreducible and $\lambda \in \mathfrak{h}^*$. Suppose $L(\lambda)$ is integral and minimal. Then $V(L(\lambda)) = V(L(w_{\lambda})) = \overline{Be_{\alpha_i}}$ if and only if $w_{\lambda} \sim_R$

 $s_{\alpha_i}w_0$ for some $1 \leq i \leq n$. Moreover, an integral module can be minimal when Φ is of type A, D or E.

Theorem 1.5 (Theorem 6.1). Let Φ be irreducible and $\lambda \in \mathfrak{h}^*$. Then $L(\lambda)$ is nonintegral minimal if and only if λ is dominant regular and $\Phi_{[\lambda]}$ is given in Table 2. In this case, $\operatorname{GKdim} L(\lambda) = |\Phi^+| - |\Phi^+_{[\lambda]}|$. Moreover, a nonintegral module can be minimal when Φ is of type A, B, C, F or G.

For classical types, the characterization of minimal highest weight modules and the associated varieties of these modules are explicitly given in Theorem 7.2, Theorem 7.4, Theorem 7.6 and Theorem 7.8.

Organizations. This paper is organized as follows. In §2, we give some necessary preliminaries. In §3, we recall the algorithms of computing GK dimensions in [BXX] and some properties of **a**-functions. In §4, we determine the associated varieties of all the minimal highest weight modules. In §5, we give some characterizations for integral minimal highest weight modules. In §6, we give some characterizations for nonintegral minimal highest weight modules. In §7, we give some more explicit characterizations of minimal highest weight modules for all classical Lie algebras.

2. NOTATION AND PRELIMINARIES

Let \mathfrak{g} be a complex simple Lie algebra with adjoint group G. Write $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{h} \oplus \overline{\mathfrak{n}}$ such that \mathfrak{h} is a Cartan subalgebra and $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}$ is a Borel subalgebra of \mathfrak{g} . Here \mathfrak{n} is the nilradical of \mathfrak{b} and $\overline{\mathfrak{n}}$ is its dual space in \mathfrak{g} relative the Killing form. Let Bbe the Borel subgroup of G corresponding to \mathfrak{b} . Let Φ be the root system of $(\mathfrak{g}, \mathfrak{h})$ with the positive system Φ^+ and simple system Δ determined by \mathfrak{b} . Denote by Wthe Weyl group of Φ . Let w_0 be the longest element of W.

Note that any subset $I \subseteq \Phi$ generates a subsystem $\Phi_I \subseteq \Phi$ with corresponding Weyl group $W_I \subseteq W$. Let w_I be the longest element of W_I . When $I \subseteq \Delta$, let \mathfrak{p}_I be the standard parabolic subalgebra corresponding to I with Levi decomposition $\mathfrak{p}_I = \mathfrak{l}_I \oplus \mathfrak{u}_I$. We will frequently drop the subscript if there is no confusion.

For $\lambda \in \mathfrak{h}^*$, denote by $L(\lambda)$ the simple highest weight \mathfrak{g} -module with highest weight $\lambda - \rho$, where ρ is the half sum of positive roots. Then $L(\lambda)$ admits an central character $\chi_{\lambda} : Z(\mathfrak{g}) \to \mathbb{C}$ such that $z \cdot v = \chi_{\lambda}(z)v$ for any $z \in Z(\mathfrak{g})$ and $v \in L(\lambda)$, where $Z(\mathfrak{g})$ is the center of the universal enveloping algebra $U(\mathfrak{g})$ of \mathfrak{g} . In particular, $\chi_{\lambda} = \chi_{\mu}$ if and only if $\lambda = w\mu$ for some $w \in W$. Thus the set of central characters can be parameterized by

$$\mathfrak{D} = \left\{ \nu \in \mathfrak{h}^* \left| \begin{aligned} \operatorname{Re}\langle \nu, \alpha^{\vee} \rangle \geq 0 \text{ and whenever } \operatorname{Re}\langle \nu, \alpha^{\vee} \rangle = 0, \\ \operatorname{then } \operatorname{Im}\langle \nu, \alpha^{\vee} \rangle \geq 0, \text{ for all } \alpha \in \Delta \end{aligned} \right\}$$

where $\langle -, - \rangle$ is a bilinear form on \mathfrak{h}^* induced from the Killing form on \mathfrak{g} (e.g., [Hum08, §0.2]). We say $L(\lambda)$ has *infinitesimal character* ν when $\chi_{\lambda} = \chi_{\nu}$ for some $\nu \in \mathfrak{D}$. Put $L_w := L(-w\rho)$ for $w \in W$. They form all the simple highest weight modules with infinitesimal character ρ .

We say $\lambda \in \mathfrak{h}^*$ is regular if $\langle \lambda, \alpha^{\vee} \rangle \neq 0$ for all $\alpha \in \Phi$, otherwise λ is called singular. We say λ is dominant (resp. anti-dominant) if $\langle \lambda, \alpha^{\vee} \rangle \notin \mathbb{Z}_{<0}$ (resp. $\mathbb{Z}_{>0}$) for all $\alpha \in \Phi$. Thus any $\nu \in \mathcal{D}$ is dominant.

2.1. **GK dimensions and associated varieties.** Let M be a finite generated $U(\mathfrak{g})$ -module M. Choose a finite dimensional generating subspace M_0 of M. Let $\{U_n(\mathfrak{g})\}_{n\geq 0}$ be the standard filtration of $U(\mathfrak{g})$. Write $M_n = U_n(\mathfrak{g}) \cdot M_0$. Set

$$\operatorname{gr}(M) = \bigoplus_{n=0}^{\infty} \operatorname{gr}_n M,$$

where $\operatorname{gr}_n M = M_n/M_{n-1}$. Then $\operatorname{gr}(M)$ is a graded module of $\operatorname{gr}(U(\mathfrak{g})) \simeq S(\mathfrak{g})$.

Definition 2.1. The *Gelfand-Kirillov dimension* of *M* is defined by

$$\operatorname{GKdim} M = \frac{\lim_{n \to \infty} \log \dim(U_n(\mathfrak{g})M_0)}{\log n}$$

The associated variety of M is defined by

$$V(M) := \{ X \in \mathfrak{g}^* \mid p(X) = 0 \text{ for all } p \in \operatorname{Ann}_{S(\mathfrak{g})}(\operatorname{gr} M) \}.$$

In particular, we have (see e.g., [NOT01])

(2.1)
$$\dim V(M) = \operatorname{GKdim} M.$$

Since we can identify \mathfrak{g}^* with \mathfrak{g} via the Killing form on \mathfrak{g} , V(M) can also be viewed as a subvariety of \mathfrak{g} .

One can also define the associated variety for a two-sided ideal of $U(\mathfrak{g})$.

Definition 2.2. Let *I* be a two-sided ideal in $U(\mathfrak{g})$. Then $\operatorname{gr}(U(\mathfrak{g})/I) \simeq S(\mathfrak{g})/\operatorname{gr} I$ is a graded $S(\mathfrak{g})$ -module with annihilator $\operatorname{gr} I$. The associated variety of *I* is

$$V(I) := V(U(\mathfrak{g})/I) = \{ X \in \mathfrak{g}^* \mid p(X) = 0 \text{ for all } p \in \operatorname{gr} I \}.$$

For a highest weight module M, we have (see [Jos78, Cor. 2.8])

(2.2)
$$\dim V(\operatorname{Ann}(M)) = 2 \dim V(M).$$

If M_0 is a-invariant for a subalgebra a of g, then (see [Vog78, (1.5)(b)])

(2.3)
$$V(M) \subseteq V(\operatorname{Ann}(M)) \cap (\mathfrak{g}/\mathfrak{a})^*.$$

In particular, if M can be generated by a finite dimensional \mathfrak{b} -invariant space, then $V(M) \subseteq (\bar{\mathfrak{n}})^* \simeq \mathfrak{n}$, where the last isomorphism is induced from the Killing form.

Lemma 2.3. For every $\lambda \in \mathfrak{h}^*$, the associated variety $V(\operatorname{Ann}(L(\lambda)))$ is the closure $\overline{\mathcal{O}}$ of a nilpotent orbit \mathcal{O} .

This result can be found in [Jos85, 3.10] or [CM93, Theorem 10.2.2]. Suppose that $\mathcal{O} \subseteq \mathfrak{g}$ is a nilpotent *G*-orbit with closure $\overline{\mathcal{O}}$ [CM93]. Each irreducible component of $\overline{\mathcal{O}} \cap \mathfrak{n}$ is called an *orbital variety*, which can be written as $\mathcal{V}(w) := \overline{B(\mathfrak{n} \cap w\mathfrak{n})}$ for some $w \in W$ [Tan88].

The following result is known as Spaltenstein-Steinberg equality [Jos84, 3.1].

Lemma 2.4. Let \mathbb{O} be an orbital variety of a nilpotent orbit \mathcal{O} . Then dim $\mathbb{O} = \frac{1}{2} \dim \mathcal{O}$.

The next result is due to Joseph [Jos84, 4.7] and Borho-Brylinski [BB85].

Lemma 2.5. Let $\lambda \in \mathfrak{h}^*$. Then $V(L(\lambda))$ is a union of some orbital varieties contained in $\overline{\mathcal{O}} \cap \mathfrak{n}$, where $\overline{\mathcal{O}} = V(\operatorname{Ann}(L(\lambda)))$. Moreover, $\mathcal{V}(w) \subseteq V(L_w)$.

2.2. Minimal orbits and minimal modules. There is a unique nonzero nilpotent orbit \mathcal{O}_{min} of minimal dimension for any simple Lie algebra \mathfrak{g} [CM93]. Then each irreducible component of $\overline{\mathcal{O}}_{min} \cap \mathfrak{n}$ is called a *minimal orbital variety* by Joseph [Jos98].

Definition 2.6. We call a simple $U(\mathfrak{g})$ -module M minimal if

(2.4)
$$V(\operatorname{Ann}(M)) = \overline{\mathcal{O}}_{\min}$$

Remark 2.7. In [GS05], a Harish-Chandra module M is called minimal if Ann(M)is the Joseph ideal, which also satisfies (2.4). In our Definition 2.6, the annihilator ideal $\operatorname{Ann}(M)$ is not necessarily completely prime.

The following result is an easy consequence of Lemma 2.5.

Lemma 2.8. If $L(\lambda)$ is a minimal highest weight module, then $V(L(\lambda))$ is a union of some minimal orbital varieties.

An orbital variety is called *weakly quantizable* [Jos98] if it is the associated variety of a highest weight module. By Joseph $[Jos98, \S4.12]$, the following proposition holds.

Proposition 2.9. Let \mathbb{O} be a minimal orbital variety. Then

- (1) $\mathbb{O} = \overline{Be_{\alpha}} = \mathcal{V}(s_{\alpha}w_0)$ for some long simple root $\alpha \in \Delta$; (2) \mathbb{O} is weakly quantizable except when $\mathbb{O} = \overline{Be_{\alpha_i}}$ for $2 \leq i \leq n-2$ in type $B_n (n \ge 4).$

We will see that minimal modules always exist for any fixed simple Lie algebra g. Hence, according to (2.1), (2.2) and Lemma 2.3, the minimal highest weight modules can be characterized as those $L(\lambda)$ with the minimal (nonzero) GK dimension, which is equal to $\frac{1}{2} \dim \mathcal{O}_{\min} = (\rho, \beta^{\vee})$ by [Wan99], where $\beta \in \Phi^+$ is the highest root (e.g., [Hum78, §13.4]). The strategy for finding them is simple: we investigate the highest weight modules with minimal possible GK dimension when an infinitesimal character is fixed. The following notation is useful.

Definition 2.10. Fix $\nu \in \mathfrak{D}$. We say $L(\lambda)$ is ν -minimal if $\chi_{\lambda} = \chi_{\nu}$ and $L(\lambda)$ has minimal possible nonzero GK dimension among all the simple highest weight modules with infinitesimal character ν .

Obviously, a minimal highest weight module is always ν -minimal for some $\nu \in \mathfrak{D}$, but not vice versa. The values of the minimal and ρ -minimal GK dimension for each type are given in Table 1, see [Vog81, Lus84, BXX].

	A_n	B_n	C_n	D_n	E_6	E_7	E_8	F_4	G_2
min of $\operatorname{GKdim} L(\lambda)$	n	2n - 2	n	2n - 3	11	17	29	8	3
min of $\operatorname{GKdim} L_w$	n	2n - 1	2n - 1	2n - 3	11	17	29	11	5

TABLE 1. Minimal (nonzero) GK dimensions

Minimal highest weight modules could be integral or nonintegral. For the integral case, we need another small orbit \mathcal{O}_s , which is called *minimal special* since it

is minimal among all non-trivial special nilpotent orbits (see [CM93, §6.3] for a definition of special orbit). From Humphreys [Hum16], we have dim $\mathcal{O}_s = 2h - 2 = 2(\rho, \beta_s^{\vee})$, where h is the Coxeter number and β_s is the unique highest short root. Thus $\mathcal{O}_s = \mathcal{O}_{min}$ when Φ is simply-laced.

Definition 2.11. We call a simple $U(\mathfrak{g})$ -module M special minimal if

(2.5)
$$V(\operatorname{Ann}(M)) = \overline{\mathcal{O}}_s$$

Since \mathcal{O}_s is the unique nilpotent orbit with dimension $2(\rho, \beta_s^{\vee})$, a module M is special minimal if and only if

A well-known result is that L_w is ρ -minimal if and only if it is special minimal (see Table 1). We will present a characterization of integral special minimal highest weight modules in section 5.

3. Algorithm for GK dimensions

In this section, we will describe the algorithm obtained in [BX19, BXX] for GK dimension of highest weight modules. For $\lambda \in \mathfrak{h}^*$, define

$$\Phi_{[\lambda]} := \{ \alpha \in \Phi \mid \langle \lambda, \alpha^{\vee} \rangle \in \mathbb{Z} \}$$
$$W_{[\lambda]} := \{ w \in W \mid w\lambda - \lambda \in \mathbb{Z}\Phi \}.$$

Then $\Phi_{[\lambda]}$ is a root system with Weyl group $W_{[\lambda]}$. Let $\Delta_{[\lambda]}$ be the simple system of $\Phi_{[\lambda]}$. For any $\lambda \in \mathfrak{h}^*$, there exists a unique shortest element $w_{\lambda} \in W_{[\lambda]}$ such that $\mu = w_{\lambda}^{-1}\lambda$ is anti-dominant. Thus $W_{[\lambda]} = W_{[\mu]}$ and w_{λ} is a minimal length coset representative of $W_{[\mu]}/W_J$, where $J = \{\alpha \in \Delta_{[\mu]} \mid \langle \mu, \alpha^{\vee} \rangle = 0\}$. More results about integral subsystem can be found in [Hum08], [BXX, §5.3] and [XZ22, §4.1]

Proposition 3.1 ([BX19, Prop. 3.8]). Let $\lambda \in \mathfrak{h}^*$. Then

$$\operatorname{GKdim} L(\lambda) = |\Phi^+| - \mathbf{a}_{[\lambda]}(w_\lambda),$$

where $\mathbf{a}_{[\lambda]}$ is the **a**-function on the sub Weyl group $W_{[\lambda]}$.

Here the map $\mathbf{a}: W \to \mathbb{N}$ is known as Lusztig's **a**-function [Lus03]. We will explain more about it in the following.

Now we consider the **a**-functions appeared in Proposition 3.1. A good reference is [Lus03]. Recall that the Weyl group W of \mathfrak{g} is a Coxeter group generated by $S = \{s_{\alpha} \mid \alpha \in \Delta\}$. Let $\ell(-)$ be the *length function* on W. The Hecke algebra \mathcal{H} over $\mathcal{A} := \mathbb{Z}[v, v^{-1}]$ is generated by $T_w, w \in W$ with relations

$$T_w T_{w'} = T_{ww'}$$
 if $\ell(ww') = \ell(w) + \ell(w')$,
and $(T_s + v^{-1})(T_s - v) = 0$ for any $s \in S$.

The unique elements C_w such that

 $\overline{C_w} = C_w, \qquad C_w \equiv T_w \mod \mathcal{H}_{<0}$

is known as the Kazhdan-Lusztig basis of \mathcal{H} , where $\bar{}: \mathcal{H} \to \mathcal{H}$ is the bar involution such that $\bar{q} = q^{-1}$, $\overline{T_w} = T_{w^{-1}}^{-1}$, and $\mathcal{H}_{<0} = \bigoplus_{w \in W} \mathcal{A}_{<0} T_w$ with $\mathcal{A}_{<0} = v^{-1} \mathbb{Z}[v^{-1}]$. If C_y occurs in the expansion of hC_w (resp. C_wh) with respect to the KL-basis

If C_y occurs in the expansion of hC_w (resp. C_wh) with respect to the KL-basis for some $h \in \mathcal{H}$, then we write $y \leftarrow_L w$ (resp. $y \leftarrow_R w$). Extend \leftarrow_L (resp. \leftarrow_R) to a preorder \leq_L (resp. \leq_R) on W. For $x, w \in W$, write $x \leq_{LR} w$ if there exists $x = w_0, w_1, \dots, w_n = w$ such that for every $0 \le i < n$ we have either $w_i \le_L w_{i+1}$ or $w_i \le_R w_{i+1}$. Let $\sim_L, \sim_R, \sim_{LR}$ be the equivalence relations associated with \le_L , \le_R, \le_{LR} (for example, $x \sim_L w$ if and only if $x \le_L w$ and $w \le_L x$). The equivalence classes on W for \le_L, \le_R, \le_{LR} are called *left cells*, *right cells* and *two-sided cells* respectively. The following result is straightforward.

Lemma 3.2. Let $x, y, z \in W$. If z = xy and $\ell(z) = \ell(y) + \ell(x)$, then $z \leq_L y$ and $z \leq_R x$.

Obviously {1} (resp. $\{w_0\}$) is the largest (smallest) two sided-cell of W. Let C be the set of elements $w \in W \setminus \{1\}$ such that w has a unique reduced expression. Denote $Cw_0 := \{ww_0 \mid w \in C\}$.

Lemma 3.3. Let $w \in W$, then

- (1) The sets C and Cw_0 are two-sided cells of W.
- (2) If $w \neq 1, w_0$, then $x \leq_{LR} w \leq_{LR} y$ for any $x \in \mathcal{C}w_0$ and $y \in \mathcal{C}$.

Proof. (1) is an easy consequence of [Lus83, Prop. 3.8] and [KL79, Remark 3.3]. (2) Since $w \neq 1$, there exists $s \in S$ such that $\ell(sw) < \ell(w)$. Lemma 3.2 implies that $w \leq_R s$. Thus $w \leq_{LR} y$ in view of $s \in S \subseteq C$ and (1). Similarly, we can show that $x \leq_{LR} w$.

Write $C_x C_y = \sum_{z \in W} h_{x,y,z} C_z$ with $h_{x,y,x} \in \mathcal{A}$. Then $\mathbf{a} : W \to \mathbb{N}$ is defined by $\mathbf{a}(z) = \max\{\deg h_{x,y,z} \mid x, y \in W\}$ for $z \in W$.

The following lemma is an easy consequence of Lusztig's results [Lus03, §14.2].

Lemma 3.4. Let $x, w \in W$. Then

- (1) $\mathbf{a}(w) = \mathbf{a}(w^{-1}).$
- (2) If $x \leq_{LR} w$, then $\mathbf{a}(x) \geq \mathbf{a}(w)$. Hence $\mathbf{a}(x) = \mathbf{a}(w)$ whenever $x \sim_{LR} w$.
- (3) If w_I is the longest element of the parabolic subgroup of W generated by simply reflections of $I \subseteq \Delta$, then $\mathbf{a}(w_I)$ is equal to the length $\ell(w_I)$ of w_I .
- (4) If W is a direct product of Coxeter subgroups W_1 and W_2 , then

$$\mathbf{a}(w) = \mathbf{a}(w_1) + \mathbf{a}(w_2)$$

for $w = (w_1, w_2) \in W_1 \times W_2 = W$.

Lemma 3.5. Let $w \in W$. If $w \notin C \cup Cw_0 \cup \{1, w_0\}$, then

$$\mathbf{a}(1) = 0 < 1 = \mathbf{a}(\mathcal{C}) < \mathbf{a}(w) < \mathbf{a}(\mathcal{C}w_0) < \mathbf{a}(w_0) = \ell(w_0) = |\Phi^+|.$$

Proof. In view of [Lus03, Prop. 13.7, 13.8], we get $\mathbf{a}(1) = 0 < \mathbf{a}(\mathcal{C})$ and $\mathbf{a}(\mathcal{C}w_0) < \ell(w_0) = \mathbf{a}(w_0)$. Lemma 3.4 implies $\mathbf{a}(\mathcal{C}) = \mathbf{a}(s) = 1$ for any $s \in S$, while Lemma 3.3 yields $\mathbf{a}(\mathcal{C}) \leq \mathbf{a}(w) \leq \mathbf{a}(\mathcal{C}w_0)$. If $\mathbf{a}(\mathcal{C}) = \mathbf{a}(w)$, we obtain $w \in \mathcal{C}$ by [Lus03, §14.2 P11] and [BV83, Corollary 2.15], a contradiction. We can get a similar contradiction when $\mathbf{a}(w) = \mathbf{a}(\mathcal{C}w_0)$.

Remark 3.6. From Lemma 3.5, an integral module M is special minimal if and only if

(3.1)
$$\operatorname{GKdim} M = \frac{1}{2} \operatorname{dim} \mathcal{O}_s = (\rho, \beta_s^{\vee}) = |\Phi^+| - \mathbf{a}(w_0 \mathcal{C}).$$

4. Associated varieties of highest weight modules

In this section, we determine the associated varieties of all the minimal highest weight modules (Theorem 4.2). We will also recall the relation between associated varieties and cells.

4.1. Associated varieties of minimal highest weight modules. Recall that any subset $I \subseteq \Delta$ generates a subsystem $\Phi_I \subseteq \Phi$ with corresponding parabolic subalgebra $\mathfrak{p} = \mathfrak{l} \oplus \mathfrak{u} \supset \mathfrak{b}$ of \mathfrak{g} .

We say $\lambda \in \mathfrak{h}^*$ is Φ_I^+ -dominant if and only if $\langle \lambda, \alpha^{\vee} \rangle \in \mathbb{Z}_{>0}$ for all $\alpha \in I$. Let $\bar{\mathfrak{u}}$ be the nilpotent subalgebra opposite to \mathfrak{u} . Then $\bar{\mathfrak{n}} = \bar{\mathfrak{u}} \oplus \bar{\mathfrak{n}}_I$ and $\mathfrak{p} = \bar{\mathfrak{n}}_I \oplus \mathfrak{b}$ where $\bar{\mathfrak{n}}_I := \bar{\mathfrak{n}} \cap \mathfrak{l}$. The following result is a generalization of [Jos84, Lem. 6.1].

Proposition 4.1. Let \mathfrak{g} be a simple complex Lie algebra and $\lambda \in \mathfrak{h}^*$. Then λ is Φ_I^+ -dominant if and only if $V(L(\lambda)) \subseteq \mathfrak{u}$.

Proof. Denote by $F(\lambda)$ the simple \mathfrak{l} -module with highest weight $\lambda - \rho$. Then $F(\lambda)$ is finite dimensional if and only if λ is Φ_I^+ -dominant.

First assume that λ is Φ_I^+ -dominant. Since $F(\lambda)$ can be viewed as a p-module with trivial u-action, we obtain a surjective homomorphism

$$\varphi: U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} F(\lambda) \to L(\lambda).$$

Then $L(\lambda)$ is generated by $\varphi(F(\lambda)) \simeq F(\lambda)$, which is p-invariant. Thus (2.3) implies $V(L(\lambda)) \subseteq (\mathfrak{g}/\mathfrak{p})^* \simeq (\bar{\mathfrak{u}})^* \simeq \mathfrak{u}$.

Next assume that $V(L(\lambda)) \subseteq \mathfrak{u}$. Let v^+ be a nonzero highest weight vector of $M = L(\lambda)$. We claim that $U(\bar{\mathfrak{n}}_I) \cdot v^+$ is finite-dimensional. If this does hold, then $F(\lambda) \subseteq U(\bar{\mathfrak{n}}_I) \cdot v^+$ is finite-dimensional and λ is Φ_I^+ -dominant. Now we prove the claim. Start with $M_0 = \mathbb{C}v^+$, which is a 1-dimensional \mathfrak{b} -invariant generating space of M. It gives a \mathfrak{b} -invariant filtration $M_n = U_n(\mathfrak{g}) \cdot M_0 = U_n(\bar{\mathfrak{n}}) \cdot M_0$. This implies that $V(M) \subseteq (\mathfrak{g}/\mathfrak{b})^* \simeq \mathfrak{n}$ in view of (2.3). Since the subvariety $\mathfrak{u} \simeq (\bar{\mathfrak{u}})^* \subseteq \mathfrak{n}$ is defined by the ideal $\bar{\mathfrak{n}}_I S(\bar{\mathfrak{n}})$, the condition $V(M) \subseteq \mathfrak{u}$ yields that the $\bar{\mathfrak{n}}_I$ -action on grM is nilpotent. In other words, there is a positive integer j with $U_j(\bar{\mathfrak{n}}_I) \cdot M_i \subseteq M_{i+j-1}$ for all $i \in \mathbb{N}$. Fix $\alpha \in I$. When i = 0, one has

(4.1)
$$e^{j}_{-\alpha} \cdot v^{+} = u \cdot v^{+}$$

for a $u \in U_{j-1}(\bar{\mathbf{n}})$. If $u \neq 0$, the weight of u must be of the form $\beta_1 + \cdots + \beta_k$ for a $k \leq j-1$, where $-\beta_i \in \Phi^+$. Taking the weights of both vectors in (4.1), we get $j\alpha = -(\beta_1 + \cdots + \beta_k)$ for a $k \leq j-1$. This can not hold for simple root $\alpha \in I$. We obtain $e_{-\alpha}^j \cdot v^+ = 0$. Hence $U(\bar{\mathbf{n}}_I)v^+$ is finite-dimensional. \Box

Recall that $I_{\lambda} = \{ \alpha \in \Delta \mid \langle \lambda, \alpha^{\vee} \rangle \in \mathbb{Z}_{>0} \}$ in (1.1). Obviously λ is $\Phi^+_{I_{\lambda}}$ -dominant. **Theorem 4.2.** Let $\lambda \in \mathfrak{h}^*$. If $L(\lambda)$ is a minimal highest weight module, then

$$V(L(\lambda)) = \bigcup_{\alpha \in \Delta \setminus I_{\lambda}} \overline{Be_{\alpha}}.$$

Moreover, I_{λ} contains all the simple short roots.

Proof. Set $I = I_{\lambda}$. By Lemma 2.8 and Proposition 2.9, we can write $V(L(\lambda)) = \bigcup_{\alpha \in T} \overline{Be_{\alpha}}$ for a subset $T \subseteq \Delta$ which contains no short root. Since λ is Φ_I^+ -dominant, Proposition 4.1 yields $e_{\alpha} \in \mathfrak{u}_I$ for $\alpha \in T$. This forces $T \subseteq \Delta \setminus I$. On the other hand, with $e_{\alpha} \in \mathfrak{u}_{\Delta \setminus T}$ for $\alpha \in T$, one has $V(L(\lambda)) \subseteq \mathfrak{u}_{\Delta \setminus T}$. Thus λ is $\Phi_{\Delta \setminus T}^+$ -dominant by Proposition 4.1. We obtain $\Delta \setminus T \subseteq I$. Hence $T = \Delta \setminus I$.

4.2. Right cells and associated varieties. Denote $I_w = \operatorname{Ann}(L_w)$. By Borho-Brylinski [BB82] and Joseph [Jos85], we know that $V(I_w) = V(U(\mathfrak{g})/I_w)$ is the closure of a single special (in the sense of Lusztig [Lus79]) nilpotent orbit. From [KL79] and [BV83], there is a bijection between special nilpotent orbits of \mathfrak{g} and two-sided cells of the Weyl group W, see also [Tan88]. The special orbits are listed in Collingwood-McGovern [CM93]. For a nilpotent orbit \mathcal{O} , denote by $\operatorname{Irr}(\overline{\mathcal{O}} \cap \mathfrak{n})$ the set of irreducible components in $\overline{\mathcal{O}} \cap \mathfrak{n}$.

The following result was conjectured by Tanisaki [Tan88, Conj.3.4]. The case of type A was proved by Borho-Brylinski [BB85], while the case of the other classical types was proved by McGovern [McG00].

Theorem 4.3. Suppose that \mathfrak{g} is classical. Let \mathscr{C} be the two-sided cell corresponding to a special nilpotent orbit \mathcal{O} . Denote by \mathscr{C}_R the set of right cells contained in \mathscr{C} . Then there exists a bijection from \mathscr{C}_R to $\operatorname{Irr}(\overline{\mathcal{O}} \cap \mathfrak{n})$ $(w \to Y_w)$ and an ordering \prec on $\operatorname{Irr}(\overline{\mathcal{O}} \cap \mathfrak{n})$ such that $V(L_w) = Y_w \cup \tilde{Y}_w$, where \tilde{Y}_w is a union of some orbital varieties Y in $\operatorname{Irr}(\overline{\mathcal{O}} \cap \mathfrak{n})$ with $Y \prec Y_w$.

Remark 4.4. In Tanisaki's original conjecture [Tan88, Conj.3.4], \mathfrak{g} is not necessarily classical, where the cases of type E_6 and G_2 were proved by himself. For the case of type F_4 , Tanisaki showed that the conjecture is true for nine special nilpotent orbits among all the eleven ones. The cases of type E_7 , E_8 , F_4 are still unknown.

Proposition 4.5. Suppose \mathfrak{g} is of classical type. Let $w, y \in W$, then $w \sim_R y$ if and only if $V(L_w) = V(L_y)$.

Proof. If $w \sim_R y$, we get $V(L_w) = V(L_y)$ by [Jos84, Lem. 6.6] and [BB85, Cor. 6.3]. Now assume that $V(L_w) = V(L_y)$. In view of [BB82, Cor. 4.11] and [CM93, Thm. 10.2.2], $V(I_w) = V(I_y)$ is the closure of a nilpotent orbit \mathcal{O} . Let \mathscr{C} be the two-sided cell corresponding to \mathcal{O} in the Springer correspondence. With $V(L_w) = V(L_y)$, Theorem 4.3 yields $Y_w \cup \tilde{Y}_w = Y_y \cup \tilde{Y}_y$. This forces $Y_w = Y_y$. Hence $w \sim_R y$.

Remark 4.6. The proposition holds for all the simple Lie algebras once Tanisaki's original conjecture [Tan88, Conj. 3.4] is proved completely.

From [BBM89, Lem. 5.2], we know that the associated variety of a simple \mathfrak{g} -module is invariant under the corresponding translation functor. Recall that $\mathcal{V}(w) := \overline{B(\mathfrak{n} \cap w\mathfrak{n})}$ is the orbital variety for $w \in W$. Combined with Lemma 2.5 and [BX19, Cor. 3.3], we have the following result.

Proposition 4.7. Let $\lambda \in \mathfrak{h}^*$ be an integral weight. Then

$$V(L(\lambda)) = V(L_{w_{\lambda}}) \supseteq \mathcal{V}(w_{\lambda}).$$

Theorem 4.8. Suppose that \mathfrak{g} is classical. Let $\lambda, \mu \in \mathfrak{h}^*$ be integral. Then $V(L(\lambda)) = V(L(\mu))$ if and only if $w_{\lambda} \sim_R w_{\mu}$.

5. INTEGRAL MINIMAL HIGHEST WEIGHT MODULES

Recall that an integral minimal highest weight module must be special minimal. In this section, we will prove the following result about special minimal highest weight modules. **Theorem 5.1.** Let $\lambda \in \mathfrak{h}^*$ be integral. Then $L(\lambda)$ is special minimal if and only if there exist $\gamma_1, \gamma_2 \cdots, \gamma_m \in \Delta \ (m \ge 1)$ such that

- (1) λ is regular or semiregular.
- (2) $s_{\gamma_m} \cdots s_{\gamma_2} s_{\gamma_1} \lambda$ is dominant.
- (3) For any $k \leq m$, γ_k is the unique simple root with $\langle s_{\gamma_{k-1}} \cdots s_{\gamma_1} \lambda, \gamma_k^{\vee} \rangle \leq 0$. If λ is semiregular, the equality holds if and only if k = m.

In this case, $\operatorname{GKdim} L(\lambda) = |\Phi^+| - \mathbf{a}(w_0 \mathcal{C}) = (\rho, \beta_s^{\vee})$. Moreover, a special minimal module is minimal when Φ is of type A, D or E.

Here we say an integral weight $\lambda \in \mathfrak{h}^*$ is *semiregular* if there is the unique positive root $\gamma \in \Phi^+$ such that $\langle \lambda, \gamma^{\vee} \rangle = 0$.

Example 5.2. Let $\Phi = E_6$ and $\lambda = \rho - 2\alpha_1 = (-1, 2, 3, 4, 5, -3, -3, 3)$ (here we adopt the notation in [Hum78]). Then $\langle \lambda, \alpha_1^{\vee} \rangle = -3$ and $\langle \lambda, \alpha_i^{\vee} \rangle > 0$ for $i \neq 1$. Moreover, $s_{\alpha_1}\lambda = \rho + \alpha_1$ is dominant and semiregular (since $\langle \rho + \alpha_1, \alpha_3^{\vee} \rangle = 0$). By Theorem 5.1, $L(\lambda)$ is special minimal with GKdim $L(\lambda) = (\rho, \beta_s^{\vee}) = 11$. In view of Table 1 and Theorem 4.2, it is also minimal with associated variety $\overline{Be_{\alpha_1}}$. This is compatible with [BXX, Exam. 7.2].

Firstly we have the following proposition.

Proposition 5.3. Let $\nu \in \mathfrak{D}$ and $\lambda \in W\nu$ be integral. If ν is not regular, then the following conditions are equivalent:

- (1) GKdim $L(\lambda) = |\Phi^+| \mathbf{a}(w_0 w_J);$
- (2) $w_{\lambda} \sim_L w_0 w_J$ in W;
- (3) $L(\lambda)$ is ν -minimal.

Moreover, $L(\lambda)$ is special minimal if and only if |J| = 1, i.e., λ is semiregular.

Proof. Since $\mu = w_0 \nu$ is anti-dominant, the set $J = \{\alpha \in \Delta \mid \langle \mu, \alpha^{\vee} \rangle = 0\} \neq 0$ and thus $w_J \neq 1$ when ν is not regular.

Recall that w_{λ} is a shortest representative in W/W_J . There exists $x \in W$ such that $xw_{\lambda}w_J = w_0$ and $\ell(x) + \ell(w_{\lambda}) + \ell(w_J) = \ell(w_0)$. In view of Lemma 3.2, one has $w_0w_J \leq_L w_{\lambda}$ since $xw_{\lambda} = w_0w_J$ and $\ell(x) + \ell(w_{\lambda}) = \ell(w_0w_J) = \ell(w_0) - \ell(w_J)$. It follows from Lemma 3.4(2) that $\mathbf{a}(w_0w_J) \geq \mathbf{a}(w_{\lambda})$. Combined with Proposition 3.1 and Lemma 3.5, one has

$$\operatorname{GKdim} L(\lambda) = |\Phi^+| - \mathbf{a}(w_\lambda) \ge |\Phi^+| - \mathbf{a}(w_0 w_J) > 0$$

By [Lus03, §14.2 P9], the equality holds if and only if $w_{\lambda} \sim_L w_0 w_J$ or $L(\lambda)$ is ν -minimal.

For the second statement, Lemma 3.5 implies that $\mathbf{a}(w_0w_J)$ is maximal if and only if $w_J \in \mathcal{C}$, which means w_J has a unique reduced expression. This happens only when |J| = 1. In this case, $\operatorname{GKdim} L(\lambda) = |\Phi^+| - \mathbf{a}(w_0\mathcal{C})$ and thus $L(\lambda)$ is special minimal. \Box

The proof of Theorem 5.1. It suffices to prove the first statement since the second and the third one are easy consequence of Table 1 and (3.1).

First assume that $L(\lambda)$ is special minimal. Recall that w_{λ} is the shortest element in W with $w_{\lambda}^{-1}\lambda$ anti-dominant. Set $w := w_0 w_{\lambda}^{-1}$. Thus it is the longest element such that $\nu := w\lambda$ is dominant. Proposition 5.3 implies that λ is either regular or semiregular. Proposition 3.1 and Lemma 3.5 yield GKdim $L(\lambda) = |\Phi^+| - \mathbf{a}(w_0 \mathcal{C})$ and $w_{\lambda}^{-1} \in w_0 \mathcal{C}$. Therefore $w = w_0 w_{\lambda}^{-1} \in \mathcal{C}$ has a unique reduced expression, which we denoted by $w = s_{\gamma_m} \cdots s_{\gamma_2} s_{\gamma_1}$ $(m \ge 1)$. If λ is semiregular, then $\langle \nu, \alpha^{\vee} \rangle = 0$ for some $\alpha \in \Delta$. This forces $\gamma_m = \alpha$ since w is longest. Then (2) and (3) follows easily.

Conversely, set $w = s_{\gamma_m} \cdots s_{\gamma_2} s_{\gamma_1}$ and $\nu = w\lambda \in \mathcal{D}$. Then (3) means w has the unique reduced expression, i.e., $w \in \mathcal{C}$. If λ is regular, we obtain $w_{\lambda}^{-1} = w_0 w$ and GKdim $L(\lambda) = |\Phi^+| - \mathbf{a}(w_0 \mathcal{C})$ is special minimal, in view of Proposition 3.1 and Lemma 3.5. Now suppose that λ is semiregular and $J = \{\alpha\}$. Then (3) yields $\alpha = \gamma_m$. Hence $w_{\lambda}^{-1} = w_0 w$ and the remaining argument is similar to the regular case.

The above proof also implies the following useful result.

Proposition 5.4. Let $\lambda \in \mathfrak{h}^*$ be integral. Then $L(\lambda)$ is special minimal if and only if $w_0 w_{\lambda}^{-1} \neq 1$ and has the unique reduced expression, i.e., $w_0 w_{\lambda}^{-1} \in \mathcal{C}$.

From Theorem 4.2, Theorem 5.1 and Proposition 5.4, we have the following.

Corollary 5.5. Suppose $L(\lambda)$ is integral and minimal. Then $V(L(\lambda)) = V(L(w_{\lambda})) = \overline{Be_{\alpha_i}} = \mathcal{V}(s_{\alpha_i}w_0)$ if and only if $w_{\lambda} \sim_R s_{\alpha_i}w_0$ for some $1 \leq i \leq n$. Moreover, an integral module can be minimal when Φ is of type A, D or E.

6. Nonintegral minimal highest weight modules

Recall that any $\lambda \in \mathfrak{h}^*$ gives an integral subsystem $\Phi_{[\lambda]}$ with simple system $\Delta_{[\lambda]}$ and Weyl group $W_{[\lambda]}$. In this section, we prove the following result about nonintegral minimal modules.

Theorem 6.1. Let Φ be irreducible and $\lambda \in \mathfrak{h}^*$. Then $L(\lambda)$ is nonintegral minimal if and only if λ is dominant regular and $\Phi_{[\lambda]}$ is given in Table 2. In this case, $\operatorname{GKdim} L(\lambda) = |\Phi^+| - |\Phi^+_{[\lambda]}|$. Moreover, a nonintegral minimal module is minimal when Φ is of type A, B, C, F or G.

Φ	A_n	B_n	C_n	D_n	E_6	E_7	E_8	F_4	G_2
$\Phi_{[\lambda]}$	A_{n-1}	$B_1 \times B_{n-1}$	D_n	D_{n-1}	D_5	E_6	$A_1 \times E_7$	C_4	A_2

TABLE 2. Maximal nonintegral systems

Here we say a highest weight module M is *nonintegral minimal* if GKdim M is minimal among all nonintegral highest weight modules.

Example 6.2. Let $\Phi = F_4$ with $\Delta = \{\varepsilon_2 - \varepsilon_3, \varepsilon_3 - \varepsilon_4, \varepsilon_4, \frac{1}{2}(\varepsilon_1 - \varepsilon_2 - \varepsilon_3 - \varepsilon_4)\}$ (see e.g., §6.1). Choose $\lambda = (4, 1, \frac{3}{2}, \frac{1}{2})$. It is easy to verify that $\Phi_{[\lambda]}$ is a subsystem with simple roots $\{\varepsilon_3 - \varepsilon_4, \varepsilon_4, \frac{1}{2}(\varepsilon_1 - \varepsilon_2 - \varepsilon_3 - \varepsilon_4), \varepsilon_2\}$. Therefore $\Phi_{[\lambda]} \simeq C_4$ and λ is dominant. By Theorem 6.1, $L(\lambda)$ is nonintegral minimal with GKdim $L(\lambda) = |\Phi^+| - |\Phi_{[\lambda]}| = 24 - 16 = 8$. It is minimal with associated variety $\overline{Be_{\varepsilon_2 - \varepsilon_3}}$ in view of Table 1 and Theorem 4.2.

Now we recall some definitions and properties about closed subsystem and maximal subsystem. Most results can be found in [Kan01, §12].

Let Φ be an irreducible root system. We adopt the notation in [Hum78] and write $\Delta = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$, where roots and weights are realized as vectors of vector

spaces with orthonormal basis ε_i . In particular, the simple roots for $\Phi = A_n, B_n, C_n$ and D_n are chosen to be $\alpha_1 = \varepsilon_1 - \varepsilon_2, \, \alpha_2 = \varepsilon_2 - \varepsilon_3, \, \cdots, \, \alpha_{n-1} = \varepsilon_{n-1} - \varepsilon_n$ with one more root $\alpha_n = \varepsilon_n - \varepsilon_{n+1}$, ε_n , $2\varepsilon_n$ and $\varepsilon_{n-1} + \varepsilon_n$ respectively. The simple roots of E_8 are $\{\frac{1}{2}(\varepsilon_1+\varepsilon_8-(\varepsilon_2+\cdots+\varepsilon_7)), \varepsilon_1+\varepsilon_2, \varepsilon_2-\varepsilon_1, \varepsilon_3-\varepsilon_2, \varepsilon_4-\varepsilon_3, \varepsilon_5-\varepsilon_4, \varepsilon_6-\varepsilon_5, \varepsilon_7-\varepsilon_6\},\$ while the first 6 (resp. 7) simple roots generate E_6 (resp. E_7). The simple roots for F_4 and G_2 are $\{\varepsilon_2 - \varepsilon_3, \varepsilon_3 - \varepsilon_4, \varepsilon_4, \frac{1}{2}(\varepsilon_1 - \varepsilon_2 - \varepsilon_3 - \varepsilon_4)\}$ and $\{\varepsilon_1 - \varepsilon_2, -2\varepsilon_1 + \varepsilon_2 + \varepsilon_3\}$ respectively.

Denote $\Phi^{\vee} = \{\beta^{\vee} \mid \beta \in \Phi\}$. It is a root system, which is called the *dual system* of Φ . One always has $\Phi^{\vee\vee} = \Phi$. If Φ is simply-laced, then $\Phi^{\vee} = \Phi$. In the non simply-laced cases, we have $B_n^{\vee} = C_n$, $F_4^{\vee} \simeq F_4$ and $G_2^{\vee} \simeq G_2$, although the root lengths might be different.

Definition 6.3. We say a subsystem $\Phi' \subseteq \Phi$ is *closed* if and only if $\alpha + \beta \in \Phi'$ whenever $\alpha, \beta \in \Phi'$ and $\alpha + \beta \in \Phi$.

Example 6.4. Let $\Phi = B_2 = \{\pm(\varepsilon_1 + \varepsilon_2), \pm(\varepsilon_1 - \varepsilon_2), \pm\varepsilon_1, \pm\varepsilon_2\}$, then $\Phi' =$ $\{\pm \varepsilon_1, \pm \varepsilon_2\} \simeq A_1 \times A_1$ form a subsystem, which is not closed.

The following result is an immediate consequence of Definition 6.3.

Lemma 6.5. Let $\lambda \in \mathfrak{h}^*$. Then $\Phi_{[\lambda]}^{\vee}$ is a closed subsystem of Φ^{\vee} .

Let α_0 be the highest root of Φ and write

$$\alpha_0 = \sum_{i=1}^n h_i \alpha_i.$$

Theorem 6.6 (Borel-de Siebenthal). Let Φ be irreducible. Then the maximal proper closed subsystem of Φ (up to the action of W) are those with simple systems

(1) $\{\alpha_1, \alpha_2, \cdots, \hat{\alpha}_i, \cdots, \alpha_n\}$ with $h_i = 1$;

(2) $\{-\alpha_0, \alpha_1, \cdots, \hat{\alpha}_i, \cdots, \alpha_n\}$ with $h_i = p$ being prime;

where $\alpha_0 = \sum_{i=1}^n h_i \alpha_i$ is the highest root of Φ .

Proposition 6.7. Let Φ be irreducible and $\lambda \in \mathfrak{h}^*$ be nonintegral. Then $|\Phi_{[\lambda]}|$ is maximal when the following conditions are satisfied:

- (1) If $\Phi = A_n$, then $\Delta_{[\lambda]} \simeq \{\varepsilon_1 \varepsilon_2, \varepsilon_2 \varepsilon_3, \cdots, \varepsilon_{n-1} \varepsilon_n\};$
- (2) If $\Phi = B_n$, then $\Delta_{[\lambda]} \simeq \{-\varepsilon_1, \varepsilon_2 \varepsilon_3, \varepsilon_3 \varepsilon_4, \cdots, \varepsilon_{n-1} \varepsilon_n, \varepsilon_n\};$
- (3) If $\Phi = C_n$, then $\Delta_{[\lambda]} \simeq \{-(\varepsilon_1 + \varepsilon_2), \varepsilon_2 \varepsilon_3, \varepsilon_3 \varepsilon_4, \cdots, \varepsilon_{n-1} \varepsilon_n\};$
- (4) If $\Phi = D_n$, then $\Delta_{[\lambda]} \simeq \{\varepsilon_2 \varepsilon_3, \varepsilon_3 \varepsilon_4, \cdots, \varepsilon_{n-1} \varepsilon_n, \varepsilon_{n-1} + \varepsilon_n\};$
- (5) If $\Phi = E_6$, then $\Delta_{[\lambda]} \simeq \{\varepsilon_1 + \varepsilon_2, \varepsilon_2 \varepsilon_1, \varepsilon_3 \varepsilon_2, \cdots, \varepsilon_5 \varepsilon_4\};$
- (6) If $\Phi = E_7$, then $\Delta_{[\lambda]} \simeq \{\alpha_1, \varepsilon_1 + \varepsilon_2, \varepsilon_2 \varepsilon_1, \cdots, \varepsilon_5 \varepsilon_4\};$
- (7) If $\Phi = E_8$, then $\Delta_{[\lambda]} \simeq \{-(\varepsilon_7 + \varepsilon_6), \alpha_1, \varepsilon_1 + \varepsilon_2, \cdots, \varepsilon_6 \varepsilon_5\};$
- (8) If $\Phi = F_4$, then $\Delta_{[\lambda]} \simeq \{-\varepsilon_1, \varepsilon_3 \varepsilon_4, \varepsilon_4, \frac{1}{2}(\varepsilon_1 \varepsilon_2 \varepsilon_3 \varepsilon_4)\}$ (9) If $\Phi = G_2$, then $\Delta_{[\lambda]} \simeq \{\varepsilon_2 + \varepsilon_3, \varepsilon_1 \varepsilon_2\}$

The corresponding $\Phi_{[\lambda]}$ are given in Table 2.

Proof. The argument is easy. We only consider $\Phi = B_n$ with

$$\Delta = \{\alpha_1, \alpha_2, \cdots, \alpha_n\} = \{\varepsilon_1 - \varepsilon_2, \cdots, \varepsilon_{n-1} - \varepsilon_n, \varepsilon_n\}$$

as a demonstration. In this case $\Phi^{\vee} \simeq C_n$ with simple roots $\{\alpha_1, \cdots, \alpha_{n-1}, 2\alpha_n\}$. The highest root is $\alpha_0 = 2\varepsilon_1 = 2\alpha_1 + \cdots + 2\alpha_{n-1} + (2\alpha_n)$. By Lemma 6.5, $\Phi_{[\lambda]}^{\vee}$ is a proper closed subsystem of C_n . If $\Phi_{[\lambda]}^{\vee}$ is maximal, its simple system should isomorphic to $\{\alpha_1, \alpha_2, \cdots, \alpha_{n-1}\}$ or $\{-\alpha_0, \alpha_1, \cdots, \hat{\alpha}_i, \cdots, \alpha_{n-1}, 2\alpha_n\}$ for $1 \leq i < n$ by Theorem 6.6. It follows that $\Phi_{[\lambda]}^{\vee}$ is isomorphic to A_{n-1} or $C_i \times C_{n-i}$ for $1 \leq i < n$. Hence $|\Phi_{[\lambda]}|$ is maximal if and only if $\Phi_{[\lambda]} \simeq C_1^{\vee} \times C_{n-1}^{\vee} \simeq B_1 \times B_{n-1}$ and $\Delta_{[\lambda]} \simeq \{-\varepsilon_1, \varepsilon_2 - \varepsilon_3, \varepsilon_3 - \varepsilon_4, \cdots, \varepsilon_{n-1} - \varepsilon_n, \varepsilon_n\}.$

Proof of Theorem 6.1. By Proposition 3.1, one has $\operatorname{GKdim} L(\lambda) = |\Phi^+| - |\Phi^+|$ $\mathbf{a}_{[\lambda]}(w_{\lambda})$. Lemma 3.5 implies that $\mathbf{a}_{[\lambda]}(w_{\lambda})$ achieves its maximal value $|\Phi_{[\lambda]}^+|$ if and only if w_{λ} is longest in $W_{[\lambda]}$, which means λ is dominant regular. At last, we apply Proposition 6.7 and get Table 2. Combined with Table 1, we can easily find that nonintegral minimal module is minimal if Φ is of type A, B, C, F or G.

From the above arguments and the proof in Proposition 5.3, we have the following.

Corollary 6.8. Let $w_{[\lambda]}$ be the longest element in $W_{[\lambda]}$. Let $\nu \in \mathfrak{D}$ and $\lambda \in W\nu$. Let δ be the unique dominant weight in $W_{[\lambda]}\lambda$. If ν is singular or nonintegral, then the following conditions are equivalent:

- (1) $L(\lambda)$ is ν -minimal;
- (2) GKdim $L(\lambda) =$ GKdim $L(\delta) = |\Phi^+| \mathbf{a}_{[\lambda]}(w_{[\lambda]}w_J);$
- (3) $w_{\lambda} \sim_L w_{[\lambda]} w_J$ in $W_{[\lambda]}$.

Note that when λ is regular, we will have $J = \emptyset$ and $w_J = 1$.

7. Minimal highest weight modules of classical types

In order to verify whether a highest weight module is minimal, we apply Theorem 5.1 for types A, D, E and Theorem 6.1 for types A, B, C, F, G. These criteria are quite practical. In this section, we intend to investigate more explicit characterization of minimal highest weight modules based on these criteria in the case of classical Lie algebras.

We continue to follow the notation in [Hum78]. Note that for $\Phi = A_{n-1}, B_n, C_n$ and D_n , a weight $\lambda \in \mathfrak{h}^*$ will be identified with a sequence $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{C}^n$ such that $\lambda = \sum_{i=1}^n \lambda_i \varepsilon_i$.

Definition 7.1. Following Mathieu [Mat00], a sequence $x = (x_1, x_2, \cdots, x_m) \in \mathbb{C}^m$ is said to be *ordered* if all differences $x_i - x_{i+1}$ are positive integers.

7.1. **Type** A_n .

Theorem 7.2. Let $\mathfrak{g} = \mathfrak{sl}(n+1,\mathbb{C})$. The simple module $L(\lambda)$ is minimal if and only if the length of the longest ordered subsequence of λ is n. In this case, let t be the smallest integer such that $(\lambda_1, \dots, \hat{\lambda_t}, \dots, \lambda_{n+1})$ is ordered. Then

- (1) If λ is not integral, then $V(L(\lambda)) = \overline{Be_{\alpha_{t-1}}} \cup \overline{Be_{\alpha_t}}$, where $e_{\alpha_0} = e_{\alpha_{n+1}} = 0$; (2) If λ is integral, then $V(L(\lambda)) = \overline{Be_{\alpha_{t-1}}}$ when $\lambda_{t-1} \leq \lambda_t$ and $V(L(\lambda)) =$ $\overline{Be_{\alpha_t}}$ when $\lambda_t \leq \lambda_{t+1}$.

Proof. It suffices to consider the first assertion since results about associated varieties are evident consequences of Theorem 4.2.

If λ is nonintegral, the first statement follows from Theorem 6.1. Indeed, the module $L(\lambda)$ is minimal if and only if λ is regular dominant and $\Phi_{\lambda} \simeq A_{n-1}$, which is equivalent to the required ordered subsequence.

Now suppose that λ is integral. Set $w = w_0 w_{\lambda}^{-1}$. It is the longest element such that $w\lambda$ is dominant. In this case, Proposition 5.4 shows that $L(\lambda)$ is minimal if and only if $w \in \mathcal{C}$. Recall that $\alpha_j = \varepsilon_j - \varepsilon_{j+1}$ for $1 \leq j \leq n$.

If $(\lambda_1, \dots, \hat{\lambda_t}, \dots, \lambda_{n+1})$ is ordered and λ is not, then $(\lambda_{t-1} - \lambda_t)(\lambda_t - \lambda_{t+1}) \leq 0$. If $\lambda_t \geq \lambda_{t-1}$, there exists $m \geq 1$ such that $\lambda_{t-m-1} > \lambda_t \geq \lambda_{t-m}$ (set $\lambda_0 = +\infty$). Evidently

$$(\lambda_1, \cdots, \lambda_{t-m-1}, \lambda_t, \lambda_{t-m}, \cdots, \lambda_{t-1}, \lambda_{t+1}, \cdots, \lambda_{n+1}) = s_{\alpha_{t-m}} \cdots s_{\alpha_{t-1}} \lambda_{t+1}$$

is dominant and $w = s_{\alpha_{t-m}} \cdots s_{\alpha_{t-1}} \in \mathcal{C}$. If $\lambda_t \leq \lambda_{t+1}$, the argument is similar.

Conversely, if $L(\lambda)$ is minimal, by Proposition 5.4, we can assume that $w \in \mathcal{C}$. Since $w \neq 1$ has the unique reduced expression, there exist $i, m \geq 1$ such that

$$w = s_{\alpha_{i+m-1}} \cdots s_{\alpha_{i+1}} s_{\alpha_i} \text{ or } s_{\alpha_{i-m+1}} \cdots s_{\alpha_{i-1}} s_{\alpha_i},$$

Since the arguments are similar, we only consider the former case, for which

$$w\lambda = (\lambda_1, \cdots, \lambda_{i-1}, \lambda_{i+1}, \cdots, \lambda_{i+m}, \lambda_i, \lambda_{i+m+1}, \cdots, \lambda_{n+1})$$

is dominant and thus $\lambda_{i-1} \geq \lambda_{i+1}$. It follows from Theorem 5.1(3) that $\langle \lambda, \alpha_i^{\vee} \rangle =$ $\lambda_j - \lambda_{j+1} \in \mathbb{Z}_{>0}$ for $j \neq i$. If $\lambda_{i-1} > \lambda_{i+1}$, then λ is ordered by removing λ_i . If $\lambda_{i-1} = \lambda_{i+1}$, then $\langle s_{\alpha_i} \lambda, \alpha_{i-1}^{\vee} \rangle = 0$. Theorem 5.1(3) yields $\gamma_m = \alpha_{i+m-1} = \alpha_{i-1}$, a contradiction.

Remark 7.3. We set $s_i = s_{\alpha_i}$ for $1 \le i \le n$ and $s_{n+1} = 1$. Let $\lambda^0 = (\frac{n}{2} + \frac{k}{n+1}, \frac{n-2}{2} + \frac{k}{n+1}, \cdots, -\frac{n-2}{2} + \frac{k}{n+1}, -\frac{n}{2} - \frac{kn}{n+1})$ and $\lambda^i = s_{i+1}s_{i+2}\cdots s_{n+1}\lambda^0$ for $1 \le i \le n$. Let $\mu^0 = (\frac{n}{2} - \frac{1}{n+1}, \frac{n-2}{2} - \frac{1}{n+1}, \cdots, -\frac{n-2}{2} - \frac{1}{n+1}, -\frac{n-2}{2} - \frac{1}{n+1})$ and $\mu^i = s_i s_{i+1} \cdots s_n \mu^0$ for $1 \le i \le n$. The above Theorem 7.2 implies that $L(\lambda^i)$ $(1 \leq i \leq n)$ and $L(\mu^i)$ $(1 \leq i \leq n)$ are both minimal and the integer t = i + 1for $L(\lambda^i)$ and t = i for $L(\mu^i)$. This recovers the result in [Jos98, §4.5]. Suppose $L(w\mu)$ is minimal, from Joseph [Jos98], we also have that Ann $(L(w\mu))$ is completely prime if and only if the infinitesimal character $\mu = \lambda^0$ with $k + 1 \in \mathbb{C} \setminus \mathbb{N}$ and $w \in \{s_{i+1}s_{i+2}\cdots s_{n+1} \mid 1 \le i \le n\}$ or $\mu = \mu^0$ and $w \in \{s_is_{i+1}\cdots s_n \mid 1 \le i \le n\}$ (equivalently $w \sim_L s_n$).

7.2. **Type** B_n .

Theorem 7.4. Let $\mathfrak{g} = \mathfrak{so}(2n+1,\mathbb{C})$. The simple module $L(\lambda)$ is minimal if and only if the following conditions are satisfied:

- (1) There is $1 \leq t \leq n$ such that $(\lambda_1, \dots, \lambda_n)$ is ordered after λ_t is removed; (2) $2\lambda_i \in \mathbb{Z}_{>0}$ for $1 \leq i \leq n$, while $2\lambda_t$ and $2\lambda_i$ $(i \neq t)$ have different parity.

In this case, $V(L(\lambda)) = \overline{Be_{\alpha_{t-1}}} \cup \overline{Be_{\alpha_t}}$ for t < n and $V(L(\lambda)) = \overline{Be_{\alpha_{n-1}}}$ for t = n.

Proof. By Theorem 5.1 and 6.1, $L(\lambda)$ is minimal if and only if λ is nonintegral dominant regular and $\Phi_{[\lambda]} \simeq B_1 \times B_{n-1}$. This yields (1) and (2). The last assertion follows from Theorem 4.2.

Remark 7.5. We set $s_i = s_{\alpha_i}$ for $1 \le i \le n$ and $s_0 = 1$. Let $\lambda^1 = (1, n - \frac{3}{2}, n - \frac{5}{2}, \dots, \frac{1}{2})$ and $\lambda^i = (n - \frac{3}{2}, \dots, n - \frac{2i-1}{2}, 1, n - \frac{2i+1}{2}, \dots, \frac{1}{2})$ for $2 \le i \le n - 1$. The above Theorem 7.4 implies that $L(\lambda^i)$ are minimal and $V(L(\lambda^1)) = \overline{Be_{\alpha_1}}$, $V(L(\lambda^n)) = \overline{Be_{\alpha_{n-1}}}$ and $V(L(\lambda^i)) = \overline{Be_{\alpha_{i-1}}} \cup \overline{Be_{\alpha_i}}$ for $2 \le i \le n-1$. This recovers the result of Joseph [Jos98, §4.8]. Suppose $L(w\mu)$ is minimal, from Joseph [Jos98], we also have that $Ann(L(w\mu))$ is completely prime if and only if the infinitesimal character $\mu = \lambda^1$ and $w \in D_{\lambda} = \{w \in W \mid w\alpha > 0 \text{ for any } \alpha > 0 \text{ and } \langle \lambda, \alpha^{\vee} \rangle > 0$ $0\} = \{s_{i-1}s_{i-2}\cdots s_0 \mid 1 \le i \le n\}.$

7.3. **Type** C_n .

Theorem 7.6. Let $\mathfrak{g} = \mathfrak{sp}(n, \mathbb{C})$. The simple module $L(\lambda)$ is minimal if and only if the following conditions are satisfied:

(1) λ is not integral and $\lambda_i \in \frac{1}{2} + \mathbb{Z}$ for all $1 \leq i \leq n$;

(2) $(\lambda_1, \cdots, \lambda_{n-1}, |\lambda_n|)$ is ordered.

In this case, $V(L(\lambda)) = \overline{Be_{\alpha_n}}$.

Proof. This follows easily from Theorem 6.1 and 4.2.

Remark 7.7. When $\mathfrak{g} = \mathfrak{sp}(n, \mathbb{C})$, Britten-Lemire [BL99] described those highest weight modules with uniformly bounded weight spaces, while Sun [Sun08] classified the minimal lowest (equivalently, highest) weight Harish-Chandra (\mathfrak{g}, K)-modules. As a consequence of their work, we know that a highest weight module of type C has uniformly bounded weight spaces if and only if it is minimal.

Let $\lambda^1 = (n + \frac{1}{2}, n - \frac{1}{2}, \dots, \frac{3}{2})$ and $\lambda^2 = (n + \frac{1}{2}, n - \frac{1}{2}, \dots, \frac{5}{2}, -\frac{3}{2})$. The above Theorem 7.6 implies that $L(\lambda^i)$ are minimal and $V(L(\lambda^i)) = \overline{Be_{\alpha_n}}$. This recovers the result of Joseph [Jos98, §4.14]. Suppose $L(w\mu)$ is minimal, from Joseph [Jos98], we also have that $\operatorname{Ann}(L(w\mu))$ is completely prime if and only if the infinitesimal character $\mu = \lambda^1$ and w = 1 or s_{α_n} .

7.4. **Type** D_n .

Theorem 7.8. Let $\Phi = D_n$ and $\lambda \in \mathfrak{h}^*$. Then $L(\lambda)$ is minimal if and only if the following conditions are satisfied:

- (1) λ is integral and contains at most one 0-entry;
- (2) There is the unique $1 \le k \le n$ such that $\langle \lambda, \alpha_k^{\vee} \rangle \le 0$;
- (3) If $1 \leq k \leq n-2$, then $(\lambda_1, \dots, \hat{\lambda}_l, \dots, \lambda_{n-1}, |\lambda_n|)$ is ordered for l = k or k+1 and $\lambda_l > -|\lambda_n|$;
- (4) If k = n 1, then $\lambda_{n-2} > |\lambda_{n-1}|$ or $\lambda_{n-3} > -\lambda_n > -\lambda_{n-1} > |\lambda_{n-2}|$;
- (5) If k = n, then $\lambda_{n-2} > |\lambda_{n-1}|$ or $\lambda_{n-3} > \lambda_n > -\lambda_{n-1} > |\lambda_{n-2}|$.

In this case, $V(L(\lambda)) = \overline{Be_{\alpha_k}}$.

Proof. Set $w = w_0 w_{\lambda}^{-1}$. Then w is the longest element such that $w\lambda$ is dominant. By Theorem 5.1, 6.1 and Proposition 5.4, $L(\lambda)$ is minimal if and only if λ is

integral and $w \in \mathcal{C}$. Let $m \ge 1$ be the length of w. We list all the elements in \mathcal{C} .

- (a) $w = s_{\alpha_{k-m+1}} \cdots s_{\alpha_k}$ with $k \le n-2$;
- (b) $w = s_{\alpha_{k+m-1}} \cdots s_{\alpha_k}$ with $k \le n-2$ and m < n-k;
- (c) $w = s_{\alpha_{n-1}} s_{\alpha_{n-2}} \cdots s_{\alpha_k}$ with $k \le n-2$ and m = n-k;
- (d) $w = s_{\alpha_n} s_{\alpha_{n-2}} \cdots s_{\alpha_k}$ with $k \le n-2$ and m = n-k;
- (e) $w = s_{\alpha_{n-m}} \cdots s_{\alpha_{n-2}} s_{\alpha_{n-1}}$ with k = n-1;
- (f) $w = s_{\alpha_n} s_{\alpha_{n-2}} s_{\alpha_{n-1}}$ with k = n 1;
- (g) $w = s_{\alpha_{n-m}} \cdots s_{\alpha_{n-2}} s_{\alpha_n}$ with k = n;
- (h) $w = s_{\alpha_{n-1}} s_{\alpha_{n-2}} s_{\alpha_n}$ with k = n;

This makes a case-by-case proof possible. For example, assume that (1), (2) and (3) are true with l = k in (3). Then (2) and (3) yield $\lambda_{k+1} \ge \lambda_k > -|\lambda_n|$ and

$$\lambda_1 > \cdots > \lambda_{k-1} > \lambda_{k+1} > \cdots > \lambda_{n-1} > |\lambda_n|.$$

If $\lambda_k \leq |\lambda_n|$ with $\lambda_n \leq 0$, then

(7.1)
$$s_{\alpha_n} s_{\alpha_{n-2}} \cdots s_{\alpha_k} \lambda = (\lambda_1, \cdots, \lambda_{k-1}, \lambda_{k+1}, \cdots, \lambda_{n-1}, -\lambda_n, -\lambda_k)$$

16

is dominant. It is evident that $w = s_{\alpha_n} s_{\alpha_{n-2}} \cdots s_{\alpha_k}$. This gives the case (d), while $\lambda_k \leq |\lambda_n|$ with $\lambda_n > 0$ corresponds to (c). Similarly, if $\lambda_{n-1} \geq \lambda_k > |\lambda_n|$ or $\lambda_{k+m} \geq \lambda_k > \lambda_{k+m+1}$ for some m < n-k-1, we arrive at (b).

Conversely, if $L(\lambda)$ is minimal and (d) holds, then (1), (2) are true and $w\lambda$ is dominant, in view of Theorem 5.1. With (7.1), we get $\lambda_{k-1} > \lambda_{k+1}$, $-\lambda_n > -\lambda_k$ and $-\lambda_n \ge \lambda_k$ (by Theorem 5.1(3), the equality holds only when λ is semiregular). Since $\langle \lambda, \alpha_i \rangle > 0$ for $i \ne k$, the sequence $(\lambda_1, \dots, \hat{\lambda}_l, \dots, \lambda_{n-1}, |\lambda_n|)$ is ordered for l = k and $\lambda_k > \lambda_n = -|\lambda_n|$, which implies (3).

The arguments for other cases which we omit are similar.

Remark 7.9. We set $s_i = s_{\alpha_i}$ for $1 \leq i \leq n$ and $s_0 = 1$. Let $\lambda^1 = (1, n - 2, n - 3, \dots, 2, 1, 0)$ and $\lambda^i = s_{i-1}s_{i-2}\cdots s_0\lambda^1 = (n-2, n-3, \dots, n-i, 1, n-i-1, \dots, 1, 0)$ for $2 \leq i \leq n-2$, $\lambda^{n-1} = s_{n-1}\lambda^{n-2} = (n-2, n-3, \dots, 3, 2, 1, 0, 1)$ and $\lambda^n = s_n\lambda^{n-2} = (n-2, n-3, \dots, 3, 2, 1, 0, 1)$ and $\lambda^n = s_n\lambda^{n-2} = (n-2, n-3, \dots, 3, 2, 1, 0, -1)$. The above Theorem 7.8 implies that $L(\lambda^i)$ is minimal and the integer k = i for $L(\lambda^i)$. This recovers the result of Joseph [Jos98, Table 3]. Suppose $L(w\mu)$ is minimal, from Joseph [Jos98], we also have that $\operatorname{Ann}(L(w\mu))$ is completely prime if and only if the infinitesimal character $\mu = \lambda^1$ and $w \in \{s_{i-1}s_{i-2}\cdots s_0 \mid 1 \leq i \leq n-2\} \cup \{s_{n-1}s_{n-3}\cdots s_0, s_ns_{n-3}\cdots s_0\}$.

Acknowledgments. We would like to thank Toshiyuki Tanisaki for very helpful conversations and many useful comments for an earlier version of the manuscript. We also would like to thank Binyong Sun for inspiring us this problem. Z.Bai was supported in part by NSFC Grant No. 12171344 and the National Key R & D Program of China (No. 2018YFA0701700 and No. 2018YFA0701701). W. Xiao was supported in part by NSFC Grant No. 11701381 and Guangdong Natural Science Foundation (Grant No. 2017A030310138). X. Xie was supported in part by NSFC Grant No. 11801031 and No. 12171030.

References

- [BB82] W. Borho and J.-L. Brylinski. Differential operators on homogeneous spaces. I. Irreducibility of the associated variety for annihilators of induced modules. *Invent. Math.*, 69(3):437–476, 1982. 3, 10
- [BB85] W. Borho and J.-L. Brylinski. Differential operators on homogeneous spaces. III. Characteristic varieties of Harish-Chandra modules and of primitive ideals. *Invent. Math.*, 80(1):1–68, 1985. 1, 5, 10
- [BBL97] G. Benkart, D. Britten, and F. Lemire. Modules with bounded weight multiplicities for simple Lie algebras. Math. Z., 225(2):333–353, 1997. 2
- [BBM89] W. Borho, J.-L. Brylinski, and R. MacPherson. Nilpotent orbits, primitive ideals, and characteristic classes, volume 78 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 1989. A geometric perspective in ring theory. 10
- [Ber71] I. N. Bernštein. Modules over a ring of differential operators. An investigation of the fundamental solutions of equations with constant coefficients. *Funkcional. Anal. i Priložen.*, 5(2):1–16, 1971. 1
- [BG80] J. N. Bernstein and S. I. Gel'fand. Tensor products of finite- and infinite-dimensional representations of semisimple Lie algebras. *Compositio Math.*, 41(2):245–285, 1980. 2
- [BL99] D. Britten and F. Lemire. On modules of bounded multiplicities for the symplectic algebras. Trans. Amer. Math. Soc., 351(8):3413–3431, 1999. 16
- [BV83] D. Barbasch and D.A. Vogan. Primitive ideals and orbital integrals in complex exceptional groups. J. Algebra, 80(2):350–382, 1983. 8, 10
- [BX19] Z. Bai and X. Xie. Gelfand-Kirillov dimensions of highest weight Harish-Chandra modules for SU(p,q). Int. Math. Res. Not. IMRN, pages 4392–4418, 2019. 2, 7, 10

- [BXX] Z. Bai, W. Xiao, and X. Xie. Gelfand-kirillov dimensions and associated varieties of highest weight modules. arXiv:2005.11536v2. 2, 4, 6, 7, 11
- [CM93] D. H. Collingwood and W. M. McGovern. Nilpotent orbits in semisimple Lie algebras. Van Nostrand Reinhold Mathematics Series. Van Nostrand Reinhold Co., New York, 1993. 5, 6, 7, 10
- [GS04] W. T. Gan and G. Savin. Uniqueness of Joseph ideal. Math. Res. Lett., 11(5-6):589–597, 2004. 2
- [GS05] W. T. Gan and G. Savin. On minimal representations definitions and properties. Represent. Theory, 9:46–93, 2005. 2, 6
- [HKM14] J. Hilgert, T. Kobayashi, and J. Möllers. Minimal representations via Bessel operators. J. Math. Soc. Japan, 66(2):349–414, 2014. 2
- [Hum78] J. E. Humphreys. Introduction to Lie algebras and representation theory, volume 9 of Graduate Texts in Mathematics. Springer-Verlag, New York-Berlin, 1978. Second printing, revised. 6, 11, 12, 14
- [Hum08] J. E. Humphreys. Representations of semisimple Lie algebras in the BGG category \$\mathcal{O}\$, volume 94 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2008. 4, 7
- [Hum16] J.E. Humphreys. Notes on the minimal special nilpotent orbit. Available at http://people.math.umass.edu/jeh/pub/minimal.pdf, 2016. 7
- [Jos76] A. Joseph. The minimal orbit in a simple Lie algebra and its associated maximal ideal. Ann. Sci. École Norm. Sup. (4), 9(1):1–29, 1976. 2
- [Jos78] A. Joseph. Gelfand-Kirillov dimension for the annihilators of simple quotients of Verma modules. J. London Math. Soc. (2), 18(1):50–60, 1978. 5
- [Jos84] A. Joseph. On the variety of a highest weight module. J. Algebra, 88(1):238–278, 1984. 5, 9, 10
- [Jos85] A. Joseph. On the associated variety of a primitive ideal. J. Algebra, 93(2):509–523, 1985. 1, 3, 5, 10
- [Jos98] A. Joseph. Orbital varietes of the minimal orbit. Ann. Sci. École Norm. Sup. (4), 31(1):17-45, 1998. 2, 6, 15, 16, 17
- [Kan01] R. Kane. Reflection groups and invariant theory, volume 5 of CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC. Springer-Verlag, New York, 2001. 12
- [KL79] D. Kazhdan and G. Lusztig. Representations of Coxeter groups and Hecke algebras. Invent. Math., 53(2):165–184, 1979. 3, 8, 10
- [Kob11] T. Kobayashi. Algebraic analysis of minimal representations. Publ. Res. Inst. Math. Sci., 47(2):585–611, 2011. 2
- [Li00] J.-S. Li. Minimal representations & reductive dual pairs. In Representation theory of Lie groups (Park City, UT, 1998), volume 8 of IAS/Park City Math. Ser., pages 293–340. Amer. Math. Soc., Providence, RI, 2000. 2
- [Lus79] G. Lusztig. A class of irreducible representations of a Weyl group. Nederl. Akad. Wetensch. Indag. Math., 41(3):323–335, 1979. 3, 10
- [Lus83] G. Lusztig. Some examples of square integrable representations of semisimple p-adic groups. Trans. Amer. Math. Soc., 277(2):623–653, 1983. 8
- [Lus84] G. Lusztig. Characters of reductive groups over a finite field, volume 107 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1984. 6
- [Lus03] G. Lusztig. Hecke algebras with unequal parameters, volume 18 of CRM Monograph Series. American Mathematical Society, Providence, RI, 2003. 7, 8, 11
- [Mat00] O. Mathieu. Classification of irreducible weight modules. Ann. Inst. Fourier (Grenoble), 50(2):537–592, 2000. 2, 14
- [McG00] W. M. McGovern. A triangularity result for associated varieties of highest weight modules. Comm. Algebra, 28(4):1835–1843, 2000. 10
- [Mel93] A. Melnikov. Irreducibility of the associated varieties of simple highest weight modules in sl(n). C. R. Acad. Sci. Paris Sér. I Math., 316(1):53–57, 1993. 1
- [Men11] G. Meng. Euclidean Jordan algebras, hidden actions, and J-Kepler problems. J. Math. Phys., 52(11):112104, 35, 2011. 2
- [NOT01] K. Nishiyama, H. Ochiai, and K. Taniguchi. Bernstein degree and associated cycles of Harish-Chandra modules—Hermitian symmetric case. Number 273, pages 13–80. 2001. Nilpotent orbits, associated cycles and Whittaker models for highest weight representations. 5

- [Sun08] B. Sun. Lowest weight modules of $\widetilde{\text{Sp}_{2n}(\mathbb{R})}$ of minimal Gelfand-Kirillov dimension. J. Algebra, 319(7):3062–3074, 2008. 2, 16
- [Tam] T. Tamori. Classification of irreducible (g, t)-modules associated to the ideals of minimal nilpotent orbits for simple lie groups of type a. arXiv:2109.03432v1. 2
- [Tan88] T. Tanisaki. Characteristic varieties of highest weight modules and primitive quotients. In Representations of Lie groups, Kyoto, Hiroshima, 1986, volume 14 of Adv. Stud. Pure Math., pages 1–30. Academic Press, Boston, MA, 1988. 1, 3, 5, 10
- [Vog78] D. A. Vogan. Gelfand-Kirillov dimension for Harish-Chandra modules. Invent. Math., 48(1):75–98, 1978. 5
- [Vog81] D. A. Vogan. Singular unitary representations. In Noncommutative harmonic analysis and Lie groups (Marseille, 1980), volume 880 of Lecture Notes in Math., pages 506–535. Springer, Berlin-New York, 1981. 6
- [Wan99] W. Q. Wang. Dimension of a minimal nilpotent orbit. Proc. Amer. Math. Soc., 127(3):935–936, 1999. 1, 6
- [Wil15] G. Williamson. A reducible characteristic variety in type A. In Representations of reductive groups, volume 312 of Progr. Math., pages 517–532. Birkhäuser/Springer, Cham, 2015. 1
- [XZ22] Wei Xiao and Ailin Zhang. Jantzen coefficients and simplicity of parabolic Verma modules. J. Algebra, 611:24–64, 2022. 7

(Bai) School of Mathematical Sciences, Soochow University, Suzhou 215006, P. R. China

Email address: zqbai@suda.edu.cn

(Ma) DEPARTMENT OF MATHEMATICS, SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN 361005, P. R. CHINA

Email address: hoxide@gmail.com

(Xiao) College of Mathematics and statistics, Shenzhen Key Laboratory of Advanced Machine Learning and Applications, Shenzhen University, Shenzhen 518060, Guangdong, P. R. China

Email address: xiaow@szu.edu.cn

(Xie) School of Mathematics and Statistics, Beijing Institute of Technology, Beijing 100081, P. R. China

Email address: xieg7@163.com