

INTERTWINING OPERATOR FOR $Sp(4, \mathbb{R})$ AND ORTHOGONAL POLYNOMIALS

ZHUOHUI ZHANG

ABSTRACT. Following the same method of [BM17] and [Zha19], we calculate the $(\mathfrak{g}, K)$ module structure for the principal series representation of $Sp(4, \mathbb{R})$. Furthermore, we introduced a hypergeometric generating function together with an inverse Mellin transform technique as an improvement to the method to calculate the intertwining operators. We have shown that the matrix entries of the simple intertwining operators for $Sp(4, \mathbb{R})$ -principal series are Hahn polynomials, and the matrix entries of the long intertwining operator can be expressed as the constant term of the Laurent expansion of some hypergeometric function.

1. AN INTRODUCTION TO THE RESULTS

The intertwining operators for principal series representation was defined in [KS80] and discussed in [VW90]. [KZ76] and [Mui09] applied intertwining operators to study the decomposition of certain classes of principal series representations. In this paper, we have shown the following new result for long intertwining operator for the minimal principal series of $Sp(4, \mathbb{R})$:

Theorem 1.1. *If the induction parameter δ of $Sp(4, \mathbb{R})$ principal series satisfies $\delta \in \{(0, 0), (1, 1)\}$ and $n - m_1 \equiv \delta_1$ and $n + m_2 \equiv \delta_2 \pmod{2}$ for $i \in \{1, 2\}$, the matrix entries $[A(\lambda)]_{m_1, m_2}^{j, n}$ for the long intertwining operator is the constant Laurent series coefficient of*

$$\begin{aligned}
& [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) = \\
& \frac{(-1)^n ((2j)!)^2}{\left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{j+n-\epsilon_\delta^{j,n}}{2}\right)} \left(\frac{\lambda_1+1}{2}\right)^{\left(-\frac{j+n-\epsilon_\delta^{j,n}}{2}\right)} \left(\frac{\lambda_1-\lambda_2+1}{2}\right)^{(j)} \left(\frac{\lambda_1+\lambda_2+1}{2}\right)^{(j)}} \times \\
& \frac{1}{c_{m_1}^j c_{m_2}^j} \frac{\left(\frac{\lambda_1-\lambda_2}{2}\right)^{\left(\frac{j+m_1-\epsilon_\delta^{j,n}}{2}\right)} \left(\frac{\lambda_1+\lambda_2}{2}\right)^{\left(\frac{-j-m_2+\epsilon_\delta^{j,n}}{2}\right)} \Gamma\left(\frac{1-\epsilon_\delta^{j,n}+j-m_1}{2}\right)}{\left(\frac{\lambda_2+1}{2}\right)^{\frac{m_2-n}{2}} \left(\frac{\lambda_2+1}{2}\right)^{-\frac{m_2-n}{2}} \Gamma\left(\frac{1-\epsilon_\delta^{j,n}+j-m_2}{2}\right)} \times \\
& (1-t_1)^{\frac{-1+\epsilon_\delta^{j,n}-j+m_1}{2}} (1-t_2)^{\frac{-1-\epsilon_\delta^{j,n}+j-m_2}{2}} t_2^{\epsilon_\delta^{j,n}-2j} t_1^{-\epsilon_\delta^{j,n}} \times \\
& {}_2F_1\left(-j+m_1, \frac{\lambda_1-\lambda_2-2j-1}{2}; t_1\right) {}_2F_1\left(-j-m_2, \frac{\lambda_1+\lambda_2-2j-1}{2}; t_2\right) \times \\
& (1.1) \quad {}_5F_4\left(\begin{matrix} -j+\epsilon_\delta^{j,n}, 1, \frac{-j+m_2+1+\epsilon_\delta^{j,n}}{2}, \frac{-j-n-\lambda_1+1+\epsilon_\delta^{j,n}}{2}, \frac{-j-m_2+\lambda_1+\lambda_2+\epsilon_\delta^{j,n}}{2} \\ -j+\epsilon_\delta^{j,n}, \frac{-j+m_1+1+\epsilon_\delta^{j,n}}{2}, \frac{-j-n+\lambda_1+1+\epsilon_\delta^{j,n}}{2}, \frac{-j-m_1-\lambda_1+\lambda_2+\epsilon_\delta^{j,n}}{2}+1 \end{matrix}; \frac{t_2^2(1-t_1)}{t_1^2(1-t_2)}\right).
\end{aligned}$$

where $\epsilon_\delta^{j,n} = \begin{cases} 0 & j-n \equiv \delta_i \pmod{2} \\ 1 & j-n \not\equiv \delta_i \pmod{2} \end{cases}$.

The notations in the theorem above will be specified in the following sections.

This paper is the second part of my Ph.D thesis. I would like to thank my doctoral advisor Stephen D. Miller, Siddhartha Sahi, James Lepowsky and Doron Zeilberger for instructions and inspiring ideas. I also would like to thank Oleg Marichev from Wolfram Research for his support and generous sharing of his symbolic integration package.

2. THE LIE GROUP $Sp(4, \mathbb{R})$

2.1. The Structure of the Group $Sp(4, \mathbb{R})$. The real symplectic group $G = Sp(4, \mathbb{R})$ is the subgroup of $SL(4, \mathbb{R})$ consisting of all elements $g \in SL(4, \mathbb{R})$ such that $g^t J g = J$, where

$$J = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}.$$

The real Lie algebra of the symplectic group is

$$\mathfrak{g} = \mathfrak{sp}(4, \mathbb{R}) = \{X \in \mathfrak{sl}(4, \mathbb{R}) | X^t J + J X = 0\}.$$

A root space decomposition for the complexified Lie algebra $\mathfrak{g}_{\mathbb{C}}$ and its Chevalley basis are determined by the following choice of data:

- (1) A Cartan subalgebra $\mathfrak{h}_{\mathbb{C}}$ generated by $H_1 = E_{1,1} - E_{3,3}, H_2 = E_{2,2} - E_{4,4}$;
- (2) The simple roots α_1 and α_2 sending $t_1 H_1 + t_2 H_2$ to

$$\alpha_1(t_1 H_1 + t_2 H_2) = t_1 - t_2, \quad \alpha_2(t_1 H_1 + t_2 H_2) = 2t_2$$

- (3) The simple coroots $\check{\alpha}_1$ and $\check{\alpha}_2$ sending $t_1 H_1 + t_2 H_2$ to

$$\check{\alpha}_1(t_1 H_1 + t_2 H_2) = t_1 - t_2, \quad \check{\alpha}_2(t_1 H_1 + t_2 H_2) = t_2$$

- (4) The fundamental weights ϖ_1 and ϖ_2 in $\mathfrak{h}_{\mathbb{C}}^*$ satisfying $\langle \varpi_i, \check{\alpha}_j \rangle = \delta_{ij}$;
- (5) The set of positive roots $\Delta^+(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\}$;
- (6) $\rho_{\mathbb{C}} = \frac{1}{2} \sum_{\alpha \in \Delta^+(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})} \alpha = \frac{4\alpha_1 + 3\alpha_2}{2} = \varpi_1 + \varpi_2$;
- (7) A basis for positive root spaces $\mathfrak{g}_{\alpha}$ for $\alpha \in \Delta^+(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$

$$\begin{aligned} X_{\alpha_1} &= E_{1,2} - E_{4,3} & X_{\alpha_2} &= E_{2,4} \\ X_{\alpha_1 + \alpha_2} &= E_{2,3} + E_{1,4} & X_{2\alpha_1 + \alpha_2} &= E_{1,3} \end{aligned}$$

with the corresponding negative root vector given by $X_{-\alpha} = X_{\alpha}^t$;

We will also represent an element $\lambda \in \mathfrak{h}_{\mathbb{C}}^*$ by a pair of complex numbers (λ_1, λ_2) where $\lambda_i = \lambda(H_i)$. In this coordinate, $\rho_{\mathbb{C}}$ is represented by $(2, 1)$.

On the real Lie group $Sp(4, \mathbb{R})$, there is a Cartan involution $\theta(g) = (g^t)^{-1}$ which determines a subgroup of fixed points $K = G^{\theta}$. The Lie algebra $\mathfrak{k} \subset \mathfrak{g}$ of the maximal compact subgroup $K \subset G$ is:

$$\mathfrak{k} = \left\{ \begin{pmatrix} A & B \\ -B & A \end{pmatrix} \mid A \text{ antisymmetric}, B \in \mathfrak{sym}(2) \right\}$$

The map $\begin{pmatrix} A & B \\ -B & A \end{pmatrix} \mapsto A + iB$ from $\mathfrak{k}$ to 2×2 matrices identifies the Lie algebra $\mathfrak{k}$ with $\mathfrak{u}(2)$. The 4-dimensional Lie algebra $\mathfrak{k}$ have generators U_0, U_1, U_2, U_3 corresponding

to the 4 infinitesimal generators $\gamma_0, \gamma_1, \gamma_2, \gamma_3$ of $\mathfrak{u}(2)$:

$$U_0 = \frac{1}{2} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}, U_1 = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$$

$$U_2 = \frac{1}{2} \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}, U_3 = \frac{1}{2} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

They satisfy a commutation relation:

$$[U_0, U_i] = 0, [U_i, U_j] = -\epsilon_{ijk} U_k$$

where the *Levi-Civita symbol* ϵ_{ijk} takes value 1 if (ijk) is an even permutation of (123) , -1 if (ijk) is an odd permutation of (123) , and 0 if two or more elements in $\{i, j, k\}$ are equal.

2.2. The Cartan Subgroups of $Sp(4, \mathbb{R})$.

2.2.1. The Maximally Compact Cartan Subalgebra. A Cartan subalgebra of $K \cong U(2)$ is $\mathfrak{k} = \mathbb{R}(U_0 + U_3) \oplus \mathbb{R}(U_0 - U_3)$. The basis vectors $U_0 \pm U_3$ are deliberately chosen so that they can be related to H_{α_1} and H_{α_2} by Cayley transforms. Let the simple roots β_1, β_2 act on an element $t_1(U_0 + U_3) + t_2(U_0 - U_3) \in \mathfrak{k}_{\mathbb{C}}$ by

$$\beta_1(t_1(U_0 + U_3) + t_2(U_0 - U_3)) = i(t_1 - t_2),$$

$$\beta_2(t_1(U_0 + U_3) + t_2(U_0 - U_3)) = 2it_2.$$

In the Vogan diagram (see [Kna16, Section VI.8]), the shorter root is named to be the compact root. In this situation, all roots are imaginary:

$$\begin{array}{c} \beta_1 \quad \beta_2 \\ \circ \longleftarrow \bullet \end{array}$$

We can decompose the set of positive roots $\Delta^+(\mathfrak{g}_{\mathbb{C}}, \mathfrak{k}_{\mathbb{C}})$ into the union of the set of compact roots and noncompact roots:


$$\Delta_c^+(\mathfrak{g}_{\mathbb{C}}, \mathfrak{k}_{\mathbb{C}}) = \{\beta_1\}$$

$$\Delta_{nc}^+(\mathfrak{g}_{\mathbb{C}}, \mathfrak{k}_{\mathbb{C}}) = \{\beta_2, \beta_1 + \beta_2, 2\beta_1 + \beta_2\};$$

The compact root vectors are

$$v_{\pm\beta_1} = \frac{1}{i}(U_1 \pm iU_2).$$

These roots are displayed in the following picture, where the gray color stands for noncompact roots, and the light gray color stands for compact roots:



The subspace $\mathfrak{t}_{\mathbb{C}} \oplus \mathfrak{g}_{\beta_1} \oplus \mathfrak{g}_{-\beta_1}$ of $\mathfrak{g}_{\mathbb{C}}$ generated by the vectors $\{v_{\beta_1}, v_{-\beta_1}, U_0 + U_3, U_0 - U_3\}$ is isomorphic to $\mathfrak{gl}(2, \mathbb{C})$. The embedding of this Lie subalgebra into $\mathfrak{g}_{\mathbb{C}}$ corresponds to the embedding of $U(2)$ into G as K .

The -1 eigenspace $\mathfrak{p}_{\mathbb{C}}$ for θ is generated by the following noncompact root vectors:

$$(2.1) \quad v_{\pm(2\beta_1+\beta_2)} = \frac{1}{2}(H_1 \pm i(X_{2\alpha_1+\alpha_2} + X_{-2\alpha_1-\alpha_2}))$$

$$(2.2) \quad v_{\pm\beta_2} = \frac{1}{2}(H_2 \pm i(X_{\alpha_2} + X_{-\alpha_2}))$$

$$(2.3) \quad v_{\pm(\beta_1+\beta_2)} = \frac{1}{2}(X_{\alpha_1+\alpha_2} + X_{-\alpha_1-\alpha_2} \mp i(X_{\alpha_1} + X_{-\alpha_1})).$$

2.2.2. *A Cayley transform and the maximal noncompact Cartan subalgebra.* There are 4 conjugacy classes of Cartan subgroups of $Sp(4, \mathbb{R})$ displayed in the diagram below, connected by Cayley transforms corresponding to the roots β_2 and $2\beta_1 + \beta_2$:

$$(2.4) \quad \begin{array}{ccc} & \mathbb{S}^1 \times \mathbb{S}^1 & \\ \swarrow c_{\beta_2} & & \searrow c_{2\beta_1+\beta_2} \\ \mathbb{R}^\times \times \mathbb{S}^1 & & \mathbb{S}^1 \times \mathbb{R}^\times \\ \searrow c_{2\beta_1+\beta_2} & & \swarrow c_{\beta_2} \\ & \mathbb{R}^\times \times \mathbb{R}^\times & \end{array}$$

Consider the *Cayley* transforms $c_\beta = \text{Ad exp}(\frac{\pi}{4}(\overline{v_\beta} - v_\beta))$ for a noncompact root β , when $\beta = \beta_2$ or $2\beta_1 + \beta_2$, the Cayley transform is nontrivial:

$$\begin{aligned} c_{\beta_2} &= \text{Ad exp}\left(\frac{\pi}{4}(\overline{v_{\beta_2}} - v_{\beta_2})\right) \\ c_{2\beta_1+\beta_2} &= \text{Ad exp}\left(\frac{\pi}{4}(\overline{v_{2\beta_1+\beta_2}} - v_{2\beta_1+\beta_2})\right). \end{aligned}$$

The two Cayley transforms c_{β_2} and $c_{2\beta_1+\beta_2}$ commute with each other. If we start from the real Cartan subalgebra $\mathfrak{t}$, the other real Cartan subalgebras of $Sp(4, \mathbb{R})$ can be obtained by applying the Cayley transforms c_{β_2} and $c_{2\beta_1+\beta_2}$:

$$\begin{array}{ccc} & \mathbb{R}(U_0 + U_3) \oplus \mathbb{R}(U_0 - U_3) & \\ \swarrow c_{\beta_2} & & \searrow c_{2\beta_1+\beta_2} \\ \mathbb{R}(U_0 + U_3) \oplus \mathbb{R}iH_2 & & \mathbb{R}iH_1 \oplus \mathbb{R}(U_0 - U_3) \\ \searrow c_{2\beta_1+\beta_2} & & \swarrow c_{\beta_2} \\ & \mathbb{R}iH_1 \oplus \mathbb{R}iH_2 & \end{array}$$

This diagram shows how the generators of the Cartan subalgebras are mapped to each other by the Cayley transforms shown in the diagram (2.4). We define the *maximally noncompact Cartan subalgebra* $\mathfrak{a} = \mathbb{R}H_1 \oplus \mathbb{R}H_2 \subset \mathfrak{p}$. The roots α_i acts on $\mathfrak{h}_{\mathbb{C}} = \mathfrak{a} \otimes_{\mathbb{R}} \mathbb{C}$, and they can be related to the roots β_i on the compact Cartan subgroup by

$$\alpha_i \circ (c_{\beta_2} c_{2\beta_1+\beta_2}) = \beta_i.$$

Moreover, we can apply the composite Cayley transform on the root vectors v_β :

$$c_{\beta_2} c_{2\beta_1+\beta_2} v_\beta = c_{2\beta_1+\beta_2} c_{\beta_2} v_\beta = \begin{cases} iX_{\beta \circ (c_{\beta_2} c_{2\beta_1+\beta_2})^{-1}} & \text{if } \beta \in \{\beta_2, 2\beta_1 + \beta_2\} \\ X_{\beta \circ (c_{\beta_2} c_{2\beta_1+\beta_2})^{-1}} & \text{if } \beta \in \{\beta_1 + \beta_2\} \end{cases}.$$

2.2.3. Weyl Group. From the embeddings $\Phi_{\alpha_1}, \Phi_{\alpha_2}$ of $SL(2, \mathbb{R})$ into $Sp(4, \mathbb{R})$ given by the simple roots α_1, α_2 , the simple Weyl reflections corresponding to these two roots are:

$$w_{\alpha_i} = \Phi_{\alpha_i} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \exp \left(\frac{\pi}{2} (X_{\alpha_i} - X_{-\alpha_i}) \right).$$

The Lie algebra $\mathfrak{k}$ has basis $\{X_\alpha - X_{-\alpha}\}_{\alpha \in \Delta^+}$, and the relationship between this basis and the basis $\{U_i\}$ is:

$$\begin{aligned} U_0 + U_3 &= X_{2\alpha_1+\alpha_2} - X_{-2\alpha_1-\alpha_2} & 2U_1 &= X_{\alpha_1+\alpha_2} - X_{-\alpha_1-\alpha_2} \\ 2U_2 &= X_{\alpha_1} - X_{-\alpha_1} & U_0 - U_3 &= X_{\alpha_2} - X_{-\alpha_2}. \end{aligned}$$

Therefore, the simple reflections w_{α_i} can be expressed as

$$(2.5) \quad w_{\alpha_1} = \exp(\pi U_2), w_{\alpha_2} = \exp \left(\frac{\pi}{2} (U_0 - U_3) \right).$$

Under the basis $\{\alpha_1, \alpha_2\}$, the actions of the simple reflections on an element $n_1\alpha_1 + n_2\alpha_2 \in \mathfrak{h}_{\mathbb{C}}^*$ are:

$$\begin{aligned} w_{\alpha_1}(n_1\alpha_1 + n_2\alpha_2) &= (2n_2 - n_1)\alpha_1 + n_2\alpha_2 \\ w_{\alpha_2}(n_1\alpha_1 + n_2\alpha_2) &= n_1\alpha_1 + (n_1 - n_2)\alpha_2. \end{aligned}$$

If we represent $\lambda \in \mathfrak{h}_{\mathbb{C}}^*$ by (λ_1, λ_2) , the actions of w_{α_i} on (λ_1, λ_2) where $\lambda_i = \lambda(H_i)$ is

$$\begin{aligned} w_{\alpha_1}(\lambda_1, \lambda_2) &= (\lambda_2, \lambda_1) \\ w_{\alpha_2}(\lambda_1, \lambda_2) &= (\lambda_1, -\lambda_2). \end{aligned}$$

The action of the simple reflections on the nilpotent radical satisfies:

$$\bar{\mathfrak{n}} \cap \text{Ad}(w_{\alpha_i})^{-1} \mathfrak{n} = \mathbb{R} X_{-\alpha_i}.$$

2.2.4. Harish-Chandra isomorphism. Since $\mathfrak{sp}(4, \mathbb{C})$ is rank 2, there are two Casimir elements Ω_2 and Ω_4 with degree 2 and degree 4 respectively. They generate the center $Z(\mathfrak{g}_{\mathbb{C}})$ of the complexified universal enveloping algebra. Consider the adjoint representation ad on $\mathfrak{g}_{\mathbb{C}}$, then the corresponding element in the center is [Yan11]:

$$\begin{aligned} \Omega_2 &= \sum_{i,j} \text{Tr}(\text{ad}(X_i)\text{ad}(X_j)) \tilde{X}_i \tilde{X}_j = \sum_{i,j} B(X_i, X_j) \tilde{X}_i \tilde{X}_j \\ &= \frac{1}{12} (H_1^2 + H_2^2 + 4H_1 + 2H_2 + 2(X_{-\alpha_1} X_{\alpha_1} + 2X_{-\alpha_2} X_{\alpha_2} \\ &\quad + X_{-\alpha_1-\alpha_2} X_{\alpha_1+\alpha_2} + 2X_{-2\alpha_1-\alpha_2} X_{2\alpha_1+\alpha_2})) \end{aligned}$$

where $\{\tilde{X}_i\}$ is a dual basis of the basis $\{X_i\}$ of $\mathfrak{g}_{\mathbb{C}}$ under the Killing form on $\mathfrak{g}$. The central element Ω_4 corresponding to the standard representation is a quartic element in $U(\mathfrak{g}_{\mathbb{C}})$ which we will only write down the $U(\mathfrak{h}_{\mathbb{C}})$ part explicitly:

$$\begin{aligned} \Omega_4 &= \frac{1}{5184} ((H_1 + H_2 + 3)^4 - 2(2H_1 H_2 + 2H_1 + 4H_2 + 7)((H_1 + 2)^2 + \\ &\quad (H_2 + 1)^2) - 11) + \text{other terms} \end{aligned}$$

Since $\rho = \varpi_1 + \varpi_2$, the Harish-Chandra homomorphism γ' will map each H_i to:

$$\gamma'(H_1) = H_1 - 2, \gamma'(H_2) = H_2 - 1$$

The image of Ω_2 and Ω_4 under the Harish-Chandra isomorphism γ' are:

$$\begin{aligned} \gamma'(\Omega_2) &= \frac{1}{12}(H_1^2 + H_2^2 - 5) \\ \gamma'(\Omega_4) &= \frac{1}{5184}(H_1^4 + H_2^4 + 6H_1^2 H_2^2 - 6(H_1^2 + H_2^2) - 11). \end{aligned}$$

3. IWASAWA DECOMPOSITION ON THE GROUP LEVEL

We would like to compute the Iwasawa decomposition $kman$ of an element $\bar{n} \in \overline{N}$. For a root α with $\chi_{-\alpha}(t) = \exp(tX_{-\alpha}) \in \overline{N}$, the corresponding Lie group $SL(2, \mathbb{C})$ embedded into $G_{\mathbb{C}}$ has generators:

$$\chi_{\alpha}(t) = \exp(tX_{\alpha}), \chi_{-\alpha}(t) = \exp(tX_{-\alpha}), h_{\alpha}(t) = \exp(tH_{\alpha})$$

In terms of these matrix generators of $SL(2, \mathbb{C})$, the Iwasawa decomposition of $\chi_{-\alpha}(t)$ is

$$(3.1) \quad \chi_{-\alpha}(t) = \kappa_{\alpha}(t) h_{\alpha} \left(\sqrt{1+t^2} \right) \chi_{\alpha} \left(\frac{t}{1+t^2} \right)$$

where

$$\kappa_{\alpha}(t) = \exp(\arctan(-t)(X_{\alpha} - X_{-\alpha}))$$

is an element in the maximal compact subgroup $K_0 \cong SO(2, \mathbb{R})$. In general, for any simple real root α for $G = Sp(4, \mathbb{R})$, we would like to study at the embedding ϕ_{α} of a Lie algebra $\mathfrak{sl}(2, \mathbb{R})$ into $\mathfrak{g}$, and the corresponding homomorphism of a Lie group $\Phi_{\alpha} : SL(2, \mathbb{R}) \rightarrow G$. The image of Φ_{α} is fixed by θ , and the embedding for the Lie algebra satisfies:

- (1) $\phi_{\alpha} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = X_{\alpha}, \phi_{\alpha} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = X_{-\alpha}, \phi_{\alpha} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = H_{\alpha},$
- (2) $\Phi_{\alpha} \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} = \chi_{\alpha}(t), \Phi_{\alpha} \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix} = \chi_{-\alpha}(t), \Phi_{\alpha} \begin{pmatrix} t & 0 \\ 0 & 1/t \end{pmatrix} = h_{\alpha}(t).$

We can see that the homomorphisms Φ_{α} and ϕ_{α} respect the Cartan involution: if we denote θ' as the Cartan involution on $SL(2, \mathbb{R})$ and $\mathfrak{sl}(2, \mathbb{R})$, we can see that $\Phi_{\alpha} \circ \theta' = \theta \circ \Phi_{\alpha}$ and $\phi_{\alpha} \circ \theta' = \theta \circ \phi_{\alpha}$. Therefore, $\text{Im}(\Phi_{\alpha}) \cap K = \Phi_{\alpha}(K_0)$ where K_0 is the maximal compact subgroup $SO(2, \mathbb{R})$ in $SL(2, \mathbb{R})$.

Recall that under any simple reflection w_{α_i} , there is a root vector $X_{-\alpha_i}$ such that $\bar{n} \cap w_{\alpha_i}^{-1} \bar{n} w_{\alpha_i} = \mathbb{R} X_{-\alpha_i}$. We can factor the nilpotent group $w_{\alpha_i}^{-1} N w_{\alpha_i}$ into

$$w_{\alpha_i}^{-1} N w_{\alpha_i} = N_{-\alpha} N' \text{ where } N' \subset N \text{ and } N_{-\alpha} = \{e^{tX_{-\alpha_i}} | t \in \mathbb{R}\}.$$

Using this factorization of the group $w_{\alpha_i}^{-1} N w_{\alpha_i}$, we can find a coordinate system on $w_{\alpha_i}^{-1} N w_{\alpha_i}$ which is consistent with the Iwasawa decomposition. Namely, for any $n \in N$, there is a $t \in \mathbb{R}$ and $n' \in N'$ such that

$$w_{\alpha_i}^{-1} n w_{\alpha_i} = \kappa_{\alpha}(t) h_{\alpha} \left(\sqrt{1+t^2} \right) \chi_{\alpha} \left(\frac{t}{1+t^2} \right) n'.$$

4. PARABOLIC SUBGROUPS OF $Sp(4, \mathbb{R})$

4.1. Minimal Parabolic Subgroup and Jacobi Parabolic Subgroup. Denote P_0 as the *minimal parabolic* subgroup of $Sp(4, \mathbb{R})$, with the Lie algebra having a Levi decomposition:

$$\mathfrak{p}_0 = \mathfrak{a} \oplus \mathfrak{n}_0, \text{ where } \mathfrak{n}_0 = \bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_\alpha.$$

For any real root α of $\Delta(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$, consider the homomorphism $\phi_\alpha : \mathfrak{sl}(2, \mathbb{R}) \rightarrow \mathfrak{g}$, which exponentiates to a homomorphism $\Phi_\alpha : \mathrm{SL}(2, \mathbb{C}) \rightarrow G_{\mathbb{C}}$. Letting γ_α be the image of $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$, then

$$\gamma_\alpha = \exp \frac{2\pi i H_\alpha}{|\alpha|^2} = \exp \pi (X_\alpha - \theta X_\alpha)$$

Recalling the statements in [Kna13, Section VII.4], every nonidentity element of the centralizer M of $\mathfrak{a}$ in K has order 2, and is generated by γ_α for all real roots α . In the case of $Sp(4, \mathbb{R})$,

$$M = \{\gamma_{\alpha_2}^{\epsilon_1} \gamma_{2\alpha_1 + \alpha_2}^{\epsilon_2} | \epsilon_i = 0 \text{ or } 1\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2,$$

The generators $\gamma_{\alpha_2}, \gamma_{2\alpha_1 + \alpha_2}$ of M can also be expressed in terms of U_i :

$$\gamma_{\alpha_2} = \exp(\pi(U_0 - U_3)), \gamma_{2\alpha_1 + \alpha_2} = \exp(\pi(U_0 + U_3)).$$

The minimal parabolic subgroup P_0 has a Levi decomposition $P_0 = MA_0N_0$ M is the centralizer of $\mathfrak{a}$ in K , and A_0 and N_0 are the analytic subgroups formed by exponentiating $\mathfrak{a}$ and $\mathfrak{n}_0$, respectively.

There are two proper standard parabolic subgroups that contain P_0 . The parabolic subgroup P_1 with an abelian nilpotent radical is called a *Siegel* subgroup. The Levi decomposition $P_1 = M_1N_1$ has the form:

$$M_1 = \left\{ \begin{pmatrix} A & 0 \\ 0 & (A^t)^{-1} \end{pmatrix}, A \in GL(2, \mathbb{R}) \right\} \cong GL(2, \mathbb{R})$$

$$N_1 = \left\{ \begin{pmatrix} 1 & 0 & x_4 & x_3 \\ 0 & 1 & x_3 & x_2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, x_i \in \mathbb{R} \right\} \cong \mathbb{R}^3.$$

Their Lie algebras have restricted root space decompositions:

$$\mathfrak{m}_1 = \mathfrak{a} \oplus \mathfrak{g}_{\alpha_1} \oplus \mathfrak{g}_{-\alpha_1}$$

$$\mathfrak{n}_1 = \bigoplus_{\alpha \in \{\alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\}} \mathfrak{g}_\alpha.$$

There is another class of parabolic subgroup called the *Jacobi* (or *Heisenberg* and in some literature also called *Klingen*) parabolic subgroup $P_2 = M_2N_2$, having a Levi decomposition:

$$M_2 = \left\{ \begin{pmatrix} h_1 & 0 & 0 & 0 \\ 0 & a & 0 & b \\ 0 & 0 & 1/h_1 & 0 \\ 0 & c & 0 & d \end{pmatrix}, \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{R}), h_1 \in \mathbb{R}^\times \right\} \cong SL(2, \mathbb{R}) \times \mathbb{R}^\times$$

$$N_2 = \left\{ \begin{pmatrix} 1 & x_1 & x_4 & x_3 \\ 0 & 1 & x_3 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -x_1 & 1 \end{pmatrix}, x_i \in \mathbb{R} \right\}.$$

The Lie algebras of each subgroup have restricted root space decompositions:

$$\begin{aligned}\mathfrak{m}_2 &= \mathfrak{a} \oplus \mathfrak{g}_{\alpha_2} \oplus \mathfrak{g}_{-\alpha_2} \\ \mathfrak{n}_2 &= \bigoplus_{\alpha \in \{\alpha_1, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\}} \mathfrak{g}_\alpha.\end{aligned}$$

Thus $\mathfrak{n}_2 = \bigoplus_{\alpha \in \{\alpha_1, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\}} \mathbb{R}X_\alpha$. Since the root vectors X_α satisfy the commutation relations

$$\begin{aligned}[X_\alpha, X_{2\alpha_1 + \alpha_2}] &= 0 \text{ for } \alpha \in \{\alpha_1, \alpha_1 + \alpha_2\} \\ [X_{\alpha_1}, X_{\alpha_1 + \alpha_2}] &= 2X_{2\alpha_1 + \alpha_2},\end{aligned}$$

the group N_2 is isomorphic to the Heisenberg group $\mathcal{H}_3$.

4.2. Induction from the Minimal Parabolic Subgroup. In this section consider the minimal principal series obtained by induction from the minimal parabolic subgroup P_0 . Consider the Cartan involution $\theta(g) = (g^t)^{-1}$ on $Sp(2n, \mathbb{R})$, which acts on the split Cartan subalgebra

$$\mathfrak{h}_{\mathbb{R}} = \bigoplus_{i=1}^n \mathbb{C}(E_{i,i} - E_{n+i,n+i})$$

by sending all elements to their negatives. In this case, $\theta = -\text{id}$ on the root lattice X . Therefore,

$$X/(1 - \theta)X = \bigoplus_{i=1}^n \mathbb{Z}/2\mathbb{Z}\varpi_i$$

and -1 eigenspace of θ on X is $X^{-\theta} = X$. The discrete character δ and the continuous character λ can thus be determined by two vectors

$$(4.1) \quad \delta = \sum_{i=1}^n \delta_i \varpi_i \longleftrightarrow (\delta_1, \dots, \delta_n) \quad \delta_i \in \mathbb{Z}/2\mathbb{Z}$$

$$(4.2) \quad \lambda = \sum_{i=1}^n \lambda_i \varpi_i \longleftrightarrow (\lambda_1, \dots, \lambda_n) \quad \lambda \in \mathbb{C}.$$

If all of the λ_i 's are integers satisfying $\lambda_i \equiv \delta_i \pmod{2}$, then there exists an integral character $\chi_{\delta, \lambda} = \sum_{i=1}^n \lambda_i \varpi_i$ such that δ is the image of χ in the quotient $X/(1 - \theta)X$, and $\lambda = \left(\frac{1-\theta}{2}\right) \chi$.

In the case of $Sp(2n, \mathbb{R})$, a minimal principal series representation $I_{P_0}(\chi_{\delta, \lambda})$ of $Sp(4, \mathbb{R})$ is determined by the following data:

- (1) A continuous character $\lambda : \mathfrak{a} \rightarrow \mathbb{C}$ represented by the pair (λ_1, λ_2) with $\lambda_i = \lambda(H_i)$. This character can be extended linearly to a character on $\mathfrak{a} \otimes \mathbb{C} = \mathfrak{h}$, also denoted by λ ;
- (2) A character $\delta : M \rightarrow \{\pm 1\}$ represented by a pair (δ_1, δ_2) where $\delta_i \in \{0, 1\}$ such that $\delta(\gamma_{\alpha_2}^{\epsilon_1} \gamma_{2\alpha_1 + \alpha_2}^{\epsilon_2}) = (-1)^{\delta_1 \epsilon_1 + \delta_2 \epsilon_2}$.

If the numbers λ_i are integers, with $\delta_i \equiv \lambda_i \pmod{2}$, the parameters defined in (4.1)-(4.2) combine to an algebraic character $\chi_{\delta, \lambda}$ on the complex Cartan subalgebra $\mathfrak{a}_{\mathbb{C}} = \mathbb{C}H_1 \oplus \mathbb{C}H_2$.

The exponential map from $\mathfrak{a}$ to the split torus A_0 is a bijection. Every element $a \in A_0$ can be written as $a = \exp H_a$ for some $H_a \in A_0$. For any $\nu \in X$, we

introduce the notation $a^\nu = \exp \nu(H_a)$. The algebraic character $\chi_{\delta, \lambda}$ exponentiates to a character $\chi_{\delta, \lambda} : MA_0 \rightarrow \mathbb{C}^*$ given by

$$(\chi_{\delta, \lambda + \rho})(ma) = \delta(m)a^{\lambda + \rho}.$$

We define the *principal series* representation $I_{P_0}(\delta, \lambda)$ induced from the minimal parabolic subgroup $P \subset G$ as the following vector space of functions on G :

$$(4.3) \quad I_P(\chi_{\delta, \lambda}) = \{f : G \rightarrow \mathbb{C} \mid f(gman) = (\chi_{\delta, \lambda + \rho})^{-1}(ma)f(g)\}.$$

The action $\pi_{P_0}(\chi_{\delta, \lambda})$ of G on $f \in I_{P_0}(\chi_{\delta, \lambda})$ is given by the *left* regular representation

$$(\pi_{P_0}(\chi_{\delta, \lambda})(g)f)(h) = f(g^{-1}h).$$

The $(\mathfrak{g}, K)$ -module of the principal series representation $I_{P_0}(\chi_{\delta, \lambda})$ embeds into the space

$$C_\delta(K) = \{f : K \rightarrow \mathbb{C} \mid f \text{ smooth and } f(km) = \delta(m)^{-1}f(k)\}.$$

This space is isomorphic to the space of smooth global sections of the line bundle $K \otimes_M \mathbb{C}_\delta$, where $\mathbb{C}_\delta$ is the vector space on which M acts by δ^{-1} . To get a basis for the space $C_\delta(K)$ from the Wigner D -functions, we consider the right action by $m \in M$ on an arbitrary function $f \in C_\delta(K)$. f can be written as the finite linear combination of Wigner D -functions with coefficients $a_{m_1, m_2}^{(j, n)}$:

$$f(k) = \sum_{\substack{j, n \\ -j \leq m_i \leq j}} a_{m_1, m_2}^{(j, n)} W_{m_1, m_2}^{(j, n)}(k).$$

The action by $m \in M$ on the right gives

$$f(km) = \sum_{\substack{j, n \\ -j \leq m_i \leq j}} a_{m_1, m_2}^{(j, n)} W_{m_1, m_2}^{(j, n)}(km).$$

Recall that the Wigner D -functions are matrix coefficients of $U(2)$ -representations. Their values on the product of two elements km come from the multiplication of two matrices:

$$W_{m_1, m_2}^{(j, n)}(km) = \sum_{m_3} W_{m_1, m_3}^{(j, n)}(k) W_{m_3, m_2}^{(j, n)}(m).$$

The action of a general element

$$m = \gamma_{\alpha_2}^{\epsilon_1} \gamma_{2\alpha_1 + \alpha_2}^{\epsilon_2} = e^{\pi(\epsilon_1 + \epsilon_2)U_0} e^{\pi(-\epsilon_1 + \epsilon_2)U_3} \in M$$

on the Wigner D -functions is diagonal:

$$W_{m_3, m_2}^{(j, n)}(e^{\pi(\epsilon_1 + \epsilon_2)U_0} e^{\pi(-\epsilon_1 + \epsilon_2)U_3}) = (-1)^{-(n-m_2)\epsilon_1 - (n+m_2)\epsilon_2} \delta_{m_3, m_2}.$$

Therefore, $f(km)$ can be written as

$$f(km) = \sum_{\substack{j, n \\ -j \leq m_i \leq j}} a_{m_1, m_2}^{(j, n)} (-1)^{-(n-m_2)\epsilon_1 - (n+m_2)\epsilon_2} W_{m_1, m_2}^{(j, n)}(k).$$

Because of the linear independence of different Wigner D -functions, the equality $f(km) = \delta(m)^{-1}f(k) = (-1)^{-\delta_1\epsilon_1 - \delta_2\epsilon_2}f(k)$ holds for all $k \in K$ if and only if j, n, m_1, m_2 satisfy the compatibility condition

$$(-1)^{(n-m_2)\epsilon_1 + (n+m_2)\epsilon_2} = (-1)^{\delta_1\epsilon_1 + \delta_2\epsilon_2}.$$

Therefore, the space $C_\delta(K)$ can be written as the direct sum

$$(4.4) \quad C_\delta(K) = \bigoplus_{(j,n) \in \mathbf{KTypes}(\delta_1, \delta_2)} \bigoplus_{\substack{m_1 \in \{-j, -j+1, \dots, j\} \\ m_2 \in \mathbf{M}(j, n; \delta_1, \delta_2)}} \mathbb{C}W_{m_1, m_2}^{(j, n)}$$

in which the two sets of admissible j, n, m_1, m_2 are defined as:

$$(4.5) \quad \mathbf{KTypes}(\delta_1, \delta_2) = \{(j, n) \in \frac{1}{2}\mathbb{Z}_{\geq 0} \times \frac{1}{2}\mathbb{Z} \mid 2j \equiv 2n \equiv \delta_1 + \delta_2 \pmod{2}\}$$

$$(4.6)$$

$$\mathbf{M}(j, n; \delta_1, \delta_2) = \{m_2 \in \{-j, -j+1, \dots, j\} \mid n - m_2 \equiv \delta_1 \text{ and } n + m_2 \equiv \delta_2 \pmod{2}\}.$$

Similarly to the case of $SU(2, 1)$, for each $(j, n) \in \mathbf{KTypes}(\delta_1, \delta_2)$, we can denote

$$(4.7) \quad \tau^{(j, n)} = \bigoplus_{\substack{m_1 \in \{-j, -j+1, \dots, j\} \\ m_2 \in \mathbf{M}(j, n; \delta_1, \delta_2)}} \mathbb{C}W_{m_1, m_2}^{(j, n)}$$

as the K -isotypic subspace of $I_{P_0}(\delta, \lambda)$ which decomposes into copies of irreducible K -representations of highest weight (j, n) . The restriction of the $(\mathfrak{g}, K)$ -module of the principal series $I_{P_0}(\delta, \lambda)$ to K can be decomposed as a direct sum of the K -isotypic spaces $\tau^{(j, n)}$:

$$(4.8) \quad I_{P_0}(\delta, \lambda) = \bigoplus_{(j, n) \in \mathbf{KTypes}(\delta_1, \delta_2)} \tau^{(j, n)}.$$

The different copies of irreducible K -representations are distinguished by the index m_2 , and the action of $\mathfrak{u}(2)$ raising and lowering operators $U_1 \pm iU_2$ moves each m_1 to $m_1 \pm 1$. For each K -isotypic space $\tau^{(j, n)}$, the cardinality of the set $\mathbf{M}(j, n; \delta_1, \delta_2)$ is equal to the multiplicity of K -types with highest weight (j, n) . In fact, if we assume

$$(\delta_1, \delta_2) \in \{(0, 0), (1, 1)\},$$

the set $\mathbf{M}(j, n; \delta_1, \delta_2)$ is

$$(4.9) \quad \mathbf{M}(j, n; \delta_1, \delta_2) = \begin{cases} \{\dots, j-4, j-2, j\} & j-n+\delta_1 \equiv 0 \pmod{2} \\ \{\dots, j-5, j-3, j-1\} & j-n+\delta_1 \equiv 1 \pmod{2} \end{cases}$$

and its cardinality is

$$(4.10) \quad |\mathbf{M}(j, n; \delta_1, \delta_2)| = \begin{cases} j+1 & j-n+\delta_1 \equiv 0 \pmod{2} \\ \lfloor j-\frac{1}{2} \rfloor + 1 & j-n+\delta_1 \equiv 1 \pmod{2} \end{cases}.$$

We will make use of these facts to calculate the long intertwining operator for the principal series of $Sp(4, \mathbb{R})$.

5. THE $(\mathfrak{g}, K)$ -MODULE STRUCTURE

5.1. Normalization of basis and the Iwasawa decomposition. The set of positive and negative noncompact roots are

$$\Delta_{nc}^+ = \{\beta_2, \beta_1 + \beta_2, 2\beta_1 + \beta_2\} \quad \Delta_{nc}^- = \{-\beta_2, -\beta_1 - \beta_2, -2\beta_1 - \beta_2\}.$$

We define the vectors $u_\beta \in \mathfrak{p}_{\mathbb{C}}$ by multiplying the v_β defined in (2.1)-(2.3) by a factor:

$$u_\beta = \begin{cases} \sqrt{2}i v_\beta & \beta \in \{\beta_2, 2\beta_1 + \beta_2, -\beta_2, -2\beta_1 - \beta_2\} \\ v_\beta & \text{otherwise} \end{cases}.$$

Under this normalization, the Lie algebra Iwasawa decomposition of the basis vectors u_β is:

$$(5.1) \quad u_{\pm(2\beta_1+\beta_2)} = \frac{1}{\sqrt{2}}(\pm(U_0 + U_3) + iH_1 \mp 2X_{2\alpha_1+\alpha_2})$$

$$(5.2) \quad u_{\pm\beta_2} = \frac{1}{\sqrt{2}}(\pm(U_0 - U_3) + iH_2 \mp 2X_{\alpha_2})$$

$$(5.3) \quad u_{\pm(\beta_1+\beta_2)} = -(U_1 \mp iU_2) + X_{\alpha_1+\alpha_2} \mp iX_{\alpha_1}$$

The representation of $K = U(2)$ on $\mathfrak{p}_{\mathbb{C}}$ decomposes into two irreducible subrepresentations

$$\mathfrak{p}_{\mathbb{C}}^+ = \oplus_{\beta \in \Delta_{nc}^+} \mathbb{C}u_\beta \quad \mathfrak{p}_{\mathbb{C}}^- = \oplus_{\beta \in \Delta_{nc}^-} \mathbb{C}u_\beta.$$

The u_β 's are the normalized weight vectors on which the adjoint action by $U_1 \pm iU_2$ acts as

$$\text{ad}(U_1 \pm iU_2)u_\beta = q_{\beta, \beta \pm \beta_1} u_{\beta \pm \beta_1},$$

where the coefficient $q_{\beta, \beta \pm \beta_1}$ turns out to be

$$q_{\beta, \beta \pm \beta_1} = \begin{cases} -i\sqrt{2} & \text{if } \beta \pm \beta_1 \in \Delta(\mathfrak{g}_{\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}) \\ 0 & \text{otherwise} \end{cases}.$$

5.2. Left Action of $\mathfrak{p}_{\mathbb{C}}$. For any noncompact root β , define two integers:

$$(5.4) \quad m_\beta = -i\beta(U_3) \in \{-1, 0, 1\}$$

$$(5.5) \quad n_\beta = -i\beta(U_0) \in \{-1, 1\}$$

Each weight β is uniquely determined by the pair of integers (m_β, n_β) specified in the following chart:

	$m_\beta = -1$	$m_\beta = 0$	$m_\beta = 1$
$n_\beta = -1$	$-2\beta_1 - \beta_2$	$-\beta_1 - \beta_2$	$-\beta_2$
$n_\beta = 1$	β_2	$\beta_1 + \beta_2$	$2\beta_1 + \beta_2$

TABLE 1. Correspondence between (m_β, n_β) and noncompact roots

By embedding the principal series representation $I(\chi_{\delta, \lambda})$ into $C_\delta(K)$ (compare to the identical method in [BM17] and the $SU(2, 1)$ case in Section 5), we would like to understand the action of

$$dl(u_\beta) = dr(-\text{Ad}(k^{-1})u_\beta)$$

on any basis vector $W_{m_1, m_2}^{(j, n)}(k)$ in $C_\delta(K)$. By the definition of the principal series in (4.3), the right action on any vector $f \in I(\chi_{\delta, \lambda})$ by H_i is always a scalar multiplication by $\lambda_i + \rho(H_i)$

$$(5.6) \quad dr(H_i)f = -(\lambda_i + \rho(H_i))f,$$

and the right action by any element from N annihilates f . We can therefore use the decomposition (5.1)-(5.3) and the differential operators (A.7)-(A.9) in $\mathfrak{g}_{\mathbb{C}}$ of u_β

to calculate the right action of u_β on the basis vectors $W_{m_1, m_2}^{(j, n)}$ of $I(\chi_{\delta, \lambda})$. Also, recall from 6.1.1 that $\rho(H_1) = 2, \rho(H_2) = 1$, we have

$$\begin{aligned} \text{dr}(u_{\pm(2\beta_1 + \beta_2)})W_{m_1, m_2}^{(j, n)} &= \frac{i}{\sqrt{2}}(\mp n \mp m_2 - (\lambda_1 + \rho(H_1)))W_{m_1, m_2}^{(j, n)} \\ &= \frac{i(\mp n \mp m_2 - (\lambda_1 + 2))}{\sqrt{2}}W_{m_1, m_2}^{(j, n)} \end{aligned} \quad (5.7)$$

$$\begin{aligned} \text{dr}(u_{\pm\beta_2})W_{m_1, m_2}^{(j, n)} &= \frac{i}{\sqrt{2}}(\mp n \pm m_2 - (\lambda_2 + \rho(H_2)))W_{m_1, m_2}^{(j, n)} \\ &= \frac{i(\mp n \pm m_2 - (\lambda_2 + 1))}{\sqrt{2}}W_{m_1, m_2}^{(j, n)} \end{aligned} \quad (5.8)$$

$$\text{dr}(u_{\pm(\beta_1 + \beta_2)})W_{m_1, m_2}^{(j, n)} = -i\sqrt{(j \pm m_2)(j \mp m_2 + 1)}W_{m_1, m_2 \mp 1}^{(j, n)}. \quad (5.9)$$

Recalling the correspondence between the weight β and (m_β, n_β) discussed in (5.4)-(5.5) and Table 1, since the irreducible constituents of $\mathfrak{p}_\mathbb{C}^\pm$ have highest weights $(j, n) = (1, \pm 1)$ respectively, the right action of u_β with $\beta \in \Delta_{nc}^\pm$ can be transferred to the left by observing

$$\text{Ad}(k^{-1})u_\beta = \sum_{\nu \in \Delta_{nc}^\pm} \overline{W_{m_\nu, m_\beta}^{(1, \pm 1)}(k^{-1})}u_\nu = \sum_{\nu \in \Delta_{nc}^\pm} W_{m_\beta, m_\nu}^{(1, \pm 1)}(k)u_\nu.$$

Based on the correspondence in Table 1 between the weights of u_β and the pair of integers (m_β, n_β) , it is clear that the left action of u_β for $\beta \in \Delta_{nc}^\pm$ on the Wigner D -functions $W_{m_1, m_2}^{(j, n)} \in \mathcal{C}_\delta(K)$ can be written explicitly as follows,

$$\begin{aligned} \text{dl}(u_\beta)W_{m_1, m_2}^{(j, n)} &= \\ &\left(W_{m_\beta, \mp 1}^{(1, \pm 1)}\text{dr}(u_{\pm\beta_2}) + W_{m_\beta, 0}^{(1, \pm 1)}\text{dr}(u_{\pm(\beta_1 + \beta_2)}) + W_{m_\beta, \pm 1}^{(1, \pm 1)}\text{dr}(u_{\pm(2\beta_1 + \beta_2)}) \right) W_{m_1, m_2}^{(j, n)}. \end{aligned} \quad (5.10)$$

According to (5.7)-(5.9), we apply the right action $\text{dr}(u_\beta)$ to $W_{m_1, m_2}^{(j, n)}$ and get

$$\begin{aligned} \text{dl}(u_\beta)W_{m_1, m_2}^{(j, n)} &= \\ i \left(\frac{\mp n \pm m_2 - (\lambda_2 + \rho(H_2))}{\sqrt{2}}W_{m_\beta, \mp 1}^{(1, \pm 1)} + \frac{\mp n \mp m_2 - (\lambda_1 + \rho(H_1))}{\sqrt{2}}W_{m_\beta, \pm 1}^{(1, \pm 1)} \right) W_{m_1, m_2}^{(j, n)} \\ &- i\sqrt{(j \pm m_2)(j \mp m_2 + 1)}W_{m_\beta, 0}^{(1, \pm 1)}W_{m_1, m_2 \mp 1}^{(j, n)}. \end{aligned} \quad (5.11)$$

We can replace the products of Wigner D -functions by a linear combination of Wigner D -functions with Clebsch-Gordan coefficients as described in formula (A.10) of Section A. The left action of u_β for $\beta \in \Delta_{nc}^\pm$ on Wigner D -functions can

thus be expressed as

$$\begin{aligned}
 & dl(u_\beta) W_{m_1, m_2}^{(j, n)} \\
 = & i \sum_{j_0 \in \{j-1, j, j+1\}} \binom{j+j_0, m_1+m_\beta}{j, m_1, 1, m_\beta} \left(\frac{\mp n \mp m_2 - (\lambda_1 + \rho(H_1))}{\sqrt{2}} \binom{j+j_0, m_2 \pm 1}{j, m_2, 1, \pm 1} W_{m_1+m_\beta, m_2 \pm 1}^{(j+j_0, n \pm 1)} + \right. \\
 & \left. \left(-\sqrt{(j \pm m_2)(j \mp m_2 + 1)} \binom{j+j_0, m_2 \mp 1}{j, m_2 \mp 1, 1, 0} + \frac{\mp n \pm m_2 - (\lambda_2 + \rho(H_2))}{\sqrt{2}} \binom{j+j_0, m_2 \mp 1}{j, m_2, 1, \mp 1} \right) \right) \\
 & W_{m_1+m_\beta, m_2 \mp 1}^{(j+j_0, n \pm 1)} \quad (5.12)
 \end{aligned}$$

After computing all the Clebsch-Gordan coefficients using the formulas listed in Table 2, the action of weight vectors u_β of $\mathfrak{p}_\mathbb{C}$ on the left when $\beta \in \Delta_{nc}^\pm$ can be expressed as the following linear combination:

$$(5.13) \quad dl(u_\beta) W_{m_1, m_2}^{(j, n)} = \frac{i}{2} \sum_{\substack{j_0 \in \{-1, 0, 1\} \\ \epsilon = \pm 1}} \binom{j+j_0, m_1+m_\beta}{j, m_1, 1, m_\beta} C_{j+j_0} q_{j_0, \epsilon} \kappa_{\pm, j_0, \epsilon}(j, n, m_1; \lambda) W_{m_1+m_\beta, m_2+\epsilon}^{(j+j_0, n \pm 1)}$$

with the coefficients given in the tables:

	$\frac{C_{j+j_0}}{j_0 = -1 \quad j^{-\frac{1}{2}}(2j+1)^{-\frac{1}{2}}}$	
	$j_0 = 0 \quad j^{-\frac{1}{2}}(j+1)^{-\frac{1}{2}}$	
	$j_0 = 1 \quad (j+1)^{-\frac{1}{2}}(2j+1)^{-\frac{1}{2}}$	
$q_{j_0, \epsilon}$	$\epsilon = -1$	$\epsilon = 1$
$j_0 = -1$	$\sqrt{(j+m_2-1)(j+m_2)}$	$\sqrt{(j-m_2-1)(j-m_2)}$
$j_0 = 0$	$\sqrt{(j+m_2)(j-m_2+1)}$	$\sqrt{(j-m_2)(j+m_2+1)}$
$j_0 = 1$	$\sqrt{(j-m_2+1)(j-m_2+2)}$	$\sqrt{(j+m_2+1)(j+m_2+2)}$
$\kappa_{+, j_0, \epsilon}$	$\epsilon = -1$	$\epsilon = 1$
$j_0 = -1$	$2+2j-m_2-n-(\lambda_2+\rho(H_2))$	$-n-m_2-(\lambda_1+\rho(H_1))$
$j_0 = 0$	$2-n-m_2-(\lambda_2+\rho(H_2))$	$n+m_2+(\lambda_1+\rho(H_1))$
$j_0 = 1$	$-2j-m_2-n-(\lambda_2+\rho(H_2))$	$-n-m_2-(\lambda_1+\rho(H_1))$
$\kappa_{-, j_0, \epsilon}$	$\epsilon = -1$	$\epsilon = 1$
$j_0 = -1$	$n+m_2-(\lambda_1+\rho(H_1))$	$2+2j+m_2+n-(\lambda_2+\rho(H_2))$
$j_0 = 0$	$n+m_2-(\lambda_1+\rho(H_1))$	$-2-n-m_2+(\lambda_2+\rho(H_2))$
$j_0 = 1$	$n+m_2-(\lambda_1+\rho(H_1))$	$-2j+m_2+n-(\lambda_2+\rho(H_2))$

The $(\mathfrak{g}, K)$ -action on $Sp(4, \mathbb{R})$ principal series $I(\chi_{\delta, \lambda})$ is completely determined by formula (5.13) and the four tables above.

6. INTERTWINING OPERATORS

The longest element $w_0 = w_{\alpha_2} w_{\alpha_1} w_{\alpha_2} w_{\alpha_1}$ in the Weyl group of $Sp(4, \mathbb{R})$ corresponds to the long intertwining operator:

$$A(w_0, \chi_{\delta, \lambda}) f(k) = \int_{N \cap w^{-1} N w} f(k w \bar{n}) d\bar{n}.$$

Applying the Langlands' Lemma on factorization of intertwining operators in [Sha10], $A(w_0, \chi_{\delta, \lambda})$ can be factored into 4 intertwining operators corresponding to simple

reflections:

$$(6.1) \quad A(w_0, \chi_{\delta, \lambda}) = A(w_{\alpha_2}, w_{\alpha_1} w_{\alpha_2} w_{\alpha_1} \chi_{\delta, \lambda}) A(w_{\alpha_1}, w_{\alpha_2} w_{\alpha_1} \chi_{\delta, \lambda}) A(w_{\alpha_2}, w_{\alpha_1} \chi_{\delta, \lambda}) \times A(w_{\alpha_1}, \chi_{\delta, \lambda}).$$

Since there is an embedding of the Harish-Chandra module of $I(\chi_{\delta, \lambda})$ into the space $C_\delta(K)$, we can express any function $f(k)$ in the principal series under such embedding as a linear combination of Wigner D -functions:

$$f(k) = \sum_{(j, n)} \sum_{(m_1, m_2)} a_{m_1, m_2}^{(j, n)} W_{m_1, m_2}^{(j, n)}(k).$$

Therefore, it is sufficient to compute the matrix coefficients of the intertwining operator on the basis $W_{m_1, m_2}^{(j, n)}$ of the space $C_\delta(K)$:

$$\begin{aligned} A(w, \chi_{\delta, \lambda}) W_{m_1, m_2}^{(j, n)}(k) &= \int_{\bar{N} \cap N^w} W_{m_1, m_2}^{(j, n)}(kw\bar{n}) d\bar{n} \\ &= \int_{\bar{N} \cap N^w} a(\bar{n})^{-(\lambda+\rho)} W_{m_1, m_2}^{(j, n)}(kwk(\bar{n})) d\bar{n} \\ &= \sum_{m_3} \left(\int_{\bar{N} \cap N^w} a(\bar{n})^{-(\lambda+\rho)} W_{m_3, m_2}^{(j, n)}(wk(\bar{n})) d\bar{n} \right) W_{m_1, m_3}^{(j, n)}(k), \end{aligned}$$

and the matrix coefficients of the intertwining operator $A(w, \delta, \lambda)$ are given by the formula

$$(6.2) \quad \langle W_{m_1, m_3}^{(j, n)}, A(w, \chi_{\delta, \lambda}) W_{m_1, m_2}^{(j, n)} \rangle = \int_{\bar{N} \cap N^w} a(\bar{n})^{-(\lambda+\rho)} W_{m_3, m_2}^{(j, n)}(wk(\bar{n})) d\bar{n}.$$

We will use this formula to compute the intertwining operators explicitly for the group $Sp(4, \mathbb{R})$ in this section. Combining the Langlands' Lemma and (6.2), we have the following proposition:

Proposition 6.1. *Let $w_0 = w_{\alpha_2} w_{\alpha_1} w_{\alpha_2} w_{\alpha_1}$ be the longest element in the Weyl group W of $Sp(4, \mathbb{R})$. The matrix for the long intertwining operator $A(w_0, \chi_{\delta, \lambda})$ under the basis $W_{m_1, m_2}^{(j, n)}$ of $C_\delta(K)$ has a factorization:*

$$A(w_0, \chi_{\delta, \lambda})|_{j, n} = A_4(\nu) \cdot A_3(\lambda) \cdot A_2(\lambda) \cdot A_1(\lambda)$$

If we define

$$(6.3) \quad Q(z, n) = \frac{\pi 2^{2-2z} \Gamma(2z-1)}{\Gamma(z+n) \Gamma(z-n)},$$

and let

$$\begin{aligned} S_{m_3, m_2}^{j, n}(z) &= \sum_{-j \leq m_4 \leq j} i^{-2m_4} M_{m_3, m_4}^{j, n} N_{m_4, m_2}^{j, n} Q(z, m_4) \\ T_{m_1}^n(z) &= i^{-n+m_1} Q\left(z, \frac{m_1-n}{2}\right) \end{aligned}$$

where $N_{m_1, m_3}^{j, n}$ is the inverse matrix of $M_{m_1, m_3}^{j, n}$, with

$$M_{m_3, m_4}^{j, n} = c_{m_3}^j c_{m_4}^j \begin{cases} \frac{i^{m_3-m_4} (-1)^{2j} 2^{-j}}{(j-m_3)! (m_3-m_4)! (j+m_4)!} {}_2F_1\left(\begin{smallmatrix} -j+m_3, -j-m_4 \\ 1+m_3-m_4 \end{smallmatrix}; -1\right) & m_3 > m_4 \\ \frac{i^{m_4-m_3} (-1)^{2j} 2^{-j}}{(j+m_3)! (m_4-m_3)! (j-m_4)!} {}_2F_1\left(\begin{smallmatrix} -j-m_3, -j+m_4 \\ 1-m_3+m_4 \end{smallmatrix}, -1\right) & m_3 \leq m_4 \end{cases}$$

then the operators $A_i(\lambda)$ act as

$$(6.4) \quad A_1(\lambda)W_{m_1, m_2}^{(j, n)} = \sum_{m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)} S_{m_3, m_2}^{j, n} \left(\frac{\lambda_1 - \lambda_2 + 1}{2} \right) W_{m_1, m_3}^{(j, n)}$$

$$(6.5) \quad A_2(\lambda)W_{m_1, m_2}^{(j, n)} = T_{m_2}^n \left(\frac{\lambda_1 + 1}{2} \right) W_{m_1, m_2}^{(j, n)}$$

$$(6.6) \quad A_3(\lambda)W_{m_1, m_2}^{(j, n)} = \sum_{m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)} S_{m_3, m_2}^{j, n} \left(\frac{\lambda_1 + \lambda_2 + 1}{2} \right) W_{m_1, m_3}^{(j, n)}$$

$$(6.7) \quad A_4(\lambda)W_{m_1, m_2}^{(j, n)} = T_{m_2}^n \left(\frac{\lambda_2 + 1}{2} \right) W_{m_1, m_2}^{(j, n)}.$$

We will prove the Proposition 6.1 in the following two sections 6.1 and 6.2.

Remark 6.1. The simple reflection w_{α_1} sends the character $\delta = (\delta_1, \delta_2)$ on M to (δ_2, δ_1) . For the w_{α_1} intertwining operator, it is important to recall that the parity condition $\mathbb{M}(j, n; \delta_2, \delta_1)$ of m_3 restricts the allowed Wigner D -functions $W_{m_1, m_3}^{(j, n)}$ in $I_{P_0}(w_{\alpha_1} \chi_{\delta, \lambda})$ as in (4.7) and (4.8).

6.1. Rank 1 Intertwining Operators. Starting from any character λ on $\mathfrak{a}_{\mathbb{C}}$, the rank 1 intertwining operator $A(w_{\alpha}, \mu)$ associated to a simple reflection w_{α} can be written as

$$A(w_{\alpha}, \lambda)f(k) = \int_{\overline{N} \cap w_{\alpha}^{-1} N w_{\alpha}} f(kw_{\alpha} \bar{n}) d\bar{n} = \int_{-\infty}^{\infty} f(kw_{\alpha} \exp(tX_{-\alpha})) dt.$$

By the Iwasawa decomposition of $\exp(tX_{-\alpha})$ given in (3.1),

$$\exp(tX_{-\alpha}) = \kappa_{\alpha}(t) h_{\alpha}(\sqrt{1+t^2}) \chi_{\alpha} \left(\frac{t}{1+t^2} \right),$$

where $\kappa_{\alpha}(t) = \exp(\arctan(-t)(X_{\alpha} - X_{-\alpha}))$. If f is any vector in the principal series representation $I(\chi_{\delta, \lambda})$, the action of $\exp(tX_{-\alpha})$ on the right is

$$\begin{aligned} f(kw_{\alpha} \exp(tX_{-\alpha})) &= f \left(kw_{\alpha} \kappa_{\alpha}(t) h_{\alpha}(\sqrt{1+t^2}) \chi_{\alpha} \left(\frac{t}{1+t^2} \right) \right) \\ &= (1+t^2)^{-\frac{(\tilde{\alpha}, \lambda + \rho)}{2}} f \left(kw_{\alpha} e^{\arctan(-t)(X_{\alpha} - X_{-\alpha})} \right). \end{aligned}$$

Denoting by $\theta(t) = \arctan(-t)$, since $X_{\alpha_1} - X_{-\alpha_1} = 2U_2$, $X_{\alpha_2} - X_{-\alpha_2} = U_0 - U_3$, and recalling from (2.5) the expressions of simple reflections w_{α_i} in terms of Euler angles, the action of $\exp(tX_{-\alpha_1})$ and $\exp(tX_{-\alpha_2})$ on a Wigner D -function $f = W_{m_1, m_2}^{(j, n)}$ is thus

$$(6.8) \quad \begin{aligned} &W_{m_1, m_2}^{(j, n)}(kw_{\alpha_1} e^{\theta(t)(X_{\alpha_1} - X_{-\alpha_1})}) \\ &= \sum_{-j \leq m_3 \leq j} W_{m_1, m_3}^{(j, n)}(k) W_{m_3, m_2}^{(j, n)}(0, 0, -\pi - 2\theta(t), 0) \end{aligned}$$

$$(6.9) \quad \begin{aligned} &W_{m_1, m_2}^{(j, n)}(kw_{\alpha_2} e^{\theta(t)(X_{\alpha_2} - X_{-\alpha_2})}) \\ &= \sum_{-j \leq m_3 \leq j} W_{m_1, m_3}^{(j, n)}(k) W_{m_3, m_2}^{(j, n)} \left(-\frac{\pi}{2} - \theta(t), \frac{\pi}{2} + \theta(t), 0, 0 \right). \end{aligned}$$

Therefore, the simple intertwining operators $A(w_{\alpha_1}, \lambda)W_{m_1, m_2}^{(j, n)}$ and $A(w_{\alpha_2}, \lambda)W_{m_1, m_2}^{(j, n)}$ can be expressed in terms of integrals involving Wigner D -functions:

$$(6.10) \quad \left(A(w_{\alpha_1}, \lambda)W_{m_1, m_2}^{(j, n)} \right) (k) =$$

$$\sum_{-j \leq m_3 \leq j} W_{m_1, m_3}^{(j, n)}(k) \int_{-\infty}^{\infty} (1+t^2)^{-\frac{\langle \alpha_1, \lambda + \rho \rangle}{2}} W_{m_3, m_2}^{(j, n)}(0, 0, -\pi - 2\theta(t), 0) dt$$

$$(6.11) \quad \left(A(w_{\alpha_2}, \lambda)W_{m_1, m_2}^{(j, n)} \right) (k) =$$

$$\sum_{-j \leq m_3 \leq j} W_{m_1, m_3}^{(j, n)}(k) \int_{-\infty}^{\infty} (1+t^2)^{-\frac{\langle \alpha_2, \lambda + \rho \rangle}{2}} W_{m_3, m_2}^{(j, n)}\left(-\frac{\pi}{2} - \theta(t), \frac{\pi}{2} + \theta(t), 0, 0\right) dt$$

Observe that $\theta(t)$ is an odd function. It is important to mention that, from the formula of the Wigner D -function (A.3), we have

$$d_{m_3, m_2}^{(j, n)}(-\pi - 2\theta(t)) = \sum_{p=\max(0, m_3-m_2)}^{\min(j-m_2, j+m_3)} \frac{(-1)^{2j+m_2-m_3+p}}{(j+m_3-p)!p!(m_2-m_3+p)!(j-m_2-p)!} \cos^{m_2-m_3+2p}(\theta(t)) \sin^{2j+m_3-m_2-2p}(\theta(t)).$$

If $2j+m_3-m_2 \equiv 1 \pmod{2}$, the integrand of (6.10) is an odd function, which makes the integral (6.10) zero. If $W_{m_1, m_2}^{(j, n)} \in I_{P_0}(\chi_{\delta, \lambda})$, we must have

$$2j \equiv \delta_1 + \delta_2 \pmod{2}$$

and

$$m_2 \in \mathbb{M}(j, n; \delta_1, \delta_2).$$

The set $\mathbb{M}(j, n; \delta_1, \delta_2)$ has been defined in (4.6). In order to make the integral (6.10) nonzero, the function $d_{m_3, m_2}^{(j, n)}(-\pi - 2\theta(t))$ must be an even function. In this case, the exponent $2j+m_3-m_2 \equiv 0 \pmod{2}$. Therefore,

$$(6.12) \quad 2j+m_3-m_2 \equiv \delta_1 + \delta_2 + m_3 - m_2 \equiv 0 \pmod{2}.$$

Thus, if m_2 satisfies $n - m_2 \equiv \delta_1 \pmod{2}$ and $n + m_2 \equiv \delta_2 \pmod{2}$, by (6.12), we must have

$$n - m_3 \equiv n - m_2 + \delta_1 + \delta_2 \equiv \delta_2 \pmod{2}$$

$$n + m_3 \equiv n + m_2 + \delta_1 + \delta_2 \equiv \delta_1 \pmod{2}.$$

Thus the parity condition m_3 is given by the set $\mathbb{M}(j, n; \delta_2, \delta_1)$, with δ_1 and δ_2 flipped from the parity condition of m_2 . In particular, $A(w_{\alpha_1}, \lambda)W_{m_1, m_2}^{(j, n)}$ lies in the space $I_{P_0}(w_{\alpha_1}\chi_{\delta, \lambda})$. Hence the sum in (6.10) is in fact a sum over $m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)$:

$$(6.13) \quad \left(A(w_{\alpha_1}, \lambda)W_{m_1, m_2}^{(j, n)} \right) (k) =$$

$$\sum_{m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)} W_{m_1, m_3}^{(j, n)}(k) \int_{-\infty}^{\infty} (1+t^2)^{-\frac{\langle \alpha_1, \lambda + \rho \rangle}{2}} W_{m_3, m_2}^{(j, n)}(0, 0, -\pi - 2\theta(t), 0) dt.$$

Note that the Wigner D -function $W_{m_3, m_2}^{(j, n)}(0, 0, -\pi - 2\theta(t), 0)$ in (6.8) and (6.13) has a nonzero U_2 -Euler angle. In the following section, we will diagonalize the matrix

of Wigner D -functions and transform the U_2 –Euler angle to a U_3 –Euler angle in order to compute the intertwining operators more easily.

6.2. Diagonalization of the intertwining operators. From the commutation relation of Pauli matrices, U_2 and U_3 can be related in the following way:

$$(6.14) \quad U_2 = \text{Ad}(e^{-\frac{3\pi}{2}U_1})U_3.$$

We can use this relation to diagonalize the action of

$$e^{(\pi+2\theta(t))U_2} = \text{Ad}(e^{-\frac{3\pi}{2}U_1})e^{(\pi+2\theta(t))U_3}$$

appearing in the Iwasawa decomposition of $\exp(tX_{-\alpha_1})$ of the Wigner D -functions. By the multiplicativity of the Wigner D -function, we have

$$\begin{aligned} & W_{m_1, m_2}^{(j, n)}(k\omega_{\alpha_1} e^{2\theta(t)U_2}) = W_{m_1, m_2}^{(j, n)}(ke^{(\pi+2\theta(t))U_2}) \\ &= \sum_{m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)} W_{m_1, m_3}^{(j, n)}(k) W_{m_3, m_2}^{(j, n)}(e^{-\frac{3\pi}{2}U_1} e^{(\pi+2\theta(t))U_3} \exp \frac{3\pi}{2}U_1) \\ &= \sum_{m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)} W_{m_1, m_3}^{(j, n)}(k) \sum_{m_4, m_5} W_{m_3, m_4}^{(j, n)}(e^{-\frac{3\pi}{2}U_1}) W_{m_5, m_2}^{(j, n)}(e^{\frac{3\pi}{2}U_1}) W_{m_4, m_5}^{(j, n)}(e^{(\pi+2\theta(t))U_3}) \\ &= \sum_{m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)} W_{m_1, m_3}^{(j, n)}(k) \sum_{m_4, m_5} W_{m_3, m_4}^{(j, n)}(e^{-\frac{3\pi}{2}U_1}) W_{m_5, m_2}^{(j, n)}(e^{\frac{3\pi}{2}U_1}) \times \\ & (6.15) \quad W_{m_4, m_5}^{(j, n)}(0, -\pi - 2\theta(t), 0, 0). \end{aligned}$$

We define the function $S_{m_3, m_2}'^{(j, n)}(z)$ by

$$(6.16) \quad \begin{aligned} S_{m_3, m_2}'^{(j, n)}(z) &= \sum_{m_4, m_5} W_{m_3, m_4}^{(j, n)}(e^{-\frac{3\pi}{2}U_1}) W_{m_5, m_2}^{(j, n)}(e^{\frac{3\pi}{2}U_1}) \\ & \int_{-\infty}^{\infty} (1+t^2)^{-z} W_{m_4, m_5}^{(j, n)}(0, -\pi - 2\theta(t), 0, 0). \end{aligned}$$

Change-of-basis matrix. Similarly to the relation (6.14) between U_2 and U_3 , we can relate U_1 and U_2 by conjugating a multiple of U_3 :

$$(6.17) \quad U_1 = \text{Ad}(e^{\frac{\pi}{2}U_3})U_2.$$

Recalling the notation from (A.2) and (A.3) that $c_m^j = \sqrt{(j+m)!(j-m)!}$, the change-of-basis matrices $W_{m_3, m_4}^{(j, n)}(e^{-\frac{3\pi}{2}U_1})$ and $W_{m_5, m_2}^{(j, n)}(e^{\frac{3\pi}{2}U_1})$ can be expressed in terms of the value of a Wigner D -function:

$$(6.18) \quad \begin{aligned} & W_{m_3, m_4}^{(j, n)}(e^{-\frac{3\pi}{2}U_1}) = W_{m_3, m_4}^{(j, n)}(e^{\frac{\pi}{2}U_3} e^{-\frac{3\pi}{2}U_2} e^{-\frac{\pi}{2}U_3}) = W_{m_3, m_4}^{(j, n)}\left(-\frac{\pi}{2}, \frac{3\pi}{2}, \frac{\pi}{2}\right) \\ &= c_{m_3}^j c_{m_4}^j \begin{cases} \frac{i^{m_3-m_4} (-1)^{2j} 2^{-j}}{(j-m_3)!(m_3-m_4)!(j+m_4)!} {}_2F_1\left(\begin{smallmatrix} -j+m_3, -j-m_4 \\ 1+m_3-m_4 \end{smallmatrix}; -1\right) & m_3 > m_4 \\ \frac{i^{m_4-m_3} (-1)^{2j} 2^{-j}}{(j+m_3)!(m_4-m_3)!(j-m_4)!} {}_2F_1\left(\begin{smallmatrix} -j-m_3, -j+m_4 \\ 1-m_3+m_4 \end{smallmatrix}; -1\right) & m_3 \leq m_4 \end{cases}. \end{aligned}$$

We define

$$(6.19) \quad M_{m_3, m_4}^j = W_{m_3, m_4}^{(j, n)}(e^{-\frac{3\pi}{2}U_1})$$

$$(6.20) \quad N_{m_5, m_2}^j = W_{m_5, m_2}^{(j, n)}(e^{\frac{3\pi}{2}U_1}).$$

As we have seen in (A.4), the entries of M_{m_3, m_4}^j and N_{m_3, m_4}^j are related to values $P_{j-m_1}^{m_1-m_4, m_1+m_4}(0)$ of Jacobi polynomials $P_n^{\alpha, \beta}(x)$:

$$(6.21) \quad M_{m_3, m_4}^j = \frac{c_{m_4}^j}{c_{m_3}^j} (-1)^{2j} i^{-m_3+m_4} 2^{-m_4} P_{j-m_4}^{-m_3+m_4, m_3+m_4}(0)$$

$$(6.22) \quad N_{m_5, m_2}^j = \frac{c_{m_2}^j}{c_{m_5}^j} (-1)^{2j} i^{m_5-m_2} 2^{-m_2} P_{j-m_2}^{-m_5+m_2, m_5+m_2}(0).$$

After introducing these notations, the function $S_{m_3, m_2}^{\prime(j, n)}(z)$ defined in (6.16) becomes

$$(6.23) \quad S_{m_3, m_2}^{\prime(j, n)}(z) = \sum_{m_4, m_5} M_{m_3, m_4}^{(j, n)} N_{m_5, m_2}^{(j, n)} \int_{-\infty}^{\infty} (1+t^2)^{-z} W_{m_4, m_5}^{(j, n)}(0, -\pi - 2\theta(t), 0, 0) dt.$$

Among the two equivalent definitions (A.5) from [AS67] and (A.6) in [Res08], for the sake of simplicity in expressions we choose the definition (A.6) using hypergeometric functions

$$P_n^{\alpha, \beta}(x) = \binom{n+\alpha}{n} \left(\frac{x+1}{2} \right)^n {}_2F_1 \left(\begin{matrix} -n, -n-\beta \\ \alpha+1 \end{matrix}; \frac{x-1}{x+1} \right).$$

These Jacobi polynomials have a generating function [SM84]

$$(6.24) \quad \sum_{n=0}^{\infty} P_n^{\alpha-n, \beta-n}(x) t^n = \left(1 + \frac{1}{2}(x+1)t \right)^{\alpha} \left(1 + \frac{1}{2}(x-1)t \right)^{\beta}$$

for $|x| < 1$. The series on the left hand side converges absolutely for $|t| < 1$. For any meromorphic function $f(t)$, we denote $[f(t)]_0$ as the zeroth Laurent series coefficient of $f(t)$. The change-of-basis matrices $M_{m_3, m_4}^j, N_{m_5, m_2}^j$ can thus be expressed as the constant term in Laurent series,

$$(6.25) \quad M_{m_3, m_4}^j = \left[(-1)^{2j} \frac{c_{m_4}^j}{c_{m_3}^j} i^{-m_3+m_4} 2^{-m_4} t^{m_4-j} \left(1 + \frac{t}{2} \right)^{j-m_3} \left(1 - \frac{t}{2} \right)^{j+m_3} \right]_0$$

$$(6.26) \quad N_{m_5, m_2}^j = \left[(-1)^{2j} \frac{c_{m_2}^j}{c_{m_5}^j} i^{m_5-m_2} 2^{-m_2} t^{m_2-j} \left(1 + \frac{t}{2} \right)^{j-m_5} \left(1 - \frac{t}{2} \right)^{j+m_5} \right]_0.$$

The Singular Integrals. In (6.2), we have described the procedure of calculating the matrix entries

$$(6.27) \quad [A(w, \lambda)]_{m_3, m_2}^{(j, n)} = \langle W_{m_1, m_3}^{(j, n)}, A(w, \lambda) W_{m_1, m_2}^{(j, n)} \rangle$$

of the intertwining operator $A(w, \lambda)$. The right hand side of (6.27) is independent of m_1 . The calculation of simple intertwining operators reduces to the calculation of integrals (6.13) and (6.11). Combining the diagonalized operator (6.15) with (6.10) and (6.11), the problem of the calculation of the intertwining operators $A(w, \lambda)$ reduces to the following two integrals:

$$(6.28) \quad \int_{-\infty}^{\infty} (1+t^2)^{-\frac{\langle \alpha_1, \lambda + \rho \rangle}{2}} W_{m_4, m_5}^{(j, n)}(0, -\pi - 2\theta(t), 0, 0) dt$$

$$(6.29) \quad \int_{-\infty}^{\infty} (1+t^2)^{-\frac{\langle \alpha_2, \lambda + \rho \rangle}{2}} W_{m_3, m_2}^{(j, n)} \left(-\frac{\pi}{2} - \theta(t), \frac{\pi}{2} + \theta(t), 0, 0 \right) dt$$

Recall from the definition of Wigner D -functions and their values on Euler angles that

$$W_{m_1, m_2}^{(j, n)}(\zeta, \psi, 0, 0) = e^{im\zeta + im_1\psi} \delta_{m_1, m_2}.$$

The Wigner D -function part of the integrands of the above two integrals (6.28) and (6.29) are

$$\begin{aligned} W_{m_4, m_5}^{(j, n)}(0, -\pi - 2\theta(t), 0, 0) &= i^{-2m_4} (1 + it)^{m_4} (1 - it)^{-m_4} \delta_{m_4, m_5} \\ W_{m_3, m_2}^{(j, n)}\left(-\frac{\pi}{2} - \theta(t), \frac{\pi}{2} + \theta(t), 0, 0\right) &= i^{-n+m_2} (1 + it)^{\frac{n-m_2}{2}} (1 - it)^{-\frac{n-m_2}{2}} \delta_{m_3, m_2} \end{aligned}$$

The integral $\int_{-\infty}^{\infty} (1 + it)^{s_1} (1 - it)^{s_2} dt$ is convergent for $\text{Re}(s_1 + s_2) < -1$, and can be meromorphically continued to the whole complex plane as

$$(6.30) \quad \int_{-\infty}^{\infty} (1 + it)^{s_1} (1 - it)^{s_2} dt = \pi 2^{s_1+s_2+2} \frac{\Gamma(-s_1 - s_2 - 1)}{\Gamma(-s_1)\Gamma(-s_2)}.$$

We can use (6.30) to express the two integrals (6.28) and (6.29) in terms of Γ -functions. Thus the integral (6.28) becomes

$$\begin{aligned} & \int_{-\infty}^{\infty} (1 + t^2)^{-\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2}} W_{m_4, m_5}^{(j, n)}(0, -\pi - 2\theta(t), 0, 0) dt \\ &= i^{-2m_4} \delta_{m_4, m_5} \int_{-\infty}^{\infty} (1 - it)^{-\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2} - m_4} (1 + it)^{-\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2} + m_4} dt \\ &= i^{-2m_4} \pi 2^{-\langle \check{\alpha}_1, \lambda + \rho \rangle + 2} \frac{\Gamma(\langle \check{\alpha}_1, \lambda + \rho \rangle - 1)}{\Gamma\left(\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2} - m_4\right) \Gamma\left(\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2} + m_4\right)} \delta_{m_4, m_5} \end{aligned}$$

and the integral (6.29) is

$$\begin{aligned} & \int_{-\infty}^{\infty} (1 + t^2)^{-\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2}} W_{m_3, m_2}^{(j, n)}\left(-\frac{\pi}{2} - \theta(t), \frac{\pi}{2} + \theta(t), 0, 0\right) dt \\ &= i^{-n+m_2} \delta_{m_3, m_2} \int_{-\infty}^{\infty} (1 - it)^{-\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2} - \frac{n-m_2}{2}} (1 + it)^{-\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2} + \frac{n-m_2}{2}} dt \\ &= i^{-n+m_2} \pi 2^{-\langle \check{\alpha}_2, \lambda + \rho \rangle + 2} \frac{\Gamma(\langle \check{\alpha}_2, \lambda + \rho \rangle - 1)}{\Gamma\left(\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2} - \frac{n-m_2}{2}\right) \Gamma\left(\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2} + \frac{n-m_2}{2}\right)} \delta_{m_3, m_2}. \end{aligned}$$

We have defined the function $Q(z, n) = \frac{\pi 2^{2-2z} \Gamma(2z-1)}{\Gamma(z+n)\Gamma(z-n)}$ in (6.3), so the integrals (6.28) and (6.29) can be expressed as:

$$(6.31) \quad \begin{aligned} & \int_{-\infty}^{\infty} (1 + t^2)^{-\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2}} W_{m_4, m_5}^{(j, n)}(0, -\pi - 2\theta(t), 0, 0) dt \\ &= i^{-2m_4} Q\left(\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2}, m_4\right) \delta_{m_4, m_5} \end{aligned}$$

$$(6.32) \quad \begin{aligned} & \int_{-\infty}^{\infty} (1 + t^2)^{-\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2}} W_{m_3, m_2}^{(j, n)}\left(-\frac{\pi}{2} - \theta(t), \frac{\pi}{2} + \theta(t), 0, 0\right) dt \\ &= i^{-n+m_2} Q\left(\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2}, \frac{m_2 - n}{2}\right) \delta_{m_3, m_2}. \end{aligned}$$

If we denote by $T_{m_2}^n(z) = i^{-n+m_2} Q\left(z, \frac{m_2-n}{2}\right)$, the result of the integral in (6.32) is equal to $T_{m_2}^n\left(\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2}\right) = i^{-n+m_2} Q\left(\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2}, \frac{m_2-n}{2}\right)$. By collecting the results

from (6.18) and (6.31), it turns out that the $S_{m_3, m_2}'^{(j, n)}(z)$ defined in (6.16) and (6.23) is exactly the same as the $S_{m_3, m_2}^{(j, n)}(z)$ in Proposition 6.1:

$$\begin{aligned} S_{m_3, m_2}^{(j, n)}\left(\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2}\right) &= S_{m_3, m_2}'^{(j, n)}\left(\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2}\right) \\ &= \sum_{m_4} M_{m_3, m_4}^{(j, n)} N_{m_4, m_2}^{(j, n)} i^{-2m_4} Q\left(\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2}, m_4\right). \end{aligned}$$

Therefore, by (6.13), the simple intertwining operator $A(w_{\alpha_1}, \lambda)$ acts on Wigner D -functions by

$$A(w_{\alpha_1}, \lambda) W_{m_1, m_2}^{(j, n)} = \sum_{-j \leq m_3 \leq j} S_{m_3, m_2}^{(j, n)}\left(\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2}\right) W_{m_1, m_3}^{(j, n)}$$

From (6.11) and (6.32), we can see that the simple intertwining operator $A(w_{\alpha_2}, \lambda)$ acts by

$$A(w_{\alpha_2}, \lambda) W_{m_1, m_2}^{(j, n)} = T_{m_2}^n \left(\frac{\langle \check{\alpha}_2, \lambda + \rho \rangle}{2} \right) W_{m_1, m_2}^{(j, n)}.$$

By the Langlands' lemma (6.1)

$$A(w_0, \lambda) = A(w_{\alpha_2}, w_{\alpha_1} w_{\alpha_2} w_{\alpha_1} \lambda) A(w_{\alpha_1}, w_{\alpha_2} w_{\alpha_1} \lambda) A(w_{\alpha_2}, w_{\alpha_1} \lambda) A(w_{\alpha_1}, \lambda),$$

we can replace λ in the formula above by $w_{\alpha_1} \lambda$, $w_{\alpha_2} w_{\alpha_1} \lambda$ and $w_{\alpha_1} w_{\alpha_2} w_{\alpha_1} \lambda$ to calculate the other stages in the composition of simple intertwining operators. Recall that if we express λ by the pair (λ_1, λ_2) as in Section 6.1.1, where $\lambda_i = \lambda(H_i)$, the composition of actions by the Weyl group action will send the pair (λ_1, λ_2) to:

$$(\lambda_1, \lambda_2) \xrightarrow{w_{\alpha_1}} (\lambda_2, \lambda_1) \xrightarrow{w_{\alpha_2}} (\lambda_2, -\lambda_1) \xrightarrow{w_{\alpha_1}} (-\lambda_1, \lambda_2) \xrightarrow{w_{\alpha_2}} (-\lambda_1, -\lambda_2).$$

The pairing of $\lambda + \rho$ with the coroots $\check{\alpha}_1$ and $\check{\alpha}_2$ are given by

$$\begin{aligned} \langle \check{\alpha}_1, \lambda + \rho \rangle &= \lambda_1 - \lambda_2 + 1 \\ \langle \check{\alpha}_2, \lambda + \rho \rangle &= \lambda_2 + 1 \end{aligned}$$

Therefore, we can conclude that the matrix coefficients for the simple intertwining operators are

$$\begin{aligned} A(w_{\alpha_1}, \lambda) W_{m_1, m_2}^{(j, n)} &= \sum_{m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)} S_{m_3, m_2}^{(j, n)}\left(\frac{\langle \check{\alpha}_1, \lambda + \rho \rangle}{2}\right) W_{m_1, m_3}^{(j, n)} \\ A(w_{\alpha_2}, w_{\alpha_1} \lambda) W_{m_1, m_2}^{(j, n)} &= T_{m_2}^n \left(\frac{\langle \check{\alpha}_2, w_{\alpha_1} \lambda + \rho \rangle}{2} \right) W_{m_1, m_2}^{(j, n)} \\ A(w_{\alpha_1}, w_{\alpha_2} w_{\alpha_1} \lambda) W_{m_1, m_2}^{(j, n)} &= \sum_{m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)} S_{m_3, m_2}^{(j, n)}\left(\frac{\langle \check{\alpha}_1, w_{\alpha_2} w_{\alpha_1} \lambda + \rho \rangle}{2}\right) W_{m_1, m_3}^{(j, n)} \\ A(w_{\alpha_2}, w_{\alpha_1} w_{\alpha_2} w_{\alpha_1} \lambda) W_{m_1, m_2}^{(j, n)} &= T_{m_2}^n \left(\frac{\langle \check{\alpha}_2, w_{\alpha_1} w_{\alpha_2} w_{\alpha_1} \lambda + \rho \rangle}{2} \right) W_{m_1, m_2}^{(j, n)} \end{aligned}$$

Collecting the above result with the change-of-basis matrix defined in (6.18), we have finished the proof of the Proposition 6.1.

6.3. Expression of $S_{m_1, m_4}^{j, n}(z)$ as Hypergeometric Functions. In this section, we assume

$$(\delta_1, \delta_2) = (0, 0) \text{ or } (1, 1),$$

in which case j, n are integers. We also introduce the rising factorial or the *Pochhammer symbol*

$$(6.33) \quad (a)^{(n)} = \frac{\Gamma(a+n)}{\Gamma(a)}$$

Definition 6.1. *Part I of [Obe12].*

(1) *The Mellin transform of a function $f(x)$ is formally defined as*

$$(6.34) \quad \mathcal{M}(f(x))(z) = \int_0^\infty f(x)x^{z-1}dx.$$

(2) *Consider a function $F(z)$ of one complex variable $z = \sigma + i\tau$, such that*

(a) *$F(z)$ is holomorphic on the strip $S = \{z \in \mathbb{C} | a < \sigma < b\}$, such that $F(z) \rightarrow 0$ uniformly in the strip $S_\epsilon = \{z \in \mathbb{C} | a + \epsilon < \sigma < b - \epsilon\}$ for arbitrarily small $\epsilon > 0$, and*

(b) *$\int_{-\infty}^\infty |F(\sigma + i\tau)|d\tau < \infty$ for all $\sigma \in (a, b)$.*

We define the inverse Mellin transform of the function $F(z)$ as

$$(6.35) \quad \mathcal{M}^{-1}(F(z))(x) = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} F(z)x^{-z}dz,$$

for $x > 0$ and some fixed $\gamma \in (a, b)$. It satisfies the property that

$$(6.36) \quad \mathcal{M}(\mathcal{M}^{-1}(F))(z) = F(z).$$

The function $F(z)$ to our interest is the ratio of Γ -functions $\frac{\Gamma(z)}{\Gamma(\frac{1+z-\nu}{2})\Gamma(\frac{1+z+\nu}{2})}$. We would like to calculate the integral

$$(6.37) \quad \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{\Gamma(z)}{\Gamma(\frac{1+z-\nu}{2})\Gamma(\frac{1+z+\nu}{2})} x^{-z} dz$$

along some well chosen contour on which the integral converges. Though this integral doesn't satisfy the assumption on $F(z)$ above, we still can take the inverse Mellin transform of this function, resulting in a $\mathcal{M}^{-1}(F(z))(x)$ with discontinuities. We refer to [Bat55, Page 49, Section 1.19] for a general *Mellin-Barnes integral*

$$(6.38) \quad \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{\prod_{i=1}^m \Gamma(a_i + A_i z) \prod_{j=1}^n \Gamma(b_j - B_j z)}{\prod_{s=1}^p \Gamma(c_s + C_s z) \prod_{t=1}^q \Gamma(d_t - D_t z)} x^{-z} dz,$$

where $\gamma \in \mathbb{R}$ and $A_i, B_j, C_s, D_t > 0$. If $x \in \mathbb{R}$, the integrand is asymptotically $e^{-\frac{1}{2}\alpha\pi|y|}|y|^{\beta\gamma+\lambda}|\frac{x}{\rho}|^{-\gamma}$, where the numbers $\alpha, \beta, \lambda, \rho$ are defined by

$$\begin{aligned}\alpha &= \sum_{i=1}^m A_i + \sum_{j=1}^n B_j - \sum_{s=1}^p C_s - \sum_{t=1}^q D_t \\ \beta &= \sum_{i=1}^m A_i - \sum_{j=1}^n B_j - \sum_{s=1}^p C_s + \sum_{t=1}^q D_t \\ \lambda &= \operatorname{Re}\left(\sum_{i=1}^m a_i + \sum_{j=1}^n b_j - \sum_{s=1}^p c_s - \sum_{t=1}^q d_j\right) - \frac{m+n-p-q}{2} \\ \rho &= \prod_{i=1}^m A_i^{A_i} \prod_{j=1}^n B_j^{-B_j} \prod_{s=1}^p C_s^{-C_s} \prod_{t=1}^q D_t^{D_t},\end{aligned}$$

and $y = \operatorname{Im}(z)$. Our integral (6.37) satisfies $\alpha = \beta = 0$ and $\lambda = -\frac{1}{2}$, $\rho = 2$, falling into the fourth type in the description on the convergence of the Mellin-Barnes integral in [Bat55], which states that the integral (6.37) conditionally converges to an analytic function in x on the intervals $|x| < 2$ and $|x| > 2$, with points of discontinuity at $x = \pm 2$. In [Obe12, Section II, (5.21)], for $\operatorname{Re}(z) > 0$, we have an explicit formula for (6.37):

$$(6.39) \quad \mathcal{M}^{-1}\left(\frac{\Gamma(z)}{\Gamma\left(\frac{1+z-\nu}{2}\right)\Gamma\left(\frac{1+z+\nu}{2}\right)}\right)(x) = \begin{cases} 2\pi^{-1}(4-x^2)^{-1/2} \cos(\nu \arccos(x/2)) & |x| < 2 \\ 0 & |x| > 2 \end{cases}$$

In the rest of this chapter, we will only interchange finite sums with this conditionally convergent integral, and thus there are no analytic issues justifying the calculation in the remaining portion of this thesis. In (6.3) we have defined the function $Q(z, n) = \frac{\pi 2^{2-2z} \Gamma(2z-1)}{\Gamma(z+n) \Gamma(z-n)}$. We would like to compute the inverse Mellin transform $\mathcal{M}^{-1}(Q(z, m_3))(x)$ of $Q(z, m_3)$:

$$\begin{aligned}(6.40) \quad \mathcal{M}^{-1}\left(\frac{\pi 2^{2-2z} \Gamma(2z-1)}{\Gamma(z+m_3) \Gamma(z-m_3)}\right)(x) &= \frac{4\pi}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{\Gamma(2z-1)}{\Gamma(z+m_3) \Gamma(z-m_3)} (4x)^{-z} dz \\ &= \frac{2\pi}{2\pi i} \int_{2\gamma-1-i\infty}^{2\gamma-1+i\infty} \frac{\Gamma(z)}{\Gamma\left(\frac{z+1+2m_3}{2}\right) \Gamma\left(\frac{z+1-2m_3}{2}\right)} (4x)^{-\frac{z+1}{2}} dz \\ &= \frac{\pi}{2\pi i \sqrt{x}} \int_{2\gamma-1-i\infty}^{2\gamma-1+i\infty} \frac{\Gamma(z)}{\Gamma\left(\frac{z+1+2m_3}{2}\right) \Gamma\left(\frac{z+1-2m_3}{2}\right)} (2\sqrt{x})^{-z} dz \\ &= \frac{\pi}{\sqrt{x}} \mathcal{M}^{-1}\left(\frac{\Gamma(z)}{\Gamma\left(\frac{1+z-2m_3}{2}\right) \Gamma\left(\frac{1+z+2m_3}{2}\right)}\right)(2\sqrt{x}) \\ &= \frac{\pi}{\sqrt{x}} \begin{cases} \pi^{-1}(1-x)^{-1/2} \cos(2m_3 \arccos(\sqrt{x})) & 0 < x < 1 \\ 0 & x > 1 \end{cases} \\ &= \frac{1}{\sqrt{x(1-x)}} \begin{cases} \cos(2m_3 \arcsin(\sqrt{1-x})) & |x| < 1 \\ 0 & |x| > 1 \end{cases}.\end{aligned}$$

If we define the *Heaviside step function* $\theta(x) = \begin{cases} 0 & x \leq 0 \\ 1 & x > 0 \end{cases}$, the result above can be written as

$$\begin{aligned} \mathcal{M}^{-1}(Q(z, m_3))(x) &= \frac{1 - \theta(|x| - 1)}{\sqrt{(1-x)x}} \cos(2m_3 \arcsin(\sqrt{1-x})) \\ (6.41) \quad &= \frac{1 - \theta(|x| - 1)}{\sqrt{(1-x)x}} \left(\frac{1}{2} (\sqrt{x} + i\sqrt{1-x})^{-2m_3} + \frac{1}{2} (\sqrt{x} + i\sqrt{1-x})^{2m_3} \right) \end{aligned}$$

Therefore, the inverse Mellin transform $\mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x)$ of the matrix entries of intertwining operators $S_{m_3, m_2}^{j, n}(z)$ is

$$\begin{aligned} \mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x) &= \frac{1 - \theta(|x| - 1)}{2\sqrt{(1-x)x}} \sum_{m_3=-j}^j M_{m_1, m_3}^j N_{m_3, m_4}^j \\ (6.42) \quad & i^{-2m_3} \left((\sqrt{x} + i\sqrt{1-x})^{-2m_3} + (\sqrt{x} + i\sqrt{1-x})^{2m_3} \right). \end{aligned}$$

By (6.25) and (6.26), the change-of-basis matrix M_{m_1, m_3}^j and N_{m_1, m_3}^j are the zeroth Laurent coefficient of some analytic functions. By substituting the matrix entries M_{m_1, m_3}^j and N_{m_1, m_3}^j in (6.42) with the corresponding Laurent series (6.25) and (6.26), the sum over m_3 from $-j$ to j as in (6.42) actually gives rise to a geometric series. Recalling formulas (6.25) and (6.26) and defining

$$\begin{aligned} M_{m_1, m_3}^j(t) &= (-1)^{2j} \frac{c_{m_3}^j}{c_{m_1}^j} i^{-m_1+m_3} 2^{-m_3} t^{m_3-j} \left(1 + \frac{t}{2}\right)^{j-m_1} \left(1 - \frac{t}{2}\right)^{j+m_1} \\ N_{m_3, m_4}^j(s) &= (-1)^{2j} \frac{c_{m_4}^j}{c_{m_3}^j} i^{m_3-m_4} 2^{-m_4} s^{m_4-j} \left(1 + \frac{s}{2}\right)^{j-m_3} \left(1 - \frac{s}{2}\right)^{j+m_3}, \end{aligned}$$

where s, t are formal variables, we define

$$\begin{aligned} \sigma_1(s, t) &= \sum_{m_3=-j}^j M_{m_1, m_3}^j(t) N_{m_3, m_4}^j(s) i^{-2m_3} (\sqrt{x} + i\sqrt{1-x})^{-2m_3} \\ &= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-m_4} \left(1 - \frac{t}{2}\right)^{j+m_1} \left(1 + \frac{t}{2}\right)^{j-m_1} s^{m_4-j} t^{-j} \left(1 + \frac{s}{2}\right)^j \left(1 - \frac{s}{2}\right)^j \times \\ &\quad \sum_{m_3=-j}^j \left(\frac{t(1-\frac{s}{2})}{2(1+\frac{s}{2})} \right)^{m_3} (\sqrt{x} + i\sqrt{1-x})^{-2m_3} \\ &= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-m_4} \left(1 - \frac{t}{2}\right)^{j+m_1} \left(1 + \frac{t}{2}\right)^{j-m_1} s^{m_4-j} t^{-j} \left(1 + \frac{s}{2}\right)^j \left(1 - \frac{s}{2}\right)^j \times \\ (6.43) \quad & \frac{\left(\frac{(1-s/2)t}{2(1+s/2)(i\sqrt{1-x}+\sqrt{x})^2} \right)^{-j} - \left(\frac{(1-s/2)t}{2(1+s/2)(i\sqrt{1-x}+\sqrt{x})^2} \right)^{j+1}}{1 - \frac{(1-s/2)t}{2(1+s/2)(i\sqrt{1-x}+\sqrt{x})^2}} \end{aligned}$$

and

$$\begin{aligned}
\sigma_2(s, t) &= \sum_{m_3=-j}^j M_{m_1, m_3}^j(t) N_{m_3, m_4}^j(s) i^{-2m_3} (\sqrt{x} + i\sqrt{1-x})^{2m_3} \\
&= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-m_4} \left(1 - \frac{t}{2}\right)^{j+m_1} \left(1 + \frac{t}{2}\right)^{j-m_1} s^{m_4-j} t^{-j} \left(1 + \frac{s}{2}\right)^j \left(1 - \frac{s}{2}\right)^j \times \\
&\quad \sum_{m_3=-j}^j \left(\frac{t(1-\frac{s}{2})}{2(1+\frac{s}{2})}\right)^{m_3} (\sqrt{x} + i\sqrt{1-x})^{2m_3} \\
&= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-m_4} \left(1 - \frac{t}{2}\right)^{j+m_1} \left(1 + \frac{t}{2}\right)^{j-m_1} s^{m_4-j} t^{-j} \left(1 + \frac{s}{2}\right)^j \left(1 - \frac{s}{2}\right)^j \times \\
(6.44) \quad &\frac{\left(\frac{(1-s/2)t(i\sqrt{1-x}+\sqrt{x})^2}{2(1+s/2)}\right)^{-j} - \left(\frac{(1-s/2)t(i\sqrt{1-x}+\sqrt{x})^2}{2(1+s/2)}\right)^{j+1}}{1 - \frac{(1-s/2)t(i\sqrt{1-x}+\sqrt{x})^2}{2(1+s/2)}}.
\end{aligned}$$

If we denote by $[\sigma_i(s, t)]_{0,0}$ for the constant term in the Laurent series expansion of $\sigma_i(s, t)$, then (6.42) can be expressed as

$$(6.45) \quad \mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x) = \frac{1 - \theta(|x| - 1)}{2\sqrt{(1-x)x}} ([\sigma_1(s, t) + \sigma_2(s, t)]_{0,0}).$$

To make the calculation simpler, and since we only care about the *constant* term of the Laurent series $\sigma_i(s, t)$, we can feel free to make change of variables so that $\sigma_1(s, t)$ and $\sigma_2(s, t)$ have the same denominators. Two such choices are

$$\begin{aligned}
&\sigma_1(-s, 2/t) \\
&= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-j-m_4} (-1)^{m_4-j} \left(1 - \frac{1}{t}\right)^{j+m_1} \left(1 + \frac{1}{t}\right)^{j-m_1} s^{m_4-j} t^j \left(1 + \frac{s}{2}\right)^j \left(1 - \frac{s}{2}\right)^j \times \\
&\quad \frac{\left(\frac{(1+s/2)}{t(1-s/2)(i\sqrt{1-x}+\sqrt{x})^2}\right)^{-j} - \left(\frac{(1+s/2)}{t(1-s/2)(i\sqrt{1-x}+\sqrt{x})^2}\right)^{j+1}}{1 - \frac{(1+s/2)}{t(1-s/2)(i\sqrt{1-x}+\sqrt{x})^2}} \\
&= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-j-m_4} (-1)^{m_4-j} (t-1)^{j+m_1} (t+1)^{j-m_1} s^{m_4-j} t^{-j} \left(1 + \frac{s}{2}\right)^j \left(1 - \frac{s}{2}\right)^j \times \\
(6.46) \quad &\frac{\left(\frac{(1-s/2)t(i\sqrt{1-x}+\sqrt{x})^2}{(1+s/2)}\right)^{-j} - \left(\frac{(1-s/2)t(i\sqrt{1-x}+\sqrt{x})^2}{(1+s/2)}\right)^{j+1}}{1 - \frac{(1-s/2)t(i\sqrt{1-x}+\sqrt{x})^2}{(1+s/2)}}
\end{aligned}$$

and

$$\begin{aligned}
 & \sigma_2(s, 2t) \\
 &= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-j-m_4} (-1)^{j+m_1} (t-1)^{j+m_1} (1+t)^{j-m_1} s^{m_4-j} t^{-j} \left(1 + \frac{s}{2}\right)^j \left(1 - \frac{s}{2}\right)^j \times \\
 (6.47) \quad & \frac{\left(\frac{(1-s/2)t(i\sqrt{1-x}+\sqrt{x})^2}{(1+s/2)}\right)^{-j} - \left(\frac{(1-s/2)t(i\sqrt{1-x}+\sqrt{x})^2}{(1+s/2)}\right)^{j+1}}{1 - \frac{(1-s/2)t(i\sqrt{1-x}+\sqrt{x})^2}{(1+s/2)}}
 \end{aligned}$$

After taking the constant Laurent series coefficient of the function $\sigma_1(-s, 2/t) + \sigma_2(s, 2t)$, we will get the following expression for $\mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))$:

$$\begin{aligned}
 \mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x) &= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-j-m_4} \frac{(-1)^{j+m_1} + (-1)^{m_4-j}}{2} \\
 & \frac{1 - \theta(|x| - 1)}{\sqrt{(1-x)x}} \left[\frac{1}{1 - (i\sqrt{1-x} + \sqrt{x})^2 \frac{(1-\frac{s}{2})t}{1+\frac{s}{2}}} \right. \\
 & \left(\left(1 + \frac{s}{2}\right)^{2j} (1-t)^{j+m_1} (1+t)^{j-m_1} s^{m_4-j} t^{-2j} (i\sqrt{1-x} + \sqrt{x})^{-2j} \right. \\
 & \left. - \left(1 - \frac{s}{2}\right)^{2j+1} (1-t)^{j+m_1} (1+t)^{j-m_1} \left(1 + \frac{s}{2}\right)^{-1} s^{m_4-j} t \right. \\
 (6.48) \quad & \left. \left. (i\sqrt{1-x} + \sqrt{x})^{2j+2} \right) \right]_{0,0}
 \end{aligned}$$

Notice that in the Laurent series expansion of the function in s, t in (6.48), the exponent of t in the Laurent expansion of the second term

$$\left(1 - \frac{s}{2}\right)^{2j+1} \left(1 - \frac{t}{2}\right)^{j+m_1} \left(1 + \frac{t}{2}\right)^{j-m_1} \left(1 + \frac{s}{2}\right)^{-1} s^{m_4-j} t$$

in the parentheses is always strictly greater than 0. Therefore it does not contribute to the zeroth Laurent series coefficient in the expansion of (6.48):

$$\begin{aligned}
 \mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x) &= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-j-m_4} \frac{(-1)^{j+m_1} + (-1)^{m_4-j}}{2} \\
 & \frac{1 - \theta(|x| - 1)}{\sqrt{(1-x)x}} \left[\frac{1}{1 - (i\sqrt{1-x} + \sqrt{x})^2 \frac{(1-\frac{s}{2})t}{1+\frac{s}{2}}} \right. \\
 & \left(\left(1 + \frac{s}{2}\right)^{2j} (1-t)^{j+m_1} (1+t)^{j-m_1} s^{m_4-j} t^{-2j} (i\sqrt{1-x} + \sqrt{x})^{-2j} \right) \Big]_{0,0} \\
 &= \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} 2^{-j-m_4} \frac{(-1)^{j+m_1} + (-1)^{m_4-j}}{2} \frac{1 - \theta(|x| - 1)}{\sqrt{(1-x)x}} \\
 & \left[\sum_{q \geq 0, q \in \mathbb{Z}} \left(\left(1 + \frac{s}{2}\right)^{2j-q} \left(1 - \frac{s}{2}\right)^q (1-t)^{j+m_1} (1+t)^{j-m_1} \times \right. \right. \\
 (6.49) \quad & \left. \left. s^{m_4-j} t^{q-2j} (i\sqrt{1-x} + \sqrt{x})^{2(q-j)} \right) \right]_{0,0},
 \end{aligned}$$

which expands to

$$(6.50) \quad \mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x) = \frac{c_{m_4}^j}{c_{m_1}^j} i^{-m_1-m_4} \frac{(-1)^{2j+m_1-m_4} + 1}{2} \frac{1 - \theta(|x| - 1)}{\sqrt{(1-x)x}} \times \\ \sum_{q, \kappa_1, \kappa_2 \in \mathbb{Z}} \binom{2j-q}{j-m_4-\kappa_1} \binom{q}{\kappa_1} \binom{j+m_1}{\kappa_2} \binom{j-m_1}{2j-q-\kappa_2} (-1)^{\kappa_1+\kappa_2} (i\sqrt{1-x} + \sqrt{x})^{2(q-j)}.$$

The summations over κ_1 and κ_2 can be expressed as hypergeometric values

$$(6.51) \quad \sum_{\kappa_1 \in \mathbb{Z}} \binom{2j-q}{\kappa_1} \binom{q}{j+m_4-\kappa_1} (-1)^{\kappa_1} = \binom{2j-q}{j-m_4} {}_2F_1\left(\begin{matrix} -q, -j+m_4 \\ 1+j-q+m_4 \end{matrix}; -1\right)$$

$$(6.52) \quad \sum_{\kappa_2 \in \mathbb{Z}} \binom{j+m_1}{\kappa_2} \binom{j-m_1}{2j-q-\kappa_2} (-1)^{\kappa_2} = \binom{j-m_1}{2j-q} {}_2F_1\left(\begin{matrix} -2j+q, -j-m_1 \\ 1-j+q-m_1 \end{matrix}; -1\right).$$

The sum in (6.50) becomes

$$(6.53) \quad \sum_{0 \leq q \leq 2j, q \in \mathbb{Z}} \binom{2j-q}{j-m_4} \binom{j-m_1}{2j-q} {}_2F_1\left(\begin{matrix} -q, -j+m_4 \\ 1+j-q+m_4 \end{matrix}; -1\right) {}_2F_1\left(\begin{matrix} -2j+q, -j-m_1 \\ 1-j+q-m_1 \end{matrix}; -1\right) (i\sqrt{1-x} + \sqrt{x})^{2(q-j)}.$$

Again, by writing the values of hypergeometric function ${}_2F_1$ at -1 in terms of Jacobi polynomials, we have

$$(6.54) \quad {}_2F_1\left(\begin{matrix} -q, -j+m_4 \\ 1+j-q+m_4 \end{matrix}; -1\right) = 2^{j-m_4} \frac{(j-m_4)!(j+m_4-q)!}{(2j-q)!} P_{j-m_4}^{j+m_4-q, -j+m_4+q}(0) \\ (6.55) \quad {}_2F_1\left(\begin{matrix} -2j+q, -j-m_1 \\ 1-j+q-m_1 \end{matrix}; -1\right) = 2^{j+m_1} \frac{(j+m_1)!(-j-m_1+q)!}{q!} P_{j+m_1}^{j-m_1-q, -j-m_1+q}(0),$$

the sum (6.50) is equal to

$$(6.56) \quad \sum_{0 \leq q \leq 2j, q \in \mathbb{Z}} 2^{2j+m_1-m_4} \frac{(j-m_1)!(j+m_1)!}{q!(2j-q)!} P_{j-m_4}^{j+m_4-q, -j+m_4+q}(0) P_{j+m_1}^{j-m_1-q, -j-m_1+q}(0) \times \\ (i\sqrt{1-x} + \sqrt{x})^{2(q-j)}.$$

Again by (6.24), (6.50) is the constant term of the Laurent series expansion of the following function in s and t

$$(6.57) \quad s^{m_4-j} t^{-j-m_1} \sum_{0 \leq q \leq 2j, q \in \mathbb{Z}} \frac{(j-m_1)!(j+m_1)!}{q!(2j-q)!} (1+s)^{2j-q} (1-s)^q (1-t)^{2j-q} \times \\ (1+t)^q (i\sqrt{1-x} + \sqrt{x})^{2(q-j)} \\ = s^{m_4-j} t^{-j-m_1} \frac{(j-m_1)!(j+m_1)!}{(2j)!} \times \\ \left((i\sqrt{1-x} + \sqrt{x})^{-1} (1+s)(1-t) + (i\sqrt{1-x} + \sqrt{x}) (1-s)(1+t) \right)^{2j}.$$

For simplicity of notations, we set $\xi = i\sqrt{1-x} + \sqrt{x}$. The parenthesis of (6.57) can be reorganized as

$$\begin{aligned}
 & (\xi^{-1}(1+s)(1-t) + \xi(1-s)(1+t))^{2j} \\
 &= (\xi + \xi^{-1}) + (\xi^{-1} - \xi)s - (\xi^{-1} - \xi)t - st(\xi + \xi^{-1})^{2j} \\
 (6.58) \quad &= \sum_{\substack{\nu_1, \nu_2, \nu_3, \nu_4 \\ \sum \nu_i = 2j}} \frac{(2j)!}{\nu_1! \nu_3! \nu_2! \nu_4!} (\xi + \xi^{-1})^{\nu_1 + \nu_4} (\xi^{-1} - \xi)^{\nu_3 + \nu_2} (-1)^{\nu_2 + \nu_4} s^{\nu_3 + \nu_4} t^{\nu_2 + \nu_4}.
 \end{aligned}$$

Therefore, the constant term of (6.57) can be rewritten into a sum involving multinomial coefficients, noting that $2j + m_1 - m_4 + \nu_1 - \nu_4 = 2j$:

$$\begin{aligned}
 & (-1)^{j+m_1} \sum_{\nu_1, \nu_4} \frac{(j-m_1)!(j+m_1)!}{\nu_1!(j-m_4-\nu_4)!\nu_4!(j+m_1-\nu_4)!} (\xi + \xi^{-1})^{\nu_1 + \nu_4} (\xi^{-1} - \xi)^{2j+m_1-m_4-2\nu_4} \\
 (6.59) \quad &= (-1)^{j+m_1} \sum_{\nu_1, \nu_4} \frac{(j-m_1)!(j+m_1)!}{\nu_1!(j-m_4-\nu_4)!\nu_4!(j+m_1-\nu_4)!} (\xi + \xi^{-1})^{\nu_1 + \nu_4} (\xi^{-1} - \xi)^{2j-\nu_1-\nu_4}.
 \end{aligned}$$

Reorganizing,

$$\begin{aligned}
 & \left(\frac{c_{m_1}^j}{c_{m_4}^j} \right)^2 (-1)^{j+m_1} \sum_{\nu_1, \nu_4} \frac{(j-m_4)!(j+m_4)!}{\nu_1!(j-m_4-\nu_4)!\nu_4!(j+m_4-\nu_1)!} (\xi + \xi^{-1})^{\nu_1 + \nu_4} (\xi^{-1} - \xi)^{2j-\nu_1-\nu_4} \\
 (6.60) \quad &= \left(\frac{c_{m_1}^j}{c_{m_4}^j} \right)^2 (-1)^{j+m_1} \sum_{\nu_1, \nu_4} \frac{(j-m_4)!(j+m_4)!}{\nu_1!(j-m_4-\nu_4)!\nu_4!(j+m_4-\nu_1)!} (\sqrt{x})^{\nu_1 + \nu_4} (-i\sqrt{1-x})^{2j-\nu_1-\nu_4}
 \end{aligned}$$

We can observe that the function in the summation above can be simplified to the zeroth Laurent coefficient of a function in one variable u :

$$\begin{aligned}
 & \sum_{\nu_1, \nu_4} \frac{(j-m_4)!(j+m_4)!}{\nu_1!(j-m_4-\nu_4)!\nu_4!(j+m_4-\nu_1)!} (\sqrt{x})^{\nu_1 + \nu_4} (-i\sqrt{1-x})^{2j-\nu_1-\nu_4} \\
 &= \left[u^{-j-m_1} \sum_{\nu_1, \nu_4} \binom{j+m_4}{\nu_1} \binom{j-m_4}{\nu_4} (-i\sqrt{1-x}u)^{j+m_4-\nu_1} (-i\sqrt{1-x})^{j-m_4-\nu_2} u^{\nu_4} (\sqrt{x})^{\nu_1 + \nu_4} \right]_0 \\
 (6.61) \quad &= [u^{-j-m_1} (\sqrt{x} - iu\sqrt{1-x})^{j+m_4} (u\sqrt{x} - i\sqrt{1-x})^{j-m_4}]_0
 \end{aligned}$$

Plugging the formula above back into (6.50),

$$\begin{aligned}
 & \mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x) = \frac{c_{m_1}^j}{c_{m_4}^j} i^{-m_1-m_4} \frac{(-1)^{2j+m_1-m_4} + 1}{2} \frac{1 - \theta(|x| - 1)}{\sqrt{(1-x)x}} \times \\
 (6.62) \quad & (-i)^{2j} (-1)^{j+m_1} [u^{-j-m_1} (\sqrt{x} + u\sqrt{1-x})^{j+m_4} (iu\sqrt{x} + \sqrt{1-x})^{j-m_4}]_0,
 \end{aligned}$$

and recall that $j, m_1, m_4 \in \mathbb{Z}$ and $m_1 - m_4$ is even, we finally have

$$\begin{aligned}
 & \mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x) = \frac{c_{m_1}^j}{c_{m_4}^j} \frac{((-1)^{2j+m_1-m_4} + 1)(-1)^{\frac{m_1-m_4}{2}} (1 - \theta(|x| - 1))}{2\sqrt{(1-x)x}} \\
 (6.63) \quad & \left[\frac{(u\sqrt{1-x} + i\sqrt{x})^{j+m_4} (\sqrt{1-x} + iu\sqrt{x})^{j-m_4}}{u^{j+m_1}} \right]_0.
 \end{aligned}$$

From the generating function of Jacobi polynomials which we introduced in (6.24), this inverse Mellin transform is in fact

$$(6.64) \quad \begin{aligned} & \mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x) \\ &= \frac{c_{m_1}^j}{c_{m_4}^j} \frac{((-1)^{2j+m_1} i^{2j} + (-1)^{j+m_4})(1 - \theta(|x| - 1))}{2(1-x)^{\frac{m_1+m_4+1}{2}} x^{-\frac{2j+m_1+m_4-1}{2}}} P_{j+m_1}^{(-m_1-m_4, -2j-1)} \left(\frac{2}{x} - 1 \right). \end{aligned}$$

We refer to formulas 05.06.26.0002.01 and 07.23.17.0055.01 of [Res08],

$$(6.65) \quad P_n^{(a, b)}(z) = \frac{\Gamma(a+n+1)}{\Gamma(a+1)\Gamma(n+1)} {}_2F_1 \left(\begin{matrix} -n, a+b+n+1 \\ a+1 \end{matrix}; \frac{1-z}{2} \right)$$

$$(6.66) \quad {}_2F_1(a, b; c; z) = (1-z)^{-a} {}_2F_1 \left(\begin{matrix} a, c-b \\ c \end{matrix}; \frac{z}{z-1} \right).$$

Applying these two formulas to (6.64), we get

$$(6.67) \quad \begin{aligned} & \mathcal{M}^{-1}(S_{m_1, m_4}^{j, n}(z))(x) \\ &= \frac{c_{m_1}^j}{c_{m_4}^j} \frac{((-1)^{2j} + (-1)^{m_4-m_1})(1 - \theta(|x| - 1))}{2(1-x)^{\frac{m_1+m_4+1}{2}} x^{\frac{m_1-m_4+1}{2}}} P_{j+m_1}^{(m_4-m_1, -m_1-m_4)} (1-2x) \end{aligned}$$

According to formula 9.43 in [Obe12], we can apply the Mellin transform $\mathcal{M}$ on the Jacobi polynomials,

$$(6.68) \quad \begin{aligned} & \mathcal{M} \left(\left\{ \begin{matrix} (b-x)^{\mu-1} P_n^{(\alpha, \beta)}(1-\gamma x) & x < b \\ 0 & x > b \end{matrix} \right\} (z) \right) \\ &= \frac{\Gamma(\alpha+n+1)\Gamma(\mu)\Gamma(z)b^{z+\mu-1}}{n!\Gamma(1+\alpha)\Gamma(\mu+z)} {}_3F_2 \left(\begin{matrix} -n, 1+n+\alpha+\beta, z \\ 1+\alpha, \mu+z \end{matrix}; \frac{1}{2}\gamma b \right) \end{aligned}$$

After applying the Mellin transform, the matrix entries $S_{m_1, m_4}^{j, n}(z)$ can be transformed to

$$(6.69) \quad \begin{aligned} & S_{m_1, m_4}^{j, n}(z) = \\ & \frac{(-1)^{2j} + (-1)^{m_4-m_1}}{2} \frac{c_{m_1}^j}{c_{m_4}^j} \frac{\Gamma\left(\frac{-m_1-m_4+1}{2}\right) \Gamma\left(z + \frac{-m_1+m_4-1}{2}\right)}{\Gamma(-m_1+m_4+1) \Gamma(z-m_1)} \\ & {}_3F_2 \left(\begin{matrix} -j-m_1, j-m_1+1, z-\frac{m_1}{2}+\frac{m_4}{2}-\frac{1}{2} \\ z-m_1, -m_1+m_4+1 \end{matrix}; 1 \right). \end{aligned}$$

According to 07.27.17.0042.01 of [Res08], we have the following transformation property of ${}_3F_2$ value at 1:

$$(6.70) \quad {}_3F_2(-n, b, c; d, e; 1) = \frac{(b)^{(n)}(-b-c+d+e)^{(n)}}{(d)^{(n)}(e)^{(n)}} {}_3F_2 \left(\begin{matrix} d-b, e-b, -n \\ -b-c+d+e, -b-n+1 \end{matrix}; 1 \right).$$

Apply this transform to the formula for $S_{m_1, m_4}^{j, n}(z)$,

$$\begin{aligned} & S_{m_1, m_4}^{j, n}(z) \\ &= \frac{((-1)^{2j} + (-1)^{m_4-m_1})\pi(2j)! \sec\left(\frac{m_1+m_4}{2}\pi\right) \Gamma\left(\frac{2z-m_1+m_4-1}{2}\right)}{2c_{m_1}^j c_{m_4}^j \Gamma\left(\frac{-2j-m_1+m_4+1}{2}\right) \Gamma(j+z)} \\ & {}_3F_2 \left(\begin{matrix} -j+z-1, -j-m_1, m_4-j \\ -2j, -j-\frac{m_1}{2}+\frac{m_4}{2}+\frac{1}{2} \end{matrix}; 1 \right). \end{aligned}$$

We also need to consider that m_1 and m_4 satisfy the same parity condition, that $m_1, m_4 \in \mathbb{M}(j, n; \delta_2, \delta_1)$, and therefore $\frac{m_1+m_4}{2}$ is an integer. Hence $S_{m_1, m_4}^{j, n}$ can be expressed in a simpler form in terms of hypergeometric function ${}_3F_2$:

$$(6.71) \quad S_{m_1, m_4}^{j, n}(z) = \frac{(-1)^{\frac{m_1+m_4}{2}} (2j)! \pi}{c_{m_1}^j c_{m_4}^j} \frac{\Gamma\left(\frac{2z-m_1+m_4-1}{2}\right)}{\Gamma\left(\frac{-2j-m_1+m_4+1}{2}\right) \Gamma(j+z)} {}_3F_2\left(\begin{matrix} -j+z-1, -j-m_1, m_4-j \\ -2j, -j-\frac{m_1}{2}+\frac{m_4}{2}+\frac{1}{2} \end{matrix}; 1\right)$$

6.4. Normalizations. The paper [KS80] discussed the normalization of intertwining operators in detail. There is a scalar valued meromorphic function $\eta(w, \chi_{\delta, \lambda})$ in λ such that:

$$(6.72) \quad A(w_0, \chi_{\delta, \lambda}) A(w_0^{-1}, w_0 \chi_{\delta, \lambda}) = \eta(w_0, \chi_{\delta, \lambda}) I.$$

The meromorphic function $\eta(w_0, \chi_{\delta, \lambda})$ factor into rank-one factors:

$$\eta(w_0, \chi_{\delta, \lambda}) = \prod_{\substack{\alpha \text{ reduced} \\ \langle \alpha_i, \alpha \rangle > 0 \text{ for positive } \alpha_i}} \eta_{\delta, \alpha} \left(\frac{\langle \lambda, \rho^{(\alpha)} \rangle}{\langle \rho^{(\alpha)}, \rho^{(\alpha)} \rangle} \right).$$

Moreover, these $\eta_{\delta, \alpha}(z)$'s factor into meromorphic functions $\gamma_{\delta, \alpha}(z)$:

$$(6.73) \quad \eta_{\delta, \alpha}(z) = \gamma_{\delta, \alpha}(z) \overline{\gamma_{\delta, \alpha}(-\bar{z})}.$$

In terms of the long intertwining operator $A(w_0, w_0 \chi_{\delta, \lambda})$ corresponding to w_0 , it is related to the adjoint of the intertwining operator corresponding to w_0 :

$$A(w_0, w_0 \chi_{\delta, \lambda})^* = A(w_0, \chi_{\delta, -\bar{\lambda}}).$$

We can thus normalize the intertwining operator $A(w_0, \chi_{\delta, \lambda})$ by dividing it by the meromorphic factor:

$$\gamma(w_0, \chi_{\delta, \lambda}) = \prod_{\substack{\alpha \text{ reduced} \\ \langle \alpha_i, \alpha \rangle > 0 \text{ for positive } \alpha_i}} \gamma_{\delta, \alpha} \left(\frac{\langle \lambda, \rho^{(\alpha)} \rangle}{\langle \rho^{(\alpha)}, \rho^{(\alpha)} \rangle} \right).$$

We denote the new normalized intertwining operator by

$$A'(w_0, \chi_{\delta, \lambda}) = \gamma(w_0, \chi_{\delta, \lambda})^{-1} A(w_0, \chi_{\delta, \lambda}).$$

In the case of $Sp(4, \mathbb{R})$ principal series, according to (6.72) and (6.73), there is a normalization factor $\gamma(w_{\alpha_1}, \lambda)$ of the intertwining operator entries $S_{m_1, m_4}^{j, n}(z)$, such that the operator $A(w_{\alpha_1}, \lambda)$ has the property

$$A(w_{\alpha_1}, w_{\alpha_1} \lambda) A(w_{\alpha_1}, \lambda) = \gamma(w_{\alpha_1}, \lambda) \gamma(w_{\alpha_1}, w_{\alpha_1} \lambda) I.$$

We define the normalized intertwining operator as $A'(w_{\alpha_1}, \lambda) = \frac{1}{\gamma(w_{\alpha_1}, \lambda)} A(w_{\alpha_1}, \lambda)$. Since we have assumed that $(\delta_1, \delta_2) \in \{(0, 0), (1, 1)\}$, the j and n are integers. We can check that for $m_1, m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)$,

$$(6.74) \quad \begin{aligned} & \sum_{m_2 \in \mathbb{M}(j, n; \delta_2, \delta_1)} S_{m_1, m_2}^{j, n}(z) S_{m_2, m_3}^{j, n}(1-z) \\ &= \sum_{\substack{m_2 \in \mathbb{M}(j, n; \delta_2, \delta_1) \\ -j \leq m_4, m_5 \leq j}} (-1)^{m_4+m_5} M_{m_1, m_4}^{j, n} N_{m_4, m_2}^{j, n} M_{m_2, m_5}^{j, n} N_{m_5, m_3}^{j, n} Q(z, m_4) Q(1-z, m_5). \end{aligned}$$

Since $\frac{1+(-1)^k}{2}$ is 0 for $k \equiv 1 \pmod{2}$, and is 1 for $k \equiv 0 \pmod{2}$, the sum on m_2 can be reduced to

$$(6.75) \quad \sum_{m_2 \in \mathbb{M}(j, n; \delta_2, \delta_1)} N_{m_4, m_2}^{j, n} M_{m_2, m_5}^{j, n} = \sum_{-j \leq m_2 \leq j} N_{m_4, m_2}^{j, n} M_{m_2, m_5}^{j, n} \frac{1 + (-1)^{n-m_2-\delta_1}}{2} \frac{1 + (-1)^{n+m_2-\delta_2}}{2}$$

Taking into account the condition $\delta_1 = \delta_2 \in \{0, 1\}$ and its consequence that $m_3, n \in \mathbb{Z}$, we can expand the terms

$$\begin{aligned} \frac{1 + (-1)^{n-m_2-\delta_1}}{2} \frac{1 + (-1)^{n+m_2-\delta_2}}{2} &= \frac{1}{2} + \frac{(-1)^{n-\delta_1}}{2} \frac{(-1)^{m_2} + (-1)^{-m_2}}{2} \\ &= \frac{1}{2} + \frac{(-1)^{n-\delta_1}}{2} (-1)^{m_2} \\ &= \frac{1}{2} + \frac{(-1)^{n-\delta_1}}{2} e^{im_2\pi}. \end{aligned}$$

Since $W_{m_2, m_2}^{(j, n)}(0, -\pi, 0, 0) = e^{im_2\pi}$, by (6.14) and the definition (6.19) and (6.20) of $M_{m_2, m_5}^{j, n}$ and $N_{m_4, m_2}^{j, n}$, we have

$$\sum_{m_2 \in \mathbb{M}(j, n; \delta_2, \delta_1)} N_{m_4, m_2}^{j, n} M_{m_2, m_5}^{j, n} W_{m_2, m_2}^{(j, n)}(0, -\pi, 0, 0) = W_{m_4, m_5}^{(j, n)}(0, 0, -\pi, 0).$$

The sum (6.75) can thus be written as

$$\begin{aligned} \sum_{m_2 \in \mathbb{M}(j, n; \delta_2, \delta_1)} N_{m_4, m_2}^{j, n} M_{m_2, m_5}^{j, n} &= \frac{1}{2} \delta_{m_4, m_5} + \frac{(-1)^{n-\delta_1}}{2} W_{m_4, m_5}^{(j, n)}(e^{\pi U_2}) \\ &= \frac{1}{2} \delta_{m_4, m_5} + \frac{(-1)^{j-m_4-\delta_1+n}}{2} \delta_{-m_4, m_5}, \end{aligned}$$

and the sum (6.74) becomes

$$\begin{aligned} &\sum_{m_2 \in \mathbb{M}(j, n; \delta_2, \delta_1)} S_{m_1, m_2}^{j, n}(z) S_{m_2, m_3}^{j, n}(1-z) \\ &= \sum_{-j \leq m_4, m_5 \leq j} (-1)^{m_4+m_5} M_{m_1, m_4}^{j, n} N_{m_5, m_3}^{j, n} Q(z, m_4) Q(1-z, m_5) \frac{\delta_{m_4, m_5}}{2} + \\ &\quad \sum_{-j \leq m_4, m_5 \leq j} (-1)^{m_4+m_5} M_{m_1, m_4}^{j, n} N_{m_5, m_3}^{j, n} Q(z, m_4) Q(1-z, m_5) \frac{(-1)^{j-m_4-\delta_1+n}}{2} \delta_{-m_4, m_5} \\ &= \frac{1}{2} \sum_{-j \leq m_4 \leq j} M_{m_1, m_4}^{j, n} N_{m_4, m_3}^{j, n} Q(z, m_4) Q(1-z, m_4) + \frac{1}{2} \sum_{-j \leq m_4 \leq j} M_{m_1, m_4}^{j, n} N_{-m_4, m_3}^{j, n} \\ &\quad (-1)^{j-m_4-\delta_1+n} Q(z, m_4) Q(1-z, -m_4). \end{aligned} \quad (6.76)$$

We observe that the product of the Q -functions are simply products of Γ -functions:

$$(6.77) \quad Q(z, n) Q(1-z, n) = Q(z, n) Q(1-z, -n) = \pi \frac{\Gamma(z-1/2) \Gamma(-z+1/2)}{\Gamma(z) \Gamma(1-z)},$$

hence the formula (6.76) is in fact

$$\begin{aligned}
 & \sum_{m_2 \in \mathbb{M}(j, n; \delta_2, \delta_1)} S_{m_1, m_2}^{j, n}(z) S_{m_2, m_3}^{j, n}(1-z) \\
 &= \frac{\pi}{2} \frac{\Gamma(z-1/2)}{\Gamma(z)} \frac{\Gamma(-z+1/2)}{\Gamma(1-z)} \sum_{-j \leq m_4 \leq j} \left(M_{m_1, m_4}^{j, n} N_{m_4, m_3}^{j, n} + (-1)^{j-m_4-\delta_1+n} M_{m_1, m_4}^{j, n} N_{-m_4, m_3}^{j, n} \right) \\
 &= \frac{\pi}{2} \frac{\Gamma(z-1/2)}{\Gamma(z)} \frac{\Gamma(-z+1/2)}{\Gamma(1-z)} (\delta_{m_1, m_3} + (-1)^{-m_1-\delta_1+n} \delta_{m_1, m_3}) \\
 & \quad (6.78) \\
 &= \pi \frac{\Gamma(z-1/2)}{\Gamma(z)} \frac{\Gamma(-z+1/2)}{\Gamma(1-z)} \frac{1 + (-1)^{-m_1-\delta_1+n}}{2} \delta_{m_1, m_3}.
 \end{aligned}$$

Therefore, it is natural to consider be the action of the intertwining operators on the K -types with $j = 0$:

$$(6.79) \quad S_{0,0}^{(0,n)}(z) = \frac{\sqrt{\pi} \Gamma(z-1/2)}{\Gamma(z)}$$

and set the normalization factor $\gamma(w_\alpha, \lambda)$ to be

$$\gamma(w_\alpha, \lambda) = S_{0,0}^{(0,n)} \left(\frac{\langle \check{\alpha}, \lambda + \rho \rangle}{2} \right).$$

The matrix entries of the normalized intertwining operator $A'(w_{\alpha_1}, \lambda)$ are given by the function

$$(6.80) \quad \mathcal{S}_{m_1, m_2}^{j, n}(z) := \frac{S_{m_1, m_2}^{j, n}(z)}{S_{0,0}^{(0,n)}(z)} = \left(\frac{\sqrt{\pi} \Gamma(z-1/2)}{\Gamma(z)} \right)^{-1} S_{m_1, m_2}^{j, n}(z),$$

and by (6.71) we can further simplify the expression of the matrix entries in terms of Pochhammer symbols:

$$(6.81) \quad \mathcal{S}_{m_1, m_2}^{j, n}(z) = \frac{(-1)^{\frac{m_1+m_2}{2}} (2j)! \sqrt{\pi}}{c_{m_1}^j c_{m_2}^j \Gamma\left(\frac{1-2j-m_1+m_2}{2}\right)} \frac{(z-1/2)^{(-\frac{m_1-m_2}{2})}}{(z)^{(j)}} {}_3F_2 \left(\begin{matrix} -j+z-1, -j-m_1, -j+m_2 \\ -2j, -j-\frac{m_1-m_2}{2}+\frac{1}{2} \end{matrix}; 1 \right).$$

According to (6.78), if $m_1, m_3 \in \mathbb{M}(j, n; \delta_1, \delta_2)$, the normalized intertwining operator entries $\mathcal{S}_{m_1, m_2}^{j, n}(z)$ satisfy the property

$$(6.82) \quad \sum_{m_2 \in \mathbb{M}(j, n; \delta_2, \delta_1)} \mathcal{S}_{m_1, m_2}^{j, n}(z) \mathcal{S}_{m_2, m_3}^{j, n}(1-z) = \delta_{m_1, m_3}.$$

If $n \geq 0$, the n -th Laurent series coefficient of the function in terms of t is the ${}_3F_2$ hypergeometric function (see [SM84, Page 94]):

$$(6.83) \quad [(1-t)^{-\lambda} {}_2F_1 \left(\begin{matrix} a, b \\ c \end{matrix}; zt \right)]_n = \frac{\Gamma(\lambda+n)}{\Gamma(n+1)\Gamma(\lambda)} {}_3F_2 \left(\begin{matrix} -n, a, b \\ c, 1-n-\lambda \end{matrix}; z \right).$$

Therefore, $\mathcal{S}_{m_1, m_2}^{j, n}(z)$ is the constant Laurent series coefficient of either of the following two functions:

$$(6.84) \quad H_{m_1, m_2}^j(z; t) = \frac{(2j)!(j+m_1)!}{c_{m_1}^j c_{m_2}^j \sqrt{\pi}} \Gamma\left(\frac{1-m_1-m_2}{2}\right) \frac{(z-1/2)^{(-\frac{m_1-m_2}{2})}}{(z)^{(j)}} \times \frac{(1-t)^{\frac{-1+m_1+m_2}{2}}}{(-t)^{j+m_1}} {}_2F_1\left(\begin{matrix} -j+m_2, -1-j+z \\ -2j \end{matrix}; t\right)$$

$$(6.85) \quad G_{m_1, m_2}^j(z; t) = \frac{(2j)!(j-m_2)!}{c_{m_1}^j c_{m_2}^j \sqrt{\pi}} \Gamma\left(\frac{1+m_1+m_2}{2}\right) \frac{(z-1/2)^{(-\frac{m_1-m_2}{2})}}{(z)^{(j)}} \times \frac{(1-t)^{\frac{-1-m_1-m_2}{2}}}{(-t)^{j-m_2}} {}_2F_1\left(\begin{matrix} -j-m_1, -1-j+z \\ -2j \end{matrix}; t\right).$$

6.5. Computing the Product Matrix. In this section we compute the product of the four matrices:

$$A'(w_0, \chi_{\delta, \lambda}) = A_4(\lambda) \cdot A_3(\lambda) \cdot A_2(\lambda) \cdot A_1(\lambda)$$

when $\delta = (0, 0)$ or $(1, 1)$. We would also like to normalize each individual matrices $A_i(\lambda)$ such that they satisfy the condition:

$$A_i(-\lambda)A_i(\lambda) = I.$$

Recalling the parity conditions (4.5) and (4.6) satisfied by j, n, m_1, m_2 , since $(\delta_1, \delta_2) \in \{(0, 0), (1, 1)\}$, the pair (j, n) are integers, and m_1, m_2 satisfy

$$\begin{matrix} n-m_i+\delta_1 \equiv 0 \\ n+m_i+\delta_2 \equiv 0 \end{matrix} \pmod{2}.$$

Despite the difficulty of calculating the product of the 4 matrices, a lot of terms in the sum can be reduced using the parity condition. We replace the function $T_{m_1}^n(z)$ by the normalized function defined by

$$\mathcal{T}_{m_1}^n(z) = \frac{i^{-n+m_1}}{(z)^{(\frac{m_1-n}{2})}(z)^{(\frac{-m_1+n}{2})}}.$$

The matrix entry $[A'(\lambda)]_{m_1, m_2}^{j, n}$ of the normalized intertwining operator can be expressed as the constant term of the Laurent series in t_1, t_2 of the sum:

$$(6.86) \quad [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) = \mathcal{T}_{m_2}^n\left(\frac{\lambda_2+1}{2}\right) \times \sum_{m_3 \in \mathbb{M}(j, n; \delta_1, \delta_2)} G_{m_2, m_3}^j\left(\frac{\lambda_1+\lambda_2+1}{2}, t_2\right) \mathcal{T}_{m_3}^n\left(\frac{\lambda_1+1}{2}\right) H_{m_3, m_1}^j\left(\frac{\lambda_1-\lambda_2+1}{2}; t_1\right).$$

Plugging (6.84) and (6.85) into (6.86), we have

$$\begin{aligned}
 & [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) \\
 &= \frac{((2j)!)^2}{c_{m_1}^j c_{m_2}^j \pi} \frac{i^{-n+m_2}}{\left(\frac{\lambda_2+1}{2}\right)^{\left(\frac{m_2-n}{2}\right)} \left(\frac{\lambda_2+1}{2}\right)^{\left(\frac{-m_2+n}{2}\right)}} \frac{1}{\left(\frac{\lambda_1-\lambda_2+1}{2}\right)^{(j)} \left(\frac{\lambda_1+\lambda_2+1}{2}\right)^{(j)}} \times \\
 & {}_2F_1\left(-j+m_1, -1-j+\frac{\lambda_1-\lambda_2+1}{2}; t_1\right) {}_2F_1\left(-j-m_2, -1-j+\frac{\lambda_1+\lambda_2+1}{2}; t_2\right) \times \\
 & \sum_{m_3 \in \mathbb{M}(j, n; \delta_1, \delta_2)} \frac{i^{-n+m_3}}{\left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{m_3-n}{2}\right)} \left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{-m_3+n}{2}\right)}} \Gamma\left(\frac{1+m_3+m_2}{2}\right) \Gamma\left(\frac{1-m_1-m_3}{2}\right) \times \\
 & \left(\frac{\lambda_1-\lambda_2}{2}\right)^{\left(\frac{m_1-m_3}{2}\right)} \left(\frac{\lambda_1+\lambda_2}{2}\right)^{\left(\frac{m_3-m_2}{2}\right)} \frac{(1-t_2)^{\frac{-1-m_3-m_2}{2}} (1-t_1)^{\frac{-1+m_1+m_3}{2}}}{(-t_2)^{j-m_3} (-t_1)^{j+m_3}}.
 \end{aligned}$$

Based on the parity of $j - m_3$, we can separate the calculation into two cases:

6.5.1. *Summation when $j - n \equiv \delta_i \pmod{2}$.* In the situation that $j - n \equiv \delta_i \pmod{2}$, the set of $m_3 \in \mathbb{M}(j, n; \delta_2, \delta_1)$'s are:

$$m_3 = j - 2p \text{ where } p \in \{0, 1, 2, \dots, j\}.$$

Thus we can replace m_3 by $j - 2p$, and the sum reduces to a sum of finitely many Pochhammer symbols:

$$\begin{aligned}
 [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) &= \frac{((2j)!)^2}{c_{m_1}^j c_{m_2}^j \pi} \frac{i^{-n+m_2}}{\left(\frac{\lambda_2+1}{2}\right)^{\left(\frac{m_2-n}{2}\right)} \left(\frac{\lambda_2+1}{2}\right)^{\left(\frac{-m_2-n}{2}\right)}} \times \\
 & {}_2F_1\left(-j+m_1, -j+\frac{\lambda_1-\lambda-1}{2}; t_1\right) {}_2F_1\left(-j-m_2, -j+\frac{\lambda_1+\lambda-1}{2}; t_2\right) \times \\
 & \sum_{p=0}^j \frac{i^{-n+j-2p}}{\left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{j-n-2p}{2}\right)} \left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{-j-n-2p}{2}\right)}} \Gamma\left(\frac{1+j+m_2}{2} - p\right) \Gamma\left(\frac{1-j-m_1}{2} + p\right) \times \\
 & (6.87) \\
 & \left(\frac{\lambda_1-\lambda_2}{2}\right)^{\left(\frac{-j+m_1}{2}+p\right)} \left(\frac{\lambda_1+\lambda_2}{2}\right)^{\left(\frac{j-m_2}{2}-p\right)} (1-t_1)^{\frac{-1+j+m_1}{2}-p} (1-t_2)^{\frac{-1-j-m_2}{2}+p} t_1^{-2j+2p} t_2^{-2p}.
 \end{aligned}$$

By definition of Pochhammer symbols, we have the formula

$$(6.88) \quad (x)^{(l \pm p)} = (x)^{(l)} (x+l)^{(\pm p)}$$

which we can apply to the Pochhammer symbols in the summation (6.87) to single out the $\pm p$'s in the Pochhammer exponents. Also, by applying the formula

$$(6.89) \quad (x)^{(-p)} (1-x)^{(p)} = (-1)^p \text{ for } p \in \mathbb{Z}$$

to the Pochhammer symbols with $-p$ as exponent, we can manage to change all the exponents of the Pochhammer symbols in (6.87) to (p) . The explicit set of rules is

the following

$$\begin{aligned}
 & \left(\frac{\lambda_1 + 1}{2} \right)^{\left(\frac{j-n-2p}{2} \right)} \rightarrow \left(\frac{\lambda_1 + 1}{2} \right)^{\left(\frac{j-n}{2} \right)} \left(\frac{\lambda_1 + 1 + j - n}{2} \right)^{(-p)} \\
 (6.90) \quad & \rightarrow (-1)^p \left(\frac{\lambda_1 + 1}{2} \right)^{\left(\frac{j-n}{2} \right)} \left(1 - \frac{\lambda_1 + 1 + j - n}{2} \right)^{(p)}
 \end{aligned}$$

$$(6.91) \quad \left(\frac{\lambda_1 + 1}{2} \right)^{\left(-\frac{j-n-2p}{2} \right)} \rightarrow \left(\frac{\lambda_1 + 1}{2} \right)^{\left(-\frac{j-n}{2} \right)} \left(\frac{\lambda_1 + 1 - j + n}{2} \right)^{(p)}$$

$$\begin{aligned}
 & \Gamma \left(\frac{1+j+m_2}{2} - p \right) \rightarrow \Gamma \left(\frac{1+j+m_2}{2} \right) \left(\frac{1+j+m_2}{2} \right)^{(-p)} \\
 (6.92) \quad & \rightarrow (-1)^p \Gamma \left(\frac{1+j+m_2}{2} \right) \left(1 - \frac{1+j+m_2}{2} \right)^{(p)}
 \end{aligned}$$

$$(6.93) \quad \Gamma \left(\frac{1-j-m_1}{2} + p \right) \rightarrow \Gamma \left(\frac{1-j-m_1}{2} \right) \left(\frac{1-j-m_1}{2} \right)^{(p)}$$

$$\begin{aligned}
 (6.94) \quad & \left(\frac{\lambda_1 - \lambda_2}{2} \right)^{\left(-\frac{j-m_1-2p}{2} \right)} \rightarrow \left(\frac{\lambda_1 - \lambda_2}{2} \right)^{\left(-\frac{j-m_1}{2} \right)} \left(\frac{\lambda_1 - \lambda_2 - j - m_1}{2} \right)^{(p)} \\
 & \left(\frac{\lambda_1 + \lambda_2}{2} \right)^{\left(\frac{j-m_2-2p}{2} \right)} \rightarrow \left(\frac{\lambda_1 + \lambda_2}{2} \right)^{\left(\frac{j-m_2}{2} \right)} \left(\frac{\lambda_1 + \lambda_2 + j - m_2}{2} \right)^{(-p)} \\
 (6.95) \quad & \rightarrow (-1)^p \left(\frac{\lambda_1 + \lambda_2}{2} \right)^{\left(\frac{j-m_2}{2} \right)} \left(1 - \frac{\lambda_1 + \lambda_2 + j - m_2}{2} \right)^{(p)}.
 \end{aligned}$$

After applying (6.90)-(6.95) and bringing out the factors not involved in the summation, we have

$$\begin{aligned}
 & [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) = \\
 & \frac{j! + m_2 - 2n ((2j)!)^2 \left(\frac{\lambda_1 - \lambda_2}{2} \right)^{\left(\frac{-j+m_1}{2} \right)} \left(\frac{\lambda_1 + \lambda_2}{2} \right)^{\left(\frac{j-m_2}{2} \right)}}{\pi c_{m_1}^j c_{m_2}^j \left(\frac{\lambda_1 + 1}{2} \right)^{\left(\frac{-j+n}{2} \right)} \left(\frac{\lambda_1 + 1}{2} \right)^{\left(\frac{j-n}{2} \right)}} \times \\
 & \frac{\Gamma \left(\frac{1+j+m_2}{2} \right) \Gamma \left(\frac{1-j-m_1}{2} \right)}{\left(\frac{\lambda_1 + \lambda_2 + 1}{2} \right)^{(j)} \left(\frac{\lambda_1 - \lambda_2 + 1}{2} \right)^{(j)} \left(\frac{\lambda_2 + 1}{2} \right)^{\left(\frac{n-m_2}{2} \right)} \left(\frac{\lambda_2 + 1}{2} \right)^{\left(\frac{-n+m_2}{2} \right)}} \times \\
 & {}_2F_1 \left(-j+m_1, \frac{\lambda_1 - \lambda_2 - 2j-1}{-2j}; t_1 \right) {}_2F_1 \left(-j-m_2, \frac{\lambda_1 + \lambda_2 - 2j-1}{-2j}; t_2 \right) \times \\
 (6.96) \quad & \gamma_{m_1, m_2}^{j, n}(\lambda; t_1, t_2),
 \end{aligned}$$

where the function $\gamma_{m_1, m_2}^{j, n}(\lambda; t_1, t_2)$ in two variables t_1, t_2 is defined as the hypergeometric sum

$$(6.97) \quad \gamma_{m_1, m_2}^{j, n}(\lambda; t_1, t_2) = (1 - t_1)^{\frac{-1+j+m_1}{2}} (-t_1)^{-2j} (1 - t_2)^{\frac{-1-j-m_2}{2}} \times \\ \sum_{p=0}^j \frac{\left(\frac{1-j-m_1}{2}\right)^{(p)} \left(\frac{-j+m_1+\lambda_1-\lambda_2}{2}\right)^{(p)} \left(\frac{1-j+n-\lambda_1}{2}\right)^{(p)} (1)^{(p)}}{\left(\frac{1-j-m_2}{2}\right)^{(p)} \left(1 - \frac{j-m_2+\lambda_1+\lambda_2}{2}\right)^{(p)} \left(\frac{1-j+n+\lambda_1}{2}\right)^{(p)}} \frac{1}{p!} \left(\frac{t_1^2(1-t_2)}{t_2^2(1-t_1)}\right)^p.$$

We can use a partial sum formula for hypergeometric series from 16.2.4 of [OLBC10]:

$$(6.98) \quad \sum_{k=0}^m \frac{(\mathbf{a})^{(k)}}{(\mathbf{b})^{(k)}} \frac{z^k}{k!} = \frac{(\mathbf{a})^{(m)} z^m}{(\mathbf{b})^{(m)} m!} {}_qF_p \left(\begin{matrix} -m, 1, 1-m-\mathbf{b} \\ 1-m-\mathbf{a} \end{matrix}; \frac{(-1)^{p+q+1}}{z} \right),$$

where $\mathbf{a} = (a_1, \dots, a_p)$ and $\mathbf{b} = (b_1, \dots, b_q)$ are p and q dimensional vectors of complex numbers, and the Pochhammer symbol $(\mathbf{a})^{(k)}$ and $(\mathbf{b})^{(k)}$ for these two vectors are defined as the product $(\mathbf{a})^{(k)} = (a_1)^{(k)} \dots (a_p)^{(k)}$ and $(\mathbf{b})^{(k)} = (b_1)^{(k)} \dots (b_q)^{(k)}$. Then the summation term in $\gamma_{m_1, m_2}^{j, n}(\lambda; t_1, t_2)$ can be simplified into a single compact form hypergeometric function ${}_5F_4$:

$$\begin{aligned} & \sum_{p=0}^j \frac{\left(\frac{1-j-m_1}{2}\right)^{(p)} \left(\frac{-j+m_1+\lambda_1-\lambda_2}{2}\right)^{(p)} \left(\frac{1-j+n-\lambda_1}{2}\right)^{(p)} (1)^{(p)}}{\left(\frac{1-j-m_2}{2}\right)^{(p)} \left(1 - \frac{j-m_2+\lambda_1+\lambda_2}{2}\right)^{(p)} \left(\frac{1-j+n+\lambda_1}{2}\right)^{(p)}} \frac{1}{p!} \left(\frac{t_1^2(1-t_2)}{t_2^2(1-t_1)}\right)^p \\ &= \frac{\left(\frac{1-j-m_1}{2}\right)^{(j)} \left(\frac{-j+m_1+\lambda_1-\lambda_2}{2}\right)^{(j)} \left(\frac{-j+n-\lambda_1+1}{2}\right)^{(j)}}{\left(\frac{1-j-m_2}{2}\right)^{(j)} \left(\frac{-j+m_2-\lambda_1-\lambda_2}{2} + 1\right)^{(j)} \left(\frac{-j+n+\lambda_1+1}{2}\right)^{(j)}} \left(\frac{t_1^2(1-t_2)}{t_2^2(1-t_1)}\right)^j \times \\ & {}_5F_4 \left(\begin{matrix} -j, 1, \frac{1-j+m_2}{2}, \frac{-j-n-\lambda_1+1}{2}, \frac{-j-m_2+\lambda_1+\lambda_2}{2} \\ \frac{1-j+m_1}{2}, \frac{1-j-n+\lambda_1}{2}, \frac{-j-m_1-\lambda_1+\lambda_2}{2} + 1, -j \end{matrix}; \frac{t_2^2(1-t_1)}{t_1^2(1-t_2)} \right) \end{aligned}$$

and if we replace the sum in (6.102) by the expression above involving ${}_5F_4$, also we recover the Pochhammer symbols from the transforms (6.90)-(6.95) based on (6.88) and (6.89), the original function (6.87) can be expressed as

$$(6.99) \quad [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) = \frac{(-1)^n ((2j)!)^2}{\left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{j+n}{2}\right)} \left(\frac{\lambda_1+1}{2}\right)^{\left(-\frac{j+n}{2}\right)} \left(\frac{\lambda_1-\lambda_2+1}{2}\right)^{(j)} \left(\frac{\lambda_1+\lambda_2+1}{2}\right)^{(j)}} \times \\ \frac{1}{c_{m_1}^j c_{m_2}^j} \frac{\left(\frac{\lambda_1-\lambda_2}{2}\right)^{\left(\frac{j+m_1}{2}\right)} \left(\frac{\lambda_1+\lambda_2}{2}\right)^{\left(\frac{-j-m_2}{2}\right)} \Gamma\left(\frac{1+j-m_1}{2}\right)}{\left(\frac{\lambda_2+1}{2}\right)^{\left(\frac{m_2-n}{2}\right)} \left(\frac{\lambda_2+1}{2}\right)^{\left(-\frac{m_2-n}{2}\right)} \Gamma\left(\frac{1+j-m_2}{2}\right)} \times \\ (1 - t_1)^{\frac{-1-j+m_1}{2}} (1 - t_2)^{\frac{-1+j-m_2}{2}} t_2^{-2j} \times \\ {}_2F_1 \left(\begin{matrix} -j+m_1, \frac{\lambda_1-\lambda_2-2j-1}{2} \\ -2j \end{matrix}; t_1 \right) {}_2F_1 \left(\begin{matrix} -j-m_2, \frac{\lambda_1+\lambda_2-2j-1}{2} \\ -2j \end{matrix}; t_2 \right) \times \\ {}_5F_4 \left(\begin{matrix} -j, 1, \frac{-j+m_2+1}{2}, \frac{-j-n-\lambda_1+1}{2}, \frac{-j-m_2+\lambda_1+\lambda_2}{2} \\ -j, \frac{-j+m_1+1}{2}, \frac{-j-n+\lambda_1+1}{2}, \frac{-j-m_1-\lambda_1+\lambda_2}{2} + 1 \end{matrix}; \frac{t_2^2(1-t_1)}{t_1^2(1-t_2)} \right).$$

The $-j$'s as parameters to the function ${}_5F_4$ don't necessarily cancel since they are non-positive. They play an important role in making the function ${}_5F_4$ rational.

6.5.2. *Summation when $j - n \not\equiv \delta_i \pmod{2}$.* On the other hand, if $j - n \not\equiv \delta_i \pmod{2}$,

$$m = j - 2p - 1 \text{ where } p \in \{0, 1, \dots, j - 1\},$$

we have

$$\begin{aligned} [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) &= \frac{((2j)!)^2}{c_{m_1}^j c_{m_2}^j \pi} \frac{i^{-n+m_2}}{\left(\frac{\lambda_2+1}{2}\right)^{\left(\frac{m_2-n}{2}\right)} \left(\frac{\lambda_2+1}{2}\right)^{\left(-\frac{m_2-n}{2}\right)}} \times \\ &{}_2F_1\left(-j+m_1, -j+\frac{\lambda_1-\lambda-1}{2}; t_1\right) {}_2F_1\left(-j-m_2, -j+\frac{\lambda_1+\lambda-1}{2}; t_2\right) \times \\ &\sum_{p=0}^{j-1} \frac{i^{-n+j-2p-1}}{\left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{j-n-2p-1}{2}\right)} \left(\frac{\lambda_1+1}{2}\right)^{\left(-\frac{j-n-2p-1}{2}\right)}} \Gamma\left(\frac{j+m_2}{2} - p\right) \Gamma\left(\frac{2-j-m_1}{2} + p\right) \times \\ &\left(\frac{\lambda_1-\lambda_2}{2}\right)^{\left(\frac{1-j+m_1}{2}+p\right)} \left(\frac{\lambda_1+\lambda_2}{2}\right)^{\left(\frac{j-m_2-1}{2}-p\right)} (1-t_1)^{\frac{-2+j+m_1}{2}-p} \times \\ (6.100) \quad &(1-t_2)^{\frac{-j-m_2}{2}+p} t_1^{-2j+2p+1} t_2^{-2p-1}. \end{aligned}$$

Comparing with the even case, after applying (6.88) and (6.89), the summation will change to

$$\begin{aligned} [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) &= \\ &\frac{i^{j+m_2-2n} ((2j)!)^2 \left(\frac{\lambda_1-\lambda_2}{2}\right)^{\left(\frac{-j+m_1+1}{2}\right)} \left(\frac{\lambda_1+\lambda_2}{2}\right)^{\left(\frac{j-m_2-1}{2}\right)}}{\pi c_{m_1}^j c_{m_2}^j \left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{-j+n+1}{2}\right)} \left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{j-n-1}{2}\right)}} \times \\ &\frac{\Gamma\left(\frac{j+m_2}{2}\right) \Gamma\left(\frac{2-j-m_1}{2}\right)}{\left(\frac{\lambda_1+\lambda_2+1}{2}\right)^{(j)} \left(\frac{\lambda_1-\lambda_2+1}{2}\right)^{(j)} \left(\frac{\lambda_2+1}{2}\right)^{\left(\frac{n-m_2}{2}\right)} \left(\frac{\lambda_2+1}{2}\right)^{\left(\frac{-n+m_2}{2}\right)}} \times \\ &{}_2F_1\left(-j+m_1, \frac{\lambda_1-\lambda_2-2j-1}{2}; t_1\right) {}_2F_1\left(-j-m_2, \frac{\lambda_1+\lambda_2-2j-1}{2}; t_2\right) \times \\ (6.101) \quad &\gamma_{m_1, m_2}^{'j, n}(\lambda; t_1, t_2), \end{aligned}$$

where

$$\begin{aligned} \gamma_{m_1, m_2}^{'j, n}(\lambda; t_1, t_2) &= (1-t_1)^{\frac{j+m_1-2}{2}} t_2^{-2j+1} t_1^{-1} (1-t_2)^{\frac{-j-m_2}{2}} \times \\ &{}_2F_1\left(-j+m_1, \frac{\lambda_1-\lambda_2-2j-1}{2}; t_1\right) {}_2F_1\left(-j-m_2, \frac{\lambda_1+\lambda_2-2j-1}{2}; t_2\right) \times \\ (6.102) \quad &\sum_{p=0}^{j-1} \frac{\left(\frac{2-j-m_1}{2}\right)^{(p)} \left(\frac{-j+m_2+\lambda_1-\lambda_2}{2}\right)^{(p)} \left(1-\frac{j-n+\lambda_1}{2}\right)^{(p)} (1)^{(p)}}{\left(\frac{2-j-m_2}{2}\right)^{(p)} \left(\frac{3}{2}-\frac{j-m_2-\lambda_1-\lambda_2}{2}\right)^{(p)} \left(\frac{2-j+n+\lambda_1}{2}\right)^{(p)}} \frac{1}{p!} \left(\frac{t_1^2(1-t_2)}{t_2^2(1-t_1)}\right)^p. \end{aligned}$$

By the same formula (6.98) and using the relation between Pochhammer symbols and Γ -functions $(a)^{(n)} = \Gamma(a+n)/\Gamma(a)$ to absorb the extra Γ factors in $\gamma_{m_1, m_2}^{'j, n}$, the

matrix entries for the long intertwining operator has an expression

$$\begin{aligned}
 [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) = & \frac{(-1)^n ((2j)!)^2}{\left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{j+n-1}{2}\right)} \left(\frac{\lambda_1+1}{2}\right)^{\left(-\frac{j+n-1}{2}\right)} \left(\frac{\lambda_1-\lambda_2+1}{2}\right)^{(j)} \left(\frac{\lambda_1+\lambda_2+1}{2}\right)^{(j)}} \times \\
 & \frac{1}{c_{m_1}^j c_{m_2}^j} \frac{\left(\frac{\lambda_1-\lambda_2}{2}\right)^{\left(\frac{j+m_1-1}{2}\right)} \left(\frac{\lambda_1+\lambda_2}{2}\right)^{\left(\frac{-j-m_2+1}{2}\right)} \Gamma\left(\frac{j-m_1}{2}\right)}{\left(\frac{\lambda_2+1}{2}\right)^{\frac{m_2-n}{2}} \left(\frac{\lambda_2+1}{2}\right)^{-\frac{m_2-n}{2}} \Gamma\left(\frac{j-m_2}{2}\right)} \times \\
 & (1-t_1)^{\frac{-j+m_1}{2}} (1-t_2)^{\frac{j-m_2-2}{2}} t_2^{1-2j} t_1^{-1} \times \\
 & {}_2F_1\left(-j+m_1, \frac{\lambda_1-\lambda_2-2j-1}{-2j}; t_1\right) {}_2F_1\left(-j-m_2, \frac{\lambda_1+\lambda_2-2j-1}{-2j}; t_2\right) \times \\
 (6.103) \quad & {}_5F_4\left(\begin{matrix} -j+1, 1, \frac{-j+m_2+2}{2}, \frac{-j-n-\lambda_1+2}{2}, \frac{-j-m_2+\lambda_1+\lambda_2+1}{2} \\ -j+1, \frac{-j+m_1+2}{2}, \frac{-j-n+\lambda_1+2}{2}, \frac{-j-m_1-\lambda_1+\lambda_2+1}{2} \end{matrix}; \frac{t_2^2(1-t_1)}{t_1^2(1-t_2)}\right).
 \end{aligned}$$

Therefore, if we set $\epsilon_\delta^{j, n} = \begin{cases} 0 & j-n \equiv \delta_i \pmod{2} \\ 1 & j-n \not\equiv \delta_i \pmod{2} \end{cases}$, we can summarize the above two cases (6.99), (6.103) into one single formula:

$$\begin{aligned}
 [A(\lambda)]_{m_1, m_2}^{j, n}(t_1, t_2) = & \frac{(-1)^n ((2j)!)^2}{\left(\frac{\lambda_1+1}{2}\right)^{\left(\frac{j+n-\epsilon_\delta^{j, n}}{2}\right)} \left(\frac{\lambda_1+1}{2}\right)^{\left(-\frac{j+n-\epsilon_\delta^{j, n}}{2}\right)} \left(\frac{\lambda_1-\lambda_2+1}{2}\right)^{(j)} \left(\frac{\lambda_1+\lambda_2+1}{2}\right)^{(j)}} \times \\
 & \frac{1}{c_{m_1}^j c_{m_2}^j} \frac{\left(\frac{\lambda_1-\lambda_2}{2}\right)^{\left(\frac{j+m_1-\epsilon_\delta^{j, n}}{2}\right)} \left(\frac{\lambda_1+\lambda_2}{2}\right)^{\left(\frac{-j-m_2+\epsilon_\delta^{j, n}}{2}\right)} \Gamma\left(\frac{1-\epsilon_\delta^{j, n}+j-m_1}{2}\right)}{\left(\frac{\lambda_2+1}{2}\right)^{\frac{m_2-n}{2}} \left(\frac{\lambda_2+1}{2}\right)^{-\frac{m_2-n}{2}} \Gamma\left(\frac{1-\epsilon_\delta^{j, n}+j-m_2}{2}\right)} \times \\
 & (1-t_1)^{\frac{-1+\epsilon_\delta^{j, n}-j+m_1}{2}} (1-t_2)^{\frac{-1-\epsilon_\delta^{j, n}+j-m_2}{2}} t_2^{\epsilon_\delta^{j, n}-2j} t_1^{-\epsilon_\delta^{j, n}} \times \\
 & {}_2F_1\left(-j+m_1, \frac{\lambda_1-\lambda_2-2j-1}{-2j}; t_1\right) {}_2F_1\left(-j-m_2, \frac{\lambda_1+\lambda_2-2j-1}{-2j}; t_2\right) \times \\
 (6.104) \quad & {}_5F_4\left(\begin{matrix} -j+\epsilon_\delta^{j, n}, 1, \frac{-j+m_2+1+\epsilon_\delta^{j, n}}{2}, \frac{-j-n-\lambda_1+1+\epsilon_\delta^{j, n}}{2}, \frac{-j-m_2+\lambda_1+\lambda_2+\epsilon_\delta^{j, n}}{2} \\ -j+\epsilon_\delta^{j, n}, \frac{-j+m_1+1+\epsilon_\delta^{j, n}}{2}, \frac{-j-n+\lambda_1+1+\epsilon_\delta^{j, n}}{2}, \frac{-j-m_1-\lambda_1+\lambda_2+\epsilon_\delta^{j, n}}{2} \end{matrix}; \frac{t_2^2(1-t_1)}{t_1^2(1-t_2)}\right).
 \end{aligned}$$

Thus we have finished the proof of Theorem 1.1.

APPENDIX A. AN INTRODUCTION TO WIGNER D -MATRICES AND HYPERGEOMETRIC FUNCTIONS

This appendix is a summary of [Zha19, Section 2]. The Lie algebra $\mathfrak{sl}(2, \mathbb{C})$ is generated by the *Pauli matrices* :

$$\sigma_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

and its compact real form $\mathfrak{u}(2)$ is generated by $\gamma_i = \frac{i}{2}\sigma_i$:

$$\gamma_0 = \begin{pmatrix} \frac{i}{2} & 0 \\ 0 & \frac{i}{2} \end{pmatrix}, \gamma_1 = \begin{pmatrix} 0 & \frac{i}{2} \\ \frac{i}{2} & 0 \end{pmatrix}, \gamma_2 = \begin{pmatrix} 0 & \frac{i}{2} \\ -\frac{i}{2} & 0 \end{pmatrix}, \gamma_3 = \begin{pmatrix} \frac{i}{2} & 0 \\ 0 & -\frac{i}{2} \end{pmatrix}.$$

These generators exponentiate to an *Euler angle* parametrization of the compact Lie group $U(2)$:

(A.1)

$$e^{-\zeta\gamma_0}\mathcal{U}(\psi, \theta, \phi) = e^{-\zeta\gamma_0}e^{-\psi\gamma_3}e^{-\theta\gamma_2}e^{-\phi\gamma_3} = \begin{pmatrix} e^{\frac{1}{2}(-\zeta-\phi-\psi)} \cos \frac{\theta}{2} & -e^{\frac{1}{2}(-\zeta+\phi-\psi)} \sin \frac{\theta}{2} \\ e^{\frac{1}{2}(-\zeta-\phi+\psi)} \sin \frac{\theta}{2} & e^{\frac{1}{2}(-\zeta+\phi+\psi)} \cos \frac{\theta}{2} \end{pmatrix}.$$

An arbitrary irreducible representation $\pi_{j,n}$ of $U(2)$ can be realized on the space $\text{Sym}^{2j}W \otimes \det^{(j+n)}$, on which $U(2)$ acts by right regular action on any degree $2j$ homogeneous polynomial $f \in \text{Sym}^{2j}(W) \otimes \det^{(j+n)}$ in 2 variables z_1, z_2 :

$$\pi_{j,n}(g)f(z) = (\det g)^{j+n}f(g^{-1}z).$$

Let $m \in \frac{1}{2}\mathbb{Z}$ such that $-j \leq m \leq j$ and $j \pm m$ are integers, the weight basis $\{v_m^j\}_{-j \leq m \leq j}$ for $\text{Sym}^{2j}W \otimes \det^{(j+n)}$ is defined as:

$$v_m^j = \frac{z_1^{j-m} z_2^{j+m}}{\sqrt{(j-m)!(j+m)!}}.$$

There is a hermitian inner product $\langle v_{m_1}^j, v_{m_2}^j \rangle = \delta_{m_1, m_2}$ on $\text{Sym}^{2j}W \otimes \det^{(j+n)}$. The *Wigner D-functions* $W_{m_1, m_2}^{(j, n)}(\zeta, \psi, \theta, \phi)$ are the matrix coefficients of the irreducible representation $\pi_{j,n}$ under this hermitian inner product:

$$\begin{aligned} W_{m_1, m_2}^{(j, n)}(\zeta, \psi, \theta, \phi) &= \langle v_{m_1}^j, e^{-\gamma_0\zeta}\mathcal{U}(\psi, \theta, \phi)v_{m_2}^j \rangle \\ (A.2) \quad &= c_{m_1}^j c_{m_2}^j e^{in\zeta} e^{i(m_1\psi + m_2\phi)} d_{m_1, m_2}^{(j, n)}(\theta), \end{aligned}$$

where $c_m^j = \sqrt{(j+m)!(j-m)!}$ is a normalization factor, and the function $d_{m_1, m_2}^{(j, n)}(\theta)$ is given by the trigonometric polynomial

$$\begin{aligned} d_{m_1, m_2}^{(j, n)}(\theta) &= \sum_{p=\max(0, m_1-m_2)}^{\min(j-m_2, j+m_1)} \frac{(-1)^{m_2-m_1+p}}{(j+m_1-p)!p!(m_2-m_1+p)!(j-m_2-p)!} \\ (A.3) \quad &\sin^{m_2-m_1+2p} \left(\frac{\theta}{2} \right) \cos^{2j+m_1-m_2-2p} \left(\frac{\theta}{2} \right). \end{aligned}$$

which can also be expressed in terms of Jacobi polynomials:

$$(A.4) \quad d_{m_1, m_2}^{(j, n)}(\theta) = \frac{\left(\sin \frac{\theta}{2}\right)^{m_1-m_2} \left(\cos \frac{\theta}{2}\right)^{m_1+m_2}}{(j+m_2)!(j-m_2)!} P_{j-m_1}^{(m_1-m_2, m_1+m_2)}(\cos \theta).$$

For $n \geq 0$ and for $\alpha, \beta \in \mathbb{R}$, the *Jacobi polynomials* $P_n^{(\alpha, \beta)}(z)$ are a class of orthogonal polynomials defined in [AS67] as

(A.5)

$$P_n^{(\alpha, \beta)}(z) = \frac{\Gamma(\alpha+n+1)}{n! \Gamma(\alpha+\beta+n+1)} \sum_{m=0}^n \binom{n}{m} \frac{\Gamma(\alpha+\beta+n+m+1)}{\Gamma(\alpha+m+1)} \left(\frac{z-1}{2} \right)^m.$$

In [Res08], the Jacobi polynomial is also defined as

$$(A.6) \quad P_n^{\alpha, \beta}(x) = \binom{n+\alpha}{n} \left(\frac{x+1}{2} \right)^n {}_2F_1 \left(-n, -n-\beta, \alpha+1; \frac{x-1}{x+1} \right).$$

There is a left and right regular action by γ_i on these Wigner D -matrices:

$$(A.7) \quad \begin{aligned} dr(\gamma_0)W_{m_1, m_2}^{(j, n)} &= -inW_{m_1, m_2}^{(j, n)} & dl(\gamma_0)W_{m_1, m_2}^{(j, n)} &= inW_{m_1, m_2}^{(j, n)} \\ dr(\gamma_3)W_{m_1, m_2}^{(j, n)} &= -im_2W_{m_1, m_2}^{(j, n)} & dl(\gamma_3)W_{m_1, m_2}^{(j, n)} &= im_1W_{m_1, m_2}^{(j, n)} \end{aligned}$$

$$(A.8) \quad dr(\gamma_1 \pm i\gamma_2)W_{m_1, m_2}^{(j, n)} = i\sqrt{(j \pm m_2)(j \mp m_2 + 1)}W_{m_1, m_2 \mp 1}^{(j, n)}$$

$$(A.9) \quad dl(\gamma_1 \pm i\gamma_2)W_{m_1, m_2}^{(j, n)} = -i\sqrt{(j \mp m_1)(j \pm m_1 + 1)}W_{m_1 \pm 1, m_2}^{(j, n)}.$$

The product of Wigner D -functions $W_{m_{11}, m_{12}}^{(j_1, n_1)} W_{m_{21}, m_{22}}^{(j_2, n_2)}$ can be expressed in terms of other Wigner D -matrices in terms of the *Clebsch-Gordan coefficients*:

$$(A.10) \quad W_{m_{11}, m_{12}}^{(j_1, n_1)} W_{m_{21}, m_{22}}^{(j_2, n_2)} = \sum_{\substack{|j_1 - j_2| \leq J \leq j_1 + j_2 \\ J - |j_1 - j_2| \in \mathbb{Z} \\ M_1 = m_{11} + m_{21} \\ M_2 = m_{12} + m_{22}}} \begin{pmatrix} J, M_1 \\ j_1, m_{11}, j_2, m_{21} \end{pmatrix} \begin{pmatrix} J, M_2 \\ j_1, m_{12}, j_2, m_{22} \end{pmatrix} W_{M_1, M_2}^{(J, n_1 + n_2)}.$$

The Clebsch-Gordan coefficients for $V^j \otimes V^1$ are listed in the following table:

$\begin{pmatrix} j+j_0, m_1+m_2 \\ j, m_1, 1, m_2 \end{pmatrix}$	$m_2 = -1$	$m_2 = 0$	$m_2 = 1$
$j_0 = -1$	$\sqrt{\frac{(j+m_1)(j+m_1-1)}{2j(2j+1)}}$	$-\sqrt{\frac{(j-m_1)(j+m_1)}{j(2j+1)}}$	$\sqrt{\frac{(j-m_1)(j-m_1-1)}{2j(2j+1)}}$
$j_0 = 0$	$\sqrt{\frac{(j+m_1)(j-m_1+1)}{2j(j+1)}}$	$\frac{m_1}{\sqrt{j(j+1)}}$	$-\sqrt{\frac{(j-m_1)(j+m_1+1)}{2j(j+1)}}$
$j_0 = 1$	$\sqrt{\frac{(j-m_1+1)(j-m_1+2)}{(2j+2)(2j+1)}}$	$\sqrt{\frac{(j-m_1+1)(j+m_1+1)}{(j+1)(2j+1)}}$	$\sqrt{\frac{(j+m_1+1)(j+m_1+2)}{(2j+2)(2j+1)}}$

TABLE 2. Table for Clebsch-Gordan coefficients of $V^j \otimes V^1$

REFERENCES

- [AS67] Milton Abramowitz and Irene A. Stegun. Handbook of mathematical functions: with formulas, graphs, and mathematical tables. 1967.
- [Bat55] Harry Bateman. Higher transcendental functions. *California Institute of Technology Bateman Manuscript Project, New York: McGraw-Hill, 1953-1955*, 1955.
- [BM17] Jack Buttcane and Stephen D. Miller. Weights, raising and lowering operators, and K -types for automorphic forms on $SL(3, \mathbb{R})$. *arXiv preprint arXiv:1702.08851*, 2017.
- [Kna13] Anthony W. Knap. *Lie groups beyond an introduction*, volume 140. Springer Science & Business Media, 2013.
- [Kna16] Anthony W. Knap. *Representation Theory of Semisimple Groups: An Overview Based on Examples (PMS-36)*. Princeton University Press, 2016.
- [KS80] Anthony W. Knap and Elias M. Stein. Intertwining operators for semisimple groups, II. *Inventiones mathematicae*, 60(1):9–84, 1980.
- [KZ76] Anthony W. Knap and Gregg Zuckerman. Classification of irreducible tempered representations of semisimple Lie groups. *Proceedings of the National Academy of Sciences*, 73(7):2178–2180, 1976.
- [Mui09] Goran Muić. Intertwining operators and composition series of generalized and degenerate principal series for $Sp(4, \mathbb{R})$. *Glasnik matematički*, 44(2):349–399, 2009.
- [Obe12] Fritz Oberhettinger. *Tables of Mellin transforms*. Springer Science & Business Media, 2012.
- [OLBC10] Frank W.J. Olver, D.W. Lozier, R.F. Boisvert, and C.W. Clark. NIST digital library of mathematical functions. *Online companion to [65]: http://dlmf.nist.gov*, 2010.
- [Res08] Wolfram Research. Wolfram Functions Site. <http://functions.wolfram.com>, 2008. [Online; Modification Date 2001-10-29].

- [Sha10] Freydoon Shahidi. *Eisenstein series and automorphic L -functions*, volume 58. American Mathematical Soc., 2010.
- [SM84] H Srivastava and H.L. Manocha. *Treatise on generating functions*. John Wiley & Sons, 1984.
- [VW90] D.A. Vogan and N.R. Wallach. Intertwining operators for real reductive groups. *Advances in Mathematics*, 82(2):203–243, 1990.
- [Yan11] Jae-Hyun Yang. Invariant differential operators on Siegel-Jacobi space. *arXiv preprint arXiv:1107.0509*, 2011.
- [Zha19] Zhuohui Zhang. Principal series representation of $SU(2, 1)$ and its intertwining operator. *arXiv preprint arXiv:1909.01096*, 2019.

E-mail address: zhuohui.zhang@weizmann.ac.il