

Quantum Brascamp-Lieb Dualities

Mario Berta,¹ David Sutter,² and Michael Walter^{3,4}

¹*Institute for Quantum Information, RWTH Aachen University, Aachen, Germany*

²*IBM Quantum, IBM Research Europe, Zürich, Switzerland*

³*Faculty of Computer Science, Ruhr University Bochum, Germany*

⁴*Korteweg-de Vries Institute for Mathematics,*

Institute for Theoretical Physics, Institute for Language,

Logic & Computation, and QuSoft, University of Amsterdam, Netherlands

Brascamp-Lieb inequalities are entropy inequalities which have a dual formulation as generalized Young inequalities. In this work, we introduce a fully quantum version of this duality, relating quantum relative entropy inequalities to matrix exponential inequalities of Young type. We demonstrate this novel duality by means of examples from quantum information theory—including entropic uncertainty relations, strong data-processing inequalities, super-additivity inequalities, and many more. As an application we find novel uncertainty relations for Gaussian quantum operations that can be interpreted as quantum duals of the well-known family of ‘geometric’ Brascamp-Lieb inequalities.

I. INTRODUCTION

The classical Brascamp-Lieb (BL) problem asks, given a finite sequence of surjective linear maps $L_k: \mathbb{R}^m \rightarrow \mathbb{R}^{m_k}$ and $q_k \in \mathbb{R}_+$ for $k \in [n]$, for the optimal constant $C \in \mathbb{R}$ such that [7, 10, 14, 47]

$$\int_{\mathbb{R}^m} \prod_{k=1}^n f_k(L_k x) dx \leq \exp(C) \prod_{k=1}^n \|f_k\|_{1/q_k} \quad (1)$$

holds for all non-negative functions $f_k: \mathbb{R}^{m_k} \rightarrow \mathbb{R}_+$, $k \in [n]$, where $\|\cdot\|_p$ denotes the p -norm. Many classical integral inequalities fall into this framework, such as the Hölder inequality, Young’s inequality, and the Loomis-Whitney inequality. A celebrated theorem by Lieb asserts that the optimal constant in Eq. (1) can be computed by optimizing over centred Gaussians f_k alone [47].

Remarkably, Eq. (1) has a dual, entropic formulation in terms of the *differential entropy* $H(g) := -\int g(x) \log g(x) dx$. Namely, Eq. (1) holds for all f_1, \dots, f_n as above if, and only if, for all probability densities g on \mathbb{R}^m with finite differential entropy, we have [19]

$$H(g) \leq \sum_{k=1}^n q_k H(g_k) + C. \quad (2)$$

Here, g_k denotes the marginal probability density on \mathbb{R}^{m_k} corresponding to L_k , i.e., the push-forward of g along L_k defined by $\int_{\mathbb{R}^m} \phi(L_k x) g(x) dx = \int_{\mathbb{R}^{m_k}} \phi(y) g_k(y) dy$ for all bounded, continuous functions ϕ on \mathbb{R}^{m_k} . The duality between Eq. (1) and Eq. (2) readily generalizes to arbitrary measure spaces and measurable maps [19].

Of particular interest is the so-called *geometric* case where each L_k is a surjective partial isometry and $\sum_{k=1}^n q_k L_k^\dagger L_k = \mathbb{1}_{\mathbb{R}^m}$ [2–7]. In this case, Eq. (1) and Eq. (2) hold with $C = 0$. This setup includes the Hölder and Loomis-Whitney inequalities. Equivalently, we are given n subspaces $V_k \subseteq \mathbb{R}^m$ (the supports of the L_k) such that $\sum_{k=1}^n q_k \Pi_k = \mathbb{1}_{\mathbb{R}^m}$, where Π_k denotes

the orthogonal projection onto V_k . In this case we can think of the marginal densities g_k as functions on V_k , namely

$$g_{V_k}(y) = \int_{V_k^\perp} g(y+z) dz \quad \forall y \in V_k. \quad (3)$$

In particular, if V_k is a coordinate subspace of \mathbb{R}^m then g_{V_k} is nothing but the usual marginal probability density of the corresponding random variables, justifying our terminology. As a concrete example, let V_1, V_2 be the two coordinate subspaces of \mathbb{R}^2 and $q_1 = q_2 = 1$; then Eq. (2) amounts to the sub-additivity property of the differential entropy, which is dual to the trivial estimate $\int_{\mathbb{R}^2} f_1(x_1)f_2(x_2) dx \leq \|f_1\|_1\|f_2\|_1$. In contrast, already for three equiangular lines in \mathbb{R}^2 (a ‘Mercedes star’ configuration) and $q_1 = q_2 = q_3 = \frac{2}{3}$, neither inequality is immediate.

Recently, the BL duality has been extended on the entropic side to not only include entropy inequalities as in Eq. (2) but also relative entropy inequalities in terms of the Kullback-Leibler divergence [52]. The dual analytic form then again corresponds to generalized Young inequalities as in Eq. (1) but now for weighted p -norms. Interestingly, this extended BL duality covers many fundamental entropic statements from information theory and more. This includes, e.g., hypercontractivity inequalities, strong data processing inequalities, and transportation-cost inequalities [53].

Here, we raise the question how aforementioned BL dualities can be extended in the non-commutative setting. Our main motivation comes from quantum information theory, where quantum entropy inequalities are pivotal and dual formulations often promise new insights. BL dualities for non-commutative integration have previously been studied by Carlen and Lieb [20]. Amongst other contributions, they gave BL dualities similar to Eq. (1) - Eq. (2) leading to generalized sub-additivity inequalities for quantum entropy.

In this paper, we extend the classical duality results of [52, 53] to the quantum setting—thereby generalizing Carlen and Lieb’s BL duality to the quantum relative entropy and general quantum channel evolutions. In particular, we derive in Section II a fully quantum BL duality for quantum relative entropy and discuss its properties. In Section III we then discuss a plethora of examples from quantum information theory that are covered by our quantum BL duality. As novel inequalities, we give quantum versions of the *geometric* Brascamp-Lieb inequalities discussed above, whose entropic form can be interpreted as an uncertainty relation for certain Gaussian quantum operations (Section IIIB).

Note added: Since the first version of our manuscript, our geometric quantum Brascamp-Lieb inequalities from Section IIIB have been extended to the conditional case [50] and to more general Gaussian quantum operations [29]. We briefly mention these extensions in Section IIIB.

Notation. Let A and B be separable Hilbert spaces. We denote the set of bounded operators on A by $L(A)$, the set of trace-class operators on A by $T(A)$, the set of Hermitian operators on A by $\text{Herm}(A)$, the set of positive operators on A by $P_{\succ}(A)$, and the set of positive semi-definite operators on A by $P_{\succeq}(A)$. A *density operator* or *quantum state* is a positive semi-definite trace-class operator with unit trace; we denote the set of density operators on A by $S(A)$. The set of trace-preserving and positive maps from $T(A)$ to $T(B)$ is denoted by $\text{TPP}(A, B)$ and the set of trace-preserving and *completely* positive maps from $T(A)$ to $T(B)$ is denoted by $\text{TPCP}(A, B)$. For $\mathcal{E} \in \text{TPP}(A, B)$ the *adjoint map* \mathcal{E}^\dagger , which is a unital and positive map from $L(B)$ to $L(A)$, is defined by $\text{tr } \mathcal{E}(X)^\dagger Y = \text{tr } X^\dagger \mathcal{E}^\dagger(Y)$ for all $X \in T(A)$ and

$Y \in L(B)$. When it is clear from the context, we sometimes leave out identity operators, i.e., we may write $\rho_A \sigma_{AB} \rho_B$ for $(\rho_A \otimes \mathbb{1}_B) \sigma_{AB} (\mathbb{1}_A \otimes \rho_B)$.

The *von Neumann entropy* of a density operator $\rho \in S(A)$ is defined as¹

$$H(\rho) := -\text{tr } \rho \log \rho$$

and can be infinite (only) if A is infinite-dimensional. The *quantum relative entropy* of $\omega \in S(A)$ with respect to $\tau \in P_{\succ}(A)$ is given by

$$D(\omega \parallel \tau) := \text{tr } \omega (\log \omega - \log \tau) \quad \text{if } \omega \ll \tau \text{ and } +\infty \text{ otherwise,}$$

where $\omega \ll \tau$ denotes that the support of ω is contained in the support of τ . The von Neumann entropy can be expressed as a relative entropy, $H(\rho) = -D(\rho \parallel \mathbb{1})$, where $\mathbb{1}$ denotes the identity operator. For $\rho_{AB} \in S(A \otimes B)$ with $H(A)_\rho < \infty$, the *conditional entropy of A given B* is defined as [45]

$$H(A|B)_\rho := H(A)_\rho - D(\rho_{AB} \parallel \rho_A \otimes \rho_B),$$

where the notation $H(A)_\rho := H(\rho_A)$ refers to the entropy of the reduced density operator $\rho_A = \text{tr}_B(\rho)$ on A . For A and B finite-dimensional we can also write $H(A|B)_\rho = H(AB)_\rho - H(B)_\rho$.

Throughout this manuscript the default is that Hilbert spaces are finite-dimensional unless explicitly stated otherwise (such as in Section III B).

II. BRASCAMP-LIEB DUALITY FOR QUANTUM RELATIVE ENTROPIES

In this section, we describe our main result (Theorem II.1) and discuss some of its mathematical properties.

A. Main result

The following result establishes a version of the Brascamp-Lieb dualities of [19, 52, 53] for quantum relative entropies.

Theorem II.1 (Quantum Brascamp-Lieb duality). *Let $n \in \mathbb{N}$, $\vec{q} = (q_1, \dots, q_n) \in \mathbb{R}_+^n$, $\vec{\mathcal{E}} = (\mathcal{E}_1, \dots, \mathcal{E}_n)$ with $\mathcal{E}_k \in \text{TPP}(A, B_k)$ for $k \in [n]$, $\sigma \in P_{\succ}(A)$, $\vec{\sigma} = (\sigma_1, \dots, \sigma_n)$ with $\sigma_k \in P_{\succ}(B_k)$ for $k \in [n]$, and $C \in \mathbb{R}$. Then, the following two statements are equivalent:*

$$\sum_{k=1}^n q_k D(\mathcal{E}_k(\rho) \parallel \sigma_k) \leq D(\rho \parallel \sigma) + C \quad \forall \rho \in S(A), \quad (4)$$

$$\text{tr} \exp \left(\log \sigma + \sum_{k=1}^n \mathcal{E}_k^\dagger(\log \omega_k) \right) \leq \exp(C) \prod_{k=1}^n \left\| \exp(\log \omega_k + q_k \log \sigma_k) \right\|_{1/q_k} \quad \forall \omega_k \in P_{\succ}(B_k), \quad (5)$$

where $\|L\|_p := (\text{tr} |L|^p)^{\frac{1}{p}}$ is the Schatten p -norm for $p \in [1, \infty]$ and an anti-norm for $p \in (0, 1]$.² Moreover, Eq. (5) holds for all $\omega_k \in P_{\succ}(B_k)$ if and only if it holds for all $\omega_k \in S(B_k)$ with full support.

¹ The case when ρ_A does not have full support is covered by the convention $0 \log 0 = 0$. Unless specified otherwise, we choose to leave the basis of the logarithm function $\log(\cdot)$ unspecified and write $\exp(\cdot)$ for its inverse function.

² An *anti-norm* is a non-negative function on $P_{\succ}(A)$ that is homogeneous ($\|\alpha\omega\| = \alpha \|\omega\|$ for $\alpha > 0$) and super-additive ($\|\omega + \omega'\| \geq \|\omega\| + \|\omega'\|$) for $\omega, \omega' \in P_{\succ}(A)$ [13]. NB: $\|\cdot\|_1$ is both a norm and an anti-norm.

We refer to Eq. (4) as a quantum Brascamp-Lieb inequality in *entropic form*, and to Eq. (5) as a quantum Brascamp-Lieb inequality in *analytic form*. The latter can be understood as a quantum version of a Young-type inequality. The two formulations in Eq. (4) and Eq. (5) encompass a large class of concrete inequalities, as we will see in Section III below; we are also often interested in identifying the smallest constant $C \in \mathbb{R}$ such that either inequality holds. To this end, both directions of Theorem II.1 are of interest:

1. To prove quantum entropy inequalities, Theorem II.1 allows us to alternatively work with matrix exponential inequalities in the analytic form. That this approach can give crucial insights was already discovered in the original proof of strong sub-additivity of the von Neumann entropy [49], which relied on Lieb's triple matrix inequality for the exponential function (see also [31, 60] for more recent works). We discuss similar examples in Section III C.
2. In the commutative setting, we know that for deriving Young-type inequalities it can be beneficial to work in the entropic form [19, 21]. As the quantum relative entropy has natural properties mirroring its classical counterpart, this translates to the non-commutative setting. We discuss corresponding examples in Section III A and Section III B.

The proof of Theorem II.1 relies on the following formula for the Legendre transform of the quantum relative entropy and its dual.

Fact II.2 (Variational formula for quantum relative entropy [56]). *Let $\sigma \in \mathcal{P}_{>}(A)$. Then:*

- *For all $\rho \in \mathcal{S}(A)$ we have*

$$D(\rho\|\sigma) = \sup_{\omega \in \mathcal{P}_{\geq}(A)} \{ \text{tr } \rho \log \omega - \log \text{tr} \exp(\log \omega + \log \sigma) \} . \quad (6)$$

Furthermore, the supremum is attained for $\omega = \exp(\log \rho - \log \sigma) / \text{tr} \exp(\log \rho - \log \sigma)$.

- *For all $H \in \text{Herm}(A)$, we have*

$$\log \text{tr} \exp(H + \log \sigma) = \sup_{\omega \in \mathcal{S}(A)} \{ \text{tr } H \omega - D(\omega\|\sigma) \} . \quad (7)$$

Furthermore, the supremum is attained for $\omega = \exp(H + \log \sigma) / \text{tr} \exp(H + \log \sigma)$.

These variational formulas are powerful on their own for proving quantum entropy inequalities, as, e.g., the first term in Eq. (6) only depends on ρ (but not on σ) and the second term only on σ (but not on ρ). We refer to [60] for a more detailed discussion.

We mention that Carlen-Lieb use the variational characterization of the von Neumann entropy to derive Brascamp-Lieb dualities and [20, bottom of page 564] commented that their proof strategy extends to the relative entropy via Petz's variational expression for the relative entropy (Fact II.2), which is what is done here.

Proof of Theorem II.1. We first show that Eq. (4) implies Eq. (5). Let $H_k := \log \omega_k$ and define $H \in \text{Herm}(A)$ and $\rho \in \mathcal{S}(A)$ by

$$H := \sum_{k=1}^n \mathcal{E}_k^\dagger(H_k) \quad \text{and} \quad \rho := \frac{\exp(H + \log \sigma)}{\text{tr} \exp(H + \log \sigma)} , \quad (8)$$

respectively. Then,

$$\begin{aligned}
\log \operatorname{tr} \exp \left(\log \sigma + \sum_{k=1}^n \mathcal{E}_k^\dagger(H_k) \right) &= \log \operatorname{tr} \exp(H + \log \sigma) \\
&= \operatorname{tr} H \rho - D(\rho \| \sigma) \\
&= \sum_{k=1}^n \operatorname{tr} \mathcal{E}_k^\dagger(H_k) \rho - D(\rho \| \sigma) \\
&\leq C + \sum_{k=1}^n q_k \left(\operatorname{tr} \frac{H_k}{q_k} \mathcal{E}_k(\rho) - D(\mathcal{E}_k(\rho) \| \sigma_k) \right) \\
&\leq C + \sum_{k=1}^n q_k \log \operatorname{tr} \exp \left(\frac{H_k}{q_k} + \log \sigma_k \right),
\end{aligned}$$

where we used Eq. (7) in both the second and the last step and Eq. (4) in the penultimate step. By substituting $H_k = \log \omega_k$ and taking the exponential on both sides we obtain Eq. (5).

We now show that, conversely, Eq. (5) implies Eq. (4). Let $\omega = \exp(H)$, with H defined as in Eq. (8) in terms of $H_k = \log(\omega_k)$ for $\omega_k \in P_{\succ_{\sigma_k}}(B_k)$ that we will choose later. Then, using Eq. (6),

$$\begin{aligned}
D(\rho \| \sigma) &\geq \operatorname{tr} \rho \log \omega - \log \operatorname{tr} \exp(\log \omega + \log \sigma) \\
&= \sum_{k=1}^n \operatorname{tr} \rho \mathcal{E}_k^\dagger(H_k) - \log \operatorname{tr} \exp \left(\sum_{k=1}^n \mathcal{E}_k^\dagger(H_k) + \log \sigma \right) \\
&= \sum_{k=1}^n \operatorname{tr} \mathcal{E}_k(\rho) \log \omega_k - \log \operatorname{tr} \exp \left(\log \sigma + \sum_{k=1}^n \mathcal{E}_k^\dagger(\log \omega_k) \right) \\
&\geq \sum_{k=1}^n q_k \left(\operatorname{tr} \mathcal{E}_k(\rho) \frac{\log \omega_k}{q_k} - \log \operatorname{tr} \exp \left(\frac{\log \omega_k}{q_k} + \log \sigma_k \right) \right) - C \\
&= \sum_{k=1}^n q_k D(\mathcal{E}_k(\rho) \| \sigma_k) - C,
\end{aligned}$$

where the last inequality uses Eq. (5) and the final step follows from Eq. (6) provided we choose ω_k^{1/q_k} as the maximizer for the variational expression of $D(\mathcal{E}_k(\rho) \| \sigma_k)$. \square

Remark II.3. As the variational characterizations from Fact II.2 hold in the general W^* -algebra setting [56], the BL duality in Theorem II.1 extends to separable Hilbert spaces.

Remark II.4. The BL duality in Theorem II.1 can be extended to $\sigma \in P_{\succeq}(A)$ and $\vec{\sigma} = (\sigma_1, \dots, \sigma_n)$ with $\sigma_k \in P_{\succeq}(B_k)$ for $k \in [n]$ when

1. $\mathcal{E}_k(\rho) \ll \sigma_k$ for all $\rho \in S(A)$ with $\rho \ll \sigma$
2. $\mathcal{E}^\dagger(\log \omega_k) \ll \sigma$ for all $\omega_k \in P_{\succeq}(B)$ with $\omega_k \ll \sigma_k$.

Then, the BL duality still holds but for the alternative conditions

$$\rho \in S(A) \text{ with } \rho \ll \sigma \text{ in Eq. (4) \quad and \quad } \omega_k \in P_{\succ}(B_k) \text{ with } \omega_k \ll \sigma_k \text{ in Eq. (5).}$$

To see this, note that the variational formula in Eq. (6) still holds for $\sigma \in P_{\succ}(A)$ as long as $\rho \ll \sigma$ with the supremum taken over $\omega \in P_{\succ}(A)$ with $\omega \ll \sigma$. Similarly, Eq. (7) still holds for $H \in \text{Herm}(A)$ for $H \ll \sigma$ with the supremum taken over $\omega \in S(A)$ with $\omega \ll \sigma$. The proof of Theorem II.1 then also goes through in the more general form.

In many important applications, we are interested in using Theorem II.1 either in the situation that $\sigma_k = \mathcal{E}_k(\sigma)$ for all $k \in [n]$, or in a setting where $\sigma = \mathbb{1}_A$ and $\sigma_k = \mathbb{1}_{B_k}$ for all $k \in [n]$. In the latter case, Theorem II.1 specializes to the following equivalence between von Neumann entropy inequalities and Young-type inequalities:

Corollary II.5. *Let $n \in \mathbb{N}$, $\vec{q} = (q_1, \dots, q_n) \in \mathbb{R}_+^n$, $\vec{\mathcal{E}} = (\mathcal{E}_1, \dots, \mathcal{E}_n)$ with $\mathcal{E}_k \in \text{TPP}(A, B_k)$ for $k \in [n]$, and $C \in \mathbb{R}$. Then, the following two statements are equivalent:*

$$H(\rho) \leq \sum_{k=1}^n q_k H(\mathcal{E}_k(\rho)) + C \quad \forall \rho \in S(A), \quad (9)$$

$$\text{tr exp} \left(\sum_{k=1}^n \mathcal{E}_k^\dagger(\log \omega_k) \right) \leq \exp(C) \prod_{k=1}^n \|\omega_k\|_{1/q_k} \quad \forall \omega_k \in S(B_k). \quad (10)$$

Carlen and Lieb previously proved a variant of Corollary II.5 in the W^* -algebra setting assuming that the maps \mathcal{E}_k^\dagger are W^* -homomorphisms and that $q_k \in [0, 1]$ [20, Theorem 2.2]. One interesting special case is when the \mathcal{E}_k are partial trace maps. The entropic form Eq. (9) then corresponds to generalized sub-additivity inequalities for the von Neumann entropy (cf. Section III A).

B. Weighted anti-norms

In the commutative setting, the right-hand side of Eq. (5) can conveniently be understood as a product of σ_k -weighted norms or anti-norms of the operators ω_k [52, 53]. It is natural to ask whether such an interpretation also holds quantumly. To this end, given $p \in (0, 1]$ and $\sigma \in P_{\succ}(A)$, define

$$\|\omega\|_{\sigma,p} := (\text{tr exp}(\log \omega^p + \log \sigma))^{\frac{1}{p}} = \left\| \exp \left(\log \omega + \frac{1}{p} \log \sigma \right) \right\|_p,$$

for all $\omega \in P_{\succ}(A)$. The following proposition, which follows readily from [41], shows that $\|\cdot\|_{\sigma,p}$ is an anti-norm provided that $p \leq 1$. For $p > 1$, it is easy to find $\sigma \in P_{\succ}(A)$ such that the functional $\|\cdot\|_{\sigma,p}$ is neither a norm nor an anti-norm.

Proposition II.6. *For $p \in (0, 1]$ and $\sigma \in P_{\succ}(A)$, $\|\cdot\|_{\sigma,p}$ is homogeneous and concave, hence an anti-norm.*

Proof. Clearly, $\|\cdot\|_{\sigma,p}$ is homogeneous. Since moreover $p \in (0, 1]$, [41, Lemma D.1] asserts that its concavity on the set of positive matrices is equivalent to the concavity of its p -th power, i.e.,

$$\omega \mapsto \text{tr exp}(p \log \omega + H), \quad (11)$$

where $H = \log \sigma$. A well-known result of Lieb [46] states that Eq. (11) is indeed concave for any Hermitian matrix H . Thus, $\|\cdot\|_{\sigma,p}$ is concave. As a consequence of homogeneity and concavity, we obtain that $\|\cdot\|_{\sigma,p}$ is super-additive, as $\|\omega + \omega'\|_{\sigma,p} = 2 \left\| \frac{1}{2}\omega + \frac{1}{2}\omega' \right\|_{\sigma,p} \geq \|\omega\|_{\sigma,p} + \|\omega'\|_{\sigma,p}$ for all $\omega, \omega' \in P_{\succ}(A)$. We conclude that $\|\cdot\|_{\sigma,p}$ is an anti-norm. \square

Thus, the quantum Brascamp-Lieb inequality in its analytic form Eq. (5) can be written as

$$\mathrm{tr} \exp \left(\log \sigma + \sum_{k=1}^n \mathcal{E}_k^\dagger (\log \omega_k) \right) \leq \exp(C) \prod_{k=1}^n \|\omega_k\|_{\sigma_k, 1/q_k} \quad \forall \omega_k \in \mathcal{S}(B_k), \quad (12)$$

where, assuming that all $q_k \geq 1$, the right-hand side can be interpreted in terms of anti-norms, pleasantly generalizing Eq. (10).

C. Convexity and tensorization

For fixed $n \in \mathbb{N}$, $\vec{\mathcal{E}} = (\mathcal{E}_1, \dots, \mathcal{E}_n)$ with $\mathcal{E}_k \in \mathrm{TPP}(A, B)$, $\sigma \in \mathcal{P}_{\succ}(A)$, and $\vec{\sigma} = (\sigma_1, \dots, \sigma_n)$ with $\sigma_k \in \mathcal{P}_{\succ}(B_k)$, we define the *Brascamp-Lieb (BL) set* as

$$\mathrm{BL}(\vec{\mathcal{E}}, \vec{\sigma}, \sigma) := \left\{ (\vec{q}, C) \in \mathbb{R}_+^n \times \mathbb{R} : \text{Eq. (4)/Eq. (5) holds} \right\}.$$

We record the following elementary property.

Proposition II.7 (Convexity). *The set $\mathrm{BL}(\vec{\mathcal{E}}, \vec{\sigma}, \sigma)$ is convex.*

Proof. We use the characterization using the entropic form Eq. (4). Let $(\vec{q}^{(i)}, C^{(i)}) \in \mathrm{BL}(\vec{\mathcal{E}}, \vec{\sigma}, \sigma)$ for $i \in \{1, 2\}$. Let $\theta \in [0, 1]$ and (\vec{q}, C) the corresponding convex combination, i.e., $\vec{q} := \theta \vec{q}^{(1)} + (1 - \theta) \vec{q}^{(2)}$ and $C := \theta C^{(1)} + (1 - \theta) C^{(2)}$. Then, for all $\rho \in \mathcal{S}(A)$,

$$\begin{aligned} \sum_{k=1}^n q_k D(\mathcal{E}_k(\rho) \| \sigma_k) &= \theta \sum_{k=1}^n q_k^{(1)} D(\mathcal{E}_k(\rho) \| \sigma_k) + (1 - \theta) \sum_{k=1}^n q_k^{(2)} D(\mathcal{E}_k(\rho) \| \sigma_k) \\ &\leq \theta \left(D(\rho \| \sigma) + C^{(1)} \right) + (1 - \theta) \left(D(\rho \| \sigma) + C^{(2)} \right) = D(\rho \| \sigma) + C. \end{aligned}$$

Thus, $(\vec{q}, C) \in \mathrm{BL}(\vec{\mathcal{E}}, \vec{\sigma}, \sigma)$. □

In the commutative case, the BL set satisfies a *tensorization property* [53, Section V.B], and we can ask if a similar property holds in the non-commutative case as well. Namely, do we have that for $(\vec{q}, C^{(i)}) \in \mathrm{BL}(\vec{\mathcal{E}}^{(i)}, \vec{\sigma}^{(i)}, \sigma^{(i)})$ with $i \in \{1, 2\}$ and

$$\vec{\mathcal{E}} := \left(\mathcal{E}_1^{(1)} \otimes \mathcal{E}_1^{(2)}, \dots, \mathcal{E}_n^{(1)} \otimes \mathcal{E}_n^{(2)} \right) \quad \text{as well as} \quad \vec{\sigma} := \left(\sigma_1^{(1)} \otimes \sigma_1^{(2)}, \dots, \sigma_n^{(1)} \otimes \sigma_n^{(2)} \right)$$

that

$$\left(\vec{q}, C^{(1)} + C^{(2)} \right) \stackrel{?}{\in} \mathrm{BL} \left(\vec{\mathcal{E}}, \vec{\sigma}, \sigma^{(1)} \otimes \sigma^{(2)} \right). \quad (13)$$

As we will see in several examples (Section III), tensorization does in general not hold due to the potential presence of entanglement. Indeed, the problem of deciding in which case Eq. (13) holds can be understood as a general information-theoretic *additivity problem*, which contains the (non-)additivity for the minimum output entropy as a special case (cf. Eq. (39) in Section IIID).

III. APPLICATIONS OF QUANTUM BRASCAMP-LIEB DUALITY

The purpose of this section is to present examples from quantum information theory where the duality from Theorem II.1 is applicable. The majority of examples concern entropy inequalities that are of interest from an operational viewpoint. Theorem II.1 then shows that all entropy inequalities of suitable structure have a dual formulation as an analytic inequality, and vice versa. Depending on the scenario, one form may be easier to prove than the other, and we find that these reformulations often give additional insight.

A. Generalized (strong) sub-additivity

In this section, we discuss entropy inequalities that generalize the sub-additivity and strong sub-additivity properties of the von Neumann entropy. Recall that the latter states that $H(AB) + H(BC) \geq H(ABC) + H(B)$ for $\rho_{ABC} \in \mathcal{S}(A \otimes B \otimes C)$ [49].

We first state the following result from [20, Theorem 1.4 & Theorem 3.1], which gives generalized sub-additivity relations and their dual analytic form. Here, the second argument in the relative entropy is always equal to the identity. Throughout this section, all quantum channels are given by partial trace channels.

Corollary III.1 (Quantum Shearer and Loomis-Whitney inequalities, [20]). *Let S_1, \dots, S_n be non-empty subsets of $[m]$ such that every $s \in [m]$ belongs to at least p of those subsets. Then, the following inequalities hold and are equivalent:*

$$H(A_1 \dots A_m) \leq \frac{1}{p} \sum_{k=1}^n H(\{A_s\}_{s \in S_k}) \quad \forall \rho \in \mathcal{S}(A_1 \otimes \dots \otimes A_m), \quad (14)$$

$$\mathrm{tr} \exp \left(\sum_{k=1}^n \mathbb{1}_{\bar{S}_k} \otimes \log \omega_{S_k} \right) \leq \prod_{k=1}^n \|\omega_{S_k}\|_p \quad \forall \omega_{S_k} \in \mathcal{S}(\otimes_{s \in S_k} A_s), \quad (15)$$

where \bar{S} denotes the complement of a subset S of $[m]$.

Inequalities in the form of Eq. (14) have been termed *quantum Shearer's inequalities* and their analytic counterparts as in Eq. (15) are known as *quantum Loomis-Whitney inequalities*. Interestingly, and as explained in [20, Section 1.3], the latter cannot directly be deduced from standard matrix trace inequalities such as Golden-Thompson combined with Cauchy-Schwarz. That Eq. (14) and Eq. (15) are equivalent follows from Corollary II.5 by choosing $C = 0$, $q_k = \frac{1}{p}$, and $\mathcal{E}_k(\cdot) = \mathrm{tr}_{\bar{S}_k}(\cdot)$. The following result provides a conditional version of the quantum Shearer inequality with side information.

Proposition III.2 (Conditional quantum Shearer inequality). *Let S_1, \dots, S_n be non-empty subsets of $[m]$ such that every $s \in [m]$ belongs to exactly p of those subsets. Then,*

$$H(A_1 \dots A_m | B) \leq \frac{1}{p} \sum_{k=1}^n H(\{A_s\}_{s \in S_k} | B) \quad \forall \rho \in \mathcal{S}(A_1 \otimes \dots \otimes A_m \otimes B). \quad (16)$$

For $n = 2$, $S_1 = \{1\}$, $S_2 = \{2\}$, $p = 1$, Eq. (16) reduces to $H(A_1 A_2 | B) \leq H(A_1 | B) + H(A_2 | B)$, which is equivalent to the strong sub-additivity of von Neumann entropy.³

Note that, in contrast to Corollary III.1, in the conditional case it is not enough to assume that every $s \in [m]$ belongs to at least p of the subsets. This is clear from the following proof. For a concrete counterexample, note that for $n = 2$, $S_1 = S_2 = \{1\}$, $S_3 = \{2\}$, $p = 1$, Eq. (16) is violated for, e.g., a maximally entangled state between A_1 and B .

Proof of Corollary III.1 and Proposition III.2. We adapt the argument of [20] to the conditional case. If S and T are two subsets of $[m]$ then strong sub-additivity implies that

$$H(\{A_s\}_{s \in S \cup T} | B) + H(\{A_s\}_{s \in S \cap T} | B) \leq H(\{A_s\}_{s \in S} | B) + H(\{A_s\}_{s \in T} | B).$$

This means that we obtain a stronger version of Eq. (16) if we replace any two subsets S_k , S_l by $S_k \cup S_l$, $S_k \cap S_l$. Moreover, each such replacement preserves the number of times that any $s \in [m]$ is contained in the subsets S_1, \dots, S_n . We can successively apply replacement steps until we arrive at the situation where $S_k \subseteq S_l$ or $S_l \subseteq S_k$ for any two subsets. Without loss of generality, this means that it suffices to prove Corollary III.1 and Proposition III.2 in the case that $S_1 \supseteq \dots \supseteq S_n$. In this case, $S_1 = \dots = S_p = [m]$, since each $s \in [m]$ is contained in at least p of the subsets. The corresponding inequality Eq. (16) can thus be simplified to

$$0 \leq \sum_{k=p+1}^n H(\{A_s\}_{s \in S_k} | B).$$

If $B = \emptyset$, as in Corollary III.1, this inequality holds since the von Neumann entropy is never negative. And if each $s \in [m]$ belongs to *exactly* p of the subsets, as in Proposition III.2, then $S_{p+1} = \dots = S_n = \emptyset$, so the inequality holds trivially. \square

Remark III.3. Corollary III.1 and Proposition III.2 also hold for separable Hilbert spaces, as the variational characterizations from Fact II.2 hold in the general W^* -algebra setting [56].

B. Brascamp-Lieb inequalities for Gaussian quantum operations

In this section, we present quantum versions of the classical Brascamp-Lieb inequalities as in Eq. (1) and Eq. (2), where probability distributions on \mathbb{R}^m are replaced by quantum states on $L^2(\mathbb{R}^m)$, the Hilbert space of square-integrable wave functions on \mathbb{R}^m . We focus on the *geometric* case discussed in the introduction. The marginal distribution with respect to a subspace $X \subseteq \mathbb{R}^m$ has the following natural quantum counterpart. Define a TPCP map \mathcal{E}_X as the composition of the unitary $L^2(\mathbb{R}^m) \cong L^2(X) \otimes L^2(X^\perp)$ with the partial trace over the second tensor factor. Given a density operator ρ on $L^2(\mathbb{R}^m)$, we can think of

$$\rho_X = \mathcal{E}_X(\rho)$$

as the *reduced density operator corresponding to X* . This is the natural quantum version of the marginal probability density in Eq. (3) of the introduction. Indeed, if we identify ρ with its

³ Our quantum BL duality (Theorem II.1) does not directly provide a dual analytic form for the strong sub-additivity of von Neumann entropy or more generally Eq. (16). Rather, in Section III E we provide a dual analytic form for the (a priori more general) *data processing inequality* of the quantum relative entropy.

kernel in $L^2(\mathbb{R}^m \times \mathbb{R}^m)$, and likewise for ρ_k , then we have the completely analogous formula

$$\rho_k(y, y') = \int_{X^\perp} \rho(y + z, y' + z) dz \quad \forall y, y' \in X.$$

This definition is very similar in spirit to the quantum addition operation in the quantum entropy power inequality of [43] (see also [28, 44]) and in fact contains the latter as a special case. In linear optical terms, ρ_X can be interpreted as the reduced state of $\dim X$ many output modes obtained by subjecting ρ to a network of beamsplitters with arbitrary transmissivities.

The following result establishes quantum versions of the Brascamp-Lieb dualities as in Eq. (1) and Eq. (2) for the geometric case.

Proposition III.4 (Geometric quantum Brascamp-Lieb inequalities). *Let $X_1, \dots, X_n \subseteq \mathbb{R}^m$ be subspaces and let $q_1, \dots, q_n \geq 0$ such that $\sum_{k=1}^n q_k \Pi_k = \mathbb{1}_{\mathbb{R}^m}$, where Π_k denotes the orthogonal projection onto X_k . Then, for all $\rho \in S(L^2(\mathbb{R}^m))$ with finite second moments,*

$$H(\rho) \leq \sum_{k=1}^n q_k H(\rho_{X_k}). \quad (17)$$

Furthermore, for all $\omega_{X_k} \in S(L^2(X_k))$ such that $\exp\left(\sum_{k=1}^n \mathbb{1}_{X_k^\perp} \otimes \log \omega_{X_k}\right)$ has finite second moments, it holds that⁴

$$\mathrm{tr} \exp\left(\sum_{k=1}^n \mathbb{1}_{X_k^\perp} \otimes \log \omega_{X_k}\right) \leq \prod_{k=1}^n \|\omega_{X_k}\|_{1/q_k}. \quad (18)$$

Note that if X_k is spanned by a subset $S_k \subseteq [m]$ of the m coordinates of \mathbb{R}^m , then ρ_{X_k} is nothing but the reduced density matrix of subsystems S_k , which appears on the right-hand side of the quantum Shearer inequality Eq. (14). Thus, Proposition III.4 implies Corollary III.1 in the case that all $s \in [m]$ are contained in *exactly* p of the subsets S_k .

To establish Proposition III.4, we will first prove the entropic form Eq. (17) using a quantum version of the heat flow approach from [8, 21] (cf. the recent works [22–24] on entropy inequalities for quantum Markov semigroups). We assume some familiarity with Gaussian quantum systems (see, e.g., [40]) and follow the framework of König and Smith [43], which holds under regularity assumptions on the quantum state, which were subsequently removed by De Palma and Trevisan [28].

Let $X \subseteq \mathbb{R}^m$ be a subspace and m_X its dimension. For all $x \in X$, define position and momentum operators on $L^2(X)$ by $(Q_{X,x}\psi)(y) := (x \cdot y)\psi(y)$ and $P_{X,x} := -i\partial_x$. Denote by \mathcal{N}_t^X the *non-commutative heat flow* or *heat semigroup* [28, 43], which is a one-parameter semi-group, meaning $\mathcal{N}_0^X = \mathbb{1}$ and $\mathcal{N}_s^X \circ \mathcal{N}_t^X = \mathcal{N}_{s+t}^X$ for $s, t \geq 0$. On a suitable domain it is generated by

$$\mathcal{L}_X := -\frac{1}{4} \sum_{j=1}^{m_X} [Q_{X,e_j}, [Q_{X,e_j}, \rho]] + [P_{X,e_j}, [P_{X,e_j}, \rho]],$$

⁴ As we only prove Eq. (17) for states with finite second moments, we can not apply Theorem II.1 directly to obtain Eq. (18) (see end of proof). Removing this assumption is an interesting open problem.

where $\{e_j\}_{j=1}^{m_X}$ is an arbitrary orthonormal basis of X (but we will not directly use this specific form). For every $t \geq 0$, \mathcal{N}_t^X is a Gaussian TPCP map, hence fully determined by its action on covariance matrices and mean vectors,⁵ which is given by

$$\Sigma \mapsto \Sigma + t\mathbb{1} \quad \text{and} \quad \mu \mapsto \mu. \quad (19)$$

In particular, the heat flow is independent of the choice of orthonormal basis in X . The generalized partial trace maps $\mathcal{E}_X: \rho \mapsto \rho_X$ defined above are also Gaussian and act by

$$\Sigma \mapsto \Sigma|_X \quad \text{and} \quad \mu \mapsto \mu|_X, \quad (20)$$

where $\mu|_X$ denotes the restriction of μ onto $X \oplus X$ and likewise for $\Sigma|_X$. The non-commutative heat flow is compatible with the maps \mathcal{E}_X , i.e.,

$$\mathcal{E}_X \circ \mathcal{N}_t^{\mathbb{R}^m} = \mathcal{N}_t^X \circ \mathcal{E}_X.$$

Indeed, since both channels (and hence their composition) are Gaussian, it suffices to verify that the action commutes on the level of mean vectors and covariance matrices, and the latter is clear from Eq. (19) and Eq. (20). See also [28, Lemma 2]. Thus, we may unambiguously introduce the notation

$$\rho_X^{(t)} := \mathcal{E}_X(\mathcal{N}_t^{\mathbb{R}^m}(\rho)) = \mathcal{N}_t^X(\mathcal{E}_X(\rho)) = \mathcal{N}_t^X(\rho_X) \quad (21)$$

for the reduced density operator on $L^2(X)$ at time t . Similarly, we may show that \mathcal{E}_X is compatible with phase-space translations (cf. [43, Lemma XI.1]). For $x \in X$, define the unitary one-parameter groups

$$\mathcal{Q}_{X,x}^{(t)}(\rho) := e^{-itP_{X,x}} \rho e^{itP_{X,x}} \quad \text{and} \quad \mathcal{P}_{X,x}^{(t)}(\rho) := e^{itQ_{X,x}} \rho e^{-itQ_{X,x}}.$$

They are Gaussian, leave the covariance matrices invariant, and send mean vectors $\mu \mapsto \mu + t(x^T, 0)$ and $\mu + t(0, x^T)$, respectively. By comparing with Eq. (20), we find that

$$\mathcal{Q}_{X,x}^{(t)} \circ \mathcal{E}_X = \mathcal{E}_X \circ \mathcal{Q}_{\mathbb{R}^m,x}^{(t)} \quad \text{and} \quad \mathcal{P}_{X,x}^{(t)} \circ \mathcal{E}_X = \mathcal{E}_X \circ \mathcal{P}_{\mathbb{R}^m,x}^{(t)}. \quad (22)$$

In the following we shall make use of two crucial properties of the heat flow that will allow us to ‘linearize’ the proof of the entropy inequality: First, the entropy of $\rho_X^{(t)}$ grows logarithmically as $t \rightarrow \infty$ and becomes *asymptotically independent* of the state ρ , as proved in [43, Corollary III.4] and [28, Theorem 5]:

$$|H(\rho_X^{(t)})/m_X - (1 - \log 2 + \log t)| \rightarrow 0.$$

In particular, this implies that any valid inequality of the form Eq. (17) must satisfy the inequality

$$\sum_{k=1}^n q_k m_{X_k} \geq m, \quad (23)$$

⁵ The *mean vector* of a quantum state ρ on $L^2(X)$ is the linear form μ on $X \oplus X$ given by $\mu(v) = \text{tr } \rho R_{X,v}$, where $R_{X,v} = Q_{X,x} + P_{X,y}$ for $v = (x, y)$; the *covariance matrix* of ρ is the quadratic form Σ defined by $\Sigma(v, v') = \text{tr } \rho \{R_{X,v} - \mu(v), R_{X,v'} - \mu(v')\}$.

since this is precisely equivalent to the validity of Eq. (17) as $t \rightarrow \infty$. For us, this condition follows by taking the trace on both sides of our assumption that $\sum_{k=1}^n q_k \Pi_k = \mathbb{1}_{\mathbb{R}^m}$. To state the second property of the heat flow that we will need, we momentarily assume sufficient regularity of the states under consideration, following [43]. Then, the *Fisher information* of a one-parameter family of states $\{\sigma^{(s)}\}$ is defined as

$$J(\{\sigma^{(s)}\}) := \partial_{s=0}^2 D(\sigma^0 \| \sigma^{(s)}).$$

It satisfies the following version of the data processing inequality [43, Theorem IV.4]: For any TPCP map \mathcal{E} ,

$$J(\{\mathcal{E}(\sigma^{(s)})\}) \leq J(\{\sigma^{(s)}\}). \quad (24)$$

For a covariant family of the form $\sigma^{(s,K)} := e^{isK} \sigma e^{-isK}$, the Fisher information can be computed as [43, Lemma IV.5]

$$J(\{\sigma^{(s,K)}\}) = \text{tr} \sigma [K, [K, \log \sigma]]. \quad (25)$$

We can now state the *quantum de Bruijn identity* [43, Theorem V.1], which computes the derivative of the entropy along the heat flow in terms of the Fisher information:

$$\partial_t H(\rho_X^{(t)}) = \frac{1}{4} J(\rho_X^{(t)}), \quad (26)$$

where the *total Fisher information* $J(\sigma_X)$ of a state σ_X on $L^2(X)$ is defined by

$$J(\sigma_X) := \sum_{j=1}^{m_X} J(\{\sigma^{(s, Q_{X, e_j})}\}) + J(\{\sigma^{(s, P_{X, e_j})}\}), \quad (27)$$

for an arbitrary orthonormal basis $\{e_j\}_{j=1}^{m_X}$ of X . While above we assumed regularity, the Fisher information $J(\sigma_X)$ can be defined for any state with finite second moments, and the de Bruijn identity (26) generalizes as well [28, Definition 7 & Proposition 1].⁶

Proof of Proposition III.4. We first prove the entropic Eq. (17) by considering $\rho^{(t)} := \rho_{\mathbb{R}^m}^{(t)}$ for $t \geq 0$. As $t \rightarrow \infty$, Eq. (17) holds up to arbitrarily small error, as explained below Eq. (23). To show its validity at $t = 0$, we would therefore like to argue that $\partial_t H(\rho^{(t)}) \geq \sum_{k=1}^n q_k \partial_t H(\rho_{X_k}^{(t)})$ for all $t \geq 0$. In view of the de Bruijn identity in Eq. (26) and Eq. (21), it suffices to establish the following *super-additivity* property of the Fisher information for all states σ on $L^2(\mathbb{R}^m)$ with finite second moment:

$$\sum_{k=1}^n q_k J(\sigma_{X_k}) \leq J(\sigma). \quad (28)$$

⁶ The key idea is to first define an *integral* version of the Fisher information [28, Definition 6]. In the setting without side information, this is defined for a state σ_X on $L^2(X)$ and for $t > 0$ by $\Delta(\sigma_X)(t) := I(X : V)_{\sigma_{XV}(t)}$, where $\sigma_{XV}(t)$ denotes the classical-quantum state with V is a multivariate Gaussian random variable with covariance matrix $t(I_X \oplus I_X)$ and $\sigma_{X|V=v} = \mathcal{D}_{X,v} \sigma_X \mathcal{D}_{X,v}^\dagger$, with $\mathcal{D}_{X,v} = \mathcal{Q}_{X,x}^{(1)} \circ \mathcal{P}_{X,y}^{(1)}$ for $v = (x, y) \in X \oplus X$; one also sets $\Delta_X(\sigma_X)(0) := 0$. This is well-defined for any state σ_X and satisfies a finitary version of the de Bruijn identity [28, Theorem 1]. Moreover, if σ_X is a state with finite energy then $\Delta(\sigma_X)(t)$ is continuous, increasing, and concave as a function of $t \geq 0$. Hence, for such states one can define the Fisher information $J(\sigma_X)$ as the (right) derivative of $\Delta(\sigma_X)(t)$ at $t = 0$, that is, as $J(\sigma_X) = \lim_{t \downarrow 0} \frac{\Delta(\sigma_X)(t)}{t}$ [28, Definition 7]. Then the de Bruijn identity (26) follows directly from its finitary version [28, Proposition 1].

We first prove this under the regularity assumptions of [43], so that Eq. (25) applies. We will abbreviate $Q_j := Q_{\mathbb{R}^m, e_j}$ and $P_j := P_{\mathbb{R}^m, e_j}$, where $\{e_j\}_{j=1}^m$ is the standard basis of \mathbb{R}^m . For all $x \in X_k$, it holds that

$$\begin{aligned}
J(\{\sigma_{X_k}^{(s, Q_{X_k, x})}\}) &= J(\{\mathcal{P}_{X_k, x}^{(s)}(\mathcal{E}_{X_k}(\sigma))\}) \\
&= J(\{\mathcal{E}_{X_k}(\mathcal{P}_{\mathbb{R}^m, x}^{(s)}(\sigma))\}) \\
&\leq J(\{\mathcal{P}_{\mathbb{R}^m, x}^{(s)}(\sigma)\}) \\
&= \text{tr } \sigma [Q_{\mathbb{R}^m, x}, [Q_{\mathbb{R}^m, x}, \log \sigma]] \\
&= \sum_{j, j'=1}^m x_j x_{j'} \text{tr } \sigma [Q_j, [Q_{j'}, \log \sigma]] \\
&= \sum_{j, j'=1}^m (xx^T)_{j, j'} \text{tr } \sigma [Q_j, [Q_{j'}, \log \sigma]]
\end{aligned}$$

where the second step is by the compatibility of phase-space translations and generalized partial trace (22), the third step uses the data-processing inequality for the Fisher information Eq. (24), and the fourth step follows from Eq. (25). If we apply the same argument to $J(\{\sigma_{X_k}^{(s, P_{X_k, x})}\})$ and sum both inequalities over an orthonormal basis $\{x\}$ of X_k , we obtain

$$J(\sigma_{X_k}) \leq J(\sigma, \Pi_k), \quad (29)$$

where we used the shortcut $J(\sigma, A) := \sum_{j, j'=1}^m A_{j, j'} (\text{tr } \sigma [Q_j, [Q_{j'}, \log \sigma]] + \text{tr } \sigma [P_j, [P_{j'}, \log \sigma]])$ for any positive semidefinite $m \times m$ matrix A , which is linear in A . Thus, our assumption that $\sum_{k=1}^n q_k \Pi_k = \mathbb{1}_{\mathbb{R}^m}$ (with all $q_k \geq 0$) implies the desired inequality:

$$\sum_k q_k J(\sigma_{X_k}) \leq \sum_k q_k J(\sigma, \Pi_k) = J(\sigma, \sum_k q_k \Pi_k) = J(\sigma, \mathbb{1}_{\mathbb{R}^m}) = J(\sigma). \quad (30)$$

This establishes Eq. (28) and hence Eq. (17) for states that are sufficiently regular. While Eq. (25) need not apply in general, the Fisher information $J(\sigma)$ and the de Bruijn identity (26) have been generalized to arbitrary states with finite second moments [28], as discussed above. The quantity $J(\sigma, A)$ can be defined in the same manner so that Eqs. (29) and (30) hold verbatim, see [29, Definition 6, Propositions 6 & 9].⁷

The analytic form in Eq. (18) then follows from a slight extension of Theorem II.1, or more specifically the special case discussed in Corollary II.5. Namely, we need to incorporate on the entropic side the finite second moment assumption from Eq. (17). By inspection, the variational formulae from Fact II.2 applied to operators with finite second moment still hold for the respective suprema only taken over operators with finite second moment. Hence, following the proof of the BL duality in Theorem II.1, we can still go from the entropic to the analytic form when assuming that the operator in exponential form on the left hand side of the analytic form has finite second moment. \square

⁷ The idea is the same in Footnote 6. One first defines an integral quantity $\Delta(\sigma, A)$ just like for the Fisher information, except that the multivariate Gaussian random variable now has covariance matrix $A \oplus A$ [29, Definition 5]. If σ_X is a state with finite energy then $t \mapsto \Delta(\sigma, tA)$ is again continuous, increasing, and concave [29, Proposition 5], and hence one can define $J(\sigma, A) := \lim_{t \downarrow 0} \frac{\Delta(\sigma, tA)}{t}$ [29, Definition 6]. Then $J(\sigma, A)$ satisfies a Stam inequality [29, Prop. 9] that implies (29), and it is still a linear function of A (for nonnegative linear combinations) [29, Prop. 6], which is what we used in (30).

While the preceding discussion restricted to the geometric case, we can also consider the general case of surjective linear map $L_k: \mathbb{R}^m \rightarrow \mathbb{R}^{m_k}$, as in Section I. For this, write L_k as the composition of an invertible map $M_k \in \text{GL}(m)$ and the projection onto the first m_k coordinates. Define a unitary operator U_k on $L^2(\mathbb{R}^m)$ by $(U_k g)(x) := g(M_k^{-1}x)/\sqrt{|\det M_k|}$. Then, $\mathcal{E}_k(\rho) := \text{tr}_{m_k+1, \dots, m}(U_k \rho U_k^\dagger)$ defines a TPCP map that is the natural quantum version of the marginalization $g \mapsto g_k$ (same notation as in Eq. (2)). We leave it for future work to determine under which conditions such quantum Brascamp-Lieb inequalities hold in general.

Note added: In follow-up work, Eq. (17) from Proposition III.4 has been extended to the conditional case with side information [50, Theorem 7.3] for Gaussian states, based on [44]. Subsequently, the latter assumption was removed by De Palma and Trevisan [29], who further generalized Proposition III.4 and also fully resolved the aforementioned question.

C. Entropic uncertainty relations

In this section, we explain how the duality of Theorem II.1 and Corollary II.5 offers an elegant way to prove entropic uncertainty relations (cf. the related work [57]). In order to compare our uncertainty bounds with the previous literature, we work in the current subsection with the explicit logarithm function relative to base two.

Example III.5 (Maassen-Uffink). For $\rho_A \in \mathcal{S}(A)$ the *Maassen-Uffink entropic uncertainty relation* [54] for two arbitrary basis measurements,

$$M_{\mathbb{X}}(\cdot) = \sum_x \langle x | \cdot | x \rangle |x\rangle \langle x|_X \quad \text{and} \quad M_{\mathbb{Z}}(\cdot) = \sum_z \langle z | \cdot | z \rangle |z\rangle \langle z|_Z,$$

asserts in its strengthened form [11] that

$$H(X) + H(Z) \geq -\log c(X, Z) + H(A) \quad \text{with } c(X, Z) := \max_{x,z} |\langle x | z \rangle|^2. \quad (31)$$

The constant $c(X, Z)$ is tight in the sense that there exist quantum states that achieve equality for certain measurement maps. Eq. (10) of Corollary II.5 for $n = 2$, $q_1 = q_2 = 1$, $\mathcal{E}_1 = M_{\mathbb{X}}$, and $\mathcal{E}_2 = M_{\mathbb{Z}}$ then immediately gives the equivalent analytic form

$$\text{tr exp} \left(M_{\mathbb{X}}^\dagger(\log \omega_1) + M_{\mathbb{Z}}^\dagger(\log \omega_2) \right) \leq c(X, Z) \quad \forall \omega_1, \omega_2 \in \mathcal{S}(A). \quad (32)$$

In other words, in order to prove Eq. (31) it suffices to show Eq. (32). Now, since the logarithm is operator concave and $M_{\mathbb{X}}^\dagger$ is a unital map, the operator Jensen inequality [36] implies

$$M_{\mathbb{X}}^\dagger(\log X_1) \leq \log M_{\mathbb{X}}^\dagger(X_1).$$

Together with the monotonicity of $X \mapsto \text{tr exp}(X)$ [18, Theorem 2.10] and the Golden-Thompson inequality⁸ [34, 61], this establishes the analytic form of Eq. (32)

$$\text{tr exp} \left(M_{\mathbb{X}}^\dagger(\log \omega_1) + M_{\mathbb{Z}}^\dagger(\log \omega_2) \right) \leq \text{tr exp} \left(\log M_{\mathbb{X}}^\dagger(\omega_1) + \log M_{\mathbb{Z}}^\dagger(\omega_2) \right)$$

⁸ The Golden-Thompson inequality ensures that for all Hermitian matrices H_1 and H_2 we have $\text{tr exp}(H_1 + H_2) \leq \text{tr exp}(H_1) \text{tr exp}(H_2)$.

$$\begin{aligned} &\leq \text{tr } M_{\mathbb{X}}^{\dagger}(\omega_1) M_{\mathbb{Z}}^{\dagger}(\omega_2) \\ &\leq c(X, Z). \end{aligned}$$

Thus, the entropic Maassen-Uffink relation Eq. (31) follows from our Corollary II.5.

We note that the approach of proving entropic uncertainty relations via the Golden-Thompson inequality was pioneered by Frank & Lieb [31] and is conceptually different from the original proofs that are either based on complex interpolation theory for Schatten p -norms [54] or the monotonicity of quantum relative entropy [26]. We refer to [25] for a review on entropic uncertainty relations. As a possible extension one could choose non-trivial pre-factors $q_k \neq 1$ and study the optimal uncertainty bounds in that setting as well (as done in [57] without the $H(A)$ term). Another natural extension is to general quantum channels instead of measurements (as detailed in [12, 32]). The constant $c(X, Z)$ from Eq. (31) is *multiplicative* for tensor product measurements. However, we might ask more generally if for given measurements the optimal lower bound in Eq. (31) becomes multiplicative for tensor product measurements. This amounts to an instance of the tensorization question from Eq. (13) and we refer to [32, 57] for a discussion.

An advantage of our BL analysis is that it suggests tight generalizations to multiple measurements by means of the multivariate extension of the Golden-Thompson inequality [60]. A basic example is as follows.

Example III.6 (Six-state [27]). For $\rho_A \in \mathcal{S}(A)$ with $\dim(A) = 2$ and measurement maps $M_{\mathbb{X}}, M_{\mathbb{Y}}, M_{\mathbb{Z}}$ of the Pauli matrices $\sigma_X, \sigma_Y, \sigma_Z$ we have

$$H(X) + H(Y) + H(Z) \geq 2 + H(A). \quad (33)$$

Moreover, this relation is tight in the sense that there exist quantum states that achieve equality. Note that applying the Maassen-Uffink relation Eq. (31) for any two of the three Pauli measurements only yields the weaker bound

$$H(X) + H(Y) + H(Z) \geq \frac{3}{2} + \frac{3}{2}H(A).$$

The equivalent analytic form of Eq. (33) is given by Corollary II.5 as

$$\text{tr} \exp \left(M_{\mathbb{X}}^{\dagger}(\log \omega_1) + M_{\mathbb{Y}}^{\dagger}(\log \omega_2) + M_{\mathbb{Z}}^{\dagger}(\log \omega_3) \right) \leq \frac{1}{4} \quad \forall \omega_1, \omega_2, \omega_3 \in \mathcal{S}(A).$$

The same steps as in the proof of the Maassen-Uffink relation, together with Lieb's triple matrix inequality [46] then yield the upper bound⁹

$$\begin{aligned} &\int_0^\infty \text{tr } M_{\mathbb{X}}(\omega_1) \frac{1}{M_{\mathbb{Z}}(\omega_3)^{-1} + t} M_{\mathbb{Y}}(\omega_2) \frac{1}{M_{\mathbb{Z}}(\omega_3)^{-1} + t} dt \\ &= \sum_{x,y} \langle x | \omega_1 | x \rangle \langle y | \omega_2 | y \rangle \int_0^\infty |\langle x | \frac{1}{M_{\mathbb{Z}}(\omega_3)^{-1} + t} | y \rangle|^2 dt \\ &\leq \max_{x,y} \int_0^\infty |\langle x | \frac{1}{M_{\mathbb{Z}}(\omega_3)^{-1} + t} | y \rangle|^2 dt. \end{aligned}$$

⁹ Lieb's triple matrix inequality corresponds to the three matrix Golden-Thompson inequality from [60].

In the penultimate step we used that

$$\begin{aligned}
M_{\mathbb{X}}(\omega) &= \sum_{x \in \{x_0, x_1\}} \langle x | \omega | x \rangle |x\rangle\langle x| \quad \text{where } \left\{ |x_0\rangle = \frac{1}{\sqrt{2}}(1, 1)^T, |x_1\rangle = \frac{1}{\sqrt{2}}(1, -1)^T \right\} \\
M_{\mathbb{Y}}(\omega) &= \sum_{y \in \{y_0, y_1\}} \langle y | \omega | y \rangle |y\rangle\langle y| \quad \text{where } \left\{ |y_0\rangle = \frac{1}{\sqrt{2}}(1, i)^T, |y_1\rangle = \frac{1}{\sqrt{2}}(1, -i)^T \right\} \\
M_{\mathbb{Z}}(\omega) &= \sum_{z \in \{z_0, z_1\}} \langle z | \omega | z \rangle |z\rangle\langle z| \quad \text{where } \left\{ |z_0\rangle = (1, 0)^T, |z_1\rangle = (0, 1)^T \right\}.
\end{aligned} \tag{34}$$

As $(M_{\mathbb{Z}}(\omega_3)^{-1} + t)^{-1} = \sum_z \frac{1}{\langle z | \omega_3 | z \rangle^{-1} + t} |z\rangle\langle z|$, we get

$$\left| \langle x | \frac{1}{M_{\mathbb{Z}}(\omega_3)^{-1} + t} | y \rangle \right|^2 = \left| \sum_z \frac{1}{\langle z | \omega_3 | z \rangle^{-1} + t} \langle x | z \rangle \langle z | y \rangle \right|^2.$$

Together with $\langle x | z_0 \rangle \langle z_0 | y \rangle = \frac{1}{2}$ and $\langle x | z_1 \rangle \langle z_1 | y \rangle = \pm \frac{1}{2}$ for all $x \in \{x_0, x_1\}$, $y \in \{y_0, y_1\}$ we find the upper bound

$$\frac{1}{4} \int_0^\infty \left((\langle z_0 | \omega_3 | z_0 \rangle^{-1} + t)^{-2} + (\langle z_1 | \omega_3 | z_1 \rangle^{-1} + t)^{-2} \right) dt = \frac{1}{4} (\langle z_0 | \omega_3 | z_0 \rangle + \langle z_1 | \omega_3 | z_1 \rangle) = \frac{1}{4}.$$

This then concludes the proof of the six-state entropic uncertainty relation Eq. (33).

D. Minimum output entropy

The Brascamp-Lieb duality from Theorem II.1 and Corollary II.5 is also applied usefully to general quantum channels. Recall that the *minimum output entropy* of a map $\mathcal{E} \in \text{TPCP}(A, B)$ is defined by

$$H_{\min}(\mathcal{E}) := \min_{\rho \in \mathcal{S}(A)} H(\mathcal{E}(\rho)). \tag{35}$$

The computation of minimum output entropy is in general NP-complete [9]. Nevertheless, it is a fundamental information measure [58] that has been used, e.g., to prove super-additivity of the Holevo information [38]. Corollary II.5 for $n = 2$, $q_1 = q_2 = 1$, $\mathcal{E}_1 = \mathcal{I}$, and $\mathcal{E}_2 = \mathcal{E}$ gives the following result.

Corollary III.7 (Minimum output entropy). *For $\mathcal{E} \in \text{TPCP}(A, B)$ and $C \in \mathbb{R}$, the following two statements are equivalent:*

$$C \leq H(\mathcal{E}(\rho)) \quad \forall \rho \in \mathcal{S}(A), \tag{36}$$

$$\text{tr} \exp(\log \omega_1 + \mathcal{E}^\dagger(\log \omega_2)) \leq \exp(-C) \quad \forall \omega_1 \in \mathcal{S}(A), \omega_2 \in \mathcal{S}(B). \tag{37}$$

Moreover, we have

$$H_{\min}(\mathcal{E}) = - \max_{\omega \in \mathcal{S}(B)} \lambda_{\max}(\mathcal{E}^\dagger(\log \omega)). \tag{38}$$

It is unclear if the form Eq. (38) could give new insights on the tensorization question of when the minimal output entropy of tensor product channels becomes additive. That is, for which $\mathcal{E}, \mathcal{F} \in \text{TPCP}(A, B)$ do we have

$$H_{\min}(\mathcal{E} \otimes \mathcal{F}) \stackrel{?}{=} H_{\min}(\mathcal{E}) + H_{\min}(\mathcal{F}). \quad (39)$$

We note that probabilistic counterexamples are known [38], which shows that the tensorization question Eq. (13) is in general answered in the negative.

Proof of Corollary III.7. We give two proofs of Eq. (38), one based on the variational characterization of the relative entropy from Eq. (6), and the other based on the dual formulation from Eq. (37). Using the former approach, we see that

$$\begin{aligned} H_{\min}(\mathcal{E}) &= \min_{\rho \in \mathcal{S}(A)} H(\mathcal{E}(\rho)) = \min_{\rho \in \mathcal{S}(A)} -D(\mathcal{E}(\rho) \| \mathbb{1}) \\ &= \min_{\rho \in \mathcal{S}(A)} - \left(\max_{\omega \in \mathcal{P}_{\succ}(B)} \text{tr } \mathcal{E}(\rho) \log \omega - \log \text{tr } \omega \right) \\ &= \min_{\rho \in \mathcal{S}(A), \omega \in \mathcal{S}(B)} - \text{tr } \rho \mathcal{E}^\dagger(\log \omega) \\ &= - \max_{\rho \in \mathcal{S}(A), \omega \in \mathcal{S}(B)} \text{tr } \rho \mathcal{E}^\dagger(\log \omega) \\ &= - \max_{\omega \in \mathcal{S}(B)} \lambda_{\max}(\mathcal{E}^\dagger(\log \omega)), \end{aligned}$$

where the final step follows from the variational formula of the largest eigenvalue.

Alternatively we can verify Eq. (38) in the analytic picture. To see this, we note that using the equivalence between Eq. (36) and Eq. (37) as well as the monotonicity of the logarithm,

$$H_{\min}(\mathcal{E}) = - \max_{\omega_1 \in \mathcal{S}(A), \omega_2 \in \mathcal{S}(B)} \log \text{tr} \exp(\log \omega_1 + \mathcal{E}^\dagger(\log \omega_2)). \quad (40)$$

Next, note that, for any Hermitian H , the Golden-Thompson inequality gives

$$\max_{\omega_1 \in \mathcal{S}(A)} \text{tr} \exp(\omega_1 + H) \leq \max_{\omega_1 \in \mathcal{S}(A)} \text{tr } \omega_1 \exp(H) = \lambda_{\max}(\exp(H)) = \exp(\lambda_{\max}(H)),$$

where the second step uses again the variational formula for the largest eigenvalue. This inequality is in fact an equality, since the upper-bound is attained if we choose ω_1 to be a projector onto an eigenvector of H with largest eigenvalue (any such ω_1 commutes with H). If we use this to evaluate Eq. (40), then we obtain the desired result. \square

Example III.8 (Qubit depolarizing channel). The minimal output entropy of the qubit depolarizing channel

$$\mathcal{E}_p: X \mapsto (1-p)X + p \frac{\mathbb{1}_{\mathbb{C}^2}}{2} \text{tr } X \quad \text{for } p \in [0, 1] \quad (41)$$

is given by $H_{\min}(\mathcal{E}_p) = h(p/2)$ with $h(x) := -x \log x - (1-x) \log(1-x)$ is the *binary entropy function*. In the entropic picture, this follows as the concavity of the entropy ensures that the optimizer in Eq. (35) can always be taken to be a pure state; the unitary covariance property

of the depolarizing channel then implies that we only need to evaluate the output entropy for a single arbitrary pure state. In the analytic picture, we can use Eq. (38) to see that

$$H_{\min}(\mathcal{E}_p) = - \max_{\omega \in \mathcal{S}(B)} \lambda_{\max}(\mathcal{E}^\dagger(\log \omega)) = - \max_{t \in [0,1]} \left\{ \left(1 - \frac{p}{2}\right) \log t + \frac{p}{2} \log(1-t) \right\} = h\left(\frac{p}{2}\right),$$

where the second step follows from unitary covariance and the final step uses that $t^* = 1 - p/2$ is the optimizer.

E. Data-processing inequality

The examples given so far employed Corollary II.5, but in this section we give an example that demonstrates Theorem II.1 in its full strength (with $\sigma, \sigma_k \neq \mathbb{1}$). The *data-processing inequality* (DPI) for the quantum relative entropy is a cornerstone in quantum information theory [51, 55, 62]. It states that, for $\rho \in \mathcal{S}(A)$ and $\sigma \in \mathcal{P}_\succ(A)$, the quantum relative entropy cannot increase when applying a channel $\mathcal{E} \in \text{TPP}(A, B)$ to both arguments, i.e.,

$$D(\mathcal{E}(\rho) \parallel \mathcal{E}(\sigma)) \leq D(\rho \parallel \sigma).$$

The DPI is mathematically equivalent to many other fundamental results, including the strong sub-additivity of quantum entropy [48, 49]. Our Brascamp-Lieb duality framework fits the DPI. That is, Theorem II.1 applied for $n = 1$, $q_1 = 1$, $\sigma_1 = \mathcal{E}(\sigma)$, and $C = 0$ implies the following duality.

Corollary III.9 (DPI duality). *For $\sigma \in \mathcal{P}_\succ(A)$ and $\mathcal{E} \in \text{TPP}(A, B)$ the following inequalities hold and are equivalent:*

$$\begin{aligned} D(\mathcal{E}(\rho) \parallel \mathcal{E}(\sigma)) &\leq D(\rho \parallel \sigma) \quad \forall \rho \in \mathcal{S}(A), \\ \text{tr exp}(\log \sigma + \mathcal{E}^\dagger(\log \omega)) &\leq \text{tr exp}(\log \omega + \log \mathcal{E}(\sigma)) \quad \forall \omega \in \mathcal{P}_\succ(B). \end{aligned} \quad (42)$$

As a simple example for $\text{tr } \sigma \leq \text{tr } \rho = 1$, one can immediately see that $D(\rho \parallel \sigma) \geq 0$ by considering the trace map $\mathcal{E}(\cdot) = \text{tr}(\cdot)$. Namely, data processing for the trace map takes the trivial analytic form $\text{tr } \log \omega \leq 0$ for quantum states $\omega \in \mathcal{S}(A)$.

Given that the DPI is quite powerful, we suspect that Eq. (42) may be of interest too. We note that Eq. (42) does not immediately follow from existing results and thus seems novel. For example, employing the operator concavity of the logarithm, the operator Jensen inequality, and the Golden-Thompson inequality we get

$$\text{tr exp}(\log \sigma + \mathcal{E}^\dagger(\log \omega)) \leq \text{tr exp}(\log \sigma + \log \mathcal{E}^\dagger(\omega)) \leq \text{tr } \mathcal{E}^\dagger(\omega) \sigma = \text{tr } \omega \mathcal{E}(\sigma). \quad (43)$$

This immediately implies Hansen's multivariate Golden-Thompson inequality [35, Inequality (1)], but is in general still weaker than Eq. (42) as the Golden-Thompson inequality applied to the right-hand side of Eq. (42) likewise gives

$$\text{tr exp}(\log \omega + \log \mathcal{E}(\sigma)) \leq \text{tr } \omega \mathcal{E}(\sigma). \quad (44)$$

Only when $\sigma = \mathbb{1}$ and \mathcal{E} is unital does Eq. (42) simplify to $\text{tr exp}(\mathcal{E}^\dagger(\log \omega)) \leq \text{tr } \omega$, reducing to Eq. (43).¹⁰

¹⁰ Alternatively, this also follows directly via Jensen's trace inequality [37].

F. Strong data-processing inequalities

It is a natural to study potential strengthenings of the DPI inequality and a priori it is possible to seek for additive or multiplicative improvements. Additive strengthenings of the DPI have recently generated interest in quantum information theory [30, 42, 59, 60]. Here, we consider multiplicative improvements of the DPI, which have been called *strong data-processing inequalities* in the literature. To this end, define the *contraction coefficient* of $\mathcal{E} \in \text{TPCP}(A, B)$ at $\sigma \in \mathcal{S}(A)$ as

$$\eta(\sigma, \mathcal{E}) := \sup_{\mathcal{S}(A) \ni \rho \neq \sigma} \frac{D(\mathcal{E}(\rho) \| \mathcal{E}(\sigma))}{D(\rho \| \sigma)}. \quad (45)$$

The data-processing inequality then ensures that $\eta(\sigma, \mathcal{E}) \leq 1$, and we say that \mathcal{E} satisfies a strong data-processing inequality at σ if $\eta(\sigma, \mathcal{E}) < 1$. Theorem II.1 for $n = 1$, $C = 0$, $\sigma_1 = \mathcal{E}(\sigma)$, and $q_1 = \eta(\sigma, \mathcal{E})^{-1}$ implies the following equivalence.

Corollary III.10 (Strong DPI duality). *For $\mathcal{E} \in \text{TPP}(A, B)$, $\sigma \in \mathcal{P}_{\succ}(A)$, and $\eta > 0$, the following two statements are equivalent:*

$$D(\mathcal{E}(\rho) \| \mathcal{E}(\sigma)) \leq \eta D(\rho \| \sigma) \quad \forall \rho \in \mathcal{S}(A), \quad (46)$$

$$\text{tr} \exp(\log \sigma + \mathcal{E}^\dagger(\log \omega)) \leq \left\| \exp\left(\log \omega + \frac{1}{\eta} \log \mathcal{E}(\sigma)\right) \right\|_\eta \quad \forall \omega \in \mathcal{S}(B). \quad (47)$$

Thus, to determine $\eta(\sigma, \mathcal{E})$, we aim to find the smallest constant $\eta \in [0, 1]$ such that Eq. (46) or, equivalently, Eq. (47) holds. For unital \mathcal{E} and maximally mixed $\sigma = \mathbb{1}/d$, $d := \dim(A)$, the duality in Corollary III.10 simplifies to

$$\log d - H(\mathcal{E}(\rho)) \leq \eta (\log d - H(\rho)) \quad \forall \rho \in \mathcal{S}(A), \quad (48)$$

$$\text{tr} \exp(\mathcal{E}^\dagger(\log \omega)) \leq d^{\frac{\eta-1}{\eta}} \|\omega\|_\eta \quad \forall \omega \in \mathcal{S}(B). \quad (49)$$

Often we are also interested in the global *contraction coefficient* of \mathcal{E} , obtained by optimizing $\eta(\sigma, \mathcal{E})$ over all $\sigma \in \mathcal{S}(A)$, i.e.,

$$\eta(\mathcal{E}) := \sup_{\sigma \in \mathcal{S}(A)} \eta(\sigma, \mathcal{E}). \quad (50)$$

Example III.11 (Qubit depolarizing channel). For the qubit depolarizing channel \mathcal{E}_p from Eq. (41), which is unital, we claim that

$$\eta\left(\frac{\mathbb{1}_{\mathbb{C}^2}}{2}, \mathcal{E}_p\right) = (1-p)^2. \quad (51)$$

To prove this in the entropic picture we start by recalling that $\eta(\mathcal{E}_p) = (1-p)^2$ [39], which already gives $\eta(\frac{\mathbb{1}_{\mathbb{C}^2}}{2}, \mathcal{E}_p) \leq (1-p)^2$. Thus, it suffices to find states $\rho \in \mathcal{S}(A)$ such that

$$(1-p)^2 \leq \frac{D(\mathcal{E}_p(\rho) \| \frac{\mathbb{1}_{\mathbb{C}^2}}{2})}{D(\rho \| \frac{\mathbb{1}_{\mathbb{C}^2}}{2})} = \frac{1 - H(\mathcal{E}_p(\rho))}{1 - H(\rho)} \quad (52)$$

up to arbitrarily small error. The states $\rho_\varepsilon = \text{diag}(\frac{1}{2} + \varepsilon, \frac{1}{2} - \varepsilon)$ satisfy this condition in the limit $\varepsilon \rightarrow 0$. Indeed,

$$\lim_{\varepsilon \rightarrow 0} \frac{1 - H(\mathcal{E}_p(\rho_\varepsilon))}{1 - H(\rho_\varepsilon)} = \lim_{\varepsilon \rightarrow 0} \frac{1 - h((1-p)(1/2 + \varepsilon) + p/2)}{1 - h(1/2 + \varepsilon)} = (1-p)^2, \quad (53)$$

as follows from the Taylor expansion of the binary entropy function $h(\cdot)$.

In the analytic form of Eq. (49), the statement of Eq. (51) is equivalent to the claim that $\eta = (1-p)^2$ is the smallest $\eta \in [0, 1]$ such that

$$\text{tr} \exp((1-p) \log \omega + \frac{p}{2} \mathbf{1}_{\mathbb{C}^2} \text{tr} \log \omega) \leq 2^{\frac{\eta-1}{\eta}} \|\omega\|_\eta \quad \text{for all } \omega \in \mathcal{S}(B). \quad (54)$$

Without loss of generality we can assume that $\omega = \text{diag}(t, 1-t)$ for $t \in [0, 1]$. Then, the statement above simplifies to showing that $\eta = (1-p)^2$ is the smallest $\eta \in [0, 1]$ such that

$$(t(1-t))^{\frac{p}{2}} (t^{1-p} + (1-t)^{1-p}) \leq 2^{\frac{\eta-1}{\eta}} (t^\eta + (1-t)^\eta)^{\frac{1}{\eta}} \quad \text{for all } t \in [0, 1]. \quad (55)$$

G. Super-additivity of relative entropy

Another type of strengthening of the DPI is as follows. The quantum relative entropy is *super-additive* for product states in the second argument. That is, for $\rho_{AB}, \sigma_{AB} \in \mathcal{S}(A \otimes B)$ we have

$$D(\rho_{AB} \| \sigma_A \otimes \sigma_B) \geq D(\rho_A \| \sigma_A) + D(\rho_B \| \sigma_B). \quad (56)$$

This directly follows from the non-negativity of the relative entropy, since $D(\rho_{AB} \| \sigma_A \otimes \sigma_B) - D(\rho_A \| \sigma_A) - D(\rho_B \| \sigma_B) = D(\rho_{AB} \| \rho_A \otimes \rho_B) \geq 0$. If the state in the second argument is not a product state we can apply the DPI twice and find

$$D(\rho_{AB} \| \sigma_{AB}) \geq t D(\rho_A \| \sigma_A) + (1-t) D(\rho_B \| \sigma_B) \quad \text{for all } t \in [0, 1]. \quad (57)$$

A natural question is thus to find parameters $\alpha(\sigma_{AB}), \beta(\sigma_{AB})$ with $\alpha(\sigma_A \otimes \sigma_B) = \beta(\sigma_A \otimes \sigma_B) = 1$ such that¹¹

$$D(\rho_{AB} \| \sigma_{AB}) \geq \alpha(\sigma_{AB}) D(\rho_A \| \sigma_A) + \beta(\sigma_{AB}) D(\rho_B \| \sigma_B). \quad (58)$$

Recently, it was shown [17] that Eq. (58) indeed holds for

$$\alpha(\sigma_{AB}) = \beta(\sigma_{AB}) = \left(1 + 2 \left\| \sigma_A^{-\frac{1}{2}} \otimes \sigma_B^{-\frac{1}{2}} \sigma_{AB} \sigma_A^{-\frac{1}{2}} \otimes \sigma_B^{-\frac{1}{2}} - \mathbf{1}_{AB} \right\|_\infty \right)^{-1}. \quad (59)$$

Applying Theorem II.1 for $n = 2$, $\sigma_1 = \sigma_A$, $\sigma_2 = \sigma_B$, $C = 0$, $\mathcal{E}_1 = \text{tr}_B$, $\mathcal{E}_2 = \text{tr}_A$, $q_1 = \alpha$, and $q_2 = \beta$ gives the following BL duality.

Corollary III.12 (Duality for super-additivity of relative entropy). *For $\sigma_{AB} \in \mathcal{P}_\succ(A \otimes B)$ with $\text{tr} \sigma_{AB} = 1$, $\alpha > 0$, and $\beta > 0$, the following two statements are equivalent:*

$$\alpha D(\rho_A \| \sigma_A) + \beta D(\rho_B \| \sigma_B) \leq D(\rho_{AB} \| \sigma_{AB}) \quad \forall \rho_{AB} \in \mathcal{S}(A \otimes B), \quad (60)$$

$$\text{tr} \exp(\log \sigma_{AB} + \log \omega_A + \log \omega_B) \leq \|\exp(\log \omega_A + \alpha \log \sigma_A)\|_{\frac{1}{\alpha}} \|\exp(\log \omega_B + \beta \log \sigma_B)\|_{\frac{1}{\beta}} \quad \forall \omega_A \in \mathcal{S}(A), \omega_B \in \mathcal{S}(B). \quad (61)$$

We leave it as an open question to find parameters $\alpha(\sigma_{AB})$ and $\beta(\sigma_{AB})$ different from Eq. (59), satisfying Eq. (60) or equivalently Eq. (61).

¹¹ We might also ask for $\alpha(\sigma_{AB}) + \beta(\sigma_{AB}) \geq 1$.

IV. CONCLUSION

Our fully quantum Brascamp-Lieb dualities raise a plethora of possible extensions to study. Taking inspiration from the commutative case [53], this could include, e.g., Gaussian optimality questions, hypercontractivity inequalities, transportation cost inequalities, strong converses in Shannon theory, entropy power inequalities [1], or algorithmic and complexity-theoretic questions [15, 16, 33]. For some of these applications it seems that an extension of Barthe’s reverse Brascamp-Lieb duality [7] to the non-commutative setting would be useful.

Acknowledgements. MB and DS thank the Stanford Institute for Theoretical Physics for their hospitality during the time this project was initiated. MW would like to thank Graeme Smith and JILA for their hospitality. MW gratefully acknowledges Misha Gromov for his hospitality at IHES and for suggesting the problem discussed in Section III B. We thank Eric Carlen for corresponding with us about Brascamp-Lieb inequalities for relative entropies. We thank Ernest Tan for informing us about an error in Section III C in a previous version of this manuscript. DS acknowledges support from the Swiss National Science Foundation via the NCCR QSIT as well as project No. 200020_165843. MB acknowledges funding by the European Research Council (ERC Grant Agreement No. 948139). MW acknowledges support by the NWO through Veni grant no. 680-47-459 and grant OCENW.KLEIN.267, by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy - EXC 2092 CASA - 390781972, by the BMBF through project Quantum Methods and Benchmarks for Resource Allocation (QuBRA), and by the European Research Council (ERC) through ERC Starting Grant 101040907-SYMOPTIC.

-
- [1] V. Anantharam, V. Jog, and C. Nair. Unifying the Brascamp-Lieb inequality and the entropy power inequality. *IEEE International Symposium on Information Theory (ISIT)*, pages 1847–1851, 2019. DOI: [10.1109/ISIT.2019.8849711](https://doi.org/10.1109/ISIT.2019.8849711). Extended version available at [arXiv:1901.06619](https://arxiv.org/abs/1901.06619).
 - [2] K. Ball. *Volumes of sections of cubes and related problems*, pages 251–260. Springer Berlin Heidelberg, Berlin, Heidelberg, 1989. DOI: [10.1007/BFb0090058](https://doi.org/10.1007/BFb0090058).
 - [3] K. Ball. Shadows of convex bodies. *Transactions of the American Mathematical Society*, 327(2):891–901, 1991. DOI: [10.1090/S0002-9947-1991-1035998-3](https://doi.org/10.1090/S0002-9947-1991-1035998-3).
 - [4] K. Ball. Volume ratios and a reverse isoperimetric inequality. *Journal of the London Mathematical Society*, s2-44(2):351–359, 1991. DOI: [10.1112/jlms/s2-44.2.351](https://doi.org/10.1112/jlms/s2-44.2.351).
 - [5] K. Ball. An elementary introduction to modern convex geometry. *Flavors of geometry*, 31:1–58, 1997. Available online: <http://library.msri.org/books/Book31/files/ball.pdf>.
 - [6] K. Ball. *Convex geometry and functional analysis*, volume 1, chapter 4, pages 161–194. Elsevier, 2001. Available online: <https://www.ceremade.dauphine.fr/~lehec/gdl/handbook.pdf>.
 - [7] F. Barthe. On a reverse form of the Brascamp-Lieb inequality. *Inventiones mathematicae*, 134(2):335–361, 1998. DOI: [10.1007/s002220050267](https://doi.org/10.1007/s002220050267).
 - [8] F. Barthe and D. Cordero-Erausquin. *Inverse Brascamp-Lieb Inequalities along the Heat Equation*, pages 65–71. Springer Berlin Heidelberg, Berlin, Heidelberg, 2004.
 - [9] S. Beigi and P. W. Shor. On the complexity of computing zero-error and Holevo capacity of quantum channels, 2007. [arXiv:0709.2090](https://arxiv.org/abs/0709.2090).
 - [10] J. Bennett, A. Carbery, M. Christ, and T. Tao. The Brascamp-Lieb inequalities: Finiteness, structure and extremals. *Geometric and Functional Analysis*, 17(5):1343–1415, 2008. DOI: [10.1007/s00039-007-0619-6](https://doi.org/10.1007/s00039-007-0619-6).

- [11] M. Berta, M. Christandl, R. Colbeck, J. M. Renes, and R. Renner. The uncertainty principle in the presence of quantum memory. *Nature Physics*, 6, 2010. DOI: [10.1038/nphys1734](https://doi.org/10.1038/nphys1734).
- [12] M. Berta, F. Furrer, and V. B. Scholz. The smooth entropy formalism for von Neumann algebras. *Journal of Mathematical Physics*, 57:015213, 2016. DOI: [10.1063/1.4936405](https://doi.org/10.1063/1.4936405).
- [13] J.-C. Bourin and F. Hiai. Norm and anti-norm inequalities for positive semi-definite matrices. *International Journal of Mathematics*, 22(08):1121–1138, 2011. DOI: [10.1142/S0129167X1100715X](https://doi.org/10.1142/S0129167X1100715X).
- [14] H. J. Brascamp and E. H. Lieb. Best constants in Young’s inequality, its converse, and its generalization to more than three functions. *Advances in Mathematics*, 20(2):151–173, 1976. DOI: [10.1016/0001-8708\(76\)90184-5](https://doi.org/10.1016/0001-8708(76)90184-5).
- [15] P. Bürgisser, C. Franks, A. Garg, R. Oliveira, M. Walter, and A. Wigderson. Efficient algorithms for tensor scaling, quantum marginals, and moment polytopes. In *2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 883–897. IEEE, 2018. DOI: [10.1109/FOCS.2018.00088](https://doi.org/10.1109/FOCS.2018.00088).
- [16] P. Bürgisser, C. Franks, A. Garg, R. Oliveira, M. Walter, and A. Wigderson. Towards a theory of non-commutative optimization: geodesic 1st and 2nd order methods for moment maps and polytopes. In *2019 IEEE 60th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 845–861. IEEE, 2019. DOI: [10.1109/FOCS.2019.00055](https://doi.org/10.1109/FOCS.2019.00055).
- [17] A. Capel. Superadditivity of quantum relative entropy for general states. *IEEE Transactions on Information Theory*, 64(7):4758–4765, 2018. DOI: [10.1109/TIT.2017.2772800](https://doi.org/10.1109/TIT.2017.2772800).
- [18] E. Carlen. *Trace Inequalities and Quantum Entropy: An Introductory Course*. Contemporary Mathematics, 2009. DOI: [10.1090/conm/529](https://doi.org/10.1090/conm/529).
- [19] E. A. Carlen and D. Cordero-Erausquin. Subadditivity of the entropy and its relation to Brascamp–Lieb type inequalities. *Geometric and Functional Analysis*, 19(2):373–405, 2009. DOI: [10.1007/s00039-009-0001-y](https://doi.org/10.1007/s00039-009-0001-y).
- [20] E. A. Carlen and E. H. Lieb. Brascamp-Lieb inequalities for non-commutative integration. *Documenta Mathematica*, 13:553–584, 2008. Available online: <https://www.math.uni-bielefeld.de/documenta/vol-13/16.pdf>.
- [21] E. A. Carlen, E. H. Lieb, and M. Loss. A sharp analog of Young’s inequality on SN and related entropy inequalities. *The Journal of Geometric Analysis*, 14(3):487–520, 2004. DOI: [10.1007/BF02922101](https://doi.org/10.1007/BF02922101).
- [22] E. A. Carlen and J. Maas. An analog of the 2-Wasserstein metric in non-commutative probability under which the Fermionic Fokker–Planck equation is gradient flow for the entropy. *Communications in Mathematical Physics*, 331(3):887–926, 2014. DOI: [10.1007/s00220-014-2124-8](https://doi.org/10.1007/s00220-014-2124-8).
- [23] E. A. Carlen and J. Maas. Gradient flow and entropy inequalities for quantum Markov semigroups with detailed balance. *Journal of Functional Analysis*, 273(5):1810–1869, 2017. DOI: [10.1016/j.jfa.2017.05.003](https://doi.org/10.1016/j.jfa.2017.05.003).
- [24] E. A. Carlen and J. Maas. Non-commutative calculus, optimal transport and functional inequalities in dissipative quantum systems. *Journal of Statistical Physics*, 178(2):319–378, 2020. DOI: [10.1007/s10955-019-02434-w](https://doi.org/10.1007/s10955-019-02434-w).
- [25] P. J. Coles, M. Berta, M. Tomamichel, and S. Wehner. Entropic uncertainty relations and their applications. *Rev. Mod. Phys.*, 89:015002, 2017. DOI: [10.1103/RevModPhys.89.015002](https://doi.org/10.1103/RevModPhys.89.015002).
- [26] P. J. Coles, R. Colbeck, L. Yu, and M. Zwolek. Uncertainty relations from simple entropic properties. *Physical Review Letters*, 108:210405, 2012. DOI: [10.1103/PhysRevLett.108.210405](https://doi.org/10.1103/PhysRevLett.108.210405).
- [27] P. J. Coles, L. Yu, V. Gheorghiu, and R. B. Griffiths. Information-theoretic treatment of tripartite systems and quantum channels. *Physical Review A*, 83:062338, 2011. DOI: [10.1103/PhysRevA.83.062338](https://doi.org/10.1103/PhysRevA.83.062338).
- [28] G. De Palma and D. Trevisan. The conditional entropy power inequality for bosonic quantum systems. *Communications in Mathematical Physics*, 360(2):639–662, 2018. DOI: [10.1007/s00220-017-3082-8](https://doi.org/10.1007/s00220-017-3082-8).
- [29] G. De Palma and D. Trevisan. The generalized strong subadditivity of the von Neumann entropy for bosonic quantum Gaussian systems, 2021. [arXiv:2105.05627](https://arxiv.org/abs/2105.05627).
- [30] O. Fawzi and R. Renner. Quantum conditional mutual information and approxi-

- mate Markov chains. *Communications in Mathematical Physics*, 340(2):575–611, 2015. DOI: [10.1007/s00220-015-2466-x](https://doi.org/10.1007/s00220-015-2466-x).
- [31] R. L. Frank and E. H. Lieb. Extended quantum conditional entropy and quantum uncertainty inequalities. *Communications in Mathematical Physics*, 323(2):487–495, 2013. DOI: [10.1007/s00220-013-1775-1](https://doi.org/10.1007/s00220-013-1775-1).
- [32] L. Gao, M. Junge, and N. LaRacuente. Uncertainty principle for quantum channels. *IEEE International Symposium on Information Theory (ISIT)*, pages 996–1000, 2018. DOI: [10.1109/ISIT.2018.8437730](https://doi.org/10.1109/ISIT.2018.8437730).
- [33] A. Garg, L. Gurvits, R. Oliveira, and A. Wigderson. Algorithmic and optimization aspects of Brascamp-Lieb inequalities, via operator scaling. *Geometric and Functional Analysis*, 28(1):100–145, 2018. DOI: [10.1007/s00039-018-0434-2](https://doi.org/10.1007/s00039-018-0434-2).
- [34] S. Golden. Lower bounds for the Helmholtz function. *Physical Review*, 137:B1127–B1128, 1965. DOI: [10.1103/PhysRev.137.B1127](https://doi.org/10.1103/PhysRev.137.B1127).
- [35] F. Hansen. Multivariate extensions of the Golden-Thompson inequality. *Annals of Functional Analysis*, 6(4):301–310, 2015. DOI: [10.15352/afa/06-4-301](https://doi.org/10.15352/afa/06-4-301).
- [36] F. Hansen and G. K. Pedersen. Jensen’s operator inequality. *Bulletin of the London Mathematical Society*, 35(4):553–564, 2003. DOI: [10.1112/S0024609303002200](https://doi.org/10.1112/S0024609303002200).
- [37] F. Hansen and G. K. Pedersen. Jensen’s trace inequality in several variables. *International Journal of Mathematics*, 14(06):667–681, 2003. DOI: [10.1142/S0129167X03001983](https://doi.org/10.1142/S0129167X03001983).
- [38] M. B. Hastings. Superadditivity of communication capacity using entangled inputs. *Nature Physics*, 5(4):255–257, 2009. DOI: [10.1038/nphys1224](https://doi.org/10.1038/nphys1224).
- [39] F. Hiai and M. B. Ruskai. Contraction coefficients for noisy quantum channels. *Journal of Mathematical Physics*, 57(1):015211, 2016. DOI: [10.1063/1.4936215](https://doi.org/10.1063/1.4936215).
- [40] A. S. Holevo. *Quantum Systems, Channels, Information*. De Gruyter Studies in Mathematical Physics 16, 2012. DOI: [10.1515/9783110273403](https://doi.org/10.1515/9783110273403).
- [41] D. Huang. Generalizing lieb’s concavity theorem via operator interpolation. *Advances in Mathematics*, 369:107208, 2020. DOI: [10.1016/j.aim.2020.107208](https://doi.org/10.1016/j.aim.2020.107208).
- [42] M. Junge, R. Renner, D. Sutter, M. M. Wilde, and A. Winter. Universal recovery maps and approximate sufficiency of quantum relative entropy. *Annales Henri Poincaré*, 19(10):2955–2978, 2018. DOI: [10.1007/s00023-018-0716-0](https://doi.org/10.1007/s00023-018-0716-0).
- [43] R. König and G. Smith. The entropy power inequality for quantum systems. *IEEE Transactions on Information Theory*, 60(3):1536–1548, 2014. DOI: [10.1109/TIT.2014.2298436](https://doi.org/10.1109/TIT.2014.2298436).
- [44] R. König and G. Smith. Corrections to “The entropy power inequality for quantum systems”. *IEEE Transactions on Information Theory*, 62(7):4358–4359, 2016. DOI: [10.1109/TIT.2016.2563438](https://doi.org/10.1109/TIT.2016.2563438).
- [45] A. Kuznetsova. Conditional entropy for infinite-dimensional quantum systems. *Theory of Probability & Its Applications*, 55(4):709–717, 2011. DOI: [10.1137/S0040585X97985121](https://doi.org/10.1137/S0040585X97985121).
- [46] E. H. Lieb. Convex trace functions and the Wigner-Yanase-Dyson conjecture. *Advances in Mathematics*, 11(3):267 – 288, 1973. DOI: [10.1016/0001-8708\(73\)90011-X](https://doi.org/10.1016/0001-8708(73)90011-X).
- [47] E. H. Lieb. Gaussian kernels have only Gaussian maximizers. *Inventiones Mathematicae*, 102(1):179–208, 1990. DOI: [10.1007/BF01233426](https://doi.org/10.1007/BF01233426).
- [48] E. H. Lieb and M. B. Ruskai. A fundamental property of quantum-mechanical entropy. *Physical Review Letters*, 30:434–436, 1973. DOI: [10.1103/PhysRevLett.30.434](https://doi.org/10.1103/PhysRevLett.30.434).
- [49] E. H. Lieb and M. B. Ruskai. Proof of the strong subadditivity of quantum-mechanical entropy. *Journal of Mathematical Physics*, 14(12):1938–1941, 1973. DOI: [10.1063/1.1666274](https://doi.org/10.1063/1.1666274).
- [50] L. Ligthart. Linear quantum entropy inequalities beyond strong subadditivity and their applications. *MSc thesis, University of Amsterdam & Vrije Universiteit Amsterdam*, 2020.
- [51] G. Lindblad. Completely positive maps and entropy inequalities. *Communications in Mathematical Physics*, 40(2):147–151, 1975. DOI: [10.1007/BF01609396](https://doi.org/10.1007/BF01609396).
- [52] J. Liu, T. A. Courtade, P. Cuff, and S. Verdú. Brascamp-Lieb inequality and its reverse: An information theoretic view. *IEEE International Symposium on Information Theory (ISIT)*, pages 1048–1052, 2016. DOI: [10.1109/ISIT.2016.7541459](https://doi.org/10.1109/ISIT.2016.7541459).
- [53] J. Liu, T. A. Courtade, P. Cuff, and S. Verdu. Information-theoretic perspectives on Brascamp-Lieb

- inequality and its reverse, 2017. [arXiv:1702.06260](#).
- [54] H. Maassen and J. B. M. Uffink. Generalized entropic uncertainty relations. *Physical Review Letters*, 60:1103–1106, 1988. DOI: [10.1103/PhysRevLett.60.1103](#).
 - [55] A. Müller-Hermes and D. Reeb. Monotonicity of the quantum relative entropy under positive maps. *Annals of Henri Poincaré*, 2017. DOI: [10.1007/s00023-017-0550-9](#).
 - [56] D. Petz. A variational expression for the relative entropy. *Communications in Mathematical Physics*, 114(2):345–349, 1988. DOI: [10.1007/BF01225040](#).
 - [57] R. Schwonnek. Additivity of entropic uncertainty relations. *Quantum*, 2:59, 2018. DOI: [10.22331/q-2018-03-30-59](#).
 - [58] P. W. Shor. Equivalence of additivity questions in quantum information theory. *Communications in Mathematical Physics*, 246(3):473–473, 2004. DOI: [10.1007/s00220-003-0981-7](#).
 - [59] D. Sutter. *Approximate Quantum Markov Chains*. Springer International Publishing, 2018. DOI: [10.1007/978-3-319-78732-9_5](#).
 - [60] D. Sutter, M. Berta, and M. Tomamichel. Multivariate trace inequalities. *Communications in Mathematical Physics*, 352(1):37–58, 2017. DOI: [10.1007/s00220-016-2778-5](#).
 - [61] C. J. Thompson. Inequality with applications in statistical mechanics. *Journal of Mathematical Physics*, 6(11):1812–1813, 1965. DOI: [10.1063/1.1704727](#).
 - [62] A. Uhlmann. Relative entropy and the Wigner-Yanase-Dyson-Lieb concavity in an interpolation theory. *Communications in Mathematical Physics*, 54(1):21–32, 1977. DOI: [10.1007/BF01609834](#).