

Stationary states of weak coupling limit type Markov generators and quantum transport models

Álvaro Hernández-Cervantes¹ and Roberto Quezada²

January 28, 2022

Universidad Autónoma Metropolitana, Iztapalapa Campus, Av. San Rafael Atlixco 186, Col. Vicentina, 09340 Iztapalapa D.F., México.
E-mail: ahc_89@hotmail.com¹, roqb@xanum.uam.mx²

Abstract

We prove that every stationary state in the annihilator of all Kraus operators of a weak coupling limit type Markov generator consists of two pieces, one of them supported on the interaction-free subspace and the second one on its orthogonal complement. In particular, we apply the previous result to describe in detail the structure of a slightly modified quantum transport model due to Arefeva, Kozyrev and Volovich (modified AKV's model) studied first in Ref.[16], in terms of generalized annihilation and creation operators.

1 Introduction

The main aim of this work is two-fold. We describe the structure of stationary states in the annihilator of all Kraus (or noise) operators of the class of weak coupling limit type Markov generators (WCLT generators) introduced in Ref.[1] and, in particular, we describe in detail the structure of the stationary states of the modified transport AKV's model, studied first in Ref.[16], in terms of generalized annihilation and creation operators.

A natural space to search for stationary states of WCLT generators is the annihilator of all Kraus or noise operators defined as

$$Ann(D) := \{\rho : \text{tr}(\rho D_\omega) = 0, \text{ for every Bohr frequency } \omega\}$$

that contains (sometimes properly) the commutant $\{H\}'$ of the reference Hamiltonian. We show that stationary states in $\text{Ann}(D)$ consist of a piece supported on the intersection of the kernels of all Kraus operators and its adjoints, denoted by W_D and called interaction-free subspace, and another part supported on W_D^\perp . While the part supported on W_D belongs to the fixed points sub-algebra of the semigroup, the second is much more interesting and include detailed balance as well as non-detailed balance but in the class of local detailed balance stationary states.

For the modified AKV's model, stationary states supported on W_D^\perp , exhibit a very interesting structure described explicitly by using an operator Z , called interference operator in Ref.[16], and its adjoint Z^* . Operator Z maps the support subspace $P_2(\mathbf{h})$ of the degenerate second spectral projection of the reference Hamiltonian H , into the support subspace $P_3(\mathbf{h})$ of the third spectral projection. Operators Z and Z^* perform transitions between $P_2(\mathbf{h})$ and $P_3(\mathbf{h})$, similar to birth and death transitions in classical stochastic processes or creation and annihilation operators in the quantum setting. We show in Theorem 34 that any stationary state of the modified AKV's model, is a mixture of a state σ supported on a subspace of $P_2(\mathbf{h})$ and its conjugation by Z , $Z\sigma Z^*$, that is supported on a subspace of $P_3(\mathbf{h})$. It turns out that the slightly modification of AKV's transport model performed in Ref.[16] yields another quantum transport model, indeed: the total probability mass of any initial state of the modified transport model can be redistributed as $t \rightarrow \infty$, choosing appropriate values of the parameters of the model (the Γ 's), so that most of the total probability of the final state is concentrated on the portion supported on P_3 , even though the support of the final state is not contained in $P_3(\mathbf{h})$ (there is a non-zero trace on $P_2(\mathbf{h})$), see Remark 36. Moreover, the evolution increases the energy of the small system when it is initially in a state supported on a distinguished subspace of $P_2(\mathbf{h})$, so that when $t \rightarrow \infty$ there is an energy gain proportional to $(N - 1) = \dim P_2(\mathbf{h})$, see Remark 37.

The paper is organized as follows. After some preliminaries, in Section 3 we describe the structure of stationary state WCLT Markov generators as a convex combination of a state supported on the interaction-free subspace W_D and another state supported on the orthogonal complement W_D^\perp . In Section 4 we apply the results of the previous section to study the stationary state of the modified AKV's model by a method considerable shorter than the one used in Ref.[16] In contradistinction with the previous reference, where the assumption that any stationary state belongs to $\{H\}'$ is used, we start with the computation of subharmonic projections and prove that any stationary state belongs to $\{H\}'$. Moreover we study the approach to equilibrium property for this semigroup and characterize the attraction domain of any stationary state.

2 Preliminaries

Following Ref.[1], we call Markov generator (or simply generator) any formal expression with the GKSL structure from which one can construct a Quantum Markov Semigroup. In particular we shall consider Markov generators of weak coupling limit type \mathcal{L} , associated with a positive self-adjoint operator (reference Hamiltonian) with discrete spectral decomposition

$$H = \sum_{\varepsilon_m \in \text{Sp}(H)} \varepsilon_m P_{\varepsilon_m} \quad (1)$$

Where ε_m are the eigenvalues and P_{ε_m} is the spectral projection on the eigenspace associated with ε_m .

A weak coupling limit type Markov generator has the structure

$$\mathcal{L} = \sum_{\omega \in B_+} \mathcal{L}_\omega \quad (2)$$

The subset of ordered pairs of eigenvalues (Arveson spectrum)

$$B_+ := \{(\varepsilon_n, \varepsilon_m) : \varepsilon_n - \varepsilon_m > 0\} \quad (3)$$

is the set of Bohr frequencies and

$$B_{+, \omega} := \{(\varepsilon_n, \varepsilon_m) \in B_+ : \varepsilon_n - \varepsilon_m = \omega\} \quad (4)$$

is the set of pairs of eigenvalues associated with the frequency ω .

For every Bohr frequency ω and $x \in \mathcal{B}(\mathfrak{h})$ the corresponding generator \mathcal{L}_ω has the Gorini-Kossakowski-Sudarshan and Lindblad (GKSL) canonical form

$$\begin{aligned} \mathcal{L}_\omega(x) &:= i[\Delta_\omega, x] \\ &- \Gamma_{-, \omega} \left(\frac{1}{2} \{D_\omega^* D_\omega, x\} - D_\omega^* x D_\omega \right) - \Gamma_{+, \omega} \left(\frac{1}{2} \{D_\omega D_\omega^*, x\} - D_\omega x D_\omega^* \right) \end{aligned} \quad (5)$$

with partial interaction (or Kraus) operators

$$D_\omega := \sum_{(\varepsilon_n, \varepsilon_m) \in B_{+, \omega}} P_{\varepsilon_m} D P_{\varepsilon_n}, \quad (6)$$

with $D \in \mathcal{B}(\mathfrak{h})$ an interaction operator. The effective Hamiltonian

$$\Delta_\omega = \Delta_\omega^* \in \{H\}' \quad (7)$$

$$\Gamma_{+,\omega} = c_\omega \frac{1}{e^{\beta(\omega)\omega} - 1} \quad ; \quad \Gamma_{-,\omega} = c_\omega \frac{e^{\beta(\omega)\omega}}{e^{\beta(\omega)\omega} - 1} \quad ; \quad c_\omega \geq 0 \quad (8)$$

β a regular enough function (inverse temperature) with

$$\beta(\omega) > 0 \quad ; \quad \forall \omega \in B_+ \quad (9)$$

The predual generator has the form

$$\mathcal{L}_* = \sum_{\omega \in B_+} \mathcal{L}_{\omega*}$$

where for each Bohr frequency ω ,

$$\begin{aligned} \mathcal{L}_{\omega*}(\rho) = & \left(-\frac{\Gamma_{-,\omega}}{2} \{D_\omega^* D_\omega, \rho\} + \Gamma_{+,\omega} D_\omega^* \rho D_\omega \right) \\ & + \left(-\frac{\Gamma_{+,\omega}}{2} \{D_\omega D_\omega^*, \rho\} + \Gamma_{-,\omega} D_\omega \rho D_\omega^* \right) - i[\Delta_\omega, \rho] \end{aligned} \quad (10)$$

From now on we will write simply P_n instead P_{ϵ_n} whenever there is no confusion.

3 Stationary states in the annihilator of all Kraus operators

In this section we study stationary states of WCLT generators that belong to the annihilator of all Kraus operators defined as

$$Ann(D) := \{\rho : \text{tr}(\rho D_\omega) = 0, \text{ for every Bohr frequency } \omega\}$$

Notice that

$$\{H\}'' \subseteq \{H\}'$$

with equality if H is non degenerate. Moreover, for the class of WCLT Markov generators the inequality

$$\{H\}' \subset Ann(D)$$

holds. Indeed, for any $\rho \in \{H\}'$,

$$\text{Tr}(\rho \sum_{(\epsilon_n, \epsilon_m) \in B_{+, \omega}} P_m D P_n) = \sum_{(\epsilon_n, \epsilon_m) \in B_{+, \omega}} \text{Tr}(P_n P_m \rho D) = 0,$$

since $n \neq m$. But one can find WCLT generators with invariant states $\rho \in Ann(D) \setminus \{H\}'$. See Corollary 27, below.

3.1 Stationary states supported on the interaction-free subspace

We start by describing the simplest class of stationary states in the annihilator of all Kraus operators, which are characterized by the stronger condition $\rho D_\omega = 0 = D_\omega \rho$ for all ω . The support subspace of these stationary states, defined as the closure of its range, is contained in the intersection of the kernels of all Kraus operators; moreover, these stationary states do not depend on the coefficients $\Gamma_{\pm, \omega}$.

Definition 1. (i) We call singular any subset $\Theta \subseteq \mathcal{B}(\mathfrak{h})$ satisfying the condition

$$\ker \Theta := \bigcap_{\theta \in \Theta} \ker \theta \neq \{0\}$$

(ii) The interaction-free subspace is defined by

$$W_D := \bigcap_{\omega \in \mathbf{B}_+} (\ker D_\omega \cap \ker D_\omega^*)$$

Remark 2. The relation of the interaction-free subspace W_D with the notion of decoherence for quantum Markov semigroups[6, 7, 8] and in particular with the decoherence-free subspace[19, 18, 2] will be discussed in forthcoming work.

The proof of the following proposition is straightforward.

Proposition 3. Let \mathcal{L} be a WCLT generator and $D \in \mathcal{B}(\mathfrak{h})$ an interaction operator. Then,

(i) $W_D \neq \{0\}$ if and only if the subset

$$\Theta := \{P_l D P_k, P_k D^* P_l : l < k\}$$

is singular.

(ii) $0 \neq u \in W_D$ if and only if $u = \sum_l P_l u = \sum_l u_l$, with

$$u_l \in V_l := \left(\bigcap_{k < l} \ker(P_k D P_l) \bigcap_{k > l} \ker(P_k D^* P_l) \right) \cap P_l(\mathfrak{h})$$

for every l .

(iii) $V_k \perp V_l$ if $k \neq l$ and $W_D = \bigoplus_l V_l$.

Theorem 4. *With the notations in the above proposition, let $\mathcal{L}_* := \mathcal{L}_{*0} + i\delta_\Delta$ be a Markov generator of WCLT with associated reference Hamiltonian $H \in \mathcal{B}(\mathfrak{h})$. Let $D \in \mathcal{B}(\mathfrak{h})$ be an interaction operator satisfying $W_D \neq \{0\}$. Then for every $u_j \in W_D$, $\|u_j\| = 1$ and $\sum_j \lambda_j = 1$, the state*

$$\rho := \sum_j \lambda_j |u_j\rangle\langle u_j| \quad (11)$$

*satisfies $\mathcal{L}_{*0}(\rho) = 0$. Moreover, if $\delta_{\Delta_\omega}(\rho) = [\Delta_\omega, \rho]$, where*

$$\Delta_\omega = \zeta_{-, \omega} D_\omega^* D_\omega + \zeta_{+, \omega} D_\omega D_\omega^* \quad (12)$$

and $\zeta_{-, \omega}, \zeta_{+, \omega} \in \mathbb{R}$, then ρ is an \mathcal{L}_ -invariant state.*

Proof. The generator \mathcal{L}_{*0} is a closed operator on the Banach space of finite-trace operators on \mathfrak{h} . Being ρ the limit of a sequence of finite-rank approximations $\rho_N \in \text{dom}(\mathcal{L}_{*0})$, the identity $\mathcal{L}_{*0}(\rho_N) = 0$ for all $n \geq 0$, implies that $\rho \in \text{dom}(\mathcal{L}_{*0})$ and $\mathcal{L}_{*0}(\rho) = 0$. Moreover, condition $u_j \in W_D$ for every j , implies that Δ_ω commutes with ρ . \square

Remark 5. (i) Notice that condition $u \in W_D$, implies that the pure state $|u\rangle\langle u|$ is an extremal stationary state. Moreover this operator belongs to the fixed points $\mathcal{F}(\mathcal{T})$ of the direct semigroup $(\mathcal{T}_t)_{t \geq 0}$.

(ii) In the finite-dimensional case, one can write $\mathcal{B}(\mathbb{C}^d) = \{H\}' \oplus (\{H\}')^\perp$, with respect to the Hilbert-Schmidt inner product. Then for every $D \in \mathcal{B}(\mathbb{C}^d)$, there exist unique $D_1 \in \{H\}'$ and $D_2 \in (\{H\}')^\perp$ such that $D = D_1 + D_2$. Hence, for each Bohr frequency $\omega := \epsilon_n - \epsilon_m > 0$ we have

$$\begin{aligned} D_\omega &:= \sum_{(\epsilon_n, \epsilon_m) \in \mathbf{B}_{+, \omega}} P_m D P_n = \sum_{(\epsilon_n, \epsilon_m) \in \mathbf{B}_{+, \omega}} P_m (D_1 + D_2) P_n \\ &= \sum_{(\epsilon_n, \epsilon_m) \in \mathbf{B}_{+, \omega}} P_m D_1 P_n + \sum_{(\epsilon_n, \epsilon_m) \in \mathbf{B}_{+, \omega}} P_m D_2 P_n \\ &= \sum_{(\epsilon_n, \epsilon_m) \in \mathbf{B}_{+, \omega}} P_m D_2 P_n \end{aligned}$$

This implies that the interesting interaction operators are those belonging to $(\{H\}')^\perp$.

Our next theorem gives a characterization of states supported on W_D in terms of its product with Kraus operators.

Theorem 6. *An invariant state ρ has the form (11) for some $u_j \in W_D$ and $\lambda_j > 0$ if and only if*

$$\rho D_\omega = 0 = D_\omega \rho \quad (13)$$

for every Bohr frequency ω .

Proof. If ρ has the form (11) then it immediately follows that $\rho D_\omega = 0 = D_\omega \rho$ since the support of ρ is a subset of W_D .

Conversely, let ρ be a state, then $\rho = \sum_k \rho_k |u_k\rangle\langle u_k|$ for some complete orthonormal system (u_k) of \mathfrak{h} and $\rho_k > 0$. Moreover, if for every Bohr frequency $\rho D_\omega = 0 = D_\omega \rho$ then $\sum_k \rho_k |D_\omega u_k\rangle\langle u_k| = D_\omega \rho = 0$. Therefore for every u_l we have that $0 = \sum_k \rho_k |D_\omega u_k\rangle\langle u_k| u_l = \rho_l D_\omega u_l$, which implies that $u_l \in \ker D_\omega$ for every frequency ω . Likewise, one proves that $u_l \in \ker D_\omega^*$ for every ω . We can conclude that $u_l \in W_D$ for every l and, hence, ρ has the form (11). This finishes the proof. \square

3.2 Local detailed balance stationary states

The stationary states of the form (11) in Theorem 4, are the simplest invariant states of a WCLT generator but not all invariant states belong to this class. In the case of a degenerate reference Hamiltonian, the stationary states exhibit a block structure, with blocks that may depend on the Γ 's, see Section 4 and examples in Subsection 3.3.

Given a state ρ and a subspace V , by " ρ is supported on V " we mean that " ρ is a subspace of V ". We start with a generalization of the theorem in previous section.

Theorem 7. *With the notations in Theorem 4, let $\mathcal{L}_* = \mathcal{L}_{*0} + i\delta_\Delta$ be a WCLT generator associated with (possibly degenerate) H . Then*

- (i) *a state ρ commutes with the orthogonal projection on the subspace W_D , P_{W_D} , if and only if ρ has the structure*

$$\rho = P_{W_D} \rho P_{W_D} + P_{W_D^\perp} \rho P_{W_D^\perp} \quad (14)$$

where $P_{W_D^\perp}$ is the orthogonal projection on W_D^\perp .

- (ii) *If moreover ρ is invariant for \mathcal{L}_* , then*

$$\rho = \theta \rho_{W_D} + (1 - \theta) \rho_{W_D^\perp}, \quad \theta \in [0, 1] \quad (15)$$

with ρ_{W_D} an invariant state supported on W_D and $\rho_{W_D^\perp}$ an invariant state supported on W_D^\perp .

Proof. We have that, $P_{W_D} + P_{W_D^\perp} = \mathbb{1}$, then for any state ρ ,

$$P_{W_D}\rho + P_{W_D^\perp}\rho = \rho = \rho P_{W_D} + \rho P_{W_D^\perp}$$

Hence, ρ commutes with P_{W_D} if and only if it commutes with $P_{W_D^\perp}$.

The assumption ρ commutes with P_{W_D} and simple computations show that

$$\rho = (P_{W_D} + P_{W_D^\perp})\rho(P_{W_D} + P_{W_D^\perp}) = P_{W_D}\rho P_{W_D} + P_{W_D^\perp}\rho P_{W_D^\perp}$$

Conversely, if ρ has the above structure, then the subspaces W_D and W_D^\perp are invariant under the action of ρ and this implies that ρ commutes with the orthogonal projection P_{W_D} . This proves (i).

If $\rho = \sum_j \rho_j |u_j\rangle\langle u_j|$ for some $u_j \in \mathfrak{h}$ commutes with P_{W_D} , then

$$P_{W_D}\rho = P_{W_D}\rho P_{W_D} = \sum_j \rho_j |P_{W_D}u_j\rangle\langle P_{W_D}u_j|$$

that has the form (11) and, hence, its is invariant. Consequently, if ρ is invariant then $P_{W_D^\perp}\rho P_{W_D^\perp} = \rho - P_{W_D}\rho P_{W_D}$ is invariant due to the linearity of the infinitesimal generator \mathcal{L}_* . Moreover, the support of $P_{W_D}\rho P_{W_D} = P_{W_D}\rho$ is a subset of W_D and the support of $P_{W_D^\perp}\rho P_{W_D^\perp} = P_{W_D^\perp}\rho$ is orthogonal with the former. Item (ii) readily follows after normalization. \square

Theorem 8. *Let \mathcal{L}_* be a WCLT generator associated with a (possible degenerate) reference Hamiltonian H and let ρ be a state. If for every Bohr frequency ω the following condition holds*

$$\rho D_\omega = c_\omega D_\omega \rho, \tag{16}$$

for some $c_\omega \neq 0$ that may depend on the Bohr frequencies, then,

(i) ρ commutes with $D_\omega D_\omega^*$ and $D_\omega^* D_\omega$ for every ω .

(ii) ρ has the structure (14) in the above theorem

(iii) if moreover $c_\omega = \frac{\Gamma_{-,\omega}}{\Gamma_{+,\omega}}$, $\forall \omega$, then ρ is a stationary state.

In particular, any state ρ satisfying (16) belongs to the annihilator of all Kraus operators.

Proof. Item (i), follows immediate from (16).

If (16) holds for every Bohr frequency, then $u \in \ker D_\omega$ implies that $0 = \rho D_\omega u = c_\omega D_\omega \rho u$, meaning that $\rho u \in \ker D_\omega$. Likewise for every $u \in \mathfrak{h}$ one has that $\rho D_\omega^* u = c_\omega^{-1} D_\omega^* \rho u$, then $v = D_\omega^* u \in \text{Im} D_\omega^* = (\ker D_\omega)^\perp$ implies that $\rho v \in \text{Im} D_\omega^*$. This implies that the subspaces $\ker D_\omega$ and $(\ker D_\omega)^\perp$ are ρ -invariant. Therefore ρ commutes with each orthogonal projection $P_{\ker D_\omega}$, consequently, commutes with $P_{W_D} = \Pi_{\omega \in B_+} P_{\ker D_\omega} P_{\ker D_\omega^*}$. As a consequence of item (i) in the above theorem, ρ has the required structure (14) and this proves (ii).

To prove (iii) observe that for any state satisfying (16) with $c_\omega = \frac{\Gamma_{-, \omega}}{\Gamma_{+, \omega}}$ for all ω we have

$$\begin{aligned} \mathcal{L}_{\omega^*}(\rho) &= \left(-\frac{\Gamma_{-, \omega}}{2} \{D_\omega^* D_\omega, \rho\} + \Gamma_{+, \omega} D_\omega^* \rho D_\omega \right) \\ &\quad + \left(-\frac{\Gamma_{+, \omega}}{2} \{D_\omega D_\omega^*, \rho\} + \Gamma_{-, \omega} D_\omega \rho D_\omega^* \right) - i[\Delta_\omega, \rho] \\ &= (-\Gamma_{-, \omega} + c_\omega \Gamma_{+, \omega}) D_\omega^* D_\omega \rho + (-\Gamma_{+, \omega} + c_\omega^{-1} \Gamma_{-, \omega}) \rho D_\omega^* D_\omega \\ &\quad - i\rho \left(\zeta_{-, \omega} D_\omega^* D_\omega + \zeta_{+, \omega} D_\omega D_\omega^* - c_\omega^{-1} c_\omega \zeta_{-, \omega} D_\omega^* D_\omega \right. \\ &\quad \left. - \zeta_{+, \omega} c_\omega c_\omega^{-1} D_\omega D_\omega^* \right) = 0 \end{aligned} \quad (17)$$

for every Bohr frequency. This proves that ρ is invariant.

Finally, inequality $\Gamma_{-, \omega} > \Gamma_{+, \omega}$ implies that $c_\omega \neq 1$ for all ω , then condition $\rho D_\omega = c_\omega D_\omega \rho$ yields $\text{tr}(\rho D_\omega) = c_\omega \text{tr}(D_\omega \rho) = c_\omega \text{tr}(\rho D_\omega)$, which in turn implies that $\text{tr}(\rho D_\omega) = 0$. Consequently ρ belongs to the annihilator of all Kraus operators. This finishes the proof. \square

Definition 9. (i) Any \mathcal{L}_* -invariant state satisfying (16) will be called *local detailed balance invariant state*.

(ii) If moreover, in (16) $c_\omega = \frac{\Gamma_{-, \omega}}{\Gamma_{+, \omega}}$, $\forall \omega$, the corresponding \mathcal{L}_* -invariant state will be called *detailed balance invariant state*.

Remark 10. Any detailed balance invariant state is local detailed balance, but there exists local detailed balance states that are not detailed balance, see for instance Ref.[9, 17]

The result of the following lemma is essentially contained in the proof of Proposition 7 in Ref.[5]

Lemma 11. $W_D = \bigcap_{\omega \in B_+} \ker \left(D_\omega D_\omega^* + D_\omega^* D_\omega \right)$.

Proof. Inequality $W_D \subset \bigcap_{\omega \in B_+} \ker (D_\omega D_\omega^* + D_\omega^* D_\omega)$ is evident.

Now, if $D_\omega D_\omega^* u + D_\omega^* D_\omega u = 0$, then identity

$$0 = \langle u, D_\omega D_\omega^* u \rangle + \langle u, D_\omega^* D_\omega u \rangle = \|D_\omega^* u\|^2 + \|D_\omega u\|^2$$

imply that $\|D_\omega^* u\|^2 = 0$ and $\|D_\omega u\|^2 = 0$, hence $u \in \ker D_\omega \cap \ker D_\omega^* \subset W_D$. \square

Proposition 12. *The orthogonal projection P_{W_D} on W_D , belongs to the commutant of H .*

Proof. By Lemma 11, we have that

$$W_D = \bigcap_{\omega \in B_+} \ker (D_\omega D_\omega^* + D_\omega^* D_\omega) = \bigcap_{\omega \in B_+} \ker (D_\omega D_\omega^*) \cap \ker (D_\omega^* D_\omega) \quad (18)$$

where for every $\omega = \epsilon_n - \epsilon_m$,

$$D_\omega = \sum_{(\epsilon_n, \epsilon_m) \in B_{+, \omega}} P_{\epsilon_m} D P_{\epsilon_n} \quad (19)$$

We affirm that each spectral projection P_{ϵ_k} commutes with P_{W_D} , i.e., $P_{W_D} \in \{H\}'$. Since

$$P_{W_D} = \Pi_{\omega \in B_+} P_{\ker(D_\omega D_\omega^*)} P_{\ker(D_\omega^* D_\omega)}$$

it suffices to prove that every spectral projection commutes with $P_{\ker(D_\omega D_\omega^*)}$ and $P_{\ker(D_\omega^* D_\omega)}$ for all $\omega \in B_+$.

Due to Lemma 3.1 in Ref.[1] condition $(\epsilon_n, \epsilon_m), (\epsilon_{n'}, \epsilon_m) \in B_{+, \omega}$ implies $\epsilon_n = \epsilon_{n'}$. Hence, each P_{ϵ_m} appears on the left of only one summand of (19). Moreover, one can order the set $B_{+, \omega} := \{\epsilon_m : \epsilon_m + \omega \in \text{Sp}(H)\}$ so that $B_\omega = \{\epsilon_0(\omega) < \epsilon_1(\omega) < \dots\}$. From now on we write simply ϵ_k instead $\epsilon_k(\omega)$ and we can write $D_\omega = \sum_{\epsilon_m \in B_{+, \omega}} P_{\epsilon_m} D P_{\epsilon_m + \omega}$. Now, after simple computations we get

$$\begin{aligned} D_\omega D_\omega^* &= \sum_{\epsilon_m \in B_{+, \omega}} \sum_{\epsilon_{m'} \in B_{+, \omega}} P_{\epsilon_m} D P_{\epsilon_m + \omega} P_{\epsilon_{m'} + \omega} D^* P_{\epsilon_{m'}} \\ &= \sum_{\epsilon_m \in B_{+, \omega}} P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m} \end{aligned} \quad (20)$$

Orthogonality of ranges of the above summands readily implies that

$$P_{\ker(D_\omega D_\omega^*)} = \Pi_{\epsilon_m \in B_{+, \omega}} P_{\ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m})}$$

Clearly $\ker P_{\epsilon_m} \subseteq \ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m})$, hence for every $m \geq 0$,

$$P_{\ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m})} = P_{\ker P_{\epsilon_m}} + P_{\ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m}) \cap \text{Im} P_{\epsilon_m}}$$

with the second summand being a subprojection of P_{ϵ_m} . Therefore we have that

$$\begin{aligned} P_{\epsilon_l} P_{\ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m})} &= P_{\epsilon_l} = P_{\ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m})} P_{\epsilon_l}, \quad l \neq m \\ P_{\epsilon_m} P_{\ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m})} &= P_{\ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m}) \cap \text{Im} P_{\epsilon_m}} \\ &= P_{\ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m})} P_{\epsilon_m} \end{aligned} \quad (21)$$

This proves the commutativity of each spectral projection with every

$P_{\ker(P_{\epsilon_m} D P_{\epsilon_m + \omega} D^* P_{\epsilon_m})}$ and, consequently, with $P_{\ker(D_\omega D_\omega^*)}$.

The commutativity with $P_{\ker(D_\omega^* D_\omega)}$ is proved in a similar way. \square

Remark 13. As a consequence of the above Proposition, one can add the term $i[\Delta_1, \rho]$ to any WCLT generator, with Δ_1 any real function of the Hamiltonian, and if $\rho \in \{H\}'$, still have that $P_{W_D} \rho P_{W_D}$ is invariant.

Remark 14. (Quantum detailed balance) In the case when a detailed balance state ρ is faithful, condition (16) yields a privileged GKS representation of \mathcal{L} , in the sense of Ref.[12] Therefore, in this case, our detailed balance condition implies the quantum detailed balance condition $\mathcal{L} - \tilde{\mathcal{L}} = 2i[\Delta_\omega, \cdot]$ where $\tilde{\mathcal{L}}$ is the 0-dual generator. Moreover, in this case,

- The 0-dual \mathcal{T} is a quantum Markov semigroup and
- The Markov generator \mathcal{L} commutes with σ_{-i} , where (σ_t) is the associated modular group $\sigma_t(x) = \rho^{it} x \rho^{-it}$, see Ref.[12]

Remark 15. As we have shown in Theorem 8, condition (16) implies that the state ρ belongs to $\text{Ann}(D)$. The reciprocal is not true, indeed, taken $H = \epsilon_0 |e_0\rangle\langle e_0| + \epsilon_1 (|e_1\rangle\langle e_1| + |e_2\rangle\langle e_2|)$ and $D = |e_0\rangle\langle e_1| + |e_1\rangle\langle e_2|$, one can easily see that the state $\rho = \frac{1}{2} |e_0\rangle\langle e_0| + \frac{1}{4} |e_1 + e^{i\theta} e_2\rangle\langle e_1 + e^{i\theta} e_2|$ belongs to $\text{Ann}(D)$, but doesn't exist a constant c_ω such that $\rho D_\omega = c_\omega D_\omega \rho$ for all ω .

3.3 Examples

Example 16. (Degenerate H) Consider a (possible degenerate) reference Hamiltonian with the property that for every Bohr frequency ω there exists a unique pair of eigenvalues (ϵ_n, ϵ_m) such that $\omega = \epsilon_n - \epsilon_m$ (if moreover H is non-degenerate then it is called generic). Let \mathcal{L} be a Markov generator associated with H , then $D_\omega = P_m D P_n$ if $\omega = \epsilon_n - \epsilon_m$, with not necessarily

rank-one spectral projections P_n . Let ρ be a faithful invariant state belonging to $\{H\}' \subset \{H\}''$, since for every ω , $\rho D_\omega = \frac{\rho_m}{\rho_n} D_\omega \rho$, the detailed balance condition in the above theorem holds true iff $\frac{\rho_m}{\rho_n} = \frac{\Gamma_{-, \omega}}{\Gamma_{+, \omega}}$ for every ω ; consequently, the state is invariant and has the (block) structure given in Theorem 8. Moreover, this condition yields the (block) Gibbs state

$$\rho = \frac{1}{Z} e^{-\beta(H)H}$$

Example 17. (Dark states in quantum photosynthesis (KV's model))

The invariant states of a WCLT generator modeling quantum photosynthesis, proposed by Kozyrev and Volovich[11] in the context of stochastic limit approach of degenerate quantum open systems, were studied in Ref.[16] We will show that the so called dark stationary states of that model correspond with states supported on the interaction-free subspace.

In this case the reference Hamiltonian has the spectral decomposition

$$H = \epsilon_0 P_0 + \epsilon_1 P_1 + \epsilon_2 P_2$$

where $P_0 = |e_0\rangle\langle e_0|$, $P_1 = |e_1\rangle\langle e_1|$, $P_2 = \sum_{j=2}^N |e_j\rangle\langle e_j|$ and $\epsilon_2 > \epsilon_1 > \epsilon_0 = 0$. The set of Bohr frequencies is $B_+ = \{\omega_1 = \epsilon_2, \omega_2 = \epsilon_2 - \epsilon_1, \omega_3 = \epsilon_1\}$. At level two there are two linearly dependent maximal entangled vectors: the bright photonic vector $\chi = \sum_{j=2}^N e_j$ and the bright phononic vector $\Psi = e^{i\theta}\chi$, $\theta \in (0, 2\pi) \setminus \{\pi\}$, so that the corresponding pure states coincide $|\chi\rangle\langle\chi| = |\Psi\rangle\langle\Psi|$.

Up to some constants the interaction operators are:

$$\begin{aligned} D_{\omega_1} &= |e_0\rangle\langle\chi| \\ D_{\omega_2} &= |e_1\rangle\langle\Psi| \\ D_{\omega_3} &= |e_0\rangle\langle e_1| \end{aligned} \tag{22}$$

Therefore $W_D = \{e_0, e_1, \chi\}^\perp$. Consequently, any invariant state supported on the interaction-free subspace W_D is orthogonal with the bright photonic state $|\chi\rangle\langle\chi|$ and also orthogonal with the pure states $|e_0\rangle\langle e_0|$ and $|e_1\rangle\langle e_1|$, hence, it is a global dark state in the sense of Ref.[11], i.e., dark with respect to the full generator \mathcal{L} . This result coincides with the result found in Ref.[16], where it was proved that any invariant state has the spectral decomposition

$$\rho = \lambda \left(r_0 P_0 + r_1 P_1 + \frac{1}{N-1} |\chi\rangle\langle\chi| \right) + (1 - \lambda(r_0 + r_1 - 1)) \rho_{W_D} \tag{23}$$

where ρ_{W_D} is a dark state, r_0, r_1 are functions of the Γ 's and $0 \leq \lambda \leq (1 + r_0 + r_1)^{-1}$. Indeed, assuming that ρ is invariant and defining

$$(1 - \lambda(r_0 + r_1 - 1)) \rho_{W_D} := \rho - \lambda \left(r_0 P_0 + r_1 P_1 + \frac{1}{N-1} |\chi\rangle\langle\chi| \right)$$

one has that ρ_{W_D} is:

- supported on (just complete the diagonalization of ρ) $W_D = \{e_0, e_1, \chi\}^\perp$ consequently, it is invariant, and
- orthogonal w.r.t. the Hilbert-Schmidt product with P_0 , P_1 and $|\chi\rangle\langle\chi|$, then it is global dark.

4 Invariant states of a modified AKV's model

The invariant states of a slightly modification of a quantum transport model due to Aref'eva, Kozyrev and Volovich[4], were characterized in Ref.[16] in terms of some conditions including commutation relations with remarkable operators of the model.

In this section we will give a full description of the set of invariant states of the model, including their parametrization in terms of states supported on the range of $|Z|^2 = Z^*Z$, where Z is the so called interference operator. It turns out that Z plays the role of a “generalized annihilation operator” mapping the second eigen-space of the reference Hamiltonian H into the third one, this analogy allows us to compute all invariant states by a simple method and notably less computations than those performed in the previous reference. Moreover we identify the fast recurrent subspace and study the approach to equilibrium property.

4.1 The modified AKV's quantum transport model

The corresponding generator belongs to the class of WCLT and the associated reference Hamiltonian is given by

$$H = 0P_0 + \epsilon_1 P_1 + \epsilon_2 P_2 + \epsilon_3 P_3$$

with $P_0 = |e_0\rangle\langle e_0|$, $P_1 = |e_1\rangle\langle e_1|$, $P_2 = \sum_{j=2}^N |e_j\rangle\langle e_j|$, $P_3 = \sum_{j=N+1}^{N+M} |e_j\rangle\langle e_j|$, $M \leq N - 1$ and the eigenvalues satisfy the inequalities $0 < \epsilon_3 < \epsilon_2 < \epsilon_1$. The interaction operator is

$$D = |Te_{N+1}\rangle\langle Te_1| + Z + |Te_0\rangle\langle Te_{N-1}|$$

where Z is an interference operator performing transitions between levels ϵ_2 and ϵ_3

$$Z : P_2(\mathbf{h}) \mapsto P_3(\mathbf{h})$$

defined in terms of some interference coefficients g_{ab} (see Remark 2.2 in Ref.[16]) as

$$Z = \frac{1}{\sqrt{N-1}} \sum_{b=N+1}^{N+M} \sum_{a=2}^N g_{ab} |b\rangle\langle a|$$

and $T = P_0 + P_1 + Z + Z^*$. The maximally entangled states are defined, up to normalization, as $|\psi\rangle\langle\psi|$ with $\psi = \sum_{a=2}^N |a\rangle$ and $|\psi'\rangle\langle\psi'|$ with $\psi' = \sum_{b=N+1}^{N+M} |b\rangle$. Writing simply T_a instead $T|a\rangle$, one can observe that $\sqrt{N-1}Z_{N-1} = \psi'$ and $\sqrt{N-1}Z_{N+1}^* = \psi$.

Hence, there exist six Bohr frequencies: $\omega_1 = \epsilon_1 - \epsilon_2$, $\omega_2 = \epsilon_2 - \epsilon_3$, $\omega_3 = \epsilon_3$, $\omega_4 = \epsilon_1$, $\omega_5 = \epsilon_2$, $\omega_6 = \epsilon_1 - \epsilon_3$. But only three frequencies have associated non-trivial generators, so that

$$\mathcal{L} = \mathcal{L}_{\omega_1} + \mathcal{L}_{\omega_2} + \mathcal{L}_{\omega_3}$$

$$\text{and } \mathcal{L}_{\omega_4} = \mathcal{L}_{\omega_5} = \mathcal{L}_{\omega_6} = 0.$$

Remark 18. The relations $T = T^*$, $Z^2 = 0$ and $Z^{*2} = 0$ immediately follow. Moreover, by direct computation one can show that $|Z|$ is an orthogonal projection, $TP_2T = P_3 = ZZ^*$ and $TP_3T = |Z|$, see Corollary 2.1 in Ref.[16]

One can see that

$$\begin{aligned} D_{\omega_1} &= \frac{1}{\sqrt{N-1}} |\psi\rangle\langle e_1| \\ D_{\omega_2} &= Z \\ D_{\omega_3} &= \frac{1}{\sqrt{N-1}} |e_0\rangle\langle\psi'| \end{aligned} \tag{24}$$

So that the Kraus operators are

$$\begin{aligned} L_1^{\omega_1} &= \sqrt{2\Gamma_{Re,-,\omega_1}} |\psi\rangle\langle e_1| \\ L_1^{\omega_2} &= \sqrt{2(N-1)\Gamma_{Re,-,\omega_2}} Z \\ L_1^{\omega_3} &= \sqrt{2\Gamma_{Re,-,\omega_3}} |e_0\rangle\langle\psi'| \\ L_2^{\omega_1} &= \sqrt{2\Gamma_{Re,+, \omega_1}} |e_1\rangle\langle\psi| \\ L_2^{\omega_2} &= \sqrt{2(N-1)\Gamma_{Re,+, \omega_2}} Z^* \\ L_2^{\omega_3} &= 0 \end{aligned} \tag{25}$$

with all coefficients Γ_{Re,\pm,ω_l} , $1 \leq l \leq 2$ positive, $\Gamma_{Re,-,\omega_3} > 0$ and $\Gamma_{Re,+, \omega_3} = 0$.

The effective Hamiltonian is give by

$$\begin{aligned} H_{eff} &= \Gamma_{Im,+, \omega_1} |\psi\rangle\langle\psi| - (N-1)\Gamma_{Im,-, \omega_1} |e_1\rangle\langle e_1| \\ &\quad + (N-1)\Gamma_{Im,+, \omega_2} P_3 - (N-1)\Gamma_{Im,-, \omega_2} |Z| \\ &\quad - \Gamma_{Im,-, \omega_3} |\psi'\rangle\langle\psi'| \end{aligned} \tag{26}$$

with all coefficients $\Gamma_{Im, \pm, \omega_l}$, $1 \leq l \leq 3$, positive.

Direct computation shows that

$$\begin{aligned} W_D &= \{e_1, \psi, \psi'\}^\perp \cap \ker Z \cap \ker Z^* \\ &= \{e_1, \psi, \psi'\}^\perp \cap \{Z^* e_j : j = N+1, \dots, N+M\}^\perp \cap \{Z e_j : j = 2, \dots, N\}^\perp \end{aligned} \quad (27)$$

The vectors ψ and $Z^* \psi'$ are fixed points of the orthogonal projection $|Z|$. Indeed,

$$|Z|\psi = Z^* Z \sqrt{N-1} Z^* e_{N+1} = \sqrt{N-1} Z^* e_{N+1} = \psi$$

and

$$|Z|Z^* \psi' = Z^* Z Z^* \psi' = Z^* \psi'$$

where we have used the identity $ZZ^* = P_3$. Hence we can write

$$\text{Im}|Z| = \text{span}\{\psi, Z^* \psi'\} \oplus \text{Im}|Z| \cap \{\psi, Z^* \psi'\}^\perp$$

Moreover, identity $P_2(\mathbf{h}) = \text{Im}|Z| \oplus \ker |Z|$ yields the decomposition

$$P_2(\mathbf{h}) = \text{span}\{\psi, Z^* \psi'\} \oplus \left(\text{Im}|Z| \cap \{\psi, Z^* \psi'\}^\perp \right) \oplus \ker |Z|$$

Similarly one can see that ψ' , $Z\psi$ are fixed points of P_3 and the decomposition

$$P_3(\mathbf{h}) = \text{span}\{\psi', Z\psi\} \oplus \text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$$

holds true.

4.2 Subharmonic projections and structure of stationary states

A positive operator a is subharmonic (resp. superharmonic, resp. harmonic) for a quantum Markov semigroup $(\mathcal{T})_{t \geq 0}$ if $\mathcal{T}_t(a) \geq a$ (resp. $\mathcal{T}_t(a) \leq a$, resp. $\mathcal{T}_t(a) = a$) for all $t \geq 0$. Subharmonic projections play a fundamental role in the study of irreducibility of the semigroup, see Ref.[13], in particular, orthogonal projections onto the support subspaces of invariant states are subharmonic. We will compute harmonic and subharmonic projections of the modified AKV's model as a first step in the characterization of invariant states.

Proposition 19. *Each rank-one projection $|w\rangle\langle w|$ with $w \in W_D$ is harmonic for the AKV's semigroup, consequently P_{W_D} and $P_{W_D^\perp}$ are by itself harmonic projections.*

Proof. It suffices to prove that every rank-one projection $|w\rangle\langle w|$ with $w \in W_D$ commutes with the effective Hamiltonian $H_{eff} = H_{\omega_1} + H_{\omega_2} + H_{\omega_3}$. But clearly $|w\rangle\langle w|$ commutes with H_{ω_1} since $w \in \{e_1, \psi\}^\perp$. Similarly it commutes with H_{ω_3} since $w \perp \psi'$. Now, for every $v \in P_3(\mathbf{h})$ identity $Z^*v = 0$ implies that $v = P_3v = ZZ^*v = 0$, hence $\ker Z^* = \ker P_3$, it readily follows that $P_3w = 0$. Moreover, $\ker |Z| = \ker Z$, hence $w \in \ker Z$. This implies that $|w\rangle\langle w|$ commute with H_{ω_2} and, hence, commute with H_{eff} . We can conclude that every $|w\rangle\langle w|$ is harmonic. Being a linear combination of the above class of rank-one projections, P_{W_D} is harmonic projection by itself. Moreover, due to the conservativity, we have that

$$\mathcal{T}_t(P_{W_D^\perp}) = \mathcal{T}_t(I) - \mathcal{T}_t(P_{W_D}) = I - P_{W_D} = P_{W_D^\perp}$$

This finishes the proof. \square

But apart from the above given projections, the AKV's semigroup has another subharmonic projections, as we show in the following.

Proposition 20. *The orthogonal projection p_V onto the subspace $V = \{e_0, e_1, \psi, \psi', Z\psi, Z^*\psi'\}^\perp$ is subharmonic for the quantum Markov semigroup of the modified AKV's model.*

Proof. By Theorem III.1 in Ref.[13], it suffices to prove that the orthogonal projection p_V satisfies that $Ran(p_V)$ is invariant under the action of the semigroup generated by the effective Hamiltonian and satisfy

$$L_k^{\omega_j}u = p_V L_k^{\omega_j}u, \quad 0 \leq j \leq 3, \quad 1 \leq k \leq 2, \quad \forall u \in Ran(p_V) \quad (28)$$

These conditions holds true if and only if $V = Ran(p_V)$ is invariant under the action of the semigroup generated by the effective Hamiltonian $H_{eff} = H_{\omega_1} + H_{\omega_2} + H_{\omega_3}$ and the following relations hold for all $u \in V$:

- (i) $\langle e_1, u \rangle \psi = \langle e_1, u \rangle p_V \psi,$
- (ii) $\langle \psi, u \rangle e_1 = \langle \psi, u \rangle p_V e_1$
- (iii) $p_V Z u = Z u,$
- (iv) $p_V Z^* u = Z^* u$
- (v) $\langle \psi', u \rangle e_0 = \langle \psi', u \rangle p_V e_0.$

Due to the explicit form of $H_{eff} = H_{\omega_1} + H_{\omega_2} + H_{\omega_3}$, it turns out that conditions (i) – (v) imply that V is invariant under the action of the unitary

group generated by this operator. Indeed, the only nontrivial fact is to show that p_V commutes with

$$H_{\omega_2} = (N-1)\Gamma_{Im,+,\omega_2}TP_2T - (N-1)\Gamma_{Im,-,\omega_2}TP_3T$$

But $TP_2T = P_3$ and $TP_3T = |Z|$ and for $u \in V$ one has $P_3u = ZZ^*u = Zp_VZ^*u = p_VZp_VZ^*u = p_VZZ^*u = p_VP_3u$, hence $P_3p_V = p_VP_3$. Similarly, for $u \in V$, using that $|Z|$ is a projection one has $|Z|u = Z^*Zu = Z^*p_VZu = p_VZ^*p_VZu = p_V|Z|u$, i.e., $|Z|p_V = p_V|Z|$.

Now, if p_V is the orthogonal projection on the subspace V conditions (i), (ii), (v) holds true for every $u \in V$. To show that conditions (iii) and (iv) hold, it suffices to prove that $Zu, Z^*u \in V$ for every $u \in V$. But for every $u \in V$, it readily follows that $Zu, Z^*u \in \{e_0, e_1, \psi, \psi'\}^\perp$. Moreover, notice that $\langle Zu, Z\psi \rangle = \langle u, |Z|^2\psi \rangle = \langle u, \psi \rangle = 0$ since $|Z|^2\psi = \sqrt{N-1}Z^*ZZ^*e_{N+1} = \sqrt{N-1}Z^*P_3e_{N+1} = \sqrt{N-1}Z^*e_{N+1} = \psi$ (i.e., $|Z|\psi = \psi$), and $\langle Zu, Z^*\psi' \rangle = \langle Z^2u, \psi' \rangle = 0$, since $Z^2 = 0$, hence $Zu \in V$. Finally, $Z^2 = 0$ implies that $\langle Z^*u, Z\psi \rangle = \langle u, Z^2\psi \rangle = 0$ and $\langle Z^*u, Z^*\psi' \rangle = \langle u, ZZ^*\psi' \rangle = \langle u, P_3\psi' \rangle = \langle u, \psi' \rangle = 0$, this shows that $Z^*u \in V$ and finishes the proof. \square

4.3 Invariant states

Now we look for invariant states associated with the non-trivial subharmonic projections. As a consequence of our Theorem 4, every state supported on a subspace of W_D is necessarily invariant. The existence of invariant states with support contained in W_D^\perp is a question that naturally arises. To answer this question we start with the following.

Lemma 21. *A state ρ supported on a subspace of $V \cap W_D^\perp$ is invariant if and only if $\{e_1, \psi, \psi'\} \in \ker \rho$, ρ belongs to the commutant $\{H\}'$ of the reference Hamiltonian, commute with $|Z|$ and satisfies*

$$Z^*\rho Z = \frac{\Gamma_{Re,-,\omega_2}}{\Gamma_{Re,+,\omega_2}}\rho|Z| \quad (29)$$

Proof. Let ρ be a state supported on a subspace of $V \cap W_D^\perp$, hence $\{e_1, \psi, \psi'\} \subset \ker \rho$. It readily follows that $\rho \in \ker \mathcal{L}_{*\omega_1} \cap \ker \mathcal{L}_{*\omega_3}$. Therefore, if ρ is invariant then

$$\begin{aligned} 0 = \mathcal{L}_{\omega_2*}(\rho) &= -i\left(\Gamma_{Im,+,\omega_2}P_3\rho - \Gamma_{Im,-,\omega_2}|Z|\rho\right) \\ &\quad + i\left(\Gamma_{Im,+,\omega_2}\rho P_3 - \Gamma_{Im,-,\omega_2}\rho|Z|\right) \\ &\quad - \Gamma_{Re,-,\omega_2}\left(|Z|\rho - 2Z\rho Z^* + \rho|Z|\right) \\ &\quad - \Gamma_{Re,+,\omega_2}\left(P_3\rho - 2Z^*\rho Z + \rho P_3\right) \end{aligned} \quad (30)$$

Therefore after direct computations we get

$$0 = P_2 \mathcal{L}_{\omega_2*}(\rho) P_3 = \left(-\Gamma_{Re,-,\omega_2} + i\Gamma_{Im,-,\omega_2} - \Gamma_{Re,+, \omega_2} + i\Gamma_{Im,+, \omega_2} \right) |Z| \rho P_3 \\ + \left(-\Gamma_{Re,+, \omega_2} + i\Gamma_{Im,+, \omega_2} \right) (P_2 - |Z|) \rho P_3 \quad (31)$$

Due to the positivity of the gammas and orthogonality of ranges we get that $|Z| \rho P_3 = 0$ and $(P_2 - |Z|) \rho P_3 = 0$. It follows that $P_2 \rho P_3 = 0$ and taking adjoints we get that also $P_3 \rho P_2 = 0$. But using that ρ is supported on a subspace of $W_D^\perp \cap V \subset (P_2 + P_3)(\mathfrak{h})$ we get,

$$[\rho, P_3] = (P_2 + P_3)[\rho, P_3](P_2 + P_3) = P_2 \rho P_3 - P_3 \rho P_2 = 0 \quad (32)$$

Hence ρ commutes with P_3 .

On the other hand, using the commutativity of ρ with P_3 from (30) and direct computations we get

$$0 = P_3 \mathcal{L}_{\omega_2*}(\rho) P_3 = \Gamma_{Re,-,\omega_2} Z \rho Z^* - \Gamma_{Re,+, \omega_2} P_3 \rho \quad (33)$$

It follows that

$$Z \rho Z^* = \frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,-,\omega_2}} P_3 \rho \quad (34)$$

Multiplication in (34) by Z^* on the left and by Z on the right yields

$$Z^* \rho Z = \frac{\Gamma_{Re,-,\omega_2}}{\Gamma_{Re,+, \omega_2}} |Z| \rho |Z| \quad (35)$$

To prove (29) it remains to prove that ρ commutes with $|Z|$.

Now, once again from (30) we get using (35)

$$0 = P_2 \mathcal{L}_{\omega_2*}(\rho) P_2 \\ = i\Gamma_{Im,-,\omega_2} (|Z| \rho P_2 - P_2 \rho |Z|) - \Gamma_{Re,-,\omega_2} (|Z| \rho P_2 + P_2 \rho |Z|) + 2\Gamma_{Re,+, \omega_2} Z^* \rho Z \\ = i\Gamma_{Im,-,\omega_2} (|Z| \rho P_2 - P_2 \rho |Z|) - \Gamma_{Re,-,\omega_2} (|Z| \rho P_2 + P_2 \rho |Z|) + 2\Gamma_{Re,-,\omega_2} |Z| \rho |Z| \\ = -\Gamma_{Re,-,\omega_2} |Z| \rho (P_2 - |Z|) - \Gamma_{Re,-,\omega_2} (P_2 - |Z|) \rho |Z| \\ + i\Gamma_{Im,-,\omega_2} |Z| \rho (P_2 - |Z|) - i\Gamma_{Im,-,\omega_2} (P_2 - |Z|) \rho |Z|$$

Due to orthogonality of ranges it follows that $|Z| \rho (P_2 - |Z|) = 0$ and $(P_2 - |Z|) \rho |Z| = 0$ consequently, $|Z| \rho P_2 = |Z| \rho |Z| = P_2 \rho |Z|$. But, after direct computations we get

$$[\rho, P_2 - |Z|] = (P_2 + P_3)[\rho, P_2 - |Z|](P_2 + P_3) = |Z| \rho P_2 - P_2 \rho |Z| = 0. \quad (36)$$

This proves that ρ commutes with $P_2 - |Z|$. Similarly, using that ρ commutes with P_3 , we get

$$[\rho, |Z|] = (P_2 + P_3)[\rho, |Z|](P_2 + P_3) = P_2\rho|Z| - |Z|\rho P_2 = 0. \quad (37)$$

Therefore ρ commutes with P_2 and with $|Z|$. The commutation of ρ with P_0 and P_1 is immediate. Hence we can conclude that $\rho \in \{H\}'$.

Conversely, if a state ρ is supported on a subspace of $V \cap W_D^\perp$, belongs to $\{H\}'$, commutes with $|Z|$ and satisfies (35), then $\rho \in \ker \mathcal{L}_{\omega_1*} \cap \ker \mathcal{L}_{\omega_2*} \cap \ker \mathcal{L}_{\omega_3*}$. This finishes the proof. \square

Now we are ready to characterize the set of invariant states of the model.

Theorem 22. *A state τ is invariant for the modified AKV's model if and only if it has the structure*

$$\tau = \lambda \rho_{W_D} + (1 - \lambda) \left(\frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}} \sigma + \frac{\Gamma_{Re, -, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}} Z \sigma Z^* \right) \quad (38)$$

where ρ_{W_D} is an invariant state supported on W_D , $\sigma : \text{Im}|Z| \mapsto \text{Im}|Z|$ is a state supported on a subspace of $V \cap \text{Im}|Z|$ and $0 \leq \lambda \leq 1$.

Proof. It suffices to show that a state ρ supported on a subspace of $V \cap W_D^\perp$ is invariant if and only if has the structure (38) with $\lambda = 0$. But, as a consequence of Lemma 21, any invariant state ρ supported on $V \cap W_D^\perp$ belongs to $\{H, |Z|\}'$ and can be written in the form

$$\begin{aligned} \rho &= (|Z| + P_3)\rho(|Z| + P_3) = |Z|\rho|Z| + P_3\rho P_3 \\ &= \frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}} \sigma + \frac{\Gamma_{Re, -, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}} Z \sigma Z^* \end{aligned} \quad (39)$$

where $\sigma = \frac{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}}{\Gamma_{Re,+, \omega_2}} |Z|\rho|Z|$ and we have used (34). Clearly, up to normalization, $\sigma : \text{Im}|Z| \mapsto \text{Im}|Z|$ is a state supported on a subspace of $V \cap \text{Im}|Z|$.

Conversely, if ρ is of the form (38) with $\lambda = 0$, then $e_1 \in \ker \rho$ and $\rho\psi = \frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}} \sigma\psi$, then for any $u \in \mathfrak{h}$ we have that $\langle u, \sigma\psi \rangle = \langle \sigma u, \psi \rangle = 0$ since $\sigma u \in V \cap \text{Im}|Z|$, hence $\sigma\psi = 0$, proving that $\psi \in \ker \rho$. Similarly one can see that $\psi' \in \ker \rho$. It readily follows that $\rho \in \ker \mathcal{L}_{\omega_1*} \cap \ker \mathcal{L}_{\omega_3*}$. Now by direct computations one can see that ρ commutes with H and $|Z|$, indeed, since σ commute with P_2 , then

$$\begin{aligned} P_2\rho &= \frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}} P_2\sigma = \rho P_2, \quad \text{and} \\ P_3\rho &= \frac{\Gamma_{Re, -, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}} Z \sigma Z^* = \rho P_3 \end{aligned} \quad (40)$$

Similarly,

$$|Z|\rho = \frac{\Gamma_{\text{Re},+,\omega_2}}{\Gamma_{\text{Re},+,\omega_2} + \Gamma_{\text{Re},-,\omega_2}}\sigma = \rho|Z|$$

Therefore, Lemma 21 implies that ρ is invariant. This finishes the proof. \square

Remark 23. Due to the above theorem, if τ is an invariant state for the modified AKV's model, then

$$\tau = \lambda\tau_{W_D} + (1 - \lambda)\tau_{W_D^\perp}, \quad 0 \leq \lambda \leq 1$$

with

$$\tau_{W_D^\perp} = \left(\frac{\Gamma_{\text{Re},+,\omega_2}}{\Gamma_{\text{Re},+,\omega_2} + \Gamma_{\text{Re},-,\omega_2}}\sigma + \frac{\Gamma_{\text{Re},-,\omega_2}}{\Gamma_{\text{Re},+,\omega_2} + \Gamma_{\text{Re},-,\omega_2}}Z\sigma Z^* \right)$$

The following result describes the extremal points of the subset of invariant states.

Theorem 24. *An invariant state ρ is extremal for the modified AKV's model if and only if it has one of the following structures:*

$$(i) \quad \rho = |u\rangle\langle u| \text{ with } u \in W_D = \{e_1, \psi, \psi'\}^\perp \cap \ker Z \cap \ker Z^*$$

(ii)

$$\rho = \frac{\Gamma_{\text{Re},+,\omega_2}}{\Gamma_{\text{Re},+,\omega_2} + \Gamma_{\text{Re},-,\omega_2}}|u\rangle\langle u| + \frac{\Gamma_{\text{Re},-,\omega_2}}{\Gamma_{\text{Re},+,\omega_2} + \Gamma_{\text{Re},-,\omega_2}}Z|u\rangle\langle u|Z^* \quad (41)$$

with $u \in V \cap \text{Im}|Z|$.

Proof. Due to Theorem 4, any state of the form $|u\rangle\langle u|$ with $u \in W_D$ is invariant and extremal. Conversely, if ρ is an extremal invariant state supported on W_D , then it is necessarily a pure state, i.e., $\rho = |u\rangle\langle u|$ for some $u \in W_D$.

Clearly any state ρ given by (41), is supported on $V \cap W_D^\perp$. In view of the above Theorem 22, ρ is invariant since it has the structure (38) with $\lambda = 0$ and $\sigma = |u\rangle\langle u|$.

Now, if

$$\rho = \lambda\rho_1 + (1 - \lambda)\rho_2 \quad (42)$$

with ρ_i , $i = 1, 2$ invariant states supported on $V \cap W_D^\perp$, let us say

$$\rho_i = \frac{\Gamma_{\text{Re},+,\omega_2}}{\Gamma_{\text{Re},+,\omega_2} + \Gamma_{\text{Re},-,\omega_2}}\sigma_i + \frac{\Gamma_{\text{Re},-,\omega_2}}{\Gamma_{\text{Re},+,\omega_2} + \Gamma_{\text{Re},-,\omega_2}}Z\sigma_i Z^*, \quad i = 1, 2$$

with σ_i supported on $V \cap \text{Im}|Z|$, $i = 1, 2$.

After multiplication by $|Z|$ in (42) we get,

$$|u\rangle\langle u| = \lambda\sigma_1 + (1 - \lambda)\sigma_2$$

Therefore, either $\lambda = 1$ and $\sigma_1 = |u\rangle\langle u|$ or $\lambda = 0$ and $\sigma_2 = |u\rangle\langle u|$, since $|u\rangle\langle u|$ is a pure state. This proves that ρ is extremal. Conversely, if ρ is an extremal invariant state supported on $V \cap W_D^\perp$, then it has the structure (38) with $\lambda = 0$. Hence, necessarily

$$\rho = \frac{\Gamma_{\text{Re},+, \omega_2}}{\Gamma_{\text{Re},+, \omega_2} + \Gamma_{\text{Re},-, \omega_2}}\sigma + \frac{\Gamma_{\text{Re},-, \omega_2}}{\Gamma_{\text{Re},+, \omega_2} + \Gamma_{\text{Re},-, \omega_2}}Z\sigma Z^* \quad (43)$$

with σ a state supported on $\text{Im}|Z| \cap V$, let us say $\sigma = \sum_j \sigma_j |u_j\rangle\langle u_j|$ with $(u_j)_j \subset \text{Im}|Z| \cap V$ the orthonormal basis of σ and $\sum_j \sigma_j = 1$. Consequently,

$$\rho = \sum_j \sigma_j \left(\frac{\Gamma_{\text{Re},+, \omega_2}}{\Gamma_{\text{Re},+, \omega_2} + \Gamma_{\text{Re},-, \omega_2}} |u_j\rangle\langle u_j| + \frac{\Gamma_{\text{Re},-, \omega_2}}{\Gamma_{\text{Re},+, \omega_2} + \Gamma_{\text{Re},-, \omega_2}} |Zu_j\rangle\langle Zu_j| \right) \quad (44)$$

But ρ is extremal, then there is only one non-trivial summand in (44). This finishes the proof. \square

Corollary 25. *All invariant states of modified AKV's model are detailed balance.*

Proof. Let ρ be an invariant state. Clearly the part of ρ supported on W_D satisfies a detailed balance condition, so that we can assume that ρ is supported on $V \cap W_D^\perp$. Recalling that $\{e_0, e_1, \psi, \psi'\} \subset \ker \rho$ if ρ is an invariant state supported on $V \cap W_D^\perp$ and $|Z|\psi = \psi$. By direct computations one can see that $\rho D_{\omega_1} = 0 = D_{\omega_1} \rho$, and $\rho D_{\omega_3} = 0 = D_{\omega_3} \rho$. Moreover,

$$\rho D_{\omega_2} = \frac{\Gamma_{\text{Re},-, \omega_2}}{\Gamma_{\text{Re},+, \omega_2}} Z \rho, \quad \text{and} \quad D_{\omega_2} \rho = Z \rho$$

hence, $\rho D_{\omega_2} = \frac{\Gamma_{\text{Re},-, \omega_2}}{\Gamma_{\text{Re},+, \omega_2}} D_{\omega_2} \rho$. In view of our Definition (9) we can conclude that ρ is a detailed balance state. \square

4.4 Fast recurrent subspace

The fast recurrent subspace of a quantum Markov semigroup is defined as

$$\mathcal{R} := \sup\{s(\rho) : s(\rho) \text{ is the support of an invariant state}\}$$

In this section we will characterize the fast recurrent subspace of the modified AKV's model.

Lemma 26. *With the same notations as above, we have that*

- (i) $P_{W_D} = P_0 + (P_2 - |Z|)$
- (ii) $P_{W_D^\perp} = P_1 + |Z| + P_3$
- (iii) $W_D \setminus \{e_0\} \subset V$
- (iv) $V \cap W_D^\perp = \text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp \oplus \text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$

Proof. To prove (i) it suffices to prove the identity $W_D = \{u_0e_0 + (P_2 - |Z|)v : v \in \mathfrak{h}, u_0 \in \mathbb{C}\}$. If $u = u_0e_0 + (P_2 - |Z|)v$ with $v \in \mathfrak{h}, u_0 \in \mathbb{C}$, clearly $u \perp e_1$ and $u \perp \psi'$ since $\psi' \in \text{Im}P_3$. After simple computations we get

$$\langle \psi, u \rangle = \langle \psi, (P_2 - |Z|)v \rangle = \langle (P_2 - |Z|)\psi, v \rangle = 0$$

since $P_2\psi = \psi = |Z|\psi$, hence $u \perp \psi$. Moreover, we have that $|Z|u = |Z|(P_2 - |Z|)v = 0$, then $u \in \ker |Z|$ and clearly $Z^*u = 0$. This proves that $\{u_0e_0 + (P_2 - |Z|)v : v \in \mathfrak{h}, u_0 \in \mathbb{C}\} \subset W_D$. To prove the oposite inequality observe that any $u \in \mathfrak{h}$ may be written in the form $u = P_0u + P_1u + P_2u + P_3u$, but if in addition $u \in W_D$, then $P_1u = 0$. Condition $u \in \ker |Z|$, $u \neq P_0u$ and identity $P_2 = \text{Im}|Z| \oplus \ker |Z|$, imply that $u = (P_2 - |Z|)v$ for some $v \in \mathfrak{h}$ and, hence, $P_3u = 0$. This proves (i).

Item (ii) is an immediate consequence of (i). If $u \in W_D \setminus \{e_0\}$, let us say $u = (P_2 - |Z|)v$, $v \in \mathfrak{h}$, then clearly $u \perp \{e_0, e_1\}$ and using that $P_2\psi = \psi = |Z|\psi$, we get

$$\langle \psi, u \rangle = \langle \psi, (P_2 - |Z|)v \rangle = \langle (P_2 - |Z|)\psi, u \rangle = 0$$

i.e., $u \perp \psi$. Similarly, identities $P_2Z^*\psi' = Z^*\psi' = |Z|Z^*\psi'$, imply that $u \perp Z^*\psi'$. Moreover, $u \in \{\psi', Z\psi\}^\perp$, since $\psi', Z\psi \in \text{Im}P_3$. This proves that $W_D \setminus \{e_0\} \subset V$.

Now, using (ii) we have that

$$\begin{aligned} V \cap W_D^\perp &= \{e_0, e_1, \psi, \psi', Z\psi, Z^*\psi'\}^\perp \cap (P_1 + |Z| + P_3)(\mathfrak{h}) \\ &= \{\psi, \psi', Z\psi, Z^*\psi'\}^\perp \cap (|Z| + P_3)(\mathfrak{h}) \\ &= \text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp \oplus \text{Im}P_3 \cap \{\psi', Z\psi\}^\perp \end{aligned} \tag{45}$$

This proves (iv) and finishes the proof of the lemma. \square

Corollary 27. *If $M < N - 1$, there exists invariant states of the modified AKV's model in $\text{Ann}(D) \setminus \{H\}'$.*

Proof. Let $u = u_0 e_0 + (P_2 - |Z|)v \in W_D$, with $u_0 \in \mathbb{C}$ and $v \in \mathfrak{h}$. Then we have,

$$P_0|u\rangle\langle u| - |u\rangle\langle u|P_0 = u_0|(P_2 - |Z|)v\rangle\langle e_0| - \bar{u}_0|e_0\rangle\langle(P_2 - |Z|)v| \quad (46)$$

Therefore, the invariant state $|u\rangle\langle u|$ does not belong to $\{H\}'$ if $u_0 \neq 0$, $v \neq 0$ and $|Z| < P_2$ (equivalently $M < N - 1$, see Proposition 2.2 in Ref.[16]). Nevertheless due to result of Theorem 6, the state belongs to the annihilator $\text{Ann}(D)$. \square

Remark 28. Notice that in contradistinction with Ref.[16], in the case when $|Z| < P_2$ (equivalently, $M < N - 1$) we have *found* invariant states outside $\{H\}'$, but in $\text{Ann}(D)$.

Proposition 29. (i) *The restrictions of the maps $Z : \text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp \mapsto \text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$ and $Z^* : \text{Im}P_3 \cap \{\psi', Z\psi\}^\perp \mapsto \text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$ are unitaries. Consequently $\dim(\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp) = \dim(\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp)$.*

(ii) *The state*

$$\rho = \frac{1}{\text{tr}(q)} \left(\frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}} q + \frac{\Gamma_{Re, -, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re, -, \omega_2}} ZqZ^* \right)$$

where $q = |Z|P_{\{\psi, Z^*\psi'\}^\perp}$ is the orthogonal projection on $\text{Im}|Z| \cap V$, is invariant for the modified AKV's model, with support projection $P_{W_D^\perp \cap V} = q + ZqZ^*$. Hence, ρ is faithful as an state in the hereditary subalgebra $P_{W_D^\perp \cap V} \mathcal{B}(\mathfrak{h}) P_{W_D^\perp \cap V}$.

Proof. Notice that Z maps $\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$ into $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$. Indeed, if $u \in \text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$ then $P_3 Zu = Zu$, $\langle \psi', Zu \rangle = \langle Z^*\psi', u \rangle = 0$ and $\langle Z\psi, Zu \rangle = \langle |Z|\psi, u \rangle = \langle \psi, u \rangle = 0$, hence $Zu \in \text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$. Moreover, $\langle Zu, Zv \rangle = \langle |Z|u, v \rangle = \langle u, v \rangle$, meaning that Z is unitary. Analogously one shows that Z^* maps $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$ into $\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$ and is unitary. This proves (i).

Let q be the orthogonal projection on the subspace $\text{Im}|Z| \cap V$. Notice that $q = |Z|$ on $\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$. Indeed, since ψ and $Z^*\psi'$ are fixed points of $|Z|$, this operator maps $\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$ into itself and condition $|Z|u = 0$, $u \in \text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$, implies that $u = |Z|u = 0$, then it is injective. Consequently, the map ZqZ^* maps $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$ into itself and it is injective since $ZqZ^*u = 0$ and $u \in \text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$, imply $0 = Z|Z|Z^*u = P_3u = u$. Moreover ZqZ^* is a projection since $ZqZ^*ZqZ^* = Zq|Z|qZ^* =$

ZqZ^* , and for any $u \in P_3(\mathbf{h}) = ZZ^*(\mathbf{h})$ we have that $ZqZ^*u = ZqZ^*ZZ^*u = Zq|Z|Z^*u = ZZ^*ZZ^*u = u$, hence, coincides with the orthogonal projection on $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$. Consequently, by item (iv) of the above Lemma 26 we conclude that $P_{W_D^\perp \cap V} = q + ZqZ^*$. The result of the proposition readily follows from Theorem 22. \square

Corollary 30. *The fast recurrent subspace of the modified AKV's model is*

$$\mathcal{R} = W_D \oplus W_D^\perp \cap V$$

Proof. By Theorem 22, every invariant state is supported on a subspace of \mathcal{R} . The same theorem and Proposition 29 imply that there exists an invariant state with support equal to \mathcal{R} . \square

4.5 Approach to equilibrium and attraction domains

The decoherence free sub-algebra $\mathcal{N}(\mathcal{T})$ of the QMS, \mathcal{T} , of the AKV's model was studied and characterized in Ref.[3] It was proved there that $\mathcal{N}(\mathcal{T}) \subset \mathcal{F}(\mathcal{T})$ where $\mathcal{F}(\mathcal{T})$ is the subset of fixed point of \mathcal{T} , consequently, $\mathcal{N}(\mathcal{T}) = \mathcal{F}(\mathcal{T})$ since the opposite inequality always holds true. If there exists a faithful invariant state in $\mathcal{B}(\mathbf{h})$, from a result of Frigerio and Verri[14, 15] one can conclude that for any normal state $\eta \in \mathcal{B}(\mathbf{h})$ there exists the limit $\tau = \lim_{t \rightarrow \infty} \mathcal{T}_{*t}(\eta)$. Unfortunately, as a consequence of Theorem 22, any invariant state τ for the AKV's model is singular, indeed $V^\perp \setminus \{e_0\} = \text{Span}\{e_1, \psi, \psi', Z\psi, Z^*\psi'\} \subset \ker(\tau)$. Then, to analyze the approach to equilibrium property and attraction domains, it is necessary to restrict our attention to the evolution on hereditary sub-algebras $\mathcal{A}_{W_D} = P_{W_D}\mathcal{B}(\mathbf{h})P_{W_D}$ and $\mathcal{A}_{W_D^\perp \cap V} = P_{W_D^\perp \cap V}\mathcal{B}(\mathbf{h})P_{W_D^\perp \cap V}$.

Lemma 31. *Let $\mathcal{T}_{W_D^\perp \cap V, t} = P_{W_D^\perp \cap V}\mathcal{T}_tP_{W_D^\perp \cap V}$ be the hereditary semigroup acting on the subalgebra $\mathcal{A}_{W_D^\perp \cap V}$. Then $\mathcal{N}(\mathcal{T}_{W_D^\perp \cap V}) \subset \mathcal{F}(\mathcal{T}_{W_D^\perp \cap V})$.*

Proof. Let a be a selfadjoint element $a \in \mathcal{A}_{W_D^\perp \cap V}$, then a is supported on a subspace of V , meaning that $\{e_1, \psi, \psi', Z\psi, Z^*\psi'\} \subset \ker a$. Also $e_0 \in \ker a$ since a is supported on W_D^\perp . Then a commutes with P_0 and P_1 . If moreover $a \in \mathcal{N}(\mathcal{T}_{W_D^\perp \cap V})$ then $a \in \mathcal{C}'_0$, since by Theorem 3.2 in Ref.[10] $\mathcal{N}(\mathcal{T}_{W_D^\perp \cap V}) \subset \mathcal{C}'_0$, where

$$\begin{aligned} \mathcal{C}_0 &= \{D_{\omega_\ell}, D_{\omega_\ell}^* : 1 \leq \ell \leq 3\} \\ &= \{|\psi\rangle\langle e_1|, |e_1\rangle\langle\psi|, Z, Z^*, |e_0\rangle\langle\psi'|, |\psi'\rangle\langle e_0|\} \end{aligned} \tag{47}$$

consists of the Kraus operators of the model along with their adjoints, restricted to $\mathcal{A}_{W_D^\perp \cap V}$. In particular a commutes with Z and Z^* , hence $a(\ker |Z|) \subset$

$\ker |Z|$ and $a(\text{Im}|Z|) \subset \text{Im}|Z|$. Then a commutes with $P_2 = \text{Im}|Z| \oplus \ker |Z|$, an consequently, it commutes also with $P_3 = I - P_0 - P_1 - P_2$. In conclusion a commutes with H .

Now, condition $\{e_0, e_1, \psi, \psi', Z\psi, Z^*\psi'\} \subset \ker a$ implies that $a \in \ker \mathcal{L}_{\omega_1} \cap \ker \mathcal{L}_{\omega_3}$. Moreover since a commutes with P_3 and with $|Z|$, $[H_{\omega_2}, a] = 0$. Commutation of a with Z, Z^* implies that $a \in \ker \mathcal{L}_{\omega_2}$. Consequently a is a fixed point of $\mathcal{T}_{W_D^\perp \cap V}$. Since any operator is a linear combination of four selfadjoint elements, the same result holds true for any element $a \in \mathcal{A}_{W_D^\perp \cap V}$. This finishes the proof. \square

Corollary 32. *For any initial state $\eta \in \mathcal{A}_{W_D^\perp \cap V}$ there exists an invariant state $\eta_\infty = \lim_{t \rightarrow \infty} \mathcal{T}_{W_D^\perp \cap V, *t}(\eta) \in \mathcal{A}_{W_D^\perp \cap V}$.*

Proof. This is a consequence of Lemma 31, Proposition 29 and the previously cited result of Frigerio and Verri[14, 15]. \square

Now we shall give a full description of the evolution of any state supported on $W_D^\perp \cap V$. To do so, we start by describing explicitly the evolution of states supported either on $\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$ or on $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$.

Lemma 33. *(i) Let σ be a state supported on $\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$. Then $\mathcal{T}_{*t}(\sigma)$ converges to*

$$\frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} \sigma + \frac{\Gamma_{Re,-, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} Z\sigma Z^*$$

as $t \rightarrow \infty$.

*(ii) Let η be a state supported on $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$. Then $\mathcal{T}_{*t}(\eta)$ converges to*

$$\frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} Z^*\eta Z + \frac{\Gamma_{Re,-, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} \eta$$

as $t \rightarrow \infty$

Proof. For any observable $x \in \mathcal{B}(\mathfrak{h})$ define $x_t = \text{tr}(x\mathcal{T}_{*t}(\sigma))$ and $y_t = \text{tr}(x\mathcal{T}_{*t}(Z\sigma Z^*))$. Then after derivation with respect to t we get that

$$\begin{aligned} \dot{x}_t &= -ax_t + ay_t \\ \dot{y}_t &= bx_t - by_t \end{aligned} \tag{48}$$

with $a = 2(N-1)\Gamma_{Re,-, \omega_2}$ and $b = 2(N-1)\Gamma_{Re,+, \omega_2}$. Therefore (x_t, y_t) is a classical random walk with Q -matrix

$$Q = \begin{pmatrix} -a & a \\ b & -b \end{pmatrix} \tag{49}$$

and stationary probability distribution $\pi = (\frac{b}{a+b}, \frac{a}{a+b})$. Indeed, after solving system (48) one gets,

$$x_t = \text{tr} \left(x \left(\frac{b + ae^{-t(a+b)}}{a+b} \sigma + \frac{a - ae^{-t(a+b)}}{a+b} Z \sigma Z^* \right) \right)$$

which implies that

$$\mathcal{T}_{*t}(\sigma) = \frac{b + ae^{-t(a+b)}}{a+b} \sigma + \frac{a - ae^{-t(a+b)}}{a+b} Z \sigma Z^* \mapsto \frac{b}{a+b} \sigma + \frac{a}{a+b} Z \sigma Z^*$$

as $t \rightarrow \infty$. This proves (i).

To proof (ii) take $\sigma = Z^* \eta Z$ and apply (i). This finishes the proof. \square

Theorem 34. *Let η be an initial state supported on $W_D^\perp \cap V$. Then*

$$\lim_{t \rightarrow \infty} \mathcal{T}_{*t}(\eta) = \frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} \sigma + \frac{\Gamma_{Re,-, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} Z \sigma Z^* \quad (50)$$

with $\sigma = \frac{1}{\text{tr}(q\eta)} q\eta q + \frac{1}{\text{tr}(P_3\eta)} Z^* \eta Z$.

Proof. By Corollary 32 there exists an invariant state

$$\eta_\infty = \frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} \sigma + \frac{\Gamma_{Re,-, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} Z \sigma Z^* \quad (51)$$

where σ is a state supported on $\text{Im}|Z| \cap \{\psi, Z^* \psi'\}^\perp$, such that $\lim_{t \rightarrow \infty} \mathcal{T}_{*t}(\eta) = \eta_\infty$.

Every initial state η supported on $W_D^\perp \cap V$ may be written in the form

$$\eta = (q + ZqZ^*)\eta(q + ZqZ^*) = q\eta q + ZqZ^* \eta ZqZ^* + q\eta ZqZ^* + ZqZ^* \eta q \quad (52)$$

The state $\frac{1}{\text{tr}(q\eta)} q\eta q$ is supported on $\text{Im}|Z| \cap \{\psi, Z^* \psi'\}^\perp$ and due to the above Lemma 33, it is driven by the semigroup to the limit state

$$\frac{1}{\text{tr}(q\eta)} \left(\frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} q\eta q + \frac{\Gamma_{Re,-, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} Zq\eta q Z^* \right) \quad (53)$$

as $t \rightarrow \infty$. Analogously, due to the same Lemma 33, the state $\frac{1}{\text{tr}(P_3\eta)} ZqZ^* \eta ZqZ^* = \frac{1}{\text{tr}(P_3\eta)} P_3 \eta P_3$ is supported on $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$ and

$$\begin{aligned} \lim_{t \rightarrow \infty} \mathcal{T}_{*t} \left(\frac{1}{\text{tr}(P_3\eta)} P_3 \eta P_3 \right) &= \frac{1}{\text{tr}(P_3\eta)} \left(\frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} Z^* P_3 \eta P_3 Z \right. \\ &\quad \left. + \frac{\Gamma_{Re,-, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} P_3 \eta P_3 \right) \end{aligned} \quad (54)$$

Comparing (51) with (53) and (54), we conclude that

$$\lim_{t \rightarrow \infty} \mathcal{T}_{*t}(q\eta ZqZ^* + ZqZ^*\eta q) = 0,$$

$$\sigma = \frac{1}{\text{tr}(q\eta)} q\eta q + \frac{1}{\text{tr}(P_3\eta)} Z^* P_3 \eta P_3 Z$$

and consequently,

$$Z\sigma Z^* = \frac{1}{\text{tr}(q\eta)} Zq\eta qZ^* + \frac{1}{\text{tr}(P_3\eta)} P_3\eta P_3$$

This finishes the proof. \square

Corollary 35. *The attraction domain of the invariant state*

$$\rho = \frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} \sigma + \frac{\Gamma_{Re,-, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}} Z\sigma Z^*$$

where σ is a state supported on $\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$, consists of all those initial states η such that

$$\sigma = \frac{1}{\text{tr}(q\eta)} q\eta q + \frac{1}{\text{tr}(P_3\eta)} Z^* P_3 \eta P_3 Z$$

Proof. Immediate from the above Theorem. \square

Remark 36. (Transport of states) Due to the result of Theorem 32 the total probability of an initial state η is redistributed in the limit as $t \rightarrow \infty$; so that the probability in the portion of the final state η_∞ supported on $\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$ is

$$\text{tr}(q\eta_\infty) = \frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}}$$

and the probability in the portion of the final state η_∞ supported on $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$ is

$$\text{tr}(P_3\eta_\infty) = \frac{\Gamma_{Re,-, \omega_2}}{\Gamma_{Re,+, \omega_2} + \Gamma_{Re,-, \omega_2}}$$

By adjusting the values of the Γ 's, so that $\frac{\Gamma_{Re,+, \omega_2}}{\Gamma_{Re,-, \omega_2}} = e^{-\beta(\omega_2)\omega_2} \rightarrow 0$ (equivalently $\beta(\omega_2) \rightarrow \infty$), one can “transport” any initial state η to a limit state η_∞ concentrated (but non-necessarily supported) on the subspace $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$. In particular any initial state supported on $\text{Im}|Z| \cap \{\psi, Z^*\psi'\}^\perp$ can be transported, as $t \rightarrow \infty$, to a limit state concentrated on the subspace $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$. In this sense our modified AKV's model allows us to “transport” probability mass from any initial state to a state concentrated on $\text{Im}P_3 \cap \{\psi', Z\psi\}^\perp$.

Remark 37. (Energy gain) A simple computation shows that if at $t = 0$ the system is in any initial state ρ supported on the subspace $\text{Im}|Z| \cap \{\psi, Z^*\psi\}^\perp$, then as $t \rightarrow \infty$ the energy of the system increases and the *energy gain* during the process is given by

$$\text{tr}(\rho_\infty H_{eff}) - \text{tr}(\rho H_{eff}) = (N - 1) \frac{\Gamma_{\text{Re}, -, \omega_2}}{\Gamma_{\text{Re}, +, \omega_2} + \Gamma_{\text{Re}, -, \omega_2}} (\Gamma_{\text{Im}, +, \omega_2} + \Gamma_{\text{Im}, -, \omega_2})$$

which is independent of ρ . This is an indication that degenerate open systems (with a degenerate reference Hamiltonian) seems to be appropriate for modeling effective quantum energy transfer in photosynthesis, see Ref.[20] and the reference therein. We will continue the analysis of energy transfer when the initial state of the system is arbitrary in forthcoming work.

Acknowledgment

The authors are grateful to Franco Fagnola who suggested to work in this problem and gave valuable insight. The financial support from CONACYT-Mexico (Grant 221873) and PRODEP Red de Analisis Italia-UAM, is gratefully acknowledged

References

- [1] Accardi L., Fagnola F. and Quezada R., *On three new principles in non-equilibrium statistical mechanics and Markov semigroups of weak coupling limit type*, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **19** (2) (2016) 1650009 (37 pages), DOI: 10.1142/S0219025716500090.
- [2] J. Agredo, F. Fagnola and R. Rebolledo, *Decoherence-free subspaces of a quantum Markov semigroup*, *J. Math. Phys.* **55**, (2014), 11–32.
- [3] Agredo J., García J.C. and Quezada R., *Decoherence in an excitation energy transport model*, Submitted to Proceedings of the 38th International Conference on Quantum Probability, Tokyo University of Science, 2019.
- [4] Aref'eva, Y., Volovich, I. and Kozyrev, S. 2015. *Stochastic Limit Method and Interference in Quantum Many-particles Systems*. Theoretical and Mathematical Physics **183**(3) (2015) 782–799.

- [5] Bolaños-Servín J.R. and Fagnola F, *On the structure of quantum Markov semigroups of weak coupling limit type*, IOP Conf. Series: Journal of Physics: Conf. Series 819 (2017) 012003, doi:10.1088/1742-6596/819/1/012003
- [6] Ph. Blanchard, and R. Olkiewicz, *Decoherence induced transition from quantum to classical dynamics*, *Rev. Math. Phys.*, **15**, (2003), 217–243 .
- [7] Ph. Blanchard, and R. Olkiewicz, *Decoherence as irreversible dynamical process in open quantum systems*, *Open quantum systems. III, Lectures Notes in Math.*, 1882, Springer, Berlin, (2006).
- [8] R. Carbone, E. Sasso, and V. Umanità, *Decoherence for Quantum Markov Semigroups on Matrix Algebras*, *Annales Henri Poincaré*, **14**, (2013), 17–37.
- [9] Cruz de la Rosa M.A. and Quezada R., *Non-equilibrium steady states of a Markov generator of weak coupling limit type modeling absorption-emission of m and n photons*, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **19** (4) (2016) 1650023 (17 pages), DOI: 10.1142/S0219025716500235
- [10] A. Dhahri, F. Fagnola and R. Rebolledo, *The decoherence-free subalgebra of a quantum Markov semigroup with unbounded generator*, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, **13**, (2010), 413–433.
- [11] Kozyrev S. and Volovich I., *Dark states in quantum photosynthesis*, arXiv:1603.07182v1 [physics.bio-ph] (2016).
- [12] Fagnola F. and Umanità V., *Generators of detailed balance quantum Markov semigroups*, *Infinite Dimensional Analysis, Quantum Probab. Relat. Top.* **10** (3) (2007) 335–363.
- [13] Fagnola F. and Rebolledo R, *Subharmonic projections for a quantum Markov semi- group*, *J. Math. Phys.* **43** (2002) 1074–1082.
- [14] A.Frigerio, *Quantum dynamical semigroups and approach to equilibrium*, *Lett. Math. Phys.* 2 (1977) 79–87.
- [15] A. Frigerio and M. Verri, *Long-time asymptotic properties of dynamical semigroups on W^* -algebras*, *Math. Z.* 180 (1982) 275–286.
- [16] García J.C., Gliouez S., Guerrero F. and Quezada R., *Entangled and dark stationary states of excitation energy transport models in quantum*

many-particles systems and photosynthesis, Infinite Dimensional Analysis, Quantum Probab. Relat. Top. **21** (3) (2018) 1850018 (21 pages), DOI: 10.1142/S0219025718500182

- [17] García J.C. and Guerrero F., *Some non-equilibrium invariant states for the asymmetric exclusion QMS at level one*, Infinite Dimensional Analysis, Quantum Probab. Relat. Top. **18** (01) (2015) 550007, doi.org/10.1142/S0219025715500071
- [18] I.L. Chuang, D.A. Lidar and K.B. Whaley, *Decoherence-free Subspaces for Quantum Computation*, Phys. Rev. Lett, **81**, (1998), 2594.
- [19] D. Lidar, *Review of Decoherence Free Subspaces, Noiseless Subsystems, and Dynamical Decoupling*, Adv. Chem. Phys. **154**, (2014), 295–354.
- [20] G. D. Scholes, G. R. Fleming, A. Olaya-Castro, and R. van Grondelle, *Lessons from nature about solar light harvesting*, Nature Chem. **3**, (2011), 763–774. DOI: 10.1038/NCHEM.1145