

CLASSIFICATION OF TORIC PROJECTIVE VARIETIES UP TO PROJECTIVE AUTOMORPHISMS

FRIEDRICH KNOP AND RAINER SINN

ABSTRACT. Toric subvarieties of projective space are classified up to projective automorphisms.

1. THE RESULT

A closed subvariety $X \subseteq \mathbb{P}^N = \mathbb{P}^N(\mathbb{C})$ will be called *toric* if it is the orbit closure of a subtorus $T \subseteq \mathrm{PGL}(N+1)$. The aim of this note is to classify toric subvarieties of \mathbb{P}^N as abstract subvarieties, i.e., without T being part of the structure. More precisely, we give a description of all toric subvarieties up to projective equivalence where two subvarieties $X, X' \subseteq \mathbb{P}^N$ will be called *projectively equivalent* if there is $g \in \mathrm{PGL}(N+1)$ (possibly not commuting with T) such that $g(X) = X'$.

Toric subvarieties can be constructed as follows: For any $d \in \mathbb{N}$, we choose some non-empty subset $S \subseteq \mathbb{Z}^d$ with $|S| \leq N+1$. Let $S = \{m_0, \dots, m_r\}$ (where $r \leq N$, by assumption). Then $X(S) = \overline{Tx}$ is a toric subvariety of \mathbb{P}^N where T is the image of

$$(1) \quad (\mathbb{C}^*)^d \rightarrow \mathrm{PGL}(N+1) : t \mapsto \mathrm{diag}(t^{m_0}, \dots, t^{m_r}, 1, \dots, 1)$$

(with $N-r$ ones) and $x := [1 : \dots : 1 : 0 : \dots : 0]$ (with $N-r$ zeros). Observe that $X(S)$ depends on the choice of an enumeration of S but clearly any two choices lead to projective equivalent subvarieties.

We call a subset $S \subseteq \mathbb{Z}^d$ *affinely generating* if \mathbb{Z}^d is generated as a group by the differences $m-m'$ with $m, m' \in S$. Two subsets $S, S' \subseteq \mathbb{Z}^d$ are *affinely equivalent* if $S' = \Phi(S)$ where Φ is an affine transformation of the form $\Phi(m) = Am + b$ with $A \in \mathrm{GL}(d, \mathbb{Z})$ and $b \in \mathbb{Z}^d$.

Our main result is the following:

Theorem 1. For any $d, N \in \mathbb{N}$ with $d \leq N$ the map $S \mapsto X(S)$ induces a bijection between

- affinely generating subsets $S \subseteq \mathbb{Z}^d$ with $|S| \leq N+1$ up to affine equivalence and
- toric subvarieties $X \subseteq \mathbb{P}^N$ of dimension d up to projective equivalence.

Our main application is to toric varieties attached to lattice polytopes. Recall that a *lattice polytope* P is the convex hull inside \mathbb{R}^d of a

finite subset of \mathbb{Z}^d . Thus, $S_P := P \cap \mathbb{Z}^d$ is finite and $P = \text{conv } S_P$. Lattice polytopes are in one-to-one correspondence with pairs of complete (abstract) toric varieties together with a complete and ample linear series. We call a lattice polytope \mathbb{Z} -solid if S_P affinely generates \mathbb{Z}^d . Clearly every \mathbb{Z} -solid lattice polytope is solid, i.e., $\dim P = d$. The converse is not true. Counterexamples are provided at the end of the paper (Example 2.6). Every lattice polytope with the integer decomposition property is \mathbb{Z} -solid and in particular, every lattice polygon is \mathbb{Z} -solid.

For a lattice polytope P , put $X(P) := X(S_P)$. Then applying Theorem 1 to $X(P)$ we get:

Corollary 1.1. *Let $P, P' \subseteq \mathbb{R}^d$ be two \mathbb{Z} -solid lattice polytopes with $|S_P|, |S_{P'}| \leq N + 1$. Then $X(P)$ and $X(P')$ are projectively equivalent if and only if P and P' are affinely equivalent. In particular, it follows that $X(P) = X(P')$.*

We cannot simply drop the assumption that P and P' are \mathbb{Z} -solid.

Example 1.2. Consider the two full-dimensional tetrahedra

$$P = \text{conv}\{0, e_1, e_2, e_3\} \text{ and } P' = \text{conv}\{0, e_1, e_2, e_1 + e_2 + 2e_3\}$$

in \mathbb{R}^3 . Clearly, the lattice points in P generate the $\mathbb{Z}^3 \subset \mathbb{R}^3$ as a group. On the other hand, the lattice polytope P' is not \mathbb{Z} -solid. The toric varieties corresponding to these two polytopes with 4 lattice points is \mathbb{P}^3 in both cases. In particular, they are projectively equivalent, whereas the lattice polytopes are not affinely equivalent.

One can also wonder when two toric varieties of the form $X(S)$ are projectively equivalent if S is not necessarily affinely generating. In this case, let $M(S) \subseteq \mathbb{Z}^d$ be the subgroup generated by all $m - m'$, $m, m' \in S$. Choose any isomorphism $\Phi : M(S) \rightarrow \mathbb{Z}^e$ and let $\tilde{S} := \Phi(S - m)$ where m is any element of S . Then \tilde{S} affinely generates \mathbb{Z}^e .

Corollary 1.3. *Let $S \subseteq \mathbb{Z}^d$ be a subset with $|S| \leq N + 1$. Then $X(S)$ is projectively equivalent to $X(\tilde{S})$.*

Remark. *In this paper, we work for convenience over \mathbb{C} only. It is not difficult to see, that all results hold over an arbitrary algebraically closed ground field. In positive characteristic, only the proof of Lemma 2.2 requires a slight modification.*

2. THE PROOF

As a general reference for toric varieties from the algebraic geometry point of view, we refer to [1]. The basic facts from representation theory that we use throughout the paper can be found in [2].

The proof of Theorem 1 proceeds in several steps. First we observe that the map $S \mapsto X(S)$ is well defined:

Lemma 2.1. *If $S, S' \subseteq \mathbb{Z}^d$ are affinely equivalent then $X(S), X(S') \subseteq \mathbb{P}^N$ are equal, so in particular projectively equivalent.*

Proof. We show that any affine linear isomorphism $m \mapsto Am + b$ of \mathbb{Z}^d does not change $Tx \subset \mathbb{P}^N$. Indeed, A results simply in a reparametrization of $(\mathbb{C}^*)^d$ while b corresponds to a multiplication by a non-zero constant factor which drops out by homogeneity. \square

Next we show that dimensions are preserved:

Lemma 2.2. *Let $S \subseteq \mathbb{Z}^d$ be affinely generating. Then $\chi : (\mathbb{C}^*)^d \rightarrow X(S) : t \mapsto tx$ is an open embedding. Hence $\dim X(S) = d$.*

Proof. The map χ is dominant by definition. Hence the image is open. It suffices to show that $\chi^{-1}(x) = 1$. We have

$$\begin{aligned} \chi^{-1}(x) &= \{t \in (\mathbb{C}^*)^d \mid t^{m_0} = \dots = t^{m_r}\} = \\ &= \{t \in (\mathbb{C}^*)^d \mid t^{m_1 - m_0} = \dots = t^{m_r - m_0} = 1\}. \end{aligned}$$

The latter group is trivial since \mathbb{Z}^d is affinely generated by S , so that these characters separate the torus. \square

To show surjectivity, we first investigate arbitrary homomorphisms $\varrho : T \rightarrow \mathrm{PGL}(N+1)$, where T is a torus. The image is contained in a maximal torus of $\mathrm{PGL}(N+1)$, which is conjugate to the maximal torus $\{\mathrm{diag}(1, x) : x \in (\mathbb{C}^*)^N\} \subset \mathrm{PGL}(N+1)$. This torus can clearly be lifted to $\mathrm{GL}(N+1)$ and hence also ϱ can be lifted to a homomorphism $\tilde{\varrho} : T \rightarrow \mathrm{GL}(N+1)$. Any two lifts differ by a homomorphism of T into the scalars. Via this lift $\tilde{\varrho}$, the torus T acts on \mathbb{C}^n , which decomposes into simultaneous eigenspaces and the eigenvalues corresponding to an eigenspace are given by the corresponding eigencharacter on T . Thus the set of eigencharacters $E(\varrho; \mathbb{P}^N) \subseteq M_T$ is well defined up to translation by an element of the lattice of characters M_T .

Lemma 2.3. *For every toric subvariety $X \subseteq \mathbb{P}^N$ of dimension d there is a affinely generating subset $S \subseteq \mathbb{Z}^d$ such that X is projectively equivalent to $X(S)$.*

Proof. Let $X = \overline{Tx}$ with $T \subset \mathrm{PGL}(N+1)$ a torus. Lift the inclusion ϱ to a homomorphism $\tilde{\varrho} : T \rightarrow \mathrm{GL}(N+1)$. Let $E := E(\varrho; \mathbb{P}^N)$ be the set of its eigencharacters and let $\mathbb{C}^{N+1} = \bigoplus_{\chi \in E} V_\chi$ be the corresponding decomposition in eigenspaces. Lift also x to a vector $v \in \mathbb{C}^{N+1}$ and decompose it: $v = \sum_{\chi \in E} v_\chi$. Put $S_0 := \{\chi \in E \mid v_\chi \neq 0\}$ and choose an eigenbasis whose first $r+1$ members are the v_χ with $\chi \in S_0$. After a projective transformation we may assume that this eigenbasis is the standard basis. If $S_0 = \{\chi_0, \dots, \chi_r\}$ we see that then

$$Tx = \{[\chi_0(t) : \dots : \chi_r(t) : 0 : \dots : 0] \in \mathbb{P}^N \mid t \in T\}$$

Let $M \subseteq M_T$ be the subgroup generated by all ratios $\chi_i \chi_j^{-1}$ and let $\chi'_i := \chi_i \chi_0^{-1}$. Then M is affinely generated by $S := \{\chi'_i \mid 0 \leq i \leq j\}$.

Fix an isomorphism $M \cong \mathbb{Z}^e$ for some $e \in \mathbb{N}$ such that S becomes an affinely generating subset of \mathbb{Z}^e . By construction, we have $X = \overline{Tx} = X(S)$. Lemma 2.2 then implies $e = d$. \square

Next we prove the key lemma. For any subvariety $X \subseteq \mathbb{P}^N$ let $G(X) \subseteq \text{Aut } X$ be the group of all automorphisms which can be extended to an automorphism of \mathbb{P}^N . If $X = \overline{Tx}$ is now toric, then restriction to X yields a homomorphism $\varrho : T \rightarrow G(X)$.

Lemma 2.4. $T_0 := \varrho(T)$ is a maximal torus of $G(\overline{Tx})$.

Proof. Set $X = \overline{Tx}$. Choose a maximal torus $T_1 \subseteq G(X)$ containing T_0 . We are going to show that every $t_1 \in T_1$ lies in T_0 . Since t_1 normalizes T_0 , it permutes the T_0 -orbits of X . The open T_0 -orbit Tx is unique, hence it is preserved by t_1 . Thus, there is $t_0 \in T_0$ with $t_1x = t_0x$, implying that $t' := t_1t_0^{-1}$ fixes x . Since t' centralizes T_0 , it will fix every point of Tx . Thus it acts trivially on the closure X . This shows $t' = 1$ and therefore $t_1 = t_0 \in T_0$. \square

The following Lemma finishes the proof of Theorem 1.

Lemma 2.5. Let $S, S' \subseteq \mathbb{Z}^d$ be affinely generating and assume that $X := X(S)$, $X' := X(S')$ are projectively equivalent. Then S, S' are affinely equivalent.

Proof. Let $\langle X \rangle \subseteq \mathbb{P}^N$ be the smallest linear subspace containing X . Then the linear independence of characters implies that $\langle X \rangle = \mathbb{P}^r \subseteq \mathbb{P}^N$ where $r + 1 = |S|$.

By assumption there is $g \in \text{PGL}(N+1)$ with $g(X) = X'$. Since then $g\langle X \rangle = \langle X' \rangle$ we deduce that $\dim \langle X \rangle = r = \dim \langle X' \rangle$ and that g is an automorphism of \mathbb{P}^r . Thus we may replace \mathbb{P}^N by \mathbb{P}^r and hence assume that $r = N$. Since, in this case, the only element of $\text{PGL}(N+1)$ which acts as identity on X is the identity, we see that $G(X)$ (and likewise $G(X')$) is a subgroup of $\text{PGL}(N+1)$.

Let T_0 and T'_0 be the image of $T = (\mathbb{C}^*)^d$ in $\text{PGL}(N+1)$ corresponding to S and S' , respectively, via the map defined in (1). Then, by Lemma 2.4, both T_0 and $T_1 := g^{-1}T'_0g$ are maximal tori of $G(X)$. It follows that there is $h \in G(X)$ with $h^{-1}T_1h = T_0$. Replacing g by gh we may assume additionally to $g(X) = X'$ that $T_0 = T_1$.

Lemma 2.2 also implies that $T \rightarrow T_0$ and $T \rightarrow T'_0$ are isomorphisms. Thus we get $\phi \in \text{Aut } T$ such that the following diagram commutes:

$$\begin{array}{ccc} T & \xrightarrow{\varrho_S} & T_0 \\ \phi \downarrow & & \downarrow \text{Ad}(g) \\ T & \xrightarrow{\varrho_{S'}} & T'_0 \end{array}$$

Write Φ for the automorphism of \mathbb{Z}^d corresponding to ϕ , so that using $\varrho_S \circ \phi^{-1} = \varrho_{\Phi(S)}$ we get

$$\varrho_{S'}(t) = g\varrho_{\Phi(S)}g^{-1}$$

This means that the sets of eigencharacters S' and $\Phi(S)$ are equal up to a translation, i.e., there is $b \in \mathbb{Z}^d$ such that $S' = \Phi(S) + b$. \square

We conclude the paper with an example of a solid but not \mathbb{Z} -solid lattice polytope.

Example 2.6. Consider the lattice $M = \mathbb{Z}^d + \mathbb{Z}\delta \subseteq \mathbb{R}^d$ where $\delta = (\frac{1}{2}, \dots, \frac{1}{2})$. Then the simplex $P := \text{conv}\{0, e_1, \dots, e_d\}$ is a lattice polytope. For $d \geq 3$ we have $\delta \notin P$. Hence P is not \mathbb{Z} -solid for $d \geq 3$. For $d \leq 2$ one can show that all solid lattice polytopes are \mathbb{Z} -solid.

REFERENCES

- [1] D. A. Cox, J. B. Little, and H. K. Schenck. *Toric varieties*, volume 124 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.
- [2] J. E. Humphreys. *Linear algebraic groups*. Springer-Verlag, New York-Heidelberg, 1975. Graduate Texts in Mathematics, No. 21.

FRIEDRICH KNOP: DEPARTMENT MATHEMATIK, FAU ERLANGEN-NÜRNBERG,
CAUERSTRASSE 11, 91058 ERLANGEN, GERMANY
E-mail address: `friedrich.knop@fau.de`

RAINER SINN: FACHBEREICH MATHEMATIK UND INFORMATIK, FU BERLIN,
ARNIMALLEE 2, 14195 BERLIN, GERMANY
E-mail address: `rsinn@zedat.fu-berlin.de`