ON THE PRESERVERS OF MAXIMALLY ENTANGLED STATES

BEN GROSSMANN AND HUGO J. WOERDEMAN

ABSTRACT. We characterize the linear maps that preserve maximally entangled states in $L(\mathcal{X} \otimes \mathcal{Y})$ in the case where $\dim(X)$ divides $\dim(Y)$.

1. INTRODUCTION

Entanglement is considered a valuable resource in quantum information theory; it is responsible for the power of quantum computing and for applications such as superdense coding and quantum teleportation. In the interest of "conserving" the resource of entanglement, we might ask the following question: which linear maps produce a maximally entangled output for any maximally entangled input?

More concretely: a quantum system with m mutually exclusive configurations can be represented by an m-dimensional inner product space, which we will call \mathcal{X} . The configuration of such a system is represented with a unit vector $u \in \mathcal{X}$. If \mathcal{X}, \mathcal{Y} are the spaces corresponding to two such systems, then a configuration of the joint system can be represented with a unit vector $u \in \mathcal{X} \otimes \mathcal{Y}$, where $\mathcal{X} \otimes \mathcal{Y}$ is the tensor product of the Hilbert spaces associated with each system. We could also represent this system with the associated "pure state" density operator $A = uu^*$. We can measure the "entropy of entanglement" of A by $S(\operatorname{tr}_{\mathcal{Y}}(A))$, where $\operatorname{tr}_{\mathcal{Y}}$ denotes a partial trace over the space \mathcal{Y} and $S(\rho) = -\operatorname{tr}(\rho \log_2 \rho)$ is the Von Neumann entropy. We refer to the set of pure state density operators that maximize this measure of entanglement "maximally entangled states," and denote the set as $\operatorname{MES}_{\mathcal{X},\mathcal{Y}}$. This set can be succinctly described as

$$\operatorname{MES}_{\mathcal{X},\mathcal{Y}} = \left\{ \frac{1}{m} \operatorname{vec}(A) \operatorname{vec}(A)^* \mid A : \mathcal{Y} \to \mathcal{X} \text{ is a coisometry} \right\},\$$

where vec stacks the rows of a matrix A (see Section 2 for details). Quantum information is generally transmitted via a "quantum channel". Mathematically, this channel is a linear map that takes density matrices over a first system to density matrices over a second system (with an additional "complete positivity" condition). With this in mind, we hope to classify the linear maps Φ : span(MES_{\mathcal{X},\mathcal{Y}}) \rightarrow span(MES_{\mathcal{X},\mathcal{Y}}) that preserve the property of maximal entanglement, which is to say that they satisfy $\Phi(MES_{\mathcal{X},\mathcal{Y}}) \subset MES_{\mathcal{X},\mathcal{Y}}$.

The problem of classifying the linear maps that "preserve" the subset $MES_{\mathcal{X},\mathcal{Y}}$ is an example of a "linear preserver problem". Linear preserver problems like this one have a long history (see, for instance, [3] and [1, Chapter 7]). Recently, there has been work done on linear preservers within the context of quantum information

¹⁹⁹¹ Mathematics Subject Classification. 15A86, 15B57, 46N50.

Key words and phrases. Linear preservers, Maximally entangled states, Projective space.

HJW is supported by Simons Foundation grant 355645.

theory. Two relevant papers to this endeavor are [2] and the more recent paper [5], whose results we generalize here.

In this paper, we prove the following result.

Theorem 1.1. Let $\mathcal{Y} = \mathcal{X}^k := \mathcal{X} \oplus \cdots \oplus \mathcal{X}$, k > 1. If $\Phi : \operatorname{span}(\operatorname{MES}_{\mathcal{X},\mathcal{Y}}) \to \operatorname{span}(\operatorname{MES}_{\mathcal{X},\mathcal{Y}})$ is an invertible linear map that preserves $\operatorname{MES}_{\mathcal{X},\mathcal{Y}}$, then Φ is of the form

$$\Phi: X \mapsto (U \otimes V) X^{\sigma} (U \otimes V)^*$$

where $U \in L(\mathcal{X}), V \in L(\mathcal{Y})$ are unitary if $A \mapsto A^{\sigma}$ denotes either the identity or transpose map.

This generalizes (the invertible case of) Poon's result from [5], which can be stated within our framework as follows:

Theorem 1.2. [5] A linear map Φ : span(MES_{X,X}) \rightarrow span(MES_{X,X}) preserves MES_{X,X} if and only if it has one of the following forms:

(1) $\Phi(A \otimes B) = (U \otimes V)(A \otimes B)^{\sigma}(U \otimes V)^*$

- (2) $\Phi(A \otimes B) = (U \otimes V)(B \otimes A)^{\sigma}(U \otimes V)^*$
- (3) $\Phi(X) = (\operatorname{tr} X)\rho$ for some $\rho \in \operatorname{MES}$

where $A \mapsto A^{\sigma}$ denotes either the identity or transpose map.

Notice that (2) does not appear in Theorem 1.1. Indeed, for (2) to make sense one needs $\mathcal{X} = \mathcal{Y}$. In addition, Poon was able to describe the non-invertible preservers, which take the form (3).

The proof can be summarized as follows. First, we show that an MES-preserving map Φ induces a continuous map ζ over the projective space $P(\operatorname{coisom}(\mathcal{Y}, \mathcal{X}))$ (to be introduced in Section 4). Moreover, we show that for mutually orthogonal coisometries A, B, there exist mutually orthogonal coisometries C, D such that we have either

$$\zeta([\alpha A + \beta B]) = [\alpha C + \beta D]$$
 or $\zeta([\alpha A + \beta B]) = [\bar{\alpha}C + \bar{\beta}D].$

By considering the degree of ζ restricted to the span of [A] and [B], we deduce whether the $A \mapsto A^{\sigma}$ of our MES-preserving Φ is the identity or the transpose.

Having classified the possible maps ζ , we then construct an MES-preserving extension $\tilde{\Phi} : \text{MES}_{\mathcal{V},\mathcal{V}} \to \text{MES}_{\mathcal{V},\mathcal{V}}$ and apply Theorem 1.2 to this extension $\tilde{\Phi}$. By considering the restriction of this map $\tilde{\Phi}$ to a subspace identified with $\text{MES}_{\mathcal{X},\mathcal{V}}$, we conclude that the restriction Φ must be of the form described in Theorem 1.1.

The paper is organized as follows. In Section 2, we introduce definitions, particularly the definition of MES (the set of maximally entangled states) and the definition of our vectorization operator. In Section 3, we discuss some characterizations of $MES_{\mathcal{X},\mathcal{Y}}$, and how it relates to the set of coisometries $coisom(\mathcal{Y},\mathcal{X})$. In Section 4, we define the continuous maps on MES that we will rely on in order to define the extension $\tilde{\Phi}$. In Section 5, we consider how ζ behaves on pairwise-orthogonal coisometries. In Section 6, we use ζ to define the extension $\tilde{\Phi} : MES_{\mathcal{Y},\mathcal{Y}} \to MES_{\mathcal{Y},\mathcal{Y}}$. Finally, in Section 7 we prove Theorem 1.1, our main result.

2. Definitions

Let \mathcal{X}, \mathcal{Y} denote (finite dimensional, complex) Hilbert spaces with dim $(\mathcal{X}) = m$, dim $(\mathcal{Y}) = n$. Let $\mathcal{X} \otimes \mathcal{Y}$ denote the tensor product of these spaces, which is itself a Hilbert space of dimension mn. Let $L(\mathcal{X}, \mathcal{Y})$ denote the space of linear

 $\mathbf{2}$

transformations between \mathcal{X} and \mathcal{Y} . We will also use the shorthand $L(\mathcal{X}) = L(\mathcal{X}, \mathcal{X})$. We will use $\mathbb{1}_{\mathcal{X}}$ to denote the identity operator over the Hilbert space \mathcal{X} .

We say that an operator $A : \mathcal{X} \to \mathcal{Y}$ is an *isometry* if it satisfies $A^*A = \mathbb{1}_{\mathcal{X}}$. We will say that A is a *coisometry* if its adjoint operator $A^* : \mathcal{Y} \to \mathcal{X}$ is an isometry.

A unit vector $u \in \mathcal{X} \otimes \mathcal{Y}$ is said to be *maximally entangled* if it can be written in the form

(2.1)
$$u = \sum_{j=1}^{r} \frac{1}{\sqrt{r}} x_j \otimes y_j$$

for orthonormal sets of vectors $\{x_1, \ldots, x_r\} \subset \mathcal{X}, \{y_1, \ldots, y_r\} \subset \mathcal{Y}$ and $r = \min\{m, n\}$.

The pure state (rank-1, trace-1, positive semidefinite operator) associated with u is the projection operator $uu^* \in L(\mathcal{X} \otimes \mathcal{Y})$. Such an operator is called a maximally entangled state. That is, we define the set $MES_{\mathcal{X},\mathcal{Y}}$ of maximally entangled states over \mathcal{X} and \mathcal{Y} to be

(2.2)
$$\operatorname{MES}_{\mathcal{X},\mathcal{Y}} = \{ uu^* : u \text{ is maximally entangled} \} \subset L(\mathcal{X} \otimes \mathcal{Y}).$$

We will sometimes denote $MES_{\mathcal{X},\mathcal{Y}}$ as MES (without its subscripts) when the context is clear.

We define the vectorization operator vec : $L(\mathcal{Y}, \mathcal{X}) \to \mathcal{X} \otimes \mathcal{Y}$ to be the linear map satisfying

(2.3)
$$\operatorname{vec}(xy^T) = x \otimes y$$
 for all $x \in \mathcal{X}, y \in \mathcal{Y}$.

Notably, vec is an isomorphism of Hilbert spaces. It may be helpful to note that, in terms of matrices, vec can be written as vec : $\mathbb{C}^{m \times n} \to \mathbb{C}^{mn}$,

(2.4)
$$\operatorname{vec}([a_{ij}]_{i=1}^{m}) = (a_{11}, \dots, a_{1n}, a_{21}, \dots, a_{2n}, \dots, a_{m1}, \dots, a_{mn})^T,$$

which is to say that our vectorization operator "stacks the rows" of a matrix (in contrast to the other common convention of "stacking the columns"). We adopted this convention from [6].

Throughout the paper e_1, \ldots, e_n will denote the canonical basis for \mathbb{C}^n (for instance, $e_1 = (1, 0, \ldots, 0)^T$). We will also take $E_{ij} \in \mathbb{C}^{m \times n}$ to be the matrix $E_{ij} = e_i e_j^T$, i.e. the matrix that has a 1 as its i, j entry and zeros elsewhere.

Remark 2.1. In the remainder of the paper, we will take $\dim(\mathcal{X}) = m < \infty$ and $\dim(\mathcal{Y}) = n < \infty$, with $m \le n$.

3. Characterizing MES

Proposition 3.1. If $n \ge m$, u is maximally entangled in $\mathcal{X} \otimes \mathcal{Y}$ iff there exists a coisometry $A : \mathcal{Y} \to \mathcal{X}$ such that $u = \frac{1}{\sqrt{m}} \operatorname{vec}(A)$.

Proof: Suppose that u is a maximally entangled unit vector. Let $\{x_1, \ldots, x_m\}$ and $\{y_1, \ldots, y_n\}$ be orthonormal bases of \mathcal{X} and \mathcal{Y} such that $u = \frac{1}{\sqrt{m}} \sum_{j=1}^m x_j \otimes y_j$. It follows that

(3.1)
$$\operatorname{vec}^{-1}(u) = \frac{1}{\sqrt{m}} \sum_{j=1}^{m} \operatorname{vec}^{-1}(x_j \otimes y_j) = \frac{1}{\sqrt{m}} \left(\sum_{j=1}^{m} x_j y_j^T \right)$$

and we recognize that $A = \sum_{j=1}^m x_j y_j^T$ is a singular value decomposition of a coisometry.

Conversely, suppose $u = \frac{1}{\sqrt{m}} \operatorname{vec}(A)$. Then A has a singular value decomposition $A = \sum_{j=1}^{m} x_j y_j^T$ so that

(3.2)
$$u = \frac{1}{\sqrt{m}} \operatorname{vec}\left(\sum_{j=1}^{m} x_j y_j^T\right) = \frac{1}{\sqrt{m}} \sum_{j=1}^{m} x_j \otimes y_j$$

is maximally entangled.

Let $\operatorname{tr}_{\mathcal{Y}} : L(\mathcal{X} \otimes \mathcal{Y}) \to L(\mathcal{X})$ denote the "partial trace" over the space \mathcal{Y} . That is, we define $\operatorname{tr}_{\mathcal{Y}}$ to be the unique linear operator satisfying

(3.3)
$$\operatorname{tr}_{\mathcal{Y}}(A \otimes B) = \operatorname{tr}(B)A$$
 for all $A \in L(\mathcal{X}), \quad B \in L(\mathcal{Y}).$

The following lemma is stated, for instance, in Equation 1.133 of [6].

Lemma 3.2. For any operators $A, B : \mathcal{Y} \to \mathcal{X}$, we have

(3.4)
$$\operatorname{tr}_{\mathcal{Y}}(\operatorname{vec}(A)\operatorname{vec}(B)^*) = AB^*.$$

Proof: Suppose first that A and B are rank-1. We can then write $A = uv^T$ and $B = xy^T$ for $u, x \in \mathcal{X}$ and $v, y \in \mathcal{Y}$. We find that

$$tr_{\mathcal{Y}}(\operatorname{vec}(A)\operatorname{vec}(B)^*) = tr_{\mathcal{Y}}((u \otimes v)(x \otimes y)^*)$$
$$= tr_{\mathcal{Y}}((ux^*) \otimes (vy^*))$$
$$= (y^*v)ux^* = u\overline{v^*y}x^* = (uv^T)(xy^T)^* = AB^*.$$

Now, if $A = \sum_j A_j$ and $B = \sum_k B_k$ where $A_j, B_k \in L(\mathcal{Y}, \mathcal{X})$ have rank 1, then

$$\operatorname{tr}_{\mathcal{Y}}(\operatorname{vec}(A)\operatorname{vec}(B)^*) = \operatorname{tr}_{\mathcal{Y}}\left(\operatorname{vec}\left(\sum_j A_j\right)\operatorname{vec}\left(\sum_k B_k\right)^*\right)$$
$$= \sum_{j,k} \operatorname{tr}_{\mathcal{Y}}(\operatorname{vec}(A_j)\operatorname{vec}(B_k)^*) = \sum_{j,k} A_j B_k^*$$
$$= \left(\sum_j A_j\right)\left(\sum_k B_k\right)^* = AB^*.$$

Since any operator can be written as a sum of rank-1 operators, the desired conclusion holds.

Lemma 3.3. For all $M \in \text{MES}_{\mathcal{X},\mathcal{Y}}$, we have $\text{tr}_{\mathcal{Y}}(M) = \frac{1}{m} \mathbb{1}_{\mathcal{X}}$. Moreover, for $M \in \text{span}(\text{MES}_{\mathcal{X},\mathcal{Y}})$, we have $\text{tr}_{\mathcal{Y}}(M) = \frac{\text{tr}(M)}{m} \mathbb{1}_{\mathcal{X}}$.

Proof: By Proposition 3.1, there exists a coisometry $A : \mathcal{Y} \to \mathcal{X}$ such that $M = \frac{1}{m} \operatorname{vec}(A) \operatorname{vec}(A)^*$. By Lemma 3.2, we find that

(3.5)
$$\operatorname{tr}_{\mathcal{Y}}(M) = \frac{1}{m} \operatorname{tr}_{\mathcal{Y}}(\operatorname{vec}(A) \operatorname{vec}(A)^*) = \frac{1}{m} A A^* = \frac{1}{m} \mathbb{1}_{\mathcal{X}}$$

By the linearity of the partial trace, it follows that $\operatorname{tr}_{\mathcal{Y}}(M) = \frac{\operatorname{tr}(M)}{m} \mathbb{1}_{\mathcal{X}}$ for all $M \in \operatorname{span}(\operatorname{MES}_{\mathcal{X},\mathcal{Y}}).$

The following proposition generalizes Remark 2.3 in [5].

Proposition 3.4. The set of pure states in the real linear span of MES is precisely MES. That is: if M is a rank-1, trace-1, positive operator, then $M \in \text{span}(\text{MES})$ implies that $M \in \text{MES}$.

Proof: Let M be a rank-1, trace-1, positive operator in span(MES). Let u be a unit-vector such that $M = uu^*$. Let $A : \mathcal{Y} \to \mathcal{X}$ be an operator such that $u = \frac{1}{\sqrt{m}} \operatorname{vec}(A)$. By Lemma 3.3, $M \in \operatorname{span}(\operatorname{MES}_{\mathcal{X},\mathcal{Y}})$ implies that

(3.6)
$$\operatorname{tr}_{\mathcal{Y}}(M) = \frac{\operatorname{tr} M}{m} \mathbb{1}_{\mathcal{X}} = \frac{1}{m} \mathbb{1}_{\mathcal{X}}.$$

By Lemma 3.2, this implies that

(3.7)
$$\mathbb{1}_{\mathcal{X}} = m \operatorname{tr}_{\mathcal{Y}}(M) = \operatorname{tr}_{\mathcal{Y}}(\operatorname{vec}(A) \operatorname{vec}(A^*)) = AA^*$$

That is, A is a coisometry. By Proposition 3.1, we may conclude that $u = \frac{1}{\sqrt{m}} \operatorname{vec}(A)$ is maximally entangled, and that $M = uu^*$ is an element of MES.

4. Continuous Maps on MES

Remark 4.1. In the remainder of the paper, Φ will denote an invertible linear map Φ : span(MES_{\mathcal{X},\mathcal{Y}}) \rightarrow span(MES_{\mathcal{X},\mathcal{Y}}) such that Φ (MES_{\mathcal{X},\mathcal{Y}}) \subset MES_{\mathcal{X},\mathcal{Y}}.

For a (finite-dimensional) vector space V and a subset $S \subset V$, let P(S) denote complex projective space over S, the topological space of one-dimensional \mathbb{C} -subspaces of V that contain a non-zero element of S. For an element $A \in S$, we use $[A] = \{zA : z \in \mathbb{C}\}$ to denote the element of P(S) containing A.

For $A \in L(\mathcal{Y}, \mathcal{X})$, let $\pi_A \in L(\mathcal{X} \otimes Y)$ be given by

$$\pi_A = \frac{1}{\operatorname{tr}(AA^*)}\operatorname{vec}(A)\operatorname{vec}(A)^*.$$

We see that $\pi_A \in \text{MES}_{\mathcal{X},\mathcal{Y}}$ if and only if $A : \mathcal{Y} \to \mathcal{X}$ is a (complex) multiple of a coisometry (that is, if $AA^* = k \mathbb{1}_{\mathcal{X}}$ for some k > 0).

Remark 4.2. Let K be a compact subset of the finite-dimensional normed vector space $L(\mathcal{Y}, \mathcal{X})$. The map π is constant over any equivalence class in P(K), so the induced map from P(K) to $L(\mathcal{X} \otimes \mathcal{Y})$ given by $[A] \mapsto \pi_A$ is well-defined. Moreover, this induced map is a homeomorphism between P(K) and the map's image.

With $K = \text{coisom}(\mathcal{Y}, \mathcal{X})$ in particular, we see that the image of the map $[A] \mapsto \pi_A$ is precisely $\text{MES}_{\mathcal{X},\mathcal{Y}}$.

Proof: To see that the map $[A] \mapsto \pi_A$ is well defined and continuous over the quotient space P(K), it suffices to note that if $A_1, A_2 \in K$ satisfy $A_1 \sim A_2$ (i.e. if

 $A_1 = kA_2$ for some non-zero $k \in \mathbb{C}$), then $\pi_{A_1} = \pi_{A_2}$. We see moreover that this induced map is injective: that is, we observe that if $\pi(A_1) = \pi(A_2)$, then $A_1 \sim A_2$. Because K is compact, the quotient space $P(K) = K/\sim$ is also compact.

So, we have found that the induced map $[A] \mapsto \pi_A$ is a continuous, injective map from the compact quotient space K/\sim to the Hausdorff space $L(\mathcal{X} \otimes \mathcal{Y})$. Because our map is an injective, continuous map with compact domain and Hausdorff codomain, it is a homeomorphism between the map's domain and the map's image (by Theorem 26.6 of [4]), as desired.

With the above in mind, we note that the set $\operatorname{coisom}(\mathcal{Y}, \mathcal{X})$ is compact in $L(\mathcal{Y}, \mathcal{X})$. We define $\zeta : P(\operatorname{coisom}(\mathcal{Y}, \mathcal{X})) \to P(\operatorname{coisom}(\mathcal{Y}, \mathcal{X}))$ to be the unique linear map such that for all $A \in \operatorname{coisom}(\mathcal{Y}, \mathcal{X})$, we have $\pi_{\mathcal{L}[A]} = \Phi(\pi_A)$.

That is, we define ζ so that the following diagram (of continuous maps) commutes:

(4.1)
$$\begin{array}{c} \operatorname{MES}_{\mathcal{X},\mathcal{Y}} & \xrightarrow{\Phi} & \operatorname{MES}_{\mathcal{X},\mathcal{Y}} \\ & & & & \\ \pi & & & & \\ P(\operatorname{coisom}(\mathcal{Y},\mathcal{X})) & \xrightarrow{\zeta} & P(\operatorname{coisom}(\mathcal{Y},\mathcal{X})) \end{array}$$

Since Φ is a continuous, invertible map on $\text{MES}_{\mathcal{X},\mathcal{Y}}$ and π is a homeomorphism, we see that ζ must also be a continuous, invertible map on $P(\text{coisom}(\mathcal{Y},\mathcal{X}))$.

This map ζ will be particularly important to the construction of $\tilde{\Phi}$ in Section 6. To begin, we show that ζ satisfies the following "preservation of subspaces" property.

Proposition 4.3. For $A_1, A_2, A_3 \in \text{coisom}(\mathcal{Y}, \mathcal{X})$: if $[A_3] \subset [A_1] + [A_2]$, then $\zeta[A_3]$ must be a subspace of $\zeta[A_1] + \zeta[A_2]$.

Proof: Equivalently, we wish to show that

 $\exists \alpha > 0$ such that $\pi_{A_1} + \pi_{A_2} - \alpha \pi_{A_3}$ is positive semidefinite \Longrightarrow

 $\exists \beta > 0$ such that $\pi_{\zeta[A_1]} + \pi_{\zeta[A_2]} - \beta \pi_{\zeta[A_3]}$ is positive semidefinite.

That is, we wish to show that

 $\exists \alpha > 0$ such that $\pi_{A_1} + \pi_{A_2} - \alpha \pi_{A_3}$ is positive semidefinite \implies

 $\exists \beta > 0$ such that $\Phi(\pi_{A_1} + \pi_{A_2} - \beta \pi_{A_3})$ is positive semidefinite.

So, suppose that there exists an $\alpha > 0$ such that $\pi_{A_1} + \pi_{A_2} - \alpha \pi_{A_3}$ is positive semidefinite. It follows that there exists an $\gamma \ge \alpha$ such that $\pi_{A_1} + \pi_{A_2} - \gamma \pi_{A_3}$ is rank 1 and positive semidefinite. Since $\pi_{A_1} + \pi_{A_2} - \gamma \pi_{A_3}$ is a rank-1 positive semidefinite element of span(MES), we may apply Proposition 3.4 to state that there exists a k > 0 and coisometry $B: \mathcal{Y} \to \mathcal{X}$ such that

$$\pi_{A_1} + \pi_{A_2} - \gamma \pi_{A_3} = k \pi_B$$

Thus, we conclude that

$$\Phi(\pi_{A_1} + \pi_{A_2} - \alpha \pi_{A_3}) = \Phi(\pi_{A_1} + \pi_{A_2} - \gamma \pi_{A_3} + (\gamma - \alpha) \pi_{A_3})$$

$$= \Phi(k\pi_B + (\gamma - \alpha)\pi_{A_3})$$
$$= k\Phi(\pi_B) + (\gamma - \alpha)\Phi(\pi_{A_3})$$

Thus, the desired condition holds with $\beta = \alpha$.

5. PAIRWISE ORTHOGONAL COISOMETRIES

We will say that two coisometries $A_1, A_2: \mathcal{Y} \to \mathcal{X}$ are orthogonal if $A_1A_2^* = 0$.

Lemma 5.1. For coisometries $A_1, A_2 : \mathcal{Y} \to \mathcal{X}$, the following are equivalent:

- (1) $A_1 A_2^* = 0.$
- (2) $A_2 A_1^* = 0.$

(3) The map $\begin{bmatrix} A_1 \\ A_2 \end{bmatrix} : \mathcal{Y} \to \mathcal{X} \oplus \mathcal{X}$ is a coisometry.

- (4) $\alpha A_1 + \beta A_2$ is a coisometry for $\alpha, \beta \in \mathbb{C}$ whenever $|\alpha|^2 + |\beta|^2 = 1$.
- (5) The images of A_1^*, A_2^* are orthogonal subspaces of \mathcal{Y} .

Proof: 1 \iff 2: It suffices to note that $(A_1A_2^*)^* = A_2A_1^*$. 1 \implies 3: Suppose that $A_2A_1^* = A_1A_2^* = 0$. We then compute

(5.1)
$$\begin{bmatrix} A_1 \\ A_2 \end{bmatrix} \begin{bmatrix} A_1 \\ A_2 \end{bmatrix}^* = \begin{bmatrix} A_1 A_1^* & A_1 A_2^* \\ A_2 A_1^* & A_2 A_2^* \end{bmatrix} = \begin{bmatrix} \mathbb{1}_{\mathcal{X}} & 0 \\ 0 & \mathbb{1}_{\mathcal{X}} \end{bmatrix} = \mathbb{1}_{\mathcal{X} \oplus \mathcal{X}}.$$

 $3 \implies 1$: Suppose that $\begin{bmatrix} A_1 \\ A_2 \end{bmatrix}$ is a coisometry. Then

(5.2)
$$\mathbb{1}_{\mathcal{X}\oplus\mathcal{X}} = \begin{bmatrix} \mathbb{1}_{\mathcal{X}} & 0\\ 0 & \mathbb{1}_{\mathcal{X}} \end{bmatrix} = \begin{bmatrix} A_1\\ A_2 \end{bmatrix} \begin{bmatrix} A_1\\ A_2 \end{bmatrix}^* = \begin{bmatrix} A_1A_1^* & A_1A_2^*\\ A_2A_1^* & A_2A_2^* \end{bmatrix}$$

from which we may conclude that $A_j A_j^* = \mathbb{1}_{\mathcal{X}}$ (that is, A_1, A_2 are coisometries) and $A_1 A_2^* = 0$ (that is, A_1 and A_2 are orthogonal).

 $1 \iff 4$: We compute

(5.3)
$$(\alpha A_1 + \beta A_2)(\alpha A_1 + \beta A_2)^* = |\alpha|^2 A_1 A_1^* + |\beta|^2 A_2 A_2^*$$

$$(5.4) \qquad \qquad + \alpha\beta A_1 A_2^* + \bar{\alpha}\beta A_2 A_1^*$$

(5.5)
$$= (|\alpha|^2 + |\beta|^2) \mathbb{1}_{\mathcal{X}} + (\alpha \bar{\beta} A_1 A_2^*) + (\alpha \bar{\beta} A_1 A_2^*)^*.$$

If $A_1A_2^* = 0$, then we have $(\alpha A_1 + \beta A_2)(\alpha A_1 + \beta A_2)^* = (|\alpha|^2 + |\beta|^2)\mathbb{1}_{\mathcal{X}}$, which is to say that $\alpha A_1 + \beta A_2$ is a coisometry whenever $|\alpha|^2 + |\beta|^2 = 1$. That is, $1 \implies 4$.

Conversely, if 4 holds, then it must be that $(\alpha \bar{\beta} A_1 A_2^*) + (\alpha \bar{\beta} A_1 A_2^*)^* = 0$ for all choices of α, β with $|\alpha|^2 + |\beta|^2 = 1$. Let $\gamma = \alpha \bar{\beta}$, and choose self-adjoint H, K such that $A_1 A_2^* = H + iK$. We then compute

(5.6)
$$(\gamma(H+iK)) + (\gamma(H+iK))^* = (\gamma+\bar{\gamma})H + i(\gamma-\bar{\gamma})K$$

and, by 4, we know that the above expression is zero for all $\gamma = \alpha \overline{\beta}$ with $\alpha, \beta \in \mathbb{C}$ satisfying $|\alpha|^2 + |\beta|^2 = 1$.

Setting $\alpha = \beta = \frac{1}{\sqrt{2}}$, we find $\gamma = \frac{1}{2}$ and we see that we must have H = 0. Setting $\alpha = \frac{i}{\sqrt{2}}$ and $\beta = \frac{1}{\sqrt{2}}$, we find $\gamma = \frac{i}{2}$ and we see that we must have K = 0. Since H = 0 and K = 0, we conclude that $A_1 A_2^* = 0$.

The proof that $5 \iff 1$ is straightforward and therefore omitted.

7

Lemma 5.2. For any two orthogonal coisometries $A_1, A_2 : \mathcal{Y} \to \mathcal{X}$: there exist orthogonal coisometries $B_1, B_2 : \mathcal{Y} \to \mathcal{X}$ with $B_i \in \zeta(A_i)$ such that we have either

(5.7)
$$\Phi\left(\sum_{i,j=1}^{2} a_{ij} \operatorname{vec}(A_i) \operatorname{vec}(A_j)^*\right) = \sum_{i,j=1}^{2} a_{ij} \operatorname{vec}(B_i) \operatorname{vec}(B_j)^* \quad on$$

(5.8)
$$\Phi\left(\sum_{i,j=1}^{2} a_{ij} \operatorname{vec}(A_i) \operatorname{vec}(A_j)^*\right) = \sum_{i,j=1}^{2} a_{ij} \operatorname{vec}(B_j) \operatorname{vec}(B_i)^*.$$

Proof: To begin, select coisometries $\tilde{B}_1 \in \zeta(A_1), \tilde{B}_2 \in \zeta(A_2)$. Because Φ is invertible, \tilde{B}_1 and \tilde{B}_2 must be linearly independent. Let V_1 and V_2 denote the subspaces

$$V_1 = \operatorname{span}\{\operatorname{vec}(A_i)\operatorname{vec}(A_j)^*: i, j = 1, 2\},\$$

$$V_2 = \operatorname{span}\{\operatorname{vec}(\tilde{B}_i)\operatorname{vec}(\tilde{B}_j)^*: i, j = 1, 2\}.$$

We observe that the elements of the form $\pi_{\alpha A_1+\beta A_2}$ span V_1 (for instance: we see that $\{\pi_{A_1}, \pi_{A_2}, \pi_{(A_1+A_2)/2}, \pi_{(A_1+iA_2)/2}\}$ forms a spanning set), and by Proposition 4.3 we see that the image of these elements must lie in V_2 . Thus, Φ takes the subspace V_1 to V_2 .

We define the map $G:\mathbb{C}^{2\times 2}\to\mathbb{C}^{2\times 2}$ so that

$$G\left(\begin{bmatrix}a_{11} & a_{12}\\a_{21} & a_{22}\end{bmatrix}\right) = \begin{bmatrix}b_{11} & b_{12}\\b_{21} & b_{22}\end{bmatrix} \iff \Phi\left(\sum_{i,j=1}^{2} a_{ij}\operatorname{vec}(A_i)\operatorname{vec}(A_j)^*\right) = \sum_{i,j=1}^{2} b_{ij}\operatorname{vec}(\tilde{B}_i)\operatorname{vec}(\tilde{B}_j)^* \iff \Phi\left(\begin{bmatrix}\operatorname{vec}(A_1)\operatorname{vec}(A_2)\end{bmatrix}\begin{bmatrix}a_{11} & a_{12}\\a_{21} & a_{22}\end{bmatrix}\begin{bmatrix}\operatorname{vec}(A_1)^*\\\operatorname{vec}(A_2)^*\end{bmatrix}\right) = \begin{bmatrix}\operatorname{vec}(\tilde{B}_1)\operatorname{vec}(\tilde{B}_2)\end{bmatrix}\begin{bmatrix}b_{11} & b_{12}\\b_{21} & b_{22}\end{bmatrix}\begin{bmatrix}\operatorname{vec}(\tilde{B}_1)^*\\\operatorname{vec}(\tilde{B}_2)^*\end{bmatrix}$$

By our definition of \tilde{B}_1, \tilde{B}_2 , we have $G(E_{11}) = E_{11}, G(E_{22}) = E_{22}$. We note that

(5.9)
$$\pi_{\alpha A_1 + \beta A_2} = \frac{1}{m} \left[\operatorname{vec}(A_1) \operatorname{vec}(A_2) \right] \left(\begin{bmatrix} \alpha \\ \beta \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix}^* \right) \left[\operatorname{vec}(A_1)^* \\ \operatorname{vec}(A_2)^* \end{bmatrix}$$

so that because Φ preserves MES, G maps the rank-1 positive semidefinite matrices in $\mathbb{C}^{2\times 2}$ to the rank-1 positive semidefinite matrices in $\mathbb{C}^{2\times 2}$. Thus, $G : \mathbb{C}^{2\times 2} \to \mathbb{C}^{2\times 2}$ is a positive map. That is: if $A \in \mathbb{C}^{2\times 2}$ is positive semidefinite, then G(A) is also positive semidefinite.

By [7], G must be decomposable. That is, Φ must have the form $G = G_1 + G_2$, where G_1 and $\tau \circ G_2$ are completely positive and τ denotes the transpose map $A \mapsto A^T$.

Let J(G) denote the Choi matrix of G. That is,

$$J(G) = \sum_{i,j=1}^{2} E_{ij} \otimes G(E_{ij}) = (\mathbb{1}_{\mathbb{C}^{2\times 2}} \otimes G)(\operatorname{vec}(\mathbb{1}_{\mathbb{C}^{2}})\operatorname{vec}(\mathbb{1}_{\mathbb{C}^{2}})^{*})).$$

So far, we may deduce that

$$J(G) = J(G_1) + J(G_2) = \begin{bmatrix} G(E_{11}) & G(E_{12}) \\ G(E_{21}) & G(E_{22}) \end{bmatrix} = \begin{bmatrix} 1 & 0 & g_{11} & g_{12} \\ 0 & 0 & g_{21} & g_{22} \\ \hline \bar{g}_{11} & \bar{g}_{21} & 0 & 0 \\ \bar{g}_{12} & \bar{g}_{22} & 0 & 1 \end{bmatrix}$$

Let P_i, Q_i, R_i be defined so that

$$J(G_1) = \begin{bmatrix} P_1 & Q_1 \\ Q_1^* & R_1 \end{bmatrix}, \quad J(G_2) = \begin{bmatrix} P_2 & Q_2 \\ Q_2^* & R_2 \end{bmatrix}$$

Because $G_1, \tau \circ G_2$ are completely positive, their Choi matrices $J(G_1)$ and $J(\tau \circ G_2)$ must be positive semidefinite. Thus, $P_1, P_2, R_1, R_2 \succeq 0$ and

(5.10)
$$\begin{bmatrix} P_1 & Q_1 \\ Q_1^* & R_1 \end{bmatrix} \succeq 0, \quad \begin{bmatrix} P_2^T & Q_2^T \\ Q_2 & R_2^T \end{bmatrix} \succeq 0$$

Because P_1, P_2 are positive definite matrices satisfying $P_1 + P_2 = E_{11}$, we must have $P_i = p_i E_{11}$ with $p_1 + p_2 = 1$ and $p_i \ge 0$. Similarly, $R_i = r_i E_{22}$ with $r_1 + r_2 = 1$ and $r_i \ge 0$.

By (5.10), we must have $Q_1 = q_1 E_{12}$ and $Q_2 = q_2 E_{21}$, where $|q_i| \leq \sqrt{p_i r_i}$ for i = 1, 2.

Now, since G maps rank-1 positive semidefinite matrices to rank-1 positive semidefinite matrices (as stated below (5.9)), it must be that the matrix

$$G\left(\begin{bmatrix}\alpha\\\beta\end{bmatrix}\begin{bmatrix}\alpha\\\beta\end{bmatrix}^*\right) = \begin{bmatrix}|\alpha|^2 & \alpha\bar{\beta}q_1 + \bar{\alpha}\beta\bar{q}_2\\ \bar{\alpha}\beta\bar{q}_1 + \alpha\bar{\beta}q_2 & |\beta|^2\end{bmatrix}$$

is positive semidefinite with rank 1. Thus, its determinant must be zero for all α, β . That is, we must have

$$|\alpha|^2 |\beta|^2 = |\alpha \bar{\beta} q_1 + \bar{\alpha} \beta \bar{q}_2|^2$$

for all choices $\alpha, \beta \in \mathbb{C}$. Taking $\alpha = e^{i\theta}$ and $\beta = 1$, we see that this implies

$$1 = |e^{i\theta}q_1 + e^{-i\theta}q_2|^2, \quad \theta \in \mathbb{R}.$$

Thus, for all values of θ ,

$$|e^{i\theta}q_1 + e^{-i\theta}q_2|^2 = |q_1|^2 + |q_2|^2 + 2\operatorname{Re}[q_1\bar{q}_2e^{i\theta}]$$

is constant. This only occurs if $q_1\bar{q}_2 = 0$, which is to say that $q_1 = 0$ or $q_2 = 0$. That is, we must have

$$J(G) = \begin{bmatrix} 1 & 0 & 0 & e^{-i\alpha} \\ 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\ e^{i\alpha} & 0 & 0 & 1 \end{bmatrix} \quad \text{or} \quad J(G) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & e^{i\alpha} & 0 \\ \hline 0 & e^{-i\alpha} & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

If J(G) has the first form, then we have

$$\Phi\left(\sum_{p,q=1}^{2} a_{pq} \operatorname{vec}(A_p) \operatorname{vec}(A_q)^*\right) = \sum_{p,q=1}^{2} a_{pq} e^{i\alpha(p-q)} \operatorname{vec}(\tilde{B}_p) \operatorname{vec}(\tilde{B}_q)^*$$
$$= \sum_{p,q=1}^{2} a_{pq} \operatorname{vec}(B_p) \operatorname{vec}(B_q)^*,$$

where we have taken $B_1 := \tilde{B}_1$ and $B_2 := e^{i\alpha}\tilde{B}_2$. That is, (5.7) applies, as desired.

If J(G) has the second form, then we have

$$\Phi\left(\sum_{p,q=1}^{2} a_{pq} \operatorname{vec}(A_p) \operatorname{vec}(A_q)^*\right) = \sum_{p,q=1}^{2} a_{pq} e^{i\alpha(q-p)} \operatorname{vec}(\tilde{B}_q) \operatorname{vec}(\tilde{B}_p)^*$$
$$= \sum_{p,q=1}^{2} a_{pq} \operatorname{vec}(B_q) \operatorname{vec}(B_p)^*,$$

where again, we have taken $B_1 := \tilde{B}_1$ and $B_2 := e^{i\alpha}\tilde{B}_2$. That is, (5.8) applies, as desired.

If (5.7) applies, then since Φ preserves MES, we can state by Lemmas 3.2, 3.3, and 5.1 that for all α, β with $|\alpha|^2 + |\beta|^2 = 1$, we have

$$\begin{split} \mathbb{1}_{\mathcal{X}} &= \operatorname{tr}_{\mathcal{Y}}(\Phi[(\operatorname{vec}(\alpha A_{1} + \beta A_{2}) \operatorname{vec}(\alpha A_{1} + \beta A_{2})^{*})]) \\ &= \operatorname{tr}_{\mathcal{Y}}[|\alpha|^{2} \operatorname{vec}(B_{1}) \operatorname{vec}(B_{1})^{*} + |\beta|^{2} \operatorname{vec}(B_{2}) \operatorname{vec}(B_{2})^{*} \\ &+ \alpha \bar{\beta} \operatorname{vec}(B_{1}) \operatorname{vec}(B_{2})^{*} + \bar{\alpha} \beta \operatorname{vec}(B_{2}) \operatorname{vec}(B_{1})^{*}] \\ &= |\alpha|^{2} B_{1} B_{1}^{*} + |\beta|^{2} B_{2} B_{2}^{*} + \alpha \bar{\beta} B_{1} B_{2}^{*} + \bar{\alpha} \beta B_{2} B_{1}^{*} \\ &= (|\alpha|^{2} + |\beta|^{2}) \mathbb{1}_{\mathcal{X}} + 2 \operatorname{Re}[\alpha \bar{\beta} B_{1} B_{2}^{*}] = \mathbb{1}_{\mathcal{X}} + 2 \operatorname{Re}[\alpha \bar{\beta} B_{1} B_{2}^{*}]. \end{split}$$

Thus, we have $\operatorname{Re}[\alpha \overline{\beta} B_1 B_2^*] = 0$ for all $\alpha, \beta \in \mathbb{C}$ with $|\alpha|^2 + |\beta|^2 = 1$, so that $B_1 B_2^* = 0$. Thus, B_1 and B_2 are orthogonal coisometries.

Similarly: if (5.8) applies, then for all α, β with $|\alpha|^2 + |\beta|^2 = 1$, we have

$$\begin{split} \mathbb{I}_{\mathcal{X}} &= \operatorname{tr}_{\mathcal{Y}}(\Phi[(\operatorname{vec}(\alpha A_1 + \beta A_2) \operatorname{vec}(\alpha A_1 + \beta A_2)^*)]) \\ &= \operatorname{tr}_{\mathcal{Y}}[|\alpha|^2 \operatorname{vec}(B_1) \operatorname{vec}(B_1)^* + |\beta|^2 \operatorname{vec}(B_2) \operatorname{vec}(B_2)^* \\ &+ \alpha \bar{\beta} \operatorname{vec}(B_2) \operatorname{vec}(B_1)^* + \bar{\alpha} \beta \operatorname{vec}(B_1) \operatorname{vec}(B_2)^*] \\ &= \mathbb{I}_{\mathcal{X}} + 2 \operatorname{Re}[\alpha \bar{\beta} B_2 B_1^*]. \end{split}$$

We again deduce that B_1 and B_2 are orthogonal coisometries. In either case, we have found that B_1 and B_2 are orthogonal coisometries, as desired.

We now consider maps over $P(\mathbb{C}^2)$ (which would more commonly be notated as $P\mathbb{C}^1$ in topological contexts). We will denote by $[\alpha, \beta]$ the equivalence class of $(\alpha, \beta) \in \mathbb{C}^2$. That is,

$$[\alpha, \beta] = \{k(\alpha, \beta) : k \in \mathbb{C}\} \in P(\mathbb{C}^2)$$

For coisometries A, B satisfying $AB^* = 0$, we can select coisometries $C \in \zeta([A])$ and $D \in \zeta([B])$. For such an A, B, C, D, we then define $f : P(\mathbb{C}^2) \to P(\mathbb{C}^2)$ so that for $(\gamma, \delta) \in f([\alpha, \beta])$, we have

$$\zeta([\alpha A + \beta B]) = [\gamma C + \delta D]$$

That is, if $\iota_{A,B}$ denotes the homeomorphism $[\alpha,\beta] \in P(\mathbb{C}^2) \mapsto [\alpha A + \beta B] \in P(\operatorname{span}\{A,B\})$, then

$$f_{A,B,C,D} = \iota_{C,D}^{-1} \circ \zeta \circ \iota_{A,B}$$

so that f is a well-defined, continuous map.

Lemma 5.3. For any coisometries A, B satisfying $AB^* = 0$ and coisometries $C \in \zeta([A])$ and $D \in \zeta([B])$, the map $f : P(\mathbb{C}^2) \to P(\mathbb{C}^2)$ described above is continuous.

Proof: Let $\iota_{A,B}$ denote the homeomorphism $[\alpha,\beta] \in P(\mathbb{C}^2) \to [\alpha A + \beta B] \in P(\operatorname{span}(A,B))$, and similarly define $\iota_{C,D}$. We note that π induces a homeomorphism between $P(\operatorname{span}(A,B))$ and the image of $P(\operatorname{span}(A,B))$ in $\operatorname{MES}_{\mathcal{X},\mathcal{Y}}$, and a similar observation applies to $\operatorname{span}(C,D)$. Thus, we see that f can be written as

$$f = \iota_{C,D}^{-1} \circ \pi|_{P(\operatorname{span}(C,D))}^{-1} \circ \Phi \circ \pi|_{P(\operatorname{span}(A,B))} \circ \iota_{A,B}$$

and is thus the composition of continuous functions.

Since $P(\mathbb{C}^2)$ is homeomorphic to S^2 (the two-dimensional sphere), we can consider the degree of the map $f_{A,B} : P(\mathbb{C}^2) \to P(\mathbb{C}^2)$. Since f is a homeomorphism, we must have $\deg(f) \in \{1, -1\}$.

Lemma 5.4. For a fixed MES-preserving Φ , there exists an $\epsilon \in \{1, -1\}$ such that for all choices of $A, B, C, D \in \operatorname{coisom}(\mathcal{Y}, \mathcal{X})$ with $AB^* = 0, C \in \zeta([A])$, and $D \in \zeta([B])$, we have $\operatorname{deg}(f_{A,B}) = \epsilon$.

Proof: Let f_0 be the map f corresponding to the coisometries A_0, B_0, C_0, D_0 , and let f_1 be the map corresponding to the coisometries A_1, B_1, C_1, D_1 . In order to show that f_0 and f_1 have the same degree, it suffices to show that these maps are homotopic.

By Lemma 5.1, the set of pairs (A, B) of coisometries such that $AB^* = 0$ is path connected. Thus, there exist paths $A, B : [0, 1] \rightarrow \text{coisom}(\mathcal{Y}, \mathcal{X})$ such that $A(0) = A_0, A(1) = A_1, B(0) = B_0, B(1) = B_1$, and for all $t \in [0, 1]$ we have $A(t)B^*(t) = 0$.

Let $\Gamma(t) = \zeta([A(t)])$ and $\Delta(t) = \zeta([B(t)])$. Γ, Δ are paths in $P(\text{coisom}(\mathcal{Y}, \mathcal{X}))$ connecting $\zeta([A_0])$ to $\zeta([A_1])$ and $\zeta([B_0])$ to $\zeta([B_1])$.

If we consider the tautological vector bundle over $P(\operatorname{coisom}(\mathcal{Y}, \mathcal{X}))$, it suffices to find a non-vanishing (constant magnitude) vector-field over the image Γ connecting $(C_0, \zeta([A_0]))$ to $(C(1), \zeta([A_1]))$ and a non-vanishing vector-field over the image of Δ connecting $(D_0, \zeta([B_0]))$ to $(D_1, \zeta([B_1]))$. These vector fields give us paths C(t), D(t) in $\operatorname{coisom}(\mathcal{Y}, \mathcal{X})$.

We can construct such a vector field as follows. To begin, select a continuous $v, w : [0, 1] \to \mathcal{X} \otimes \mathcal{Y}$ such that for all $t \in [0, 1], \pi_{\zeta([A_t])}v(t) \neq 0$ and $\pi_{\zeta([B_t])}w(t) \neq 0$. With that, take $c(t) = \text{vec}^{-1}(\pi_{\zeta([A_t])}v(t))$. Finally, define $C(t) = e^{i(p+qt)}\frac{c(t)}{\|c(t)\|}$, where $p, q \in \mathbb{R}$ are chosen so that $C(0) = C_0$ and $C(1) = C_1$. Similarly, take $d(t) = \text{vec}^{-1}(\pi_{\zeta([B_t])}w(t))$. Define $D(t) = e^{i(r+st)}\frac{d(t)}{\|d(t)\|}$, where $r, s \in \mathbb{R}$ are chosen so that $D(0) = D_0$ and $D(1) = D_1$.

With that, we see that f_t for $t \in [0, 1]$ is a homotopy of the maps f_0 and f_1 .

Combining Lemmas 5.2 and 5.4 lets us deduce the following.

Proposition 5.5. Let A_1, A_2 be coisometries such that $A_1A_2^* = 0$. Then there exist coisometries $B_1 \in \zeta([A_1]), B_2 \in \zeta([A_2])$ such that for any $\alpha, \beta \in \mathbb{C}$ with $|\alpha|^2 + |\beta|^2 = 1$, we have

$$\zeta([\alpha A_1 + \beta A_2]) = [\alpha B_1 + \beta B_2]$$

in the case that $\epsilon = +1$, or

$$\zeta([\alpha A_1 + \beta A_2]) = [\overline{\alpha} B_1 + \overline{\beta} B_2]$$

in the case that $\epsilon = -1$.

Proof: In the case that (5.7) holds for some coisometries A_1, A_2, B_1, B_2 , we find that $\deg(f_{A_1,A_2,B_1,B_2}) = +1$, so that we have $\epsilon = +1$ globally. Applying (5.7), we find that

$$\Phi(\pi_{\alpha A_{1}+\beta A_{2}}) = \Phi\left[\frac{1}{m}(|\alpha|^{2}\operatorname{vec}(A_{1})\operatorname{vec}(A_{1})^{*} + |\beta|^{2}\operatorname{vec}(A_{2})\operatorname{vec}(A_{2})^{*} + \alpha\bar{\beta}\operatorname{vec}(A_{1})\operatorname{vec}(A_{2})^{*} + \bar{\alpha}\beta\operatorname{vec}(A_{2})\operatorname{vec}(A_{1})^{*})\right]$$
$$= \frac{1}{m}(|\alpha|^{2}\operatorname{vec}(B_{1})\operatorname{vec}(B_{1})^{*} + |\beta|^{2}\operatorname{vec}(B_{2})\operatorname{vec}(B_{2})^{*} + \alpha\bar{\beta}\operatorname{vec}(B_{1})\operatorname{vec}(B_{2})^{*} + \bar{\alpha}\beta\operatorname{vec}(B_{2})\operatorname{vec}(B_{1})^{*})$$
$$= \pi_{\alpha B_{1}+\beta B_{2}}$$

so that we indeed have $\zeta([\alpha A_1 + \beta A_2]) = [\alpha B_1 + \beta B_2].$

Similarly: in the case that (5.8) holds for some coisometries A_1, A_2, B_1, B_2 , we find that $\deg(f_{A_1,A_2,B_1,B_2}) = -1$, so that we have $\epsilon = -1$ globally. Applying (5.8), we find that

$$\Phi(\pi_{\alpha A_{1}+\beta A_{2}}) = \Phi\left[\frac{1}{m}(|\alpha|^{2}\operatorname{vec}(A_{1})\operatorname{vec}(A_{1})^{*} + |\beta|^{2}\operatorname{vec}(A_{2})\operatorname{vec}(A_{2})^{*} + \alpha\bar{\beta}\operatorname{vec}(A_{1})\operatorname{vec}(A_{2})^{*} + \bar{\alpha}\beta\operatorname{vec}(A_{2})\operatorname{vec}(A_{1})^{*})\right]$$
$$= \frac{1}{m}(|\alpha|^{2}\operatorname{vec}(B_{1})\operatorname{vec}(B_{1})^{*} + |\beta|^{2}\operatorname{vec}(B_{2})\operatorname{vec}(B_{2})^{*} + \alpha\bar{\beta}\operatorname{vec}(B_{2})\operatorname{vec}(B_{1})^{*} + \bar{\alpha}\beta\operatorname{vec}(B_{1})\operatorname{vec}(B_{2})^{*})$$
$$= \pi_{\bar{\alpha}B_{1}+\bar{\beta}B_{2}}$$

so that we indeed have $\zeta([\alpha A_1 + \beta A_2]) = [\bar{\alpha}B_1 + \bar{\beta}B_2].$

Remark 5.6. Rather than using the degree of the maps $f_{A,B}$ to discriminate between the possible forms of ζ , we could also use the det J(G), where J(G) denotes the Choi matrix used in the proof of Lemma 5.2. In particular: in the $\epsilon = +1$ case we would compute det J(G) = 0, and in the $\epsilon = -1$ case we would compute det J(G) = -1. By the continuity of the determinant, one may argue that given an invertible MES-preserving map Φ , we must either have det J(G) = 0 for all constructions of G, or det J(G) = -1 for all constructions of G.

6. Constructing an Extension

To begin, we generalize Proposition 5.5.

Proposition 6.1. Let $A_1, \ldots, A_k : \mathcal{Y} \to \mathcal{X}$ be a collection of mutually orthogonal coisometries. Then there exist mutually orthogonal coisometries B_1, \ldots, B_k such that for any $\alpha_1, \ldots, \alpha_k \in \mathbb{C}$ with $|\alpha_1|^2 + \cdots + |\alpha_k|^2 = 1$, we have

$$\zeta([\alpha_1 A_1 + \dots + \alpha_k A_k]) = [\alpha_1 B_1 + \dots + \alpha_k B_k]$$

in the case that $\epsilon = +1$, or

$$\zeta([\alpha_1 A_1 + \dots + \alpha_k A_k]) = [\bar{\alpha}_1 B_1 + \dots + \bar{\alpha}_k B_k]$$

in the case that $\epsilon = -1$.

Proof: In the $\epsilon = 1$ case, we can apply Lemma 5.2 to show that there exist coisometries B_1, \ldots, B_k and $\theta_{pq} \in \mathbb{R}$ for $p, q = 1, \ldots, k$ such that $\theta_{1,q} = 0$ for all q, $\theta_{qp} = -\theta_{pq}$, and we have

$$\Phi\left(\sum_{p,q=1}^{k} a_{pq} \operatorname{vec}(A_p) \operatorname{vec}(A_q)^*\right) = \sum_{p,q=1}^{k} a_{pq} e^{i\theta_{pq}} \operatorname{vec}(B_p) \operatorname{vec}(B_q)^*.$$

Because Φ is MES-preserving, the matrix $[e^{i\theta_{pq}}a_{pq}]_{p,q=1}^k$ must be rank-one and positive semidefinite whenever $[a_{pq}]_{p,q=1}^n$ is rank-one and positive semidefinite. By considering the case where $a_{pq} = 1$ for all p, q, we see that this can only occur if $M = [e^{i\theta_{pq}}]_{p,q}$ has rank one. If M is a rank-one matrix whose first row and column are all 1s, M must be the matrix whose entries are all 1s. Thus, we have $e^{i\theta_{pq}} = 1$ for all p, q. That is, we have

$$\Phi\left(\sum_{p,q=1}^{k} a_{pq} \operatorname{vec}(A_p) \operatorname{vec}(A_q)^*\right) = \sum_{p,q=1}^{k} a_{pq} \operatorname{vec}(B_p) \operatorname{vec}(B_q)^*,$$

which is equivalent to the desired statement,

$$\zeta([\alpha_1 A_1 + \dots + \alpha_k A_k]) = [\alpha_1 B_1 + \dots + \alpha_k B_k].$$

Similarly, in the $\epsilon = -1$ case, we can apply Lemma 5.2 to show that there exist coisometries B_1, \ldots, B_k and $\theta_{pq} \in \mathbb{R}$ for $p, q = 1, \ldots, k$ such that $\theta_{1,q} = 0$ for all q, $\theta_{qp} = -\theta_{pq}$, and we have

$$\Phi\left(\sum_{p,q=1}^{k} a_{pq} \operatorname{vec}(A_p) \operatorname{vec}(A_q)^*\right) = \sum_{p,q=1}^{k} a_{pq} e^{i\theta_{pq}} \operatorname{vec}(B_q) \operatorname{vec}(B_p)^*.$$

Because Φ is MES-preserving, the matrix $[e^{i\theta_{pq}}a_{pq}]_{p,q=1}^k$ must be rank-one and positive semidefinite whenever $[a_{pq}]_{p,q=1}^n$ is rank-one and positive semidefinite. Applying the same analysis as above, we conclude $e^{i\theta_{pq}} = 1$ for all p, q. That is, we have

$$\Phi\left(\sum_{p,q=1}^{k} a_{pq} \operatorname{vec}(A_p) \operatorname{vec}(A_q)^*\right) = \sum_{p,q=1}^{k} a_{pq} \operatorname{vec}(B_q) \operatorname{vec}(B_p)^*,$$

which is equivalent to the desired statement,

$$\zeta([\alpha_1A_1 + \dots + \alpha_kA_k]) = [\bar{\alpha}_1B_1 + \dots + \bar{\alpha}_kB_k].$$

We will also need the following lemma, which is the polarization identity in one of its forms.

Lemma 6.2. Given matrices $A, B \in \mathbb{C}^{p \times q}$, we have

$$AB^* = \frac{1}{4} \sum_{k=0}^{3} i^k (A + i^k B) (A + i^k B)^*$$

Proof: We compute

$$\begin{split} \sum_{k=0}^{3} i^{k} (A+i^{k}B)(A+i^{k}B)^{*} &= \sum_{k=0}^{3} i^{k} (AA^{*}+i^{-k}AB^{*}+i^{k}BA^{*}+BB^{*}) \\ &= \sum_{k=0}^{3} (i^{k}AA^{*}+AB^{*}+i^{2k}BA^{*}+i^{k}BB^{*}) \\ &= 0+4AB^{*}+0+0 \end{split}$$

Thus, we have $AB^* = \frac{1}{4} \sum_{k=0}^{3} i^k (A + i^k B) (A + i^k B)^*$ as desired.

Proposition 6.3. Suppose that $\mathcal{Y} = \mathcal{X}^k := \mathcal{X} \oplus \cdots \oplus \mathcal{X}$. Let $\Phi : \operatorname{span}(\operatorname{MES}_{\mathcal{X},\mathcal{Y}}) \to \operatorname{span}(\operatorname{MES}_{\mathcal{X},\mathcal{Y}})$ denote an invertible linear map such that $\Phi(\operatorname{MES}_{\mathcal{X},\mathcal{Y}}) \subset \operatorname{MES}_{\mathcal{X},\mathcal{Y}}$. Then Φ can be extended to a map $\tilde{\Phi} : \operatorname{span}(\operatorname{MES}_{\mathcal{Y},\mathcal{Y}}) \to \operatorname{span}(\operatorname{MES}_{\mathcal{Y},\mathcal{Y}})$ which satisfies $\tilde{\Phi}(\operatorname{MES}_{\mathcal{Y},\mathcal{Y}}) \subset \operatorname{MES}_{\mathcal{Y},\mathcal{Y}}$.

Proof: Let $\zeta : P(\operatorname{coisom}(\mathcal{Y}, \mathcal{X})) \to P(\operatorname{coisom}(\mathcal{Y}, \mathcal{X}))$ be the map induced by Φ , as defined in Section 4. Let $U : \mathcal{Y} \to \mathcal{Y}$ be a unitary transformation given by $U = [U_j]_{j=1}^k$ (where $U_j : \mathcal{Y} \to \mathcal{X}$). Note that U_p, U_q are coisometries satisfying $U_p U_q^* = 0$ for $1 \le p, q \le k$.

Define Φ : MES_{*Y*,*Y*} \to MES_{*Y*,*Y*} as follows: For j = 1, ..., k let $V_j \in \zeta([U_j])$ be such that $\zeta([\alpha U_p + \beta U_q]) = [\alpha V_p + \beta V_q]$ or $\zeta([\alpha U_p + \beta U_q]) = [\bar{\alpha} V_p + \bar{\beta} V_q]$ for $1 \leq p, q \leq k$, as guaranteed by Proposition 6.1. Then, we take

$$\tilde{\Phi}(\pi_{[U_j]_j}) = \tilde{\Phi}\left(\frac{1}{km}[\operatorname{vec}(U_p)\operatorname{vec}(U_q)^*]_{p,q=1}^k\right)$$
$$= \frac{1}{km}[\operatorname{vec}(V_p)\operatorname{vec}(V_q)^*]_{p,q=1}^k$$

Equivalently, we have defined $\tilde{\Phi}$ so that

$$\tilde{\Phi}(\pi_{[U_j]_j}) = \pi_{[V_j]_j}$$

By Proposition 6.1, $[V_j]_{j=1}^k$ is a coisometry, so that $\pi_{[V_j]_j} \in \text{MES}_{\mathcal{Y},\mathcal{Y}}$. Thus, we see from the above that $\tilde{\Phi}$ preserves $\text{MES}_{\mathcal{Y},\mathcal{Y}}$. It remains to be shown, however, that $\tilde{\Phi}$ as defined above is a linear map.

In the case of $\epsilon = 1$, Lemma 6.2 yields

$$\operatorname{vec}(U_p)\operatorname{vec}(U_q)^* = \frac{1}{4} \sum_{k=0}^3 i^k \operatorname{vec}(U_p + i^k U_q) \operatorname{vec}(U_p + i^k U_q)^*$$
$$\Phi[\operatorname{vec}(U_p)\operatorname{vec}(U_q)^*] = \frac{1}{4} \sum_{k=0}^3 i^k \Phi[\operatorname{vec}(U_p + i^k U_q) \operatorname{vec}(U_p + i^k U_q)^*]$$
$$= \frac{1}{4} \sum_{k=0}^3 i^k \operatorname{vec}(V_p + i^k V_q) \operatorname{vec}(V_p + i^k V_q)^*$$
$$= \operatorname{vec}(V_p)\operatorname{vec}(V_q)^*$$

which means that our extension can be written as

$$\tilde{\Phi}(\pi_{[U_j]_j}) = \tilde{\Phi}\left(\frac{1}{km} [\operatorname{vec}(U_p)\operatorname{vec}(U_q)^*]_{p,q=1}^k\right)$$
$$= \frac{1}{km} [\Phi(\operatorname{vec}(U_p)\operatorname{vec}(U_q)^*)]_{p,q=1}^k.$$

So, for every $M \in \operatorname{MES}_{\mathcal{Y},\mathcal{Y}} \subset L(\mathcal{Y} \otimes \mathcal{Y}) = L(\mathbb{C}^k) \otimes L(\mathcal{X} \otimes Y)$, we have $\tilde{\Phi}(M) = (\mathbb{1}_{\mathbb{C}^{k \times k}} \otimes \Phi(M))$. That is, $\tilde{\Phi}|_{\operatorname{MES}_{\mathcal{Y},\mathcal{Y}}}$ is the restriction of a linear map, and is therefore linear.

In the case of $\epsilon = -1$, Lemma 6.2 yields

$$\begin{aligned} \operatorname{vec}(U_p) \operatorname{vec}(U_q)^* &= \frac{1}{4} \sum_{k=0}^3 i^k \operatorname{vec}(U_p + i^k U_q) \operatorname{vec}(U_p + i^k U_q)^* \\ \Phi[\operatorname{vec}(U_p) \operatorname{vec}(U_q)^*] &= \frac{1}{4} \sum_{k=0}^3 i^k \Phi[\operatorname{vec}(U_p + i^k U_q) \operatorname{vec}(U_p + i^k U_q)^*] \\ &= \frac{1}{4} \sum_{k=0}^3 i^k \operatorname{vec}(V_p + i^{-k} V_q) \operatorname{vec}(V_p + i^{-k} V_q)^*] \\ &= \frac{1}{4} \sum_{k=0}^3 i^k \left(i^{-k} \operatorname{vec}(i^k V_p + V_q) \right) \left(i^{-k} \operatorname{vec}(i^k V_p + V_q)^* \right)] \\ &= \frac{1}{4} \sum_{k=0}^3 i^k \operatorname{vec}(V_q + i^k V_p) \operatorname{vec}(V_q + i^k V_p)^*] \\ &= \operatorname{vec}(V_q) \operatorname{vec}(V_p)^* \end{aligned}$$

which means that our extension can be written as

$$\tilde{\Phi}(\pi_{[U_j]_j}) = \tilde{\Phi}\left(\frac{1}{km} [\operatorname{vec}(U_p) \operatorname{vec}(U_q)^*]_{p,q=1}^k\right)$$
$$= \frac{1}{km} [\Phi(\operatorname{vec}(U_q) \operatorname{vec}(U_p)^*)]_{p,q=1}^k$$

So, for every $M \in \operatorname{MES}_{\mathcal{Y},\mathcal{Y}} \subset L(\mathcal{Y} \otimes \mathcal{Y}) = L(\mathbb{C}^k) \otimes L(\mathcal{X} \otimes \mathcal{Y})$, we have $\tilde{\Phi}(M) = (\tau_{\mathbb{C}^{k \times k}} \otimes \Phi)(M)$. That is, $\tilde{\Phi}|_{\operatorname{MES}_{\mathcal{Y},\mathcal{Y}}}$ is the restriction of a linear map, and is therefore linear.

7. The preservers of MES

We now consider the possible forms of this extension, using Theorem 1 from [5], which is to say Theorem 1.2 from the introduction. We note in particular that if $\Phi : \operatorname{span}(\operatorname{MES}_{\mathcal{Y},\mathcal{Y}}) \to \operatorname{span}(\operatorname{MES}_{\mathcal{Y},\mathcal{Y}})$ is invertible, then it must be of the form (1) or (2).

Let ad_U denote the map $\operatorname{ad}_U : A \mapsto UAU^{-1}$. We make the following observation regarding these extended maps:

Lemma 7.1. If $B \in L(\mathcal{X} \otimes \mathcal{Y})$ satisfies BM = MB for all $M \in MES_{\mathcal{X},\mathcal{Y}}$, then $B = c\mathbb{1}_{\mathcal{X} \otimes \mathcal{Y}}$ for some $c \in \mathbb{C}$.

Proof: For every maximally entangled vector $u \in \mathcal{X} \otimes \mathcal{Y}$, we have

$$Buu^* = uu^*B \implies (Buu^*)u = (uu^*B)u \implies Bu = (u^*Bu)u.$$

That is, every maximally entangled vector u is an eigenvector of B with associated eigenvalue u^*Bu . Thus, for every maximally entangled vector u, u^*Bu is an eigenvalue of B. Because the set of maximally entangled vectors is connected and the map $u \mapsto u^*Bu$ is continuous, the set

 $\Omega = \{ u^* B u : u \in \mathcal{X} \otimes \mathcal{Y} \text{ is maximally entangled} \}$

is connected. Also, Ω is a subset of the spectrum of B, which is a finite set. Since Ω is connected and finite, it must be a singleton; thus there is a $c \in \mathbb{C}$ such that $\Omega = \{c\}$. Consequently, Bu = cu for all maximally entangled vectors u. Since the maximally entangled vectors span $\mathcal{X} \otimes \mathcal{Y}$, we may conclude that $B = c \mathbb{1}_{\mathcal{X} \otimes \mathcal{Y}}$ as desired.

Lemma 7.2. If $U_1, U_2 \in L(\mathcal{X})$ and $V_1, V_2 \in L(\mathcal{Y})$ are unitary operators such that

$$\operatorname{ad}_{U_1\otimes V_1}(M) = \operatorname{ad}_{U_2\otimes V_2}(M) \quad \text{for all } M \in \operatorname{MES}_{\mathcal{X},\mathcal{Y}}$$

then we must have $U_1 \otimes V_1 = c U_2 \otimes V_2$ for some $c \in \mathbb{C}$.

Proof: We may rewrite the above condition as

$$(U_2 \otimes V_2)^* (U_1 \otimes V_1) M = M (U_2 \otimes V_2)^* (U_1 \otimes V_1) \quad \text{for all } M \in \text{MES}_{\mathcal{X}, \mathcal{Y}}.$$

Lemma 7.1 now gives $(U_2 \otimes V_2)^* (U_1 \otimes V_1) = c \mathbb{1}_{\mathcal{X} \otimes \mathcal{Y}}$ for some $c \in \mathbb{C}.$

Remark 7.3. From here on, we suppose that $\mathcal{Y} = \mathcal{X}^k$, for some integer $k \geq 2$. We also identify $\mathcal{Y} = \mathcal{X}^k$ with $\mathbb{C}^k \otimes \mathcal{X}$, where it is convenient to do so.

We now observe that $\tilde{\Phi}$ has the following properties:

Lemma 7.4. Define

$$P_{j} = [\mathbb{1}_{\mathbb{C}^{k}} - 2E_{jj}] \otimes \mathbb{1}_{\mathcal{X}} \in L(\mathbb{C}^{k} \otimes \mathcal{X}) = L(\mathcal{Y}),$$
$$Q_{pq} = (T_{pq} \otimes \mathbb{1}_{\mathcal{X}}) \otimes \mathbb{1}_{\mathcal{Y}} \in L((\mathbb{C}^{k} \otimes \mathcal{X}) \otimes \mathcal{Y}) = L(\mathcal{Y} \otimes \mathcal{Y}).$$

where T_{pq} is the permutation matrix corresponding to the transposition of the p, q entries of a vector in \mathbb{C}^k .

Then $\tilde{\Phi} \circ \operatorname{ad}_{P_i \otimes \mathbb{1}_{\mathcal{V}}} = \operatorname{ad}_{P_i \otimes \mathbb{1}_{\mathcal{V}}} \circ \tilde{\Phi}$, and $\tilde{\Phi} \circ \operatorname{ad}_{Q_{pq}} = \operatorname{ad}_{Q_{pq}} \circ \tilde{\Phi}$.

Proof: In order to avoid cumbersome notation, we prove the result for P_j with j = 1 and for Q_{pq} with p = 1, q = 2. However, the same proof can be applied for arbitrary indices.

Let M be an element of $MES_{\mathcal{Y},\mathcal{Y}}$, with

$$M = [M_{p,q}]_{p,q=1}^k, \quad M_{pq} \in L(\mathcal{X} \otimes \mathcal{Y}).$$

We compute

$$\begin{split} \mathrm{ad}_{P_1\otimes \mathbb{1}_{\mathcal{Y}}}(\tilde{\Phi}(M)) &= \tilde{\Phi}(\mathrm{ad}_{P_1\otimes \mathbb{1}_{\mathcal{Y}}}(M)) \\ &= \begin{cases} \begin{pmatrix} \Phi(M_{11}) & -\Phi(M_{12}) & -\Phi(M_{13}) & \cdots \\ -\Phi(M_{21}) & \Phi(M_{22}) & \Phi(M_{23}) & \cdots \\ -\Phi(M_{31}) & \Phi(M_{32}) & \Phi(M_{33}) & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} & \epsilon = 1 \\ \\ & \\ & \\ \begin{pmatrix} \Phi(M_{11}) & -\Phi(M_{21}) & -\Phi(M_{31}) & \cdots \\ -\Phi(M_{12}) & \Phi(M_{22}) & \Phi(M_{32}) & \cdots \\ -\Phi(M_{13}) & \Phi(M_{23}) & \Phi(M_{33}) & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} & \epsilon = -1 \end{split}$$

And similarly,

$$\begin{split} \mathrm{ad}_{Q_{12}}(\tilde{\Phi}(M)) &= \tilde{\Phi}(\mathrm{ad}_{Q_{12}}(M)) \\ & = \begin{cases} \begin{pmatrix} \Phi(M_{22}) & \Phi(M_{21}) & \Phi(M_{23}) & \cdots \\ \Phi(M_{12}) & \Phi(M_{11}) & \Phi(M_{13}) & \cdots \\ \Phi(M_{32}) & \Phi(M_{31}) & \Phi(M_{33}) & \cdots \\ \vdots & \vdots & \vdots & \ddots \\ \end{pmatrix} & \epsilon = 1 \\ \\ & = \begin{cases} \begin{pmatrix} \Phi(M_{22}) & \Phi(M_{12}) & \Phi(M_{32}) & \cdots \\ \Phi(M_{21}) & \Phi(M_{11}) & \Phi(M_{31}) & \cdots \\ \Phi(M_{23}) & \Phi(M_{13}) & \Phi(M_{33}) & \cdots \\ \vdots & \vdots & \vdots & \ddots \\ \end{pmatrix} & \epsilon = -1 \end{split}$$

Lemma 7.5. Fix any unitary $U, V \in L(\mathcal{Y})$, and let $\mathbf{S} : L(\mathcal{Y} \otimes \mathcal{Y}) \to L(\mathcal{Y} \otimes \mathcal{Y})$ be the switch-operator, defined by $A \otimes B \mapsto B \otimes A$. Then for unitary $A \in L(\mathcal{Y})$, we have

$$\mathbf{S}(\pi_A) = \pi_{A^T},$$
$$(\pi_A)^T = \pi_{\overline{A}},$$
$$\mathrm{ad}_{U \otimes V}(\pi_A) = \pi_{U A V^T}.$$

Lemma 7.6. We have $\mathcal{Y} = \mathcal{X}^k$, with $k \geq 2$. Suppose that $\Psi : \operatorname{span}(\operatorname{MES}_{\mathcal{Y},\mathcal{Y}}) \to \operatorname{span}(\operatorname{MES}_{\mathcal{Y},\mathcal{Y}})$ is a linear, invertible, MES-preserving map that commutes with $\operatorname{ad}_{P_j \otimes \mathbb{1}_{\mathcal{Y}}}$ for all j. Then we must have

$$\Psi(M) = (U \otimes V)M^{\sigma}(U \otimes V)^*,$$

where σ is either the transpose or identity map. Moreover, either U must be block diagonal (in other words, each copy of $\mathcal{X} \subset \mathcal{Y}$ is an invariant subspace of U), or k = 2 and $U = \begin{bmatrix} 0 & U_{12} \\ U_{21} & 0 \end{bmatrix}$ with $U_{12}, U_{21} \in L(\mathcal{X})$ unitary.

Proof: Suppose that Ψ is a map of the form (1) from Theorem 1.2, with σ the identity map. That is, $\Psi = \operatorname{ad}_{U\otimes V}$. Since Ψ commutes with $\operatorname{ad}_{P_j\otimes \mathbb{1}_{\mathcal{Y}}}$ for all j, we have $\operatorname{ad}_{(P_j\otimes \mathbb{1}_{\mathcal{Y}})(U\otimes V)} = \operatorname{ad}_{(U\otimes V)(P_j\otimes \mathbb{1}_{\mathcal{Y}})}$. By Lemma 7.2, we have $(P_jU)\otimes V = c(UP_j)\otimes V$ for some $c \in \mathbb{C}$. That is, if $U = [U_{p,q}]_{p,q=1}^k$, then we have in the case of j=2

$$\begin{bmatrix} U_{11} & U_{12} & U_{13} & \cdots \\ -U_{21} & -U_{22} & -U_{23} & \cdots \\ U_{31} & U_{32} & U_{33} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \otimes V = c \begin{bmatrix} U_{11} & -U_{12} & U_{13} & \cdots \\ U_{21} & -U_{22} & U_{23} & \cdots \\ U_{31} & -U_{32} & U_{33} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \otimes V.$$

When k > 2, we conclude that $U_{pq} = U_{qp} = 0$ whenever $p \neq q$, as desired. When k = 2 we either have that c = 1 and $U_{12} = U_{21} = 0$, or that c = -1 and $U_{11} = U_{22} = 0$.

Similarly, suppose that Ψ is a map of the form (1) with σ the transpose map, which is to say that $\Psi = \operatorname{ad}_{U \otimes V} \circ \tau$. Since Ψ commutes with $\operatorname{ad}_{P_i \otimes \mathbb{1}_{\mathcal{Y}}}$, we have

$$\begin{aligned} \mathrm{ad}_{P_{j}\otimes\mathbb{1}_{\mathcal{Y}}}\circ\mathrm{ad}_{U\otimes V}\circ\tau &=\mathrm{ad}_{U\otimes V}\circ\tau\circ\mathrm{ad}_{P_{j}\otimes\mathbb{1}_{\mathcal{Y}}} \Longrightarrow \\ \mathrm{ad}_{P_{j}\otimes\mathbb{1}_{\mathcal{Y}}}\circ\mathrm{ad}_{U\otimes V}\circ\tau &=\mathrm{ad}_{U\otimes V}\circ\mathrm{ad}_{P_{j}\otimes\mathbb{1}_{\mathcal{Y}}}\circ\tau \Longrightarrow \\ \mathrm{ad}_{P_{j}\otimes\mathbb{1}_{\mathcal{Y}}}\circ\mathrm{ad}_{U\otimes V} &=\mathrm{ad}_{U\otimes V}\circ\mathrm{ad}_{P_{j}\otimes\mathbb{1}_{\mathcal{Y}}}. \end{aligned}$$

By Lemma 7.2, we have $(P_jU) \otimes V = c(UP_j) \otimes V$ for some $c \in \mathbb{C}$, which again allows us to conclude that U is block-diagonal, or k = 2 and $U = \begin{bmatrix} 0 & U_{12} \\ U_{21} & 0 \end{bmatrix}$.

Now, suppose that Ψ is of the form (2) from Theorem 1.2, with σ the identity map. That is, suppose that $\Psi = \operatorname{ad}_{U\otimes V} \circ \mathbf{S}$; we wish to show that Ψ must fail to commute with $\operatorname{ad}_{P_j\otimes \mathbb{1}_{\mathcal{Y}}}$ for some j. By Lemma 7.5, we find that for unitary $A \in L(\mathcal{Y})$, we have

$$(\mathrm{ad}_{P_j \otimes \mathbb{1}_{\mathcal{Y}}} \circ \Psi)(\pi_A) = \pi_{P_j U A^T V^T}$$
$$(\Psi \circ \mathrm{ad}_{P_i \otimes \mathbb{1}_{\mathcal{Y}}})(\pi_A) = \pi_{U A^T P_i V^T}.$$

Thus, it suffices to find unitary $A \in L(\mathcal{Y})$ such that for some j, $P_j U A^T \neq c U A^T P_j$ for all $c \in \mathbb{C}$. Thus, we see that one $M \in \text{MES}_{\mathcal{Y},\mathcal{Y}}$ for which $(\text{ad}_{P_j \otimes \mathbb{1}_{\mathcal{Y}}} \circ \Psi)(M) \neq$ $(\Psi \circ \text{ad}_{P_j \otimes \mathbb{1}_{\mathcal{Y}}})(M)$ is given by $M = \pi_A$, where A is chosen so that $P_j U A^T$ and $U A^T P_j$ are not multiples. For instance, for j = 1, we can take

$$UA^{T} = \frac{1}{\sqrt{2}} \begin{bmatrix} \mathbb{1}_{\mathcal{X}} & \mathbb{1}_{\mathcal{X}} \\ \mathbb{1}_{\mathcal{X}} & -\mathbb{1}_{\mathcal{X}} \end{bmatrix} \oplus \mathbb{1}_{\mathcal{X}^{k-2}}$$

Similarly, suppose that Ψ is of the form (2) from Theorem 1.2, with σ the transpose map. That is, suppose that $\Psi = \operatorname{ad}_{U\otimes V} \circ \tau \circ \mathbf{S}$. By Lemma 7.5, we find

that for unitary $A \in L(\mathcal{Y})$, we have

$$(\mathrm{ad}_{P_j \otimes \mathbb{1}_{\mathcal{Y}}} \circ \Psi)(\pi_A) = \pi_{P_j U A^* V^T}$$
$$(\Psi \circ \mathrm{ad}_{P_i \otimes \mathbb{1}_{\mathcal{Y}}})(\pi_A) = \pi_{U A^* P_i V^T}.$$

Thus, it suffices to find unitary $A \in L(\mathcal{Y})$ such that for some j, $P_jUA^* \neq cUA^*P_j$ for all $c \in \mathbb{C}$. Thus, we see that one $M \in \text{MES}_{\mathcal{Y},\mathcal{Y}}$ for which $(\text{ad}_{P_j \otimes \mathbb{1}_{\mathcal{Y}}} \circ \Psi)(M) \neq (\Psi \circ \text{ad}_{P_j \otimes \mathbb{1}_{\mathcal{Y}}})(M)$ is given by $M = \pi_A$, where A is chosen so that P_jUA^* and UA^*P_j are not multiples. For instance, we can take

$$UA^* = \frac{1}{\sqrt{2}} \begin{bmatrix} \mathbb{1}_{\mathcal{X}} & \mathbb{1}_{\mathcal{X}} \\ \mathbb{1}_{\mathcal{X}} & -\mathbb{1}_{\mathcal{X}} \end{bmatrix} \oplus \mathbb{1}_{\mathcal{X}^{k-2}}.$$

Lemma 7.7. Let $U, V \in L(\mathcal{Y})$ be unitary with $U \in L(\mathcal{Y}) = L(\mathbb{C}^k \otimes \mathcal{X})$ block diagonal. If Ψ defined by

$$\Psi(M) = (U \otimes V) M^{\sigma} (U \otimes V)^*, \ M \in \operatorname{MES}_{\mathcal{Y}, \mathcal{Y}},$$

commutes with $\operatorname{ad}_{Q_{pq}}$ for $1 \leq p, q \leq k$, then U either has the form

 $U=\mathbb{1}_{\mathbb{C}^k}\otimes W,$

for some unitary $W: \mathcal{X} \to \mathcal{X}$, or k = 2 and $U = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \otimes W$ for some unitary $W: \mathcal{X} \to \mathcal{X}$.

Proof: As in the last proof (invoking Lemma 7.1), we see that if Ψ commutes with $\operatorname{ad}_{Q_{pq}}$, then we must have $Q_{pq}(U \otimes V) = c_{pq}(U \otimes V)Q_{pq}$ for some $c_{pq} \in \mathbb{C}$. Let $\operatorname{diag}[U_j]_{j=1}^k$ denote the block-diagonal matrix with U_j on the *j*th diagonal. Then the above tells us that $U_p = c_{pq}U_q$, $U_q = c_{pq}U_p$ and $U_j = c_{pq}U_j$ for $j \neq p, q$.

When k > 2, this implies that $c_{pq} = 1$ and $U_p = U_q$. As this holds for all pairs p, q, we may conclude that $U_j = U_\ell$ for all $1 \le j, \ell \le k$.

When k = 2, we get from $U_1 = c_{12}U_2 = c_{12}^2U_1$, that $c_{12} = \pm 1$. Thus either $U_1 = U_2$ or $U_1 = -U_2$.

We can now finally prove our main result.

Proof of Theorem 1.1: By Proposition 6.3, Φ has an extension $\tilde{\Phi} : L(\mathcal{Y} \otimes \mathcal{Y}) \to L(\mathcal{Y} \otimes \mathcal{Y})$, as stated there. By Lemma 7.4, this extension commutes with $\mathrm{ad}_{P_j \otimes \mathbb{1}_{\mathcal{Y}}}$ and $\mathrm{ad}_{Q_{pq}}$ for all $1 \leq j, p, q \leq k$. By Lemma 7.6, it follows that $\tilde{\Phi}$ is a map of the form $\tilde{\Phi}(M) = (\tilde{U} \otimes V)M^{\sigma}(\tilde{U} \otimes V)^*$, where σ is either the transpose or identity map and with \tilde{U} either block diagonal or, when k = 2, of the form $\tilde{U} = \begin{bmatrix} 0 & U_{12} \\ U_{21} & 0 \end{bmatrix}$.

We first show that we can discard the last possibility. Indeed, from the proof of Proposition 6.3 we obtain that

$$\tilde{\Phi}(\begin{bmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{bmatrix}) = \begin{bmatrix} \Phi(M_{11}) & \Phi(M_{12}) \\ \Phi(M_{21}) & \Phi(M_{22}) \end{bmatrix} \text{ or } \tilde{\Phi}(\begin{bmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{bmatrix}) = \begin{bmatrix} \Phi(M_{11}) & \Phi(M_{21}) \\ \Phi(M_{12}) & \Phi(M_{22}) \end{bmatrix}$$

If $\tilde{U} = \begin{bmatrix} 0 & U_{12} \\ U_{21} & 0 \end{bmatrix}$, then we obtain that for all $M = [M_{pq}]_{p,q=1}^2 \in \text{MES}_{\mathcal{Y},\mathcal{Y}}$ either

$$\Phi(M_{11}) = (U_{12} \otimes V)M_{22}(U_{12} \otimes V)^* \text{ or } \Phi(M_{11}) = (U_{12} \otimes V)M_{22}^T(U_{12} \otimes V)^*.$$

When dim X > 1 (and thus dim Y > 2) this leads to a contradiction as one can vary M_{22} without changing M_{11} .

If k = 2 and dim $\mathcal{X} = 1$, we have $\mathcal{X} \otimes \mathcal{Y} \cong \mathcal{Y} = \mathbb{C}^2$, and $\Phi : L(\mathcal{X} \otimes Y) \to L(\mathcal{X} \otimes \mathcal{Y})$ defines a map on 2×2 matrices that preserves all rank-one orthogonal projections. This gives that $\Phi(X) = VX^{\sigma}V^*$, where σ is either the transpose or the identity map and V is unitary. Thus Φ is of the desired form.

Going back to the general case of $k \geq 2$, we can conclude that \tilde{U} is block diagonal. Next, by Lemma 7.7, we find that

$$\tilde{U} = \mathbb{1}_{\mathbb{C}^k} \otimes U$$
, or $k = 2$ and $\tilde{U} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \otimes U$,

for some unitary $U : \mathcal{X} \to \mathcal{X}$. We now compute for $M = [M_{pq}]_{p,q=1}^k \in \text{MES}_{\mathcal{Y},\mathcal{Y}}$ that

$$\Phi(M_{11}) = [e_1^T \otimes \mathbb{1}_{\mathcal{X}} \otimes \mathbb{1}_{\mathcal{Y}}] \tilde{\Phi}(M) [e_1^T \otimes \mathbb{1}_{\mathcal{X}} \otimes \mathbb{1}_{\mathcal{Y}}]^*$$

= $[e_1^T \otimes \mathbb{1}_{\mathcal{X}} \otimes \mathbb{1}_{\mathcal{Y}}] (\tilde{U} \otimes V) M^{\sigma} (\tilde{U} \otimes V)^* [e_1 \otimes \mathbb{1}_{\mathcal{X}} \otimes \mathbb{1}_{\mathcal{Y}}]$
= $(U \otimes V) M_{11}^{\sigma} (U \otimes V)^*,$

as desired.

References

- Matej Brešar, Mikhail A. Chebotar, and Wallace S. Martindale, III. Functional identities. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2007.
- [2] Shmuel Friedland, Chi-Kwong Li, Yiu-Tung Poon, and Nung-Sing Sze. The automorphism group of separable states in quantum information theory. J. Math. Phys., 52(4):042203, 8, 2011.
- [3] Alexander Guterman, Chi-Kwong Li, and Peter Šemrl. Some general techniques on linear preserver problems. *Linear Algebra Appl.*, 315(1-3):61–81, 2000.
- [4] James R. Munkres. Topology. Prentice Hall, Inc., Upper Saddle River, NJ, 2000. Second Edition.
- [5] Edward Poon. Preservers of maximally entangled states. *Linear Algebra Appl.*, 468:122–144, 2015.
- [6] John Watrous. The theory of quantum information. Cambridge University Press, 2018.
- [7] S. L. Woronowicz. Positive maps of low dimensional matrix algebras. Rep. Math. Phys., 10(2):165–183, 1976.

DEPARTMENT OF MATHEMATICS, DREXEL UNIVERSITY, 3141 CHESTNUT ST., PHILADELPHIA, PA, 19104

E-mail address: benwgrossmann@gmail.com,hugo@math.drexel.edu