

CONTINUOUS QUIVERS OF TYPE A (I) FOUNDATIONS

KIYOSHI IGUSA, JOB D. ROCK, AND GORDANA TODOROV

ABSTRACT. We generalize type A quivers to continuous type A quivers and prove initial results about pointwise finite-dimensional (pwf) representations. We classify the indecomposable pwf representations and provide a decomposition theorem, recovering results of Botnan and Crawley-Boevey [12, 5]. We also classify the indecomposable pwf projective representations. Finally, we prove that many of the properties of finite-dimensional type A_n representations are present in finitely generated pwf representations. This is the self-contained foundational part of a series of works to study a generalization of continuous clusters categories and their relationship to other type A cluster structures.

CONTENTS

Introduction	1
History	1
Contributions	2
Acknowledgements	3
1. Continuous Quivers of Type A	3
1.1. Quiver of Continuous Type A : $A_{\mathbb{R}}$	4
1.2. Representations of $A_{\mathbb{R}}$: $\text{Rep}_k(A_{\mathbb{R}})$	5
1.3. The Subcategories $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ and $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$	6
2. Classification of Indecomposable Pwf Representations	7
2.1. Projectives	7
2.2. Sufficient Conditions for Indecomposables	12
2.3. Filtrations	13
2.4. Necessary Conditions and Decomposition Theorem	16
2.5. More on $P_{(a)}$, P_a , and the Pointwise Finite Requirement	21
2.6. Relation to Decomposition Theorems in Persistent Homology	22
3. Finitely Generated Representations: $\text{rep}_k(A_{\mathbb{R}})$	22
3.1. Requisites and Definition	23
3.2. Properites of $\text{rep}_k(A_{\mathbb{R}})$	24
3.3. Existence of Some Auslander–Reiten Sequences	26
4. Other Papers in this Series	28
References	28

INTRODUCTION

History. The indecomposable, finite-dimensional representations of type A quivers and were classified by Gabriel in [16]. In particular this yielded an understanding of all finite-dimensional representations of type A quivers.

Date: March 18, 2022.

First author supported by the Simons Foundation.

Representations of quivers, and in particular type A quivers, have been used extensively in persistent homology. Persistent homology has recently been used to study fractal dimension [1, 25] and has been shown to be effective in recovering some signals in noise [20]. Persistent homology has been applied to 3D shape classifications [10], the study of plant root systems [13], identification of breast cancer subtypes [21], and many other real world applications.

Representations of \mathbb{R} and of the infinite zigzag are generalizations of type A_n quiver representations. The first decomposition theorem for representations of \mathbb{R} was proved by Crawley-Boevey in [12]. It states that every pointwise finite representation of \mathbb{R} is a sum of indecomposable representations which are supported on intervals in \mathbb{R} . For general representations of \mathbb{R} the support intervals can be any interval: (a, b) , $(a, b]$, $[a, b)$ or $[a, b]$. Carlsson, de Silva, and Mozorov introduced zigzag persistent homology in [8] and Botnan proved a similar decomposition theorem to Crawley-Boevey's for infinite zigzag persistence in [4].

Representations of quivers have also been used to understand cluster algebras via the construction of cluster categories. Cluster algebras were introduced by Fomin and Zelevinsky in order to better understand scattering diagrams in particle physics [15]. Cluster algebras come equipped with a set of *cluster variables*, sets of cluster variables called *clusters*, and a *mutation* process to move from one cluster to another. Buan, Marsh, Reineke, Rietsin, and the third author constructed a cluster category whose indecomposable objects correspond to cluster variables, maximally rigid sums of indecomposables correspond to clusters, and mutation of clusters was given by homological approximations. In particular, a type A_n cluster algebra can be studied via the cluster category built from finite-dimensional type A_n representations. The first and third author generalized this construction to a continuous version in [19].

Contributions. We generalize type A_n quivers to continuous quivers of type A and study their representations. These generalize representations of the real line which are the basis for the continuous cluster category of [19]. The present paper is a self-contained foundational paper with a focus on representation theoretic techniques. Our goal is to study continuous quivers of type A , representations of such quivers, a generalization of the continuous cluster category, and what these continuous constructions tell us about the corresponding constructions for A_n .

We first consider an alternating orientation on \mathbb{R} given by a discrete subset $S = \{\dots < s_k < s_{k+1} < \dots\} \subset \mathbb{R}$ and a partial ordering \preceq on \mathbb{R} given by $x \prec y$ if $s_{2k} \leq x < y \leq s_{2k+1}$ for some k or if $s_{2k-1} \leq y < x \leq s_{2k}$. The elements $x, y \in \mathbb{R}$ are not related if there is an element of S in the open interval (x, y) . This is the continuous version of the zig-zag which is the quiver with vertex set \mathbb{Z} with one arrow either $i \rightarrow i+1$ or $i+1 \rightarrow i$ between successive integers (see [26]).

Let $A_{\mathbb{R}}$ denote the real line with alternating orientation given by a subset S . For any interval, i.e. connected subset, $I \subseteq \mathbb{R}$, we will construct a pointwise one dimensional representation M_I with support equal to I , called interval indecomposable representations. (See Definition 2.2.2.) The first theorem takes two representations of $A_{\mathbb{R}}$ known to be indecomposable (Proposition 2.2.1) and tells us when they are isomorphic. We allow for any alternating orientation so long as S does not have accumulation points and provide a theorem about indecomposable pwf representations and the decomposition of pwf representations: Theorems A and B, respectively. In Section 2.6 we discuss the relationship between Theorem B, decomposition results in [4, 5, 12], and the choice of method of proof in the present paper.

Theorem A (Theorem 2.3.2). *The representations M_I are indecomposable and any pointwise one-dimensional indecomposable representation of $A_{\mathbb{R}}$ is isomorphic to M_I for some interval $I \subseteq \mathbb{R}$. Let V and V' be indecomposable representations of a continuous type A quiver. Then $V \cong V'$ if and only if $\text{supp } V = \text{supp } V'$.*

Theorem B (Theorem 2.4.15). *Let V be a pointwise finite-dimensional representation of a continuous type A quiver. Then V is a direct sum of interval indecomposables.*

The proofs of Theorems A and B use Theorems 2.1.3 and 2.1.16, which classify the interval projective representations in the category of pointwise finite-dimensional representations, denoted $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. In particular we provide this characterization before we prove our decomposition theorem. Combined with Theorem B, one obtains a complete description of indecomposable projective objects in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ as Theorem C.

Theorem C (Theorems 2.1.3, 2.1.16, and 2.4.15 and Remark 2.4.16). *Any indecomposable projective object in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ is isomorphic to one of the following, where a may be $\pm\infty$.*

- (1) P_a (for $-\infty < a < +\infty$) given by

$$P_a(x) = \begin{cases} k & x \preceq a \\ 0 & \text{otherwise} \end{cases} \quad P_a(x, y) = \begin{cases} 1_k & y \preceq x \preceq a \\ 0 & \text{otherwise.} \end{cases}$$

- (2) P_a given by

$$P_a(x) = \begin{cases} k & x \preceq a, x < a \\ 0 & \text{otherwise} \end{cases} \quad P_a(x, y) = \begin{cases} 1_k & y \preceq x \preceq a, y \leq x < a \\ 0 & \text{otherwise.} \end{cases}$$

- (3) $P_{(a)}$ given by

$$P_{(a)}(x) = \begin{cases} k & x \preceq a, a < x \\ 0 & \text{otherwise} \end{cases} \quad P_{(a)}(x, y) = \begin{cases} 1_k & y \preceq x \preceq a, a < x \leq y \\ 0 & \text{otherwise.} \end{cases}$$

In Section 3 we prove properties about the category of finitely generated representations (Definition 3.1.3) over any continuous quiver of type A, denoted $\text{rep}_k(A_{\mathbb{R}})$. In the A_n case, finitely generated and finite-dimensional representations coincide. Since only finite sums of simple representations of a continuous quiver are finite-dimensional, we instead consider finitely generated representations in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. Theorem D highlights some similarities and differences between $\text{rep}(A_n)$ and $\text{rep}_k(A_{\mathbb{R}})$. Some of the properties extended to pointwise finite-dimensional representations and bounded-dimensional representations (Definition 1.3.1), denoted $\text{Rep}_k^b(A_{\mathbb{R}})$.

Theorem D (Theorem 3.0.1). *Let $A_{\mathbb{R}}$ be a continuous quiver of type A. Then the following hold.*


- (1) *For indecomposable representations M_I and M_J in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, $\text{Rep}_k^b(A_{\mathbb{R}})$, or $\text{rep}_k(A_{\mathbb{R}})$, we have $\text{Hom}(M_I, M_J) \cong k$ or $\text{Hom}(M_I, M_J) = 0$ (Proposition 3.1.2).*
- (2) *Every morphism $f : V \rightarrow W$ in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, $\text{Rep}_k^b(A_{\mathbb{R}})$, or $\text{rep}_k(A_{\mathbb{R}})$ has a kernel, a cokernel, and coinciding image and coimage in that category. (Lemma 3.1.4)*
- (3) *The category $\text{rep}_k(A_{\mathbb{R}})$ Krull-Schmidt, but not artinian (Lemma 3.1.5, Proposition 3.1.7).*
- (4) *The global dimension of $\text{rep}_k(A_{\mathbb{R}})$ is 1 (Proposition 3.2.5).*
- (5) *The Ext space of two indecomposables M_I and M_J in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, $\text{Rep}_k^b(A_{\mathbb{R}})$, or $\text{rep}_k(A_{\mathbb{R}})$ is either isomorphic to k or is 0 (Proposition 3.2.6).*
- (6) *While some Auslander–Reiten sequences exist (Proposition 3.3.2), some indecomposables have neither a left nor a right Auslander–Reiten sequence (Proposition 3.3.3).*

Acknowledgements. The authors would like to thank Ralf Schiffler for organizing the Cluster Algebra School at the University of Connecticut and Shijie Zhu for helpful discussions. They would also like to thank Magnus B. Botnan, Bill Crawley-Boevey, Bernhard Keller, and Francesco Sala for references to related work. The second author would also like to thank Eric Hanson for helpful discussions.

1. CONTINUOUS QUIVERS OF TYPE A

We let k denote a field for the entirety of this paper.

1.1. Quiver of Continuous Type A : $A_{\mathbb{R}}$. The goal of this section is to generalize the definition of type A quivers to a continuous setting. The set \mathbb{R} will serve as the vertices in our quiver. We will choose a set of sinks and sources, which will induce the orientation on the continuous quiver by indicating which vertices have paths to which others. The picture below gives an intuitive idea of the result of choosing a continuous type A quiver and the definition follows.



Definition 1.1.1. A quiver of continuous type A , denoted by $A_{\mathbb{R}}$, is a triple (\mathbb{R}, S, \preceq) , where:

- (1) (a) $S \subset \mathbb{R}$ is a discrete subset, possibly empty, with no accumulation points.
 (b) Order on $S \cup \{\pm\infty\}$ is induced by the order of \mathbb{R} , and $-\infty < s < +\infty$ for $\forall s \in S$.
 (c) Elements of $S \cup \{\pm\infty\}$ are indexed by a subset of $\mathbb{Z} \cup \{\pm\infty\}$ so that s_n denotes the element of $S \cup \{\pm\infty\}$ with index n . The indexing must adhere to the following two conditions:
 - i1 There exists $s_0 \in S \cup \{\pm\infty\}$.
 - i2 If $m \leq n \in \mathbb{Z} \cup \{\pm\infty\}$ and $s_m, s_n \in S \cup \{\pm\infty\}$ then for all $p \in \mathbb{Z} \cup \{\pm\infty\}$ such that $m \leq p \leq n$ the element s_p is in $S \cup \{\pm\infty\}$.
- (2) New partial order \preceq on \mathbb{R} , which we call the orientation of $A_{\mathbb{R}}$, is defined as:
 - p1 The \preceq order between consecutive elements of $S \cup \{\pm\infty\}$ does not change.
 - p2 Order reverses at each element of S .
 - p3 If n is even s_n is a sink.
 - p3' If n is odd s_n is a source.

Definition 1.1.2. Let $A_{\mathbb{R}} = (\mathbb{R}, S, \preceq)$ be a quiver of continuous type A . Then the associated continuous path algebra $kA_{\mathbb{R}}$ is the associative algebra over k (without unity) whose basis consists of pairs (x, y) , where $y \preceq x$. Multiplication on the pairs is given by

$$(w, x)(y, z) = \begin{cases} (w, z) & x = y \\ 0 & x \neq y. \end{cases}$$

Remark 1.1.3. The indexing requirements on S have the following immediate consequences.


- If S is empty then either (i) $s_{-1} = -\infty$ and $s_0 = +\infty$ or (ii) $s_0 = -\infty$ and $s_1 = +\infty$.
- If S is unbounded above (below) then $+\infty = s_{+\infty}$ ($-\infty = s_{-\infty}$).
- If S is bounded above (below) then there is no $s_{+\infty}$ ($s_{-\infty}$) in \tilde{S} .

The rules for the partial order have the following consequences. If $x < y \in \mathbb{R}$ and there is some $s_n \in S$ such that $x < s_n < y$ then $x \not\preceq y$ and $y \not\preceq x$. If $x \leq y \in \mathbb{R}$ and there exists $s_n, s_{n+1} \in \tilde{S}$ such that $s_n \leq x \leq y \leq s_{n+1}$ then:


$$\begin{aligned} x &\preceq y \text{ if } n \text{ is even} \\ y &\preceq x \text{ if } n \text{ is odd.} \end{aligned}$$

Example 1.1.4. We provide four examples of S and the induced partial order \preceq on \mathbb{R} .


- (1) A finite example: $S = \{\frac{1}{2}, \pi\}$, $\bar{S} = \{-\infty, \frac{1}{2}, \pi, +\infty\}$, $s_{-2} = -\infty$, $s_{-1} = \frac{1}{2}$, $s_0 = \pi$, and $s_1 = +\infty$.



- (2) A “half” unbounded example: $S = \{2n : n \in \mathbb{N}\}$, $s_{-1} = -\infty$, $s_n = 2n$ when $n \geq 0$, and $s_{+\infty} = +\infty$.



- (3) An unbounded example: $S = \{\frac{n}{2} : n \in \mathbb{Z}\}$, $s_{-\infty} = -\infty$, $s_n = \frac{n}{2}$, and $s_{+\infty} = +\infty$.



- (4) One of the two $S = \emptyset$ possibilities: $S = \emptyset$, $s_0 = -\infty$, and $s_1 = +\infty$. This causes \preceq to coincide with \leq .

Remark 1.1.5. It is important to note that the choice which element of S becomes s_0 determines the entire indexing of S and thus the entire partial order \preceq . Additionally, given a set \bar{S} there are exactly two partial orders \preceq possible no matter which element of S is chosen to be s_0 . The two partial orders are opposites of each other.

Remark 1.1.6. From now on, whenever we refer to $A_{\mathbb{R}}$, we are implicitly assuming some S with indexing and \preceq have been set. By ‘the straight descending orientation’ we mean the one where $S = \emptyset$, $s_0 = -\infty$, and $s_1 = +\infty$ as in Example 1.1.4. This is the case where \preceq coincides with \leq .

1.2. Representations of $A_{\mathbb{R}}$: $\text{Rep}_k(A_{\mathbb{R}})$.

Definition 1.2.1. A representation V of $A_{\mathbb{R}}$ is a module over the path algebra $kA_{\mathbb{R}}$. Explicitly, one assigns to each real number x a vector space $V(x)$ and to each pair (x, y) , where $y \preceq x$, a linear transformation $V(x, y) : V(x) \rightarrow V(y)$ such that $V(y, z) \circ V(x, y) = V(x, z)$ whenever such a composition is defined. The support of a representation V is the set of all $x \in \mathbb{R}$ such that $V(x) \neq 0$. We denote the support of a representation V by $\text{supp } V$.

A simple representation at x is a representation V such that $V(x) \cong k$ and if $y \neq x$ then $V(y) = 0$. The linear map $V(x, x)$ is the identity and $V(y, z) = 0$ if $y \neq x$ or $z \neq x$.

Definition 1.2.2. A morphism $f : V \rightarrow W$ of representations of $A_{\mathbb{R}}$ is a morphism of $kA_{\mathbb{R}}$ modules. Explicitly, it is a collection of linear maps $f(x) : V(x) \rightarrow W(x)$, for all $x \in \mathbb{R}$, making the following squares commute for each pair $x, y \in \mathbb{R}$ where $y \preceq x$:

$$\begin{array}{ccc} V(x) & \xrightarrow{V(x,y)} & V(y) \\ f(x) \downarrow & & \downarrow f(y) \\ W(x) & \xrightarrow{W(x,y)} & W(y). \end{array}$$

Since we’re working with modules over an associative algebra, and associative algebras are in particular rings, the category of k -representations of $A_{\mathbb{R}}$, denoted $\text{Rep}_k(A_{\mathbb{R}})$, is abelian.

Propositions 1.2.3 and 1.2.4 can be proved almost the exact same way as they would for discrete quivers of type A.

Proposition 1.2.3. *A morphism of representations $f : V \rightarrow W$ in $\text{Rep}_k(A_{\mathbb{R}})$ is an isomorphism if and only if $f(x)$ is an isomorphism for each $x \in \mathbb{R}$.*

Proposition 1.2.4. *Let V and V' be representations of $A_{\mathbb{R}}$ such that $V \cong V'$. Then $\text{supp } V = \text{supp } V'$.*

1.3. The Subcategories $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ and $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$. In this subsection we define the pointwise finite and bounded subcategories of $\text{Rep}_k(A_{\mathbb{R}})$. We provide examples of representations in each subcategory and highlight the differences between them.

Definition 1.3.1. The category of pointwise finite representations, denoted $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, is the full subcategory of $\text{Rep}_k(A_{\mathbb{R}})$ consisting of representations V such that for all $x \in \mathbb{R}$, $\dim V(x) < \infty$.

The category of bounded representations, denoted $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$, is the full subcategory of $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ whose objects are representations V such that there exists $n \in \mathbb{N}$ and for all $x \in \mathbb{R}$, $\dim V(x) < n$.

It is important to note that the conditions in Definition 1.3.1 are not related to the *support* of any representation. I.e. there exist representations in both $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ and $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$ with unbounded support. Such examples are provided below.

Example 1.3.2. We now give some examples of representations in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ and $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$. Each representation will be over $A_{\mathbb{R}}$ with the straight descending orientation (see Remark 1.1.6).

- (1) We give an example of a representation in $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$ with unbounded support. A representation in $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$ is V :

$$V(x) = \begin{cases} k & x \geq 0 \\ 0 & x < 0 \end{cases} \quad V(x, y) = \begin{cases} 1_k & 0 \leq y \leq x \\ 0 & \text{otherwise} \end{cases}$$

Notice that the *support* of V is unbounded. This is fine. The *dimension* of all the $V(x)$ vector spaces is bounded above by 1.

- (2) We now give an example of an infinite coproduct that is still in $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$. Let $M = \bigoplus M_{\{z\}}$ where for each $z \in \mathbb{R}$, $M_{\{z\}}$ be the following representation of $A_{\mathbb{R}}$:

$$M_{\{z\}}(x) = \begin{cases} k & x = z \\ 0 & \text{otherwise} \end{cases} \quad M_{\{z\}}(x, y) = \begin{cases} 1_k & x = y = z \\ 0 & \text{otherwise} \end{cases}$$

That is, $M_{\{z\}}$ is the simple representation at z , which is in $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$. However, $M = \bigoplus_{z \in \mathbb{R}} M_{\{z\}}$ is also in $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$ since $\dim M(x) = 1$ for all $x \in \mathbb{R}$.

- (3) We now give an example of a representation in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ *but not* in $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$. Let W be the representation of $A_{\mathbb{R}}$ where $W(x)$ is k^n where $n = 0$ if $x < 1$ and n is the largest integer less than or equal to x otherwise. I.e., $W(10.4) = k^{10}$. Let $W(x, y)$ be 0 if $y < 1$ or $x < 1$. Otherwise, $W(x, y)$ is the projection of the first $\dim W(y)$ coordinates of $k^{\dim W(x)}$ using the standard basis. For example, $W(10, 4)$ is the projection of k^{10} onto the first 4 coordinates. While $W(x)$ is finite-dimensional for all $x \in \mathbb{R}$, there is no n such that $\dim W(x) \leq n$ for all $x \in \mathbb{R}$.

Originally, the authors only attempted to prove a version of Theorem 2.4.15 for $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$. However, it was noted that nearly all the proof techniques relied on finite-dimensional vector spaces, not on the dimension of the vector spaces being bounded. In the category $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$ the authors discovered projective indecomposable objects that are not projective in $\text{Rep}_k(A_{\mathbb{R}})$. Further study revealed these objects to also be projective in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. See Section 2.1 for details on these new projective objects. These new projectives in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ are necessary to obtain a category of finitely generated representations (Definition 3.1.3, denoted $\text{rep}_k(A_{\mathbb{R}})$) which has all the reasonable properties one could expect from a continuous version of finitely generated representations.

In contrast to the apparent superiority of $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, the category is simply too big to even have all projective covers. While pathological examples of representations without projective covers can be constructed in both $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ and $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$, the more well-behaved examples of representations without projective covers exist only in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. See Example 2.1.17 in Section 2.1. Such a representation does not exist in $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$ and so this can be considered the first step towards finitely generated representations.

2. CLASSIFICATION OF INDECOMPOSABLE PWF REPRESENTATIONS

In this section we provide a complete classification of indecomposable pointwise finite-dimensional representations of a continuous quiver of type A (Theorem 2.4.15). We focus on representation theoretic techniques and provide a self-contained approach. In particular, we characterize projective indecomposables (Theorems 2.1.3 and 2.1.16) before our decomposition theorem and obtain the completeness of the classification (Remark 2.4.16) as a result. Additionally, our proof of the decomposition theorem is algorithmic. We discuss related results by Botnan and Crawley-Boevey in Section 2.6.

2.1. Projectives. We will construct all pointwise finite-dimensional projective representations in the category $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. There are two types of indecomposable projectives: projectives P_c generated at one point as in Definition 2.1.4, which are quite similar to the projectives for finite quivers. These kinds of projectives are actually projective in both $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ and in $\text{Rep}_k(A_{\mathbb{R}})$. The new kind of projectives $P_{(c)}$ and $P_{(c)}$ will be projectives which have half open intervals as supports as in Definition 2.1.13. These representations are projective in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ but are not projective in $\text{Rep}_k(A_{\mathbb{R}})$ (see Example 2.5.3). We start with the case of a projective generated at one point.

Definition 2.1.1. Given any point $c \in \mathbb{R}$ and any vector space X over k , let $(PX)_c$ be the representation defined as follows.

$$(PX)_c(x) = \begin{cases} X & \text{if } x \preceq c \\ 0 & \text{otherwise} \end{cases}$$

and $(PX)_c(x, y) = \text{id}_X$ if $y \preceq x \preceq c$.

Lemma 2.1.2. For any representation V in $\text{Rep}_k(A_{\mathbb{R}})$ (not necessarily pointwise finite) and any k -vector space X we have:

$$\text{Hom}((PX)_c, V) = \text{Hom}_k(X, V(c)),$$

i.e., the functor which takes X to $(PX)_c$ is left adjoint to the evaluation functor $V \mapsto V(c)$.

Proof. Given any morphism $f : (PX)_c \rightarrow V$, let $f_c : (PX)_c(c) = X \rightarrow V(c)$ be the restriction of f to the point c . Then, for any $x \preceq c$, the commutativity of the diagram:

$$\begin{array}{ccc} (PX)_c(c) = X & \xrightarrow{\text{id}_X} & X = (PX)_c(x) \\ f_c \downarrow & & \downarrow f_x \\ V(c) & \xrightarrow{V(c, x)} & V(x) \end{array}$$

forces the map $f_x : (PX)_c(x) \rightarrow X(x)$ to be equal to $V(c, x) \circ f_c$. Conversely, any linear map $g : X \rightarrow V(c)$ extends to a morphism $\bar{g} : (PX)_c \rightarrow V$ by the same formula ($\bar{g}(x) = V(c, x) \circ g : (PX)_c(x) = X \rightarrow X(x)$). \square

Theorem 2.1.3. For any vector space X and any $c \in \mathbb{R}$, the representation $(PX)_c$ is projective in $\text{Rep}_k(A_{\mathbb{R}})$.

Proof. Let $p : V \rightarrow W$ be an epimorphism and let $f : (PX)_c \rightarrow W$ be any morphism. Then $p_c : V(c) \rightarrow W(c)$ is an epimorphism. So, the linear map $f_c : X \rightarrow W(c)$ lifts to a map $g : X \rightarrow V(c)$ which, by Lemma 2.1.2, extends to a morphism $\bar{g} : (PX)_c \rightarrow V$. Since $p \circ \bar{g}$ and $f : (PX)_c \rightarrow W$ agree at c , they are equal by Lemma 2.1.2. So, $(PX)_c$ is projective. \square

Note that $(PX)_c$ is indecomposable if and only if X is one-dimensional as it is in the following definition. In this case the indecomposable projective is denoted simply by P_c .

Definition 2.1.4. Let $c \in \mathbb{R}$. Define the representation P_c in $\text{Rep}_k(A_{\mathbb{R}})$ as:

$$P_c(x) = \begin{cases} k & \text{if } x \preceq c \\ 0 & \text{otherwise} \end{cases}, \quad P_c(x, y) = \begin{cases} 1_k & \text{if } y \preceq x \preceq c \\ 0 & \text{otherwise} \end{cases}$$

The rest of this subsection is devoted to the construction of all pointwise finite-dimensional projective representation, including objects $P_{(a)}$ and $P_{(b)}$ for $s_{2n-1} < a < s_{2n} < b < s_{2n+1}$ with supports $(a, s_{2n}]$ and $[s_{2n}, b)$ respectively.

In order to describe these new types of projective representations in the category of pointwise finite-dimensional representations of $A_{\mathbb{R}}$ we need to set up notation of “image filtration” (Definition 2.1.5) and “support intervals” (Definition 2.1.7). Recall s_n is a sink if n is even and a source if n is odd.

Definition 2.1.5. Let V be a pointwise finite-dimensional representation of $A_{\mathbb{R}}$ such that $\text{supp } V \subset [s_0, s_1]$. (If $s_0 = -\infty$ or $s_1 = +\infty$ then $\text{supp } V$ is a subset of $(s_0, s_1]$, $[s_0, s_1)$, or (s_0, s_1) , whichever applies.) Let $b = s_0$ or $b \in (s_0, s_1)$.

Let b be the greatest lower bound of $\text{supp } V$. When $b \in \text{supp } V$ we set $V^{\bullet}(b) = V(b)$. The image filtration of $V^{\bullet}(b)$ is the set of distinct subspaces of the form $V(x, b)(V(x))$. Let $V^{\circ}(b)$ be the colimit of the vector spaces $V(x)$, for $b < x < s_1$, with the linear maps $V(x, y)$, for $b < y \leq x < s_1$. Since each $V(x)$ is finite-dimensional, $V^{\circ}(b)$ is at most countably infinite dimensional. Denote by $V^{\circ}(x, b)$ the colimit linear map from $V(x)$ to $V^{\circ}(b)$. The image filtration of $V^{\circ}(b)$ is the set of distinct (finite-dimensional) subspaces of the form $V^{\circ}(x, b)(V(x))$.

When $b \in \text{supp } V$ we take I to be $[b, c]$ or $[b, c)$. When $b \notin \text{supp } V$ we take I to be $(b, c]$ or (b, c) . For all such I and when $b \in \text{supp } V$, let

$$(1) \quad V_I^{\bullet}(b) := \bigcap_{x \in I} V(x, b)(V(x)) \subset V^{\bullet}(b)$$

Whether or not $b \in \text{supp } V$, we let

$$(2) \quad V_{I \setminus \{b\}}^{\circ}(b) := \bigcap_{x \in I \setminus \{b\}} V^{\circ}(x, b)(V(x)) \subset V^{\circ}(b).$$

Then $V_I^{\bullet}(b)$ and $V_{I \setminus \{b\}}^{\circ}(b)$ are members of the image filtrations of $V^{\bullet}(b)$ and $V^{\circ}(b)$, respectively. In particular, there exists x_0 in I such that $V_I^{\bullet}(b) = V(x_0, b)(V(x_0))$ or $x_0 \in I \setminus b$ such that $V_{I \setminus \{b\}}^{\circ}(b) = V^{\circ}(x_0, b)(V(x_0))$. Whenever $b \in \text{supp } V$, $V^{\bullet}(b)$ is finite-dimensional and so the image filtration is finite. Since $V^{\circ}(b)$ may not be finite-dimensional and the dimension of the vector spaces $V(x)$ are not bounded the filtration on $V^{\circ}(b)$ may be infinite but still countable with a minimal term. In fact, $V(s_1, b)(b)$ and $V^{\circ}(s_1, b)$ are the minimal objects in the filtrations of $V^{\bullet}(b)$ and $V^{\circ}(b)$, respectively.

Remark 2.1.6. (a) If $I = [b, c]$ ($c \in I$) then

$$\begin{aligned} V_I^{\bullet}(b) &= V_{[b, c]}^{\bullet}(b) = V(c, b)(V(c)) \\ V_{I \setminus \{b\}}^{\circ}(b) &= V_{[b, c]}^{\circ}(b) = V^{\circ}(c, b)(V(c)). \end{aligned}$$

(b) If $I = [b, c)$ ($c \notin I$) then whenever $c \in \mathbb{R}$ we have

$$\begin{aligned} V_I^{\bullet}(b) &= V_{[b, c)}^{\bullet}(b) \supset V(c, b)(V(c)) \text{ and} \\ V_{I \setminus \{b\}}^{\circ}(b) &= V_{(b, c)}^{\circ}(b) \supset V^{\circ}(c, b)(V(c)) \end{aligned}$$

but in both cases the subspaces may be different.

- (c) For any $x < z$ in I , the term $V(x, b)(V(x))$ is redundant in the intersection (1) since $V(x, b)(V(x)) \supset V(z, b)(V(z))$. Thus,

$$V_I^\bullet(b) = \bigcap_{x \in I} V(x, b)(V(x)) = \bigcap_{y \in I, y \geq z} V(z, b)V(y, z)(V(y)) = V(z, b)V_{I \cap [z, s_1]}^\bullet(z)$$

- (c') For any $x < z$ in I , the term $V^\circ(x, b)(V(x))$ is redundant in the intersection (2) since $V^\circ(x, b)(V(x)) \supset V^\circ(z, b)(V(z))$. Thus,

$$V_I^\circ(b) = \bigcap_{x \in I} V^\circ(x, b)(V(x)) = \bigcap_{y \in I, y \geq z} V^\circ(z, b)V^\circ(y, z)(V(y)) = V^\circ(z, b)V_{I \cap [z, s_1]}^\circ(z)$$

Definition 2.1.7. Let $W \subset V^\bullet(b)$ be a subspace. Define I_W as:

$$I_W = \{x \geq b \in \text{supp } V \mid W \subset V(x, b)(V(x))\}$$

Such $\{I_W\}$ are called support intervals for $V^\bullet(b)$.

Let $W \subset V^\circ(b)$ be a finite-dimensional subspace. Then we define I_W similarly for $b \notin I$:

$$I_W = \{x > b \in \text{supp } V \mid W \subset V^\circ(x, b)(V(x))\}$$

These $\{I_W\}$ are also called support intervals for $V^\circ(b)$.

Proposition 2.1.8. (a) *There is a 1-1 correspondence between support intervals for $V^\bullet(b)$ and the terms in the image filtration of $V^\bullet(b)$ given by $I \mapsto V_I^\bullet \subset V^\bullet(b)$ and $W \mapsto I_W$.*

- (a') *There is also a 1-1 correspondence between the support intervals for $V^\circ(b)$ and the terms in the image filtration of $V^\circ(b)$ given by $I \mapsto V_I^\circ \subset V^\circ(b)$ and $W \mapsto I_W$.*

Proof. We first prove (a). If $W = V(x_0, b)(V(x_0))$ then I_W contains x_0 and $V_{I_W}^\bullet$ is the intersection of $V(x_0, b)(V(x_0)) = W$ and the subspaces $V(x, b)(V(x))$ which all contains W by definition. So, $V_{I_W}^\bullet = W$.

If $I = I_W$ then $W \subset V_I^\bullet(b)$ by definition. If $V(x, b)(V(x))$ contains $V_I^\bullet(b)$, it contains W . So, $I_{V_I^\bullet(b)} \subset I_W = I$. But $I \subset I_{V_I^\bullet(b)}$. So, $I = I_{V_I^\bullet(b)}$ for any support interval I .

The proof of (a) as stated works for (a') if we replace V_I^\bullet with V_I° . \square

Remark 2.1.9. The image filtration of $V^\bullet(b)$ can be written:

$$V^\bullet(b) \supsetneq V(x_n, b)(V(x_n)) \supsetneq V(x_{n-1}, b)(V(x_{n-1})) \supsetneq \cdots \supsetneq V(x_1, b)(V(x_1)).$$

By Proposition Proposition 2.1.8, we see this is actually a filtration

$$(\bullet*) \quad V^\bullet(b) \supsetneq V_{I_n}^\bullet \supsetneq V_{I_{n-1}}^\bullet \supsetneq \cdots \supsetneq V_{I_1}^\bullet.$$

where each x_i in the first form is an element of I_i in the second form.

For the image filtration of $V^\circ(b)$, we have the following equivalent forms, where each x_i in the first form is an element of I_i in the second form:

$$V^\circ(b) \cdots \supsetneq V^\circ(x_n, b)(V(x_n)) \supsetneq V^\circ(x_{n-1}, b)(V(x_{n-1})) \supsetneq \cdots \supsetneq V^\circ(x_1, b)(V(x_1)).$$

By Proposition 2.1.8, we see this is actually a filtration

$$(\circ*) \quad V^\circ(b) \cdots \supsetneq V_{I_n}^\circ \supsetneq V_{I_{n-1}}^\circ \supsetneq \cdots \supsetneq V_{I_1}^\circ.$$

Lemma 2.1.10. (a) *Let $W' \subsetneq W$ be consecutive terms in the image filtration $(\bullet*)$ of $V^\bullet(b)$. Then $I_W \subsetneq I_{W'}$ and, for any element $x \in (I_{W'} - I_W)$, we have $V(x, b)(V(x)) = W'$ in $V^\bullet(b)$.*

- (a') *Let $W' \subsetneq W$ be consecutive terms in the image filtration $(\circ*)$ of $V^\circ(b)$. Then $I_W \subsetneq I_{W'}$ and, for any element $x \in (I_{W'} - I_W)$, we have $V^\circ(x, b)(V(x)) = W'$ in $V^\circ(b)$.*

Proof. We prove (a) first. Since $x \notin I_W$ it follows from Remark 2.1.6(c) that W is not a subset of $V(x, b)(V(x))$ in $V^\bullet(b)$. Since $x \in I_{W'}$ it follows that $W' \subset V(x, b)(V(x))$. Thus $W' \subset V(x, b)(V(x)) \subsetneq W$. Since W, W' are consecutive in the image filtration, $V(x, b)(V(x)) = W'$ as claimed. The proof of (a) as stated works for (a') by replacing V_I^\bullet with V_I° . \square

Lemma 2.1.11. *Let $W \subset V^\bullet(b)$ (resp. $W \subset V^\circ(b)$) be a finite-dimensional subspace and let $x_1 < x_2 \in I_W$. Let $I_1 = I_W \cap [x_1, s_1]$ and $I_2 = I_W \cap [x_2, s_1]$. (If $s_1 = +\infty$ then, for each i , $I_i = I_W \cap [x_i, s_1)$). Then $V_{I_1}^\bullet(x_1) \subset V(x_2, x_1)(V_{I_2}^\bullet(x_2))$ (resp. $V_{I_1}^\circ(x_1) \subset V(x_2, x_1)(V_{I_2}^\circ(x_2))$).*

Proof. This is a special case of Remarks 2.1.6(c) and 2.1.6(c'). \square

Lemma 2.1.12.

- (a) *For any interval I of the form $[b, c]$ or $(b, c]$ and any element $v \in V_I^\bullet(b)$, there is a collection of elements $\{V_x^\bullet \in V(x)\}_{x \in I}$ so that $V_b^\bullet = v$ and $V(y, x)(V_y^\bullet) = V_x^\bullet$ for all $b \leq x \leq y \in I$.*
- (a') *For any interval I of the form $(b, c]$ or (b, c) and any element $v \in V_I^\circ(b)$, there is a collection of elements $\{v_x \in V(x)\}_{x \in I}$ so that $v_b = v$ and $V(y, x)(v_y) = v_x$ and $V^\circ(x, b)(v_x) = v_b$ for all $b < x \leq y \in I$.*

Proof. If $I = [b, c]$ then $V_I^\bullet(b) = V(c, b)(V(c))$ and we can choose one element $w \in V^\bullet(c)$ so that $V(c, b)(w) = v$ and let $v_x = V(c, x)(w)$ for all $x \in [b, c]$. Similarly, if $I = (b, c]$ then $V_I^\circ(b) = V^\circ(c, b)(V(c))$ and we make a similar choice.

Otherwise, $I = [b, c)$ or $I = (b, c)$ for some $c > b$. In this case, choose an increasing sequence of real numbers $b < x_1 < x_2 < x_3 < \dots$ in (b, c) converging to c . Let $J_0 = I$, for each $l > 0$, let $J_l = I \cap [x_l, c)$ and additionally, let $v_l \in V_{J_l}^\bullet(x_l)$ (resp. $v_l \in V_{J_l}^\circ(x_l)$) be chosen recursively as follows.

- (1) Set $v_0 = v \in V_I^\bullet(b)$, (resp. $v_0 = v \in V_I^\circ(b)$).
- (2) Given v_k in $V_{J_k}^\bullet(x_k)$ (resp. in $V_{J_k}^\circ(x_k)$), by Lemma 2.1.11, there exists v_{l+1} in $V_{J_{l+1}}^\bullet(x_{l+1})$ (resp. $V_{J_{l+1}}^\circ(x_{l+1})$) so that $V(x_{k+1}, x_k)(v_{k+1}) = v_k$ (resp. $V^\circ(x_{l+1}, x_l)(v_{l+1}) = v_k$).

After this sequence of elements v_k is chosen, the vector v_x for any $x \in I$ is given by $v_x = V(x_l, x)(v_k)$ for any $x_l > x$. This is well defined by condition (2) in the case of $V^\bullet(b)$ and by condition (2) combined with the universal property of $V^\circ(b)$ in that case. \square

Definition 2.1.13. Let s_0 be a sink or $-\infty$ and let $s_1 > s_0$ be the next source or $+\infty$. Let $s_0 < a < s_1$. For $I = [s_0, a]$ or $[s_0, a)$ let P_I , also written $P_a = P_{[s_0, a]}$ or $P_a = P_{[s_0, a)}$, denote the representation with support I so that $P_I(x)$ is one-dimensional with generator v_x for all $x \in I$ and $P(y, x)(v_y) = v_x$ for all $x < y \in I$.

For $a = s_1$, define P_a as before. However, when $a = s_1$, P_a is not defined this way. If $s_0 = -\infty$ then P_a and P_a are instead $P_{(s_0, a]}$ and $P_{(s_0, a)}$, respectively.

Proposition 2.1.14. *P_a and P_a as in Definition 2.1.13 are projective in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$.*

Proof. We first assume that $s_0 \in \mathbb{R}$. To show that P_I is projective it suffices to show that any epimorphism $p : E \rightarrow P_I$ has a section. Let $W \subset E^\bullet(s_0)$ be the smallest term in the image filtration of $E^\bullet(s_0)$ which maps onto $P_I^\bullet(s_0)$.

Claim: I_W contains I and thus $W \subset E_I^\bullet(s_0)$.

Proof: For each $x \in I$, there is a $w \in E(x)$ so that $p_x(w) = v_x \in P_I(x)$. But then $p_{s_0} E^\bullet(x, s_0)(w) = P_I^\bullet(x, s_0)(v_x) = v_{s_0} \neq 0$. So, $W \subset E^\bullet(x, s_0)(E(x))$ which implies $x \in I_W$. Since this holds for all $x \in I$ we get that $I \subset I_W$.

By construction of W , there is a $w \in W \subset E_I^\bullet(s_0)$ so that $p(w) = v_{s_0}$. By Lemma 2.1.12 there are elements $w_x \in E(x)$ for all $x \in I$ so that $E(x, y)(w_x) = w_y$ for all $y \leq x \in I$. Then, a section $s : P \rightarrow E$ is given by $s(v_x) = w_x$ for all $x \in I$.

If we instead assume $s_0 = -\infty$ then above we replace E^\bullet with E° and P^\bullet with P° where appropriate. By the universal property of colimits, the map on representations induces a map $E^\circ(-\infty) \rightarrow P_I^\circ(-\infty)$. Then the rest of the proof holds as stated. \square

Lemma 2.1.15. *Suppose $s_0 = -\infty$ and P is a pointwise finite-dimensional representation with support (s_0, a) , for $a \leq s_1$, or $(s_0, a]$, for $a < s_1$. Either $P^\circ(s_0)$ is finite-dimensional or $P^\circ(s_0) \cong k^\infty$.*

Proof. Since P is pointwise finite-dimensional, if the dimension of $P(x)$ is bounded by some n for all $x \in \mathbb{R}$ then $P^\circ(s_0)$ is also bounded by n .

Now suppose $P^\circ(s_0)$ is not finite-dimensional. For each $i > 0$, let $n_i = \dim P_{I_i}^\circ$. Let $e_i \in k^\infty$ denote the unit vector with a 1 in the i th coordinate. For a choice of basis of $P^\circ(s_0)$, we note that since each morphism is a monomorphism and the image filtration $(\circ*)$ of $P^\circ(s_0)$ has a minimal element, we may inductively choose a basis on $P^\circ(s_0)$. We do this by first choosing a basis of $P_{I_1}^\circ$, then completing it to a basis of $P_{I_2}^\circ$ and so on. Since each $P_{I_i}^\circ$ is finite-dimensional this is well defined.

Since we have a consistent choice of bases, map the chosen basis of each $P_{I_i}^\circ$ to the collection $\{e_i\} \subset k^\infty$ in a consistent way. Since each $P_{I_i}^\circ \cong P(x_i)$ this induces a map $P^\circ(s_0) \rightarrow k^\infty$.

To see the map is surjective take any element w of k^∞ ; w has finitely many nonzero coordinates. Thus it is some linear combination of finitely many e_j 's. Then there is a $P_{I_i}^\circ$ whose basis contains enough elements to surject on to the e_j 's. Thus there is an element v in $P_{I_i}^\circ$ such that $v \mapsto w$ and so there is an element $\tilde{v} \in P^\circ(s_0)$ that maps to w . The map is injective since if $\tilde{v} \neq \tilde{v}'$ in $P^\circ(s_0)$ then there is a pair $v \neq v'$ in a $P_{I_i}^\circ$ such that $v \mapsto \tilde{v}$ and $v' \mapsto \tilde{v}'$. We know v and v' map to different elements in k^∞ so \tilde{v} and \tilde{v}' must also. Therefore, $P^\circ(s_0) \cong k^\infty$. \square

The following theorem will give a characterization of one sided projective objects in $\text{Rep}_k^{\text{pwf}}(A_\mathbb{R})$.

Theorem 2.1.16. *Let $s_0 \leq a < s_1$ with s_0 a sink and s_1 the next source. Let P be a pointwise finite-dimensional representation of $A_\mathbb{R}$ with $\text{supp } P \subset [s_0, a]$.*

- (1) *Then P is projective in $\text{Rep}_k^{\text{pwf}}(A_\mathbb{R})$ if and only if all maps $P(x, s_0) : P(x) \rightarrow P(s_0)$ are injective for all $x \in \text{supp } P$.*
- (2) *Every projective representation in $\text{Rep}_k^{\text{pwf}}(A_\mathbb{R})$ with support in $[s_0, a]$ is a finite direct sum of representations of the forms P_b and P_b for $s_0 \leq b \leq a$.*
- (2') *Every projective representation in $\text{Rep}_k^{\text{pwf}}(A_\mathbb{R})$ with support in $(s_0, a]$ (i.e., $s_0 = -\infty$) is a possibly infinite direct sum of representations of the forms P_b and P_b for $s_0 < b \leq a$.*

Proof. When $a = s_0$, statements (1) are (2) are trivially true and statement (2') does not apply.

(1) Suppose that there is some $x_0 \in [b, a] \subset [s_0, a]$ so that $P(x_0, s_0) : P(x_0) \rightarrow P(s_0)$ is not injective. Then we will show that P is not projective. Indeed consider the quotient object Q given $Q(x) = P(x)$ for all $x \geq x_0$ and $Q(x) = 0$ for all $x < x_0$. We have an epimorphism $\pi : P \rightarrow Q$. Let \tilde{Q} be the representation given by $\tilde{Q}(x) = P(x)$ for $x \geq x_0$ and $\tilde{Q}(x) = P(x_0)$ for all $x \leq x_0$ with $\tilde{Q}(y, x) = \text{Id}$, the identity, when $x, y \leq x_0$ and $\tilde{Q}(y, x) = P(y, x_0)$ when $x \leq x_0 < y$. Let $p : \tilde{Q} \rightarrow Q$ be the projection map. Claim: the quotient map $\pi : P \rightarrow Q$ does not lift to \tilde{Q} , i.e. there is no $\gamma : P \rightarrow \tilde{Q}$ such that $p \circ \gamma = \pi$. Proof of claim: Since $\pi_{x_0} = \text{Id} : P(x_0) = Q(x_0)$ and $p_{x_0} = \text{Id} : \tilde{Q}(x_0) \rightarrow Q(x_0)$ we would have $\gamma_{x_0} = \text{Id}$. But that gives a contradiction to the basic property of maps between representations: $\gamma_{s_0} \circ P(x_0, s_0) = \tilde{Q}(x_0, s_0) \circ \gamma_{x_0} = \text{Id}$, but $\gamma_{s_0} \circ P(x_0, s_0)$ is not injective by assumption. Therefore, P is not projective.

Conversely, suppose that all morphisms $P(x, s_0)$ are monomorphisms. Choose a basis B for $P(s_0)$ compatible with the image filtration. Thus, a subset B_i of B is a basis for each subspace $P_{J_i}(s_0)$ in the image filtration of $P(s_0)$ where $J_i = I_{P_{J_i}(s_0)}$ are ordered by inclusion: $J_1 \subsetneq J_2 \subsetneq \cdots \subsetneq J_n$. Then $P_{J_1}(s_0) \supsetneq P_{J_2}(s_0) \supsetneq \cdots \supsetneq P_{J_n}(s_0)$.

(2) By Lemma 2.1.12, every $v \in B_i - B_{i+1}$ lifts to a compatible system of elements $v_x \in P(x)$ for all $x \in J_i$. By Lemma 2.1.10, $P(x, s_0)(P(x)) = P_{J_i}(s_0)$ for all $x \in J_i - J_{i-1}$ (where $J_0 = \emptyset$). Since $P(x, s_0)$ is a monomorphism, the liftings $v_x \in P(x)$ for all $v \in B_i$ form a basis for $P(x)$

for all $x \in J_i - J_{i-1}$. For each $v \in B_i$, the lifting v_x of v generate a pointwise one-dimensional subrepresentation Q_v of P with support in J_i , i.e. P_v has the form P_b or $P_{\bar{b}}$ depending on whether $J_i = [s_0, b]$ or $[s_0, \bar{b})$. The Q_v , for $v \in B$, are disjoint and generate all of $P(x)$ for every $x \in [s_0, a]$. Thus P is a direct sum of the Q_v as claimed.

Below is an example of such a decomposition for (2).

$$\begin{aligned} [s_0, b_1) &= \text{supp } Q_{v_1} \bullet \circ \\ [s_0, b_2] &= \text{supp } Q_{v_2} \bullet \text{---} \bullet \\ [s_0, b_3] &= \text{supp } Q_{v_3} \bullet \text{---} \bullet \\ [s_0, b_4) &= \text{supp } Q_{v_4} \bullet \text{---} \circ \\ [s_0, b_5] &= \text{supp } Q_{v_5} \bullet \text{---} \bullet \end{aligned}$$

With the exception of choosing a basis, we may apply all of the argument for statement (2) to statement (2'). If $P^\circ(s_0)$ is finite-dimensional we get a basis and apply the argument for (2). By Lemma 2.1.15, if $P^\circ(s_0)$ is infinite-dimensional then it is isomorphic to k^∞ and by the proof of the same lemma we have a basis that respects the filtration. We then apply the argument for (2). \square

Example 2.1.17. Let $A_{\mathbb{R}}$ have the straight descending orientation and for each positive integer n let V_n be the following representation:

$$V_n(x) = \begin{cases} k & n \leq x \\ 0 & \text{otherwise} \end{cases} \quad V_n(x, y) = \begin{cases} 1_k & n \leq y \leq x \\ 0 & \text{otherwise} \end{cases}$$

Using Theorem 2.1.16 we see that the projective cover of each V_n is the projective indecomposable with support $\mathbb{R} = (-\infty, +\infty)$.

Note that $V = \bigoplus V_n$ is still pointwise finite. One can check it is isomorphic to the representation W (item (3) in Example 1.3.2). However, the projective cover is infinitely many copies of the indecomposable projective with support $(-\infty, +\infty)$, which is not pointwise finite-dimensional. Therefore, this rather tame example does not have a projective cover in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$.

However, the dually constructed representation V' (each V'_n has support $(-\infty, n]$) is its own projective cover by Theorem 2.1.16 and so does have a projective cover. While V and V' exist in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, neither exists in $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$. So, this type of asymmetry does not happen in $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$.

2.2. Sufficient Conditions for Indecomposables. Here we give sufficient conditions for pointwise finite-dimensional representations to be indecomposable. In Section 2.4 we will show that these conditions are also necessary.

Proposition 2.2.1. *Let V be a representation of $A_{\mathbb{R}}$ such that*

- (1) $\dim V(x) \leq 1$ for all $x \in \mathbb{R}$,
- (2) if $V(x) \neq 0 \neq V(z)$ and $x \leq y \leq z$ in \mathbb{R} then $V(y) \neq 0$, and
- (3) if $V(x) \neq 0 \neq V(y)$ and $x \preceq y$ then $V(y, x)$ is an isomorphism.

Then V is indecomposable.

Proof. Suppose V is not indecomposable for contradiction. Then $V \cong W_1 \oplus W_2$ with $W_1 \neq 0 \neq W_2$. Then $\text{supp } W_1 \cap \text{supp } W_2 = \emptyset$ and $\text{supp } W_1 \cup \text{supp } W_2 = \text{supp } V$. Since $W_1, W_2 \neq 0$ there exist $x_1 \in \text{supp } W_1$ and $x_2 \in \text{supp } W_2$. By symmetry we may assume $x_1 < x_2$.

Claim 1: There are only finitely many elements of S in the open interval (x_1, x_2) .

Pf: This follows from the fact that (x_1, x_2) is a bounded interval, i.e., $[x_1, x_2]$ is compact. If $S \cap [x_1, x_2]$ were infinite, it would contain a converging sequence (any infinite subset of a compact set contains a converging sequence). By definition, S does not contain a converging sequence. So, $S \cap [x_1, x_2]$ is finite. A fortiori, $S \cap (x_1, x_2)$ is finite.

Claim 2: There exist $x_1 \in \text{supp } W_1$ and $x_2 \in \text{supp } W_2$ such that $S \cap (x_1, x_2) = \emptyset$.

Pf: Let $n = \# \{S \cap (x_1, x_2)\}$. If $n \geq 1$ we will find another pair $x'_1 < x'_2$ in the respective supports of W_1, W_2 so that $\# \{S \cap (x'_1, x'_2)\} < \# \{S \cap (x_1, x_2)\} = n$. This will imply that $n = 0$.

To find this second pair x'_1, x'_2 choose any element s_k in $S \cap (x_1, x_2)$ which is nonempty by assumption that $n \geq 1$. Then s_k is in the support of W_1 or W_2 . In the first case, $x'_1 = s_k, x'_2 = x_2$ gives the desired pair. Indeed, in this case, $S \cap (x'_1, x'_2) \subset S \cap (x_1, x_2)$ since $s_k \in S \cap (x_1, x_2)$ but $s_k \notin S \cap (x'_1, x'_2)$. Also, $x'_1 = s_k$ is in the support of W_1 by assumption and $x'_2 = x_2$ is in the support of W_2 . The second case is similar. In both cases, the value of n can be reduced if it is positive. So, the minimal value of n is 0.

By Claim 2 we may assume there are no elements of S between x_1 and x_2 , i.e. the \prec orientation of \mathbb{R} is constant in the closed interval $[x_1, x_2]$ and either $V(x_2, x_1)$ or $V(x_1, x_2)$ is an isomorphism. In the first case, we consider the projection $V \rightarrow W_1$ and in the second case we consider the other projection $f : V \rightarrow W_2$. By symmetry, we may take the first case, i.e. $V(x_2, x_1)$ is an isomorphism. Then we have the following commuting diagram:

$$\begin{array}{ccc} V(x_1) & \xleftarrow[V(x_2, x_1)]{\cong} & V(x_2) \neq 0 \\ \cong \downarrow f_{x_1} & & \downarrow f_{x_2} \\ W_1(x_1) & \longleftarrow & W_1(x_2) = 0 \end{array}$$

Since $x_1 \in \text{supp } W_1$, it follows that $f_{x_1} : V(x_1) \rightarrow W_1(x_1)$ is an isomorphism. But $W_1(x_2) = 0$ since $x_2 \in \text{supp } W_2$ and $x_2 \notin \text{supp } W_1$. The commutativity of the diagram then gives a contradiction. Thus V is indecomposable. \square

Definition 2.2.2. For any interval I in \mathbb{R} let M_I be the representation of $A_{\mathbb{R}}$ given as follows.

$$M_I(x) = \begin{cases} k & x \in I \\ 0 & \text{otherwise} \end{cases} \quad M_I(x, y) = \begin{cases} 1_k & y \preceq x \text{ and } x, y \in I \\ 0 & \text{otherwise} \end{cases}$$

The conditions of Proposition 2.2.1 are satisfied immediately. So, M_I is indecomposable. If a representation $V \cong M_I$ we call V an interval indecomposable or interval indecomposable representation.

Corollary 2.2.3. Let V be an indecomposable representation which is pointwise one-dimensional (satisfies the conditions of Proposition 2.2.1). Let $J \subseteq \text{supp } V$ be a connected subset and let V_J be the restriction of V to J , i.e. $V_J(x) = V(x) = k$ for all $x \in J$ and $V_J(x) = 0$ for all $x \notin J$, and $V_J(y, x) = V(y, x)$ for all $x, y \in J$. Then V_J is an indecomposable representation.

Proof. We will show that conditions (1), (2) and (3) of Proposition 2.2.1 are satisfied by the representation V_J .

(1) By definition of V_J it follows that $\dim_k V_J(x) \leq 1$.

(2) This follows since J is connected subset of $\text{supp } V$.

(3) Suppose there is $x \preceq y$ with $x, y \in J$ such that $V_J(y, x) = V(y, x)$ not an isomorphism. Since $\dim_k V_J(x) \leq 1$, this is equivalent to $V_J(y, x) = 0$.

Let $I = \{t \mid x \prec t \preceq y \text{ such that } V(y, t) \neq 0\}$. Then $x \in (\text{supp } V) \setminus I = J_1 \cup J_2$ where $J_1 \cap J_2 = \emptyset$ and we may assume $x \in J_1$. Then V_{J_1} is a subrepresentation of V but is also a quotient of V since the map $\pi : V \rightarrow V_{J_1}$ defined as $\pi_x = \text{Id}_{V_J(x)}$ for $x \in J_1$ and $\pi_x = 0$ for $x \notin J_1$ is a representation homomorphism using the fact that $V(t_2, t_1) = 0$ for all $t_2 \in (\text{supp } V) \setminus J_1$ and all $t_1 \in J_1$. Actually π is a splitting for the inclusion $V_{J_1} \rightarrow V$, contradicting the assumption that V is indecomposable. Therefore (3) holds for V_J .

So by Proposition 2.2.1 it follows that V_J is indecomposable. \square

2.3. Filtrations. In this section will provide some lemmas necessary for Section 2.4. In both this section and in Section 2.4 we will be using notation $\text{Hom}(-, -)$ for $\text{Hom}_{\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})}(-, -)$ and $\text{End}(-)$

for $\text{End}_{\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})}(-)$ where $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ is the full subcategory of $\text{Rep}_k(A_{\mathbb{R}})$ whose objects are all pointwise finite representations of $A_{\mathbb{R}}$.

Lemma 2.3.1. *Let V be an indecomposable pointwise one-dimensional representation. Then the endomorphism ring of V is the field k .*

Proof. Let $x_0 \in \text{supp } V$. By definition, $V(x_0) \cong k$. Choose a morphism $f(x_0) : V(x_0) \rightarrow V(x_0)$.

Claim: if $f(x_0) \neq 0$ this determines an isomorphism $V \xrightarrow{\cong} V$.

Since $V(x_0) \cong k$, $f(x_0)$ is an isomorphism. If $y \in \text{supp } V$ such that $y \preceq x_0$ then $V(x_0, y)$ is an isomorphism. So, for all $y \preceq x_0$ in $\text{supp } V$, define

$$f(y) := V(x_0, y) \circ f(x_0) \circ (V(x_0, y))^{-1}.$$

Dually, for all $y \in \text{supp } V$ such that $x_0 \preceq y$ define

$$f(y) := (V(y, x_0))^{-1} \circ f(x_0) \circ V(y, x_0).$$

If there are no sinks and sources in $\text{supp } V$, except possibly the endpoints, we have an induced morphism $V \rightarrow V$ such that $f(x)$ is an isomorphism for all $x \in \mathbb{R}$ (by setting $f(x) = 0$ when $x \notin \text{supp } V$). By Proposition 1.2.3 f is an isomorphism. Now, suppose there is a sink or source in the interior of $\text{supp } V$.

Let s_n be a source such that $x_0 \preceq s_n$. By the paragraph above we already have $f(s_n)$. For each $y \preceq s_n$ for which we do not yet have an $f(y)$ we can use the technique above and define it without making choices. By a dual argument if $s_n \preceq x_0$ we can define $f(y)$ for all y such that $s_n \preceq y$. Note that between any real number x and x_0 there are only finitely many sinks and sources between x and x_0 in the total order of \mathbb{R} . By repeated use of this technique, we have an induced isomorphism $f(x) : V(x) \xrightarrow{\cong} V(x)$ for all $x \in \mathbb{R}$. Thus, we have an induced isomorphism $f : V \xrightarrow{\cong} V$. If $g : V \rightarrow V$ is a nonzero morphism then $g(x)$ is nonzero as before is an isomorphism that determines the rest of g . Then $g(x)$ and $f(x)$ are multiplication by nonzero scalars and there exists $t \in k$ such that $tg(x) = f(x)$. Therefore, $\text{End}(V) \cong k$. \square

Theorem 2.3.2. *Let V and V' be two indecomposable pointwise one-dimensional representations of $A_{\mathbb{R}}$. Then $\text{supp } V = \text{supp } V'$ if and only if $V \cong V'$.*

Proof. We first assume $\text{supp } V = \text{supp } V'$. Let $x_0 \in \text{supp } V = \text{supp } V'$. By definition, $V(x_0) \cong k \cong V'(x_0)$. Choose an isomorphism $f(x_0) : V(x_0) \xrightarrow{\cong} V'(x_0)$ and apply the argument from Lemma 2.3.1. The reverse direction is a special case of Proposition 1.2.4. \square

Definition 2.3.3. Let $X_1 \subset X_2 \subset \cdots \subset X_n$ be a filtration of a vector space $X = X_n$. A basis B for X is said to respect the filtration if $B \cap X_j$ is a basis for X_j for each j . A direct sum decomposition $X = \bigoplus Y_i$ of X is said to respect the filtration if each X_j is a direct sum of some of the Y_i .

Lemma 2.3.4. *For any $b \in \mathbb{R}$, let \mathcal{V}_b be the full subcategory of $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ whose objects are interval indecomposables V with $b \in \text{supp}(V) \subset [b, \infty)$. Let $\mathcal{W}_b := \text{add } \mathcal{V}_b$. Then:*

- (1) *The restriction map, $\text{res} : \mathcal{V}_b \rightarrow \text{Rep}_k(\{b\})$ given by $\text{res}(V) = V(b)$ and $\text{res}(f) = f(b)$ defines a monomorphism $\text{Hom}_{\mathcal{V}_b}(V, V') \rightarrow \text{Hom}_k(V(b), V'(b))$ for all $V, V' \in \mathcal{V}_b$, i.e. restriction to b is a faithful functor on \mathcal{V}_b .*
- (2) *The restriction map, $\text{res} : \mathcal{W}_b \rightarrow \text{Rep}_k(\{b\})$ is also a faithful functor on \mathcal{W}_b .*
- (3) *There is a unique total ordering on the set of isomorphism classes of objects of \mathcal{V}_b so that:*
 - (a) $\text{Hom}_{\mathcal{V}_b}(V', V) = 0$ and $\dim_k \text{Hom}_{\mathcal{V}_b}(V, V') = 1$ whenever $V > V'$ and
 - (b) composition of nonzero maps $V \rightarrow V' \rightarrow V''$ is always nonzero.
- (4) *Any $W \in \mathcal{W}_b$ has a unique filtration $0 = W_0 \subset W_1 \subset \cdots \subset W_m$ so that each W_k/W_{k-1} lies in $\text{add } \mathcal{V}_k$ where $V_1 < V_2 < \cdots < V_m$. Evaluating at vertex b we get a filtration $W_1(b) \subset W_2(b) \subset \cdots \subset W_m(b) = W(b)$ which we call the filtration of $W(b)$ induced by the filtration of W .*

- (5) For any $W \in \mathcal{W}_b$, any direct sum decomposition $W(b) = \bigoplus X_l$ of $W(b)$ into one-dimensional subspaces which respects the filtration of $W(b)$ induced from the filtration of W extends to a direct sum decomposition of W , i.e. $W = \bigoplus Y_l$ so that $Y_l(b) = X_l$ for all l .

Proof. (1) Given V, V' in \mathcal{V}_b , the support of one of them contains the support of the other. Let $J = \text{supp } V \cap \text{supp } V' \subset [b, \infty)$. Then either $J = \text{supp } V$ or $J = \text{supp } V'$. Suppose $J = \text{supp } V$. Since V is indecomposable J is connected and $J \subseteq \text{supp } V'$. Any morphism $f : V \rightarrow V'$ induces a morphism $f_J : V_J \rightarrow V'_J$ by restricting to J . By Theorem 2.3.2, $V_J \cong V'_J$ is either V or V' . Then we have, by Lemma 2.3.1, that f_J is a scalar times a fixed isomorphism $V_J \cong V'_J$. In particular $f = 0$ if and only if f is zero at b . So, evaluation at b is faithful. (2) follows immediately from (1).

(3) Given V, V' in \mathcal{V}_b , suppose by symmetry that the support of V is properly contained in the support of V' . Then, there is some $m > b$ so that the support of V is contained in $[b, m]$. There are only finitely many elements of S inside this compact set. Without loss of generality we may assume that $b \in S$. Let l be maximal so that s_l is in the support of V . If s_l is a sink, then V is a sub-representation of V' . If s_l is a source, then V is a quotient representation of V' . In the first case, $\text{Hom}(V, V') = k$ and $\text{Hom}(V', V) = 0$. In the second case, $\text{Hom}(V, V') = 0$ and $\text{Hom}(V', V) = k$.

If there are nonzero morphisms $V \rightarrow V' \rightarrow V''$ then, by (1), evaluation at b gives isomorphisms $V(b) \cong V'(b) \cong V''(b)$. So, the relation of having a nonzero morphism $V \rightarrow V'$ is transitive, reflexive, antisymmetric and any two elements are related. So, this is a total ordering.

(4) Given $W \in \mathcal{W}_b$ we have by definition a direct sum decomposition $W = \bigoplus_{1 \leq i \leq m} (V_i)^{n_i}$ where we order the summands according to the total order given in (3). So, there exists a filtration $0 = W_0 \subset W_1 \subset W_2 \subset \cdots \subset W_m = W$ so that $W_i/W_{i-1} = n_i V_i$. Since $\text{Hom}(V_i, V_j) = 0$ for $i < j$, the sub-representation W_i is uniquely characterized as the trace of $V_1 \oplus V_2 \oplus \cdots \oplus V_i$ in W . So, the HN-filtration is unique.

(5) Let $n = \dim W(b)$ and let G be the subgroup of $GL(n, k)$ which preserves the filtration $W_1(b) \subset W_2(b) \subset \cdots \subset W_m(b)$. This is a block upper triangular matrix group which acts transitively on the set of all bases which respect this filtration of $W(b)$. Since $\text{Hom}(V_i, V_j) = K$ for $i \leq j$ and $\text{Hom}(V_i, V_j) = 0$ for $i > j$, we have by (3) that the restriction map $\text{Aut}(W) \rightarrow \text{Aut}(W(b)) = G$ is an isomorphism. Therefore, $\text{Aut}(W)$ acts transitively on the set of all bases for the vector space $W(b)$ which respect the given filtration of $W(b)$.

Recall we are given a direct sum decomposition $W(b) = \bigoplus X_l$ into one-dimensional subspaces that respects the induced filtration and $W = \bigoplus Y_l$ a direct sum decomposition of W into pointwise one-dimensional indecomposable representations. One such basis is given by choosing a generator $x_l \in X_l$ for each summand X_l of $W(b) = \bigoplus X_l$. A second basis is given by choosing a generator $y_l \in Y_l(b)$ where $W = \bigoplus (V_i)^{n_i} = \bigoplus Y_l$, where each Y_l is equal to some V_i , is the given decomposition of W into indecomposable representations which are one-dimensional at b . Take $\varphi \in G$ which takes (y_l) to (x_l) . Then $W = \bigoplus \varphi(Y_l)$ is the required decomposition of W extending the chosen decomposition of $W(b)$. \square

Lemma 2.3.5. *Given any two finite filtrations of a finite-dimensional vector space X , there exists a direct sum decomposition of X into one-dimensional subspaces which respects both filtrations.*

Proof. Given any two filtrations $V_1 \subset V_2 \subset \cdots \subset V_n = X$ and $W_1 \subset W_2 \subset \cdots \subset W_m = X$ of X we have the following representation of a quiver of type A_{n+m-1} :

$$M : \quad V_1 \xrightarrow{\subset} V_2 \xrightarrow{\subset} \cdots \xrightarrow{\subset} V_{n-1} \xrightarrow{\subset} X \xleftarrow{\supset} W_{m-1} \xleftarrow{\supset} \cdots \xleftarrow{\supset} W_2 \xleftarrow{\supset} W_1.$$

We have a direct sum decomposition $M = \bigoplus M_i$ where each M_i is one-dimensional at the middle vertex. This gives a direct sum decomposition of X into one-dimensional subspaces. Then it suffices to prove the following.

Claim: This decomposition $X = M(n) = \bigoplus M_i(n)$ respects both filtrations.

Proof: Since the maps in the representation M are all monomorphisms, the same holds for each indecomposable component M_i . So, each component is nonzero at vertex n (where $M(n) = X$).

For any $1 \leq j < n$, consider the set I_j of all indices i so that $M_i(j) \neq 0$. Then the sum of all $M_i(n)$ for all $i \in I_j$ is equal to V_j . Thus $\bigoplus M_i(n)$ respects the first filtration V_i of X . Similarly, $\bigoplus M_i(n)$ respects the second filtration W_j . So, it respects both filtrations. This proves the lemma. \square

2.4. Necessary Conditions and Decomposition Theorem. In this section we prove that the sufficient conditions in Proposition 2.2.1 are also necessary conditions. We then work up to Lemmas 2.4.9 and 2.4.10. Lemma 2.4.9 shows that the decomposition of certain subrepresentations may be extended to infinity. Lemma 2.4.10 then states that these subrepresentations are indeed summands. Because we may have infinitely many sinks and sources in our continuous quiver, these lemmas are an essential component of the proof of Theorem 2.4.15, the decomposition theorem. We save our discussion relating our proof of Theorem 2.4.15 to the decomposition theorems in [4, 5, 12] for Section 2.6.

Definition 2.4.1. Choose $A_{\mathbb{R}}$, a continuous quiver of type A . The opposite quiver of $A_{\mathbb{R}}$, denoted $A_{\mathbb{R}}^{\text{op}}$, is the continuous quiver of type A where $x \preceq y$ in $A_{\mathbb{R}}^{\text{op}}$ if and only if $y \preceq x$ in $A_{\mathbb{R}}$.

Let V be a pointwise finite representation of $A_{\mathbb{R}}$. The dual representation of V , denoted DV , is the pointwise finite representation of $A_{\mathbb{R}}^{\text{op}}$ given by

$$DV(x) := D(V(x)) \quad DV(y, x) := D(V(x, y))$$

Remark 2.4.2. In Definition 2.4.1, note that since V is pointwise finite we have $DDV \cong V$.

Lemma 2.4.3. *Let V be any object of $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. Then the restriction V_J of V to any closed interval $J = [a, b]$ where $s_n \leq a < b \leq s_{n+1}$ for some $n \in \mathbb{Z}$, decomposes as $V_J = A \oplus B$ where A has support in the open interval (a, b) and B is a finite direct sum of indecomposable one-dimensional representations which are nonzero at either a , b or both.*

Proof. Without loss of generality we may assume that s_n is a sink and s_{n+1} is a source. Let K be the subrepresentation of V_J given by $K(x) = \ker(V(x, n) : V(x) \rightarrow V(s_n))$ for all $x \in J = [s_n, s_{n+1}]$.

Then $P = V_J/K$ is a projective representation of J since all morphisms $P(x) \rightarrow P(s_n)$ are monomorphisms by construction of K . So $V_J \cong K \oplus P$. It is straightforward to decompose P as a direct sum of finitely many pointwise one-dimensional representations, each nonzero at n .

It remains to show that K is a direct sum of a representation with support on the open interval (s_n, s_{n+1}) and a finite number of pointwise one-dimensional representations all nonzero at s_{n+1} . This is accomplished using the dual representation DK . Since DK is a representation of the opposite quiver, the interval J with n as source and s_{n+1} as sink, using exactly the same argument as above we see that $DK = A \oplus B$ where $A_{n+1} = 0$ and B is a projective representation of J^{op} . Thus $K \cong DA \oplus DB$ where DA has support in the open interval (s_n, s_{n+1}) and DB is a finite direct sum of one-dimensional representations which are all nonzero at s_{n+1} . \square

Lemma 2.4.4. *If V is an indecomposable object of $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ with support in an interval $[s_n, s_{n+1}]$ for some $n \in \mathbb{Z}$, then V is pointwise one-dimensional.*

Proof. The support of V must be an interval $J \subseteq [s_n, s_{n+1}]$. If J contains either of its endpoints then the previous lemma applies. It remains to consider the case when $J = (a, b)$ is open. Let $c \in (a, b)$. Then applying the previous lemma to the intervals $[a, c]$ and $[c, b]$ we decompose $V_{[a, c]}$ and $V_{[c, b]}$ into a direct sum of finitely many pointwise one-dimensional representations each of which is nonzero at c . The other components of $V_{[a, c]}$ and $V_{[c, b]}$ given by the lemma must be zero since they would be components of V . This is equivalent to a representation of a finite quiver of type A_m with straight orientation. So, we can choose the decompositions of $V_{[a, c]}$ and $V_{[c, b]}$ so that they give the same decomposition of V_c . This decomposes $V = V_{[a, b]}$ into a direct sum of pointwise one-dimensional representations. Since V is indecomposable there is only one component. \square

Lemma 2.4.5. *Let V be an indecomposable object of $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. For any two integers $n < m$, the restriction of V to the closed interval $[s_n, s_m]$ is a direct sum of finitely many indecomposable pointwise one-dimensional representations.*

Proof. The proof is by induction on $m - n$. Suppose first that $m - n = 1$ and let $J = [s_n, s_m] = [s_n, s_{n+1}]$. If $\text{supp} V \subseteq J$ then V is pointwise one-dimensional by Lemma 2.4.4. If $\text{supp} V \not\subseteq J$ then, by Lemma 2.4.3, the restriction of V to J is a direct sum of pointwise one-dimensional objects plus a summand A with support on (s_n, s_{n+1}) . But such a summand would also be a summand of V by Lemma 2.4.6. Therefore, $A = 0$ and the Lemma holds for $m = n + 1$.

Now suppose $m \geq n + 2$ and take any integer k so that $n < l < m$. By induction on $m - n$, $V_{[s_n, s_l]}$ and $V_{[s_l, s_m]}$ decompose into pointwise one-dimensional components. By Lemma 2.3.4, this gives two filtrations of V_k . By Lemma 2.3.5, there is a direct sum decomposition of V_n compatible with both filtrations. This extends to compatible direct sum decompositions of $V_{[s_n, s_l]}$ and $V_{[s_l, s_m]}$ which paste together to give a decomposition of $V_{[s_n, s_m]}$ into one-dimensional representations. \square


Lemma 2.4.6. *Let V be a representation in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ and let $V_{(-\infty, b]}$ be the restriction of V to the interval $(-\infty, b]$. Then any summand W of $V_{(-\infty, b]}$ which is zero at b is a summand of V .*

Proof. Let $\pi : V_{(-\infty, b]} \rightarrow V_{(-\infty, b]}$ be the projection to W . Then $\pi_b : V(b) \rightarrow V(b)$ is zero. So, π and the zero morphism on $V_{[b, \infty)}$ agree on the overlap of their domains. So, their union is an endomorphism of V . This endomorphism is evidently the projection to W showing that W is a summand of V . \square

Construction 2.4.7. Let $A_{\mathbb{R}}$ be a continuous quiver of type A whose sinks and sources are unbounded above. I.e., for each sink or source s_n there is an s_{n+1} . Let V be a pointwise finite-dimensional representation of $A_{\mathbb{R}}$ such that, for all $n \in \mathbb{Z}$, the restriction $V_{[s_n, s_{n+1}]}$ contains no direct summands whose support is contained entirely in (s_n, s_{n+1}) (i.e., $A = 0$ in the $A \oplus B$ decomposition in Lemma 2.4.3).

Consider the restriction $V_{[s_{l-1}, s_l]}$. By assumption $V_{[s_{l-1}, s_l]}$ is a finite direct sum of indecomposables, all of whose support includes s_l or s_{l-1} . Let $V_{0,0}^l$ be the direct sum of all those summands that include *only* s_l , not s_{l-1} . Now consider $V_{[s_{l-1}, s_{l+1}]}$. By assumption $V_{[s_{l-1}, s_{l+1}]}$ is a finite direct sum of indecomposables, each of whose support contains s_{l-1} , s_l or s_{l+1} . Let $V_{0,1}^l$ be the direct sum of all such indecomposables whose support contains *both* s_l and s_{l+1} , but *not* s_{l-1} . Let $V_{1,1}^l$ be the direct sum of such indecomposables whose support contains s_l but *not* s_{l+1} or s_{l-1} . We ignore those indecomposables whose support does not contain s_l .

We can continue this process for all $n \geq 0$. For each $n \geq 0$ and $0 \leq i \leq n$ we define $V_{i,n}^l$ in the following way. It is the direct sum of those summands of $V_{[s_{l-1}, s_{l+n}]}$ whose support contains exactly sinks and sources s_{l+j} for $0 \leq j \leq n - i$. Note that this never includes s_{l-1} . In particular, $V_{0,n}^l$ is the direct sum of those interval indecomposable summands of $V_{[s_{l-1}, s_{l+n}]}$ whose support contains s_l and s_{l+n} . We have three examples below, two from the previous paragraph and also the summands we consider from $V_{[s_{l-1}, s_{l+2}]}$.



We note that if $1 \leq i \leq n$ then $V_{i,n}^l = V_{i+1,n+1}^l$. Note also that the constructions can be made on $(-\infty, s_{l+1}]$ instead and those representations are denoted $V_l^{i,n}$. \diamond

Proposition 2.4.8. *Let $A_{\mathbb{R}}$ and V be as in Construction 2.4.7 for some s_l . Then, for all $n \geq 1$ and $1 \leq i \leq n$, $V_{i,n}^l$ and $V_l^{i,n}$ are split subrepresentations of V .*

Proof. We know the representation $V_{1,n}$ is a split subrepresentation of $V_{[s_{l-1}, s_{l+n+1}]}$ as a consequence of Lemma 2.4.5. We know that $V_{1,n}(s_{l-1}) = 0$ and $V_{1,n}(s_{l+n+1}) = 0$. By two uses of Lemma 2.4.6 we see that $V_{1,n}$ is a split subrepresentation of V . Finally, recall that $V_{i,n}^l = V_{i+1,n+1}^l$ when $i \geq 1$. By a similar argument $V_l^{i,n}$ is a split subrepresentation of V . \square

Lemma 2.4.9. *Let $l \in \mathbb{Z}$ and V be a representation with support contained in $[s_{l-1}, +\infty)$. Assume that for all $n \geq 0$, any indecomposable summand of $V_{[s_{l-1}, s_{l+n}]}$ has support at s_{l+n} . Then a decomposition of $V_{[s_{l-1}, s_{l+n}]}$ into interval indecomposables extends to a decomposition of $V_{[s_{l-1}, s_{l+n+1}]}$.*

Proof. Suppose $V_{[s_{l-1}, s_{l+n}]} \cong \bigoplus_i M_{I_i}$ is a decomposition. Then each I_i includes s_{l+n} .

If s_{l+n} is a sink then $V(s_{l+n}, s_{l+n+1})$ is a monomorphism. Any interval indecomposable summands of $V_{[s_{l-1}, s_{l+n+1}]}$ that do not have support at s_{l+n} are projective. In particular, they are split subrepresentations of the same restriction (combine Lemmas 2.4.3 and 2.4.6). Let U_{n+1} be the quotient of $V_{[s_{l-1}, s_{l+n+1}]}$ by these projective interval indecomposables. Since $(U_{n+1})_{[s_{l-1}, s_{l+n}]} = V_{[s_{l-1}, s_{l+n}]}$ and $U_{n+1}(s_{l+n}, s_{l+n+1})$ is an isomorphism we can extend the decomposition to U_{n+1} . Since U_{n+1} is a decomposable summand of $V_{[s_{l-1}, s_{l+n+1}]}$ and the other summand is decomposable by Theorem 2.1.16 we have extended our decomposition.

If s_{l+n} is a source then $V(s_{l+n}, s_{l+n+1})$ is an epimorphism. Any interval indecomposable summands of $V_{[s_{l-1}, s_{l+n+1}]}$ that do not have support at s_{l+n} are injective. They are split subrepresentations as before. We can now apply the same argument in the previous paragraph and extend the decomposition. \square

The assumptions in the following lemma are justified by Proposition 2.4.8.

Lemma 2.4.10. *Let $A_{\mathbb{R}}$ and V be as in Construction 2.4.7. For all $l \in \mathbb{Z}$, $n \geq 1$, and $1 \leq i \leq n$, assume $V_{i,n}^l = 0 = V_l^{i,n}$. Then V contains a summand as in Lemma 2.4.9 but whose support does not contain s_{l-1} .*

Proof. For each $n \geq 1$ and a decomposition of $V_{[s_{l-1}, s_{l+n}]}$ let K_n be the sum of interval summands whose support is nonzero at s_{l-1} . By assumption, if $s_{l-1} \leq x \leq y \leq s_{l+n}$ then $\dim K_n(x) \geq \dim K_n(y)$. Note that $\dim K_n(x) = \dim K_{n+1}(x)$ on $[s_{l-1}, s_{l+n}]$, though the decomposition of K_n is not assumed to extend exactly.

Therefore we have a function $[s_{l-1}, +\infty) \rightarrow \mathbb{N}$ that is weakly decreasing and whose initial value is finite. Therefore, the function must stabilize to some particular value. Let m be sufficiently large that $\dim K_n(s_{l+n}) = \dim K_{n+1}(s_{l+n+1})$ for all $n \geq m$. Then, by assumption, every map $V(s_{l+n}, s_{l+n+1})$ for $n \geq m$ is mono or epi. So we can use the same technique in Lemma 2.4.9 to extend a decomposition of $V_{[s_{l-1}, s_{l+m+1}]}$ to all of $V_{[s_{l-1}, +\infty)}$.

Then any summands of $V_{[s_{l-1}, +\infty)}$ with bounded support that is nonzero at s_{l-1} are split subrepresentations of $V_{[s_{l-1}, +\infty)}$ (Lemma 2.4.6). Denote those summands by U and the rest by W . Then W satisfies the hypothesis of Lemma 2.4.9 and we have a decomposition of W already. In particular, we can write $W \cong W_1 \oplus W_2$ where the summands of W_1 are nonzero at s_{l-1} and the summands at W_2 are 0 at s_{l-1} . Then by a further use of Lemma 2.4.6 we see W_2 is actually the summand of V that we desired. \square

Notation 2.4.11. Let $A_{\mathbb{R}}$ and V be as in Construction 2.4.7. For some l , let W_2 be as in the end of the proof of Lemma 2.4.10. As seen in the proof, W_2 is a direct sum of interval indecomposables. Let $V_{0,\infty}^l$ be the direct sum of those summands of W_2 who have support at s_l .

Remark 2.4.12. Construction 2.4.7, Proposition 2.4.8, Lemma 2.4.9, Lemma 2.4.10, and Notation 2.4.11 can all be performed on $(-\infty, s_{l+1}]$ instead of $[s_{l-1}, +\infty)$. These representations will be denoted $V_l^{i,n}$ and $V_l^{0,\infty}$.

Lemma 2.4.13. *Let V be a puf representation with support on an open interval which has no sinks or sources. Let c be any point in the open interval. Then $V = V_0 \oplus V_1 \oplus V_2$ where V_0 is a direct sum of finitely many interval representations having c in its support and V_1, V_2 are representations having support strictly below c , strictly above c , respectively.*

Proof. This follows from a combination of Lemmas 2.3.5 and 2.4.3. \square

In order to prove Theorem 2.4.15 we prove the following lemma, which recovers the specific case of the real line in Crawley-Boevey's theorem in [12].

Lemma 2.4.14. *Let V be as in Lemma 2.4.13. Then V is a direct sum of interval representations.*

Proof. Let (a, b) be an open subinterval of \mathbb{R} containing the support of V where a and b are allowed to be $-\infty$ and $+\infty$, respectively. Let C be the set of all points in (a, b) of the form $c(k, n) := \tan(\tan^{-1} a + k(\tan^{-1} b - \tan^{-1} a)/2^n)$ for positive integers k, n where k is odd, which is a dense subset of (a, b) .

By induction on n using Lemma 2.4.13, we get a decomposition of V in the form $V = V_\infty \oplus \bigoplus V_{k,n}$ where each $V_{k,n}$ is a direct sum of finitely many interval indecomposables having $c(k, n)$ in its support and no point in the form $c(j, m)$ in its support where $m < n$ and V_∞ has no elements of the form $c(k, n)$ in its support. In that case, V_∞ is a direct sum of simple representations since for any $a \leq c < d \leq b$, there is a number of the form $c(k, n)$ in the open interval (c, d) . So, the morphism $V_\infty(c, d) : V_\infty(c) \rightarrow V_\infty(d)$ must be zero since it factors through $V_\infty(c(k, n)) = 0$. \square

Theorem 2.4.15. *Let $A_{\mathbb{R}}$ be a continuous quiver of type A and V be a representation in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. Then V is the direct sum of interval indecomposables (Definition 2.2.2).*

Proof. Outline: We complete this proof in four parts. In Part 1, we consider the indecomposable summands whose support is contained entirely between a sink and source. In Part 2, we consider the indecomposable summands whose support contains at least one but only finitely many sinks and sources. In Part 3, we consider the indecomposable summands whose support may contain infinitely many sinks and sources, but is bounded on exactly one side. Finally, in Part 4, we concern ourselves with indecomposable summands whose support is \mathbb{R} . Since the case where $A_{\mathbb{R}}$ has no sinks or sources in \mathbb{R} has been covered by Lemma 2.4.14, we assume that $A_{\mathbb{R}}$ has at least one sink or source in \mathbb{R} .

Part 1: Let s_n and s_{n+1} be an adjacent pair of sink, source or $\pm\infty$; however, only one may be $\pm\infty$ by assumption. We use the notation $[s_n, s_{n+1}]$ even if one of the endpoints is actually $\pm\infty$. By Lemma 2.4.3, $V_{[s_n, s_{n+1}]}$ decomposes to $A_n \oplus B_n$ where the support of A_n is contained in (s_n, s_{n+1}) . By Lemma 2.4.6, A_n is a direct summand of V .

Thus, for all n where s_n or s_{n+1} is in \mathbb{R} , we have such an A_n . So we have that $V \cong (\bigoplus A_n) \oplus U$. By Lemma 2.4.14 each A_n decomposes into a direct sum of indecomposable representations.

Part 2: We now assume $V \cong U$ as in the end of Part 1. If $A_{\mathbb{R}}$ has finitely many sinks and sources then, by the proof of Lemma 2.4.5, V is a finite direct sum of indecomposable representations. So we shall now assume $A_{\mathbb{R}}$ has infinitely many sinks and sources. Choose a sink or source s_l in \mathbb{R} . By Proposition 2.4.8 we know $V_{i,n}^l$ is a split subrepresentation with bounded support for all $1 \leq i \leq n$. Thus, for all l such that $s_l \in \mathbb{R}$ we obtain such direct summands, none of which are counted twice. Thus we have $V \cong (\bigoplus_{l,n} V_{1,n}^l) \oplus U$.

Part 3: Now we assume $V \cong U$ as in the end of Part 2. Then for each $l \in \mathbb{Z}$ we apply Lemma 2.4.10 and obtain $V_{0,\infty}^l$ as in Notation 2.4.11. By Remark 2.4.12 we also obtain $V_l^{0,\infty}$ for each l . Each $V_{0,\infty}^l$ and $V_l^{0,\infty}$ decompose into interval indecomposables and so we have $V \cong (\bigoplus (V_{0,\infty}^l \oplus V_l^{0,\infty})) \oplus U$.

Part 4: We assume $V \cong U$ as in the end of Part 3. For any $s_l \in \mathbb{R}$, we know $V_{0,\infty}^l = 0$ and $V_l^{0,\infty} = 0$. Choose some sink or source s_l in \mathbb{R} and let $X = V_{[s_l, +\infty)}$ and $Y = V_{(-\infty, s_l]}$. We can then construct $X_{0,\infty}^l$ and $Y_l^{0,\infty}$. Since $V_{0,\infty}^l = 0$ and $V_l^{0,\infty} = 0$, we see $\dim X_{0,\infty}^l(s_l) = \dim Y_l^{0,\infty}(s_l)$. In particular, they are both finite.

Furthermore, $V(x, y)$ is an isomorphism for all $y \preceq x$ in \mathbb{R} . Choose a decomposition of $V_{[s_{l-1}, s_{l+1}]}$ and use the technique in Lemma 2.4.9 to extend this decomposition to all of X and all of Y . But together this yields a decomposition of V .

This will give us a bijection $V \cong \bigoplus_{\dim V(s_l)} M_{(-\infty, +\infty)}$. Thus, V is a direct sum of indecomposable representations.

Conclusion: In Parts 1–3 we decomposed V into $Z \oplus U$ and in Parts 2–4 we decomposed the previous Part's U . In Parts 1–3 we showed that the Z summand was a direct sum of indecomposables and in Part 4 we showed the final U is a direct sum of indecomposables. Therefore, given any pointwise finite-dimensional representation V of $A_{\mathbb{R}}$, it is the direct sum of indecomposable representations. If V itself is indecomposable it appears as one of described indecomposable summands, depending on its support. \square

Remark 2.4.16. The theorem above, with the aid of Theorem 2.1.16, completely classifies indecomposable projective objects in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ and $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$. They come in three forms, up to isomorphism.

(1) P_a as in Definition 2.1.4:

$$P_a(x) = \begin{cases} k & x \preceq a \\ 0 & \text{otherwise} \end{cases} \quad P_a(x, y) = \begin{cases} 1_k & y \preceq x \preceq a \\ 0 & \text{otherwise.} \end{cases}$$

(2) P_a given by

$$P_a(x) = \begin{cases} k & x \preceq a, x < a \\ 0 & \text{otherwise} \end{cases} \quad P_a(x, y) = \begin{cases} 1_k & y \preceq x \preceq a, y \leq x < a \\ 0 & \text{otherwise.} \end{cases}$$

(3) P_a given by

$$P_a(x) = \begin{cases} k & x \preceq a, a < x \\ 0 & \text{otherwise} \end{cases} \quad P_a(x, y) = \begin{cases} 1_k & y \preceq x \preceq a, a < x \leq y \\ 0 & \text{otherwise.} \end{cases}$$

Note that unless a is a source at least one of (2) or (3) will define the 0 representation. If a is a sink then both (2) and (3) will be the 0 representation.

Additionally, it is worth noting that if V is a subrepresentation of any sum of projective indecomposables then V is also projective. This follows from Theorem 2.1.16 (1). Therefore, $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ is hereditary.

Example 2.4.17. Let the set of sinks and sources $S = \{0, 1\}$, where $s_0 = 0$ is a sink and $s_1 = 1$ is a source. We provide a complete list of indecomposable projectives in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ with this orientation. The values $a, b, c \in \mathbb{R}$ below are such that $a < 0 < b < 1 < c$.

$$\begin{array}{cccccccccccc} \{ \{0\}, & (-\infty, 0], & (a, 0], & [a, 0], & [0, b), & [0, b], & [0, 1), & [0, +\infty), & (1, +\infty), & (c, +\infty), & [c, +\infty) \} \\ P_0 & P_{-\infty} & P_{(a)} & P_{[a]} & P_{(b)} & P_{[b]} & P_{(1)} & P_{[1]} & P_{(1)} & P_{(c)} & P_{[c]} \end{array}$$

Remark 2.4.18. We also have the indecomposable injective objects in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$.

(1) I_a given by:

$$I_a(x) = \begin{cases} k & a \preceq x \\ 0 & \text{otherwise} \end{cases} \quad I_a(x, y) = \begin{cases} 1_k & a \preceq y \preceq x \\ 0 & \text{otherwise} \end{cases}$$

(2) I_a given by

$$I_a = \begin{cases} k & a \preceq x, x < a \\ 0 & \text{otherwise} \end{cases} \quad I_a(x, y) = \begin{cases} 1_k & a \preceq y \preceq x, x \leq y < a \\ 0 & \text{otherwise} \end{cases}$$

(3) I_a given by

$$I_a = \begin{cases} k & a \preceq x, a < x \\ 0 & \text{otherwise} \end{cases} \quad I_a(x, y) = \begin{cases} 1_k & a \preceq y \preceq x, a < y \leq x \\ 0 & \text{otherwise} \end{cases}$$

2.5. More on P_a , P_a , and the Pointwise Finite Requirement. As mentioned in Section 2.1, the indecomposable projectives P_a and P_a , whichever are nonzero, are not projective in $\text{Rep}_k(A_{\mathbb{R}})$. They are only projective in the smaller subcategory $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. We will prove this using a specific representation, denoted \mathfrak{P} , that exists only in $\text{Rep}_k(A_{\mathbb{R}})$. I.e., it is not pointwise finite-dimensional. We will use that same representation to show why Theorem 2.4.15 can fail without the pointwise finite assumption.

Construction 2.5.1. We will denote the problematic representation by \mathfrak{P} . First, let $a \in \mathbb{R}$ such that a is not a sink. Let $p \in \mathbb{R}$ such that $p \preceq a$ and $p \neq a$. By symmetry, suppose $p < a$. Let $\{x_i\}_{i=0}^{\infty}$ be a strictly increasing sequence converging to a such that $x_0 > p$. Let $M = \bigoplus_{\{x_i\}} M_{[p, x_i]}$. Then the support of M is $[p, a)$. Let $\pi : M(p) \rightarrow k$ be a surjection given by sending each 1 in $M_{[p, x_i]}(p) = k$ to $1 \in k$.

Let \mathfrak{P} be given by

$$\mathfrak{P}(x) = \begin{cases} k & x = p \\ M(x) & x \neq p \end{cases} \quad \mathfrak{P}(x, y) = \begin{cases} 1_k & x = y = p \\ \pi \circ M(x, y) & x \neq y = p \\ M(x, y) & \text{otherwise.} \end{cases}$$

We see that \mathfrak{P} also has support $[p, a)$. ◇

Proposition 2.5.2. *Let $A_{\mathbb{R}}$ be a continuous quiver of type A. Let $p, a \in \mathbb{R}$ such that a is not a sink, $p \preceq a$, and $p < a$. Then there is no nontrivial morphism $P_a \rightarrow \mathfrak{P}$, where \mathfrak{P} is from Construction 2.5.1.*

Proof. Choose x_m in the sequence from Construction 2.5.1. Let $f(x_m) : P_a \rightarrow \mathfrak{P}(x)$ be a linear map. Since $P_a = k$, $f(x_m)$ is determined by $f(x_m)(1)$. Since $\mathfrak{P}(x) = M(x)$ for $x \neq p$, we see

$$f(1) = (\underbrace{0, \dots, 0}_{m-1}, r_m, r_{m+1}, \dots, r_n, 0, 0, \dots)$$

Then for any linear map $f(x_{n+1}) : P_a(x_{n+1}) \rightarrow \mathfrak{P}(x_{n+1})$ we know that

$$f(x_m) \circ P_a(x_{n+1}, x_m) \neq \mathfrak{P}(x_{n+1}, x_m) \circ f(x_{n+1})$$

Therefore, there is no morphism of representations $P_a \rightarrow \mathfrak{P}$. □

Proposition 2.5.3. *Let $A_{\mathbb{R}}$ be a continuous quiver of type A and $a \in \mathbb{R}$ such that a is not a sink. Then each nonzero P_a and P_a is not projective in $\text{Rep}_k(A_{\mathbb{R}})$.*

Proof. Let $p \in \mathbb{R}$ such that $p < a$ and $p \preceq a$. The other case, where $p > a$ and $p \preceq a$, is similar. Then, there is a nontrivial morphism of indecomposable representations $f : P_a \rightarrow M_{[p, a)}$. Let \mathfrak{P} be as in Construction 2.5.1. For each $x \in [p, a)$, let $f(x) : \mathfrak{P}(x) \rightarrow M_{[p, a)}(x)$ be given by $\mathfrak{P}(x, p)$. Since $\mathfrak{P}(p) = M_{[p, a)}(x)$ for all $x \in [p, a)$, this is a well-defined morphism of representations. In particular, it is an epimorphism.

So now we have an epimorphism $P_a \twoheadrightarrow M_{[p, a)}$ and an epimorphism $\mathfrak{P} \twoheadrightarrow M_{[p, a)}$. However, there is no nontrivial morphism $P_a \rightarrow \mathfrak{P}$ in $\text{Rep}_k(A_{\mathbb{R}})$, by Proposition 2.5.2. Therefore, P_a is not projective. □

Proposition 2.5.4. *Let $A_{\mathbb{R}}$ be a continuous quiver of type A and \mathfrak{P} as in Construction 2.5.1. Then \mathfrak{P} is not the direct sum of pointwise one-dimensional indecomposables.*

Proof. We saw in the proof of Proposition 2.5.3 that there is an epimorphism $\mathfrak{P} \rightarrow M_{[p,a]}$. However, just as in the proof of Proposition 2.5.2 there are no nontrivial morphisms $M_{[p,a]} \rightarrow \mathfrak{P}$. Thus, $M_{[p,a]}$ is not a direct summand of \mathfrak{P} . But if \mathfrak{P} had a direct sum decomposition, one of the components must have support $[p, a]$. But that would mean the indecomposable is $M_{[p,a]}$. Therefore, \mathfrak{P} does not decompose into a direct sum of one-dimensional indecomposables. \square

2.6. Relation to Decomposition Theorems in Persistent Homology. Theorem 2.4.15 is, in some sense, a combination of the Crawley-Boevey’s BarCode theorem from [12] and Botnan’s decomposition theorem in [4]. Part of our argument actually follows the latter paper. The BarCode theorem handles representations on the continuum but only a straight orientation. By contrast, Botnan’s decomposition handles the infinite zigzag orientation but only in the discrete setting.

One might think to use Botnan’s paper explicitly with Crawley-Boevey’s result. However, this cannot be done directly. In order to make use of the combination of theorems, several technical lemmas would still be required. In particular, one would have to argue which pwf representations can be “lifted” to a discrete quiver and then prove that the decomposition can be “pushed back down.” While intuitive, the technical details in such an argument (see [14] for a similar argument) are still involved. We avoided such a proof in order to provide a self-contained foundation of continuous type A quivers as well as an algorithmic proof of Theorem 2.4.15.

While Theorem 2.4.15 recovers a result by Botnan and Crawley-Boevey in [5], the method of proof is different. One might consider the proof presented in Section 2.4 as a “direct” proof while the proof in [5] uses representations of products of posets.

Both of [12, 4] worked with pointwise finite-dimensional representations and each displayed a non-example for a representation that is not pointwise finite-dimensional. Theorem 2.4.15 adheres to exactly the same restrictions and a relevant non-example appears in Section 2.5 as Construction 2.5.1 and Proposition 2.5.4.

3. FINITELY GENERATED REPRESENTATIONS: $\text{rep}_k(A_{\mathbb{R}})$

In this section we will prove results about the category of finitely generated representations, denoted $\text{rep}_k(A_{\mathbb{R}})$. Many of the properties one could reasonably expect to hold in a continuous version of $\text{rep}_k(A_n)$ do, in fact, hold for $\text{rep}_k(A_{\mathbb{R}})$. The properties that change due to the nature of the continuum are Auslander–Reiten sequences and descending chains of subrepresentations. We provide an incomplete list of the properties that hold or do not hold in the form of a theorem and dedicate the rest of this section to proving each of the items in the theorem.

Theorem 3.0.1. *Let $A_{\mathbb{R}}$ be a continuous quiver of type A and denote by $\text{rep}_k(A_{\mathbb{R}})$ the category of finitely generated representations (Definition 3.1.3). Then the following hold.*

- (1) *For indecomposable representations M_I and M_J in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$, or $\text{rep}_k(A_{\mathbb{R}})$, we have $\text{Hom}(M_I, M_J) \cong k$ or $\text{Hom}(M_I, M_J) = 0$ (Proposition 3.1.2).*
- (2) *Every morphism $f : V \rightarrow W$ in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$, or $\text{rep}_k(A_{\mathbb{R}})$ has a kernel, a cokernel, and coinciding image and coimage in that category. (Lemma 3.1.4)*
- (3) *The category $\text{rep}_k(A_{\mathbb{R}})$ Krull-Schmidt, but not artinian (Lemma 3.1.5, Proposition 3.1.7).*
- (4) *The global dimension of $\text{rep}_k(A_{\mathbb{R}})$ is 1 (Proposition 3.2.5).*
- (5) *The Ext space of two indecomposables M_I and M_J in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$, or $\text{rep}_k(A_{\mathbb{R}})$ is either isomorphic to k or is 0 (Proposition 3.2.6).*
- (6) *While some Auslander–Reiten sequences exist (Proposition 3.3.2), some indecomposables have neither a left nor a right Auslander–Reiten sequence (Proposition 3.3.3).*

3.1. Requisites and Definition. In this subsection we define the category of finitely generated representations of a continuous type A and prove Theorem 3.0.1 (1) – (3).

Notation 3.1.1. We may use $|$ instead of $(,)$, $[,]$ or $] , [$ to write an interval. When this happens, we mean that the endpoint may or may not be included; either we are making no assumptions about endpoints or it is clear what choice is possible from context. I.e., for all $a, b \in \mathbb{R}$, $|a, b|$ can be one of four possibilities. However, when we write our intervals, we allow $a = -\infty$ and $b = +\infty$ so long as we obtain a subset of \mathbb{R} . So, the notation $|a, b|$ will never mean $[-\infty, b|$, $|a, +\infty]$, or $[-\infty, +\infty]$.

Proposition 3.1.2. *Let V and W be indecomposable representations in $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$. Then either $\text{Hom}(V, W) \cong k$ or $\text{Hom}(V, W) = 0$.*

Proof. Suppose $\text{Hom}(V, W) \neq 0$ and choose a nontrivial $f : V \rightarrow W$. Then there is $x \in \mathbb{R}$ such that $f(x) : V(x) \rightarrow W(x)$ is not 0. Since $V(x) \cong k \cong W(x)$ we see $f(x)$ is an isomorphism. For all $y \preceq x$, $W(x, y) \circ f(x) = f(y) \circ V(x, y)$. If $V(y) \neq 0$ and $W(y) \neq 0$ then $f(y) = W(x, y) \circ f(x) \circ V(x, y)^{-1}$. For all z such that $x \preceq z$, $W(z, x) \circ f(x) = f(z) \circ V(z, x)$. Then again if the vector spaces are nontrivial we have $f(z) = W(z, x)^{-1} \circ f(x) \circ V(z, x)$.

So for the sink and source $s \preceq x \preceq s'$ we see each of $f(s)$ and $f(s')$ are either 0 or determined by x . Since the set of sinks and sources is discrete with no accumulation points we can use our arguments in the previous paragraph repeatedly and see that each nontrivial $f(y)$ is determined by $f(x)$. Since $\text{Hom}(V(x), W(x)) \cong k$ and every nontrivial $f(y)$ is determined by $f(x)$, we see $\text{Hom}(V, W) \cong k$. \square

Definition 3.1.3. We define $\text{rep}_k(A_{\mathbb{R}})$ as the full subcategory of $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ whose objects are representations V that are finitely generated by indecomposable projectives (listed in Remark 2.4.16).

Lemma 3.1.4. *Let $f : V \rightarrow W$ be a morphism in \mathcal{C} where $\mathcal{C} = \text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$, $\text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$, or $\text{rep}_k(A_{\mathbb{R}})$.*

- *f has a kernel in \mathcal{C} ,*
- *f has a cokernel in \mathcal{C} , and*
- *the image and coimage of f coincide and lie in \mathcal{C} .*

Proof. First note that f is a morphism in $\text{Rep}_k(A_{\mathbb{R}})$. By a dimension argument for $V(x)$, $W(x)$, $\ker f(x)$, and $\text{coker } f(x)$ at each $x \in \mathbb{R}$ the statement must be true for $\mathcal{C} = \text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ and $\mathcal{C} = \text{Rep}_k^{\text{b}}(A_{\mathbb{R}})$.

Now suppose $\mathcal{C} = \text{rep}_k(A_{\mathbb{R}})$. Since $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ is abelian the image and coimage of f coincide. Since $V \twoheadrightarrow \text{im } f$ and V is finitely generated, so is $\text{im } f$. Similarly, since W is finitely generated by some $\bigoplus_{i=1}^n P_i$ there is a surjection $\bigoplus_{i=1}^n P_i \twoheadrightarrow \text{coker } f$.

Suppose $g : \bigoplus Q_i \twoheadrightarrow V$ generates V . Then $\ker(f \circ g)$ is a subrepresentation of a projective; since $\text{Rep}_k^{\text{pwf}}(A_{\mathbb{R}})$ is hereditary this means $\ker(f \circ g)$ is projective. Also $\ker(f \circ g)$ maps to $\ker f$. For any $0 \neq \hat{v} \in \ker f(x)$ there is $v \in V(x)$ from the inclusion. Then there is $\tilde{v} \in \bigoplus Q_i(x)$ that maps to v .

Let $\bigoplus Q'_i = \ker(f \circ g)$. Any projective subrepresentation of a finitely generated projective is finitely generated, so $\bigoplus Q'_i$ is finitely generated. We also know that since $\tilde{v} \mapsto v \mapsto 0$, there exists $\bar{v} \in \bigoplus Q'_i(x)$ that maps to v and so maps to \hat{v} . Thus, $\bigoplus Q'_i \twoheadrightarrow \ker f$ so $\ker f$ is also finitely generated. Therefore, $\ker f$, $\text{im } f$, and $\text{coker } f$ are all generated by finitely generated and so in $\text{rep}_k(A_{\mathbb{R}})$. \square

Lemma 3.1.5. *Let V be a representation in $\text{rep}_k(A_{\mathbb{R}})$. Then V is isomorphic to a finite direct sum of interval indecomposables. Furthermore, $\text{rep}_k(A_{\mathbb{R}})$ is Krull-Schmidt.*

Proof. Suppose V is in $\text{rep}_k(A_{\mathbb{R}})$ and $\bigoplus_{i=1}^n Q_i \rightarrow V$ be a surjective morphism required by Definition 3.1.3.

Since $\dim Q_i(x) \leq 1$ for all $x \in \mathbb{R}$, $\dim V(x) \leq n$ for all $x \in \mathbb{R}$. That is, both Q and V are in $\text{Rep}_k^b(A_{\mathbb{R}})$. By Theorem 2.4.15, V is a direct sum of (*a priori* possibly infinitely many) interval indecomposables.

Since each Q_i is projective, the support of each Q_i contains at most 3 sinks and sources (1 source and 2 sinks). Then, since Q is a finite direct sum, the support of Q itself contains finitely many sinks and sources. Since Q surjects onto V , the support of V must also contain only finitely many sinks and sources.

For contradiction, suppose V is an infinite direct sum of indecomposables. Since V is pointwise finite-dimensional and its support contains finitely many sinks and sources, infinitely many summands must have support that does not contain a sink or a source; i.e. each of these indecomposable's support is bounded by an adjacent sink and source. Since there are only finitely many sinks and sources in the support of V , infinitely many must have support between the same adjacent sink and source.

For each $Q_i = P_a$ for some a (classification in Remark 2.4.16), any indecomposable hit by Q_i must contain a in its support. Since V is pointwise finite dimensional there can only be finitely many such indecomposables. Thus there must be some $Q_i = P_a$ or P_a .

If $Q_i = P_a$ then any indecomposable V_α hit by Q_i has the property that $\text{glb supp } V_\alpha \leq a$. If Q_i hit infinitely many indecomposables there must be infinitely many with support of the form (a, b_α) and the b_α must converge on a . However, V is also in $\text{Rep}_k^b(A_{\mathbb{R}})$ and so this is a contradiction as $\lim \dim V(x)$ as $x \rightarrow a$ from above would ∞ . The same argument holds if $Q_i = P_a$. Therefore, V is the direct sum of finitely many indecomposables. Combined with Theorem 2.3.2 and Lemma 2.3.1 this shows $\text{rep}_k(A_{\mathbb{R}})$ is Krull-Schmidt. \square

Remark 3.1.6. In [24], Sala and Schiffmann prove their category of coherent representations (which they call coherent persistence modules) has similar properties to Theorem 3.0.1. In their paper, *tame* representations have finitely-many places where non-isomorphisms occur in the representation. *Coherent* representations are tame with bounded support and with right continuous dimension functions. Our category $\text{rep}_k(A_{\mathbb{R}})$ is the category of tame representations of $A_{\mathbb{R}}$.

Proposition 3.1.7. *The category $\text{rep}_k(A_{\mathbb{R}})$ is not Artinian.*

Proof. Let P_a be a projective indecomposable (Remark 2.4.16) such that a is not in S . Let $b \in \bar{S}$ such that $b \preceq a$; note $b \neq a$. Then, for every $b \preceq z \preceq a$ such that $b \neq z \neq a$, $P_z \subsetneq P_a$. Furthermore, for any two such z, z' such that $z \preceq z'$, we have $P_z \subsetneq P_{z'} \subsetneq P_a$. Thus, we have an infinite (uncountable!) descending chain and so $\text{rep}_k(A_{\mathbb{R}})$ is not Artinian. \square

Example 3.1.8. Let us return to the representation M in Example 1.3.2. It is an uncountable sum and so not in the category $\text{rep}_k(A_{\mathbb{R}})$. In particular, any surjection onto M by a sum of interval indecomposables would require the source representation to be an uncountable sum as well.

3.2. Properites of $\text{rep}_k(A_{\mathbb{R}})$. We now prove Theorem 3.0.1 (4) and (5).

Proposition 3.2.1. *Let $A_{\mathbb{R}}$ and $A'_{\mathbb{R}}$ be different orientations such that the sinks and sources are unbounded above and below in both $A_{\mathbb{R}}$ and $A'_{\mathbb{R}}$. Then $\text{rep}_k(A_{\mathbb{R}}) \cong \text{rep}_k(A'_{\mathbb{R}})$.*

Proof. We'll define a bijection $F : \mathbb{R} \rightarrow \mathbb{R}$ that induces a bijection on (isomorphism classes of) indecomposables and thus an equivalence of categories. Recall S is the set of sinks and sources of $A_{\mathbb{R}}$ and S' is the set of sinks and sources of $A'_{\mathbb{R}}$. First define the bijection on $S \rightarrow S'$ to be $s_n \mapsto s'_n$. Let $x \in \mathbb{R}$ and $n \in \mathbb{Z}$ such that $s_n < x < s_{n+1}$. Then $x = t \cdot s_n + (1 - t)s_{n+1}$ for some $t \in (0, 1)$. Let $F(x) = t \cdot F(s_n) + (1 - t)F(s_{n+1})$.

This induces a bijection on indecomposables as it is a bijection on \mathbb{R} . In particular, if $x \preceq y$ then $F(x) \preceq F(y)$. If $\text{Hom}(M_{[a,b]}, M_{[c,d]}) \cong k$ in $\text{rep}_k(A_{\mathbb{R}})$ then $a \preceq c$ and $b \preceq d$. Since $F(a) \preceq F(c)$ and $F(b) \preceq F(d)$, the Hom-set from $M_{[F(a), F(b)]}$ to $M_{[F(c), F(d)]}$ is also isomorphic to k . Thus we

have an equivalence on the indecomposables. Since both categories are Krull-Schmidt we have an equivalence of categories. \square

Proposition 3.2.2. *Let P and Q be projective indecomposables in $\text{rep}_k(A_{\mathbb{R}})$ and I and J be injective indecomposables in $\text{rep}_k(A_{\mathbb{R}})$.*

- Any morphism $f : P \rightarrow Q$ is either 0 or mono.
- Any morphism $g : I \rightarrow J$ is either 0 or epi.

Proof. We will prove the first statement; the second is dual. Let $f : P \rightarrow Q$ be a map of indecomposable projectives. By Theorem 2.1.16 and Remark 2.4.16 the image $\text{im } f$ in Q is a subrepresentation and so projective. Since P surjects on to $\text{im } f$ it is a split subrepresentation of P . However, P is indecomposable so $\text{im } f = 0$ or $\text{im } f \cong P$. \square

Below, for each indecomposable representation V in $\text{rep}_k(A_{\mathbb{R}})$ we create two projective representations $P_0(V)$ and $P_1(V)$. In Proposition 3.2.5 we prove that $P_1(V) \rightarrow P_0(V) \rightarrow V$ is the minimal projective presentation of V .

Construction 3.2.3. Let V be an indecomposable in $\text{rep}_k(A_{\mathbb{R}})$ with support $|a, b|$. If V is projective let $P_0(V) = V$ and $P_1(V) = 0$.

Now suppose V is not projective. Recall S is the set of sinks and sources of $A_{\mathbb{R}}$ in \mathbb{R} . Since V is finitely generated $|a, b| \cap S$ is finite. We let $P_0(V)$ be the direct sum of the following indecomposable projectives.

- P_s for all sources s in (a, b) .
- P_a if $a \notin |a, b|$ and there exists $x \preceq a$ in $|a, b|$.
- P_a if $a \in |a, b|$ and there exists $x \preceq a$, $x \neq a$ in $|a, b|$.
- P_b if $b \notin |a, b|$ and there exists $x \preceq b$ in $|a, b|$.
- P_b if $b \in |a, b|$ and there exists $x \preceq b$, $x \neq b$ in $|a, b|$.


We let $P_1(V)$ be the direct sum of the following indecomposable projectives.

- P_s for all sources s in (a, b) .
- P_a if $a \notin |a, b|$ and there exists $a \preceq x$ in $|a, b|$.
- P_a if $a \in |a, b|$.
- P_b if $b \notin |a, b|$ and there exists $b \preceq x$ in $|a, b|$.
- P_b if $b \in |a, b|$.

If a or b is a sink and in $|a, b|$ then the summand P_a or P_b is 0, respectively. We see that both $P_0(V)$ and $P_1(V)$ are nontrivial and finitely generated, so in $\text{rep}_k(A_{\mathbb{R}})$. \diamond

Proposition 3.2.4. *Let V , $P_1(V)$, and $P_0(V)$ be as in Construction 3.2.3. Then there is an injective morphism $P_1(V) \hookrightarrow P_0(V)$ whose cokernel is V .*

Proof. If V is projective the statement is trivially true. Now suppose V is not projective. There are finitely many sinks and sources, totally ordered. So on those summands we let the maps be defined in the following way where \pm means scalar multiplication by ± 1 :



Since there is no accumulation of elements of S in \mathbb{R} , a projective indecomposable at a can only appear as a summand of $P_0(V)$ or $P_1(V)$, but not both. The similar statement is true for b . Thus, only one type of projective summand of each a or b may appear in $P_0(V)$ and $P_1(V)$. Denote whichever summands appear, if any, by P_{a*} and P_{b*} .

If P_{a*} appears in $P_1(V)$ then there is a nontrivial map from P_{a*} to $P_{s_{2n+1}}$ or P_{b*} , depending on whether or not (a, b) contains any sources. If this is the case, use scalar multiplication by -1 . In the similar case for b , use scalar multiplication by $+1$.

If P_{a*} appears in $P_0(V)$ then there is a nontrivial map from $P_{s_{2n}}$ or P_{b*} to P_{a*} , depending on whether or not (a, b) contains any sinks. If this is the case, use scalar multiplication by $+1$. In the similar case for b , use scalar multiplication by -1 .

Instead of proving that this map is injective with cokernel V , we instead note that the kernel of the surjection $P_0(V) \rightarrow V$ is $P_1(V)$. This is equivalent. \square

Proposition 3.2.5. *The following hold:*

- For any indecomposable V in $\text{rep}_k(A_{\mathbb{R}})$, $P_1(V) \hookrightarrow P_0(V) \twoheadrightarrow V$ is the minimal projective resolution and presentation of V .
- All representations in $\text{rep}_k(A_{\mathbb{R}})$ are finitely presented.
- The global dimension of $\text{rep}_k(A_{\mathbb{R}})$ is 1.

Proof. We see $P_1(V)$ is superfluous in $P_0(V)$ and $P_1(V) \hookrightarrow P_0(V) \twoheadrightarrow V$ is exact by Proposition 3.2.4. Thus the sequence is the minimal projective resolution and presentation of V . Furthermore, noting that the reversal of orientation \preceq on \mathbb{R} gives the opposite category, we see the global dimension of $\text{rep}_k(A_{\mathbb{R}})$ is 1. \square

Proposition 3.2.6. *Let V and W be indecomposables in $\text{rep}_k(A_{\mathbb{R}})$. If $\text{Ext}^1(W, V) \neq 0$ then $\text{Ext}^1(W, V) \cong k$.*

Proof. Let V and W be indecomposables in $\text{rep}_k(A_{\mathbb{R}})$. By Proposition 3.2.5 the projective resolution of V is $P_1(V) \hookrightarrow P_0(V) \twoheadrightarrow V$. By definition $\text{Ext}^i(V, W)$ is the i th homology group in the chain

$$0 \longrightarrow \text{Hom}(P_0(V), W) \longrightarrow \text{Hom}(P_1(V), W) \longrightarrow 0.$$

Suppose $\text{Ext}^1(W, V) \neq 0$.

Index the projectives in $P_0(V)$ that nontrivially map to W from 1 to m , denoted P_1, \dots, P_m , such that if $P_a = P_i$ and $P_b = P_{i+1}$ for $a, b \in \mathbb{R}$ then $a < b$. Then $\text{Hom}(P_0(V), W) \cong k^m$. Let $f : (x_1, \dots, x_m)$ be a nontrivial map $P_0(V) \rightarrow W$ and $\iota : P_1(V) \rightarrow P_0(V)$ the inclusion. Index the projectives in $P_1(V)$ that nontrivially map to W from 1 to n , similarly to the projectives in $P_0(V)$, denoted Q_1, \dots, Q_n .

Then Q_1 maps to P_1 and P_2 or just P_1 . If Q_1 only maps to P_1 then the projective Q_2 maps to both P_1 and P_2 . If Q_1 maps to both P_1 and P_2 then Q_2 maps to P_2 and P_3 . Thus, the composition $f \circ \iota$ will be one of four forms:

- $(x_1, x_1 \oplus x_2, \dots, x_{i-1} \oplus x_i),$
- $(x_1 \oplus x_2, \dots, x_{i-1} \oplus x_i, x_i),$
- $(x_1, x_1 \oplus x_2, \dots, x_{i-1} \oplus x_i, x_i),$ or
- $(x_1 \oplus x_2, \dots, x_{i-1} \oplus x_i).$

In any case, basic linear algebra shows us that $\text{Hom}(P_0(V), W) \rightarrow \text{Hom}(P_1(V), W)$ is surjective or injective and the difference in dimensions is either 0 or 1. Therefore $\dim \text{Ext}^1(W, V)$ is 0 or 1. \square

3.3. Existence of Some Auslander–Reiten Sequences. In this subsection we will show that for any orientation of a continuous type A quiver, the category $\text{rep}_k(A_{\mathbb{R}})$ contains some Auslander–Reiten sequences but not all Auslander–Reiten sequences (Theorem 3.0.1 (6)). However, we will not provide a complete classification of Auslander–Reiten sequences in this paper. Such a classification will be provided in the sequel to this paper.

In [17], Gabriel and Roiter provide a general description of Auslander–Reiten sequences of representations of linear posets. However, a specific description to this context in the contemporary language and notation of representation theory is new.

We recall the definition of an almost-split sequence, commonly called an Auslander–Reiten sequence. Such short exact sequences were originally defined by Auslander and Reiten in [2].

Definition 3.3.1. Let \mathcal{A} be an abelian category and $0 \rightarrow U \xrightarrow{f} V \xrightarrow{g} W \rightarrow 0$ a short exact sequence in \mathcal{A} . The short exact sequence is an almost split sequence, or Auslander–Reiten sequence if the following conditions hold:

- f is not a section and g is not a retraction.
- U and W are indecomposable.
- If $h : U \rightarrow X$ is a nontrivial morphism of indecomposables and $U \not\cong X$ then h factors through f .
- If $h : X \rightarrow W$ is a nontrivial morphism of indecomposables and $X \not\cong W$ then h factors through g .

In the following proposition, recall that S is the set of sinks and sources in a continuous quiver of type A and that \bar{S} includes $\pm\infty$.

Proposition 3.3.2. Let $s_n, s_{n+1} \in \bar{S}$ and $a, b \in \mathbb{R}$ such that $s_n < a < b < s_{n+1}$. One of the following is a short exact sequence and in particular an Auslander–Reiten sequence.

- If s_n is a sink then the Auslander–Reiten sequence is

$$0 \longrightarrow M_{[a,b]} \xrightarrow{\begin{bmatrix} 1 \\ 1 \end{bmatrix}} M_{[a,b]} \oplus M_{(a,b)} \xrightarrow{\begin{bmatrix} 1 & -1 \end{bmatrix}} M_{(a,b)} \longrightarrow 0$$

- If s_n is a source then the Auslander–Reiten sequence is

$$0 \longrightarrow M_{(a,b)} \xrightarrow{\begin{bmatrix} 1 \\ 1 \end{bmatrix}} M_{(a,b)} \oplus M_{[a,b]} \xrightarrow{\begin{bmatrix} 1 & -1 \end{bmatrix}} M_{[a,b]} \longrightarrow 0$$

Proof. We note the two cases are symmetric and prove the first. We see the first map is injective, the second is surjective, and that the sequence is exact at $[a, b] \oplus (a, b)$. Thus, the sequence is a short exact sequence.

Denote the map $M_{[a,b]} \rightarrow M_{[a,b]} \oplus M_{(a,b)}$ in the sequence by $h_1 \oplus h_2$. By Proposition 2.2.1 we know both $M_{[a,b]}$ and $M_{(a,b)}$ are indecomposable. Let V be another indecomposable representation in $\text{rep}_k(A_{\mathbb{R}})$. By definition the support of V is an interval $|c, d|$. If there exists $x \in |c, d|$ such that $x < a$ then any morphism $f : M_{[a,b]} \rightarrow V$ must be 0. Additionally, if there exists $x \in [a, b]$ such that $x \geq d$ and $x \notin |c, d|$ then any $f : M_{[a,b]} \rightarrow V$ must be 0. Thus, any morphism $M_{[a,b]} \oplus M_{(a,b)} \rightarrow M_{[a,b]}$ must be 0 and morphism $M_{(a,b)} \rightarrow M_{[a,b]} \oplus M_{(a,b)}$ must be 0.

Claim: If $V \not\cong M_{[a,b]}$ and $f : M_{[a,b]} \rightarrow V$ is a nonzero morphism then there exists either a nonzero morphism $g_1 : M_{[a,b]} \rightarrow V$ or $g_2 : M_{(a,b)} \rightarrow V$ such that $g_i \circ h_i = f$. Proof of claim: If $V \not\cong M_{[a,b]}$ then, by the conditions in the previous paragraph combined with Theorem 2.3.2, either $b \in |c, d|$ or $a \notin |c, d|$. If $b \in |c, d|$ Then g_1 is a nonzero morphism and so $g_1 \circ h_1 = f$. If $a \notin |c, d|$ then g_2 is a nonzero morphism and so $g_2 \circ h_2 = f$. In either case, f factors through $M_{[a,b]} \oplus M_{(a,b)}$.

Finally, if $|c, d| = [a, b]$ then by Theorem 2.3.2 f is an isomorphism. By a dual argument, a morphism from an indecomposable W to $M_{(a,b)}$ that is not an isomorphism factors through $M_{[a,b]} \oplus M_{(a,b)}$. Therefore, the given sequence is an Auslander–Reiten sequence. \square

We give an example of a representation with no left or right Auslander–Reiten sequences in the form of a proposition.

Proposition 3.3.3. Let $M_{\{a\}}$ be the indecomposable representation with support $\{a\}$ where a is neither a sink nor a source. Then there is no Auslander–Reiten sequences of either of the

following forms:

$$\begin{aligned} 0 &\longrightarrow M_{\{a\}} \longrightarrow B \longrightarrow C \longrightarrow 0 \\ 0 &\longrightarrow A \longrightarrow B \longrightarrow M_{\{a\}} \longrightarrow 0. \end{aligned}$$

Proof. Suppose $s_{2n} < a < s_{2n+1}$, where s_{2n} is a sink and s_{2n+1} is a source. The other case is similar. For any indecomposable M_I , if $\text{Hom}(M_I, M_{\{a\}}) \cong k$ then $I = [c, a]$. We note that, for each $x \in (s_0, a)$, $\text{Hom}(M_{[x, a]}, M_{\{a\}}) \cong k$ for all $i \geq 0$.

Let M_I be some indecomposable such that $\text{Hom}(M_I, M_{\{a\}}) \cong k$. For any $x \in (s_0, a)$ such that $c < x$ we have $\text{Hom}(M_I, M_{J_x}) \cong k$. Since the Hom space between any two indecomposables is either k or 0 (Proposition 3.1.2), all nontrivial maps $M_I \rightarrow M_{\{a\}}$ factor through every indecomposable $M_{[x, a]}$ for $x \in (s_0, a)$ and $x > c$. Thus, it is not possible to have an Auslander–Reiten sequence in $\text{rep}_k(A_{\mathbb{R}})$ of the form $0 \rightarrow A \rightarrow B \rightarrow M_{\{a\}} \rightarrow 0$. By a dual argument, the other form is not possible, either. \square

4. OTHER PAPERS IN THIS SERIES

In Continuous Quivers of Type A (II), the second author defines a continuous analog of the Auslander–Reiten quiver, called the Auslander–Reiten Space, for both $\text{rep}_k(A_{\mathbb{R}})$ and its bounded derived category $\mathcal{D}^b(\text{rep}_k(A_{\mathbb{R}}))$ [22]. They show that the Auslander–Reiten space exhibits many of the same properties as an Auslander–Reiten quiver, such as how to find extensions of indecomposables and Auslander–Reiten sequences. Similar results are shown about the derived category.

The authors define the new continuous cluster category in Continuous Quivers of Type A (III) and generalize cluster structures to cluster theories [18]. In particular, they define the **E**-cluster theory. They then show many existing type A cluster structures are cluster theories and embed into this new theory in a way that preserves mutation.

In Continuous Quivers of Type A (IV), the second author generalizes mutation to continuous mutation [23], further generalizing transfinite mutation in [3]. The embeddings from Part (III) are shown to be part of a chain of embeddings and the notion of an abstract cluster structure is introduced in order to understand which cluster theories are more strongly related. Part (IV) concludes with a geometric model of **E**-clusters which generalizes the triangulations of polygons and laminations of hyperbolic plane in [7] and [19], respectively.

REFERENCES

- [1] H. Adams, M. Aminian, E. Farnell, M. Kirby, J. Mirth, R. Neville, C. Peterson, P. Shipman, and C. Shonkwiler, *A fractal dimension for measures via persistent homology*, to appear in Abel Symposia 2019, preprint: <https://arxiv.org/pdf/1808.01079.pdf>
- [2] M. Auslander and I. Reiten, *Representation theory of Artin algebras. III. Almost split sequences*, Communications in Algebra **3** (1975), no. 3, 239–294, DOI: 10.1080/00927877508822046
- [3] K. Baur and S. Gratz, *Transfinite mutations in the completed infinity-gon*, Journal of Combinatorial Series A **155** (2018), 321–359, DOI: 10.1016/j.jcta.2017.11.011
- [4] M.B. Botnan, *Interval Decomposition of Infinite Zigzag Persistence Modules*, Proceedings of the American Mathematical Society **145** (2017), no. 8, 3571–3577, DOI: 10.1090/proc/13465
- [5] M.B. Botnan and W. Crawley-Boevey, *Decomposition of persistence modules*, to appear in Proceedings of the American Mathematical Society, preprint: <https://arxiv.org/pdf/1811.08946.pdf>
- [6] A. Buan, R. Marsh, M. Reineke, I. Reiten, and G. Todorov, *Tilting theory and cluster combinatorics*, Advances in Mathematics **204** (2006), no. 2, 572–618, DOI: 10.1016/j.aim.2005.06.003
- [7] P. Caldero, F. Chapoton, and R. Schiffler, *Quivers with Relations Arising From Clusters (A_n Case)*, Transactions of the American Mathematical Society, Volume 358, Number 3, 1347 – 1364
- [8] G. Carlsson, V. de Silva, and D. Morozov, *Zigzag persistent homology and real-valued functions*, Proceedings of the twenty-fifth annual symposium on computational geometry, 2009, 247–256, DOI: 10.1145/1542362.1542408
- [9] G. Carlsson, T. Ishkhanov, V. de Silva, and A. Zomorodian, *On the local behavior of spaces of natural images*, International Journal of Computer Vision **76** 2008, no. 1, 1–12, DOI: 10.1007/s11263-007-0056-x

- [10] F. Chazal, D. Cohen-Steiner, L. J. Guibas, F. Mémoli, and S. Y. Oudot. *Gromov-Hausdorff Stable Signatures for Shapes using Persistence*, Computer Graphics Forum, **28** (2009), no. 5, 1393–1403, DOI: 10.1111/j.1467-8659.2009.01516.x
- [11] F. Chazal and S. Oudot, *Towards persistence-based reconstruction in Euclidean spaces*, In Proceedings of the 24th Annual Symposium on Computational Geometry (2008), 232–241, DOI: 10.1145/1377676.1377719
- [12] W. Crawley-Boevey, *Decomposition of pointwise finite-dimensional persistence modules*, Journal of Algebra and its Applications **14** (2015), no. 5, DOI: 10.1142/S0219498815500668
- [13] H. Edelsbrunner and J. L. Harer, Computational topology: an Introduction, American Mathematical Society, 2010
- [14] E. J. Hanson and J. D. Rock, *Decomposition of Pointwise Finite-Dimensional \mathbb{S}^1 Persistence Modules*, arXiv:2006.13793 [math.RT] (2020), <https://arXiv.org/pdf/arXiv:2006.13793>
- [15] S. Fomin and A. Zelevinsky, *Cluster algebras I: Foundations*, Journal of the American Mathematical Society **15** (2002), no. 2, 497–529, DOI:10.1090/S0894-0347-01-00385-X
- [16] P. Gabriel, *Unzerlegbare Darstellungen. I*, Manuscripta Mathematica **6** (1972), 71–103, DOI:10.1007/BF01298413
- [17] P. Gabriel and A. V. Roiter, Representations of Finite-Dimensional Algebras, Springer-Verlag, Berlin Heidelberg, 1997
- [18] K. Igusa, J. D. Rock, and G. Todorov, *Continuous Quivers of Type A (III) Embeddings of Cluster Theories*, arXiv:2004.10740 [math.RT] (2020), <https://arXiv.org/pdf/2004.10740>
- [19] K. Igusa and G. Todorov, *Continuous Cluster Categories I*, Algebras and Representation Theory **18** (2015), no. 1, 65–101, DOI: 10.1007/s10468-014-9481-z
- [20] J. Jaquette and B. Schweinhart, *Fractal Dimension Estimation with Persistent Homology: A Comparative Study*, arXiv:1907.11182v2 [math.DS] (2019), <https://arxiv.org/pdf/1907.11182v2.pdf>
- [21] M. Nicolau, A. J. Levine, and G. Carlsson, *Topology based data analysis identifies a subgroup of breast cancers with a unique mutational profile and excellent survival*, Proceedings of the National Academy of Sciences **108** (2011), no. 17, 7265–7270, DOI: 10.1073/pnas.1102826108
- [22] J. D. Rock, *Continuous Quivers of Type A (II) The Auslander-Reiten Space*, arXiv:1910.04140v1 [math.RT] (2019), <https://arXiv.org/pdf/1910.04140.pdf>
- [23] —, *Continuous Quivers of Type A (IV) Continuous Mutation and Geometric Models of \mathbf{E} -clusters*, arXiv:2004.11341 [math.RT] (2020), <https://arXiv.org/pdf/2004.11341>
- [24] F. Sala and O. Schiffmann, *Fock space representation of the circle quantum group*, arXiv:1903.02813v1 [math.QA] (2019), <https://arxiv.org/pdf/1903.02813.pdf>
- [25] B. Schweinhart, *Fractal Dimension and the Persistent Homology of Random Geometric Complexes*, arXiv:1808.02196v5 [math.PR], <https://arxiv.org/pdf/1808.02196.pdf>
- [26] A. Zomorodian and G. Carlsson, *Computing Persistent Homology*, Discrete and Computational Geometry **33** (2005), no. 2, 249–274, DOI: 10.1007/s00454-004-1146-y