

ON THE INTEGRABILITY OF THE WAVE PROPAGATOR ARISING FROM THE LIOUVILLE-VON NEUMANN EQUATION

YOUNGWOON KOH, YOONJUNG LEE AND IHYEOK SEO

ABSTRACT. The Liouville-von Neumann equation describes the change in the density matrix with time. Interestingly, this equation was recently regarded as a wave equation for wave functions but not a equation for density functions. This setting leads to an extended form of the Schrödinger wave equation governing the motion of a quantum particle. In this paper we obtain the integrability of the wave propagator arising from the Liouville-von Neumann equation in this setting.

1. INTRODUCTION

The pure state of a quantum system is described by a wave function obeying the Schrödinger equation $i\partial_t\psi + \Delta\psi = 0$. For the description of mixed states, the notion of density matrix was introduced by von Neumann [17] (see also [13]). This density matrix for a pure state is equal to the product of the wave function and its complex conjugate at different arguments. Motivated by this, one can consider

$$\begin{aligned} i\frac{\partial}{\partial t}(\psi(x, t)\bar{\psi}(y, t)) &= i\frac{\partial\psi(x, t)}{\partial t}\bar{\psi}(y, t) + i\frac{\partial\bar{\psi}(y, t)}{\partial t}\psi(x, t) \\ &= -\Delta_x\psi(x, t)\bar{\psi}(y, t) + \Delta_y\bar{\psi}(y, t)\psi(x, t) \\ &= (-\Delta_x + \Delta_y)(\psi(x, t)\bar{\psi}(y, t)), \end{aligned}$$

which leads to the following equation

$$i\partial_t\Psi(x, y, t) + (\Delta_x - \Delta_y)\Psi(x, y, t) = 0 \quad (1.1)$$

where $(x, y, t) \in \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}$.

This equation is commonly called the Liouville-von Neumann equation in physical literature, giving the change in the density matrix with time ([18, 14]). However, it is more interesting to regard (1.1) as a wave equation for wave functions but not a equation for density functions. This is the key point in a recent work by Chen [3]. Contrary to Schrödinger's wave functions, the wave functions $\Psi(x, y, t)$ of the Liouville-von Neumann equation (1.1) for a single particle are *bipartite*. These *bipartite* wave functions satisfy all the basic properties of Schrödinger's wave functions which correspond to those *bipartite* wave functions of product forms. Indeed, the

2010 *Mathematics Subject Classification.* Primary: 35B45; Secondary: 35Q40.

Key words and phrases. Strichartz estimates, von Neumann-Landau equation.

Y. Koh was supported by NRF-2019R1F1A1054310. I. Seo was supported by NRF-2019R1F1A1061316.

Schrödinger equation is a special case of the equation (1.1) with the initial data of product form, $\Psi(x, y, 0) = \psi(x)\bar{\psi}(y)$, because in this case $\Psi(x, y, t) = \psi(x, t)\bar{\psi}(y, t)$ with $\psi(x, t)$ satisfying the Schrödinger equation with the initial data $\psi(x, 0) = \psi(x)$ and vice versa. This extension of Schrödinger's form establishes a mathematical expression of wave-particle duality and that von Neumann's entropy is a quantitative measure of complementarity between wave-like and particle-like behaviors. Furthermore, it suggests that collapses of Schrödinger's wave functions are just the simultaneous transition of the particle from many levels to one. See [3] for details. The equation considered as a wave function equation is also explicitly used in [12], connecting with Bose-Einstein condensation.

The problem we want to discuss in this paper is integrability of wave propagator $e^{it(\Delta_x - \Delta_y)}$ which gives a formula for the solution to the Liouville-von Neumann equation. Applying the Fourier transform to (1.1), the solution $\Psi(x, y, t)$ is indeed given by

$$e^{it(\Delta_x - \Delta_y)} f(x, y) := \frac{1}{(2\pi)^{2n}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{ix \cdot \xi + iy \cdot \xi' - it(|\xi|^2 - |\xi'|^2)} \hat{f}(\xi, \xi') d\xi d\xi',$$

where f is the initial data $\Psi(x, y, 0)$ and \hat{f} is the Fourier transform thereof. Our result is stated as follows.

Theorem 1.1. *Let $n \geq 1$. Assume that $2 \leq q \leq \infty$, $2 \leq r_2 \leq r_1 \leq \infty$ and*

$$\frac{2}{q} = n \left(1 - \frac{1}{r_1} - \frac{1}{r_2} \right). \quad (1.2)$$

Then we have

$$\|e^{it(\Delta_x - \Delta_y)} f\|_{L_t^q L_x^{r_1} L_y^{r_2}} \lesssim \|f\|_{L^2}, \quad (1.3)$$


except for $(q, r_1, r_2) = (2, \infty, \infty)$ when $n = 1$ and for $(q, r_1, r_2) = (2, \infty, 2)$ when $n = 2$.

Compared with (1.3), the space-time integrability known as *Strichartz estimates* for the Schrödinger case has been extensively studied over the last several decades and is now completely understood as follows (see [20, 8, 16, 11]):

$$\|e^{it\Delta} f\|_{L_t^q L_x^r} \lesssim \|f\|_{L^2}$$

if and only if (q, r) is n -Schrödinger-admissible, i.e., $q, r \geq 2$, $(q, r, n) \neq (2, \infty, 2)$ and $2/q + n/r = n/2$. Particularly when $r_1 = r_2$, the exponent pair (q, r_1, r_2) in the theorem becomes $2n$ -Schrödinger-admissible. In this case, (1.3) can be found in [15] and the range of (r_1, r_2) is given by the closed segment $[D, B]$ in Figure 1 below. But, it is significant in the *bipartite* form to quantify the spatial integrability differently with respect to x and y . In this regard, the main contribution of the theorem is to extend the diagonal case $r_1 = r_2$ to mixed norms $L_x^{r_1} L_y^{r_2}$, $r_1 \neq r_2$, where the range of (r_1, r_2) is given by the closed triangle with vertices A, D, B .

Notice that the condition (1.2) is necessary for (1.3) to be invariant under the scaling $(x, y, t) \rightarrow (\lambda x, \lambda y, \lambda^2 t)$, $\lambda > 0$. By the standard TT^* method, (1.3) is also equivalent to the boundedness of the time translation invariant operator $TT^* : F \rightarrow \int_{\mathbb{R}} e^{i(t-s)(\Delta_x - \Delta_y)} F(s) ds$ from $L_t^q L_x^{r_1} L_y^{r_2}$ to $L_t^{q'} L_x^{r_1'} L_y^{r_2'}$. Hence, $q \geq q'$ (i.e., $q \geq 2$) is


 FIGURE 1. The range of (r_1, r_2) for which (1.3) holds when $n \geq 3$.

required (see [10] or [9]). When $q = 2$ and $q = \infty$, $(1/r_1, 1/r_2)$ lies on the line through the points A, D and on the point B , respectively.

In the final section we also give some applications of Theorem 1.1 to nonlinear problems.

Throughout this paper, we use $\mathcal{F}f$ and \hat{f} to denote the Fourier transform of f and $\langle f, g \rangle_{x,y}$ denotes the usual inner product on $L^2_{x,y}$. We also denote $A \lesssim B$ to mean $A \leq CB$ with unspecified constant $C > 0$ which may be different at each occurrence.

2. THE NON-ENDPOINT CASE $q > 2$

In this section we prove Theorem 1.1 when $q > 2$. The endpoint case $q = 2$ will be proved in the next section.

When $q = \infty$, the estimate (1.3) follows directly from the Plancherel theorem as follows:

$$\|e^{it(\Delta_x - \Delta_y)} f\|_{L_t^\infty L_{x,y}^2} = \sup_{t \in \mathbb{R}} \|e^{-it(|\xi|^2 - |\xi'|^2)} \hat{f}\|_{L_{\xi, \xi'}^2} = \|\hat{f}\|_{L_{\xi, \xi'}^2} = \|f\|_{L_{x,y}^2}.$$

Now we only need to consider $2 < q < \infty$. By the standard TT^* argument, we may prove the following estimate

$$\left\| \int_{\mathbb{R}} e^{i(t-s)(\Delta_x - \Delta_y)} F(s) ds \right\|_{L_t^q L_x^{r_1} L_y^{r_2}} \lesssim \|F\|_{L_t^{q'} L_x^{r'_1} L_y^{r'_2}} \quad (2.1)$$

which is equivalent to (1.3). To show this, we obtain the following fixed-time estimates for the propagator $e^{it(\Delta_x - \Delta_y)}$.

Lemma 2.1. *Let $n \geq 1$ and $2 \leq r_2 \leq r_1 \leq \infty$. Then we have*

$$\|e^{it(\Delta_x - \Delta_y)} G\|_{L_x^{r_1} L_y^{r_2}} \lesssim |t|^{-n(1 - \frac{1}{r_1} - \frac{1}{r_2})} \|G\|_{L_x^{r'_1} L_y^{r'_2}}. \quad (2.2)$$

Assuming for the moment this lemma, we see that

$$\begin{aligned} \left\| \int_{\mathbb{R}} e^{i(t-s)(\Delta_x - \Delta_y)} F(s) ds \right\|_{L_t^q L_x^{r_1} L_y^{r_2}} &\leq \left\| \int_{\mathbb{R}} \|e^{i(t-s)(\Delta_x - \Delta_y)} F(s)\|_{L_x^{r_1} L_y^{r_2}} ds \right\|_{L_t^q} \\ &\lesssim \left\| \int_{\mathbb{R}} |t-s|^{-n(1-\frac{1}{r_1}-\frac{1}{r_2})} \|F(s)\|_{L_x^{r'_1} L_y^{r'_2}} ds \right\|_{L_t^q}. \end{aligned}$$

Here we use the Hardy-Littlewood-Sobolev inequality ([19], Section V.1.2) for dimension one,

$$\| |t|^{-\alpha} * g \|_{L^q} \lesssim \|g\|_{L^p}, \quad (2.3)$$

where $0 < \alpha < 1$, $1 \leq p < q < \infty$ and $1/q + 1 = 1/p + \alpha$. By applying (2.3) with $p = q'$ and $\alpha = n(1 - 1/r_1 - 1/r_2)$ to the above, we obtain the estimate (2.1) if $2 < q < \infty$, $2 \leq r_1 \leq r_2 \leq \infty$ and $2/q = n(1 - 1/r_1 - 1/r_2)$, as desired.

Proof of Lemma 2.1. It remains to prove the lemma. By the Riesz-Thorin interpolation theorem, it suffices to show (2.2) for the following three cases:

- (a) $r_1 = r_2 = \infty$,
- (b) $r_1 = \infty$ and $r_2 = 2$,
- (c) $r_1 = r_2 = 2$.

First we write

$$\begin{aligned} e^{it(\Delta_x - \Delta_y)} G(x, y) &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} K_t(x - x') K_{-t}(y - y') G(x', y') dx' dy' \\ &= K_{-t} *_{y'} (K_t *_{x'} G) \end{aligned}$$

where

$$K_t(x) = \frac{1}{(4\pi it)^{n/2}} e^{\frac{ix|^2}{4t}}$$

denotes the integral kernel for the Schrödinger propagator. Then one can see that

$$\|K_t * g\|_{L^\infty} \lesssim |t|^{-\frac{n}{2}} \|g\|_{L^1} \quad (2.4)$$

by Young's inequality and that

$$\|K_t * g\|_{L^2} = \|e^{it\Delta} g\|_{L^2} = \|g\|_{L^2} \quad (2.5)$$

by the Plancherel theorem.

Applying (2.4) repeatedly together with the Minkowski inequality, we obtain the first case (a) as follows:

$$\begin{aligned} \|e^{it(\Delta_x - \Delta_y)} G\|_{L_{x,y}^\infty} &= \left\| \|K_{-t} *_{y'} (K_t *_{x'} G)\|_{L_y^\infty} \right\|_{L_x^\infty} \\ &\lesssim |t|^{-\frac{n}{2}} \left\| \|K_t *_{x'} G\|_{L_y^1} \right\|_{L_x^\infty} \\ &\lesssim |t|^{-\frac{n}{2}} \left\| \|K_t *_{x'} G\|_{L_x^\infty} \right\|_{L_y^1} \\ &\lesssim |t|^{-n} \|G\|_{L_{x,y}^1}. \end{aligned}$$

For the second case (b), we use (2.5) and (2.4) along with the Minkowski inequality to get

$$\begin{aligned}
 \|e^{it(\Delta_x - \Delta_y)} G\|_{L_x^\infty L_y^2} &= \|\|K_{-t} * y(K_t * x G)\|_{L_y^2}\|_{L_x^\infty} \\
 &\lesssim \|\|K_t * x G\|_{L_y^2}\|_{L_x^\infty} \\
 &\lesssim \|\|K_t * x G\|_{L_x^\infty}\|_{L_y^2} \\
 &\lesssim |t|^{-\frac{n}{2}} \|\|G\|_{L_x^1}\|_{L_y^2} \\
 &\lesssim |t|^{-\frac{n}{2}} \|G\|_{L_x^1 L_y^2}.
 \end{aligned}$$

Lastly the case (c) follows directly from the Plancherel theorem as

$$\|e^{it(\Delta_x - \Delta_y)} G\|_{L_{x,y}^2} = \|e^{-it(|\xi|^2 - |\xi'|^2)} \widehat{G}\|_{L_{\xi,\xi'}^2} = \|G\|_{L_{x,y}^2}.$$

□

3. THE ENDPOINT CASE $q = 2$

It remains to prove (1.3) when $q = 2$. Following [11], we will obtain the estimate by a bilinear interpolation between the nonendpoint results and the decay estimates for a time-localized bilinear form operator. In this argument we can take advantage of the symmetry and the flexibility of the bilinear form setting.

By the standard TT^* method we may prove

$$\left\| \int_{\mathbb{R}} e^{i(t-s)(\Delta_x - \Delta_y)} F(s) ds \right\|_{L_t^q L_x^{r_1} L_y^{r_2}} \lesssim \|F\|_{L_t^{q'} L_x^{r_1'} L_y^{r_2'}},$$

and by duality this is in turn equivalent to the bilinear form estimate

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left\langle e^{-is(\Delta_x - \Delta_y)} F(s), e^{-it(\Delta_x - \Delta_y)} G(t) \right\rangle_{x,y} ds dt \lesssim \|F\|_{L_t^2 L_x^{r_1'} L_y^{r_2'}} \|G\|_{L_t^2 L_x^{r_1} L_y^{r_2}}.$$

By symmetry it suffices to restrict our attention to the retarded region

$$\Omega = \{(s, t) \in \mathbb{R}^2 : s < t\}$$

in the above double integral. Now we break Ω into a series of time-localized regions using a Whitney type decomposition (see [19] or [7]); let \mathcal{Q}_j be the family of dyadic squares in Ω whose side length is dyadic number 2^j for $j \in \mathbb{Z}$. Each square $Q = I \times J \in \mathcal{Q}_j$ has the property that

$$2^j \sim |I| \sim |J| \sim \text{dist}(I, J) \tag{3.1}$$

and $\Omega = \cup_{j \in \mathbb{Z}} \cup_{Q \in \mathcal{Q}_j} Q$ where the squares Q are essentially disjoint.

Hence we are reduced to showing the following estimate

$$\sum_{j \in \mathbb{Z}} \sum_{Q \in \mathcal{Q}_j} |T_j(F, G)| \lesssim \|F\|_{L_t^2 L_x^{r_1'} L_y^{r_2'}} \|G\|_{L_t^2 L_x^{r_1} L_y^{r_2}} \tag{3.2}$$

where

$$T_j(F, G) := \int_J \int_I \left\langle e^{-is(\Delta_x - \Delta_y)} F(s), e^{-it(\Delta_x - \Delta_y)} G(t) \right\rangle_{x,y} ds dt. \tag{3.3}$$

To get this estimate, we make use of the following two-parameter family of estimates in which the case $r_1 = r_2$ is excluded but this is harmless because the estimate (1.3) is already known for this case ([15]).

Proposition 3.1. *Let $n \geq 1$. Assume that $2 \leq r_2 < r_1 \leq \infty$ and*

$$1 = n(1 - \frac{1}{r_1} - \frac{1}{r_2}).$$

Then we have

$$\sum_{Q \in \mathcal{Q}_j} |T_j(F, G)| \lesssim 2^{-j\beta(a, \tilde{a})} \|F\|_{L_t^2 L_x^{a'} L_y^{r'_2}} \|G\|_{L_t^2 L_x^{\tilde{a}'} L_y^{r'_2}} \quad (3.4)$$

for all $j \in \mathbb{Z}$ and all $(1/a, 1/\tilde{a})$ in a neighborhood of $(1/r_1, 1/r_1)$ (see Figure 2) with

$$\beta(a, \tilde{a}) = -1 + \frac{n}{2}(2 - \frac{1}{a} - \frac{1}{\tilde{a}} - \frac{2}{r_2}).$$

Assuming for the moment this proposition which will be obtained in the next section, we now get (3.2) using a bilinear interpolation argument. Let us first consider the bilinear vector-valued operator B as

$$B(F, G) = \left\{ \sum_{Q \in \mathcal{Q}_j} T_j(F, G) \right\}_{j \in \mathbb{Z}}.$$

Then, (3.2) can be rewritten as

$$B : L_t^2 L_x^{r'_1} L_y^{r'_2} \times L_t^2 L_x^{r'_1} L_y^{r'_2} \rightarrow \ell_1^0 \quad (3.5)$$

where ℓ_p^α for $\alpha \in \mathbb{R}$ and $1 \leq p \leq \infty$ denotes the weighted sequence space with the norm

$$\|\{x_j\}_{j \geq 0}\|_{\ell_p^\alpha} = \begin{cases} (\sum_{j \geq 0} 2^{j\alpha p} |x_j|^p)^{\frac{1}{p}}, & \text{if } p \neq \infty, \\ \sup_{j \geq 0} 2^{j\alpha} |x_j|, & \text{if } p = \infty. \end{cases}$$

For a sufficient small $\varepsilon > 0$, we now choose $\frac{1}{a_0} = \frac{1}{r_1} - \varepsilon$ and $\frac{1}{a_1} = \frac{1}{r_1} + 2\varepsilon$. Note here that we cannot choose $a_0, a_1 \geq 0$ if $r_1 = \infty$ which corresponds to the cases where $(q, r_1, r_2) = (2, \infty, \infty)$ when $n = 1$ and $(q, r_1, r_2) = (2, \infty, 2)$ when $n = 2$. For this reason these cases are excluded in the theorem. Since $\beta(a_0, a_0) = 2n\varepsilon$ and $\beta(a_0, a_1) = \beta(a_1, a_0) = -n\varepsilon$, Proposition 3.1 implies

$$B : L_t^2 L_x^{a'_0} L_y^{r'_2} \times L_t^2 L_x^{a'_0} L_y^{r'_2} \rightarrow \ell_\infty^{2n\varepsilon},$$

$$B : L_t^2 L_x^{a'_0} L_y^{r'_2} \times L_t^2 L_x^{a'_1} L_y^{r'_2} \rightarrow \ell_\infty^{-n\varepsilon},$$

$$B : L_t^2 L_x^{a'_1} L_y^{r'_2} \times L_t^2 L_x^{a'_1} L_y^{r'_2} \rightarrow \ell_\infty^{-n\varepsilon}.$$

Now we apply the following bilinear interpolation lemma with $s = 1$, $p = q = r'_1$ and $\theta_0 = \theta_1 = 1/3$ to obtain

$$B : (L_t^2 L_x^{a'_0} L_y^{r'_2}, L_t^2 L_x^{a'_1} L_y^{r'_2})_{\frac{1}{3}, r'_1} \times (L_t^2 L_x^{a'_0} L_y^{r'_2}, L_t^2 L_x^{a'_1} L_y^{r'_2})_{\frac{1}{3}, r'_1} \rightarrow (\ell_\infty^{2n\varepsilon}, \ell_\infty^{-n\varepsilon})_{\frac{2}{3}, 1}. \quad (3.6)$$

Here, $(\cdot, \cdot)_{(\theta, p)}$ denotes the real interpolation functor.

Lemma 3.2. ([1], Section 3.13.5(b)) Let $A_0, A_1, B_0, B_1, C_0, C_1$ are Banach spaces and B be a bilinear operator such that $B : A_0 \times B_0 \rightarrow C_0$, $B : A_0 \times B_1 \rightarrow C_1$ and $B : A_1 \times B_0 \rightarrow C_1$. Then

$$B : (A_0, A_1)_{\theta_0, ps} \times (B_0, B_1)_{\theta_1, qs} \rightarrow (C_0, C_1)_{\theta, s}$$

if $0 < \theta_0, \theta_1 < \theta = \theta_0 + \theta_1$, $1 \leq p, q, s \leq \infty$ and $1 \leq 1/p + 1/q$.

Finally by making use of the real interpolation space identities in Lemma 3.3 below, we easily see that

$$(L_t^2 L_x^{a'_0} L_y^{r'_2}, L_t^2 L_x^{a'_1} L_y^{r'_2})_{1/3, r'_1} = L_t^2 L_x^{r'_1} L_y^{r'_2}$$

and

$$(\ell_\infty^{2n\varepsilon}, \ell_\infty^{-n\varepsilon})_{2/3, 1} = \ell_1^0$$

in (3.6). This implies (3.5) as desired.

Lemma 3.3. ([1]) Let $0 < \theta < 1$, $1 \leq p_0, p_1 \leq \infty$ and $s_0, s_1 \in \mathbb{R}$. Then

$$(L^{p_0}(A_0), L^{p_1}(A_1))_{\theta, p} = L^p((A_0, A_1)_{\theta, p})$$

for two complex Banach spaces A_0, A_1 and $1/p = (1 - \theta)/p_0 + \theta/p_1$, and

$$(\ell_{p_0}^{s_0}, \ell_{p_1}^{s_1})_{\theta, p} = \ell_p^s$$

if $s_0 \neq s_1$ and $s = (1 - \theta)s_0 + \theta s_1$.

4. PROOF OF PROPOSITION 3.1

To get (3.4), we only need to show

$$|T_j(F, G)| \lesssim 2^{-j\beta(a, \tilde{a})} \|F\|_{L_t^2(I; L_x^{a'} L_y^{r'_2})} \|G\|_{L_t^2(J; L_x^{\tilde{a}'} L_y^{r'_2})} \quad (4.1)$$


for each square $Q = I \times J \in \mathcal{Q}_j$. Using the fact that for each I there are at most a fixed finite number of intervals J which satisfy (3.1) and they are all contained in a neighborhood of I of size $O(2^j)$, we indeed get

$$\begin{aligned} \sum_{Q \in \mathcal{Q}_j} |T_j(F, G)| &\lesssim 2^{-j\beta(a, \tilde{a})} \sum_{Q \in \mathcal{Q}_j} \|F\|_{L_t^2(I; L_x^{a'} L_y^{r'_2})} \|G\|_{L_t^2(I; L_x^{\tilde{a}'} L_y^{r'_2})} \\ &\leq 2^{-j\beta(a, \tilde{a})} \left(\sum_{Q \in \mathcal{Q}_j} \|F\|_{L_t^2(I; L_x^{a'} L_y^{r'_2})}^2 \right)^{\frac{1}{2}} \cdot \left(\sum_{Q \in \mathcal{Q}_j} \|G\|_{L_t^2(I; L_x^{\tilde{a}'} L_y^{r'_2})}^2 \right)^{\frac{1}{2}} \\ &\lesssim 2^{-j\beta(a, \tilde{a})} \|F\|_{L_t^2(\mathbb{R}; L_x^{a'} L_y^{r'_2})} \|G\|_{L_t^2(\mathbb{R}; L_x^{\tilde{a}'} L_y^{r'_2})} \end{aligned}$$

as desired.

We shall now show (4.1) for the following three cases (see Figure 2):

- (a) $a = \tilde{a} = \infty$ (point E),
- (b) $r_2 \leq a < r_1$ and $\tilde{a} = 2$ (segment $(A, B]$),
- (c) $a = 2$ and $r_2 \leq \tilde{a} < r_1$ (segment $(D, C]$).

FIGURE 2. The range of (a, \tilde{a}) for which (3.4) holds when $n \geq 3$.

The proposition will then follow by interpolation with the range of (a, \tilde{a}) as in Figure 2.

To show (a), we first use Hölder's inequality in x, y and the decay estimate (2.2) with the fact that $|t - s| \sim 2^j$;

$$\begin{aligned}
 |T_j(F, G)| &= \left| \int_J \int_I \left\langle e^{i(t-s)(\Delta_x - \Delta_y)} F(s), G(t) \right\rangle_{x,y} ds dt \right| \\
 &\leq \int_J \int_I \|e^{i(t-s)(\Delta_x - \Delta_y)} F(s)\|_{L_x^\infty L_y^{r_2}} \|G(t)\|_{L_x^1 L_y^{r_2'}} ds dt \\
 &\lesssim 2^{-nj(1 - \frac{1}{r_2})} \|F\|_{L_t^1(I; L_x^1 L_y^{r_2'})} \|G\|_{L_t^1(J; L_x^1 L_y^{r_2'})},
 \end{aligned}$$

and then Hölder's inequality in each t, s gives

$$|T_j(F, G)| \leq 2^{-nj(1 - \frac{1}{r_2}) + j} \|F\|_{L_t^2(I; L_x^1 L_y^{r_2'})} \|G\|_{L_t^2(J; L_x^1 L_y^{r_2'})}$$

as desired.

For the case (b), we bring the s -integration inside the inner product in (3.3) and apply Hölder's inequality in x, y to obtain

$$\begin{aligned}
 |T_j(F, G)| &\leq \int_J \left\| \int_I e^{-is(\Delta_x - \Delta_y)} F(s) ds \right\|_{L_{x,y}^2} \|e^{-it(\Delta_x - \Delta_y)} G(t)\|_{L_{x,y}^2} dt \\
 &\leq \left\| \int_I e^{-is(\Delta_x - \Delta_y)} F(s) ds \right\|_{L_{x,y}^2} \int_J \|e^{-it(\Delta_x - \Delta_y)} G(t)\|_{L_{x,y}^2} dt. \quad (4.2)
 \end{aligned}$$

By applying the Plancherel theorem and Hölder's inequality, the second term in the right-hand side of (4.2) becomes

$$\int_J \|e^{-it(\Delta_x - \Delta_y)} G(t)\|_{L_{x,y}^2} dt \leq 2^{\frac{j}{2}} \|G\|_{L_t^2(J; L_{x,y}^2)}. \quad (4.3)$$

On the other hand, we handle the first term using the dual version of the nonendpoint estimates as follows:

$$\begin{aligned} \left\| \int_I e^{-is(\Delta_x - \Delta_y)} F(s) \, ds \right\|_{L_{x,y}^2} &= \left\| \int_{\mathbb{R}} e^{-is(\Delta_x - \Delta_y)} \chi_I(s) F(s) \, ds \right\|_{L_{x,y}^2} \\ &\lesssim \|\chi_I F\|_{L_t^{q'_a} L_x^{a'} L_y^{r'_2}} \\ &= \|F\|_{L_t^{q'_a}(I; L_x^{a'} L_y^{r'_2})}, \end{aligned}$$

where $2 < q_a \leq \infty$, $2 \leq r_2 \leq a$ and $2/q_a = n(1 - 1/a - 1/r_2)$ which imply

$$\frac{1}{r_1} = 1 - \frac{1}{r_2} - \frac{1}{n} < \frac{1}{a} \leq \frac{1}{r_2}.$$

Then Hölder's inequality in t gives

$$\begin{aligned} \left\| \int_I e^{-is(\Delta_x - \Delta_y)} F(s) \, ds \right\|_{L_{x,y}^2} &\lesssim 2^{j(\frac{1}{2} - \frac{1}{q_a})} \|F\|_{L_t^2(I; L_x^{a'} L_y^{r'_2})} \\ &= 2^{j(\frac{1}{2} - \frac{n}{2}(1 - \frac{1}{a} - \frac{1}{r_2}))} \|F\|_{L_t^2(I; L_x^{a'} L_y^{r'_2})}. \end{aligned} \quad (4.4)$$

Combining (4.2), (4.3) and (4.4), we finally get

$$|T_j(F, G)| \lesssim 2^{j(1 - \frac{n}{2}(1 - \frac{1}{a} - \frac{1}{r_2}))} \|F\|_{L_t^2(I; L_x^{a'} L_y^{r'_2})} \|G\|_{L_t^2(J; L_{x,y}^2)}$$

as desired. A similar argument gives the case (c).

5. APPLICATIONS

In this final section we present some applications of Theorem 1.1 to the following nonlinear problem:

$$\begin{cases} i\partial_t u + (\Delta_x - \Delta_y)u = \pm |u|^\alpha u, \\ u(x, y, 0) = f(x, y), \end{cases} \quad (5.1)$$

where $(x, y, t) \in \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}$ and $\alpha > 0$. Particularly when $n = 1$, this equation with $\alpha = 2$ is reduced to the two-dimensional hyperbolic nonlinear Schrödinger equation which appears in nonlinear optics ([6, 21]) and arises naturally in the study of modulation of wave trains in gravity water waves ([22, 23]). Various issues concerning the well-posedness of this special case $n = 1$ have been intensively studied until lately (see, for example, [2, 5] and references therein). Motivated by this, we address here the higher dimensional cases. Our result is the following theorem which shows that the problem (5.1) is locally well-posed:

Theorem 5.1. *Let $n \geq 3$. Then for $f \in L_x^2 H_y^1$ there exist $T > 0$ and a unique solution to (5.1) for $0 < \alpha < 2/(n-1)$,*

$$u \in C_t([0, T]; L_x^2 H_y^1) \cap L_t^q([0, T]; L_x^{r_1} W_y^{1, r_2}),$$

with (q, r_1, r_2) satisfying all the conditions given in Theorem 1.1 together with

$$\frac{\alpha(n-1)-1}{2} \leq \frac{1}{q} \leq \frac{\alpha(n-1)}{2}, \quad \frac{1}{2} - \frac{\alpha}{2} \leq \frac{1}{r_2} - \frac{\alpha}{n} \leq \frac{1}{r_1} \leq 1 - \frac{\alpha}{2}. \quad (5.2)$$

Remark 5.2. From the proof, one can see that the same result can be valid for L^2 initial data $f \in L_x^2 L_y^2$ under the diagonal case $r_1 = r_2$ with $0 < \alpha < 2/n$.

Proof of Theorem 5.1. By Duhamel's principle, the solution of (5.1) can be written as

$$\Phi(u) = e^{it(\Delta_x - \Delta_y)} f - i \int_0^t e^{i(t-s)(\Delta_x - \Delta_y)} F(u) ds \quad (5.3)$$

where $F(u) = \pm |u(\cdot, \cdot, s)|^\alpha u(\cdot, \cdot, s)$. For suitable values of $T, A > 0$, we shall show that Φ defines a contraction map on

$$X(T, A) = \left\{ u \in C_t(I; L_x^2 H_y^1) \cap L_t^q(I; L_x^{r_1} W_y^{1, r_2}) : \sup_{t \in I} \|u\|_{L_x^2 H_y^1} + \|u\|_{L_t^q(I; L_x^{r_1} W_y^{1, r_2})} \leq A \right\}$$

equipped with the distance

$$d(u, v) = \sup_{t \in I} \|u - v\|_{L_{x,y}^2} + \|u - v\|_{L_t^q(I; L_x^{r_1} L_y^{r_2})},$$

where $I = [0, T]$ and (q, r_1, r_2) is given as in Theorem 5.1.

To control the Duhamel term in (5.3), we need the following inhomogeneous estimates which are derived from the homogeneous estimates (1.3) adopting TT^* argument and the Christ-Kiselev lemma [4]:

Corollary 5.3. *Let $n \geq 1$. Assume that (q, r_1, r_2) and $(\tilde{q}, \tilde{r}_1, \tilde{r}_2)$ are given as in Theorem 1.1. Then*

$$\left\| \int_0^t e^{it(\Delta_x - \Delta_y)} F(s) ds \right\|_{L_t^q L_x^{r_1} L_y^{r_2}} \lesssim \|F\|_{L_t^{\tilde{q}'} L_x^{\tilde{r}_1'} L_y^{\tilde{r}_2'}} \quad (5.4)$$

if $q > \tilde{q}'$.

Now we first show that $\Phi(u) \in X$ for $u \in X$. Using Plancherel's theorem and then the adjoint form of (1.3), we see that

$$\begin{aligned} \sup_{t \in I} \|\Phi(u)\|_{L_x^2 H_y^1} &\leq C \|f\|_{L_x^2 H_y^1} + C \sup_{t \in I} \left\| \int_0^t e^{-is(\Delta_x - \Delta_y)} F(u) ds \right\|_{L_x^2 H_y^1} \\ &\leq C \|f\|_{L_x^2 H_y^1} + C \|F(u)\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} W_y^{1, \tilde{r}_2'})} \end{aligned}$$

for $(\tilde{q}, \tilde{r}_1, \tilde{r}_2)$ given as in Corollary 5.3. For $q > \tilde{q}'$, we assume for the moment that

$$\|F(u)\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} W_y^{1, \tilde{r}_2'})} \leq CT^{\frac{1}{\tilde{q}'} - \frac{1}{q}} \|u\|_{L_t^\infty(I; L_x^2 H_y^1)}^\alpha \|u\|_{L_t^q(I; L_x^{r_1} W_y^{1, r_2})}, \quad (5.5)$$

where

$$1/\tilde{r}_1' = \alpha/2 + 1/r_1 \quad \text{and} \quad 1/\tilde{r}_2' = \alpha(1/2 - 1/n) + 1/r_2 \quad (5.6)$$

which imply

$$1/\tilde{q}' - 1/q = (2 + \alpha - n\alpha)/2 > 0 \quad (\text{thus, } \alpha < 2/(n-1)) \quad (5.7)$$

by combining the conditions (1.2) for (q, r_1, r_2) and $(\tilde{q}, \tilde{r}_1, \tilde{r}_2)$. Hence we get for $u \in X$

$$\sup_{t \in I} \|\Phi(u)\|_{L_x^2 H_y^1} \leq C \|f\|_{L_x^2 H_y^1} + CT^{\frac{2+\alpha-n\alpha}{2}} A^{\alpha+1}.$$

The same argument together with Theorem 1.1 also implies

$$\|\Phi(u)\|_{L_t^q(I; L_x^{r_1} W_y^{1, r_2})} \leq C\|f\|_{L_x^2 H_y^1} + CT^{\frac{2+\alpha-n\alpha}{2}} A^{\alpha+1}$$

for $u \in X$. Consequently, $\Phi(u) \in X$ if

$$C\|f\|_{L_x^2 H_y^1} + CT^{\frac{2+\alpha-n\alpha}{2}} A^{\alpha+1} \leq A. \quad (5.8)$$

Next we show that for $u, v \in X$

$$d(\Phi(u), \Phi(v)) \leq \frac{1}{2}d(u, v). \quad (5.9)$$

First we use (5.4) to get

$$\begin{aligned} d(\Phi(u), \Phi(v)) &= d\left(\int_0^t e^{i(t-s)(\Delta_x - \Delta_y)} F(u) ds, \int_0^t e^{i(t-s)(\Delta_x - \Delta_y)} F(v) ds\right) \\ &\leq C\|F(u) - F(v)\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} L_y^{\tilde{r}_2'})}. \end{aligned}$$

Using the simple inequality $\|u\|^\alpha u - \|v\|^\alpha v \leq C(|u|^\alpha + |v|^\alpha)|u - v|$, and then applying Hölder's inequality in x, y, t under the condition (5.6), we obtain

$$\begin{aligned} \|F(u) - F(v)\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} L_y^{\tilde{r}_2'})} &\leq CT^{\frac{1}{\tilde{q}'} - \frac{1}{q}} (\|u\|_{L_t^\infty(I; L_x^2 L_y^{\frac{2n}{n-2}})}^\alpha + \|v\|_{L_t^\infty(I; L_x^2 L_y^{\frac{2n}{n-2}})}^\alpha) \|u - v\|_{L_t^q(I; L_x^{r_1} L_y^{r_2})}. \end{aligned}$$

Then, (5.7) and the Sobolev embedding $\dot{H}^1 \hookrightarrow L^{\frac{2n}{n-2}}$ when $n > 2$ give

$$d(\Phi(u), \Phi(v)) \leq CT^{\frac{2+\alpha-n\alpha}{2}} A^\alpha d(u, v)$$

for $u \in X$.

Finally, we choose $A = 2C\|f\|_{L_x^2 H_y^1}$ and T so that $CT^{\frac{2+\alpha-n\alpha}{2}} A^\alpha \leq 1/2$, and thus (5.8) and (5.9) hold as desired. Therefore, there exists a unique local solution $u \in C_t(I; L_x^2 H_y^1) \cap L_t^q(I; L_x^{r_1} W_y^{1, r_2})$.

It remains to show (5.5). We first see that

$$\|F(u)\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} W_y^{1, \tilde{r}_2'})} \leq C\|u\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} L_y^{\tilde{r}_2'})}^\alpha + C\|u\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} L_y^{\tilde{r}_2'})}^\alpha \|\nabla_y u\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} L_y^{\tilde{r}_2'})}.$$

By Hölder's inequality, we then get

$$\|u\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} L_y^{\tilde{r}_2'})}^\alpha \leq C\|u\|_{L_t^\infty(I; L_x^2 L_y^{\frac{2n}{n-2}})}^\alpha \|u\|_{L_t^{\tilde{q}'}(I; L_x^{r_1} L_y^{r_2})}$$

and

$$\|u\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} L_y^{\tilde{r}_2'})}^\alpha \leq C\|u\|_{L_t^\infty(I; L_x^2 L_y^{\frac{2n}{n-2}})}^\alpha \|u\|_{L_t^{\tilde{q}'}(I; L_x^{r_1} \dot{W}_y^{1, r_2})}$$

under the condition (5.6). Thus, we obtain

$$\begin{aligned} \|F(u)\|_{L_t^{\tilde{q}'}(I; L_x^{\tilde{r}_1'} W_y^{1, \tilde{r}_2'})} &\leq C\|u\|_{L_t^\infty(I; L_x^2 H_y^1)}^\alpha \|u\|_{L_t^{\tilde{q}'}(I; L_x^{r_1} W_y^{1, r_2})} \\ &\leq CT^{\frac{1}{\tilde{q}'} - \frac{1}{q}} \|u\|_{L_t^\infty(I; L_x^2 H_y^1)}^\alpha \|u\|_{L_t^q(I; L_x^{r_1} W_y^{1, r_2})} \end{aligned}$$

using $\dot{H}^1 \hookrightarrow L^{\frac{2n}{n-2}}$ and Hölder's inequality in time with $q > \tilde{q}'$, as desired.

We finish the proof with a remark that the condition (5.2) follows from the requirements on $(\tilde{q}, \tilde{r}_1, \tilde{r}_2)$. Indeed, combining (5.6) with $2 \leq \tilde{r}_2 \leq \tilde{r}_1 \leq \infty$ implies the

second condition in (5.2), while the first one in (5.2) follows from a combination with $2 \leq \tilde{q} \leq \infty$ and (1.2) for (q, r_1, r_2) and $(\tilde{q}, \tilde{r}_1, \tilde{r}_2)$. \square

REFERENCES

- [1] J. Bergh and J. Löfström, *Interpolation Spaces, An Introduction*, Springer-Verlag, Berlin-New York, 1976.
- [2] R. Carles and C. Gallo, *WKB analysis of nonelliptic nonlinear Schrödinger equations*, Commun. Contemp. Math. 22 (2020), 1950045, 19 pp.
- [3] Z. Chen, *von Neumann-Landau equation for wave functions, wave-particle duality and collapses of wave functions*, arXiv:quant-ph/0703204v2, 2007.
- [4] M. Christ and A. Kiselev, *Maximal functions associated to filtrations*, J. Funct. Anal. 179 (2001), no. 2, 409-425.
- [5] B. Dodson, J. L. Marzuola, B. Pausader and D. P. Sporn, *The profile decomposition for the hyperbolic Schrödinger equation*, Preprint, arXiv:1708.08014.
- [6] E. Dumas, D. Lannes and J. Szeftel, *Variants of the focusing NLS equation: derivation, justification, and open problems related to filamentation*, in Laser filamentation, CRM Ser. Math. Phys., Springer, Cham, 2016, 19-75.
- [7] D. Foschi, *Inhomogeneous Strichartz estimates*, J. Hyperbolic Differ. Equ. 2 (2005), 1-24.
- [8] J. Ginibre and G. Velo, *The global Cauchy problem for the nonlinear Schrödinger equation revisited*, Ann. Inst. H. Poincaré Anal. Non Linéaire 2 (1985), 309-327.
- [9] L. Grafakos, *Classical Fourier Analysis*, Springer, New York, 2014.
- [10] L. Hörmander, *Estimates for translation invariant operators in L^p spaces*, Acta Math. 104 (1960), 93-140.
- [11] M. Keel and T. Tao, *Endpoint Strichartz estimates*, Amer. J. Math. 120 (1998), 955-980.
- [12] S. Klainerman and M. Machedon, *On the uniqueness of solutions to the Gross-Pitaevskii hierarchy*, Commun. Math. Phys. 279 (2008), 169-185.
- [13] L. D. Landau, *Das Dämpfungsproblem in der Wellenmechanik*, Z. Physik, 45 (1927), 430-441.
- [14] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics: Non-relativistic Theory*, Pergamon Press, Oxford, 1987.
- [15] C. Liu and M. Liu, *On the Cauchy Problem for Von Neumann-Landau Wave Equation*, Journal of Applied Mathematics and Physics, 2 (2014), 1224-1332.
- [16] S. J. Montgomery-Smith, *Time decay for the bounded mean oscillation of solutions of the Schrödinger and wave equations*, Duke Math. J. 91 (1998), 393-408.
- [17] J. von Neumann, *Wahrscheinlichkeitstheoretischer Aufbau der Quantenmechanik*, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse 11 (1927), 245-272.
- [18] J. von Neumann, *Mathematical Foundations of Quantum Mechanics*, Princeton University Press, Princeton, 1955.
- [19] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton University Press, Princeton, New Jersey, 1970.
- [20] R. S. Strichartz, *Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations*, Duke Math. J. 44 (1977), 705-714.
- [21] C. Sulem and P.-L. Sulem, *The nonlinear Schrödinger equation. Self-focusing and wave collapse*, Applied Mathematical Sciences, 139. Springer-Verlag, New York, 1999.
- [22] N. Totz, *A justification of the modulation approximation to the 3D full water wave problem*, Comm. Math. Phys. 335 (2015), 369-443.
- [23] N. Totz and S. Wu, *A rigorous justification of the modulation approximation to the 2D full water wave problem*, Comm. Math. Phys. 310 (2012), 817-883.

DEPARTMENT OF MATHEMATICS EDUCATION, KONGJU NATIONAL UNIVERSITY, KONGJU 32588,
REPUBLIC OF KOREA

Email address: `ywkoh@kongju.ac.kr`

DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON 16419, REPUBLIC OF KO-
REA

Email address: `yjglee@skku.edu`

DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON 16419, REPUBLIC OF KO-
REA

Email address: `ihseo@skku.edu`