

A RIGHT INVERSE OF DIFFERENTIAL OPERATOR $\Delta + a$ IN WEIGHTED HILBERT SPACE $L^2(\mathbb{R}^n, e^{-|x|^2})$

SHAORYU DAI¹, YANG LIU² AND YIFEI PAN³

ABSTRACT. In this note, we prove the existence of weak solutions of a Poisson type equation in the weighted Hilbert space $L^2(\mathbb{R}^n, e^{-|x|^2})$.

1. INTRODUCTION

In this note, we study the right inverse of differential operator $\Delta + a$ in a Hilbert space by proving the following result on the existence of (global) weak solutions of a Poisson type equation in the weighted Hilbert space $L^2(\mathbb{R}^n, e^{-|x|^2})$, where a is a real constant throughout.

Theorem 1.1. *For each $f \in L^2(\mathbb{R}^n, e^{-|x|^2})$, there exists a weak solution $u \in L^2(\mathbb{R}^n, e^{-|x|^2})$ solving the equation*

$$\Delta u + au = f$$

in $\mathbb{R}^n$ with the norm estimate

$$\int_{\mathbb{R}^n} u^2 e^{-|x|^2} dx \leq \frac{1}{8n} \int_{\mathbb{R}^n} f^2 e^{-|x|^2} dx.$$

The novelty of Theorem 1.1 is that the differential operator $\Delta + a$ has a bounded right inverse

$$\begin{aligned} Q : L^2(\mathbb{R}^n, e^{-|x|^2}) &\longrightarrow L^2(\mathbb{R}^n, e^{-|x|^2}), \\ (\Delta + a)Q &= I \end{aligned}$$

with the norm estimate $\|Q\| \leq \frac{1}{\sqrt{8n}}$. In particular, the Laplace operator Δ has a bounded right inverse $Q_0 : L^2(\mathbb{R}^n, e^{-|x|^2}) \longrightarrow L^2(\mathbb{R}^n, e^{-|x|^2})$, which, to the best of our knowledge, appears to be new.

As a simple consequence of Theorem 1.1, we can obtain the following result on the existence of global weak solutions of a Poisson type equation for square integrable functions. This result seems to be classical and well-known, but we could not find an exact reference. At least it is well-known [1] that when f is compact supported in $\mathbb{R}^n$, then a solution of $-\Delta u = f$ is $\Phi * f$, where Φ is the fundamental solution of Laplace's equation.

Theorem 1.2. *For each $f \in L^2(\mathbb{R}^n)$ or $f \in L^\infty(\mathbb{R}^n)$, there exists a weak solution $u \in L^2_{loc}(\mathbb{R}^n)$ solving the equation*

$$\Delta u + au = f.$$

In particular, the Poisson equation $\Delta u = f$ has a weak solution $u \in L^2_{loc}(\mathbb{R}^n)$ for $f \in L^2(\mathbb{R}^n)$ or $f \in L^\infty(\mathbb{R}^n)$.

The proof of Theorem 1.2 follows from the observation that $L^2(\mathbb{R}^n) \subset L^2(\mathbb{R}^n, e^{-|x|^2})$, $L^\infty(\mathbb{R}^n) \subset L^2(\mathbb{R}^n, e^{-|x|^2})$ and $L^2(\mathbb{R}^n, e^{-|x|^2}) \subset L^2_{loc}(\mathbb{R}^n)$.

The method employed in this note was motivated from the Hörmander L^2 method [2] for Cauchy-Riemann equations from several complex variables.

2. SEVERAL LEMMAS

Here, we consider weighted Hilbert space

$$L^2(\mathbb{R}^n, e^{-\varphi}) = \{f \mid f \in L^2_{loc}(\mathbb{R}^n); \int_{\mathbb{R}^n} f^2 e^{-\varphi} dx < +\infty\},$$

where φ is a nonnegative function on $\mathbb{R}^n$. We denote the weighted inner product for $f, g \in L^2(\mathbb{R}^n, e^{-\varphi})$ by $\langle f, g \rangle_\varphi = \int_{\mathbb{R}^n} f g e^{-\varphi} dx$ and the weighted norm of $f \in L^2(\mathbb{R}^n, e^{-\varphi})$ by $\|f\|_\varphi = \sqrt{\langle f, f \rangle_\varphi}$. Let k be a positive integer, $C^k(\mathbb{R}^n)$ denote the set of all k -times continuously differentiable real-valued functions on $\mathbb{R}^n$, and $C_0^\infty(\mathbb{R}^n)$ denote the set of all smooth functions $\phi : \mathbb{R}^n \rightarrow \mathbb{R}$ with compact support. Let $\Delta = \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2}$ be the Laplace operator on $\mathbb{R}^n$. For $u, f \in L^2_{loc}(\mathbb{R}^n)$, we say that f is the Laplace of u in the weak sense, written $\Delta u = f$, provided $\int_{\mathbb{R}^n} u \Delta \phi dx = \int_{\mathbb{R}^n} f \phi dx$ for all test functions $\phi \in C_0^\infty(\mathbb{R}^n)$.

Let $\varphi \in C^2(\mathbb{R}^n)$. For $\forall \phi \in C_0^\infty(\mathbb{R}^n)$, we first define the following formal adjoint of Δ with respect to the weighted inner product in $L^2(\mathbb{R}^n, e^{-\varphi})$. Let $u \in L^2_{loc}(\mathbb{R}^n)$. We calculate as follows.

$$\begin{aligned} \langle \phi, \Delta u \rangle_\varphi &= \int_{\mathbb{R}^n} \phi \Delta u e^{-\varphi} dx \\ &= \int_{\mathbb{R}^n} u \Delta (\phi e^{-\varphi}) dx \\ &= \int_{\mathbb{R}^n} e^\varphi u \Delta (\phi e^{-\varphi}) e^{-\varphi} dx \\ &= \langle e^\varphi \Delta (\phi e^{-\varphi}), u \rangle_\varphi \\ &=: \langle \Delta_\varphi^* \phi, u \rangle_\varphi, \end{aligned}$$

where $\Delta_\varphi^* \phi = e^\varphi \Delta (\phi e^{-\varphi})$ is the formal adjoint of Δ with domain in $C_0^\infty(\mathbb{R}^n)$. Let $(\Delta + a)_\varphi^*$ be the formal adjoint of $\Delta + a$ with domain in $C_0^\infty(\mathbb{R}^n)$. Note that $I_\varphi^* = I$, where I is the identity operator. Then $(\Delta + a)_\varphi^* = \Delta_\varphi^* + a$.

Let ∇ be the gradient operator on $\mathbb{R}^n$. Now we give several key lemmas based on functional analysis.

Lemma 2.1. *Let $\varphi \in C^2(\mathbb{R}^n)$. For each $f \in L^2(\mathbb{R}^n, e^{-\varphi})$, there exists a global weak solution $u \in L^2(\mathbb{R}^n, e^{-\varphi})$ solving the equation*

$$\Delta u + au = f$$

in $\mathbb{R}^n$ with the norm estimate

$$\|u\|_\varphi^2 \leq c$$

if and only if

$$|\langle f, \phi \rangle_\varphi|^2 \leq c \left\| (\Delta + a)_\varphi^* \phi \right\|_\varphi^2, \quad \forall \phi \in C_0^\infty(\mathbb{R}^n),$$

where c is a constant.

Proof. Let $\Delta + a = H$. Then $(\Delta + a)_\varphi^* = H_\varphi^*$.

(Necessity) For $\forall \phi \in C_0^\infty(\mathbb{R}^n)$, from the definition of H_φ^* and Cauchy-Schwarz inequality, we have

$$|\langle f, \phi \rangle_\varphi|^2 = |\langle Hu, \phi \rangle_\varphi|^2 = \left| \langle u, H_\varphi^* \phi \rangle_\varphi \right|^2 \leq \|u\|_\varphi^2 \|H_\varphi^* \phi\|_\varphi^2 \leq c \|H_\varphi^* \phi\|_\varphi^2 = c \left\| (\Delta + a)_\varphi^* \phi \right\|_\varphi^2.$$

(Sufficiency) Consider the subspace

$$E = \{H_\varphi^* \phi \mid \phi \in C_0^\infty(\mathbb{R}^n)\} \subset L^2(\mathbb{R}^n, e^{-\varphi}).$$

Define a linear functional $L_f : E \rightarrow \mathbb{R}$ by

$$L_f(H_\varphi^* \phi) = \langle f, \phi \rangle_\varphi = \int_{\mathbb{R}^n} f \phi e^{-\varphi} dx.$$

Since

$$|L_f(H_\varphi^* \phi)| = |\langle f, \phi \rangle_\varphi| \leq \sqrt{c} \|H_\varphi^* \phi\|_\varphi,$$

then L_f is a bounded functional on E . Let $\overline{E}$ be the closure of E with respect to the norm $\|\cdot\|_\varphi$ of $L^2(\mathbb{R}^n, e^{-\varphi})$. Note that $\overline{E}$ is a Hilbert subspace of $L^2(\mathbb{R}^n, e^{-\varphi})$. So by Hahn-Banach's extension theorem, L_f can be extended to a linear functional $\tilde{L}_f$ on $\overline{E}$ such that

$$|\tilde{L}_f(g)| \leq \sqrt{c} \|g\|_\varphi, \quad \forall g \in \overline{E}. \quad (1)$$

Using the Riesz representation theorem for $\tilde{L}_f$, there exists a unique $u_0 \in \overline{E}$ such that

$$\tilde{L}_f(g) = \langle u_0, g \rangle_\varphi, \quad \forall g \in \overline{E}. \quad (2)$$

Now we prove $\Delta u_0 + a u_0 = f$. For $\forall \phi \in C_0^\infty(\mathbb{R}^n)$, apply $g = H_\varphi^* \phi$ in (2). Then

$$\tilde{L}_f(H_\varphi^* \phi) = \langle u_0, H_\varphi^* \phi \rangle_\varphi = \langle H u_0, \phi \rangle_\varphi.$$

Note that

$$\tilde{L}_f(H_\varphi^* \phi) = L_f(H_\varphi^* \phi) = \langle f, \phi \rangle_\varphi.$$

Therefore,

$$\langle H u_0, \phi \rangle_\varphi = \langle f, \phi \rangle_\varphi, \quad \forall \phi \in C_0^\infty(\mathbb{R}^n),$$

i.e.,

$$\int_{\mathbb{R}^n} H u_0 \phi e^{-\varphi} dx = \int_{\mathbb{R}^n} f \phi e^{-\varphi} dx, \quad \forall \phi \in C_0^\infty(\mathbb{R}^n),$$

i.e.,

$$\int_{\mathbb{R}^n} (Hu_0 - f)\phi e^{-\varphi} dx = 0, \quad \forall \phi \in C_0^\infty(\mathbb{R}^n).$$

Thus, $Hu_0 = f$, i.e., $\Delta u_0 + au_0 = f$.

Next we give a bound for the norm of u_0 . Let $g = u_0$ in (1) and (2). Then we have

$$\|u_0\|_\varphi^2 = |\langle u_0, u_0 \rangle_\varphi| = \left| \tilde{L}_f(u_0) \right| \leq \sqrt{c} \|u_0\|_\varphi.$$

Therefore, $\|u_0\|_\varphi^2 \leq c$.

Note that $u_0 \in \overline{E}$ and $\overline{E} \subset L^2(\mathbb{R}^n, e^{-\varphi})$. Then $u_0 \in L^2(\mathbb{R}^n, e^{-\varphi})$. Let $u = u_0$. So there exists $u \in L^2(\mathbb{R}^n, e^{-\varphi})$ such that $\Delta u + au = f$ with $\|u\|_\varphi^2 \leq c$. The proof is complete. $\square$

Lemma 2.2. *Let $\varphi \in C^4(\mathbb{R}^n)$. Then*

$$\left\| (\Delta + a)_\varphi^* \phi \right\|_\varphi^2 = \|(\Delta + a)\phi\|_\varphi^2 + \langle \phi, \Delta(\Delta_\varphi^* \phi) - \Delta_\varphi^*(\Delta\phi) \rangle_\varphi, \quad \forall \phi \in C_0^\infty(\mathbb{R}^n).$$

Proof. Let $\Delta + a = H$. Then $(\Delta + a)_\varphi^* = H_\varphi^*$. For $\forall \phi \in C_0^\infty(\mathbb{R}^n)$, we have

$$\begin{aligned} \|H_\varphi^* \phi\|_\varphi^2 &= \langle H_\varphi^* \phi, H_\varphi^* \phi \rangle_\varphi \\ &= \langle \phi, HH_\varphi^* \phi \rangle_\varphi \\ &= \langle \phi, H_\varphi^* H \phi \rangle_\varphi + \langle \phi, HH_\varphi^* \phi - H_\varphi^* H \phi \rangle_\varphi \\ &= \langle H\phi, H\phi \rangle_\varphi + \langle \phi, HH_\varphi^* \phi - H_\varphi^* H \phi \rangle_\varphi \\ &= \|H\phi\|_\varphi^2 + \langle \phi, HH_\varphi^* \phi - H_\varphi^* H \phi \rangle_\varphi \end{aligned} \tag{3}$$

Note that

$$\begin{aligned} HH_\varphi^* \phi &= (\Delta + a)(\Delta + a)_\varphi^* \phi \\ &= (\Delta + a)(\Delta_\varphi^* \phi + a\phi) \\ &= \Delta(\Delta_\varphi^* \phi) + a\Delta\phi + a\Delta_\varphi^* \phi + a^2\phi \end{aligned}$$

and

$$\begin{aligned} H_\varphi^* H \phi &= (\Delta + a)_\varphi^* (\Delta + a)\phi \\ &= (\Delta_\varphi^* + a)(\Delta\phi + a\phi) \\ &= \Delta_\varphi^*(\Delta\phi) + a\Delta_\varphi^* \phi + a\Delta\phi + a^2\phi. \end{aligned}$$

Then

$$HH_\varphi^* \phi - H_\varphi^* H \phi = \Delta(\Delta_\varphi^* \phi) - \Delta_\varphi^*(\Delta\phi). \tag{4}$$

So by (3) and (4), we have

$$\|H_\varphi^* \phi\|_\varphi^2 = \|H\phi\|_\varphi^2 + \langle \phi, \Delta(\Delta_\varphi^* \phi) - \Delta_\varphi^*(\Delta\phi) \rangle_\varphi.$$

This lemma is proved. $\square$

Lemma 2.3. Let $\varphi = |x|^2$. Then for $\forall \phi \in C_0^\infty(\mathbb{R}^n)$,

$$\langle \phi, \Delta(\Delta_\varphi^* \phi) - \Delta_\varphi^*(\Delta \phi) \rangle_\varphi = 8n \|\phi\|_\varphi^2 + 8 \|\nabla \phi\|_\varphi^2.$$

Proof. For $\forall \phi \in C_0^\infty(\mathbb{R}^n)$, by the definition of Δ_φ^* and the following formula

$$\Delta(\alpha\beta) = \beta\Delta\alpha + \alpha\Delta\beta + 2\nabla\alpha \cdot \nabla\beta, \quad \forall \alpha, \beta \in C^2(\mathbb{R}^n),$$

we have

$$\Delta_\varphi^* \phi = e^\varphi \Delta(\phi e^{-\varphi}) = \Delta\phi + \phi|\nabla\varphi|^2 - \phi\Delta\varphi - 2\nabla\phi \cdot \nabla\varphi. \quad (5)$$

From (5) we have

$$\begin{aligned} \Delta(\Delta_\varphi^* \phi) &= \Delta^2\phi + \Delta(\phi|\nabla\varphi|^2) - \Delta(\phi\Delta\varphi) - 2\Delta(\nabla\phi \cdot \nabla\varphi) \\ &= \Delta^2\phi + \Delta\phi|\nabla\varphi|^2 + \phi\Delta(|\nabla\varphi|^2) + 2\nabla\phi \cdot \nabla(|\nabla\varphi|^2) \\ &\quad - \Delta\phi\Delta\varphi - \phi\Delta^2\varphi - 2\nabla\phi \cdot \nabla(\Delta\varphi) - 2\Delta(\nabla\phi \cdot \nabla\varphi) \end{aligned}$$

and

$$\Delta_\varphi^*(\Delta\phi) = \Delta^2\phi + \Delta\phi|\nabla\varphi|^2 - \Delta\phi\Delta\varphi - 2\nabla(\Delta\phi) \cdot \nabla\varphi.$$

Then

$$\begin{aligned} \Delta(\Delta_\varphi^* \phi) - \Delta_\varphi^*(\Delta\phi) &= \phi\Delta(|\nabla\varphi|^2) + 2\nabla\phi \cdot \nabla(|\nabla\varphi|^2) - \phi\Delta^2\varphi \\ &\quad - 2\nabla\phi \cdot \nabla(\Delta\varphi) - 2\Delta(\nabla\phi \cdot \nabla\varphi) + 2\nabla(\Delta\phi) \cdot \nabla\varphi. \end{aligned} \quad (6)$$

Let $\varphi = |x|^2$. We have $\nabla\varphi = 2x$, $\Delta\varphi = 2n$, $|\nabla\varphi|^2 = 4|x|^2$, $\nabla(|\nabla\varphi|^2) = 8x$, $\Delta(|\nabla\varphi|^2) = 8n$. Then by (6) and the following formula

$$\Delta(\nabla\phi \cdot x) = \nabla(\Delta\phi) \cdot x + 2\Delta\phi, \quad \forall \phi \in C_0^\infty(\mathbb{R}^n),$$

we get

$$\Delta(\Delta_\varphi^* \phi) - \Delta_\varphi^*(\Delta\phi) = 8n\phi + 16(\nabla\phi \cdot x) - 8\Delta\phi.$$

Consequently,

$$\begin{aligned} \langle \phi, \Delta(\Delta_\varphi^* \phi) - \Delta_\varphi^*(\Delta\phi) \rangle_\varphi &= \langle \phi, 8n\phi + 16(\nabla\phi \cdot x) - 8\Delta\phi \rangle_\varphi \\ &= 8n \|\phi\|_\varphi^2 + 8 \langle \phi, 2(\nabla\phi \cdot x) - \Delta\phi \rangle_\varphi. \end{aligned}$$

Note, as the key step of the proof, that

$$\begin{aligned}
\langle \phi, 2(\nabla \phi \cdot x) - \Delta \phi \rangle_\varphi &= \int_{\mathbb{R}^n} \phi (2(\nabla \phi \cdot x) - \Delta \phi) e^{-\varphi} dx \\
&= \int_{\mathbb{R}^n} \phi \sum_{j=1}^n \left(2x_j \frac{\partial \phi}{\partial x_j} - \frac{\partial^2 \phi}{\partial x_j \partial x_j} \right) e^{-|x|^2} dx \\
&= - \int_{\mathbb{R}^n} \phi \sum_{j=1}^n \frac{\partial}{\partial x_j} \left(\frac{\partial \phi}{\partial x_j} e^{-|x|^2} \right) dx \\
&= - \sum_{j=1}^n \int_{\mathbb{R}^n} \phi \frac{\partial}{\partial x_j} \left(\frac{\partial \phi}{\partial x_j} e^{-|x|^2} \right) dx \\
&= \sum_{j=1}^n \int_{\mathbb{R}^n} \frac{\partial \phi}{\partial x_j} \left(\frac{\partial \phi}{\partial x_j} e^{-|x|^2} \right) dx \\
&= \sum_{j=1}^n \int_{\mathbb{R}^n} \left(\frac{\partial \phi}{\partial x_j} \right)^2 e^{-|x|^2} dx \\
&= \int_{\mathbb{R}^n} |\nabla \phi|^2 e^{-|x|^2} dx \\
&= \|\nabla \phi\|_\varphi^2.
\end{aligned}$$

Then

$$\langle \phi, \Delta (\Delta_\varphi^* \phi) - \Delta_\varphi^* (\Delta \phi) \rangle_\varphi = 8n \|\phi\|_\varphi^2 + 8 \|\nabla \phi\|_\varphi^2.$$

The lemma is proved. □

3. PROOF OF THEOREMS

The proof of Theorems 1.1.

Proof. Let $\varphi = |x|^2$. By Lemma 2.2 and Lemma 2.3, we have for $\forall \phi \in C_0^\infty(\mathbb{R}^n)$,

$$\begin{aligned}
\left\| (\Delta + a)_\varphi^* \phi \right\|_\varphi^2 &= \|(\Delta + a) \phi\|_\varphi^2 + \langle \phi, \Delta (\Delta_\varphi^* \phi) - \Delta_\varphi^* (\Delta \phi) \rangle_\varphi \\
&\geq \langle \phi, \Delta (\Delta_\varphi^* \phi) - \Delta_\varphi^* (\Delta \phi) \rangle_\varphi \\
&\geq 8n \|\phi\|_\varphi^2.
\end{aligned} \tag{7}$$

By Cauchy-Schwarz inequality and (7), we have for $\forall \phi \in C_0^\infty(\mathbb{R}^n)$,

$$\begin{aligned} |\langle f, \phi \rangle_\varphi|^2 &\leq \|f\|_\varphi^2 \|\phi\|_\varphi^2 \\ &= \left(\frac{1}{8n} \|f\|_\varphi^2 \right) (8n \|\phi\|_\varphi^2) \\ &\leq \left(\frac{1}{8n} \|f\|_\varphi^2 \right) \|(\Delta + a)_\varphi^* \phi\|_\varphi^2. \end{aligned}$$

Let $c = \frac{1}{8n} \|f\|_\varphi^2$. Then

$$|\langle f, \phi \rangle_\varphi|^2 \leq c \|(\Delta + a)_\varphi^* \phi\|_\varphi^2, \quad \forall \phi \in C_0^\infty(\mathbb{R}^n).$$

By Lemma 2.1, there exists a global weak solution $u \in L^2(\mathbb{R}^n, e^{-\varphi})$ solving the equation

$$\Delta u + au = f$$

in $\mathbb{R}^n$ with the norm estimate

$$\|u\|_\varphi^2 \leq c,$$

i.e.,

$$\Delta u + au = f \quad \text{with} \quad \int_{\mathbb{R}^n} u^2 e^{-|x|^2} dx \leq \frac{1}{8n} \int_{\mathbb{R}^n} f^2 e^{-|x|^2} dx.$$

The proof is complete. $\square$

Theorem 3.1. *There exists a bounded operator $Q : L^2(\mathbb{R}^n, e^{-|x|^2}) \rightarrow L^2(\mathbb{R}^n, e^{-|x|^2})$ such that*

$$(\Delta + a)Q = I \quad \text{with} \quad \|Q\| \leq \frac{1}{\sqrt{8n}},$$

where $\|Q\|$ is the norm of Q in $L^2(\mathbb{R}^n, e^{-|x|^2})$.

Proof. Let $\varphi = |x|^2$. For each $f \in L^2(\mathbb{R}^n, e^{-\varphi})$, from Theorem 1.1, there exists $u \in L^2(\mathbb{R}^n, e^{-\varphi})$ such that

$$(\Delta + a)u = f \quad \text{with} \quad \|u\|_\varphi \leq \frac{1}{\sqrt{8n}} \|f\|_\varphi.$$

Denote this u by $Q(f)$. Then $Q(f)$ satisfies

$$(\Delta + a)Q(f) = f \quad \text{with} \quad \|Q(f)\|_\varphi \leq \frac{1}{\sqrt{8n}} \|f\|_\varphi.$$

Note that f is arbitrary in $L^2(\mathbb{R}^n, e^{-\varphi})$. So $Q : L^2(\mathbb{R}^n, e^{-\varphi}) \rightarrow L^2(\mathbb{R}^n, e^{-\varphi})$ is a bounded operator such that

$$(\Delta + a)Q = I \quad \text{with} \quad \|Q\| \leq \frac{1}{\sqrt{8n}}.$$

The proof is complete. $\square$

4. FURTHER REMARKS

Remark 1. Given $\lambda > 0$ and $x_0 \in \mathbb{R}^n$, for the weight $\varphi = \lambda|x - x_0|^2$, we obtain the following corollary from Theorem 1.1.

Corollary 4.1. *For each $f \in L^2(\mathbb{R}^n, e^{-\lambda|x-x_0|^2})$, there exists a weak solution $u \in L^2(\mathbb{R}^n, e^{-\lambda|x-x_0|^2})$ solving the equation*

$$\Delta u + au = f$$

with the norm estimate

$$\int_{\mathbb{R}^n} u^2 e^{-\lambda|x-x_0|^2} dx \leq \frac{1}{8n\lambda^2} \int_{\mathbb{R}^n} f^2 e^{-\lambda|x-x_0|^2} dx.$$

Proof. From $f \in L^2(\mathbb{R}^n, e^{-\lambda|x-x_0|^2})$, we have

$$\int_{\mathbb{R}^n} f^2(x) e^{-\lambda|x-x_0|^2} dx < +\infty. \quad (8)$$

Let $x = \frac{y}{\sqrt{\lambda}} + x_0$ and $g(y) = f(x) = f\left(\frac{y}{\sqrt{\lambda}} + x_0\right)$. Then by (8), we have

$$\frac{1}{(\sqrt{\lambda})^n} \int_{\mathbb{R}^n} g^2(y) e^{-|y|^2} dy < +\infty,$$

which implies that $g \in L^2(\mathbb{R}^n, e^{-|y|^2})$. For g , applying Theorem 1.1 with a replaced by $\frac{a}{\lambda}$, there exists a weak solution $v \in L^2(\mathbb{R}^n, e^{-|y|^2})$ solving the equation

$$\Delta v(y) + \frac{a}{\lambda} v(y) = g(y) \quad (9)$$

in $\mathbb{R}^n$ with the norm estimate

$$\int_{\mathbb{R}^n} v^2(y) e^{-|y|^2} dy \leq \frac{1}{8n} \int_{\mathbb{R}^n} g^2(y) e^{-|y|^2} dy. \quad (10)$$

Note that $y = \sqrt{\lambda}(x - x_0)$ and $g(y) = f(x)$. Let $u(x) = \frac{1}{\lambda} v(y) = \frac{1}{\lambda} v(\sqrt{\lambda}(x - x_0))$. Then (9) and (10) can be rewritten by

$$\Delta u(x) + au(x) = f(x) \quad (11)$$

$$\int_{\mathbb{R}^n} u^2(x) e^{-\lambda|x-x_0|^2} dx \leq \frac{1}{8n\lambda^2} \int_{\mathbb{R}^n} f^2(x) e^{-\lambda|x-x_0|^2} dx. \quad (12)$$

(12) implies that $u \in L^2(\mathbb{R}^n, e^{-\lambda|x-x_0|^2})$. Then by (11) and (12), the proof is complete. $\square$

Remark 2. From Corollary 4.1, we can obtain the following corollary, which shows that for any choice of a , the differential operator $\Delta + a$ has a bounded right inverse in $L^2(U)$, provided U is a bounded open set. This result should be well-known, and we could not locate a reference (for a very close related result, see Theorem 6 in page 324 in [1]).

Corollary 4.2. *Let $U \subset \mathbb{R}^n$ be any bounded open set. For each $f \in L^2(U)$, there exists a weak solution $u \in L^2(U)$ solving the equation*

$$\Delta u + au = f$$

with the norm estimate $\|u\|_{L^2(U)} \leq c\|f\|_{L^2(U)}$, where the constant c depends only on the diameter of U .

Proof. Let $x_0 \in U$. For given $f \in L^2(U)$, extending f to zero on $\mathbb{R}^n \setminus U$, we have

$$\tilde{f} = \begin{cases} f, & x \in U \\ 0, & x \in \mathbb{R}^n \setminus U. \end{cases}$$

Then $\tilde{f} \in L^2(\mathbb{R}^n) \subset L^2(\mathbb{R}^n, e^{-|x-x_0|^2})$. From Corollary 4.1, there exists $\tilde{u} \in L^2(\mathbb{R}^n, e^{-|x-x_0|^2})$ such that

$$\Delta \tilde{u} + a\tilde{u} = \tilde{f} \quad \text{with} \quad \int_{\mathbb{R}^n} \tilde{u}^2 e^{-|x-x_0|^2} dx \leq \frac{1}{8n} \int_{\mathbb{R}^n} \tilde{f}^2 e^{-|x-x_0|^2} dx.$$

Then

$$\int_{\mathbb{R}^n} \tilde{u}^2 e^{-|x-x_0|^2} dx \leq \frac{1}{8n} \int_{\mathbb{R}^n} \tilde{f}^2 dx = \frac{1}{8n} \int_U f^2 dx.$$

Note that

$$\int_{\mathbb{R}^n} \tilde{u}^2 e^{-|x-x_0|^2} dx \geq \int_U \tilde{u}^2 e^{-|x-x_0|^2} dx \geq \int_U \tilde{u}^2 e^{-|U|^2} dx = e^{-|U|^2} \int_U \tilde{u}^2 dx,$$

where $|U|$ is the diameter of U . Therefore,

$$e^{-|U|^2} \int_U \tilde{u}^2 dx \leq \frac{1}{8n} \int_U f^2 dx, \quad \text{i.e.,} \quad \int_U \tilde{u}^2 dx \leq \frac{e^{|U|^2}}{8n} \int_U f^2 dx.$$

Restricting $\tilde{u}$ on U to get u , then

$$\Delta u + au = f \quad \text{with} \quad \int_U u^2 dx \leq \frac{e^{|U|^2}}{8n} \int_U f^2 dx.$$

Note that $u \in L^2(U)$ and let $c = \sqrt{\frac{e^{|U|^2}}{8n}}$. Then the proof is complete. $\square$

Remark 3. When $f \in L^2(\mathbb{R}^n)$, the solutions of $\Delta u = f$ are not necessary in $L^2(\mathbb{R}^n)$. For example: $n = 1$,

$$f(x) = \begin{cases} \frac{1}{x}, & x \geq 1, \\ x, & 0 < x < 1, \\ 0, & x \leq 0, \end{cases}$$

$$u(x) = \int_0^x (x-t)f(t)dt + c_1x + c_2 = -\frac{x}{2} + x \ln x + \frac{2}{3} + c_1x + c_2, \quad x \geq 1,$$

where c_1 and c_2 are arbitrary real constants. It is easy to see $u \notin L^2(\mathbb{R})$.

REFERENCES

- [1] L. C. Evans, Partial differential equations, Second edition (2010).
- [2] L. Hörmander, L^2 estimates and existence theorems for the $\bar{\partial}$ operator, Acta Mathematica, 113 (1965), 89-152.

1 SCHOOL OF MATHEMATICS, SOUTHEAST UNIVERSITY, NANJING, 210096, CHINA;

DEPARTMENT OF MATHEMATICS, JINLING INSTITUTE OF TECHNOLOGY, NANJING, 211169, CHINA.

E-mail address: dymdsy@163.com

2 DEPARTMENT OF MATHEMATICS, ZHEJIANG NORMAL UNIVERSITY, JINHUA, 321004, CHINA.

E-mail address: liuyang@zjnu.edu.cn

3 DEPARTMENT OF MATHEMATICAL SCIENCES, PURDUE UNIVERSITY FORT WAYNE, FORT WAYNE, 46805-1499, USA.

E-mail address: pan@pfw.edu