

# The current algebra representations of quantum many-particle Schrödinger type Hamiltonian models, their factorized structure and integrability\*

D. Prorok<sup>1</sup>, A.K. Prykarpatski<sup>2</sup>

<sup>1</sup> Department of Physics and Applied Computer Science, AGH University of Science and Technology, Krakow, Poland

<sup>2</sup> Institute of Mathematics, Department of Physics, Mathematics and Computer Science, Cracow University of Technology, Krakow 31-155, Poland

Received July 16, 2019, in final form August 6, 2019

There is developed a current algebra representation scheme for reconstructing algebraically factorized quantum Hamiltonian and symmetry operators in the Fock type space and its application to quantum Hamiltonian and symmetry operators in case of quantum integrable spatially many- and one-dimensional dynamical systems. As examples, we have studied in detail the factorized structure of Hamiltonian operators, describing such quantum integrable spatially many- and one-dimensional models as generalized oscillatory, Calogero-Sutherland, Coulomb type and nonlinear Schrödinger dynamical systems of spinless bose-particles.

**Key words:** Fock space, current algebra representations, Hamiltonian reconstruction, Bogolubov generating functional, Calogero-Moser-Sutherland model, quantum integrability, quantum symmetries

**PACS:** 11.10.Ef, 11.15.Kc, 11.10.-z, 11.15.-q, 11.10.Wx, 05.30.-d

## 1. Introduction

Given any classical Hamiltonian many-particle non-relativistic system with the standard cotangent phase space  $T^*(\mathbb{R}^3)^{\otimes N}$ , where the quantity of particles  $N \in \mathbb{Z}_+$  is fixed, there is a standard recipe for producing a quantum system by a method known as “canonical quantization”, assigning to the system a suitably constructed [1] self-adjoint Hamiltonian operator, acting in the related Hilbert space  $\mathcal{H} = L_2(\mathbb{R}^{3 \otimes N}; \mathbb{C})$ . In case when all the particles are equivalent to each other and the particle number  $N \in \mathbb{Z}_+$  can vary within the system, there is another recipe applied to this system, called the “second quantization”, producing the corresponding [2, 3] quantum self-adjoint Hamiltonian operator, acting already in a specially constructed Fock space  $\Phi_F$ , whose basis vectors are generated by means of actions of additional so-called “creation” and “annihilation” operators on a uniquely defined “vacuum” zero-particle vector state  $|0\rangle \in \Phi_F$ , whose structure in most practical cases is hidden. Even though this method appeared to be very effective for the study of numerous quantum many-particle Hamiltonian systems, some important problems related to the *a priori* non-self-adjointness of the “creation” and “annihilation” operators in the Fock space  $\Phi_F$ , urged researchers to suggest a dual quantization scheme, based strictly on physically “observable” operators in a suitably constructed cyclic Hilbert space  $\Phi$ , generated by means of the so-called “groundstate” vector  $|\Omega\rangle \in \Phi$ , and being completely different from the Fock space  $\Phi_F$ .

Several authors have been developing this idea of quantizing nonrelativistic models, making use of the local current algebra operators [4–10] as the basic dynamical variables, that is the density  $\rho(x) : \Phi \rightarrow \Phi$  and current  $J(x) : \Phi \rightarrow \Phi$  operators at spatial point  $x \in \mathbb{R}^3$ , representing, as is well known, generators of the fundamental physical symmetry group  $\text{Diff}(\mathbb{R}^3) \ltimes \mathcal{S}(\mathbb{R}^3; \mathbb{R})$ , the semidirect product of the diffeomorphism

\*To memory of Nikolai N. Bogolubov, a mathematical physics giant of the XX-th century — on his 110-th Birthday Jubilee.

group  $\text{Diff}(\mathbb{R}^3)$  of the space  $\mathbb{R}^3$  and the Schwarz space of smooth real valued functions on  $\mathbb{R}^3$ . Moreover, the corresponding quantum Hamiltonian operators of the Schrödinger type in the Hilbert space  $\Phi$ , as it appeared to be very surprising, possess a very nice factorized structure, completely determined by this groundstate vector  $|\Omega\rangle \in \Phi$ . This fact posed a very interesting and important problem of studying the related mathematical structure of these factorized operators and the correspondence to the classical Hamiltonian many-particle non-relativistic systems generating them, specified by some kinetic and inter-particle potential energy.

Analytical studies in modern mathematical physics are strongly based on the exactly solvable physical models which are of great help in understanding their mathematical and frequently hidden physical nature. Especially, the solvable models are of great importance in quantum many-particle physics, amongst which one can single out the oscillatory systems and Coulomb systems, modelling phenomena in plasma physics, the well known Calogero-Moser and Calogero-Moser-Sutherland models, describing a system of many particles on an axis, interacting pair-wise through long range potentials, modelling both some quantum-gravity and fractional statistics effects. In this work we developed investigations of local quantum current algebra symmetry representations in suitably renormalized representation Hilbert spaces, suggested and devised before by G.A. Goldin with his collaborators, having further applied their results to constructing the related factorized operator representations for secondly-quantized many-particle integrable Hamiltonian systems. The main technical ingredient of the current algebra symmetry representation approach consists in the weak equivalence of the initial many-particle quantum Hamiltonian operator to a suitably constructed quantum Hamiltonian operator in the factorized form, strictly depending only on its ground state vector. The latter makes it possible to reconstruct the initial quantum Hamiltonian operator in the case of its strong equivalence to the related factorized Hamiltonian operator form, thereby constructing, as a by-product, the corresponding  $N$ -particle groundstate vector for arbitrary  $N \in \mathbb{Z}_+$ . Being uniquely defined by means of the Bethe groundstate vector representation in the Hilbert space, the analyzed factorized operator structure of quantum completely integrable many-particle Hamiltonian systems on the axis proves to be closely related to their quantum integrability by means of the quantum inverse scattering transform. As examples we have studied in detail the factorized structure of Hamiltonian operators, describing such quantum integrable spatially many- and one-dimensional models as generalized oscillatory, Calogero-Moser-Sutherland and nonlinear Schrödinger dynamical systems of spinless bose-particles.

## 2. Fock type Hilbert space, nonrelativistic quantum current algebra and its representations

### 2.1. Preliminaries

Let  $\Phi$  be a separable Hilbert space,  $F$  be a topological real linear space and  $\mathcal{A} := \{A(f) : f \in F\}$  a family of commuting self-adjoint operators in  $\Phi$  (i.e., these operators commute in the sense of their resolutions of the identity) with dense in  $\Phi$  domain  $\text{Dom } A(f) := D_{A(f)}$ ,  $f \in F$ . Consider the Gelfand rigging [11–13] of the Hilbert space  $\Phi$ , i.e., a chain

$$\mathcal{D} \subset \Phi_+ \subset \Phi \subset \Phi_- \subset \mathcal{D}', \quad (2.1)$$

in which  $\Phi_+$  is a Hilbert space, topologically (densely and continuously) and quasi-nucleus (the inclusion operator  $i : \Phi_+ \rightarrow \Phi$  is of the Hilbert-Schmidt type) embedded into  $\Phi$ , the space  $\Phi_-$  is the dual to  $\Phi_+$  as the completion of functionals on  $\Phi_+$  with respect to the norm  $\|f\|_- := \sup_{\|u\|_+ = 1} |(f|u)_\Phi|$ ,  $f \in \Phi$ , a linear dense in  $\Phi_+$  topological space  $\mathcal{D} \subseteq \Phi_+$  is such that  $\mathcal{D} \subset D_{A(f)} \subset \Phi$  and the mapping  $A(f) : \mathcal{D} \rightarrow \Phi_+$  is continuous for any  $f \in F$ . Then, the following structural theorem [5, 7, 11, 12, 14–17] holds.

**Theorem 2.1.** *Assume that the family of operators  $\mathcal{A}$  satisfies the following conditions:*

- a) for  $A(f)$ ,  $f \in F$ , the closure of the operator  $A(f)$  in  $\Phi$  coincides with  $A(f)$  for any  $f \in F$ , that is  $\overline{A(f)} = A(f)$  on domain  $D_{A(f)}$  in  $\Phi$ ;*
- b) the range  $A(f) \subset \Phi$  for any  $f \in F$ ;*

c) for every  $|\psi\rangle \in \mathcal{D}$  the mapping  $F \ni f \rightarrow A(f)|\psi\rangle \in \Phi_+$  is linear and continuous;  
 d) there exists a strong cyclic vector  $|\Omega\rangle \in \bigcap_{f \in F} D_{A(f)}$ , such that the set of all vectors  $|\Omega\rangle$  and  $\prod_{j=1}^n A(f_j)|\Omega\rangle$ ,  $n \in \mathbb{Z}_+$ , is total in  $\Phi_+$  (i.e., their linear hull is dense in  $\Phi_+$ ).

Then, there exists a probability measure  $\mu$  on  $(F', C_\sigma(F'))$ , where  $F'$  is the dual of  $F$  and  $C_\sigma(F')$  is the  $\sigma$ -algebra generated by cylinder sets in  $F'$  such that, for  $\mu$ -almost every  $\eta \in F'$  there is a generalized joint eigenvector  $\omega(\eta) \in \Phi_-$  of the family  $\mathcal{A}$ , corresponding to the joint eigenvalue  $\eta \in F'$ , that is for any  $\varphi \in \mathcal{D} \subset \Phi_+$

$$(\omega(\eta)|A(f)\varphi)_{\Phi_- \times \Phi_+} = \eta(f)(\omega(\eta)|\varphi)_{\Phi_- \times \Phi_+} \quad (2.2)$$

with  $\eta(f) \in \mathbb{R}$  denoting here the pairing between  $F$  and  $F'$ .

The mapping

$$\mathcal{D} \ni |\varphi\rangle \rightarrow (\omega(\eta)|\varphi)_{\Phi_- \times \Phi_+} := \varphi(\eta) \in \mathbb{C} \quad (2.3)$$

for any  $\eta \in F'$  can be continuously extended to a unitary surjective operator  $\mathcal{F}_\eta : \Phi_+ \rightarrow L_2^{(\mu)}(F'; \mathbb{C})$ , where

$$\mathcal{F}_\eta |\varphi\rangle := \varphi(\eta) \quad (2.4)$$

for any  $\eta \in F'$  is a generalized Fourier transform, corresponding to the family  $\mathcal{A}$ . Moreover, the image of the operator  $A(f)$ ,  $f \in F'$ , under the  $\mathcal{F}_\eta$ -mapping is the operator of multiplication by the function  $F' \ni \eta \rightarrow \eta(f) \in \mathbb{C}$ .

Now let us assume that a Hilbert space  $\Phi := \Phi_F$  possesses the standard canonical Fock space structure [1–3, 7, 12, 18–20], that is

$$\Phi_F = \oplus_{n \in \mathbb{Z}_+} \Phi_{(s)}^{\otimes n}, \quad (2.5)$$

where subspaces  $\Phi_{(s)}^{\otimes n}$ ,  $n \in \mathbb{Z}_+$ , are the symmetrized tensor products of a Hilbert space  $\mathcal{H} \simeq L_2^{(s)}(\mathbb{R}^m; \mathbb{C})$ . If a vector  $\varphi := (\varphi_0, \varphi_1, \dots, \varphi_n, \dots) \in \Phi_F$ , its norm

$$\|\varphi\|_\Phi := \left( \sum_{n \in \mathbb{Z}_+} \|\varphi_n\|_n^2 \right)^{1/2}, \quad (2.6)$$

where  $\varphi_n \in \Phi_{(s)}^{\otimes n} \simeq L_2^{(s)}((\mathbb{R}^m)^{\otimes n}; \mathbb{C})$  and  $\|\dots\|_n$  is the corresponding norm in  $\Phi_{(s)}^{\otimes n}$  for all  $n \in \mathbb{Z}_+$ . Note here that concerning the rigging structure (2.1), there holds the corresponding rigging for the Hilbert spaces  $\Phi_{(s)}^{\otimes n}$ ,  $n \in \mathbb{Z}_+$ , that is

$$\mathcal{D}_{(s)}^n \subset \Phi_{(s),+}^{\otimes n} \subset \Phi_{(s)}^{\otimes n} \subset \Phi_{(s),-}^{\otimes n} \quad (2.7)$$

with some suitably chosen dense and separable topological spaces of symmetric functions  $\mathcal{D}_{(s)}^n$ ,  $n \in \mathbb{Z}_+$ . Concerning expansion (2.5) we obtain by means of projective and inductive limits [2, 11, 12, 14] the quasi-nucleus rigging of the Fock space  $\Phi$  in the form (2.1).

Consider now any basis vector  $|(\alpha)_n\rangle \in \Phi_{(s)}^{\otimes n}$ ,  $n \in \mathbb{Z}_+$ , which can be written [3, 12, 21–23] in the following canonical Dirac ket-form:

$$|(\alpha)_n\rangle := |\alpha_1, \alpha_2, \dots, \alpha_n\rangle, \quad (2.8)$$

where, by definition,

$$|\alpha_1, \alpha_2, \dots, \alpha_n\rangle := \frac{1}{\sqrt{n!}} \sum_{\sigma \in S_n} |\alpha_{\sigma(1)}\rangle \otimes |\alpha_{\sigma(2)}\rangle \dots |\alpha_{\sigma(n)}\rangle \quad (2.9)$$

and vectors  $|\alpha_j\rangle \in \mathcal{H}_+ \subset \Phi_{(s)}^{\otimes 1}(\mathbb{R}^m; \mathbb{C}) \simeq \mathcal{H}$ ,  $j \in \mathbb{Z}_+$ , are bi-orthogonal to each other, that is  $(\alpha_k|\alpha_j)_{\mathcal{H}} = \delta_{k,j}$  for any  $k, j \in \mathbb{Z}_+$ . The corresponding scalar product of base vectors as (2.9) is given as follows:

$$\begin{aligned} ((\beta)_n|(\alpha)_n) &= (\beta_n, \beta_{n-1}, \dots, \beta_2, \beta_1|\alpha_1, \alpha_2, \dots, \alpha_{n-1}, \alpha_n) \\ &= \sum_{\sigma \in S_n} (\beta_1|\alpha_{\sigma(1)})_{\mathcal{H}} \dots (\beta_n|\alpha_{\sigma(n)})_{\mathcal{H}} := \text{per}\{(\beta_i|\alpha_j)_{\mathcal{H}}\}_{i,j=1,n}, \end{aligned} \quad (2.10)$$

where “per” denotes the permanent of matrix and  $(\cdot|\cdot)_{\mathcal{H}}$  is the corresponding scalar product in the Hilbert space  $\mathcal{H}$ . Based now on the representation (2.8), one can define an operator  $a^+(\alpha) : \Phi_{(s)}^{\otimes n} \rightarrow \Phi_{(s)}^{\otimes(n+1)}$  for any  $|\alpha\rangle \in \mathcal{H}_-$  as follows:

$$a^+(\alpha)|\alpha_1, \alpha_2, \dots, \alpha_n\rangle := |\alpha, \alpha_1, \alpha_2, \dots, \alpha_n\rangle, \quad (2.11)$$

which is called the “creation” operator in the Fock space  $\Phi_F$ . The adjoint operator  $a(\beta) := (a^+(\beta))^* : \Phi_{(s)}^{\otimes(n+1)} \rightarrow \Phi_{(s)}^{\otimes n}$  with respect to the Fock space  $\Phi_F$  (2.1) for any  $|\beta\rangle \in \mathcal{H}_-$ , called the “annihilation” operator, acts as follows:

$$a(\beta)|\alpha_1, \alpha_2, \dots, \alpha_{n+1}\rangle := \sum_{\sigma \in S_n} (\beta|\alpha_j)_{\mathcal{H}} |\alpha_1, \alpha_2, \dots, \alpha_{j-1}, \hat{\alpha}_j, \alpha_{j+1}, \dots, \alpha_{n+1}\rangle, \quad (2.12)$$

where the “hat” over a vector denotes that it should be omitted from the sequence.

It is easy to check that the commutator relationship

$$[a(\alpha), a^+(\beta)] = (\alpha|\beta)_{\mathcal{H}} \quad (2.13)$$

holds for any vectors  $|\alpha\rangle \in \mathcal{H}$  and  $|\beta\rangle \in \mathcal{H}$ . Expression (2.13), owing to the rigged structure (2.1), can be naturally extended to the general case, when vectors  $|\alpha\rangle$  and  $|\beta\rangle \in \mathcal{H}_-$ , conserving its form. In particular, taking  $|\alpha\rangle := |\alpha(x)\rangle = \frac{1}{\sqrt{2\pi}} e^{i\langle \lambda | x \rangle} \in H_- := L_{2,-}(\mathbb{R}^m; \mathbb{C})$  for any  $x \in \mathbb{R}^m$ , one easily gets from (2.13) that

$$[a(x), a^+(y)] = \delta(x - y), \quad (2.14)$$

where we put, by definition,  $\langle \cdot | \cdot \rangle$  the usual scalar product in the  $m$ -dimensional Euclidean space  $(\mathbb{R}^m; \langle \cdot | \cdot \rangle)$ ,  $a^+(x) := a^+(a(x))$  and  $a(y) := a(a(y))$  for all  $x, y \in \mathbb{R}^m$  and denoted by  $\delta(\cdot)$  the classical Dirac delta-function.

The above construction makes it possible to observe easily that there exists the unique vacuum vector  $|0\rangle \in \oplus_{(s)}^{\otimes n}|_{n=0}$ , such that for any  $x \in \mathbb{R}^m$

$$a(x)|0\rangle = 0, \quad (2.15)$$

and the set of vectors

$$\left( \prod_{j=1}^n a^+(x_j) \right) |0\rangle \in \Phi_{(s)}^{\otimes n} \quad (2.16)$$

is total in  $\Phi_{(s)}^{\otimes n}$ , that is their linear integral hull over the functional spaces  $\Phi_{(s)}^{\otimes n}$  is dense in the Hilbert space  $\Phi_{(s)}^{\otimes n}$  for every  $n \in \mathbb{Z}_+$ . This means that for any vector  $\varphi \in \Phi_F$ , the following canonical representation

$$\varphi = \sum_{n \in \mathbb{Z}_+}^{\oplus} \frac{1}{\sqrt{n!}} \int_{(\mathbb{R}^m)^n} \varphi_n(x_1, \dots, x_n) a^+(x_1) a^+(x_2) \dots a^+(x_n) |0\rangle \quad (2.17)$$

holds with the Fourier type coefficients  $\varphi_n \in \Phi_{(s)}^{\otimes n}$  for all  $n \in \mathbb{Z}_+$  with  $\Phi_{(s)}^{\otimes 1} \simeq \mathcal{H} = L_2(\mathbb{R}^m; \mathbb{C})$ . The latter is naturally endowed with the Gelfand type quasi-nucleus rigging, dual to

$$\mathcal{H}_+ \subset \mathcal{H} \subset \mathcal{H}_-, \quad (2.18)$$

making it possible to construct a quasi-nucleus rigging of the dual Fock space  $\Phi_F := \oplus_{n \in \mathbb{Z}_+} \Phi_{(s)}^{\otimes n}$ . Thereby, the chain (2.18) generates the dual Fock space quasi-nucleolus rigging

$$\mathcal{D} \subset \Phi_{F,+} \subset \Phi_F \subset \Phi_{F,-} \subset \mathcal{D}' \quad (2.19)$$

with respect to the Fock space  $\Phi_F$ , easily following from (2.1) and (2.18).

Construct now the following self-adjoint operator  $\rho(x) : \Phi_F \rightarrow \Phi_F$  as

$$\rho(x) := a^+(x)a(x), \quad (2.20)$$

called the density operator at a point  $x \in \mathbb{R}^m$ , satisfying the commutation properties:

$$\begin{aligned} [\rho(x), \rho(y)] &= 0, \\ [\rho(x), a(y)] &= -a(y)\delta(x-y), \\ [\rho(x), a^+(y)] &= a^+(y)\delta(x-y) \end{aligned} \quad (2.21)$$

for any  $x, y \in \mathbb{R}^m$ .

Now, if we construct the following self-adjoint family  $\mathcal{R} := \left\{ \int_{\mathbb{R}^m} \rho(x)f(x)dx : f \in F \right\}$  of linear operators in the Hilbert space  $\Phi$ , where  $F := \mathcal{S}(\mathbb{R}^m; \mathbb{R})$  is the Schwartz functional space dense in  $H$ , one can derive, making use of theorem 2.1, that there exists the generalized Fourier transform (2.4), such that

$$\Phi = L_2^{(\mu)}(F'; \mathbb{C}) \simeq \int_{F'}^{\oplus} \Phi_{\eta} d\mu(\eta) \quad (2.22)$$

for some Hilbert space sets  $\Phi_{\eta}$ ,  $\eta \in F'$ , and a suitable measure  $\mu$  on  $F'$ , with respect to which the corresponding joint eigenvector  $\omega(\eta) \in \Phi_{\eta}$  for any  $\eta \in F'$  generates the Fourier transformed family  $\{\eta(f) \in \mathbb{R} : f \in F\}$ . Moreover, if  $\dim \Phi_{\eta} = 1$  for all  $\eta \in F'$ , the Fourier transformed eigenvector  $\omega(\eta) := \Omega(\eta) = 1$  for all  $\eta \in F'$ .

Now, we consider the family of self-adjoint operators  $\rho(f) : \Phi_{\eta} \rightarrow \Phi_{\eta}$ ,  $f \in F$ , as generating a unitary family  $\mathcal{U} := \{U(f) : f \in F\}$ , where the operator

$$U(f) := \exp[i\rho(f)] \quad (2.23)$$

is unitary, satisfying the abelian commutation condition

$$U(f_1)U(f_2) = U(f_1 + f_2) \quad (2.24)$$

for any  $f_1, f_2 \in F$ . Since, in general, the unitary family  $\mathcal{U}$  is defined on the Hilbert space  $\Phi_{\eta}$ , the important problem of describing its cyclic representation spaces arises, within which the factorization

$$\rho(f) = \int_{\mathbb{R}^m} a^+(x)a(x)f(x)dx \quad (2.25)$$

jointly with relationships (2.21) hold for any  $f \in F$ . This problem can be treated using mathematical tools devised both within the representation theory of  $\mathbb{C}^*$ -algebras [5, 7, 12, 21] and the Gelfand-Vilenkin [13] approach. Below we will describe the main features of the Gelfand-Vilenkin formalism, being much more suitable for the task, providing a reasonably unified framework of constructing the corresponding representations. The next definitions will be used in our construction.

**Definition 2.1.** Let  $F$  be a locally convex topological vector space,  $F_0 \subset F$  be a finite dimensional subspace of  $F$ . Let  $F^0 \subseteq F'$  be defined by

$$F^0 := \{\sigma \in F' : \sigma|_{F_0} = 0\}, \quad (2.26)$$

and called the annihilator of  $F_0$ .

The quotient space  $F'^0 := F'/F^0$  may be, evidently, identified with  $F'_0 \subset F'$ , the adjoint space of  $F_0$ .

**Definition 2.2.** Let  $Q \subseteq F'^0$ ; then, the subset

$$X_{F^0}^{(Q)} := \{\sigma \in F' : \sigma + F^0 \subset Q\} \quad (2.27)$$

is called the cylinder set with the base  $Q$  and the generating subspace  $F^0$ .

**Definition 2.3.** Let  $n = \dim F_0 = \dim F'_0 = \dim F'^0$ . One says that a cylinder set  $X^{(Q)}$  has Borel base, if  $Q$  is a Borel set, when regarded as a subset of  $\mathbb{R}^m$ .

The family of cylinder sets with Borel base forms an algebra of sets, which is a key stone for defining measurable sets in  $F'$  and the corresponding measures on  $F'$ .

**Definition 2.4.** The measurable sets in  $F'$  are the elements of the  $\sigma$ -algebra generated by the cylinder sets with Borel base.

**Definition 2.5.** A cylindrical measure in  $F'$  is a non-negative  $\sigma$ -pre-additive function  $\mu$  defined on the algebra of cylinder sets with Borel base and satisfying the conditions  $0 \leq \mu(X) \leq 1$  for any  $X$ ,  $\mu(F') = 1$  and  $\mu(\bigcup_{j \in \mathbb{Z}_+} X_j) = \sum_{j \in \mathbb{Z}_+} \mu(X_j)$ , if all sets  $X_j \subset F'$ ,  $j \in \mathbb{Z}_+$ , have a common generating subspace  $F_0 \subset F$ .

**Definition 2.6.** A cylindrical measure  $\mu$  satisfies the commutativity condition if and only if for any bounded continuous function  $\alpha : \mathbb{R}^n \rightarrow \mathbb{R}$  of  $n \in \mathbb{Z}_+$  real variables the function

$$\alpha[f_1, f_2, \dots, f_n] := \int_{F'} \alpha(\eta(f_1), \eta(f_2), \dots, \eta(f_n)) d\mu(\eta) \quad (2.28)$$

is sequentially continuous in  $f_j \in F$ ,  $j = \overline{1, m}$ .

*Remark 2.1.* It is known [5, 12, 13] that in countably normalized spaces, the properties of sequential and ordinary continuity are equivalent.

**Definition 2.7.** A cylindrical measure  $\mu$  is countably additive if and only if for any cylinder set  $X = \bigcup_{j \in \mathbb{Z}_+} X_j$ , which is the union of countable numerous mutually disjoint cylinder sets  $X_j \subset F'$ ,  $j \in \mathbb{Z}_+$ ,  $\mu(X) = \sum_{j \in \mathbb{Z}_+} \mu(X_j)$ .

The next two standard propositions [5, 12, 13, 24, 25], characterizing extensions of the measure  $\mu$  on  $X = \bigcup_{j \in \mathbb{Z}_+} X_j$ , hold.

**Proposition 2.1.** A countably additive cylindrical measure  $\mu$  can be extended to a countably additive measure on the  $\sigma$ -algebra, generated by the cylinder sets with Borel base. Such a measure will also be called a cylindrical measure.

**Proposition 2.2.** Let  $F$  be a nuclear space. Then, any cylindrical measure  $\mu$  on  $F'$ , satisfying the continuity condition, is countably additive.

Concerning the Fourier transform of a cylindrical measure  $\mu$  in  $F'$ , we will use the following natural definitions.

**Definition 2.8.** Let  $\mu$  be a cylindrical measure in  $F'$ . The Fourier transform of  $\mu$  is the nonlinear functional

$$\mathcal{L}(f) := \int_{F'} \exp[i\eta(f)] d\mu(\eta), \quad (2.29)$$

coinciding with the characteristic functional of the measure  $\mu$ .

**Definition 2.9.** The nonlinear functional  $\mathcal{L} : F \rightarrow \mathbb{C}$  on  $F$ , defined by (2.29), is called positive definite, if and only if for all  $f_j \in F$  and  $\lambda_j \in \mathbb{C}$ ,  $j = \overline{1, n}$ , the condition

$$\sum_{j,k=1}^n \bar{\lambda}_j \mathcal{L}(f_k - f_j) \lambda_k \geq 0 \quad (2.30)$$

holds for any  $n \in \mathbb{Z}_+$ .

The following important proposition, owing to Gelfand and Vilenkin [5, 13], Araki [26] and Goldin [5, 6], holds.

**Proposition 2.3.** The functional  $\mathcal{L} : F \rightarrow \mathbb{C}$  on  $F$ , defined by (2.29), is the Fourier transform of a cylindrical measure on  $F'$  if and only if it is positive definite, sequentially continuous and satisfying the condition  $\mathcal{L}(0) = 1$ .

## 2.2. The unitary family and generating functional equations

Suppose now that we have a continuous unitary representation of the unitary family  $\mathcal{U}$  on a suitable Hilbert space  $\Phi$  with a cyclic vector  $|\Omega\rangle \in \Phi$ . Then, we can put

$$\mathcal{L}(f) := (\Omega|U(f)|\Omega) \quad (2.31)$$

for any  $f \in F := \mathcal{S}$ , being the Schwartz space on  $\mathbb{R}^m$ , and observe that functional (2.31) is continuous on  $F$  owing to the continuity of the representation. Therefore, this functional is the generalized Fourier transform of a cylindrical measure  $\mu$  on  $F'$ :

$$(\Omega|U(f)|\Omega) = \int_{\mathcal{S}'} \exp[i\eta(f)] d\mu(\eta). \quad (2.32)$$

From the spectral point of view, based on theorem 2.1, there is an isomorphism between the Hilbert spaces  $\Phi$  and  $L_2^{(\mu)}(F; \mathbb{C})$ , defined by  $|\Omega\rangle \rightarrow \Omega(\eta) = 1$  and  $U(f)|\Omega\rangle \rightarrow \exp[i\eta(f)]$  and next extended by linearity upon the whole Hilbert space  $\Phi$ .

In the non-cyclic case, there exists a finite or countably infinite family of measures  $\{\mu_k : k \in \mathbb{Z}_+\}$  on  $F'$ , with  $\Phi \simeq \oplus_{k \in \mathbb{Z}_+} L_2^{(\mu_k)}(F'; \mathbb{C})$  and the unitary operator  $U(f) : \Phi \rightarrow \Phi$  for any  $f \in \mathcal{S}'$  corresponds in all  $L_2^{(\mu_k)}(F'; \mathbb{C})$ ,  $k \in \mathbb{Z}_+$ , to a multiplication operator on the exponent function  $\exp[i\eta(f)]$ . This means that there exists a single cylindrical measure  $\mu$  on  $F'$  and a  $\mu$ -measurable field of Hilbert spaces  $\Phi_\eta$  on  $F'$ , such that

$$\Phi \simeq \int_{F'}^{\oplus} \Phi_\eta d\mu(\eta), \quad (2.33)$$

with  $U(f) : \Phi \rightarrow \Phi$ , corresponding [13] to the operator of multiplication by  $\exp[i\eta(f)]$  for any  $f \in F$  and  $\eta \in F'$ . Thereby, having constructed the nonlinear functional (2.29) in an exact analytical form, one can retrieve the representation of the unitary family  $\mathcal{U}$  in the corresponding Hilbert space  $\Phi$ , making use of the suitable factorization (2.25) as follows:  $\Phi = \oplus_{n \in \mathbb{Z}_+} \Phi_n$ , where

$$\Phi_n = \prod_{j=1, n} \rho(x_j)|\Omega\rangle, \quad (2.34)$$

for all  $n \in \mathbb{Z}_+$ .

The cyclic vector  $|\Omega\rangle \in \Phi$  can be, in particular, obtained as the ground state vector of some unbounded self-adjoint positive definite Hamilton operator  $\hat{H} : \Phi \rightarrow \Phi$ , commuting with the self-adjoint non-negative particle number operator

$$N := \int_{\mathbb{R}^m} dx \rho(x), \quad (2.35)$$

that is  $[\hat{H}, N] = 0$ . Moreover, the conditions

$$\hat{H}|\Omega\rangle = 0 \quad (2.36)$$

and

$$\inf_{\varphi \in D_H} (\varphi|\hat{H}|\varphi) = (\Omega|\hat{H}|\Omega) = 0 \quad (2.37)$$

hold for the operator  $\hat{H} : \Phi \rightarrow \Phi$ , where  $D_H$  denotes its domain of definition, dense in  $\Phi$ . To find the functional (2.31), which is called the generating Bogolubov type functional for moment distribution functions

$$F_n(x_1, x_2, \dots, x_n) := (\Omega| : \rho(x_1)\rho(x_2) \dots \rho(x_n) : |\Omega\rangle), \quad (2.38)$$

where  $x_j \in \mathbb{R}^m$ ,  $j = \overline{1, n}$ , and the *normal ordering operation*  $: \cdot :$  is defined [2, 3, 5, 8, 9, 15] as

$$: \rho(x_1)\rho(x_2) \dots \rho(x_n) : = \prod_{j=1}^n \left( \rho(x_j) - \sum_{k=1}^{j-1} \delta(x_j - x_k) \right), \quad (2.39)$$

it is convenient first to choose the Hamilton operator  $H : \Phi_F \rightarrow \Phi_F$  in the following secondly quantized [3, 5, 7] representation

$$H := \frac{1}{2} \int_{\mathbb{R}^m} \langle \nabla_x a^+(x) | \nabla_x a(x) \rangle dx + V(\rho), \quad (2.40)$$

in the related Fock space  $\Phi_F$ , where the sign “ $\nabla_x$ ” means the usual gradient operation with respect to  $x \in \mathbb{R}^m$  in the Euclidean space  $\mathbb{E}^m \simeq (\mathbb{R}^m; \langle \cdot | \cdot \rangle)$ . If the energy spectrum density of the Hamiltonian operator (2.40) in the Fock space  $\Phi_F$  is bounded from below, in works done by Goldin, Grodnik, Menikov, Powers and Sharp [5, 7, 27] there was stated that this Hamiltonian, modulo the ground state energy eigenvalue, can be algebraically represented on a suitably constructed *current algebra symmetry representation Hilbert space*  $\Phi$ , as the positive definite gauge type factorized operator

$$\hat{H} = \frac{1}{2} \int_{\mathbb{R}^m} \langle (K^+(x) - A(x; \rho)) | \rho^{-1}(x) (K(x) - A(x; \rho)) \rangle dx, \quad (2.41)$$

satisfying conditions (2.36) and (2.37), where  $A(x; \rho) : \Phi \rightarrow \Phi^m$ ,  $x \in \mathbb{R}^m$ , is some specially constructed linear self-adjoint operator, satisfying the condition

$$K(x) | \Omega \rangle = A(x; \rho) | \Omega \rangle \quad (2.42)$$

for any  $x \in \mathbb{R}^m$  and the ground state  $|\Omega\rangle \in \Phi$ , corresponding to a chosen potential operators  $V(\rho) : \Phi \rightarrow \Phi$ . The singular structure of the operator (2.41) was earlier analyzed in detail in [28] where, in part, there was shown its wellposedness.

The “potential” operator  $V(\rho) : \Phi \rightarrow \Phi$  is, in general, a polynomial (or analytical) functional of the density operator  $\rho(x) : \Phi \rightarrow \Phi$  for any  $x \in \mathbb{R}^m$ , and the operator  $K(x) : \Phi \rightarrow \Phi^m$  is defined as

$$K(x) := \nabla_x \rho(x) / 2 + iJ(x), \quad (2.43)$$

where the self-adjoint “current” operator  $J(x) : \Phi \rightarrow \Phi^m$  can be naturally defined (but non-uniquely) from the continuity equality

$$\partial \rho / \partial t = -i[H, \rho(x)] = -\langle \nabla | J(x) \rangle, \quad (2.44)$$

holding for all  $x \in \mathbb{R}^m$ . Such an operator  $J(x) : \Phi \rightarrow \Phi^m$ ,  $x \in \mathbb{R}^m$ , can exist owing to the commutation condition  $[\hat{H}, \hat{N}] = 0$ , giving rise to the continuity relationship (2.44), if we additionally take into account that supports  $\rho$  of the density operator  $\rho(x) : \Phi \rightarrow \Phi$ ,  $x \in \mathbb{R}^m$ , can be chosen arbitrarily owing to the independence of (2.44) on the potential operator  $V(\rho) : \Phi \rightarrow \Phi$ , but its strict dependence on the corresponding representation (2.33).

*Remark 2.2.* Here we mention that the operator  $K(x) : \Phi \rightarrow \Phi^m$ ,  $x \in \mathbb{R}^m$ , defined by (2.43), relates to that from the work [5, 7, 27] via scaling  $K(x) \rightarrow K(x)/2$ ,  $x \in \mathbb{R}^m$ .

In particular, based on the Fock space  $\Phi_F$ , defined by (2.1) and generated by the creation-annihilation operators (2.11) and (2.12), the current operator  $J(x) : \Phi_F \rightarrow \Phi_F^m$ ,  $x \in \mathbb{R}^m$ , can be easily constructed as follows:

$$J(x) = \frac{1}{2i} [a^+(x) \nabla_x a(x) - \nabla_x a^+(x) a(x)], \quad (2.45)$$

satisfying jointly with the density operator  $\rho(x) : \Phi_F \rightarrow \Phi_F$ ,  $x \in \mathbb{R}^m$ , defined by (2.20), the following quantum current Lie algebra symmetry [4, 5, 7–9, 15, 20] relationships:

$$\begin{aligned} [J(g_1), J(g_2)] &= iJ([g_1, g_2]), \quad [\rho(f_1), \rho(f_2)] = 0, \\ [J(g_1), \rho(f_1)] &= i\rho(\langle g_1 | \nabla f_1 \rangle), \end{aligned} \quad (2.46)$$

holding for all  $f_1, f_1 \in F$  and  $g_1, g_2 \in F^m$ , where we put, by definition,

$$[g_1, g_2] := \langle g_1 | \nabla \rangle g_2 - \langle g_2 | \nabla \rangle g_1, \quad (2.47)$$

being the usual commutator of vector fields  $\langle g_1 | \nabla \rangle$  and  $\langle g_2 | \nabla \rangle$  on the configuration space  $\mathbb{R}^m$ . It is easy to observe that the current algebra (2.46) is the Lie algebra  $\mathcal{G}$ , corresponding to the Banach Lie group



$G = \text{Diff}(\mathbb{R}^m) \ltimes F$ , the semidirect product of the Banach-Lie group of diffeomorphisms  $\text{Diff}(\mathbb{R}^m)$  of the  $m$ -dimensional space  $\mathbb{R}^m$  and the abelian group  $F$  subject to the multiplicative operation Banach group of smooth functions  $F$ .

*Remark 2.3.* The self-adjointness of the operator  $A(g; \rho) : \Phi \rightarrow \Phi^m$ ,  $g \in F$ , can be stated following schemes from works [4, 7, 15] under the additional existence of such a linear anti-unitary mapping  $T : \Phi \rightarrow \Phi$  that the following invariance conditions hold:

$$T\rho(x)T^{-1} = \rho(x), \quad TJ(x)T^{-1} = -J(x), \quad T|\Omega\rangle = |\Omega\rangle \quad (2.48)$$

for any  $x \in \mathbb{R}^m$ . Thereby, owing to conditions (2.48), the following equalities

$$K(x)|\Omega\rangle = A(x; \rho)|\Omega\rangle \quad (2.49)$$

hold for any  $x \in \mathbb{R}^m$ , giving rise to the self-adjointness of the operator  $A(g; \rho) : \Phi \rightarrow \Phi^m$ ,  $g \in F$ .

It is easy to observe that the time-reversal condition (2.48) imposes the real value relationship for the ground state  $\Omega_N = \overline{\Omega}_N \in \Phi_N \simeq L_2^{(s)}(\mathbb{R}^{m \times N}; \mathbb{C})$  of the canonically represented  $N$ -particle Hamiltonian  $H_N : \Phi_N \rightarrow \Phi_N$  for arbitrary  $N \in \mathbb{Z}_+$ . Moreover, taking into account the relationship (2.49), one can easily observe that on the invariant subspace  $\Phi_N \subset \Phi$  the operator  $K(x) : \Phi \rightarrow \Phi^m$  is representable as

$$K_N(x) = \sum_{j=1, \overline{N}} \delta(x - x_j) \frac{\partial}{\partial x_j}, \quad (2.50)$$

entailing the following expression for the related operator  $A_N(x; \rho) : \Phi_N \rightarrow \Phi_N^m$  on the subspace  $\Phi_N \subset \Phi$ :

$$A_N(x; \rho) = \sum_{j=1, \overline{N}} \delta(x - x_j) \nabla_{x_j} \ln |\Omega_N(x_1, x_2, \dots, x_N)|. \quad (2.51)$$

The latter makes it possible to derive its secondly quantized [3, 12] expression as

$$A(x; \rho) = \int_{\mathbb{R}^{m \times N}} dx_2 dx_3 \dots dx_N : \rho(x) \rho(x_2) \rho(x_3) \dots \rho(x_N) : \nabla_x \ln |\Omega_N(x, x_2, \dots, x_N)|, \quad (2.52)$$

which holds for any  $x \in \mathbb{R}^m$  and arbitrary  $N \in \mathbb{Z}_+$ . Being interested in the infinite particle case when  $N \rightarrow \infty$ , the expression (2.52) can be naturally decomposed [29, 30] as

$$\begin{aligned} A(x; \rho) &:= \rho(x) \nabla \frac{\delta}{\delta \rho(x)} W(\rho) \\ &= \sum_{n \in \mathbb{Z}_+} \frac{1}{n!} \int_{\mathbb{R}^{m \times n}} dy_1 dy_2 \dots dy_n : \rho(x) \rho(y_1) \rho(y_2) \rho(y_3) \dots \rho(y_n) : \nabla_x W_{n+1}(x; y_1, y_2, \dots, y_n), \end{aligned} \quad (2.53)$$

where the corresponding real-valued coefficients  $W_n \in H_2^{(1)}(\mathbb{R}^{m \times n}; \mathbb{R})$  should be such functions that the series (2.53) were convergent in a suitably chosen representation Fock space  $\Phi_F$ , for which the resulting ground state  $\lim_{N \rightarrow \infty} \Omega_N \simeq |\Omega\rangle \in \Phi$  is necessarily cyclic in  $\Phi$  and normalized.

Based now on the above construction one easily deduces from expression (2.43) that the generating Bogolubov type functional (2.31) obeys, for all  $x \in \mathbb{R}^m$ , the following functional-differential equation:

$$[\nabla_x - i \nabla_x f] \frac{1}{i} \frac{\delta \mathcal{L}(f)}{\delta f(x)} = A \left( x; \frac{1}{i} \frac{\delta}{\delta f} \right) \mathcal{L}(f), \quad (2.54)$$

whose solutions should satisfy [31] the Fourier transform representation (2.32), and which were, in part, studied in [31]. In particular, a wide class of special so-called Poissonian white noise type solutions to the functional-differential equation (2.54) was obtained in [7, 11, 14, 15, 25, 32] by means of functional-operator methods in the following generalized form:

$$\mathcal{L}(f) = \exp \left[ \int_{\mathbb{R}^m} W \left( \frac{1}{i} \frac{\delta}{\delta f} \right) dx \right] \exp \left( \bar{\rho} \int_{\mathbb{R}^m} \{ \exp[i f(x)] - 1 \} dx \right), \quad (2.55)$$

where  $\bar{\rho} = (\Omega|\rho|\Omega) \in \mathbb{R}_+$  is a suitable Poisson process parameter and the operator  $A(x; \rho) : \Phi \rightarrow \Phi^m$ ,  $x \in \mathbb{R}^m$ , resulting from the expression (2.53) for some scalar operator  $W(\rho) : \Phi \rightarrow \Phi$ .

*Remark 2.4.* It is worth to remark here that solutions to equation (2.54) realize the suitable physically motivated representations of the abelian Banach subgroup  $F$  of the Banach group  $G = \text{Diff}(\mathbb{R}^m) \ltimes F$ , mentioned above. In the general case of this Banach group  $G$ , one can also construct [7, 8, 16] a generalized Bogolubov type functional equation, whose solutions give rise to suitable physically motivated representations of the corresponding current Lie algebra  $\mathcal{G}$ .

Recalling now the Hamiltonian operator representation (2.41), one can readily deduce that the following weak representation Hilbert space  $\Phi$  is a weak relationship

$$\left( \langle A|\rho^{-1}A \rangle - \langle K^*|\rho^{-1}A \rangle - \langle A|\rho^{-1}K \rangle \right) / 2 - V(\rho) = \epsilon_0, \quad (2.56)$$

where  $\epsilon_0 \in \mathbb{R}$  is the corresponding ground state energy density value. Thus, the main analytical problem is now reduced to constructing the expansion (2.53) corresponding to a suitable cyclic representation Hilbert space  $\Phi$  of the quantum current algebra (2.46), compatible with the Hamiltonian operator structure (2.40).

### 2.3. The Hamiltonian operator reconstruction within the cyclic current algebra representation

We assume that we are given a Banach current group  $G = \text{Diff}(\mathbb{R}^m) \ltimes F$  cyclic representation in a Hilbert space  $\Phi$  with respect to  $F$  with a cyclic vector  $|\Omega\rangle \in \Phi_+ \subset \Phi$ . Based on the well-known Araki reconstruction theorem [7, 26] for the canonical Weyl commutation relations, from (2.44) we can first readily obtain

$$[\hat{H}, U(f)] = J(\nabla f)U(f) - \frac{1}{2}\rho(\langle \nabla f_1 | \nabla f_2 \rangle)U(f), \quad (2.57)$$

where  $U(f) = \exp[i\rho(f)]$ ,  $f \in F$ , is an element of the unitary family  $\mathcal{U}$ . The expression (2.57) makes it possible to calculate the bilinear form

$$(U(f_1)\Omega|\hat{H}|U(f_2)\Omega) = (U(f_1)\Omega|J(\nabla f_1)|U(f_2)\Omega) - \frac{1}{2}(U(f_1)\Omega|\rho(\langle \nabla f_1 | \nabla f_2 \rangle)|U(f_2)\Omega) \quad (2.58)$$

for any  $f_1, f_2 \in F$ . Taking into account the symmetry properties (2.48), we finally deduce from (2.58) that for arbitrary functions  $f_1, f_2 \in F$

$$(U(f_1)\Omega|\hat{H}|U(f_2)\Omega) = \frac{1}{2}(U(f_1)\Omega|\rho(\langle \nabla f_1 | \nabla f_2 \rangle)|U(f_2)\Omega). \quad (2.59)$$

The following standard reasonings make it possible to state that the bilinear symmetric form (2.59) determines on  $\Phi$  a self-adjoint non-negative definite Hamiltonian operator  $\hat{H} : \Phi \rightarrow \Phi$ , densely defined on the domain  $D_H := \text{span}_{f \in F} \{\exp[i\rho(f)]|\Omega\rangle \in \Phi\}$ . Really, for any set of functions  $f_j \in F$ ,  $j = \overline{1, n}$ , the following inequalities

$$\sum_{j,k=\overline{1,n}} \bar{s}_j s_k \langle \nabla f_j | \nabla f_k \rangle \geq 0, \quad \sum_{j,k=\overline{1,n}} \bar{s}_j s_k (U(f_j)\Omega|\rho(x)|U(f_k)\Omega) \geq 0 \quad (2.60)$$

hold for any complex numbers  $s_j \in \mathbb{C}$ ,  $j = \overline{1, n}$ , and arbitrary  $n \in \mathbb{N}$ . Since for any non-negative definite complex matrices  $A, B \in \text{End } \mathbb{R}^n$ , the matrix  $C := \{A_{jk}B_{jk} : j, k = \overline{1, n}\} \in \text{End } \mathbb{R}^n$  proves to be non-negative definite [26, 33] too, one ensues that the bilinear form (2.60) is also non-negative definite. Then, as follows from the classical Friedrichs' theorem [17, 34–38], there exists a self-adjoint densely defined and non-negative definite operator  $\hat{H} : \Phi \rightarrow \Phi$ .

### 3. Nonrelativistic many-particle integrable quantum models, their current algebra representations and the Hamiltonian reconstruction

#### 3.1. An integrable many-particle oscillatory quantum model

As a first application of the local current algebra representation construction devised above, we consider a simple nonrelativistic oscillatory quantum model [4] of interacting bose-particles in the  $m$ -dimensional Euclidean space  $(\mathbb{R}^m; \langle \cdot | \cdot \rangle)$ ,  $m \in \mathbb{Z}_+$ , described by the secondly quantized Hamiltonian operator

$$H^{(\omega)} = \frac{1}{2} \int_{\mathbb{R}^m} dx \langle \nabla \psi^+(x) | \nabla \psi(x) \rangle + \frac{1}{2} \int_{\mathbb{R}^m} dx \langle \omega x | \omega x \rangle \psi^+(x) \psi(x), \quad (3.1)$$

acting on the corresponding Fock space  $\Phi$ , and parameterized by the positive definite frequency matrix  $\omega \in \text{End } \mathbb{R}^m$ . It is easy to check that for any  $|\Omega_0\rangle \in \Phi$ , the vector

$$|\varphi_\omega\rangle = \exp\left(-\frac{1}{2} \int_{\mathbb{R}^m} dx \rho(x) \langle x | \omega x \rangle\right) |\varphi_0\rangle \quad (3.2)$$

satisfies the identity

$$K(x) |\varphi_\omega\rangle = \exp\left(-\frac{1}{2} \int_{\mathbb{R}^m} dx \rho(x) \langle x | \omega x \rangle\right) [K(x) + \omega x \rho(x)] |\varphi_0\rangle, \quad (3.3)$$

from which one easily ensues

$$\begin{aligned} \left(H^{(\omega)} - \frac{1}{2} \text{tr } \omega N\right) |\varphi_0\rangle &= \frac{1}{2} \int_{\mathbb{R}^m} dx \langle [K^+(x) + \omega x \rho(x)] | \rho(x)^{-1} [K(x) + \omega x \rho(x)] \rangle |\varphi_0\rangle \\ &= \exp\left(\frac{1}{2} \int_{\mathbb{R}^m} dx \rho(x) \langle x | \omega x \rangle\right) H^{(0)} \exp\left(-\frac{1}{2} \int_{\mathbb{R}^m} dx \rho(x) \langle x | \omega x \rangle\right) |\varphi_0\rangle, \end{aligned} \quad (3.4)$$

meaning that operators  $(H^{(\omega)} - \frac{1}{2} \text{tr } \omega N)$  and  $H^{(0)}$  are equivalent in the Fock space  $\Phi_F$ . Thus, we can formulate the following proposition.

**Proposition 3.1.** *The quantum oscillatory Hamiltonian operator (3.1) permits on the suitable Fock space  $\Phi_F$  the factorized representation*

$$\hat{H}^{(\omega)} = \frac{1}{2} \int_{\mathbb{R}^m} dx \langle [K^+(x) + \omega x \rho(x)] | \rho(x)^{-1} [K(x) + \omega x \rho(x)] \rangle + \frac{1}{2} \text{tr } \omega N. \quad (3.5)$$

Its ground state  $|\Omega^{(\omega)}\rangle \in \Phi$  satisfies the conditions

$$\hat{H}^{(\omega)} |\Omega^{(\omega)}\rangle = \frac{1}{2} \text{tr } \omega N |\Omega^{(\omega)}\rangle, \quad [K(x) + \omega x \rho(x)] |\Omega^{(\omega)}\rangle = 0 \quad (3.6)$$

for all  $x \in \mathbb{R}^m$ .

As a consequence from the relationship (3.6), one can easily obtain that the ground state of the Hamiltonian (3.1) equals

$$|\Omega^{(\omega)}\rangle = \exp\left(-\frac{1}{2} \int_{\mathbb{R}^m} dx \rho(x) \langle x | \omega x \rangle\right) |\Omega^{(0)}\rangle, \quad (3.7)$$

where, by definition,  $K(x)|\Omega^{(0)} = 0$  for all  $x \in \mathbb{R}^m$ , and the related average energy density  $\epsilon_0 = \lim_{N \rightarrow \infty} \frac{N}{2N} \text{tr } \omega = \frac{1}{2} \text{tr } \omega$ . Moreover, as for any  $x, y \in \mathbb{R}^m$ , there holds the equality

$$[\langle D^{(\omega),+}(x)|\rho(x)^{-1}D^{(\omega)}(x)\rangle, \langle D^{(\omega),+}(y)|\rho(y)^{-1}D^{(\omega)}(y)\rangle] = 0, \quad (3.8)$$

where, by definition, the local operator

$$D^{(\omega)}(x) := K(x) + \omega x \rho(x), \quad (3.9)$$

the next operators

$$\hat{H}^{(\omega,p)} = \frac{1}{2} \int_{\mathbb{R}^m} dx \left( \langle D^{(\omega),+}(x)|\rho(x)^{-1}D^{(\omega)}(x)\rangle \right)^p \quad (3.10)$$

on the Fock space  $\Phi$  *a priori* commute to each other, that is

$$[\hat{H}^{(\omega,p)}, \hat{H}^{(\omega,q)}] = 0 \quad (3.11)$$

for any integers  $p, q \in \mathbb{Z}_+$ . Thus, we have stated the following quantum integrability proposition.

**Proposition 3.2.** *The nonrelativistic oscillatory quantum model (3.1) of interacting bose-particles in the  $m$ -dimensional space  $\mathbb{R}^m$  possesses a countable hierarchy of the commuting to each other symmetric operators (3.10) on the suitable Fock space  $\Phi_F$  and represents a quantum completely integrable model.*

The related  $N$ -particle differential expressions for the operators (3.10) can be calculated as their corresponding projections on the  $N$ -particle Fock subspace  $\Phi_N^{(s)} \subset \Phi_F$ ,  $N \in \mathbb{Z}_+$ . In particular, if a vector  $|\varphi_N\rangle \in \Phi_N$  is representable as

$$|\varphi_N\rangle := \int_{\mathbb{R}^{mN}} f_N(x_1, x_2, \dots, x_N) \prod_{j=1, \overline{N}} dx_j \psi^+(x_j) |0\rangle \quad (3.12)$$

for some coefficient function  $f_N \in L_2^{(s)}(\mathbb{R}^{mN}; \mathbb{C})$ , then, by definition,

$$\hat{H}^{(\omega,p)}|\varphi_N\rangle := |\varphi_N^{(p)}\rangle, \quad (3.13)$$

where

$$|\varphi_N^{(p)}\rangle = \int_{\mathbb{R}^{mN}} (H_N^{(\omega,p)} f_N)(x_1, x_2, \dots, x_N) \prod_{j=1, \overline{N}} dx_j \psi^+(x_j) |0\rangle \quad (3.14)$$

for a given  $p \in \mathbb{Z}_+$  and arbitrary finite  $N \in \mathbb{Z}_+$ .

The differential operators  $H_N^{(\omega,p)} : L_2^{(s)}(\mathbb{R}^{mN}; \mathbb{C}) \rightarrow L_2^{(s)}(\mathbb{R}^{mN}; \mathbb{C})$ ,  $p \in \mathbb{Z}_+$  obtained this way, respectively, also commute to each other, as this follows from (3.11), giving rise to the quantum integrability of the  $N$ -particle oscillatory Hamiltonian model  $\hat{H}_N^{(\omega)} = \hat{H}_N^{(\omega,1)} + \frac{1}{2} \text{tr } \omega N$  for arbitrary finite  $N \in \mathbb{Z}_+$ .

### 3.2. A generalized integrable many-particle oscillatory quantum model

A generalized quantum oscillatory model of bose-particles in  $\mathbb{R}^m$  is described by the  $N$ -particle Hamiltonian operator

$$H_N := \frac{1}{2} \sum_{j=1, \overline{N}} \langle \nabla_{x_j} | \nabla_{x_j} \rangle + \frac{1}{2} \sum_{j,k=1, \overline{N}} \langle \omega_N(x_j - x_k) | \omega_N(x_j - x_k) \rangle \quad (3.15)$$

on  $L_2^{(s)}(\mathbb{R}^{mN}; \mathbb{C})$ , parameterized by a positive definite interaction matrix  $\omega_N \in \text{End } \mathbb{R}^m$ ,  $N \in \mathbb{Z}_+$ . In the case when this interaction matrix depends on the particle number  $N \in \mathbb{Z}_+$  as  $\omega_N = \bar{\omega} \sqrt{N/2}$  for

some constant positive definite matrix  $\bar{\omega} \in \text{End } \mathbb{R}^m$ , the corresponding to (3.15) secondly quantized Hamiltonian operator is representable as

$$H = \frac{1}{2} \int_{\mathbb{R}^m} dx \langle \nabla \psi^+(x) | \nabla \psi(x) \rangle + \frac{N}{4} \int_{\mathbb{R}^m \times \mathbb{R}^m} dx dy \psi^+(x) \psi^+(y) \psi(y) \psi(x) \langle \bar{\omega}(x-y) | \bar{\omega}(x-y) \rangle, \quad (3.16)$$

acting on a suitably chosen Fock type representation space  $\Phi_F$ .

Consider now a quasi-local operator mapping  $D(x) : \Phi \rightarrow \Phi^m, x \in \mathbb{R}^m$ , equal to

$$D(x) := K(x) + \int_{\mathbb{R}^m} dy \langle \bar{\omega}(x-y) : \rho(x) \rho(y) : \rangle, \quad (3.17)$$

and construct the next operator expression:

$$\hat{H} = \frac{1}{2} \int_{\mathbb{R}^m} \langle D^+(x) | \rho(x)^{-1} D(x) \rangle. \quad (3.18)$$

Then, the following proposition holds.

**Proposition 3.3.** *The operator expression (3.18) is equivalent on the Fock space  $\Phi_F$  to the secondly quantized Hamiltonian operator (3.16):*

$$\hat{H} = H - \frac{\text{tr } \bar{\omega}}{2} N(N-1). \quad (3.19)$$

*Proof.* It is sufficient to successfully calculate the operator expression (3.18):

$$\begin{aligned} \hat{H} &= \frac{1}{2} \int_{\mathbb{R}^m} dx \langle \nabla \psi^+(x) | \nabla \psi(x) \rangle + \frac{1}{2} \int_{\mathbb{R}^{m \times 2}} dx dy \langle \nabla \psi^+(x) \psi(x) | \bar{\omega}(x-y) \rangle \rho(y) \\ &+ \frac{1}{2} \int_{\mathbb{R}^{m \times 2}} dx dy \rho(y) \langle \bar{\omega}(x-y) | \psi^+(x) \nabla \psi(x) \rangle + \frac{1}{2} \int_{\mathbb{R}^{m \times 3}} dx dy dz \rho(x) \rho(y) \rho(z) \langle \bar{\omega}(x-y) | \bar{\omega}(x-z) \rangle \\ &= \frac{1}{2} \int_{\mathbb{R}^m} dx \langle \nabla \psi^+(x) | \nabla \psi(x) \rangle + \frac{1}{2} \int_{\mathbb{R}^{m \times 2}} dx dy \langle \nabla(\psi^+(x) \psi(x)) | \bar{\omega}(x-y) \rangle \rho(y) \\ &+ \frac{\text{tr } \bar{\omega}}{2} \int_{\mathbb{R}^m} dx \rho(x) + \frac{1}{2} \int_{\mathbb{R}^{m \times 3}} dx dy dz \rho(x) \rho(y) \rho(z) \langle \bar{\omega}(x-y) | \bar{\omega}(x-z) \rangle \\ &= \frac{1}{2} \int_{\mathbb{R}^m} dx \langle \nabla \psi^+(x) | \nabla \psi(x) \rangle - \frac{\text{tr } \bar{\omega}}{2} \int_{\mathbb{R}^{m \times 2}} dx dy \rho(x) \rho(y) \\ &+ \frac{1}{2} \int_{\mathbb{R}^{m \times 3}} dx dy dz \rho(x) \rho(y) \rho(z) \langle \bar{\omega}(x-y) | \bar{\omega}(x-z) \rangle = \frac{1}{2} \int_{\mathbb{R}^m} dx \langle \nabla \psi^+(x) | \nabla \psi(x) \rangle - \frac{\text{tr } \bar{\omega}}{2} N(N-1) \\ &+ \frac{1}{2} \int_{\mathbb{R}^{m \times 3}} dx dy dz \rho(x) \rho(y) \rho(z) \langle \bar{\omega}(x-y) | \bar{\omega}(x-z) \rangle. \end{aligned} \quad (3.20)$$

The latter component of the operator expression (3.20) should be symmetrized to be representable as the

corresponding part of the particle interaction operator  $V(\rho) : \Phi \rightarrow \Phi$  :

$$\begin{aligned}
V(\rho) &:= \frac{1}{2} \int_{\mathbb{R}^{m \times 3}} dx dy dz : \rho(x) \rho(y) \rho(z) : \langle \bar{\omega}(x-y) | \bar{\omega}(x-z) \rangle \\
&= \frac{1}{2 \cdot 3!} \int_{\mathbb{R}^{m \times 3}} dx dy dz : \rho(x) \rho(y) \rho(z) : [\langle \bar{\omega}(x-y) | \bar{\omega}(x-z) \rangle + \langle \bar{\omega}(x-z) | \bar{\omega}(x-y) \rangle \\
&\quad + \langle \bar{\omega}(y-z) | \bar{\omega}(y-x) \rangle + \langle \bar{\omega}(y-x) | \bar{\omega}(y-z) \rangle + \langle \bar{\omega}(z-x) | \bar{\omega}(z-y) \rangle + \langle \bar{\omega}(z-y) | \bar{\omega}(z-x) \rangle] \\
&= \frac{1}{2 \cdot 3!} \int_{\mathbb{R}^{m \times 3}} dx dy dz : \rho(x) \rho(y) \rho(z) : [\langle \bar{\omega}(x-y) | \bar{\omega}(x-z) \rangle + \langle \bar{\omega}(y-z) | \bar{\omega}(y-x) \rangle \\
&\quad + \langle \bar{\omega}(x-z) | \bar{\omega}(x-y) \rangle + \langle \bar{\omega}(z-x) | \bar{\omega}(z-y) \rangle + \langle \bar{\omega}(y-x) | \bar{\omega}(y-z) \rangle + \langle \bar{\omega}(z-y) | \bar{\omega}(z-x) \rangle] \\
&= \frac{1}{2 \cdot 3!} \int_{\mathbb{R}^{m \times 3}} dx dy dz : \rho(x) \rho(y) \rho(z) : [\langle \bar{\omega}(x-y) | \bar{\omega}(x-z) - \bar{\omega}(y-z) \rangle \\
&\quad + \langle \bar{\omega}(x-z) | \bar{\omega}(x-y) - \bar{\omega}(z-y) \rangle + \langle \bar{\omega}(y-x) - \bar{\omega}(z-x) | \bar{\omega}(y-z) \rangle] \\
&= \frac{3}{2 \cdot 3!} \left( \int_{\mathbb{R}^m} dz \rho(z) \right) \int_{\mathbb{R}^{m \times 3}} dx dy : \rho(x) \rho(y) : \langle \bar{\omega}(x-y) | \bar{\omega}(x-y) \rangle \\
&= \frac{N}{4} \int_{\mathbb{R}^{m \times 3}} dx dy : \rho(x) \rho(y) : \langle \bar{\omega}(x-y) | \bar{\omega}(x-y) \rangle. \tag{3.21}
\end{aligned}$$

Having substituted the interaction potential (3.21) into the operator expression (3.20), we obtain

$$\begin{aligned}
H &= \frac{1}{2} \int_{\mathbb{R}^m} dx \langle \nabla \psi^+(x) | \nabla \psi(x) \rangle - \frac{\text{tr } \bar{\omega}}{2} N(N-1) \\
&\quad + \frac{N}{4} \int_{\mathbb{R}^m \times \mathbb{R}^m} dx dy \psi^+(x) \psi^+(y) \psi(y) \psi(x) \langle \bar{\omega}(x-y) | \bar{\omega}(x-y) \rangle, \tag{3.22}
\end{aligned}$$

that is exactly the operator relationship (3.19), proving the proposition.  $\square$

*Remark 3.1.* It is worthy to remark here that owing to its construction, the operator mappings  $\langle D^+(x) | \rho(x)^{-1} D(x) \rangle : \Phi \rightarrow \Phi$ ,  $x \in \mathbb{R}^m$ , commute to each other, that is

$$[\langle D^+(x) | \rho(x)^{-1} D(x) \rangle, \langle D^+(y) | \rho(y)^{-1} D(y) \rangle] = 0 \tag{3.23}$$

for any  $x, y \in \mathbb{R}^m$ . This naturally makes it possible to construct a countable hierarchy of commuting to each other operators  $\hat{H}^{(p)} : \Phi \rightarrow \Phi$ ,  $p \in \mathbb{Z}_+$ , where

$$\hat{H}^{(p)} := \int_{\mathbb{R}^m} dx \left( \langle D^+(x) | \rho(x)^{-1} D(x) \rangle \right)^p, \tag{3.24}$$

that is

$$[\hat{H}^{(p)}, \hat{H}^{(q)}] = 0 \tag{3.25}$$

for all  $p, q \in \mathbb{Z}_+$ . The latter, in particular, means that our generalized quantum oscillatory model (3.15) is also integrable.

Consider now the ground state vector  $|\Omega^{(\omega)}\rangle \in \Phi$ , satisfying the conditions

$$\hat{H}|\Omega^{(\omega)}\rangle = 0, \quad D(x)|\Omega^{(\omega)}\rangle = 0 \tag{3.26}$$

for any  $x \in \mathbb{R}^m$ . Then, as easily follows from (3.19), the oscillatory quantum Hamiltonian (3.16) satisfies the constraint

$$\langle \Omega^{(\omega)} | H | \Omega^{(\omega)} \rangle = \frac{\text{tr } \bar{\omega}}{2} \left( \|N|\Omega^{(\omega)}\rangle\|^2 - \|N^{1/2}|\Omega^{(\omega)}\rangle\|^2 \right) \geq 0, \tag{3.27}$$

meaning, in part, that the quantum oscillatory Hamiltonian (3.15) is bounded from below.

## 4. The spatially one-dimensional integrable quantum systems, their operator symmetries and related quantum current algebra representations

### 4.1. The Calogero-Moser-Sutherland quantum model: the current algebra representation, the Hamiltonian reconstruction and integrability

The periodic Calogero-Moser-Sutherland quantum bosonic model on the finite interval  $[0, l] \simeq \mathbb{R}/[0, l]\mathbb{Z}$  is governed by the  $N$ -particle Hamiltonian

$$H_N := - \sum_{j=1, \overline{N}} \frac{\partial^2}{\partial x_j^2} + \sum_{j \neq k=1, \overline{N}} \frac{\pi^2 \beta(\beta-1)}{l^2 \sin^2[\frac{\pi}{l}(x_j - x_k)]} \quad (4.1)$$

in the symmetric Hilbert space  $L_2^{(s)}([0, l]^N; \mathbb{C})$ , where  $N \in \mathbb{Z}_+$  and  $\beta \in \mathbb{R}$  is an interaction parameter. As it was stated in very interesting and highly speculative works [39], there exist linear differential operators

$$\mathcal{D}_j := \frac{\partial}{\partial x_j} - \frac{\pi\beta}{l} \sum_{k=1, \overline{N}, k \neq j} \cot \left[ \frac{\pi}{l}(x_j - x_k) \right] \quad (4.2)$$

for  $j = \overline{1, N}$ , such that the Hamiltonian (4.1) is factorized as the bounded from below symmetric operator

$$H_N = \sum_{j=1, \overline{N}} \mathcal{D}_j^+ \mathcal{D}_j + E_N, \quad (4.3)$$

where

$$E_N = \frac{1}{3} \left( \frac{\pi\beta}{l} \right)^2 N(N^2 - 1) \quad (4.4)$$

is the groundstate energy of the Hamiltonian operator (4.1), which means that there exists such a vector  $|\Omega_N\rangle \in L_2^{(s)}([0, l]^N; \mathbb{C})$ , satisfying for any  $N \in \mathbb{Z}_+$  the eigenfunction condition

$$H_N |\Omega_N\rangle = E_N |\Omega_N\rangle \quad (4.5)$$

and equals

$$|\Omega_N\rangle = \prod_{j < k=1, \overline{N}} \left\{ \sin \left[ \frac{\pi}{l}(x_j - x_k) \right] \right\}^\beta. \quad (4.6)$$

Being additionally interested in proving the quantum integrability of the Calogero-Moser-Sutherland model (4.1), we proceed to its second quantized representation [3, 12] and study it by means of the current algebra representation approach, described above in section 2 and devised before in [5–9, 27, 30].

The secondly quantized form of the Calogero-Moser-Sutherland Hamiltonian operator (4.1) looks as follows:

$$H = \int_0^l dx \psi_x^+(x) \psi_x(x) + \left( \frac{\pi}{l} \right)^2 \beta(\beta-1) \int_0^l dx \int_0^l dy \frac{\psi^+(x) \psi^+(y) \psi(y) \psi(x)}{\sin^2[\frac{\pi}{l}(x-y)]}, \quad (4.7)$$

acting in the corresponding Fock space  $\Phi_F := \oplus_{n \in \mathbb{Z}_+} \Phi_n^{(s)}$ ,  $\Phi_n^{(s)} \simeq L_2^{(s)}([0, l]^n; \mathbb{C})$ ,  $n \in \mathbb{Z}_+$ . To proceed to the current algebra representation of the Hamiltonian operator (4.7), it would be useful to recall the factorized representation (4.3) and preliminarily construct the following singular Dunkl type [39–41] symmetrized  $N$ -particle singular differential operator

$$D_N(x) := \sum_{j=1, \overline{N}} \delta(x - x_j) \frac{\partial}{\partial x_j} - \frac{\pi\beta}{2l} \sum_{j \neq k=1, \overline{N}} \left\{ \delta(x - x_j) \cot \left[ \frac{\pi}{l}(x_j - x_k) \right] + \delta(x - x_k) \cot \left[ \frac{\pi}{l}(x_k - x_j) \right] \right\} \quad (4.8)$$

in the Hilbert space  $L_2^{(s)}([0, l]^N; \mathbb{C})$ ,  $N \in \mathbb{Z}_+$ , parameterized by a running point  $x \in \mathbb{R}/[0, l]\mathbb{Z}$ . The corresponding secondly quantized representation of the operator (4.8) looks as follows:

$$D(x) = \psi^+(x)\psi_x(x) - \frac{\pi\beta}{l} \int_0^l dy \cot\left[\frac{\pi}{l}(x-y)\right] : \psi^+(x)\psi^+(y)\psi(y)\psi(x) : \quad (4.9)$$

for any  $x \in \mathbb{R}/[0, l]\mathbb{Z}$ , or in the current-density operator representation form as

$$D(x) = K(x) - \frac{\pi\beta}{2l} \int_0^l dy \cot\left[\frac{\pi}{l}(x-y)\right] : \rho(x)\rho(y) : . \quad (4.10)$$

Now, based on the operator (4.10), one can formulate the following proposition, first stated in [27], using a completely different approach.

**Proposition 4.1.** *The secondly quantized Hamiltonian operator (4.7) in the Fock space  $\Phi_F$  is representable in dual to (4.3) factorized form as*

$$H = \int_0^l dx D^+(x)\rho(x)^{-1}D(x) + E, \quad (4.11)$$

where the ground state energy operator  $E : \Phi \rightarrow \Phi$  equals

$$E = \frac{1}{3} \left( \frac{\pi\beta}{l} \right)^2 : N^3 : + \left( \frac{\pi\beta}{l} \right)^2 : N^2 :, \quad (4.12)$$

where

$$N := \int_0^l \rho(x) dx \quad (4.13)$$

is the particle number operator, and satisfies the determining conditions

$$(H - E)|\Omega\rangle = 0, \quad D(x)|\Omega\rangle = 0 \quad (4.14)$$

on the suitably renormalized groundstate  $|\Omega\rangle \in \Phi$  for all  $x \in \mathbb{R}/[0, l]\mathbb{Z}$ . Moreover, for any integer  $N \in \mathbb{Z}_+$ , the corresponding projected vector  $|\Omega_N\rangle := |\Omega\rangle|_{\Phi_N}$  satisfies the following eigenfunction relationships:

$$N|\Omega_N\rangle = N|\Omega_N\rangle,$$

$$\begin{aligned} E|\Omega_N\rangle &= \left( \frac{1}{3} \left( \frac{\pi\beta}{l} \right)^2 : N^3 : + \left( \frac{\pi\beta}{l} \right)^2 : N^2 : \right) |\Omega_N\rangle \\ &= \left[ \frac{1}{3} \left( \frac{\pi\beta}{l} \right)^2 (N^3 - 3N^2 + 2N) + \left( \frac{\pi\beta}{l} \right)^2 N(N-1) \right] |\Omega_N\rangle \\ &= \left[ \frac{1}{3} \left( \frac{\pi\beta}{l} \right)^2 (N^3 - 3N^2 + 2N + 3N^2 - 3N) \right] |\Omega_N\rangle \\ &= \left[ \frac{1}{3} \left( \frac{\pi\beta}{l} \right)^2 N(N^2 - 1) \right] |\Omega_N\rangle := E_N |\Omega_N\rangle, \end{aligned} \quad (4.15)$$

coinciding exactly with the result (4.4).



*Remark 4.1.* When deriving the expression (4.15), we have used the identity

$$\begin{aligned} \rho(x)\rho(y) &= : \rho(x)\rho(y) : + \rho(y)\delta(x-y), \\ \rho(x)\rho(y)\rho(z) &= : \rho(x)\rho(y)\rho(z) : + : \rho(x)\rho(y) : \delta(y-z) \\ &\quad + : \rho(y)\rho(z) : \delta(z-x) + : \rho(z)\rho(x) : \delta(x-y) + : \rho(x)\delta(y-z)\delta(z-x), \end{aligned} \quad (4.16)$$

which holds [2, 3, 7, 30] for the density operator  $\rho : \Phi \rightarrow \Phi$  at any points  $x, y, z \in \mathbb{R}/[0, l]\mathbb{Z}$ .

Observe now that the operator (4.9) can be rewritten down as follows:

$$D(x) = K(x) - A(x), \quad (4.17)$$

where

$$\begin{aligned} K(x) &:= \nabla_x \rho(x)/2 + iJ(x), \\ A(x) &:= \frac{\pi\beta}{l} \int_0^l dy \cot\left[\frac{\pi}{l}(x-y)\right] : \rho(x)\rho(y) : \end{aligned} \quad (4.18)$$

for all  $x \in \mathbb{R}/[0, l]\mathbb{Z}$ . Recalling now the second condition of (4.14), one can rewrite it equivalently as

$$K(x)|\Omega\rangle = A(x)|\Omega\rangle \quad (4.19)$$

on the renormalized groundstate vector  $|\Omega\rangle \in \Phi$  for all  $x \in \mathbb{R}/[0, l]\mathbb{Z}$ . On the other hand, owing to the expression (4.11), we obtain the searched current algebra representation

$$\hat{H} = \int_0^l dx [K^+(x) - A(x)] \rho(x)^{-1} [K(x) - A(x)] \quad (4.20)$$

of the Calogero-Moser-Sutherland Hamiltonian operator (4.1) in a suitably constructed cyclic Hilbert space  $\Phi$ , as it was already demonstrated in the work [27, 30], using the condition (4.19) in the form (2.52).

Now shortly discuss the quantum integrability of the Calogero-Moser-Sutherland model (4.1). Owing to the factorized representation (4.20) one can easily observe that for any integer  $p \in \mathbb{Z}_+$ , the operators  $D^+(x)\rho(x)^{-1}D(x) : \Phi \rightarrow \Phi$ ,  $x \in \mathbb{R}/[0, l]\mathbb{Z}$ , commute to each other and with the particle number operator  $N : \Phi \rightarrow \Phi$ , that is

$$[D^+(x)\rho(x)^{-1}D(x), D^+(y)\rho(y)^{-1}D(y)] = 0, \quad [D^+(x)\rho(x)^{-1}D(x), N] = 0 \quad (4.21)$$

for any  $x, y \in \mathbb{R}/[0, l]\mathbb{Z}$ . As a result of the commutation property (4.21), one easily obtains that, for any integer  $p \in \mathbb{Z}_+$ , the symmetric operators

$$\hat{H}^{(p)} := \int_0^l dx \left( D^+(x)\rho(x)^{-1}D(x) \right)^p \quad (4.22)$$

also commute to each other

$$[H^{(p)}, \hat{H}^{(q)}] = 0 \quad (4.23)$$

for all integers  $p, q \in \mathbb{Z}_+$ , and in particular, commute to the Calogero-Moser-Sutherland Hamiltonian operator (4.11):

$$[\hat{H}^{(p)}, \hat{H}] = 0. \quad (4.24)$$

Concerning the related  $N$ -particle differential expressions for the operators (4.22), it is sufficient to calculate their projections on the  $N$ -particle Fock subspace  $\Phi_N \subset \Phi_F$ ,  $N \in \mathbb{Z}_+$ , where one assumes that there exists a suitable embedding  $\Phi \hookrightarrow \Phi_F$ . Namely, let an arbitrary vector  $|\varphi_N\rangle \in \Phi_N$  be representable as

$$|\varphi_N\rangle := \int_{[0, l]^N} f_N(x_1, x_2, \dots, x_N) \prod_{j=1, N} dx_j \psi^+(x_j) |0\rangle \quad (4.25)$$

for some coefficient function  $f_N \in L_2^{(s)}([0, l]^N; \mathbb{C})$ . Then, by definition,

$$H^{(p)}|\varphi_N\rangle := |\varphi_N^{(p)}\rangle, \quad (4.26)$$

where

$$|\varphi_N^{(p)}\rangle = \int_{[0, l]^N} (H_N^{(p)} f_N)(x_1, x_2, \dots, x_N) \prod_{j=1, N} dx_j \psi^+(x_j) |0\rangle \quad (4.27)$$

for a given  $p \in \mathbb{Z}_+$  any  $N \in \mathbb{Z}_+$ . In particular, for  $p = 2$ , when  $H^{(2)} + E = \hat{H} : \Phi \rightarrow \Phi$ , one easily retrieves the shifted Calogero-Moser-Sutherland Hamiltonian operator (4.1):

$$H_N^{(2)} = - \sum_{j=1, N} \frac{\partial^2}{\partial x_j^2} + \sum_{j \neq k=1, N} \frac{\pi^2 \beta(\beta-1)}{l^2 \sin^2[\frac{\pi}{l}(x_j - x_k)]} - \left(\frac{\pi\beta}{l}\right)^2 \frac{N(N^2-1)}{3}. \quad (4.28)$$

Respectively, for higher integers  $p > 2$ , the resulting  $N$ -particle differential operator expressions  $H_N^{(p)} : L_2^{(s)}([0, l]^N; \mathbb{C}) \rightarrow L_2^{(s)}([0, l]^N; \mathbb{C})$ ,  $N \in \mathbb{Z}_+$ , can be obtained the above described way by means of simple yet well cumbersome calculations, and which will prove to be completely equivalent to those calculated before in the earlier cited nice work [39].

*Remark 4.2.* In the thermodynamical limit, when  $\lim_{N \rightarrow \infty, l \rightarrow \infty} N/\pi l := \bar{\rho} > 0$ , the structural operator  $D(x) : \Phi \rightarrow \Phi$ ,  $x \in \mathbb{R}/[0, l]\mathbb{Z}$ , reduces to

$$\bar{D}(x) := \lim_{N/l \rightarrow \bar{\rho}} D(x) = K(x) - \beta \int_{\mathbb{R}} dy : \frac{\rho(y)\rho(x)}{x-y} :, \quad (4.29)$$

and respectively, the operator (4.1) reduces to

$$\bar{H}_N = - \sum_{j=1, N} \frac{\partial^2}{\partial x_j^2} + \beta(\beta-1) \sum_{j \neq k=1, N} \frac{1}{(x_j - x_k)^2} \quad (4.30)$$

in the Hilbert space  $L_2^{(s)}(\mathbb{R}^N; \mathbb{C})$  for any  $N \in \mathbb{Z}_+$ , whose secondly quantized operator expression in the suitable Fock space  $\Phi$  equals

$$\bar{H} = \int_{\mathbb{R}} dx \left( \bar{D}^+(x) \rho(x)^{-1} \bar{D}(x) + \epsilon_0 \right), \quad (4.31)$$

where  $\epsilon_0 := \lim_{N/l \rightarrow \bar{\rho}} \frac{E_N}{l} = \bar{\rho}^3/3$  denotes the average energy density of the reduced Calogero-Moser-Sutherland Hamiltonian operator (4.30) as  $N \rightarrow \infty$ , exactly coinciding with the results obtained earlier in [30].

## 4.2. An integrable many-particle Coulomb type quantum model on axis

A many particle Coulomb type quantum bose model on the axis is governed by the  $N$ -particle Hamiltonian

$$H_N := - \sum_{j=1, N} \frac{\partial^2}{\partial x_j^2} + \sum_{j \neq k=1, N} \frac{\alpha}{|x_j - x_k|} + \frac{\alpha^2}{3} \sum_{j \neq k \neq s=1, N} \left[ \ln |x_j - x_k| \frac{(x_j - x_k)(x_j - x_s)}{|x_j - x_k||x_j - x_s|} \ln |x_j - x_s| \right. \\ \left. + \ln |x_k - x_j| \frac{(x_k - x_j)(x_k - x_s)}{|x_k - x_j||x_k - x_s|} \ln |x_k - x_s| + \ln |x_s - x_j| \frac{(x_s - x_j)(x_s - x_k)}{|x_s - x_j||x_s - x_k|} \ln |x_s - x_k| \right] \quad (4.32)$$

acting in the Hilbert space  $L_2^{(s)}(\mathbb{R}^N; \mathbb{C})$ ,  $N \in \mathbb{Z}_+$ , is parameterized by a real-valued interaction parameter  $\alpha \in \mathbb{R}$ , which modulates both the *binary* and *ternary* particle interactions. Its secondly quantized representation in a suitably chosen Fock space  $\Phi_F$ , looks as follows:

$$\begin{aligned} H = & \int_{\mathbb{R}} dx \psi_x^+(x) |\psi_x(x)\rangle + \int_{\mathbb{R}^2} dx dy \frac{\alpha}{|x-y|} : \rho(x) \rho(y) : \\ & + \frac{\alpha^2}{3} \int_{\mathbb{R}^3} dx dy dz : \rho(x) \rho(y) \rho(z) : \left[ \ln |x-y| \frac{(x-y)(x-z)}{|x-y||x-z|} \ln |x-z| \right. \\ & \left. + \ln |y-x| \frac{(y-z)(y-x)}{|y-z||y-x|} \ln |y-x| + \ln |z-x| \frac{(z-x)(z-y)}{|z-x||z-y|} \ln |z-y| \right], \end{aligned} \quad (4.33)$$

modulo the infinite renormalization constant operator, responsible for the coinciding points  $x = y \in \mathbb{R}$  of the Coulomb and logarithmic type interaction potentials. On the axis  $\mathbb{R}$ , one can define, for any points  $x \neq y \in \mathbb{R}$  and small positive parameter  $\varepsilon > 0$ , the functional expression  $s(x-y; \varepsilon) := \frac{(x-y)}{\varepsilon |x-y|^{1-\varepsilon}}$ , whose derivative  $\partial s(x-y; \varepsilon) / \partial x = 1/|x-y|^{1-\varepsilon}$ . Moreover, if point  $x \rightarrow y \in \mathbb{R}$ , one has  $\lim_{x \rightarrow y} s(x-y; \varepsilon) = 0$  for any  $\varepsilon > 0$ .

One can observe that for any points  $x \neq y \in \mathbb{R}$  the Hamiltonian operator (4.33) can be equivalently rewritten as

$$\begin{aligned} H = & \int_{\mathbb{R}} dx \psi_x^+(x) |\psi_x(x)\rangle + \frac{\alpha}{2} \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^2} dx dy [\partial s(x-y; \varepsilon) / \partial x + \partial s(y-x; \varepsilon) / \partial y : \rho(x) \rho(y) : \\ & + \frac{\alpha^2}{3} \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^3} dx dy dz \rho(x) \rho(y) \rho(z) : [s(x-y; \varepsilon) s(x-z; \varepsilon) \\ & + s(y-z; \varepsilon) s(y-x; \varepsilon) + s(z-x; \varepsilon) s(z-y; \varepsilon)] - \varepsilon^{-2} N(N^2 - 1) \end{aligned} \quad (4.34)$$

being a well defined quantum operator in the Fock space  $\Phi_F$ . Introduce now, at any point  $x \in \mathbb{R}$ , the quasi-local operator expression

$$D^{(\varepsilon)}(x) := K(x) - \alpha \int_{\mathbb{R}} dy : \rho(x) \rho(y) : s(x-y; \varepsilon), \quad (4.35)$$

acting in the Fock space  $\Phi$ , and construct the following operator:

$$\hat{H}^{(\varepsilon)} = \int_{\mathbb{R}} dx \langle D^{(\varepsilon),+}(x) | \rho(x)^{-1} D^{(\varepsilon)}(x) \rangle. \quad (4.36)$$

Then, one can formulate the next proposition.

**Proposition 4.2.** *The many-particle Coulomb type Hamiltonian operator (4.34) in a suitably chosen Fock space  $\Phi$  is weakly equivalent, as  $\varepsilon \rightarrow 0$ , to the operator expression (4.36), and satisfies the following regularized limiting relationship:*

$$\text{reg} \lim_{\varepsilon \rightarrow 0} \hat{H}^{(\varepsilon)} := \lim_{\varepsilon \rightarrow 0} \left[ \tilde{H}^{(\varepsilon)} - \frac{\alpha^2}{3\varepsilon^2} N(N^2 - 1) \right] = \hat{H}. \quad (4.37)$$

*Proof.* We have from (4.36) that

$$\begin{aligned}
\hat{H}^{(\varepsilon)} &= \int_{\mathbb{R}} dx \psi_x^+(x) \psi_x(x) - \alpha \int_{\mathbb{R}^2} dx dy \rho(y) s(x-y; \varepsilon) \psi^+(x) \psi_x(x) - \alpha \int_{\mathbb{R}^2} dx dy \psi_x^+(x) \psi(x) \rho(y) s(x-y; \varepsilon) \\
&+ \alpha^2 \int_{\mathbb{R}^3} dx dy dz \rho(x) \rho(y) \rho(z) s(x-y; \varepsilon) s(x-z; \varepsilon) = \int_{\mathbb{R}} dx \psi_x^+(x) \psi(x) \\
&+ \alpha \int_{\mathbb{R}^2} dx dy : \rho(x) \rho(y) : \partial s(x-y; \varepsilon) / \partial x + \frac{2\alpha^2}{3!} \int_{\mathbb{R}^3} dx dy dz \rho(x) \rho(y) \rho(z) [s(x-y; \varepsilon) s(x-z; \varepsilon) \\
&+ s(y-z; \varepsilon) s(y-x; \varepsilon) + s(z-x; \varepsilon) s(z-y; \varepsilon)] \\
&= \int_{\mathbb{R}} dx \psi_x^+(x) \psi_x(x) + \alpha \int_{\mathbb{R}^2} dx dy : \rho(x) \rho(y) : \frac{1}{|x-y|^{1-\varepsilon}} \\
&+ \frac{\alpha^2}{3} \int_{\mathbb{R}^3} dx dy dz : \rho(x) \rho(y) \rho(z) : \left[ \ln |x-y| \frac{(x-y)(x-z)}{|x-y||x-y|} \ln |x-z| \right. \\
&+ \ln |y-x| \frac{(y-z)(y-x)}{|y-z||y-x|} \ln |y-x| + \ln |z-x| \frac{(z-x)(z-y)}{|z-x||z-y|} \ln |z-y| \left. \right] \\
&+ \varepsilon^{-2} N(N^2 - 1) + O(\varepsilon). \tag{4.38}
\end{aligned}$$

□

From the relationship (4.38), one follows that

$$H^{(\varepsilon)} = \hat{H} + \frac{\alpha}{3\varepsilon^2} N(N^2 - 1) + O(\varepsilon), \tag{4.39}$$

as  $\varepsilon \rightarrow 0$ . The latter, evidently, means that the weak limit (4.37) holds, proving the proposition.

Taking into account that the quasi-operators  $D^{(\varepsilon),+}(x) \rho(x)^{-1} D^{(\varepsilon)}(x) : \Phi \rightarrow \Phi, x \in \mathbb{R}$ , commute to each other, that is

$$[D^{(\varepsilon),+}(x) \rho(x)^{-1} D^{(\varepsilon)}(x), D^{(\varepsilon),+}(y) \rho(y)^{-1} D^{(\varepsilon)}(y)] = 0 \tag{4.40}$$

for any points  $x, y \in \mathbb{R}$ , one can also construct a countable hierarchy of quantum operators

$$\hat{H}^{(\varepsilon,p)} := \int_{\mathbb{R}} dx \left( D^{(\varepsilon),+}(x) \rho(x)^{-1} D^{(\varepsilon)}(x) \right)^p \tag{4.41}$$

for all  $p \in \mathbb{Z}_+$ , also commuting to each other, that is

$$[\hat{H}^{(\varepsilon,p)}, \hat{H}^{(\varepsilon,q)}] = 0 \tag{4.42}$$

for all  $p, q \in \mathbb{Z}_+$ . Applying then to the operators (4.41) the standard weak regularization scheme, similar to that above, one can obtain, respectively, a countable hierarchy of quantum operators

$$\text{reg} \lim_{\varepsilon \rightarrow 0} \hat{H}^{(\varepsilon,p)} := \hat{H}^{(p)}, \tag{4.43}$$

commuting to each other, that is

$$[\hat{H}^{(p)}, \hat{H}^{(q)}] = 0 \tag{4.44}$$

for all  $p, q \in \mathbb{Z}_+$ . The latter makes it possible to claim that the many-particle Coulomb type quantum bose model (4.32) on the axis is completely integrable.

### 4.3. Quantum many-particle Hamiltonian dynamical system on the axis with $\delta$ -interaction, its quantum symmetries and integrability

In this section, we consider a quantum non-relativistic many-particle bose-system on the axis  $\mathbb{R}$ , governed by the Hamiltonian operator:

$$H_N := - \sum_{j=1, N} \frac{\partial^2}{\partial x_j^2} + \beta \sum_{j \neq k=1, N} \delta(x_j - x_k), \quad (4.45)$$

where  $\alpha, \beta \in \mathbb{R}$  are interaction constants, and acting in the symmetric Hilbert space  $L_2^{(s)}(\mathbb{R}^N; \mathbb{C})$ ,  $N \in \mathbb{Z}_+$ . The corresponding secondly quantized expression [1–3, 18–20] for the Hamiltonian operator (4.45) in the related Fock space  $\Phi_F \simeq \sum_{n \in \mathbb{Z}_+}^{\oplus} L_2^{(s)}(\mathbb{R}^n; \mathbb{C})$  equals

$$H = \int_{\mathbb{R}} dx (\psi_x^+ \psi_x + \beta \psi^+ \psi^+ \psi \psi), \quad (4.46)$$

where the creation  $\psi^+$ - and annihilation  $\psi$ -operators satisfy the canonical commutator relationships

$$\begin{aligned} [\psi(x), \psi^+(y)] &= \delta(x - y), \\ [\psi^+(x), \psi^+(y)] &= 0 = [\psi(x), \psi(y)] \end{aligned} \quad (4.47)$$

for any  $x, y \in \mathbb{R}$ . The Hamiltonian operator (4.46) via the Heisenberg recipe [1, 3, 42] naturally generates on the creation  $\psi^+ : \Phi_F \rightarrow \Phi_F$  and annihilation  $\psi : \Phi_F \rightarrow \Phi_F$  operators the following quantum Schrödinger type evolution flow:

$$\begin{aligned} d\psi/dt &:= \frac{1}{i} [H, \psi] = -i\psi_{xx} + 2i\beta\psi^+\psi^2, \\ d\psi^+/dt &:= \frac{1}{i} [H, \psi^+] = i\psi_{xx}^+ - 2i\beta(\psi^+)^2\psi \end{aligned} \quad (4.48)$$

with respect to the temporal parameter  $t \in \mathbb{R}$ . Subject to the quantum Schrödinger type evolution flow (4.48), the particle number operator  $N = \int_{\mathbb{R}} \rho(x) dx$  and the Hamiltonian operator (4.46) in the Fock space  $\Phi_F$  are its conservative symmetries, that is

$$\frac{d}{dt} N = 0, \quad \frac{d}{dt} H = 0 \quad (4.49)$$

for any  $t \in \mathbb{R}$ . The quantum model (4.46), as is well known [1, 18, 43], presents a completely integrable quantum Schrödinger type dynamical system, possessing an infinite hierarchy of quantum commuting to each other operators in the Fock space  $\Phi_F$ . This result was proved by means of the quantum inverse scattering transform [43], based on the existence of a special so-called Lax type quantum operator linearization in the associated operator-valued space  $C^\infty(\mathbb{R}; \text{End } \Phi_F^2)$ . In what follows below we prove the quantum integrability of the quantum Schrödinger type evolution flow (4.48), making use of the local quantum current algebra representation technique, devised in [5, 7–9, 27], similarly the way this was done in sections above.

Let us define, in the Fock space  $\Phi_F$ , the following structural operator:

$$D^{(\varepsilon)}(x) := K(x) - \beta \int_{\mathbb{R}} dy \vartheta_\varepsilon(x - y) : \rho(x) \rho(y) :, \quad (4.50)$$

where, for any  $\varepsilon > 0$ , the expression  $\vartheta_\varepsilon(x - y) := \vartheta(x - y - \varepsilon) = \{1, \text{ if } x > y - \varepsilon\} \wedge \{0, \text{ if } x \leq y + \varepsilon\}$  for  $x, y \in \mathbb{R}$  denotes the shifted classical Heaviside  $\vartheta$ -function, and construct the following quantum operator:

$$\hat{H}^{(\varepsilon)} := \int_{\mathbb{R}} dx D^{(\varepsilon),+}(x) \rho(x)^{-1} D^{(\varepsilon)}(x). \quad (4.51)$$

The next proposition (4.51) states an equivalence of the quantum Hamiltonian operator (4.46) and the weak operator limit  $\lim_{\varepsilon \rightarrow 0} \hat{H}^{(\varepsilon)}$ .

**Proposition 4.3.** *The many-particle quantum operator (4.46) in a suitably chosen Fock space  $\Phi_F$  is weakly equivalent, as  $\varepsilon \rightarrow 0$ , to the operator expression (4.51), and satisfies the following regularized limiting relationship:*

$$\text{reg} \lim_{\varepsilon \rightarrow 0} H^{(\varepsilon)} := \lim_{\varepsilon \rightarrow 0} (H^{(\varepsilon)} - \beta^2 N^3 / 3) = \hat{H}. \quad (4.52)$$

*Proof.* Having taken into account that  $\vartheta_\varepsilon(x - y)\delta(x - y) = 0 = \vartheta_\varepsilon(x - y)\delta'(x - y)$  and  $\vartheta'_\varepsilon(x - y) = \delta(x - y - \varepsilon)$  for any  $x, y \in \mathbb{R}$ , one can calculate the operator expression (4.51) and obtain:

$$\begin{aligned} H^{(\varepsilon)} &= \int_{\mathbb{R}} dx \psi_x^+(x) \psi_x(x) - \beta \int_{\mathbb{R}^2} dx dy \vartheta_\varepsilon(x - y) [\psi_x^+(x) \psi(x) \rho(y) + \rho(y) \psi^+(x) \psi_x(x)] \\ &\quad + \beta^2 \int_{\mathbb{R}^3} dx dy dz \rho(x) \rho(y) \rho(z) \vartheta_\varepsilon(x - y) \vartheta_\varepsilon(x - z) \\ &= \int_{\mathbb{R}} dx \psi_x^+(x) \psi_x(x) - \beta \int_{\mathbb{R}^2} dx dy \vartheta_\varepsilon(x - y) [\psi_x^+(x) \psi(x) + \psi^+(x) \psi_x(x)] \rho(y) \\ &\quad - \beta \int_{\mathbb{R}^2} dx dy \vartheta_\varepsilon(x - y) [\rho(y), \psi^+(x) \psi_x(x)] + \beta^2 \int_{\mathbb{R}^3} dx dy dz \rho(x) \rho(y) \rho(z) \vartheta_\varepsilon(x - y) \vartheta_\varepsilon(x - z) \\ &= \int_{\mathbb{R}} dx \psi_x^+(x) \psi_x(x) + \beta \int_{\mathbb{R}^2} dx dy \vartheta'_\varepsilon(x - y) \rho(x) \rho(y) \\ &\quad - \beta \int_{\mathbb{R}^2} dx dy [\vartheta_\varepsilon(x - y) \delta(x - y) \psi^+(x) \psi_x(x) - \rho(x) \delta'(x - y)] \\ &\quad + \beta^2 \int_{\mathbb{R}^3} dx dy dz \rho(x) \rho(y) \rho(z) \vartheta_\varepsilon(x - y) \vartheta_\varepsilon(x - z) = \int_{\mathbb{R}} dx \psi_x^+(x) \psi_x(x) \\ &\quad + \beta \int_{\mathbb{R}^2} dx dy \delta(x - y - \varepsilon) \rho(x) \rho(y) + \frac{2\beta^2}{3!} \int_{\mathbb{R}^3} dx dy dz \rho(x) \rho(y) \rho(z) [\vartheta_\varepsilon(x - y) \vartheta_\varepsilon(x - z) \\ &\quad + \vartheta_\varepsilon(y - z) \vartheta_\varepsilon(y - x) + \vartheta_\varepsilon(z - x) \vartheta_\varepsilon(z - y)] \\ &= \int_{\mathbb{R}} dx \psi_x^+(x) \psi_x(x) + \beta \int_{\mathbb{R}^2} dx \rho(x) \rho(x - \varepsilon) + \beta^2 N^3 / 3. \end{aligned} \quad (4.53)$$

□

Insomuch as from the latter expression (4.53) one easily ensues that

$$\lim_{\varepsilon \rightarrow 0} (H^{(\varepsilon)} - \beta^2 N^3 / 3) = H, \quad (4.54)$$

the weak operator relationship (4.52) in the Fock space  $\Phi$  is proved.

It is important to mention now the quasi-local operators  $D^{(\varepsilon),+}(x) \rho(x)^{-1} D^{(\varepsilon)}(x) : \Phi \rightarrow \Phi$ ,  $x \in \mathbb{R}$ , owing to their construction, commute to each other, that is

$$[D^{(\varepsilon),+}(x) \rho(x)^{-1} D^{(\varepsilon)}(x), D^{(\varepsilon),+}(y) \rho(y)^{-1} D^{(\varepsilon)}(y)] = 0 \quad (4.55)$$

for any  $x, y \in \mathbb{R}$ . The latter makes it possible to construct a countable hierarchy of operators

$$\hat{H}^{(\varepsilon,p)} := \int_{\mathbb{R}} dx \left( D^{(\varepsilon),+}(x) \rho(x)^{-1} D^{(\varepsilon)}(x) \right)^p \quad (4.56)$$

for  $p \in \mathbb{Z}_+$ , commuting to each other, that is

$$[\hat{H}^{(\varepsilon,p)}, \hat{H}^{(\varepsilon,q)}] = 0 \quad (4.57)$$

for any  $p, q \in \mathbb{Z}_+$ . Applying to the hierarchy of operators (4.56) the standard weak regularization procedure as  $\varepsilon \rightarrow 0$ , one can construct, respectively, a countable hierarchy of quantum operators

$$\hat{H}^{(p)} := \text{reg} \lim_{\varepsilon \rightarrow 0} \hat{H}^{(\varepsilon, p)} \quad (4.58)$$

on the Fock space  $\Phi$  for  $p \in \mathbb{Z}_+$ , also commuting to each other, that is

$$[\hat{H}^{(p)}, \hat{H}^{(q)}] = 0 \quad (4.59)$$

for any  $p, q \in \mathbb{Z}_+$ , as this naturally follows from the commutator relationship (4.57). The latter then means that the quantum Schrödinger type dynamical system (4.48) is integrable, the other way confirming the classical result of [44].

*Remark 4.3.* It is worth to mention that the following generalized quantum many-particle Hamiltonian bose system

$$\begin{aligned} H_N := & - \sum_{j=1, N} \frac{\partial^2}{\partial x_j^2} + \alpha \sum_{j \neq k=1, N} \delta(x_j - x_k) \\ & + i\beta \sum_{j \neq k=1, N} \left( \frac{\partial}{\partial x_j} \circ \delta(x_j - x_k) + \delta(x_j - x_k) \circ \frac{\partial}{\partial x_k} \right), \end{aligned} \quad (4.60)$$

where  $\alpha, \beta \in \mathbb{R}$  are interaction constants, and acting on the symmetric Hilbert space  $L_2^{(s)}(\mathbb{R}^N; \mathbb{C})$ ,  $N \in \mathbb{Z}_+$ , (that is with  $(\alpha\delta + \beta\delta')$ -interaction potential) generates the corresponding secondly quantized quantum Hamiltonian system

$$\begin{aligned} d\psi/dt &:= \frac{1}{i} [H, \psi] = -i\psi_{xx} + 2\alpha\psi^+\psi\psi + 2\beta\psi^+\psi\psi_x, \\ d\psi^+/dt &:= \frac{1}{i} [H, \psi^+] = i\psi_{xx}^+ - 2i\alpha(\psi^+)^2\psi + 2\beta\psi_x^+\psi\psi \end{aligned} \quad (4.61)$$

with the quantum Hamiltonian operator

$$H = \int_{\mathbb{R}} dx [\psi_x^+\psi_x + \alpha\psi^+\psi^+\psi\psi + i\beta(\psi^+\psi^+\psi_x\psi - \psi_x^+\psi^+\psi\psi)] \quad (4.62)$$

on the Fock space  $\Phi_F$ , which is completely integrable, as it was proved before in [18, 20, 42] by means of the quantum inverse scattering transform. This fact, eventually, allows us to speculate that there exists a suitable local current algebra cyclic representation space  $\Phi$ , allowing one to construct a related structural operator  $D(x) : \Phi \rightarrow \Phi$ ,  $x \in \mathbb{R}$ , factorizing the quantum Hamiltonian operator  $\hat{H} = \int_{\mathbb{R}} D^+(x)\rho(x)^{-1}D(x)$ , and reducing, up to some renormalizing constant operator, to (4.62) on the corresponding Fock space  $\Phi_F$ .

## 5. The density functional representation of the local current algebra symmetry and the factorized structure of quantum integrable many-particle Hamiltonian systems

### 5.1. The density functional representation of the local current algebra: the canonical representation reduction

We are now interested in constructing the density functional representation of the current algebra (2.46) in the Hilbert space  $\Phi \simeq \Phi_\rho$  with the cyclic vector  $|\Omega\rangle = 1 \in \Phi$ . To do this, let us consider first the “creation”  $a^+(x)$  and “annihilation” operators  $a(x)$ ,  $x \in \mathbb{R}^m$ , defined via (2.44) in the canonical Fock space  $\Phi$ , which can be formally represented as

$$a^+(x) = \sqrt{\rho(x)} \exp[-i\vartheta(x)], \quad a(x) = \exp[i\vartheta(x)] \sqrt{\rho(x)}, \quad (5.1)$$

where  $\rho(x) : \Phi \rightarrow \Phi$  is our density operator and  $\vartheta(x) : \Phi \rightarrow \Phi, x \in \mathbb{R}^m$ , is some self-adjoint operator. It is important that the operators  $\rho(x)$  and  $\vartheta(x) : \Phi \rightarrow \Phi$  realize the canonical commutation relationships

$$\begin{aligned} [\rho(x), \rho(y)] &= 0 = [\vartheta(x), \vartheta(y)], \\ [\rho(y), \vartheta(x)] &= i\delta(x - y) \end{aligned} \quad (5.2)$$

for any  $x, y \in \mathbb{R}^m$ . Concerning the current operator  $J(x) : \Phi \rightarrow \Phi, x \in \mathbb{R}^m$ , one can easily obtain its equivalent expression

$$J(x) = \rho(x) \nabla \vartheta(x). \quad (5.3)$$

Based on the canonical relationships (5.2), one can easily obtain, following [4], that

$$\vartheta(x) = \frac{1}{i} \frac{\delta}{\delta \rho(x)} + i\sigma[\rho(x)], \quad (5.4)$$

where  $\sigma[\rho(x)] : \Phi \rightarrow \Phi$  is some function of the density operator  $\rho(x) : \Phi \rightarrow \Phi, x \in \mathbb{R}^m$ . Then, respectively, the current operator (5.3) is representable in  $\Phi$  as follows:

$$J(x) = -i\rho(x) \nabla \frac{\delta}{\delta \rho(x)} + \rho(x) \nabla \sigma[\rho(x)]. \quad (5.5)$$

The functional-operator expression (5.5) proves to make sense [4, 7, 26] as operators in the Hilbert space  $\Phi_\rho \simeq \Phi$  of functional valued complex-functions on the manifold  $\mathcal{M}$ , coordinated by the density parameter  $\rho : \Phi_\rho \rightarrow \Phi_\rho$  and endowed with the scalar product  $(a|b)_{\Phi_\rho} := \int_{\mathcal{M}} \overline{a(\rho)} b(\rho) d\mu(\rho)$  subject to some measure  $\mu$  on  $\mathcal{M}$ . To calculate this measure  $\mu$  on  $\mathcal{M}$ , we present an explicit isomorphism between this Hilbert space  $\Phi_\rho$  and the corresponding Fock space  $\Phi_F$  of spinless bosonic particles in  $\mathbb{R}^m$ . First, we determine the support  $\text{supp } \mu \subset \mathcal{M}$  of the measure  $\mu$ , having assumed that the manifold

$$\mathcal{M} = \cup_{n \in \mathbb{Z}_+} \mathcal{M}_n, \quad (5.6)$$

where  $\mathcal{M}_n := \{a(\rho) : \rho(x) := \sum_{j=1}^n \delta(x - c_j) : a \in C^\infty(F'; \text{End } \Phi_\rho)\}$ , where  $c_j \in \mathbb{R}^m, j = \overline{1, n}, n \in \mathbb{N}$ , are arbitrary vector parameters. The restriction  $d\mu_n$  of the measure  $\mu$  on the submanifold  $\mathcal{M}_n$  can be presented [5, 7, 12, 15, 25] as

$$d\mu_n = \gamma_n(c_1, c_2, \dots, c_n) \prod_{j=\overline{1, n}} dc_j, \quad (5.7)$$

where functions  $\gamma_n : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}_+, n \in \mathbb{N}$ , should be determined from the condition (5.5). In accordance with the manifold structure (5.6), we can decompose the Hilbert space  $\Phi_\rho$  as follows:

$$\Phi_\rho = \oplus_{n \in \mathbb{N}} \Phi_n, \quad (5.8)$$

where the space  $\Phi_n$  depends on the mapping  $\sigma : \mathcal{M} \rightarrow \text{End } (\Phi_\rho)$  and consists of functionals that are bounded on  $\mathcal{M}_n$ , in particular, for any  $a(\rho) \in \mathcal{M}$ , the restrictions  $a(\rho)|_{\Phi_n}, n \in \mathbb{N}$ , consist of functions of vectors  $(c_1, c_2, \dots, c_n) \in \mathbb{R}^{m \times n}, n \in \mathbb{N}$ , respectively. The scalar product in  $\Phi_n, n \in \mathbb{N}$ , is suitably defined by means of the expressions (5.7). Now, we can construct the isomorphism between the Hilbert spaces  $\Phi_n, n \in \mathbb{N}$ , and the corresponding components  $\Phi_n^{(s)}, n \in \mathbb{N}$ , of the related Fock space  $\Phi_F$ , representing spinless bosonic particles in  $\mathbb{R}^m$ . In the Hilbert space  $\Phi_n := \Phi_n^{(\sigma)}, n \in \mathbb{N}$ , one can easily calculate the eigenfunctions  $\varphi_{p_1, p_2, \dots, p_n}^{(\sigma)}(\rho) \in \Phi_n^{(\sigma)}$  of the free Hamiltonian

$$H_0^{(\sigma)} := \frac{1}{2} \int_{\mathbb{R}^m} dx \langle K^+(x) | \rho^{-1}(x) K(x) \rangle \quad (5.9)$$

with structural

$$K(x) := \frac{1}{2} \nabla \rho(x) + iJ^{(\sigma)}(x), \quad K^+(x) := \frac{1}{2} \nabla \rho(x) - iJ^{(\sigma)}(x) \quad (5.10)$$



and the momentum

$$P^{(\sigma)} := \int_{\mathbb{R}^m} dx J^{(\sigma)}(x) \quad (5.11)$$

operators:

$$\begin{aligned} H_0^{(\sigma)} \varphi_{p_1, p_2, \dots, p_n}^{(\sigma)}(\rho) &= \left( \sum_{j=1, \overline{n}} E_j \right) \varphi_{p_1, p_2, \dots, p_n}^{(\sigma)}(\rho), \\ P^{(\sigma)} \varphi_{p_1, p_2, \dots, p_n}^{(\sigma)}(\rho) &= \left( \sum_{j=1, \overline{n}} p_j \right) \varphi_{p_1, p_2, \dots, p_n}^{(\sigma)}(\rho), \end{aligned} \quad (5.12)$$

where  $p_j \in \mathbb{R}^m$ ,  $j = \overline{1, n}$ , are momentums of bose-particles in  $\mathbb{R}^m$ , the operator  $H_0^{(\sigma)} : \Phi_\rho \rightarrow \Phi_\rho$  is given by the expressions (5.9), (2.43) and (5.5) and the operator  $P^{(\sigma)} : \Phi_\rho \rightarrow \Phi_\rho$  is given by the expressions (5.10) and (5.5), respectively, within which the current operator  $J^{(\sigma)}(x) : \Phi_\rho \rightarrow \Phi_\rho$  is realized under the condition  $\nabla \sigma[\rho(x)] := \sigma \rho(x)^{-1} \nabla \rho(x)$  as follows:

$$J^{(\sigma)}(x) = -i \rho(x) \nabla \frac{\delta}{\delta \rho(x)} + i \sigma \nabla \rho(x), \quad (5.13)$$

where  $\sigma \in \mathbb{R}$  is a fixed real-valued parameter. In this case the eigenfunctions  $\varphi_{p_1, p_2, \dots, p_n}^{(\sigma)}(\rho) \in \Phi_n^{(\sigma)}$ ,  $n \in \mathbb{N}$ , can be expressed [4, 10] as

$$\varphi_{p_1, p_2, \dots, p_n}^{(\sigma)}(\rho) = \frac{1}{n!} \tilde{\varphi}_0^{(\sigma)}(\rho) \left( \prod_{j=1, \overline{n}} B_{p_j}(\rho) \cdot 1 \right), \quad (5.14)$$

where

$$\begin{aligned} \tilde{\varphi}_0^{(\sigma)}(\rho) &:= \exp \left[ (\sigma - 1/2) \int_{\mathbb{R}^m} dx \rho(x) \ln \rho(x) \right], \\ B_{p_j}(\rho) &:= \int_{\mathbb{R}^m} dx \exp(i \langle p | x \rangle) \rho(x) \exp \left[ -\frac{\delta}{\delta \rho(x)} \right]. \end{aligned} \quad (5.15)$$

The corresponding  $n$ -particle Fock subspaces  $\Phi_n^{(\sigma)}$ ,  $n \in \mathbb{N}$ , can be naturally represented by means of the vectors

$$|\varphi_n^{(\sigma)}\rangle := \frac{1}{\sqrt{n!}} \int_{\mathbb{R}^{m \times n}} \prod_{j=1, \overline{n}} dp_j f_n^{(\sigma)}(p_1, p_2, \dots, p_n) a^+(p_1) a^+(p_2) \dots a^+(p_n) |0\rangle \quad (5.16)$$

with functions  $f_n^{(\sigma)} \in L_2^{(s)}(\mathbb{R}^{m \times n}, \mathbb{C})$ ,  $n \in \mathbb{N}$ , where

$$a^+(p) := \frac{1}{(2\pi)^{m/2}} \int_{\mathbb{R}^m} dx \exp(i \langle x | p \rangle) a^+(x) \quad (5.17)$$

denotes the momentum creation operator for any  $p \in \mathbb{R}^m$ .

Moreover, any functional  $\varphi_n^{(\sigma)}(\rho) \in \Phi_n^{(\sigma)}$ ,  $n \in \mathbb{N}$ , can be uniquely represented as

$$\varphi_n^{(\sigma)}(\rho) := \int_{\mathbb{R}^{m \times n}} \prod_{j=1, \overline{n}} dp_j f_n^{(\sigma)}(p_1, p_2, \dots, p_n) \varphi_{p_1, p_2, \dots, p_n}^{(\sigma)}(\rho) \quad (5.18)$$

for  $f_n^{(\sigma)} \in L_2^{(s)}(\mathbb{R}^{m \times n}; \mathbb{C})$ , since the following condition

$$\left( B_{p_{n+1}}(\rho) \prod_{j=\overline{1,n}} B_{p_j}(\rho) \cdot 1 \right) \Big|_{\rho=a^+(x)a(x)} |\varphi_n^{(\sigma)}\rangle = 0 \quad (5.19)$$

holds identically for all  $p_j \in \mathbb{R}^m, j = \overline{1, n+1}$ , and arbitrary state  $|\varphi_n^{(\sigma)}\rangle \in \Phi_F, n \in \mathbb{N}$ .

*Remark 5.1.* The condition (5.19) jointly with the constraint  $\int_{\mathbb{R}^m} \rho(x) dx = n$  in  $\Phi_n^{(\sigma)}, n \in \mathbb{N}$ , should be, in general, naturally satisfied for any current algebra representation space  $\Phi$  if and only if  $\rho(x) = \sum_{j=\overline{1,n}} \delta(x - c_j) \in \mathcal{M}_n$  for arbitrary  $n \in \mathbb{N}$ .

As a result of the above construction we can state that the Hilbert spaces  $\Phi_n^{(\sigma)}, n \in \mathbb{N}$ , and Fock subspaces  $\Phi_n^{(\sigma)}, n \in \mathbb{N}$ , are, respectively, isomorphic. As a consequence, we derive that the Hilbert space  $\Phi_\rho$  and the Fock space  $\Phi_F$  are isomorphic too.

Consider now, following [4, 7], the action of the current operator (5.13) on the basic vectors  $\varphi_n^{(\sigma)}(\rho) \in \Phi_n^{(\sigma)}, n \in \mathbb{N}$ :

$$J^{(\sigma)}(x) \varphi_n^{(\sigma)}(\rho) = \bar{\varphi}_0^{(\sigma)}(\rho) \left[ -i\rho(x) \nabla \frac{\delta}{\delta \rho(x)} + i\sigma \nabla \rho(x) \right] \varphi_n^{(\sigma)}(\rho), \quad (5.20)$$

from which one ensues easily at  $\sigma = 1/2$  its  $n$ -particle representation on the functional manifold  $\mathcal{M}_n$ :

$$J^{(1/2)}(x) \varphi_n^{(1/2)}(\rho) \Big|_{\rho(y) = \sum_{j=\overline{1,n}} \delta(y - c_j)} = \sum_{j=\overline{1,n}} \frac{1}{2} [-i\delta(x - c_j) \nabla_{c_j} + i \nabla_{c_j} \circ \delta(x - c_j)] \tilde{f}_n^{(1/2)}(c_1, c_2, \dots, c_n), \quad (5.21)$$

where we took into account that  $\bar{\varphi}_0^{(1/2)}(\rho) = 1$  for all densities  $\rho : \Phi_\rho \rightarrow \Phi_\rho$  and have put, by definition, the Fourier transform

$$\tilde{f}_n^{(1/2)}(c_1, c_2, \dots, c_n) := \int_{\mathbb{R}^{m \times n}} \prod_{j=\overline{1,n}} dp_j f_n^{(1/2)}(p_1, p_2, \dots, p_n) \exp \left( i \sum_{j=\overline{1,n}} \langle p_j | c_j \rangle \right) \quad (5.22)$$

for any fixed particle position vectors  $c_j \in \mathbb{R}^n, j = \overline{1, n}$ , and for arbitrary  $n \in \mathbb{N}$ . The expression (5.21), in particular, means that the current operator  $J^{(1/2)}(x) : \Phi_\rho \rightarrow \Phi_\rho$  is symmetric with respect to the measure  $d\mu_n^{(1/2)} := \beta_n \prod_{j=\overline{1,n}} dc_j$  on each functional submanifold  $\mathcal{M}^n$  for all  $n \in \mathbb{N}$ , where the constants  $\beta_n \in \mathbb{R}_+, n \in \mathbb{N}$ , can be determined from the normalization condition  $\|\varphi_n^{(1/2)}(\rho)\|_{\Phi_n^{(1/2)}} = (\varphi_n^{(1/2)} | \varphi_n^{(1/2)})_{\Phi_n^{(1/2)}}^{1/2}, n \in \mathbb{N}$ . The latter gives rise [5, 12, 15, 25] to the following symbolic measure expression

$$d\mu_n^{(1/2)} := \prod_{x \in \mathbb{R}^m} \delta \left( \rho(x) - \sum_{j=\overline{1,n}} \delta(x - c_j) \right) \prod_{j=\overline{1,n}} \frac{dc_j}{(2\pi)^m} \quad (5.23)$$

for all  $c_j \in \mathbb{R}^n, j = \overline{1, n}$ , and arbitrary  $n \in \mathbb{N}$ .

*Remark 5.2.* As was aptly observed in [4], the choice  $\sigma = 1/2$  makes it possible to realize the current algebra representation in the space  $\mathcal{M}$  of analytic functions, which will be *a priori* assumed further, meaning that the corresponding measure can be symbolically expressed as follows:

$$d\mu_n := \prod_{x \in \mathbb{R}^m} \delta \left( \rho(x) - \sum_{j=\overline{1,n}} \delta(x - c_j) \right) \prod_{j=\overline{1,n}} \frac{dc_j}{(2\pi)^n} \quad (5.24)$$

on the subspace  $\mathcal{M}_n$  for any  $n \in \mathbb{N}$ .

## 5.2. The quantum oscillatory model: the density functional current algebra representation and the Hamiltonian reconstruction

As a classical application of the above construction, one can consider a current algebra representation of the quantum Hamiltonian operator

$$H^{(\omega)} = \frac{1}{2} \int_{\mathbb{R}^m} \langle K(x)^+ | \rho(x)^{-1} K(x) \rangle dx + \frac{1}{2} \int_{\mathbb{R}^m} \langle \omega x | \omega x \rangle \rho(x) dx \quad (5.25)$$

in the corresponding Fock space  $\Phi$  of the generalized quantum  $N$ -particle oscillatory Hamiltonian

$$H_N^{(\omega)} = \frac{1}{2} \sum_{j=1, N} \left( \langle \nabla_{x_j} | \nabla_{x_j} \rangle + \langle \omega x_j | \omega x_j \rangle \right) \quad (5.26)$$

for  $N \in \mathbb{Z}_+$  bose-particles in the  $m$ -dimensional space  $\mathbb{R}^m$  under the external oscillatory potential, parameterized by the positive definite frequency matrix  $\omega \in \text{End } \mathbb{R}^m$ .

Having shifted the representation Hilbert space  $\Phi_\rho$  by the functional  $\tilde{\varphi}_0^{(1/2)}(\rho) := \exp[-\frac{1}{2} \int_{\mathbb{R}^m} \langle x | \omega x \rangle \times \rho(x) dx] \in \Phi_\rho$ , the corresponding current operator (5.13) becomes

$$J^{(\omega)}(x) = -i\rho(x)\nabla \frac{\delta}{\delta\rho(x)} + \frac{i}{2}\nabla\rho(x) - i\omega x\rho(x), \quad (5.27)$$

entailing simultaneously the related  $K$ -operator changing

$$K(x) = \rho(x)\nabla \frac{\delta}{\delta\rho(x)} \rightarrow K^{(\omega)}(x) = \rho(x)\nabla \frac{\delta}{\delta\rho(x)} + \omega x\rho(x) \quad (5.28)$$

for any  $x \in \mathbb{R}^m$ . The latter gives rise, respectively, to the following equivalent current algebra functional representation of the oscillatory Hamiltonian (5.25):

$$\hat{H}^{(\omega)} = \frac{1}{2} \int_{\mathbb{R}^m} \langle K^{(\omega)}(x)^+ | \rho(x)^{-1} K^{(\omega)}(x) \rangle dx + \frac{1}{2} \text{tr } \omega \int_{\mathbb{R}^m} \rho(x) dx, \quad (5.29)$$

found before in (3.5) for every  $N \in \mathbb{N}$ . The shifted current operator (5.27) makes it possible to construct the suitably deformed free particle measure

$$d\mu_1^{(\omega)}(\rho) := \exp\left(-\int_{\mathbb{R}^m} dx \rho(x) \langle x | \omega x \rangle\right) d\mu_1^{(1/2)}(\rho) \quad (5.30)$$

on the one-particle functional manifold  $\mathcal{M}_1$ , for which the following expression

$$(\Omega | \hat{H}^{(\omega)} | U(f) | \Omega) = \int_{\mathcal{M}} \exp[i\rho(f)] d\mu_1^{(\omega)}(\rho) \quad (5.31)$$

holds for any test function  $f \in F$ . The latter, jointly with the related ground state condition  $|\Omega\rangle = 1 \in \Phi_\rho$ , makes it possible to easily calculate the scalar product elements

$$(U(f_1)\Omega | \hat{H}^{(\omega)} | U(f_2)\Omega) = \int_{\mathbb{R}^m} \exp[i f_1(c) + i f_2(c)] \exp(-\langle c | \omega c \rangle) \frac{dc}{(2\pi)^m} \quad (5.32)$$

for any test functions  $f_1, f_2 \in F$ . The expression (5.32) makes it possible to successfully calculate the matrix elements  $(\rho(f_{p_1})\Omega | \hat{H}^{(\omega)} | \rho(f_{p_2})\Omega)$  of the Hamiltonian  $\hat{H}^{(\omega)} : \Phi_\rho \rightarrow \Phi_\rho$  on the corresponding eigenvectors  $\rho(f_p)\Omega \in \Phi_\rho$  for arbitrary  $p = p_1, p_2 \in \mathbb{N}$  and, therefore, to find its spectrum.

Consider now the operator (2.43) taking into account the analytical current representation (5.20) at  $\sigma = 1/2$ :

$$\begin{aligned} K(x)\varphi_n^{(1/2)}(\rho) &= \left[ \rho(x)\nabla \frac{\delta}{\delta\rho(x)} - 1/2\nabla\rho(x) \right] \varphi_n^{(1/2)}(\rho) + 1/2\nabla\rho(x)\varphi_n^{(1/2)}(\rho) \\ &= \rho(x)\nabla \frac{\delta}{\delta\rho(x)} \varphi_n^{(1/2)}(\rho) \end{aligned} \quad (5.33)$$

for any  $n \in \mathbb{N}$ . Having substituted instead of  $\varphi_n^{(1/2)}(\rho) \in \Phi_\rho^{(n)}$ ,  $n \in \mathbb{N}$ , the ground state eigenfunction  $\Omega(\rho) = 1 \in \Phi_\rho$ , we can easily retrieve the earlier derived expression (2.52). Moreover, based on the representation (5.28) and the definition (2.42), one can calculate that

$$K^{(\omega)}(x)\bar{\varphi}_0^{(1/2)}(\rho) = \left[ \rho(x)\nabla \frac{\delta}{\delta\rho(x)} + \omega x\rho(x) \right] \bar{\varphi}_0^{(1/2)}(\rho) = 0 = A^{(\omega)}(x;\rho)\bar{\varphi}_0^{(1/2)}(\rho), \quad (5.34)$$

where  $\bar{\varphi}_0^{(1/2)}(\rho) = \exp[-\frac{1}{2} \int_{\mathbb{R}^m} \langle x|\omega x \rangle \rho(x)dx] \in \Phi_\rho^{(1/2)} \simeq \Phi_\rho$ . The latter means, in particular, that the corresponding multiplication operator  $A^{(\omega)}(x;\rho) = 0$ , or, respectively,

$$K(x)\bar{\varphi}_0^{(1/2)}(\rho) := A(x;\rho)\bar{\varphi}_0^{(1/2)}(\rho) = -\omega x\rho(x)\bar{\varphi}_0^{(1/2)}(\rho), \quad (5.35)$$

where  $\bar{\varphi}_0^{(1/2)}(\rho) := |\Omega(\rho)\rangle \in \Phi_\rho$  is the corresponding ground state vector in  $\Phi_\rho$  for the oscillatory Hamiltonian operator (5.26). Making use of the operator (5.30), based on expression (2.55), one can present a special solution to the functional equation (2.54) in the form

$$\mathcal{L}(f) = \exp\left(-\int_{\mathbb{R}^m} dx \langle \omega x|x \rangle \frac{1}{2i} \frac{\delta}{\delta f(x)}\right) \exp\left(\bar{\rho} \int_{\mathbb{R}^m} \{\exp[i f(x)] - 1\} dx\right), \quad (5.36)$$

confirming similar statements from [7, 8, 30].

## 6. Conclusion

In the work we succeeded in developing an effective algebraic scheme of constructing density operator and density functional representations for the local quantum current algebra and its application to quantum Hamiltonian and symmetry operators reconstruction. We analyzed the corresponding factorization structure for quantum Hamiltonian operators, governing spatially many- and one-dimensional integrable dynamical systems. The quantum generalized oscillatory, Calogero-Moser-Sutherland and nonlinear Schrödinger models of spinless bose-particles were analyzed in detail. The central vector of the density operator current algebra representation proved to be the ground vector state of the corresponding completely integrable factorized quantum Hamiltonian system in the classical Bethe ansatz form. The latter makes it possible to classify quantum completely integrable Hamiltonian systems a priori allowing the factorized form and whose groundstate is of the Bethe ansatz from. These and related aspects of the factorized and completely integrable quantum Hamiltonian systems are planned to be studied in other place.

## 7. Acknowledgements

Authors would like to convey their warm thanks to Prof. Gerald A. Goldin for many discussions of the work and instrumental help in editing a manuscript during the XXVIII International Workshop on “Geometry in Physics”, held on 30.06–07.07.2019 in Białowieża, Poland. They also are cordially appreciated to Profs. Joel Lebowitz, Denis Blackmore and Nikolai N. Bogolubov (Jr.) for instructive discussions, useful comments and remarks on the work. A special authors’ appreciation belongs to

Prof. Joel Lebowitz for the invitation to take part in the 121-st Statistical Mechanics Conference, held on May 12–14, 2019 in the Rutgers University, New Brunswick, NJ, USA. Personal A.P.'s acknowledgement belongs to the Department of Physics, Mathematics and Computer Science of the Cracow University of Technology for a local research grant F-2/370/2018/DS.

## References

1. Takhtajan L.A., *Quantum Mechanics for Mathematicians*, American Mathematical Society, Providence, USA, 2008.
2. Berezin F.A., *The Method of Second Quantization*, Academic Press, New York, London, 1966.
3. Bogolubov N.N., Bogolubov N.N. (Jr.), *Introduction to Quantum Statistical Mechanics*, World Scientific, New Jersey, 1986.
4. Aref'eva I.Ya., *Theor. Math. Phys.*, 1972, **10**, 146–155, doi:10.1007/BF01090726.
5. Goldin G.A., *J. Math. Phys.*, 1971, **12**, 462–487, doi:10.1063/1.1665610.
6. Goldin G.A., In: *Contemporary Problems in Mathematical Physics: Proceedings of the Third International Workshop (Cotonou, Republic of Benin, 2003)*, Govaerts J., Hounkonnou M.N., Msezane A.Z. (Eds.), World Scientific, New Jersey, 2004, 3–93, doi:10.1142/9789812702487\_0001.
7. Goldin G.A., Grodnik J., Powers R.T., Sharp D.H., *J. Math. Phys.*, 1974, **15**, 88–100, doi:10.1063/1.1666513.
8. Goldin G.A., Menikoff R., Sharp D.H., *Phys. Rev. Lett.*, 1983, **51**, 2246–2249, doi:10.1103/PhysRevLett.51.2246.
9. Goldin G.A., Menikoff R., Sharp D.H., *J. Math. Phys.*, 1981, **22**, 1664–1668, doi:10.1063/1.525110.
10. Pardee W.J., Schlessinger L., Wright J., *Phys. Rev.*, 1968, **175**, 2140, doi:10.1103/PhysRev.175.2140.
11. Berezanskii Yu.M., *Expansions in Eigenfunctions of Selfadjoint Operators*, American Mathematical Society, Providence, USA, 1968.
12. Berezin F.A., Shubin M.A., *The Schrödinger Equation*, Springer Science+Business Media, Dordrecht, 2012.
13. Gelfand I., Vilenkin N., *Generalized Functions*, Academic Press, New York, 1964.
14. Berezansky Y.M., Kondratiev Y.G., *Spectral Methods in Infinite-Dimensional Analysis*, Springer Science+Business Media, Dordrecht, 1995.
15. Bogolubov N.N. (Jr.), Prykarpatsky A.K., *Phys. Elem. Part. At. Nucl.*, 1986, **17**, 791–827 (in Russian).
16. Prykarpatsky A., Bogoliubov N. (Jr.), Golenia J., Taneri U., *Int. J. Theor. Phys.*, 2008, **47**, 2882–2897, doi:10.1007/s10773-008-9721-2.
17. Reed M., Simon B., *Theory of Operators*, Academic Press, San Diego, 1978.
18. Blackmore D., Prykarpatsky A.K., Samoilenko V.Hr., *Nonlinear Dynamical Systems of Mathematical Physics: Spectral and Differential-Geometrical Integrability Analysis*, World Scientific Publishing, New Jersey, USA, 2011.
19. Faddeev L.D., Yakubovskii O.A., *Lectures on Quantum Mechanics for Mathematics Students*, American Mathematical Society, Providence, USA, 2009.
20. Prykarpatsky A., Mykytiuk I., *Algebraic Integrability of Nonlinear Dynamical Systems on Manifolds: Classical and Quantum Aspects*, Springer Science+Business Media, Dordrecht, 1998.
21. Dirac P.A.M., *The Principles of Quantum Mechanics*, Clarendon Press, Oxford, 1935.
22. Fock V., *Z. Phys.*, 1932, **75**, 622–647, doi:10.1007/BF01344458.
23. Prykarpatsky A.K., Taneri U., Bogolubov N.N. (Jr.), *Quantum Field Theory and Application to Quantum Nonlinear Optics*, World Scientific, New York, 2002.
24. Alberverio S., Kondratiev Y.G., Streit L., *How to Generalize White Noise Analysis to Non-Gaussian Measures*, Preprint Bi-Bo-S, Bielefeld, 1992.
25. Alberverio S., Daletskii A., Kondratiev Yu., Lytvynov E., *J. Geom. Phys.*, 2003, **47**, 259–302, doi:10.1016/S0393-0440(02)00221-8.
26. Araki H., *J. Math. Phys.*, 1960, **1**, 492–504, doi:10.1063/1.1703685.
27. Menikoff R., *J. Math. Phys.*, 1974, **15**, 1394–1408, doi:10.1063/1.1666822.
28. Goldin G.A., Sharp D.H., In: *Group Representations in Mathematics and Physics*, Bargmann V. (Ed.), Springer-Verlag, Berlin, 1970, 300–311.
29. Feenberg E., *Ann. Phys.*, 1974, **84**, 128–137, doi:10.1016/0003-4916(74)90296-6.
30. Menikoff R., Sharp D., *J. Math. Phys.*, 1975, **16**, 2341–2352, doi:10.1063/1.522495.
31. Goldin G.A., Sharp D.H., In: *Magic Without Magic: John Archibald Wheeler*, Klauder J.R. (Ed.), Freeman, San Francisco, 1972, 171–185.
32. Berezansky Y.M., *Rep. Math. Phys.*, 1996, **38**, 289–300, doi:10.1016/S0034-4877(97)84882-3.
33. Beckenbach E.F., Bellman R., *Inequalities*, Springer-Verlag, Berlin, 1961.

34. Friedrichs K., Math. Ann., 1934, **109**, 465–487, doi:10.1007/BF01449150.
35. Friedrichs K., Math. Ann., 1934, **109**, 685–713, doi:10.1007/BF01449164.
36. Friedrichs K., Math. Ann., 1935, **110**, 777–779, doi:10.1007/BF01448058.
37. Kato T., Perturbations Theory of Linear Operators, Springer-Verlag, Berlin, 1966.
38. Reed M., Simon B., Functional Analysis, Academic Press, San Diego, 1987.
39. Lapointe L., Vinet L., Commun. Math. Phys., 1996, **178**, 425–452, doi:10.1007/BF02099456.
40. Aniceto I., Avan J., Jevicki A., J. Phys. A: Math. Theor., 2010, **43**, 185201, doi:10.1088/1751-8113/43/18/185201.
41. Dunkl C.F., Trans. Am. Math. Soc., 1989, **311**, 167–183, doi:10.1090/S0002-9947-1989-0951883-8.
42. Mitropolsky Yu.A., Bogolubov N.N., Prykarpatsky A.K., Samoylenko V.Hr., Integrable Dynamical Systems, Naukova Dumka, Kyiv, 1987, (in Russian).
43. Sklyanin E.K., J. Sov. Math., 1982, **19**, 1546–1596, doi:10.1007/BF01091462.
44. Sklyanin E.K., Takhtadzhyan L.A., Faddeev L.D., Theor. Math. Phys., 1979, **40**, 688–706, doi:10.1007/BF01018718.

## Представлення алгебри струмів для квантових багаточастинкових гамільтонових систем типу Шредінгера, їх факторизована структура та інтегровність

Д. Пророк<sup>1</sup>, А.К. Прикарпатський<sup>2</sup>

<sup>1</sup> Факультет фізики та прикладної інформатики, Університет науки та технологій, Краків, Польща

<sup>2</sup> Факультет фізики, математики та інформатики, Краківський технологічний університет, Краків, Польща

Розвинута ефективна схема дослідження представлень квантової алгебри струмів та алгебраїчної реконструкції факторизованих квантових операторів Гамільтона та їх симетрій у просторі типу Фока. Її застосовано до квантових операторів Гамільтона та їх симетрій у випадку квантових інтегровних просторово одно- та багатовимірних динамічних систем. В якості прикладів нами детально вивчено факторизовану структуру гамільтонових операторів, що описують такі квантові інтегровні просторово багатовимірні та одновимірні моделі як узагальнені осциляторні, модель Калоджеро-Мозера-Сазерленда, типу Кулона та нелінійна динамічна система Шредінгера для системи безспінових бозе-частинок.

**Ключові слова:** *простір Фока, представлення алгебри струмів, реконструкція гамільтоніана, породжуючий функціонал Боголюбова, модель Калоджеро-Мозера-Сазерленда, квантова інтегровність, квантові симетрії*