

DEDUCING A VARIATIONAL PRINCIPLE WITH MINIMAL *A PRIORI* ASSUMPTIONS

ANDREW KRIEGER, GEORG MENZ, AND MARTIN TASSY

ABSTRACT. We study the well-known variational and large deviation principle for graph homomorphisms from \mathbb{Z}^m to \mathbb{Z} . We provide a robust method to deduce those principles under minimal *a priori* assumptions. The only ingredient specific to the model is a discrete Kirszbraum theorem i.e. an extension theorem for graph homomorphisms. All other ingredients are of a general nature not specific to the model. They include elementary combinatorics, the compactness of Lipschitz functions and a simplicial Rademacher theorem. Compared to the literature, our proof does not need any other preliminary results like e.g. concentration or strict convexity of the local surface tension. Therefore, the method is very robust and extends to more complex and subtle models, as e.g. the homogenization of limit shapes or graph-homomorphisms to a regular tree.

CONTENTS

1. Introduction	1
2. Setting and main results	7
3. Outline and discussion of proof of main results	17
4. Microscopic entropy and surface tension	20
5. Profile theorem for piecewise affine functions	27
6. Proof of the profile theorem	36
7. Proof of the variational principle	49
8. Large deviations principle	53
Acknowledgment	56
References	56

1. INTRODUCTION

Recently the study of limit shapes has attracted a lot of research. Limit shapes appear in many models, including domino tilings and

Date: January 10, 2022.

This research has been partially supported by NSF grant DMS-1712632.

dimer models (e.g. [Kas63, CEP96, CKP01]; see Figures 1a, 1b, and 1c), polymer models, lozenge tilings (e.g. [Des98, LRS01, Wil04]), Ginzburg-Landau models (e.g. [DGI00, FO04]), Gibbs models (e.g. [She05]), the Ising model (e.g. [DKS92, Cer06]), asymmetric exclusion processes (e.g. [FS06]), random matrices (e.g. [Wig59, KS99]), sandpile models (e.g. [LP08]), the six vertex model (e.g. [BCG16, CS16, RS18]), and Young tableaux (e.g. [LS77, VK77, PR07]). The appearance of limit shapes seems to be a universal phenomenon: it has been rigorously proven for many models, and it is strongly suggested by simulations for many additional models.

One can explain the appearance of limit shapes with the help of three technical results. Those are a profile theorem, a variational principle, and a large deviations principle. The profile theorem asymptotically counts the number of microscopic states, that contribute to a particular macroscopic state, i.e. a limiting profile. Given a limiting profile h , we




FIGURE 1. Examples of limit shapes.

might write the profile theorem as

$$\text{Ent}_n(\{\text{microscopic states corresponding to } h\}) \approx \text{Ent}(h),$$

where the microscopic entropy

$$\text{Ent}_n(A) := -\frac{1}{n^m} \log |A|$$

captures the number of microscopic states in a set A , and the macroscopic entropy functional is given by

$$\text{Ent}(h) = \int \text{ent}(\nabla h(x)) \, dx.$$

The integrand $\text{ent}(\nabla h(x))$ is a local quantity, depending only on the gradient ∇h at x . This article calls $\text{ent}(\cdot)$ the local surface tension (see Definition 2.12 below). See also Theorem 2.15 for a more precise statement.

The variational principle asymptotically characterizes the number of microscopic states, i.e. the microscopic entropy Ent_n , via a variational problem. For large system sizes n , the microscopic entropy of the system Ent_n is given by minimizing the macroscopic entropy $\text{Ent}(h)$ over all admissible limiting profiles h . In formula, the variational principle states (see Theorem 2.16 below)

$$\text{Ent}_n \approx \inf_h \text{Ent}(h).$$

The large deviations principle complements the variational principle. It characterizes the asymptotic fraction of microscopic states in “natural” subsets of the microscopic state space, such as (open or closed) balls around a limiting profile. Approximately speaking, the large deviations principle states that the fraction of microscopic states in a suitable subset A decays exponentially in n , with rate given by minimizing a rate function $I(h)$ over the limiting profiles corresponding to A (see Theorem 2.17 below). The rate function is, up to normalization, the same as the macroscopic entropy $\text{Ent}(h)$ above.

When the macroscopic entropy has a unique minimizer, the large deviations principle implies that almost all microscopic states must approximate this entropy-minimizing profile. Although the current article does not prove uniqueness of this minimizer, we briefly discuss uniqueness in Section 2.5, after the statement of the large deviations principle (Theorem 2.17).

The current article continues the line of research started in [MT16], which strives to develop a robust theory of variational principles and limit shapes. The intention of this article is not to provide new results on subtle and technically challenging models (as for example in [MT16]) but to provide and explain a simplified method that only relies on minimal *a priori* assumptions. In order not to be distracted by unnecessary technical difficulties we consider the most simple setting: graph homomorphisms from \mathbb{Z}^m to \mathbb{Z} .

In more detail, we consider a macroscopic (continuum) domain $R \subset \mathbb{R}^m$, and a sequence of microscopic (discrete) domains $R_n \subset \mathbb{Z}^m$ such that $\frac{1}{n}R_n \rightarrow R$ in the Hausdorff metric. A microscopic state or a height function is a graph homomorphism $h_{R_n} : R_n \rightarrow \mathbb{Z}$ for some n , and a macroscopic state or asymptotic height function is a Lipschitz function $h_R : R \rightarrow \mathbb{R}$. In the two-dimensional case, this model is equivalent to the six-vertex model with uniform weights (cf. [vB77, CPST18]). Full details of the model under study are given in Section 2.

Our method emerges from distilling the core arguments of [CKP01, She05, MT16]. Compared to those works, our method does not rely on explicit formulas for the local surface tension, strict convexity, concentration inequalities, or the FKG inequality. The robustness of this method is illustrated in the companion article [KMT17]. There, we show the homogenization of the variational principle of graph homomorphisms to \mathbb{Z} . In homogenization, homomorphisms are not chosen according to the uniform measure but instead certain heights are preferred or penalized according to a random field. Mathematically the height function is sampled from a Gibbs measure with respect to a randomized Hamiltonian. The limit shape may change drastically; for example, when the random field is unbounded, simulations show the formation of terraces. See Figures 2 and 3 for examples. We hope that the method outlined in this article can serve as a guiding principle for deducing the variational principle and related results for more complex models.

As hinted above, the three main results of this article are the profile theorem (Theorem 2.15), the variational principle (Theorem 2.16), and the large deviations principle (Theorem 2.17). The majority of the effort in this article goes into proving the profile theorem. The proof starts by proving the profile theorem in a special case (where the domain R is the union of simplices and the limiting profile h_R is piecewise affine), then bootstraps this result to the general case. The main




FIGURE 2. Quenched Aztec diamonds, 2d perspective.




FIGURE 3. Quenched Aztec diamonds, 3d perspective.

idea of each step in this proof is clear, although some care is needed to account for all the details.

We call attention to two ingredients in the proof. The first ingredient is the simplicial Rademacher theorem, so called because it approximates a Lipschitz function h_R uniformly over a large portion of its domain by a piecewise affine approximation h_K , where the “pieces” on which h_K is affine are simplices. This approximation gives control over both the direct error $|h_K - h_R|$ and over the error in the derivatives $|\nabla h_K - \nabla h_R|$. Compared to the classical Rademacher theorem

which states that h_R is almost everywhere differentiable, the simplicial Rademacher theorem is a surprisingly strong approximation result. This seems like a standard result but the authors have not found it stated in this form in the random surfaces literature, so we give details of the proof.

In order to exploit the simplicial Rademacher theorem, we need robustness of both the macroscopic and microscopic entropy, under changes both to the limiting profile and to the domain. Robustness of the macroscopic entropy follows from elementary analysis, because the simplicial Rademacher approximation has derivative ∇h_K close to ∇h_R . Robustness of the microscopic entropy rests largely upon the second ingredient that we call attention to: a Kirszbraun theorem for graph homomorphisms (see Theorem 3.1). This theorem gives conditions under which a height function $h_{R_n} : R_n \rightarrow \mathbb{Z}$ may be extended to a larger domain $\tilde{R}_n \supseteq R_n$. We expect that the main challenge in extending the method of this article to other models will be proving a comparable extension theorem.

The profile theorem can be used to prove the variational principle and the large deviations principle. Both proofs are similar, and rely on the local compactness of the space of Lipschitz functions. Moreover the two proofs are robust; once the profile theorem is proven for a model (with suitable macroscopic state space), the variational principle and large deviations principle follow automatically. We present the proof of the variational principle first and in greater detail. For the large deviations principle we highlight the differences in proof, and we also change notation (replacing symbols like Ent_n and Ent), in order to match the conventions of large deviations theory.

OVERVIEW OF REMAINING ARTICLE

In Section 2, we define the setting and formulate the main results of this article. In Section 3 we explain the main idea and the structure of the proofs of the main results. The details are then given in Sections 4 through 8.

NOTATION

- x and y usually denote points in \mathbb{R}^m .
- z usually denotes a points in \mathbb{Z}^m .

- For $x \in \mathbb{R}$, $\lfloor x \rfloor$ denotes the largest integer $\leq x$, and $\lceil x \rceil$ denotes the smallest integer $\geq x$.
- For $x = (x_1, \dots, x_m) \in \mathbb{R}^m$, $\lfloor x \rfloor := (\lfloor x \rfloor_1, \dots, \lfloor x \rfloor_m)$.
- Given a set A in some topological space, A° and \bar{A} denote the interior and closure of A respectively.
- R and R_n are “nice” domains in \mathbb{R}^m and \mathbb{Z}^m respectively (see Assumption 2.2 below).
- $h_R : R \rightarrow \mathbb{R}$ is an asymptotic height function (i.e. 1-Lipschitz function).
- $h_{R_n} : R_n \rightarrow \mathbb{Z}$ is a height function (i.e. graph homomorphism).
- $\theta(\delta)$ denotes a function with $\lim_{\delta \downarrow 0} \theta(\delta) = 0$.
- $\theta_{a,b,c}(\delta)$ denotes a function with $\lim_{\delta \downarrow 0} \theta_{a,b,c}(\delta) = 0$, with rate of convergence depending only on the parameters a, b, c .
- For a set A we denote with $|A|$ either the cardinality of A or the Lebesgue measure of A .

2. SETTING AND MAIN RESULTS

In this section we formally describe the model under study, and state the main results that we prove. In describing the model we err on the side of verbosity and explicitness. Some of the notations used are non-standard (such as the θ -notation for asymptotics described in Section 2.4), but these notations allow for relatively concise and (more importantly) precise statements of the results and proofs to follow.

In Section 2.1 we will carefully introduce the basic model, i.e. height functions on “nice” subsets of \mathbb{Z}^m . In Section 2.2 we describe a canonical family of height functions. In Section 2.3 we use these canonical height functions to define the microscopic entropy, then we go on to define the macroscopic entropy and surface tension. In Section 2.4 we introduce our asymptotic notation, as mentioned above. In Section 2.5, the main results of this article are stated.

2.1. Objects of study. Given a graph $\Gamma = (V, E)$, we write “ $v_1 \sim v_2$ ” if two vertices $v_1, v_2 \in V$ are adjacent; i.e. if $\{v_1, v_2\} \in E$. Given two graphs $\Gamma_1 = (V_1, E_1)$ and $\Gamma_2 = (V_2, E_2)$, we recall that a *graph homomorphism* is a function $\varphi : V_1 \rightarrow V_2$ such that whenever vertices $v, v' \in V_1$ are adjacent in Γ_1 , their images $\varphi(v), \varphi(v') \in V_2$ are adjacent in Γ_2 .

In this article, we specialize to the case of graph homomorphisms from certain subgraphs $R_n \subset \mathbb{Z}^m$ to a subgraph of \mathbb{Z} . For $R_n \subset \mathbb{Z}^m$, we write R_n^c for the complement of R_n in \mathbb{Z}^m , and $\partial R_n := \{z \in R_n \mid \exists z' \in R_n^c, z \sim z'\}$ for the (inner) boundary of R_n .

Definition 2.1 (Height functions). *A height function on R_n is a graph homomorphism $h_{R_n} : R_n \rightarrow \mathbb{Z}$ that preserves parity, meaning that for $z = (z_1, \dots, z_m) \in \mathbb{Z}^m$,*

$$h_{R_n}(z) \bmod 2 = z \bmod 2 \left(:= \left[\sum_{i=1}^m z_i \right] \bmod 2 \right).$$

We call a height function $h_{\partial R_n} : \partial R_n \rightarrow \mathbb{Z}$, defined on the boundary of R_n , a boundary height function.

We are interested in sequences of subgraphs $\{R_n : n \in \mathbb{N}\}$ that converge under a scaling limit to a “nice” region $R \subset \mathbb{R}^m$. More specifically, we make the following assumptions.

Assumption 2.2 (Assumptions on domains R and R_n). *We assume that $R \subset \mathbb{R}^m$ is compact and connected and that R is the closure of its interior (sets with the latter property are called regular closed sets; see e.g. [SS95]).*

We assume that $R_n \subset \mathbb{Z}^m$ and $\partial R_n \subset R$ are connected as subgraphs of \mathbb{Z}^m , and (for simplicity) we assume that $\frac{1}{n}R_n \subset R$.

We require that $\frac{1}{n}R_n \rightarrow R$ in the Hausdorff metric; that is, the metric on $\mathcal{P}(\mathbb{R}^m) := \{A \subset \mathbb{R}^m\}$ defined by

$$d_H(A, B) := \left(\sup_{x \in A} \inf_{y \in B} |x - y|_1 \right) \vee \left(\sup_{y \in B} \inf_{x \in A} |x - y|_1 \right). \quad (1)$$

Remark 2.3 (On the choice of norm in (1)). *By equivalence of norms, it does not matter which norm on \mathbb{R}^m is used in (1). Later, we will be interested primarily in the ℓ^1 norm. This is because the ℓ^1 norm is the scaling limit of the graph distance on \mathbb{Z}^m . More precisely, if $x, x' \in \mathbb{R}^m$ and if $z_n, z'_n \in \mathbb{Z}^m$ satisfy $|\frac{1}{n}z_n - x|_1 < \frac{m}{n}$ and $|\frac{1}{n}z'_n - x'|_1 < \frac{m}{n}$, then $\frac{1}{n}d_{\mathbb{Z}^m}(z_n, z'_n) \rightarrow |x - x'|_1$, where $d_{\mathbb{Z}^m}$ denotes the graph distance.*

For example, when R is compact, convex polytope, such as a hypercube or a simplex, the sets $R_n := \{z \in \mathbb{Z}^m \mid \frac{1}{n}z \in R\}$ satisfy Assumption 2.2.

Just as the microscopic domains R_n have a scaling limit, so do the microscopic height functions $h_{R_n} : R_n \rightarrow \mathbb{Z}$.

Definition 2.4 (Asymptotic height functions). *We call a function $h_R : R \rightarrow \mathbb{R}$ an asymptotic height function if h_R is Lipschitz with Lipschitz constant at most 1, with respect to the ℓ^1 -norm on \mathbb{R}^m ; that is, if*

$$\text{Lip}(h_R) := \sup_{x \neq y \in R} \frac{|h_R(x) - h_R(y)|}{|x - y|_1} \leq 1. \quad (2)$$

Likewise, if $h_{\partial R} : \partial R \rightarrow \mathbb{R}$ is 1-Lipschitz (with respect to the ℓ^1 -norm), we call $h_{\partial R}$ an asymptotic boundary height function.

We assume that $h_{\partial R_n} : \partial R_n \rightarrow \mathbb{Z}$ are boundary height functions that converge (after rescaling) to an asymptotic boundary height function $h_{\partial R} : \partial R \rightarrow \mathbb{R}$ in following sense: for each n , let $d_n = d_H(\frac{1}{n}R_n, R)$. Then, we say $\frac{1}{n}h_{\partial R_n} \rightarrow h_{\partial R}$ if and only if

$$\lim_{n \rightarrow \infty} \sup_{z \in \partial R_n} \sup_{\substack{x \in \partial R \\ |x - \frac{1}{n}z|_1 \leq d_n}} \left| \frac{1}{n}h_{\partial R_n}(z) - h_{\partial R}(x) \right| = 0.$$

Now, we define a few families of height functions and asymptotic height functions. The sets of height functions from Definition 2.5 below appear frequently in entropies, and the sets of asymptotic height functions from Definition 2.6 are important for the statement of the variational principle (Theorem 2.16).

Definition 2.5 (Sets of height functions). *Let R_n be a microscopic domain as above, let $h_{R_n} : R_n \rightarrow \mathbb{Z}$ be a boundary height function, and let $\delta > 0$. We define:*

$$\begin{aligned} M(R_n) &:= \{h_{R_n} : R_n \rightarrow \mathbb{R} \mid h_{R_n} \text{ is a height function}\} \\ M(R_n, h_{\partial R_n}) &:= \{h_{R_n} \in M(R_n) \mid h_{R_n}|_{\partial R_n} = h_{\partial R_n}\} \\ M(R_n, h_{\partial R_n}, \delta) &:= \{h_{R_n} \in M(R_n) \mid \sup_{z \in \partial R_n} |h_{R_n}(z) - h_{\partial R_n}(z)| < \delta n\}. \\ B(R_n, h_R, \delta) &:= \{h_{R_n} \in M(R_n) \mid \sup_{z \in R_n} |h_R(\frac{1}{n}z) - \frac{1}{n}h_{R_n}(z)| < \delta\}. \end{aligned}$$

In the last definition, the expression “ $h_R(\frac{1}{n}z)$ ” makes sense because of the assumption that $\frac{1}{n}R_n \subset R$ in Assumption 2.2.

Definition 2.6 (Sets of asymptotic height functions). *Let $R \subset \mathbb{R}^m$ be a domain satisfying Assumption 2.2, let $h_{\partial R} : \partial R \rightarrow \mathbb{R}$ be an asymptotic boundary height function, and let $\delta > 0$. We define:*

$$\begin{aligned} M(R) &:= \{h_R : R \rightarrow \mathbb{R} \mid h_R \text{ is an asymptotic height function}\} \\ M(R, h_{\partial R}) &:= \{h_R : R \rightarrow \mathbb{R} \mid h_R|_{\partial R} = h_{\partial R}\} \\ M(R, h_{\partial R}, \delta) &:= \{h_R : R \rightarrow \mathbb{R} \mid \forall x \in \partial R, |h_R(x) - h_{\partial R}(x)| \leq \delta\}. \end{aligned}$$

2.2. Affine height functions. Affine height functions play an important role in defining and studying the entropy of our model. For an asymptotic height function $h_R : R \rightarrow \mathbb{R}$, we mean by “affine” the usual

property: there exist $s \in [-1, 1]^m$ and $b \in \mathbb{R}$ such that $h_R(x) = s \cdot x + b$. The bounds on s ensure that h_R satisfies the Lipschitz property (2), so all such functions are indeed asymptotic height functions as per Definition 2.4.

On the microscopic domains R_n , we consider best-possible approximations to affine functions. Fix $s \in [-1, 1]^m$ and $b \in \mathbb{R}$. At a lattice point $z \in \mathbb{Z}^m$, we define $h_{R_n}^{s \cdot x + b}(z)$ to be $s \cdot z + b$, rounded to the nearest integer of correct parity (see Figure 4). In the rest of this subsection, we formalize this definition, verify that it actually does define a height function, and check that it is consistent.

Let us introduce an auxiliary notation that is used only in this subsection. Given a point $z = (z_1, \dots, z_m) \in \mathbb{Z}^m$, we say z has even or odd parity as $(\sum_{i=1}^m z_i) \in \mathbb{Z}$ has even or odd parity respectively, and we write $z \bmod 2$ for the parity of z .

Given $z \in \mathbb{Z}^m$ and $y \in \mathbb{R}$, we write $[y]_{z \bmod 2}$ for the closest integer to y that has parity $z \bmod 2$. In case of a tie, i.e. if y is an integer that has opposite parity to z , we arbitrarily choose to “round up” and set $[y]_{z \bmod 2} = y + 1 \in \mathbb{Z}$.

For example, let $z = (1, 2, 3) \in \mathbb{Z}^3$ and $z' = (4, -6, 7)$. Then z is an even point and z' is an odd point. So:

$$\begin{aligned} [5.4]_{z \bmod 2} &= 6, & [-3]_{z \bmod 2} &= -2, \\ [5.4]_{z' \bmod 2} &= 5, & [-3]_{z' \bmod 2} &= -3. \end{aligned}$$

Now, given $s \in [-1, 1]^m$ and $b \in \mathbb{R}$, we define the affine height functions $h_{R_n}^{s \cdot x + b}$ by

$$h_{R_n}^{s \cdot x + b}(z) := [s \cdot z + b]_{z \bmod 2}.$$

Note that the symbol x in the superscript of $h_{R_n}^{s \cdot x + b}$ is merely formal; “ $s \cdot x + b$ ” should be read as “the function mapping x to $s \cdot x + b$ ”. Moreover, the choice of domain R_n in the subscript does not affect the values of $h_{R_n}^{s \cdot x + b}$ at any point; for any sets $A_n, B_n \subseteq \mathbb{Z}^m$ and any point $z \in A_n \cap B_n$, one has $h_{A_n}^{s \cdot x + b}(z) = h_{B_n}^{s \cdot x + b}(z)$. An example of a function $h_{R_n}^{s \cdot x + b}$ is provided in Figure 4.

From the definition above, it is not clear that $h_{R_n}^{s \cdot x + b}$ are height functions. This is the content of Lemma 2.7.

Lemma 2.7. *Let $s \in [-1, 1]^m$ and $b \in \mathbb{R}$. For any adjacent points $z \sim z' \in \mathbb{Z}^m$, the values $h_{R_n}^{s \cdot x + b}(z)$ and $h_{R_n}^{s \cdot x + b}(z')$ differ by exactly 1.*




FIGURE 4. An affine height function $h_{R_n}^{s \cdot x + b}$ and the corresponding continuous affine function $x \mapsto s \cdot x + b$. Here $s = \frac{1}{2}$ and $b = \frac{1}{2}$.

Proof. From the definition of $h_{R_n}^{s \cdot x + b}$, we note two inequalities:

$$|h_{R_n}^{s \cdot x}(z) - (s \cdot z + b)| \leq 1, \quad (3)$$

and

$$|h_{R_n}^{s \cdot x + b}(z') - (s \cdot z' + b)| \leq 1. \quad (4)$$

Additionally, since $s \in [-1, 1]^m$, we have

$$|(s \cdot z + b) - (s \cdot z' + b)| \leq 1.$$

By the triangle inequality, $|h_{R_n}^{s \cdot x + b}(z) - h_{R_n}^{s \cdot x + b}(z')| \leq 3$. We shall show that equality cannot hold. Since the difference $h_{R_n}^{s \cdot x + b}(z) - h_{R_n}^{s \cdot x + b}(z')$ is obviously an odd integer, it will follow that the difference is ± 1 .

Suppose towards a contradiction that

$$|h_{R_n}^{s \cdot x + b}(z) - h_{R_n}^{s \cdot x + b}(z')| = 3.$$

Then (3) and (4) must be equalities. From the definition of $[\cdot]_{z \bmod 2}$, necessarily then $s \cdot z + b$ is an integer with parity opposite that of z , and so

$$h_{R_n}^{s \cdot x + b}(z) = (s \cdot z + b) + 1.$$

Likewise

$$h_{R_n}^{s \cdot x + b}(z') = (s \cdot z' + b) + 1.$$

But then

$$|h_{R_n}^{s \cdot x + b}(z) - h_{R_n}^{s \cdot x + b}(z')| = |(s \cdot z + b + 1) - (s \cdot z' + b + 1)| \leq 1.$$

□

We end this section with the following lemma. The conclusion (6) is exactly what is needed later to apply the Kirszbraun theorem (Theorem 3.1):

Lemma 2.8 (Inequality for $h_{R_n}^{s \cdot x + b}$). *Let $s, s' \in [-1, 1]^m$, $b, b' \in \mathbb{R}$, and $z, z' \in \mathbb{Z}^m$. If*

$$|(s \cdot z + b) - (s' \cdot z' + b')| \leq |z - z'|_1, \quad (5)$$

then

$$|h_{\{z\}}^{s \cdot x + b}(z) - h_{\{z'\}}^{s' \cdot x + b'}(z')| \leq |z - z'|_1. \quad (6)$$

Proof. The proof is similar to that of Lemma 2.7. By the triangle inequality and (5),

$$\begin{aligned} & |h_{\{z\}}^{s \cdot x + b}(z) - h_{\{z'\}}^{s' \cdot x + b'}(z')| \\ & \leq |h_{\{z\}}^{s \cdot x + b}(z) - (s \cdot z + b)| + |(s \cdot z + b) - (s' \cdot z' + b')| \\ & \quad + |(s' \cdot z' + b') - h_{\{z'\}}^{s' \cdot x + b'}(z')| \\ & \leq |z - z'|_1 + 2. \end{aligned} \quad (7)$$

Since $h_{\{z\}}^{s \cdot x + b}(z)$, $h_{\{z'\}}^{s' \cdot x + b'}(z')$ and $|z - z'|_1$ are all integers,

$$|h_{\{z\}}^{s \cdot x + b}(z) - h_{\{z'\}}^{s' \cdot x + b'}(z')| - |z - z'|_1 \in \{\dots, -2, -1, 0, 1, 2\}. \quad (8)$$

We want to prove that the left-hand side of (8) is ≤ 0 . By parity considerations it must be even, and we need only prove it is $\neq 2$. Assume for a contradiction that the left-hand side of (8) equals 2. Then equality holds in (7), and in particular

$$|h_{\{z\}}^{s \cdot x + b}(z) - (s \cdot z + b)| = 1 \quad \text{and} \quad |h_{\{z'\}}^{s' \cdot x + b'}(z') - (s' \cdot z' + b')| = 1.$$

As in the proof of Lemma 2.7, this implies that

$$h_{\{z\}}^{s \cdot x + b}(z) = (s \cdot z + b) + 1 \quad \text{and} \quad h_{\{z'\}}^{s' \cdot x + b'}(z') = (s' \cdot z' + b') + 1.$$

Therefore

$$|h_{\{z\}}^{s \cdot x + b}(z) - h_{\{z'\}}^{s' \cdot x + b'}(z')| = |(s \cdot z + b) - (s' \cdot z' + b')| \leq |z - z'|_1.$$

This is the desired contradiction, which completes the proof. \square

2.3. Entropies and surface tensions. In this section we make three more definitions needed for our statement of the main results. First, we define the microscopic entropy of a set of height functions. More precisely, this is the Shannon entropy of the uniform distribution over a finite set of height functions, normalized by the size of their common domain, and negated. (The negative convention is chosen so that the surface tension $\text{ent}(s)$, defined later, is convex rather than concave.)

The microscopic entropy is essentially the same as the specific free energy of [She05].

Definition 2.9 (Microscopic entropy). *Given a finite, non-empty set of height functions $A \subset M(R_n)$, we define the microscopic entropy*

$$\text{Ent}_{R_n}(A) := -\frac{1}{|R_n|} \ln |A|. \quad (9)$$

We observe that the microscopic entropy is translation invariant:

Observation 2.10 (Translation invariance). *Let $h_{\partial R_n} \in M(\partial R_n)$, $h_R \in M(R)$, $\delta > 0$, and $c \in \mathbb{R}$. Then:*

$$\begin{aligned} \text{Ent}_{R_n}(M(R_n, h_{\partial R_n} + c)) &= \text{Ent}_{R_n}(M(R_n, h_{\partial R_n})), \\ \text{Ent}_{R_n}(M(R_n, h_{\partial R_n} + c, \delta)) &= \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)), \text{ and} \\ \text{Ent}_{R_n}(B(R_n, h_R + c, \delta)) &= \text{Ent}_{R_n}(B(R_n, h_R, \delta)), \end{aligned}$$

where $h_{R_n} + c$ is the height function defined by $(h_{R_n} + c)(z) = h_{R_n}(z) + c$ for $z \in R_n$, and likewise $(h_R + c)(x) = h_R(x) + c$ for $x \in R$.

All of these sets, except for $M(R_n)$, are finite, because of the constraints they impose on h_{R_n} and the Lipschitz property of h_{R_n} . In fact, we can say more. Let us count $M(R_n, h_{\partial R_n})$, for some boundary height function $h_{\partial R_n} \in M(\partial R_n)$. The values of $h_R \in M(R_n, h_{\partial R_n})$ are fixed on ∂R_n , and for each of the $\leq |R_n|$ points x in the interior of R_n , there are at most 2 admissible values for $h_R(x)$. Therefore $|M(R_n, h_{\partial R_n})| \leq 2^{|R_n|}$. Similar logic holds for $M(R_n, h_{\partial R_n}, \delta)$ and $B(R_n, h_R, \delta)$. This leads to the following observation:

Observation 2.11 (Boundedness of Ent). *Let $h_{\partial R_n} \in M(\partial R_n)$, $h_R \in M(R)$, and $\delta > 0$. Then:*

$$\begin{aligned} \text{Ent}_{R_n}(M(R_n, h_{\partial R_n})) &\geq -\ln 2, \\ \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) &\geq -\ln 2 - \frac{\ln[2\delta n]}{|R_n|} \\ &= -\ln 2 - \delta O\left(\frac{1}{n^{m-1}}\right), \text{ and} \\ \text{Ent}_{R_n}(B(R_n, h_R, \delta)) &\geq -\ln 2 - \frac{\ln[2\delta n]}{|R_n|} \\ &= -\ln 2 - \delta O\left(\frac{1}{n^{m-1}}\right). \end{aligned}$$

Next, we define the local surface tension. There are in general many equivalent definitions of surface tension (see for example Chapter 6 of [She05]). The following definition is easiest to work with for our purposes.

Definition 2.12 (Local surface tension). *For $s \in [-1, 1]^m$, the local surface tension $\text{ent}(s)$ is defined to be the limit*

$$\text{ent}(s) := \lim_{n \rightarrow \infty} \text{ent}_n(s), \quad (10)$$

where $\text{ent}_n(s)$ is defined as

$$\text{ent}_n(s) := \text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n}^{s \cdot x + 0})) \quad (11)$$

and where $Q_n = [0, n]^m \cap \mathbb{Z}^m$ is the discrete hypercube of side length n .

The limit (10) exists by standard subadditivity arguments; we refer the interested reader to e.g. [Dur10]. In fact, by translation invariance (see Observation 2.10), we may replace $h_{\partial Q_n}^{s \cdot x + 0}$ by $h_{\partial Q_n}^{s \cdot x + b}$ in (11), for any $b \in \mathbb{R}$. Additionally, boundedness passes through the limit in (10). Therefore:

Observation 2.13 (Boundedness of $\text{ent}(s)$). *For any $s \in [-1, 1]^m$,*

$$-\ln 2 \leq \text{ent}(s) \leq 0.$$

Let us now define the macroscopic entropy.

Definition 2.14 (Macroscopic entropy). *Let $h_R : R \rightarrow \mathbb{R}$ denote an asymptotic height function. The macroscopic entropy $\text{Ent}_R(h_R)$ is defined as*

$$\text{Ent}_R(h_R) := \int_R \text{ent}(\nabla h(x)) dx.$$

2.4. Asymptotic notation. In this section we introduce a notation for asymptotic error. Compared to the Landau big- O notation, our θ -notation abstracts away the rate of convergence of the error, but makes explicit the dependence on parameters. For this purpose we write $\theta_\alpha(\delta)$ for a family of unspecified functions, parameterized by a symbol α , such that $\theta_\alpha(\delta) \rightarrow 0$ at a rate depending on the value of the parameter α . That is, for any $\varepsilon > 0$ and any admissible parameter value α , there exists $\delta_0 = \delta_0(\alpha) > 0$ such that $0 < \delta < \delta_0$ implies $\theta_\alpha(\delta) < \varepsilon$.

Extending the above notation, we frequently replace the single parameter α by a list of parameters $\alpha, \beta, \gamma, \dots$. For example, we might write an identity like

$$\begin{aligned} \min_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) &= \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) \\ &\quad + \theta_{m, R, h_{\partial R}, R_n, h_{\partial R_n}}(\delta) \\ &\quad + \theta_{m, R, h_{\partial R}, R_n, h_{\partial R_n}, \delta}(\tfrac{1}{n}). \end{aligned} \quad (12)$$

The identity states that the two entropy terms on the first line differ by a small amount; the difference vanishes as δ and $\frac{1}{n}$ go to zero, and

the rate of convergence depends on several parameters. The “ $\theta(\delta)$ ” term depends on the parameters from the setting, namely the ambient dimension m , the region R , the height function h_R of interest, and the corresponding discrete objects R_n and h_{R_n} . The “ $\theta(\frac{1}{n})$ ” term depends on these parameters along with the value of δ . We find that listing out the setting parameters m , R , h_R , R_n , and h_{R_n} makes the expression harder to read. So for the rest of the article we suppress these parameters from the subscripts of θ terms. Under this convention (12) becomes:

$$\min_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) = \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) + \theta(\delta) + \theta_\delta(\frac{1}{n}).$$

As mentioned above, the advantage of our θ notation is that it abstracts away the exact rates of convergence, but leaves explicit the dependencies between parameters. For example, suppose we want to make the error in approximation in (12) to be less than ε . We should first choose δ so that (say) $\theta(\delta) < \frac{1}{2}\varepsilon$, then choose n depending on δ (and on the suppressed parameters m , R , etc.) so that $\theta_\delta(\frac{1}{n}) < \frac{1}{2}\varepsilon$.

2.5. Main results. The main results of this article are the profile theorem, the variational principle, and the large deviations principle:

Theorem 2.15 (Profile theorem). *Under the setting explained in Section 2, for any $h_R \in M(R)$, $\delta > 0$, and $n \in \mathbb{N}$,*

$$\text{Ent}_R(h_R) = \text{Ent}_{R_n}(B(R_n, h_R, \delta)) + \theta_{h_R}(\delta) + \theta_{h_R, \delta}(\frac{1}{n}). \quad (13)$$

The second main result is the variational principle:

Theorem 2.16 (Variational principle). *Under the setting explained in Section 2,*

$$\inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) = \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) + \theta(\delta) + \theta_\delta(\frac{1}{n})$$

for any $\delta > 0$ and $n \in \mathbb{N}$.

Finally, we prove a large deviations principle for the model. We adopt the conventions of large deviations (see for example [DZ09, RAS15]).

Theorem 2.17 (Large deviations principle). *Consider the Polish space $M(R)$ of asymptotic height functions (i.e. Lipschitz functions with Lipschitz constant ≤ 1), endowed with the topology of uniform convergence (induced by the supremum norm).*

For $\delta > 0$ and $n \in \mathbb{N}$, define a probability measure $\mu_{\delta, n}$ on $M(R)$ by

$$\mu_{\delta, n}(A) := \frac{1}{|M(R_n, h_{\partial R_n}, \delta)|} \left| \{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\} \right|,$$

where \tilde{h}_{R_n} is the Lipschitz function given by rescaling and interpolating h_{R_n} so as to make it an asymptotic height function, i.e. for $z \in R_n$, $\tilde{h}_{R_n}(\frac{1}{n}z) = \frac{1}{n}h_{R_n}(z)$.

The measures $(\mu_{\delta,n})_{\delta>0, n \in \mathbb{N}}$ satisfy a large deviations principle with speed $r_{\delta,n} := |R_n|$ and tight rate function $I : M(R) \rightarrow [0, \infty]$ given by

$$I(h_R) := \begin{cases} \text{Ent}_R(h_R) - E & \text{if } h_R|_{\partial R} = h_{\partial R}, \\ \infty & \text{otherwise.} \end{cases}$$

where $E := \inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R)$. More precisely, for any Borel subset $A \subset M(R)$,

$$-\inf_{h_R \in A^\circ} I(h_R) \leq \frac{1}{r_{\delta,n}} \log \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \mu_{\delta,n}(A) \quad (14)$$

and

$$\frac{1}{r_{\delta,n}} \log \overline{\lim}_{\delta \rightarrow 0} \overline{\lim}_{n \rightarrow \infty} \mu_{\delta,n}(A) \leq -\inf_{h_R \in \overline{A}} I(h_R), \quad (15)$$

where as usual $\underline{\lim}$ and $\overline{\lim}$ denote the limit inferior and superior respectively.

Remark 2.18. It is straightforward to reduce the double limits in Theorem 2.17 to (single) sequential limits, which are more common in large deviations theory. For example, one may choose any sequences $(\delta_k)_{k \in \mathbb{N}}, (\varepsilon_k)_{k \in \mathbb{N}}$ such that $\delta_k \rightarrow 0$ and $\varepsilon_k \rightarrow 0$ as $k \rightarrow \infty$. Then, choose n_k large enough that

$$\underline{\lim}_{n \rightarrow \infty} \mu_{\delta_k, n}(A) - \varepsilon_k \leq \mu_{\delta_k, n_k}(A) \leq \overline{\lim}_{n \rightarrow \infty} \mu_{\delta_k, n}(A) + \varepsilon_k,$$

and define $\mu_k := \mu_{\delta_k, n_k}$ and $r_k := r_{\delta_k, n_k}$. Then

$$-\inf_{h_R \in A^\circ} I(h_R) \leq \underline{\lim}_{k \rightarrow \infty} \frac{1}{r_k} \log \mu_k(A) \leq \overline{\lim}_{k \rightarrow \infty} \frac{1}{r_k} \log \mu_k(A) \leq -\inf_{h_R \in \overline{A}} I(h_R).$$

For the study of limit shapes, it is useful to prove two additional results: existence and uniqueness of the minimizer of the rate function from the large deviations principle, i.e. there exists a unique $h_R^{\min} \in M(R, h_{\partial R})$ such that

$$I(h_R^{\min}) = \inf_{h_R \in M(R, h_{\partial R})} I(h_R).$$

Indeed, this holds for the simple model studied in the current article. See for example [She05] for proofs and discussion of these results. Even in more subtle models, the existence of the minimizer is often easy to show: the proof is standard as long as the local surface tension is convex and bounded below. To show uniqueness is harder. Uniqueness of the minimizer may be proved using strict convexity of the local surface

tension see for example Proposition 4.5 of [DSS08]. We do not prove these results in the current article, but rather focus on the variational principle and large deviations principle.

Once existence and uniqueness of the minimizer are established (or in the language of the current model, once it is known that the macroscopic entropy functional admits a unique minimizing height function), one can explain the appearance of a limit shape in the following way. The set of asymptotic height functions that lie within distance ε of this minimizer is an open ball in the space $M(R)$. By applying the large deviations principle on the set-theoretical complement, one sees that the percentage of microscopic height functions in $M(R_n, h_{\partial R_n}, \delta)$ that do not lie ε -close to the minimizer decays exponentially. In other words, with high probability a randomly chosen height function is close to the minimizer, and therefore the minimizer is the limit shape.

3. OUTLINE AND DISCUSSION OF PROOF OF MAIN RESULTS

In this section we briefly outline the proof of the main results and summarize some key ideas. Then we analyze the ingredients in the proof with an eye toward extending the proof to other random surface models.

In Section 4, we provide auxiliary results including basic properties of the local surface tension and microscopic entropy. A central ingredient of the overall argument is discussed in Section 5. There we prove the profile theorem in the special case of piecewise affine asymptotic height functions. In Section 6, we extend the profile theorem to general asymptotic height functions by an approximation argument, yielding the first main result Theorem 2.15. In Section 7, we use the profile theorem and a compactness argument to prove the variational principle Theorem 2.16. The argument is based on compactness of the space of asymptotic height functions with fixed boundary values $M(R, h_{\partial R})$. Finally in Section 8, we extend the proof of the variational principle in order to prove the large deviations principle Theorem 2.17.

As one can see from this outline, the main idea of the argument is to reduce the proof of the profile theorem from general domains and asymptotic height functions to simpler domains and asymptotic height functions by an approximation argument. This means that the left-hand side of (13), i.e. the macroscopic entropy, and the right-hand side, i.e. the microscopic entropy, must both be robust with respect to approximations.

The macroscopic entropy is robust because $\text{ent}(s)$ is bounded and uniformly continuous, and Lipschitz functions can be approximated

very well by linear interpolations on a simplex domain (cf. the simplicial Rademacher theorem Lemma 6.1). This approximation lemma was formulated for two dimensions in [CKP01]. The result is interesting in its own right and for the convenience of the reader we state and prove it for arbitrary dimension in Section 6.

The microscopic entropy is robust under approximations because the microscopic surface tension is very robust: even with fluctuations in the boundary values and the geometry of the boundary, one still gets the same limit in (10). This result is proved in Section 4, using a Kirszbraum theorem for graph homomorphisms stated below. This theorem gives conditions under which a graph homomorphism can be extended from a smaller domain to a larger domain. This is a discrete analogue to the classical result [Kir34], which deals with Lipschitz functions defined on subsets of \mathbb{R}^d . We also note that more general forms of the Kirszbraum theorem for graph homomorphisms are known, e.g. [CPT18].

Theorem 3.1 (Kirszbraum theorem for height functions). *Let Λ be a connected region of \mathbb{Z}^m , let $S \subset \Lambda$, and let $\bar{h} : S \rightarrow \mathbb{Z}$ be a graph homomorphism that preserves parity. There exists a graph homomorphism $h : \Lambda \rightarrow \mathbb{Z}$ such that $h = \bar{h}$ on S if and only if for all x, y in S ,*

$$|\bar{h}(x) - \bar{h}(y)| \leq |x - y|_1, \quad (16)$$

where $|x - y|_1$ is the ℓ^1 -norm in \mathbb{Z}^m .

Remark 3.2. *The parity condition is necessary in general; consider for example the function \bar{h} defined on $\{0, 2\} \subset \mathbb{Z}$ by $\bar{h}(0) = 0$, $\bar{h}(2) = 1$. The parity condition in Theorem 3.1 is the reason for the parity condition in Definition 2.1.*

Two of the authors gave a proof of a more general version of this theorem in [MT16] (see Theorem 4.1). The proof is restated below for the reader's convenience. This proof is also simplified by only addressing the model from this article, where the height functions take values in \mathbb{Z} rather than in a d -regular tree.

Proof of Theorem 3.1. Obviously if an extension h of \bar{h} exists, then \bar{h} satisfies (16). So, suppose instead that (16), and let us prove that an extension h exists. For $y \in \Lambda$, set

$$h(y) := \max\{\bar{h}(x) - |x - y|_1 \mid x \in S\}. \quad (17)$$

We must check two things: first, that $h(y) = \bar{h}(y)$ when $y \in S$, and second, that $|h(y) - h(\tilde{y})| = 1$ when $y \sim \tilde{y}$ are adjacent points in Λ .

To prove that $h|_S = \bar{h}$, let $y \in S$ and consider any point $x \in S$. By the Lipschitz property of \bar{h} ,

$$\bar{h}(x) - \bar{h}(y) \leq |\bar{h}(x) - \bar{h}(y)| \leq |x - y|_1,$$

so $\bar{h}(x) - |x - y|_1 \leq \bar{h}(y)$. Therefore the maximum in (17) is attained when $x = y$, so $h(y) = \bar{h}(y) + |y - y|_1 = \bar{h}(y)$.

To prove that h is a graph homomorphism, let $y \sim \tilde{y}$ be adjacent points in Λ , and let x, \tilde{x} be points in S that attain the maximum in (17) for y, \tilde{y} respectively, i.e. $h(y) = \bar{h}(x) - |x - y|_1$ and $h(\tilde{y}) = \bar{h}(\tilde{x}) - |\tilde{x} - \tilde{y}|_1$. Then

$$\begin{aligned} h(y) &= \max\{\bar{h}(z) + |z - y|_1 \mid z \in S\} \\ &\geq \bar{h}(\tilde{x}) - |\tilde{x} - y|_1 \\ &\geq \bar{h}(\tilde{x}) - |\tilde{x} - \tilde{y}|_1 - 1 \\ &= h(\tilde{y}) - 1, \end{aligned}$$

and likewise $h(\tilde{y}) \geq h(y) - 1$.

For every $x \in S$, the map $y \mapsto \bar{h}(x) + |x - y|_1$ preserves parity (recall the assumption that \bar{h} preserves parity), and therefore so does h . So h is a parity-preserving map such that $|h(y) - h(\tilde{y})| \leq 1$ whenever y and \tilde{y} are neighbors. This proves that h is a graph homomorphism. \square

Now, we describe further how to prove the central theorem of this article, i.e. the profile theorem in the special case of piecewise affine height functions. We derive the desired asymptotic equality by showing two inequalities. One direction of the inequality arises by overcounting the number of height functions that are close to the piecewise affine height profile; the opposite direction arises by undercounting the same set. In both directions, we subdivide the region into small blocks, so that we can compare the entropy on each block to the local surface tension (see Definition 2.12 and Figure 9). To overcount, we consider all choices of boundary values on the boundaries of the blocks, and for each boundary value function we count all possible extensions into the interior of the blocks. To undercount we have to use much smaller blocks, with boundary values fixed to match the desired affine function exactly (after rescaling, and up to rounding). The details of the proof are given in Section 5. The more difficult part of the proof is the overcounting argument, which relies on robustness of the microscopic entropy. We expect this to be a major source of difficulty when adapting our methods to other models.

As one can see, the framework of this argument is quite general and it can be adapted to more complicated models and settings. For example, the model of graph homomorphism into the infinite d -regular tree, studied by some of the current authors in [MT16], is amenable to this approach. Additionally, the authors have applied the current strategy to \mathbb{Z} -valued homomorphisms sampled according to a random environment. This means that the underlying combinatorial model

is the same as in the current article, but in the definition of the microscopic entropy, the (uniform) counting measure on $M(R_n, h_{\partial R_n})$ is replaced by a randomly perturbed measure. The conclusion is a homogenized variational principle, meaning that the microscopic entropy $\text{Ent}_n(M(R_n, h_{\partial R_n}))$, now a random variable depending on the realization of the environment, converges in probability to the minimum of the macroscopic entropy, which is still a deterministic quantity. Furthermore, we hope the method applies to other height function models, such as domino tilings (as studied in e.g. [CKP01]), and perhaps even more general tilings (as in e.g. [She01, Thu90]).

4. MICROSCOPIC ENTROPY AND SURFACE TENSION

In this section, we prove basic properties of the microscopic entropy and local surface tension. More precisely, we prove that $\text{ent}(s)$ is continuous (see Lemma 4.3), that $\text{ent}_n(s) \rightarrow \text{ent}(s)$ uniformly (see Lemma 4.4), and that Ent_{R_n} is robust under small changes to boundary values (see Lemma 4.5).

All three of these proofs split into two cases: values of the slope s that are close to 1 (that is, such that $|s|_\infty \geq 1 - \varepsilon$), where there are comparatively few possible states because of the steep slope; and slopes away from 1 (i.e. $|s|_\infty \leq 1 - \varepsilon$), where we can make arguments based on extending height functions from one domain to another via the Kirszbraun theorem (Theorem 3.1).

The first result we state is about the microscopic entropy for slopes close to 1. This lemma is used in the remainder of the section to handle the case of s close to 1.

Lemma 4.1 (Microscopic entropy for slopes near 1). *Let $\delta > 0$, let $s \in [-1, 1]^m$ with $|s|_\infty > 1 - \delta$, and let $n \in \mathbb{N}$. Consider any boundary height function $h_{\partial Q_n} \in M(\partial Q_n)$ such that*

$$\sup_{z \in \partial Q_n} |h_{\partial Q_n}(z) - h_{\partial Q_n}^s(z)| \leq \delta n. \quad (18)$$

Then,

$$\text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n})) = \theta(\delta) + \theta_\delta\left(\frac{1}{n}\right).$$

Proof. First, consider the one-dimensional case, i.e. $m = 1$. Then the problem reduces to a simple calculation. The main idea is that the large slope s forces a height function $h_{Q_n} \in M(Q_n, h_{\partial Q_n})$ to closely follow a line of slope ± 1 . By counting the number of deviating edges we overestimate the number of height functions.

Indeed, we assume without loss of generality that $\mathbb{R}^1 \ni s > 1 - \delta$ (the case $s < -(1 - \delta)$ is symmetric). We want to count height functions




FIGURE 5. A one-dimensional height function with slope $s > 1 - \delta$. Because the slope is close to 1, there cannot be many edges along which $h(x)$ decreases.

$h_{Q_n} \in M(Q_n, h_{\partial Q_n})$. The line graph $Q_n = \{-n, -(n-1), \dots, n-1, n\}$ has $2n$ edges; let us write k for the number of edges on which h_{Q_n} decreases (see Figure 5). Then the height difference $h_{Q_n}(n) - h_{Q_n}(-n)$ is exactly $(2n - k) - k$, which we simplify to $2(n - k)$. By (18), we have (ignoring rounding errors)

$$\begin{aligned} h_{Q_n}(n) - h_{Q_n}(-n) &= h_{\partial Q_n}^s(n) - h_{\partial Q_n}^s(-n) \\ &= sn - s(-n) \\ &> 2(1 - \delta)n. \end{aligned}$$

Therefore, $2(n - k) > 2(1 - \delta)n$, so $k \leq \delta n$. It follows that

$$|M(Q_n, h_{\partial Q_n})| \leq \binom{2n}{k} \leq \binom{2n}{\lceil \delta n \rceil},$$

and the limit

$$\lim_{n \rightarrow \infty} \frac{1}{2n+1} \log \binom{2n}{\lceil \delta n \rceil} = -\delta \log \delta - (1 - \delta) \log(1 - \delta) = \theta(\delta)$$

is an easy calculation using Stirling's formula.

For higher dimensions, we reduce to the one-dimensional case by treating the hypercube $\{-n, \dots, n\}^m$ as the union of $(2n+1)^{m-1}$ independent lines. In so doing we overestimate $|M(Q_n, h_{\partial Q_n})|$, because we relax the graph homomorphism condition between lines. Thus

$$|M(Q_n, h_{\partial Q_n})| \leq \binom{2n}{\lceil \delta n \rceil}^{(2n+1)^{m-1}}.$$

Taking a logarithm and dividing by $-|Q_n| = -(2n+1)^m$ yields

$$\text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n})) \geq -\frac{1}{2n+1} \log \left(\frac{2n}{\lceil \delta n \rceil} \right),$$

which completes the proof (and in particular shows why the θ error terms do not depend on the dimension m). \square

While Lemma 4.1 deals with slopes s close to 1, a different approach is needed for slopes away from 1. We use Theorem 3.1, which is a Kirszbraum theorem for graph homomorphisms. It gives a simple criterion for when a height function can be extended to larger domain. Lemma 4.2 applies the Kirszbraum theorem to derive entropy estimates. In particular, for two box sizes $n < \hat{n}$, the lemma compares $\text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n}))$ and $\text{Ent}_{Q_{\hat{n}}}(M(Q_{\hat{n}}, h_{\partial Q_{\hat{n}}}))$. The key idea is that any height function on the smaller box Q_n can be extended to a height function on $Q_{\hat{n}}$, respecting the boundary data $h_{\partial Q_{\hat{n}}}$. Therefore (up to vanishing error terms), $\text{Ent}_{Q_{\hat{n}}}(M(Q_{\hat{n}}, h_{\partial Q_{\hat{n}}})) \leq \text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n}))$.

The extension requires that the boundary data $h_{\partial Q_n}$ and $h_{\partial Q_{\hat{n}}}$ be sufficiently similar. In particular, we will assume that both boundary height functions are close to linear height functions, with slopes s and \hat{s} respectively. The parameter ε quantifies how close $h_{\partial Q_n}$ and $h_{\partial Q_{\hat{n}}}$ are to their respective linear height functions.

We also require that the slopes s and \hat{s} be close to each other, which is obviously necessary to apply the Kirszbraum theorem in our setting. Finally, we require that the two boxes sizes n and \hat{n} be not too different. In particular, we take $\hat{n} = (1 + \delta)n$, where δ is a second approximation parameter. δ also shows up in a few other bounds, and in the conclusion of the lemma as a $\theta_m(\delta)$ error term.

This is not the simplest lemma of its kind that we could state, nor is it the most general. We choose to state these conditions because they are sufficient for our applications in this section. Moreover, they are necessary in the sense that simplifying any condition, e.g. by using only a single slope s rather than two slopes, or by using linear boundary height functions without than allowing ε fluctuations, would not suffice for our purposes.

Lemma 4.2 (Entropy estimates from the Kirszbraum theorem). *Let $\delta \in (0, \frac{1}{3})$, $\varepsilon \in (0, \frac{\delta^2}{2+\delta}]$, $n, \hat{n} \in \mathbb{N}$, $s, \hat{s} \in [-1, 1]^m$, $h_{\partial Q_n} \in M(\partial Q_n)$, and $h_{\partial Q_{\hat{n}}} \in M(\partial Q_{\hat{n}})$ satisfy:*

- $\hat{n} = \lceil (1 + \delta)n \rceil$ or $\hat{n} = \lceil (1 + \delta)n \rceil + 1$,
- $|s|_{\infty} \leq 1 - 3\delta$ and $|s - \hat{s}|_{\infty} \leq \frac{\delta^2}{1+\delta}$,
- $\max_{z \in \partial Q_n} |h_{\partial Q_n}(z) - s \cdot z| \leq \varepsilon n$, and




FIGURE 6. Domains for Lemma 4.2. The smaller domain Q_n is a hypercube of side length $2n + 1$, and the larger domain $Q_{\hat{n}}$ has side length $2\hat{n} + 1$, where $\hat{n} = (1 + \delta)n$.

- $\max_{z \in \partial Q_{\hat{n}}} |h_{\partial Q_{\hat{n}}}(z) - \hat{s} \cdot z| \leq \varepsilon \hat{n}.$

See Figure 6 for a partial illustration. Then:

$$\text{Ent}_{Q_{\hat{n}}}(M(Q_{\hat{n}}, h_{\partial Q_{\hat{n}}})) \leq \text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n})) + \theta_m(\delta) + \theta_m(\frac{1}{n}). \quad (19)$$

Proof of Lemma 4.2. We apply the Kirszbraun theorem (Theorem 3.1) to construct an injection from $M(Q_n, h_{\partial Q_n})$ into $M(Q_{\hat{n}}, h_{\partial Q_{\hat{n}}})$. The existence of such an injection implies that

$$|M(Q_{\hat{n}}, h_{\partial Q_{\hat{n}}})| \geq |M(Q_n, h_{\partial Q_n})|$$

so

$$\text{Ent}_{Q_{\hat{n}}}(M(Q_{\hat{n}}, h_{\partial Q_{\hat{n}}})) \leq \frac{|Q_n|}{|Q_{\hat{n}}|} \text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n})).$$

Of course, $\frac{|Q_n|}{|Q_{\hat{n}}|} = (\frac{n}{\hat{n}})^m + \theta_m(\frac{1}{n})$ and

$$\frac{n}{\hat{n}} = \frac{n}{(1 + \delta)n} + \theta(\frac{1}{n}) = 1 + \theta(\delta) + \theta(\frac{1}{n}).$$

Since $\text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n}))$ is bounded (see Observation 2.11), the conclusion follows.

So, it remains to show that any height function $h_{Q_n} \in M(Q_n, h_{\partial Q_n})$ can be extended to a height function in $M(Q_{\hat{n}}, h_{\partial Q_{\hat{n}}})$. We want to apply the Kirszbraun theorem. The parity condition is automatic from our parity assumption in the definition of height functions (see Definition 2.1). We must verify inequality (16).

Let $z \in \partial Q_n$ and $\hat{z} \in \partial Q_{\hat{n}}$. By the triangle inequality,

$$\begin{aligned}
& |h_{\partial Q_n}(z) - h_{\partial Q_{\hat{n}}}(\hat{z})| \\
& \leq |h_{\partial Q_n}(z) - s \cdot z| + |s \cdot (z - \hat{z})| \\
& \quad + |(s - \hat{s}) \cdot \hat{z}| + |\hat{s} \cdot \hat{z} - h_{\partial Q_{\hat{n}}}(\hat{z})| \\
& \leq \varepsilon n + |s|_{\infty} |z - \hat{z}|_1 \\
& \quad + |s - \hat{s}|_{\infty} + \varepsilon \hat{n} \\
& \leq \varepsilon n + (1 - 3\delta) |z - \hat{z}|_1 \\
& \quad + \frac{\delta^2}{1+\delta} [(1 + \delta)n + 1] + \varepsilon [(1 + \delta)n + 1] \\
& = \varepsilon(2 + \delta)n + (1 - 3\delta) |z - \hat{z}|_1 + \delta^2 n \\
& \quad + \frac{\delta^2}{1+\delta} + \varepsilon \\
& \leq \frac{\delta^2}{2+\delta} (2 + \delta)n + (1 - 3\delta) |z - \hat{z}|_1 + \delta^2 n \\
& \quad + \frac{\delta^2}{1+\delta} + \frac{\delta^2}{2+\delta} \\
& \leq 2\delta^2 n + (1 - 3\delta) |z - \hat{z}|_1 + 2\delta^2.
\end{aligned}$$

By definition, $|z - \hat{z}|_1 \geq \hat{n} - n \geq \delta n$, so $2\delta^2 n \leq 2\delta |z - \hat{z}|_1$. Moreover, the leftover term $2\delta^2$ is ≤ 1 , so

$$|h_{\partial Q_n}(z) - h_{\partial Q_{\hat{n}}}(\hat{z})| \leq (1 - \delta) |z - \hat{z}|_1 + 1.$$

We assumed that $\delta |z - \hat{z}|_1 \geq \delta^2 n > 1$, so the right-hand side above is strictly less than $|z - \hat{z}|_1$. Therefore the Kirsbraun theorem applies, which completes this proof. \square

Now, we may quickly state and prove a few useful properties of the microscopic entropy and surface tension, using Lemma 4.1 and Lemma 4.2 for the proofs.

Lemma 4.3 (Continuity of $\text{ent}(s)$). *The function $s \mapsto \text{ent}(s)$, from $[-1, 1]^m$ to $[-\log 2, 0]$, is continuous. In fact, since the domain is compact, $s \mapsto \text{ent}(s)$ is uniformly continuous.*

Proof of Lemma 4.3. First, if $|s|_{\infty} = 1$, Lemma 4.1 implies $\text{ent}(s) = 0$. As $|s|_{\infty} \rightarrow 1$, the same lemma implies $\text{ent}(s) = \theta(1 - |s|_{\infty}) \rightarrow 0$. So, $s \mapsto \text{ent}(s)$ is continuous at points s with $|s|_{\infty} = 1$.

Suppose instead that $|s|_{\infty} < 1$. In the language of Lemma 4.2, let $\delta < \frac{1}{3} \wedge (1 - |s|_{\infty})$, let $\varepsilon = \frac{\delta^2}{2+\delta}$, and consider any \hat{s} satisfying $|s - \hat{s}|_{\infty} < \frac{\delta^2}{1+\delta}$. For any n, \hat{n} as in Lemma 4.2, take $h_{\partial Q_n} = h_{\partial Q_n}^s$ and $h_{\partial Q_{\hat{n}}} = h_{\partial Q_{\hat{n}}}^{\hat{s}}$. Then by Lemma 4.2,

$$\text{ent}_n(s) \geq \text{ent}_{\hat{n}}(\hat{s}) + \theta_m(\delta) + \theta_m\left(\frac{1}{n}\right).$$

Taking $n \rightarrow \infty$, we have $\text{ent}(s) \geq \text{ent}(\hat{s}) + \theta_m(\delta)$; taking $\delta \rightarrow 0$, we conclude that $\text{ent}(s) \geq \liminf_{\hat{s} \rightarrow s} \text{ent}(\hat{s})$. By exchanging the role of s and \hat{s} , we conclude that $\text{ent}(s) \leq \limsup_{\hat{s} \rightarrow s} \text{ent}(\hat{s})$. Therefore the function $s \mapsto \text{ent}(s)$ is continuous. \square

Lemma 4.4 (Uniform convergence of $\text{ent}_n(s)$). *For a fixed dimension m , the convergence of $\text{ent}_n(s) \rightarrow \text{ent}(s)$ is uniform in $s \in [-1, 1]^m$. In other words,*

$$\text{ent}(s) = \lim_{n \rightarrow \infty} \text{ent}_n(s).$$

Proof of Lemma 4.4. Let $\varepsilon > 0$. We want to find n_0 , depending only on ε and m , such that $n \geq n_0$ implies $|\text{ent}_n(s) - \text{ent}(s)| < \varepsilon$ for any $s \in [-1, 1]^m$.

By Lemma 4.1 there exists $\delta > 0$ such that $|s|_\infty \geq 1 - \delta$ implies $|\text{ent}_n(s)| < \varepsilon$ for all n . This suffices to handle the case $|s|_\infty \geq 1 - \delta$.

For the remaining case of $|s|_\infty \leq 1 - \delta$, we rely on compactness of the space $[-1 + \delta, 1 - \delta]^m$. By uniform continuity of $\text{ent}(s)$ (see Lemma 4.2, there exists $\alpha > 0$ such that $|s_1 - s_2|_\infty \leq \alpha$ implies $|\text{ent}(s_1) - \text{ent}(s_2)| < \frac{\varepsilon}{4}$. Shrink δ if necessary so that $\frac{\delta^2}{1+\delta} \leq \alpha$, then shrink δ further so that the $\theta_m(\delta)$ term from (19), the conclusion of Lemma 4.2, is less than $\frac{\varepsilon}{4}$.

Choose a finite set of slopes $s^{(1)}, \dots, s^{(k)} \in [-1 + \delta, 1 - \delta]^m$ such that for every $s^* \in [-1 + \delta, 1 - \delta]^m$ there exists some $i = 1, \dots, k$ with $|s^* - s^{(i)}|_\infty \leq \frac{\delta^2}{1+\delta}$. Finally, choose n_0 large enough that whenever $n \geq \frac{1}{1-\delta}n_0$, the $\theta_m(\frac{1}{n})$ term from (19) is less than $\frac{\varepsilon}{4}$, and for each $i = 1, \dots, k$, $|\text{ent}_n(s^{(i)}) - \text{ent}(s^{(i)})| < \frac{\varepsilon}{4}$.

Let $n \geq n_0$, let $s^* \in [-1 + \delta, 1 - \delta]^m$ be arbitrary, and fix $i = 1, \dots, k$ such that $|s^* - s^{(i)}|_\infty < \frac{\delta^2}{1+\delta}$. We apply Lemma 4.2 twice. First take $s = s^*$, $\hat{s} = s^{(i)}$, $h_{\partial Q_n} = h_{\partial Q_n}^s$, and $h_{\partial Q_{\hat{n}}} = h_{\partial Q_{\hat{n}}}^{\hat{s}}$. The conclusion is

$$\text{ent}_{\hat{n}}(s^{(i)}) \leq \text{ent}_n(s^*) + \theta_m(\delta) + \theta_m(\frac{1}{n}),$$

and by our assumptions on δ and n_0 above, the θ terms simplify to

$$\text{ent}_{\hat{n}}(s^{(i)}) \leq \text{ent}_n(s^*) + \frac{2\varepsilon}{4}.$$

By choice of n_0 ,

$$\text{ent}_{\hat{n}}(s^{(i)}) \geq \text{ent}(s^{(i)}) - \frac{\varepsilon}{4},$$

and by choice of $s^{(i)}$,

$$\text{ent}(s^{(i)}) \geq \text{ent}(s^*) - \frac{\varepsilon}{4}.$$

Combining the last three inequalities yields

$$\text{ent}_n(s) \geq \text{ent}(s) - \varepsilon.$$

For the reverse inequality, choose $s = s^{(i)}$, $\hat{s} = s^*$, and exchange the role of n and \hat{n} . Repeating the work above, we deduce the inequality

$$\text{ent}_n(s) \leq \text{ent}(s) + \varepsilon,$$

which completes the proof of Lemma 4.4. \square

Lemma 4.5 (Robustness of Ent_{Q_n}). *Let $n \in \mathbb{N}$, $\varepsilon \in (0, \frac{1}{27})$, and $s \in [-1, 1]^m$. Let $h_{\partial Q_n} \in M(\partial Q_n)$ be such that*

$$\sup_{z \in \partial Q_n} |h_{\partial Q_n}(z) - h_{\partial Q_n}^s(z)| \leq \varepsilon n.$$

Then,

$$\text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n})) = \text{ent}(s) + \theta_m(\varepsilon) + \theta_{m,\varepsilon}(\frac{1}{n}).$$

Proof. Suppose first that $|s|_\infty \geq 1 - \varepsilon^{1/2}$. Then Lemma 4.1 applies to both $h_{\partial Q_n}$ and $h_{\partial Q_n}^s$, so

$$\begin{aligned} \text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n})) &= 0 + \theta(\varepsilon) + \theta_\varepsilon(\frac{1}{n}) \\ &= \text{ent}_n(s) + \theta(\varepsilon) + \theta_\varepsilon(\frac{1}{n}) \\ &= \text{ent}(s) + \theta_m(\varepsilon) + \theta_{m,\varepsilon}(\frac{1}{n}), \end{aligned}$$

where in the last line we used Lemma 4.4 for uniform convergence of $\text{ent}_n(s) \rightarrow \text{ent}(s)$, dependent only on dimension m .

So suppose instead that $|s|_\infty \leq 1 - \varepsilon^{1/2}$. Apply Lemma 4.2 twice. Both times take $\hat{s} = s$ and $\delta = \sqrt{3}\varepsilon$. Note that then $\varepsilon = \frac{\delta^2}{3} \leq \frac{\delta}{2+\delta}$, as required by Lemma 4.2. In the first application of Lemma 4.2 take $h_{\partial Q_n} = h_{\partial Q_n}$ and $h_{\partial Q_{\hat{n}}} = h_{\partial Q_{\hat{n}}}^s$ so that

$$\text{ent}_{\hat{n}}(s) \leq \text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n})) + \theta_m(\delta) + \theta_m(\frac{1}{n}).$$

The second time exchange n and \hat{n} to derive

$$\text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n})) \leq \text{ent}_{\hat{n}}(s) + \theta_m(\delta) + \theta_m(\frac{1}{n}).$$

Since δ is determined by ε , we may replace δ by ε in the θ terms above. And as before, Lemma 4.4 implies that $\text{ent}_{\hat{n}}(s) \rightarrow \text{ent}(s)$ and $\text{ent}_n(s) \rightarrow \text{ent}(s)$ as $n \rightarrow \infty$, at a rate depending only on the dimension and on δ (since n, \hat{n} differ from n by a factor of $(1 + \delta)^{\pm 1}$). Therefore

$$\text{Ent}_{Q_n}(M(Q_n, h_{\partial Q_n})) = \text{ent}(s) + \theta_m(\varepsilon) + \theta_{m,\varepsilon}(\frac{1}{n})$$

as claimed. \square

5. PROFILE THEOREM FOR PIECEWISE AFFINE FUNCTIONS

In this section we prove a simpler version of the profile theorem, restricted to the case where the domain R is a finite union of simplices and where the asymptotic height function h_R is piecewise affine, that is, affine when restricted to a single simplex. On one hand this case is simple enough that we can prove the profile theorem directly via over- and under-counting arguments (see the proof of Theorem 5.4 below). On the other hand, this case is sufficiently powerful to approximate general domains and height functions very well (see the proof of Theorem 2.15 and especially Lemma 6.1).

We must impose some regularity assumption on the simplices chosen; in particular we need the isoperimetric ratio to be bounded above (that is, the surface area of a simplex must not be too large in comparison to its volume). For simplicity we restrict our attention to certain families of simplices. Now let us introduce a standard notation describing these simplices.

In our exposition we follow [She05]. For a point $w = (w_1, \dots, w_m) \in \mathbb{R}^m$, we recall from our list of notations that $\lfloor w \rfloor := (\lfloor w_1 \rfloor, \dots, \lfloor w_m \rfloor) \in \mathbb{Z}^m$. For a typical point $w \in \mathbb{R}$, let $s(w)$ denote the permutation of $\{1, \dots, m\}$ which rank-orders the components of $w - \lfloor w \rfloor$. In particular,

$$w_{s(1)} - \lfloor w_{s(1)} \rfloor > w_{s(2)} - \lfloor w_{s(2)} \rfloor > \dots > w_{s(m)} - \lfloor w_{s(m)} \rfloor.$$

For example, consider the point $w = (1.1, -0.5, 2.3) \in \mathbb{R}^3$. Then

$$\lfloor w \rfloor = (1, -1, 2) \quad \text{and} \quad w - \lfloor w \rfloor = (0.1, 0.5, 0.3).$$

Since the first largest coordinate in $w - \lfloor w \rfloor$ is at index 2, the second largest coordinate is at index 3, and the third largest (i.e. the smallest) is at index 1, we have $s(w) = (2 \ 3 \ 1)$.

Definition 5.1 (Simplices of scale 1). *Let S_m denote the group of permutations on $\{1, \dots, m\}$. For $v \in \mathbb{Z}^m$ and $s \in S_m$, we define $C(v, s)$ to be the closure of the set*

$$\{w \in \mathbb{R}^m \mid \lfloor w \rfloor = v, s(w) = s\}.$$

A few members of the family $\{C(v, s) \mid v \in \mathbb{Z}^m, s \in S_m\}$ are illustrated in Figure 7 in the case of dimension $m = 2$. It is an elementary observation that the $m!$ simplices $\{C(0, s) \mid s \in S_m\}$ tile the hypercube $[0, 1]^m$, i.e. $\bigcup_{s \in S_m} C(0, s) = [0, 1]^m$, and any two simplices from $\{C(0, s) \mid s \in S_m\}$ only have at an $(m - 1)$ -dimensional intersection. It follows that, the infinite family $\{C(v, s) \mid v \in \mathbb{Z}^m, s \in S_m\}$ tiles \mathbb{R}^m .

Moreover, any two simplices $C(v_1, s_1)$ and $C(v_2, s_2)$ are isometric. That is, there exists a distance-preserving bijection $f : \mathbb{R}^m \rightarrow \mathbb{R}^m$




FIGURE 7. The two simplices in dimension 2 that tile the unit square. The simplex $C(0, (1\ 2))$ is the closure of the set of points $(x, y) \in [0, 1]^2$ such that $x > y$, and $C(0, (2\ 1))$ is the closure of the points with $y > x$. The other simplices $\{C(v, s) \mid v \in \mathbb{Z}^m, s \in S_2\}$ are translates of these two simplices.




FIGURE 8. Decomposition of a unit cube into $\{C(0, s) \mid s \in S_3\}$. The simplices have been separated for a more clear figure.

such that $f(C(v_1, s_1)) = C(v_2, s_2)$. This ensures that all the simplices $C(v, s)$ have the same isoperimetric ratio. For our purposes we will also make reference to rescaled simplices.

Definition 5.2 (Simplices of scale ℓ). *For $\ell > 0$, $v \in \mathbb{Z}^m$, and $s \in S_m$, we write*

$$\ell C(v, s) := \{\ell x \mid x \in C(v, s)\}$$

for scaled copy of the simplex $C(v, s)$, scaled out from the origin.

As before, we observe that for any $\ell > 0$, the family $\{\ell C(0, s) \mid s \in S_m\}$ tiles the hypercube $[0, \ell]^m$. Therefore again, $\{\ell C(v, s) \mid v \in \mathbb{Z}^m, s \in S_m\}$ tiles \mathbb{R}^m . To approximate a general domain R that satisfies Assumption 2.2, we consider domains which are the union of simplices.

Definition 5.3 (Simplex domain of scale ℓ). *For $\ell > 0$, a simplex domain of scale ℓ is a region $K \subset \mathbb{R}^m$ that is the union of finitely many simplices of scale ℓ . We further require that simplex domains be connected, so that a simplex domain K automatically meets the requirements from Assumption 2.2.*

For example, the union of the two simplices in Figure 7 is a simplex domain of scale 1. It is clear that simplex domains can approximate more general domains $R \subset \mathbb{R}^m$; we make this observation more precise in Lemma 6.1 below. Now, let us formulate the main result of this section, the simplicial profile theorem (Theorem 5.4). It is a special case of the profile theorem for simplex domains and piecewise affine height functions; cf. the general profile theorem (Theorem 2.15).

Theorem 5.4 (Simplicial profile theorem). *Let $K = \Delta_1 \cup \dots \cup \Delta_k$ be a simplex domain of scale ℓ , in the sense of Definition 5.3. Fix a height function $h_K \in M(K)$ such that each restriction $h_K|_{\Delta_j}$, $j = 1, \dots, k$, is affine. Let $\varepsilon > 0$, let $n \in \mathbb{N}$, and let $K_n := \{z \in \mathbb{Z}^m \mid \frac{1}{n}z \in K\}$. Then for any slope $s \in [-1, 1]^m$,*

$$\begin{aligned} \text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell)) \\ = \frac{1}{|K|} \int_K \text{ent}(\nabla h_K(x)) dx + \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}\left(\frac{1}{n}\right). \end{aligned} \quad (20)$$

Remark 5.5. *In reading the proof of Theorem 5.4 for the first time, we encourage the reader to consider only a single simplex Δ rather than a simplex domain $K = \Delta_1 \cup \dots \cup \Delta_k$. The key ideas are more clear when thinking about a single simplex. In particular the simplex is decomposed into hypercubes two times, using hypercubes of a different scale each time. The two scales of hypercubes are illustrated in Figure 9. One decomposition is used to overestimate the microscopic entropy by undercounting the set $B(K_n, h_K, \varepsilon\ell)$. The other is used to underestimate the entropy by overcounting the set.*

In the more general case of a simplex domain we still decompose twice, using hypercubes of a different size each time. A typical decomposition is illustrated in Figure 10. In particular we keep only those hypercubes that lie inside a single simplex, so that h_K has a single, well-defined slope on each Q_i . Both sides of (20) are approximately additive over the simplices, but we will not explicitly prove this result here, nor do we rely on it.

Proof. As mentioned in Remark 5.5, we subdivide the region $K \subset \mathbb{R}^m$ into hypercubes Q_i (for $i = 1, \dots, k$) of equal side length q . Two different values for the side length parameter q are used at different times.




FIGURE 9. Decomposition of a single simplex into hypercubes at two scales. In both images, the shaded squares are the Q_i from the proof of Theorem 5.4. The smaller squares on the left are used when undercounting the set $B(K_n, h_K, \varepsilon\ell)$ and the larger squares on the right are used when overcounting this set.




FIGURE 10. In Theorem 5.4 K may be a simplex domain rather than a single simplex. Then only hypercubes that lie inside one of the simplices are part of the collection $\{Q_i\}$. These hypercubes are shaded. Note that there are still two different scales of hypercubes used, as illustrated in Figure 9, but only one scale is shown above. The set G_n from later in the proof is the set of grid lines contained inside the simplex domain, and U_n is the unshaded part of the simplex domain. S_n is the union of G_n and U_n .

The cubes Q_i lie in a grid with their corners on the rescaled lattice $q\mathbb{Z}^m \subset \mathbb{R}^m$. The set $\{Q_1, \dots, Q_k\}$ enumerates all such hypercubes that lie inside exactly one of the simplices $\Delta_1, \dots, \Delta_r$, as illustrated in Figure 10. This ensures that h_K has constant derivative on Q_i , which makes later arguments simpler. For $i = 1, \dots, r$ we choose $s_i \in [-1, 1]^m$

and $b_i \in \mathbb{R}$ so that $h_K|_{Q_i} = h_{Q_i}^{s_i \cdot x + b_i}$. Specifically, this means that

$$s_i := \nabla h_K(x_i) \quad \text{and} \quad b_i := h_K(x_i) - s_i \cdot x_i,$$

for an arbitrarily chosen sample point x_i from the interior of Q_i .

The hypercubes Q_i induce a decomposition of the discrete set K_n into subsets $Q_{i,n} := \{z \in \mathbb{Z}^m \mid \frac{1}{n}z \in Q_i\}$, plus a negligible remainder $K_n \setminus \bigcup_{i=1}^r Q_{i,n}$. This remainder is the unshaded part inside the triangles in Figure 10. We write qn for the side length of the discrete hypercubes $Q_{i,n}$. Technically, each $Q_{i,n}$ has an integer side length $q_{i,n} \in \mathbb{Z}$ that is equal to either $\lfloor qn \rfloor$ or $\lceil qn \rceil$, but for simplicity we elide this detail in the rest of the proof.

Let us first sketch the main idea of the proof. We start with the integral on the right-hand side of (20). Since h_K is piecewise affine, the integral reduces to a finite sum

$$\begin{aligned} \frac{1}{|K|} \int_K \text{ent}(\nabla h_K(x)) dx &= \sum_{i=1}^r \frac{1}{|K|} \int_{Q_i} \text{ent}(\nabla h_K|_{Q_i}(x)) dx + \theta_m\left(\frac{q}{\ell}\right) \\ &= \sum_{i=1}^r \frac{|Q_i|}{|K|} \text{ent}(s_i) + \theta_m\left(\frac{q}{\ell}\right) \\ &= \frac{1}{r} \sum_{i=1}^r \text{ent}(s_i) + \theta_m\left(\frac{q}{\ell}\right), \end{aligned} \tag{21}$$

where we recall that $s_i = \nabla h_K(x_i)$ for $x_i \in Q_i$. Both of the two values for the hypercube side length parameter q are chosen so that $\theta_m(\frac{q}{\ell}) = \theta_m(\varepsilon)$. The $\theta_m(\frac{q}{\ell})$ errors arise from the uncovered region $K \setminus \bigcup_{i=1}^r Q_i$, i.e. the unshaded parts of the simplex domain in Figure 10. Indeed, one simply compares the measure $|\Delta_j| = \frac{1}{m!} \ell^m$ against that of the smaller simplex Δ'_j with sides moved $\sqrt{m}q$ units inwards. Any hypercube Q_i that intersects Δ'_j must lie inside Δ_j . The $\theta_m(\frac{q}{\ell})$ error bound follows.

Now, we turn to the left-hand side of (20). Our goal is to relate $\text{Ent}_{K_n}(B(K_n, h_K, \varepsilon \ell))$ to the sum at the right-hand side of (21). Towards this end, we will under- and over-count the set of height functions $B(K_n, h_K, \varepsilon \ell)$, in order to derive the over- and under-estimates

$$\text{Ent}_{K_n}(B(K_n, h_K, \varepsilon \ell)) \leq \frac{1}{r} \sum_{i=1}^r \text{ent}(s_i) + \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}\left(\frac{1}{n}\right) \tag{22}$$

and

$$\text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell)) \geq \frac{1}{r} \sum_{i=1}^r \text{ent}(s_i) + \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}(\frac{1}{n}). \quad (23)$$

Equations (21), (22), and (23), together with the observation made above that $\theta_m(\frac{q}{\ell}) = \theta_m(\varepsilon)$, suffice to prove the theorem. In order to prove (22), we will undercount height functions in $B(K_n, h_K, \varepsilon\ell)$. We choose $q \approx \varepsilon\ell$ and consider only height functions that agree with h_{K_n} on each boundary $\partial Q_{i,n}$. These boundary data, together with the small size of $Q_{i,n}$, ensure that h_{K_n} satisfies the ℓ^∞ condition for membership in $B(K_n, h_K, \varepsilon\ell)$. Then, to prove (23), we overcount height functions. We choose a larger value $q \approx \varepsilon^{1/2}\ell$. Any height function $h_n \in B(K_n, h_K, \varepsilon\ell)$, when restricted to one of the boundary sets $\partial Q_{i,n}$ and rescaled appropriately, fluctuates away from h_K by at most $\varepsilon\ell n = \varepsilon^{1/2}qn$. When ε is small, this allows us to compare the entropy on Q_i to $\text{Ent}(M(Q_{i,n}, h_{\partial Q_{i,n}}^{s_i})) = \text{ent}_{qn}(s_i)$, where $s_i = \nabla h_K(x_i)$ for $x_i \in Q_i$.

Now, let us describe the undercounting argument in detail. We seek to derive (22), an overestimate of $\text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell))$, by undercounting the set $B(K_n, h_K, \varepsilon\ell)$. We take the side length of the hypercubes $\{Q_i\}$ to be $q = \frac{1}{4}\varepsilon\ell$.

We define an injection from the product set $\prod_i M(Q_{i,n}, h_{\partial Q_{i,n}}^{s_i \cdot x + b_i})$ into the ball $B(K_n, h_K, \varepsilon\ell)$ in the natural way: given a tuple of height functions $h_{Q_{i,n}} \in M(Q_{i,n}, h_{\partial Q_{i,n}}^{s_i \cdot x + b_i})$ for $i = 1, \dots, r$, we define $h_{K_n}(z) := h_{Q_{i,n}}(z)$ if $z \in Q_{i,n}$. For $z \in K_n \setminus \bigcup_{i=1}^r Q_{i,n}$, if $\frac{1}{n}z \in \Delta_i$ we set $h_{K_n}(z) = h_{\{z\}}^{s_i \cdot x + b_i}(z)$. It follows from Lemma 2.8 that this function h_{K_n} is a height function,

Let us check that $h_{K_n} \in B(K_n, h_K, \varepsilon\ell)$. For $z \in K_n \setminus \bigcup_{i=1}^r Q_{i,n}$, the estimate $|\frac{1}{n}h_{K_n}(z) - h_K(\frac{1}{n}z)| \leq \varepsilon\ell$ is immediate from the definition of $h_{K_n}(z)$. So, suppose that $z \in Q_{i,n}$ for some $i = 1, \dots, r$, and let $z' \in Q_{i,n}$ be a boundary point in $\partial Q_{i,n}$ that minimizes ℓ^1 distance from z . In particular, $|z - z'|_1 \leq qn$, so

$$\begin{aligned} \left| \frac{1}{n}h_{K_n}(z) - h_K(\frac{1}{n}z) \right| &= \left| \frac{1}{n}h_{Q_{i,n}}(z) - h_{Q_i}(\frac{1}{n}z) \right| \\ &\leq \left| \frac{1}{n}h_{Q_{i,n}}(z) - \frac{1}{n}h_{Q_{i,n}}(z') \right| + \left| \frac{1}{n}h_{Q_{i,n}}(z') - h_{Q_i}(\frac{1}{n}z') \right| \\ &\quad + \left| h_{Q_i}(\frac{1}{n}z') - h_{Q_i}(\frac{1}{n}z) \right| \\ &\leq \frac{1}{n}|z - z'|_1 + \frac{1}{n} + \frac{1}{n}|z - z'|_1 \\ &\leq 3q < \varepsilon\ell, \end{aligned}$$

at least for n large enough that $\frac{1}{n} \leq q = \frac{1}{4}\varepsilon\ell$. Therefore $h_{K_n} \in B(K_n, h_K, \varepsilon\ell)$ as desired. Thus

$$\left| \prod_{i=1}^r M(Q_{i,n}, h_{\partial Q_{i,n}}^{s_i \cdot x + b_i}) \right| \leq |B(K_n, h_K, \varepsilon\ell)|,$$

so

$$\sum_{i=1}^r \frac{|Q_{i,n}|}{|K_n|} \text{Ent}_{Q_{i,n}}(M(Q_{i,n}, h_{\partial Q_{i,n}}^{s_i \cdot x + b_i})) \geq \text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell)). \quad (24)$$

Now,

$$\text{Ent}_{Q_{i,n}}(M(Q_{i,n}, h_{\partial Q_{i,n}}^{s_i \cdot x + b_i})) = \text{ent}_{qn}(s_i)$$

by translation invariance (see Observation 2.10). Moreover, because $qn = \frac{1}{4}\varepsilon\ell n$, we have $\text{ent}_{qn}(s_i) \rightarrow \text{ent}(s_i)$ as $n \rightarrow \infty$ at a rate dependent on $\varepsilon\ell$; by Lemma 4.4, the convergence is uniform with respect to s_i . In other words, $\text{ent}_{qn}(s_i) = \text{ent}(s_i) + \theta_{m,\varepsilon,\ell}(\frac{1}{n})$. Therefore, recalling (24), we have

$$\text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell)) \leq \sum_{i=1}^r \frac{|Q_{i,n}|}{|K_n|} (\text{ent}(s_i) + \theta_{m,\varepsilon,\ell}(\frac{1}{n})). \quad (25)$$

Now, the difference between $\sum_{i=1}^r \frac{|Q_{i,n}|}{|K_n|}$ and $\frac{1}{r}$ is $\theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}(\frac{1}{n})$, where the first error term accounts for the unshaded part of Figure 10, and the second term is due to discretization effects. Therefore (25) simplifies to

$$\text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell)) \leq \frac{1}{r} \sum_{i=1}^r \text{ent}(s_i) + \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}(\frac{1}{n}),$$

which is exactly the overestimate (22).

Now, we turn to (23). We will overcount $B(K_n, h_K, \varepsilon\ell)$ in order to underestimate the entropy $\text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell))$. We will take side length q of the hypercubes Q_i to be $q = \varepsilon^{1/2}\ell$ for this part of the argument.

The basic idea is the following: we choose a subset $S_n \subset K_n$, and we only enforce the condition that $|\frac{1}{n}h_{K_n}(x) - h_K(\frac{1}{n}x)| < \varepsilon\ell$ from Definition 2.5 on S_n rather than on all of K_n . S_n is the complement of the (interiors of the) grid cells $Q_{i,n}$, so for any fixed height values on S_n , we can count the number of all extensions into the grid cells using a sum of entropy over the cells. There are many possible height values on S_n that satisfy the $\varepsilon\ell$ error condition, but ultimately not too many because S_n is small (compared to K_n).

Let us provide more detail. We define S_n as follows. Let G_n denote the grid formed by the boundaries of $Q_{i,n}$, i.e. the part of the grid lines

from Figure 10 that lies inside the simplex domain. Let U_n denote the points in K_n that lie outside of any hypercube $Q_{i,n}$, i.e. the unshaded part of the simplex domain in Figure (10). Let $S_n := G_n \cup U_n$. (As claimed, the complement $K_n \setminus S_n$ is the interior of the grid cells $Q_{i,n}$.) Additionally, let $\text{Adm}(S_n)$ denote the set

$$\text{Adm}(S_n) := \{ \text{“admissible” height functions on } S_n \}, \quad (26)$$

where “admissible” means those height functions $h_{S_n} \in M(S_n)$ that admit an extension to a height function in $B(K_n, h_K, \varepsilon\ell)$.

We claim that there is an injection from $B(K_n, h_K, \varepsilon\ell)$ into

$$\bigsqcup_{h_{S_n} \in \text{Adm}(S_n)} \prod_{i=1}^r M(Q_{i,n}, h_{S_n}|_{\partial Q_{i,n}}), \quad (27)$$

where “ \bigsqcup ” denotes the disjoint union (so for distinct height functions h_{S_n} and \tilde{h}_{S_n} in $\text{Adm}(S_n)$, the product sets $\prod_1^r M(Q_{i,n}, h_{S_n}|_{\partial Q_{i,n}})$ and $\prod_1^r M(Q_{i,n}, \tilde{h}_{S_n}|_{\partial Q_{i,n}})$ are considered disjoint inside the set from (27)).

Indeed, for any $h_{K_n} \in B(K_n, h_K, \varepsilon\ell)$, the function $h_{S_n} := h_{K_n}|_{S_n}$ is by definition in $\text{Adm}(S_n)$, and $(h_{K_n}|_{Q_{i,n}})_{i=1}^r$ lies in the Cartesian product from the right-hand side of (27). To see that this map is an injection, suppose that h_{K_n} and \tilde{h}_{K_n} map to the same point. Then by definition of the (purported) injection, $h_{K_n}|_{Q_{i,n}} = \tilde{h}_{K_n}|_{Q_{i,n}}$ for each hypercube $Q_{i,n}$. Additionally, since the right-hand side of (27) is a *disjoint* union, we have $h_{K_n}|_{S_n} = \tilde{h}_{K_n}|_{S_n}$. Since $K_n = \bigcup_i Q_{i,n} \cup S_n$, the two height functions h_{K_n} and \tilde{h}_{K_n} are identical. Therefore, the map is an injection, and so

$$\begin{aligned} |B(K_n, h_K, \varepsilon\ell)| &\leq \sum_{h_{S_n} \in \text{Adm}(S_n)} \prod_{i=1}^r |M(Q_{i,n}, h_{S_n}|_{\partial Q_{i,n}})| \\ &\leq |\text{Adm}(S_n)| \max_{h_{S_n} \in \text{Adm}(S_n)} \prod_{i=1}^r |M(Q_{i,n}, h_{S_n}|_{\partial Q_{i,n}})|. \end{aligned}$$

Taking logarithms and multiplying by $-\frac{1}{|K_n|}$, we see that

$$\begin{aligned}
& \text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell)) \\
& \geq \min_{h_{S_n} \in \text{Adm}(S_n)} \sum_{i=1}^r -\frac{1}{|K_n|} \log |M(Q_{i,n}, h_{S_n}|_{\partial Q_{i,n}})| \\
& \quad - \frac{\log |\text{Adm}(S_n)|}{|K_n|} \\
& = \min_{h_{S_n} \in \text{Adm}(S_n)} \sum_{i=1}^r \frac{|Q_{i,n}|}{|K_n|} \text{Ent}_{Q_{i,n}}(M(Q_{i,n}, h_{S_n}|_{\partial Q_{i,n}})) \\
& \quad - \frac{\log |\text{Adm}(S_n)|}{|K_n|}.
\end{aligned} \tag{28}$$

Now, we are almost done. We use three more asymptotic identities in the right-hand side of (28) to derive (23). First and simplest, since $\frac{|Q_{i,n}|}{|K_n|} = \frac{1}{r} + \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}(\frac{1}{n})$ and $\text{Ent}_{Q_{i,n}}$ is bounded (Observation 2.11), we replace $\frac{|Q_{i,n}|}{|K_n|}$ by $\frac{1}{r}$ in (28).

Second, we apply Lemma 4.5 to replace $\text{Ent}_{Q_{i,n}}(M(Q_{i,n}, h_{S_n}|_{\partial Q_{i,n}}))$ by $\text{ent}(s_i)$. We fix a height function h_{S_n} that achieves the minimum, then apply the lemma on each of the hypercubes $Q_{i,n}$. We recall that the hypercubes have side length $qn = \varepsilon^{1/2}\ell n$. Since h_{S_n} is admissible and since $h_K|_{Q_i} = h_{Q_i}^{s_i \cdot x + b_i}$, h_{S_n} is approximately affine, i.e.

$$|h_{S_n}(z) - h_{\partial Q_{i,n}}^{s_i \cdot x + b_i}(z)| \leq \varepsilon\ell n = \varepsilon^{1/2}qn.$$

So, Lemma 4.5 applies and yields

$$\begin{aligned}
\text{Ent}_{Q_{i,n}}(M(Q_{i,n}, h_{S_n}|_{\partial Q_{i,n}})) &= \text{ent}(s_i) + \theta_m(\varepsilon^{1/2}) + \theta_{m,\varepsilon^{1/2}}(\frac{1}{\varepsilon^{1/2}\ell n}) \\
&= \text{ent}(s_i) + \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}(\frac{1}{n}),
\end{aligned}$$

where the last line is just a matter of hiding the functions like $\varepsilon^{1/2}$ inside our θ -notation. (See Section 2.4 for the definition of θ -notation.)

Finally, we claim that $\frac{1}{|K_n|} \log |\text{Adm}(S_n)| = \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}(\frac{1}{n})$, where we recall that the set $\text{Adm}(S_n)$ was defined in (26). To see this, fix a base point $z_0 \in S_n$. There are at most $(2\varepsilon n + 1)$ choices for $h_{S_n}(z_0)$, by definition of $B(K_n, h_K, \varepsilon\ell)$. Then, since $S_n \subset \mathbb{Z}^m$ is connected (in the sense of graph theory), there are less than $2^{|S_n|}$ ways to extend h_{S_n} to the rest of S_n . So, we must estimate $|S_n|$. We recall that $S_n = G_n \cup U_n$, where G_n is the grid and U_n the unshaded region in Figure 10. Since G_n grows like n^{m-1} while $|K_n|$ grows like n^m , we have $\frac{|G_n|}{|K_n|} = \theta_{m,\varepsilon,\ell}(\frac{1}{n})$. Next, the part of $K \subset \mathbb{R}^m$ that lies outside of any hypercube Q_i , that is, the unshaded part of the simplex domain in Figure 10, is a $\theta_m(\varepsilon)$

fraction of the total volume of K . Even with discretization errors, $\frac{|U_n|}{|K_n|} = \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}(\frac{1}{n})$. Altogether,

$$\begin{aligned} \frac{1}{|K_n|} \log |\text{Adm}(S_n)| &\leq \frac{1}{|K_n|} \log \left((2\varepsilon\ell n + 1) 2^{|S_n|} \right) \\ &= \frac{\log(2\varepsilon\ell n + 1)}{|K_n|} + \frac{\log 2}{|K_n|} (|G_n| + |U_n|) \\ &= \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}(\frac{1}{n}). \end{aligned}$$

Applying the three asymptotic identities above in (28), we derive

$$\text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell)) \geq \frac{1}{r} \sum_{i=1}^r \text{ent}(s_i) + \theta_m(\varepsilon) + \theta_{m,\varepsilon,\ell}(\frac{1}{n}),$$

which is exactly (23). This completes the proof. \square

6. PROOF OF THE PROFILE THEOREM

In this section we extend Theorem 5.4, the profile theorem for piecewise affine height functions on simplex domains, to general asymptotic height theorem on general domains (subject to Assumption 2.2, as always).

The proof is an approximation argument, and we will need some auxiliary results. The most helpful is the simplicial Rademacher theorem, which states that Lipschitz functions are well-approximated by piecewise affine functions on a simplex domain. The other auxiliary results are about robustness of the microscopic and macroscopic entropies under changes in the domain and in the asymptotic height profile.

The simplicial Rademacher theorem is a general fact about Lipschitz functions. There is nothing particular to our setting, except for the use of our term “asymptotic height function” instead of “Lipschitz function.” Related results include [Sch14], which extends Lemma 6.1 from Lipschitz functions to Sobolev functions, but weakens the approximation somewhat and is therefore not suitable for our purposes here. The statement of the simplicial Rademacher theorem is adapted from Lemma 2.2 of [CKP01], and the proof is inspired by the proof there.

Lemma 6.1 (Simplicial Rademacher theorem). *Let $R \subseteq \mathbb{R}^m$ be a region satisfying Assumption 2.2, and let $h_R \in M(R, h_{\partial R})$ be an asymptotic height function on R . For any $\varepsilon > 0$ and any $\ell > 0$ sufficiently small (depending on ε), we may choose a simplex domain $K = \Delta_1 \cup \dots \cup \Delta_k \subseteq R$ of scale ℓ (see Definition 5.3) and a piecewise affine asymptotic height function $h_K : K \rightarrow \mathbb{R}$ (that is, an asymptotic*

height function such that each restriction $h_K|_{\Delta_i} : \Delta_i \rightarrow \mathbb{R}$ is affine) that satisfy the following properties:

- (a) $|R \setminus K| < \varepsilon$, where $|\cdot|$ denotes the Lebesgue measure, and $d_H(K, R) < \varepsilon$, where d_H denotes Hausdorff metric;
- (b) $\max_{x \in K} |h_K(x) - h_R(x)| < \frac{1}{2}\varepsilon$; and
- (c) on at least a $(1 - \varepsilon)$ fraction of the points in K (by Lebesgue measure), the gradients $\nabla h_K(x)$ and $\nabla h_R(x)$ agree to within ε ; more precisely, $\frac{1}{|K|}|\{x \in K \mid |\nabla h_K(x) - \nabla h_R(x)|_2 \geq \varepsilon\}| < \varepsilon$.

Remark 6.2. We recall that the Rademacher theorem states that a Lipschitz function h_R is differentiable almost everywhere. However ∇h_R may be poorly behaved. The Rademacher theorem gives no control over ∇h_R , and the Lipschitz property only implies boundedness of the derivative, not regularity. The simplicial Rademacher theorem provides an approximation both to h_R and to its derivative. Moreover the approximating function h_K has a very simple derivative, despite the potential wildness of ∇h_R . The cost is that h_K only approximates h_R well on a (large) portion of the domain rather than almost everywhere, but for our purposes this is a good trade-off.

In fact, it is not necessary that the function h_R be Lipschitz. Almost everywhere differentiability is sufficient.

Before giving the proof of Lemma 6.1, we state and prove the following lemma about the standard simplices from Definition 5.1.

Lemma 6.3. Let Δ be any of the simplices $C(v, s)$ for $v \in \mathbb{Z}^m$ and $s \in S_m$. The $m + 1$ vertices of Δ can be labelled $x^{(0)}, \dots, x^{(m)}$ in such a way that, for each $i = 1, \dots, m$,

$$x^{(i)} - x^{(i-1)} = e^{(s(i))},$$

where for $1 \leq j \leq m$, $e^{(j)}$ denotes the j -th standard basis vector (i.e., all entries of $e^{(j)}$ are 0, except the j -th entry, which is 1).

Remark 6.4. We encourage the reader to keep Figure 8 in mind (or better, in sight) while reading this proof.

Proof. For simplicity, we assume without loss of generality that $v = 0$. We use the permutation s to define a path between vertices of the simplex $C(0, s)$ starting at $(0, \dots, 0)$ and ending at $(1, \dots, 1)$. To construct the path, first observe that

$$C(0, s) = \{x = (x_1, \dots, x_m) \in [0, 1]^m \mid x_{s(i)} \geq x_{s(j)} \text{ for all } i < j\}.$$

In other words, the $s(1)$ -th component of x must be greater than the $s(2)$ -th, which is greater than or equal to the $s(3)$ -th, and so on. The path travels from $(0, \dots, 0)$ along the $s(1)$ -th axis to $e_{s(1)}$, then parallel

to the $s(2)$ -th axis to $e_{s(1)} + e_{s(2)}$, and so on up to $\sum_{i=1}^m e_i = (1, \dots, 1)$. Numbering the vertices of the path from $x^{(0)}$ to $x^{(m)}$ proves the lemma. \square

Now, we are ready for the proof of Lemma 6.1.

Proof of Lemma 6.1. Let $\ell > 0$. We choose the simplex domain $K = \Delta_1 \cup \dots \cup \Delta_k$ such that $\{\Delta_1, \dots, \Delta_k\}$ enumerates all simplices of scale ℓ (cf. Definition 5.2) that are contained in R . We define the asymptotic height function $h_K \in M(K)$ to agree with h_R on the vertices of the simplices in K , and we extend h_K into the rest of each Δ_i by linear interpolation. We will show that, once ℓ is small enough, properties (a), (b), and (c) from Lemma 6.1 all hold.

First we prove (a). The fact that $|K|$ tends to $|R|$ as $\ell \rightarrow 0$ is elementary measure theory, and we omit the proof.

Recall from (1) that we define the Hausdorff metric d_H in terms of the ℓ^1 metric on \mathbb{R}^m , for reasons explained in Remark 2.3. Therefore, for the second part of (a), it suffices to show that

$$R \subset K + B_{\ell^1}(0, \varepsilon) := \{x + y \mid x \in K, |y|_1 < \varepsilon\}.$$

We do this by constructing a subset $R' \subset R$ such that $R \subset R' + B_{\ell^1}(0, \varepsilon)$ and $R' + B_{\ell^1}(0, \alpha) \subset R$ for some $\alpha < \varepsilon$. The latter condition ensures that, for ℓ small enough, $R' \subset K$. Indeed, the ℓ^1 -diameter of a simplex of scale ℓ is $\leq m\ell$ (i.e. $\text{diam}_1 \Delta := \max\{|x - y|_1 \mid x, y \in \Delta\} \leq m\ell$), so as long as $\ell < \frac{\alpha}{m}$, every point $x \in R'$ belongs to a simplex Δ_i of scale ℓ which is part of K . Therefore $R' \subset K$, so $R \subset K + B_{\ell^1}(0, \varepsilon)$ as intended.

We proceed to construct R' . Since R is the closure of its interior, $R \subset \bigcup_{x \in R^\circ} B_{\ell^1}(x, \varepsilon)$. By compactness, choose x_1, \dots, x_k such that $R \subset B_{\ell^1}(x_1, \varepsilon) \cup \dots \cup B_{\ell^1}(x_k, \varepsilon)$. For $i = 1, \dots, k$, let

$$\alpha_i := \max\{\tilde{\alpha} > 0 \mid B_{\ell^1}(x_i, \tilde{\alpha}) \subset R\}$$

Since $x_i \in R^\circ$, each $\alpha_i > 0$; since R is compact, each $\alpha_i < \infty$. Let $\alpha = \min\{\alpha_1, \dots, \alpha_k, \frac{\varepsilon}{2}\}$, and set

$$R' = \{x \in R^\circ \mid B_{\ell^1}(x, \alpha) \subset R\}.$$

By construction, $R' + B_{\ell^1}(0, \alpha) \subset R$, and since $x_1, \dots, x_k \in R'$, $R \subset R' + B_{\ell^1}(0, \varepsilon)$. So R' satisfies the relations claimed above, which completes the proof of (a).

For later use we strengthen the volume estimate from (a). Choose ℓ smaller so that

$$|R \setminus K| < \varepsilon' := \min \left\{ \frac{\varepsilon}{2}, \frac{\varepsilon}{2|R|}, \frac{1}{2} \right\}. \quad (29)$$

In particular, this implies that

$$|R| < \frac{1}{1 - \varepsilon'} |K|. \quad (30)$$

Let us describe the key idea used to prove (b) and (c). We consider points x where h_R is differentiable, and indeed where h_R is locally approximated well by its first-order Taylor polynomial. Once the simplices Δ_i are small enough and contain a “good” point x , the vertices all lie close to x , so we can use the Taylor polynomial to estimate the values of h_R on the vertices. This yields the proof of (b) and (c).

To be more precise, we define a set S_{ρ_0} of “good” points. Recall that the Lipschitz function h_R is almost everywhere differentiable, by the Rademacher theorem. Consider any point $x \in R$ at which ∇h_R exists. Define the Taylor polynomial

$$L_x(y) := h_R(x) + \nabla h_R(x) \cdot (y - x).$$

By the definition of differentiability,

$$\lim_{y \rightarrow x} \frac{|h_R(y) - L_x(y)|}{|x - y|_2} = 0,$$

so there exists $r_0(x) > 0$ such that, for any $y \in R$ with $|y - x|_2 < r_0(x)$,

$$|h_R(y) - L_x(y)| < \left(\frac{\varepsilon}{4\sqrt{m}} \wedge \frac{\varepsilon}{2m} \right) |y - x|_2. \quad (31)$$

(Recall that $m \in \mathbb{N}$ is the dimension parameter; we could replace the parenthesized expression by $\frac{\varepsilon}{4m}$, but the expressions $\frac{\varepsilon}{4\sqrt{m}}$ and $\frac{\varepsilon}{2m}$ are useful later.) For $\rho > 0$, define the set $S_\rho \subset R$ by

$$S_\rho := \{x \in R \mid r_0(x) \geq \rho\}.$$

As $\rho \rightarrow 0$, the sets S_ρ increase to the full-measure subset of R on which h_R is differentiable. Therefore $|S_\rho| \rightarrow |R|$ as $\rho \rightarrow 0$, and in particular, there exists $\rho_0 > 0$ such that

$$|S_{\rho_0}| \geq \left(1 - \frac{\varepsilon'}{2}\right) |R|. \quad (32)$$

We choose $\ell_0 \leq \frac{\rho_0}{\sqrt{m}}$. By the Pythagorean theorem (in m dimensions), if x, y are two points that lie in a simplex Δ_i and if $x \in S_{\rho_0}$, then

$|x - y|_2 \leq \sqrt{m}\ell < \rho_0 \leq r(x)$. Therefore by (31),

$$\max_{y \in \Delta_i} |h_R(y) - L_x(y)| < \frac{\varepsilon\ell}{4} \wedge \frac{\varepsilon\ell}{2\sqrt{m}}. \quad (33)$$

There are two more steps to prove (b). First, under the assumption that $x \in \Delta_i \cap S_{\rho_0}$, we have compared $h_R|_{\Delta_i}$ to the Taylor polynomial of h_R centered at x ; we should also compare h_K to the same polynomial. Second, we show that at least $(1 - \varepsilon)k$ of the simplices have some intersection with S_{ρ_0} . Then it is straightforward to complete the proof of (b).

Regarding h_K , recall that on the vertices y_0, \dots, y_m of Δ_i , h_K agrees with h_R . Therefore by (33),

$$\max_{0 \leq i \leq m} |h_K(y_i) - L_x(y_i)| < \frac{\varepsilon\ell}{4}.$$

The function $y \mapsto h_K(y) - (h_R(x) + \nabla h_R(x) \cdot (y - x))$ is affine, so $y \mapsto |h_K(y) - (h_R(x) + \nabla h_R(x) \cdot (y - x))|$ is convex. Hence

$$\max_{y \in \Delta_i} |h_K(y) - L_x(y)| < \frac{\varepsilon\ell}{4}. \quad (34)$$

Therefore, if a simplex Δ_i has any intersection with S_{ρ_0} , then h_K satisfies the L^∞ inequality from (b) over Δ_i .

Now, let k_0 denote the number of simplices that intersect S_{ρ_0} . We claim that $k_0 \geq (1 - \varepsilon')k$. Of course, since S_{ρ_0} has large measure, it must intersect many of the simplices. More precisely, because $(k - k_0)$ is the number of simplices that do not intersect S_{ρ_0} ,

$$|S_{\rho_0}| \leq |R| - (k - k_0)|\Delta_1|.$$

(Recall that the simplices of scale ℓ are isometric, so $|\Delta_1| = \dots = |\Delta_k| = \frac{\ell^m}{m!}$.) Therefore

$$\begin{aligned} k - k_0 &\leq \frac{|R| - |S_{\rho_0}|}{|\Delta_1|} \\ &\leq \frac{\varepsilon' |R|}{2 |\Delta_1|} && (\text{By (32)}) \\ &\leq \frac{\varepsilon' (1 - \varepsilon')^{-1} |K|}{2 |\Delta_1|} && (\text{By (30)}) \\ &= \frac{\varepsilon'}{2} (1 - \varepsilon')^{-1} k && (\text{Since } K = \bigcup_{i=1}^k \Delta_i) \\ &\leq \varepsilon' k, \end{aligned}$$

since $\varepsilon' \leq \frac{1}{2}$. So, both (33) and (34) apply on $k_0 \geq (1 - \varepsilon')k$ of the simplices. We throw away the “bad” simplices, at the cost of increasing $|R \setminus K|$ by at most

$$\varepsilon' k |\Delta_i| \leq \left(\frac{\varepsilon}{2|R|} \right) k \left(\frac{|R|}{k} \right) = \frac{\varepsilon}{2}.$$

This is permissible by (29), since $|R \setminus K|$ was previously $\leq \frac{\varepsilon}{2}$. We have therefore proven (b).

For (c), we claim that if $x \in K \cap S_{\rho_0}$, then $|\nabla h_R(x) - \nabla h_K(x)| < \varepsilon$. By (30) and (32), $|K \cap S_{\rho}| > (1 - \varepsilon)|R|$, so this will suffice to prove (c).

Let $x \in K \cap S_{\rho_0}$, and fix i such that $x \in \Delta_i$. We will use Lemma 6.3 in order to make the calculations as concrete as possible. In particular, we label the vertices of Δ_i as y_0, y_1, \dots, y_m in such a way that $y_i - y_{i-1} = \ell e_{s(i)}$, where $e_{s(i)}$ is the $s(i)$ -th standard basis vector for some permutation $s \in S_m$ of $\{1, \dots, m\}$. Then by (33),

$$|h_R(y_i) - L_x(y_i)| < \frac{\varepsilon \ell}{2\sqrt{m}} \quad \text{and} \quad |h_R(y_{i-1}) - L_x(y_{i-1})| < \frac{\varepsilon \ell}{2\sqrt{m}},$$

where we recall that $L_x(y) = h_R(x) + \nabla h_R(x) \cdot (y - x)$ is the first-order Taylor polynomial of h_R at x . Combining these two inequalities,

$$\left| (h_R(y_i) - h_R(y_{i-1})) - (\nabla h_R(x) \cdot (y_i - y_{i-1})) \right| < \frac{\varepsilon \ell}{\sqrt{m}}.$$

Since $y_i - y_{i-1} = \ell e_{s(i)}$,

$$\left| \frac{h_R(y_i) - h_R(y_{i-1})}{|y_i - y_{i-1}|_2} - \nabla h_R(x) \cdot \frac{y_i - y_{i-1}}{|y_i - y_{i-1}|_2} \right| < \frac{\varepsilon}{\sqrt{m}}. \quad (35)$$

Because h_K is the linear interpolation of h_R from the vertices y_0, \dots, y_m to the rest of Δ_i , we see that the first term on the left-hand side of (35) is

$$\frac{h_R(y_i) - h_R(y_{i-1})}{|y_i - y_{i-1}|_2} = \frac{h_K(y_i) - h_K(y_{i-1})}{|y_i - y_{i-1}|_2} = \frac{\partial h_K}{\partial x_{s(i)}}(x).$$

And of course, $(y_i - y_{i-1})/|y_i - y_{i-1}|_2 = e_{s(i)}$, so the second term on the left-hand side of (35) is

$$\nabla h_R(x) \cdot \frac{y_i - y_{i-1}}{|y_i - y_{i-1}|_2} = \frac{\partial h_R}{\partial x_{s(i)}}(x).$$

The last three equations hold for all $i = 1, \dots, m$. Therefore we may drop the permutation $s(i)$ from the partial derivatives and conclude that, for every i ,

$$\left| \frac{\partial h_K}{\partial x_i}(x) - \frac{\partial h_R}{\partial x_i}(x) \right| < \frac{\varepsilon}{\sqrt{m}}.$$

Thus

$$|\nabla h_K(x) - \nabla h_R(x)|_2 < \varepsilon.$$

□

The next three lemmas regard the robustness of the macroscopic entropy and microscopic entropy to changes in domain and asymptotic height function. As seen in the simplicial Rademacher theorem (Lemma 6.1), we will change both the domain and the asymptotic height function. As long as these changes are small enough (in the appropriate senses), these lemmas show that the macroscopic entropy and microscopic entropy change by a small amount.

First, we deal with robustness of the macroscopic entropy. Because Ent_R is an integral function with continuous and bounded integrand, robustness with respect to changes in both domain and asymptotic height function is easy to prove by standard analytic arguments. The main requirement is control over the change in the derivative of the asymptotic height function, as is provided by (c) from the simplicial Rademacher theorem (see Lemma 6.1).

Lemma 6.5 (Robustness of macroscopic entropy under approximations). *Let $\varepsilon > 0$, and let $\tilde{R} \subseteq R \subset \mathbb{R}^m$ be sets meeting the assumptions from Assumption 2.2, with $|R \setminus \tilde{R}| < \varepsilon$. Let $h_{\tilde{R}} \in M(\tilde{R})$ and $h_R \in M(R)$ be such that*

$$\left| \left\{ x \in \tilde{R} \mid |\nabla h_{\tilde{R}}(x) - \nabla h_R(x)|_2 \geq \varepsilon \right\} \right| < \varepsilon.$$

Then,

$$\text{Ent}_R(h_R) = \text{Ent}_{\tilde{R}}(h_{\tilde{R}}) + \theta_m(\varepsilon).$$

Proof. Recall that $\text{Ent}_R(h_R) = \frac{1}{|R|} \int_R \text{ent}(\nabla h_R(x)) dx$ and $\text{Ent}_{\tilde{R}}(h_{\tilde{R}}) = \frac{1}{|\tilde{R}|} \int_{\tilde{R}} \text{ent}(\nabla h_{\tilde{R}}(x)) dx$. Split R into three parts.

The set $\{x \in \tilde{R} \mid |\nabla h_{\tilde{R}}(x) - \nabla h_R(x)|_2 \geq \varepsilon\}$ has measure less than ε by hypothesis. Since $\text{ent}(s)$ is bounded (see Observation 2.13), the contribution of the points in this set to $\text{Ent}_{\tilde{R}}(h_{\tilde{R}})$ is within $\theta(\varepsilon)$ of the contribution to $\text{Ent}_R(h_R)$.

Likewise, the set $R \setminus \tilde{R}$ has measure at most ε , so the contribution to $\text{Ent}_R(h_R)$ is $\theta(\varepsilon)$. Of course, this set does not contribute to $\text{Ent}_{\tilde{R}}(h_{\tilde{R}})$.

Finally, for the remaining points x , $|\nabla h_{\tilde{R}}(x) - \nabla h_R(x)|_2 < \varepsilon$. Since $\text{ent}(s)$ is uniformly continuous on its domain $s \in [-1, 1]^m$, we have $|\text{ent}(\nabla h_{\tilde{R}}(x)) - \text{ent}(\nabla h_R(x))| < \theta_m(\varepsilon)$. Since the integrands differ by at most $\theta_m(\varepsilon)$ and since the integrals are normalized by $\frac{1}{|R|}$ and $\frac{1}{|\tilde{R}|}$, the contribution from this third part of the domain is also $\theta_m(\varepsilon)$. □

Now, we turn to the microscopic entropy. Here it is easier to record two separate robustness results. The first is robustness with respect to changes in the asymptotic height function, and the second is robustness with respect to changes in domain. Robustness with respect to changes in the asymptotic height function comes immediately from the definition of the balls $B(R_n, h_R, \varepsilon)$.

Lemma 6.6 (Robustness of microscopic entropy under change in profile). *Let $\varepsilon > 0$ and $n \in \mathbb{N}$. Let $R \subset \mathbb{R}^m$ satisfy Assumption 2.2, and let $R_n \subset \mathbb{Z}^m$ satisfy $\frac{1}{n}R_n \subset R$. Let $h_R, \tilde{h}_R \in M(R)$ be two asymptotic height functions such that $\max_{x \in R} |h_R(x) - \tilde{h}_R(x)| \leq \varepsilon$. Then,*

$$\text{Ent}_{R_n}(B(R_n, h_R, 2\varepsilon)) \leq \text{Ent}_{R_n}(B(R_n, \tilde{h}_R, \varepsilon)).$$

Proof. It suffices to notice that $B(R_n, h_R, 2\varepsilon) \supseteq B(R_n, \tilde{h}_R, \varepsilon)$. This follows from the triangle inequality: for any $h_{R_n} \in B(R_n, \tilde{h}_R, \varepsilon)$ and any $z \in R_n$,

$$\begin{aligned} \left| \frac{1}{n}h_{R_n}(z) - h_R\left(\frac{1}{n}z\right) \right| &\leq \left| \frac{1}{n}h_{R_n}(z) - \tilde{h}_R\left(\frac{1}{n}z\right) \right| + \left| \tilde{h}_R\left(\frac{1}{n}z\right) - h_R\left(\frac{1}{n}z\right) \right| \\ &\leq \varepsilon + \varepsilon. \end{aligned}$$

□

More care is needed to state and prove robustness of the microscopic entropy with respect to changes in domain. The main idea is straightforward. Given two microscopic domains $\tilde{R}_n \subset R_n$, we will consider the extension map from $B(\tilde{R}_n, h_R, \varepsilon)$ to $B(R_n, h_R, \varepsilon)$ and the restriction map in the opposite direction. So long as every height function on the smaller domain admits an extension, we have $|B(\tilde{R}_n, h_R, \varepsilon)| \leq |B(R_n, h_R, \varepsilon)|$. In the opposite direction, the restriction map is not generally an injection but the pre-images are not too large; at most 2^N height functions on R_n restrict to the any specific height function on \tilde{R}_n , where $N = |R_n| \setminus |\tilde{R}_n|$.

Most of the complications arise in the extension step. Our primary extension result, namely the Kirszbraun theorem (Theorem 3.1), is insufficient. It states that a height function $h_{\tilde{R}_n} \in B(\tilde{R}_n, h_R, \varepsilon)$ admits an extension to R_n , but that extension is not necessarily in $B(R_n, h_R, \varepsilon)$. There are two ways forward: to prove a stronger extension theorem specialized to the problem under consideration, or to leverage the Lipschitz property to control the extension. For greater generality, we prefer the second method. However there are a few difficulties: the Kirszbraun theorem is subtle when the asymptotic height profile has $|\nabla h_K|_\infty$ in part of the region, and the extension cannot generally be

kept within distance ε of h_R . This leads to the following somewhat complex formulation.

Lemma 6.7 (Robustness of microscopic entropy under domain approximations). *Let $\varepsilon \in (0, 1]$ and $n \in \mathbb{N}$ with $n \geq \frac{1}{\varepsilon}$. Let $\tilde{R} \subset R \subset \mathbb{R}^m$ and $\tilde{R}_n \subset R_n \subset \mathbb{Z}^m$ satisfy these assumptions:*

$$\begin{aligned} \frac{1}{n}R_n &\subset R, & \frac{1}{n}\tilde{R}_n &\subset \tilde{R}, \\ d_H(\frac{1}{n}R_n, R) &= \theta(\varepsilon), & d_H(\frac{1}{n}\tilde{R}_n, \tilde{R}) &= \theta(\varepsilon), \end{aligned}$$

Additionally, assume that

$$\frac{|R|}{|\tilde{R}|} = 1 + \theta(\varepsilon).$$

Let $h_R \in M(R)$ be an asymptotic height function with $\text{Lip}(h_R) \leq 1 - c\varepsilon$ for some fixed $c \in (0, 1]$. Then,

$$\begin{aligned} \text{Ent}_{\tilde{R}_n}(B(\tilde{R}_n, h_R, \varepsilon)) &+ \theta(\varepsilon) + \theta_\varepsilon(\frac{1}{n}) \\ &\leq \text{Ent}_{R_n}(B(R_n, h_R, \varepsilon)) \\ &\leq \text{Ent}_{\tilde{R}_n}(B(\tilde{R}_n, h_R, \frac{c}{3}\varepsilon^2)) + \theta(\varepsilon) + \theta_\varepsilon(\frac{1}{n}). \end{aligned}$$

Remark 6.8. *The assumptions in Lemma 6.7 quantify the imprecise statements that \tilde{R} , $\frac{1}{n}R_n$, and $\frac{1}{n}\tilde{R}_n$ respectively approximate R , R , and \tilde{R} from inside.*

If we take the simplicial approximation K from Lemma 6.1 to be \tilde{R} , and its discretization $K_n := \{z \in \mathbb{Z}^m \mid \frac{1}{n}z \in K\}$ to be \tilde{R}_n , and if we recall the Assumption 2.2 about R and R_n , then the hypotheses of Lemma 6.7 are satisfied, and moreover, we may replace all instance of $\theta(\varepsilon)$ in the conclusion by $\theta(\varepsilon\ell)$.

Proof. First, we prove the following inequalities:

$$|B(R_n, h_R, \varepsilon)| \leq (2^{|R_n|})^{\theta(\varepsilon) + \theta_\varepsilon(\frac{1}{n})} |B(\tilde{R}_n, h_R, \varepsilon)| \quad (36)$$

and

$$|B(R_n, h_R, \varepsilon)| \geq |B(\tilde{R}_n, h_R, \frac{c}{3}\varepsilon^2)|. \quad (37)$$

For (36), we note that every height function $h_{R_n} \in B(R_n, h_R, \varepsilon)$ restricts to a height function $h_{R_n}|_{\tilde{R}_n} \in B(\tilde{R}_n, h_R, \varepsilon)$. Then, we claim that

$$|R_n|n^{-d} = (1 + \theta(\varepsilon))|R| \quad (38)$$

and

$$|\tilde{R}_n|n^{-d} = (1 + \theta(\varepsilon))|\tilde{R}|. \quad (39)$$

To justify (38) we argue as follows. Consider the continuum region $R_n^\square := \bigcup_{z \in \mathbb{R}_n} ([0, \frac{1}{n}]^d + \frac{1}{n}z)$, i.e. the union of hypercubes of side length $\frac{1}{n}$ translated by the points in $\frac{1}{n}R_n$. Clearly R_n^\square has Lebesgue measure equal to $|R_n|n^{-d}$, and (like $\frac{1}{n}R_n$) satisfies $d_H(R_n^\square, R) = \theta(\varepsilon)$. This implies (38). Equation (39) is analogous. Further arithmetic yields the equation

$$\frac{|R_n|}{|\tilde{R}_n|} = (1 + \theta(\varepsilon)) \left(1 + \theta_\varepsilon\left(\frac{1}{n}\right)\right), \quad (40)$$

and then

$$|R_n \setminus \tilde{R}_n| = |R_n|(\theta(\varepsilon) + \theta_\varepsilon(\frac{1}{n})).$$

Therefore the restriction map from $B(R_n, h_R, \varepsilon)$ to $B(\tilde{R}_n, h_R, \varepsilon)$ is at most $(2^{|R_n|})^{\theta(\varepsilon) + \theta_\varepsilon(\frac{1}{n})}$ -to-1. Inequality (36) follows immediately.

Now let us turn to (37). We want an injection from $B(\tilde{R}_n, h_R, \frac{c}{3}\varepsilon^2)$ into $B(R_n, h_R, \varepsilon)$. Fix a function $h_{\tilde{R}_n} \in B(\tilde{R}_n, h_R, \frac{c}{3}\varepsilon^2)$; we will construct an extension $h_{R_n} \in M(R_n)$.

Let

$$R'_n := \{z \in R_n \mid d(z, \tilde{R}_n) > \frac{\varepsilon}{3}n\},$$

where

$$d(z, \tilde{R}_n) := \min_{\tilde{z} \in \tilde{R}_n} |z - \tilde{z}|_1.$$

For $z \in R'_n$, we arrange for the extension to satisfy $|h_{R_n}(z) - nh_R(\frac{1}{n}z)| \leq 1$. When $nh_R(\frac{1}{n}z)$ is not an integer, or is an integer but has the same parity as z , this inequality uniquely determines the value of $h_{R_n}(z)$. In the remaining case, there are two candidate values; we arbitrarily choose to “round down” to the lower value. Later it is important that we consistently round down (or up).

Let us check the hypotheses of the Kirszbraun theorem. If $\tilde{z} \in \tilde{R}_n$ and if $z \in R'_n$, then $|\tilde{z} - z|_1 > \frac{\varepsilon}{3}n$. Therefore

$$\begin{aligned} & |h_{\tilde{R}_n}(\tilde{z}) - h_{R_n}(z)| \\ & \leq |h_{\tilde{R}_n}(\tilde{z}) - nh_R(\frac{1}{n}\tilde{z})| + n|h_R(\frac{1}{n}\tilde{z}) - h_R(\frac{1}{n}z)| \\ & \quad + |nh_R(\frac{1}{n}z) - h_{R_n}(z)| \\ & \leq \frac{c}{3}\varepsilon^2n + (1 - c\varepsilon)|\tilde{z} - z|_1 + 1 \\ & < c\varepsilon|\tilde{z} - z|_1 + (1 - c\varepsilon)|\tilde{z} - z|_1 + 1 \\ & < |\tilde{z} - z|_1 + 1 \\ & \leq |\tilde{z} - z|_1. \end{aligned}$$

The argument for points $z_1, z_2 \in R'_n$ is similar to the arguments made in Section 2.2. By the triangle inequality, $|h_{R_n}(z_1) - h_{R_n}(z_2)| \leq$

$|z_1 - z_2|_1 + 2$. Equality holds only if both $nh_R(\frac{1}{n}z_1)$ and $nh_R(\frac{1}{n}z_2)$ are integers of the same parity as z_1 and z_2 , respectively. In this case h_{R_n} is rounded down at both points, so the Kirszbraun inequality is still satisfied.

So, there exists an extension h_{R_n} of $h_{\tilde{R}_n}$ such that $|h_{R_n}(z) - h_R(\frac{1}{n}z)| \leq 1$ for $z \in R'_n$. We claim that $h_{R_n} \in B(R_n, h_R, \varepsilon)$. Since $\frac{c}{3}\varepsilon^2 \leq \varepsilon$ and since $1 \leq \varepsilon n$, it suffices to consider points $z \in R_n \setminus R'_n$. Fix such a z . By the definition of R'_n , there exists $\tilde{z} \in \tilde{R}_n$ such that $|\tilde{z} - z| \leq \frac{\varepsilon}{3}n$. Note that $\frac{c}{3}\varepsilon^2 \leq \frac{\varepsilon}{3}$, since $c, \varepsilon \leq 1$. By the Lipschitz property of h_R and h_{R_n} ,

$$\begin{aligned} h_{R_n}(z) &\leq h_{R_n}(\tilde{z}) + \frac{\varepsilon}{3}n \\ &= h_{\tilde{R}_n}(\tilde{z}) + \frac{\varepsilon}{3}n \\ &\leq nh_R(\frac{1}{n}\tilde{z}) + \frac{\varepsilon}{3}n + \frac{c}{3}\varepsilon^2n \\ &\leq nh_R(\frac{1}{n}\tilde{z}) + \frac{2\varepsilon}{3}n \\ &\leq nh_R(\frac{1}{n}z) + \varepsilon n. \end{aligned}$$

By symmetry, $h_{R_n}(z) \geq nh_R(\frac{1}{n}z) - \varepsilon n$, and so $h_{R_n} \in B(R_n, h_R, \varepsilon)$. This extension process defines an injection from $B(\tilde{R}_n, h_R, \frac{c}{3}\varepsilon^2)$ into $B(R_n, h_R, \varepsilon)$, which proves (37).

Finally, we derive the conclusion from (36) and (37) by taking logarithms and normalizing, using (40) to account for the difference in normalizing factors $-\frac{1}{|R_n|}$ and $-\frac{1}{|\tilde{R}_n|}$. \square

Now, let us prove the profile theorem (Theorem 2.15). The main idea is straightforward: we approximate h_R by a piecewise affine function (given by the simplicial Rademacher theorem, i.e. Lemma 6.1), for which we have already proven the simplicial profile theorem (Theorem 5.4). Then we use the robustness results (Lemma 6.5, Lemma 6.6, and Lemma 6.6) to deduce the profile theorem for h_R . However, in order to apply Lemma 6.6 we must first reduce to the case where the Lipschitz constant $\text{Lip}(h_R) := \inf\{\lambda > 0 \mid h_R \text{ is } \lambda\text{-Lipschitz}\}$ is strictly less than 1.

Proof of Theorem 2.15. First we reduce to the case where $\text{Lip}(h_R) \leq 1 - c\delta$, for a constant $c > 0$ depending only on the domain R . Then, we reduce to the piecewise affine case of Theorem 5.4.

Reduction to $\text{Lip}(h_R) \leq 1 - c\delta$. By translation invariance, we may assume that there exists $x_0 \in R$ with $h_R(x_0) = 0$. Set $c = \frac{1}{2 \text{diam}_1 R} \wedge 1$, and define

$$\tilde{h}_R := (1 - c\delta)h_R.$$

Then $\text{Lip}(\tilde{h}_R) \leq 1 - c\delta$, and for all $x \in R$, both $|h_R(x) - \tilde{h}_R(x)| \leq \frac{\delta}{2}$ and $|\nabla h_R(x) - \nabla \tilde{h}_R(x)| \leq c\delta$. Assume that the conclusion holds for \tilde{h}_R , i.e.

$$\text{Ent}_R(\tilde{h}_R) = \text{Ent}_{R_n}(B(R_n, \tilde{h}_R, \delta)) + \theta(\delta) + \theta_\delta(\frac{1}{n}). \quad (41)$$

We make two calculations. First,

$$\begin{aligned} \text{Ent}_R(h_R) &\leq \text{Ent}_R(\tilde{h}_R) + \theta(\delta) && \text{(by Lemma 6.5)} \\ &\leq \text{Ent}_{R_n}(B(R_n, \tilde{h}_R, 2\delta)) + \theta(\delta) + \theta_\delta(\frac{1}{n}) && \text{(by (41))} \\ &\leq \text{Ent}_{R_n}(B(R_n, h_R, \delta)) + \theta(\delta) + \theta_\delta(\frac{1}{n}) && \text{(by Lemma 6.6).} \end{aligned}$$

Second,

$$\begin{aligned} \text{Ent}_R(h_R) &\geq \text{Ent}_R(\tilde{h}_R) + \theta(\delta) && \text{(by Lemma 6.5)} \\ &\geq \text{Ent}_{R_n}(B(R_n, \tilde{h}_R, \frac{1}{2}\delta)) + \theta(\delta) + \theta_\delta(\frac{1}{n}) && \text{(by (41))} \\ &\geq \text{Ent}_{R_n}(B(R_n, h_R, \delta)) + \theta(\delta) + \theta_\delta(\frac{1}{n}) && \text{(by Lemma 6.6).} \end{aligned}$$

So once we prove (41) with the extra hypothesis that $\text{Lip}(\tilde{h}_R) \leq 1 - c\delta$, the general result follows.

Reduction to piecewise linear height functions. We will apply Lemma 6.1 to derive a simplex domain K and a piecewise linear height function h_K approximating R and h_R , then appeal to Theorem 5.4. In so doing we introduce two parameters: ε , which controls how well K approximates R , and ℓ , which controls the size of the simplices in K . There are a few important properties of ε and ℓ . First, $\delta = \varepsilon\ell$, so there is actually only one degree of freedom. Second, ℓ must be chosen to be sufficiently small, as is required by the simplicial Rademacher theorem (see Lemma 6.1). Third, as $\delta \rightarrow 0$ we must have $\varepsilon \rightarrow 0$, so that $\theta(\varepsilon) = \theta(\delta)$.

Let us describe explicitly how we choose ε and ℓ satisfying these constraints. We fix a sequence $\varepsilon_k \downarrow 0$ (e.g. $\varepsilon_k = \frac{1}{k}$), and for each k set

$$\ell_k := \frac{1}{2} \sup\{\ell > 0 \mid \text{Lemma 6.1 applies with } \varepsilon = \varepsilon_k\}.$$

We call attention to the fact that Lemma 6.1 is monotone in ℓ . In particular, for any $\ell \leq \ell_k$, the conclusion of the lemma holds for (ε_k, ℓ) . Now, let

$$\delta_k := \varepsilon_k \ell_k \wedge \frac{1}{2} \delta_{k-1}.$$

The sequence δ_k decreases to 0, so $\bigcup_{k=1}^{\infty} (\delta_{k+1}, \delta_k]$ is a non-trivial half-open interval with left endpoint at 0. We assume that δ lies in this interval. Fix k such that $\delta \in (\delta_{k+1}, \delta_k]$, and set $\varepsilon = \varepsilon_k$ and $\ell = \frac{\delta}{\varepsilon}$. Then

$\delta = \varepsilon \ell$ by construction, and as noted above ℓ is small enough that the simplicial Rademacher theorem (see Lemma 6.1) applies. As $\delta \rightarrow 0$ necessarily $k \rightarrow \infty$, so $\varepsilon \rightarrow 0$ as desired. Therefore this choice of ε and ℓ satisfies our criteria. Per Lemma 6.1, the corresponding simplex domain K and piecewise affine asymptotic height function h_K satisfy

$$|R \setminus K| < \varepsilon \quad \text{and} \quad d_H(K, R) < \varepsilon, \quad (42)$$

$$\max_{x \in K} |h_K(x) - h_R(x)| < \varepsilon \ell = \delta \quad (43)$$

and

$$\frac{1}{|K|} \{x \in K \mid |\nabla h_K(x) - \nabla h_R(x)|_2 \geq \varepsilon\} < \varepsilon. \quad (44)$$

All that is left is to apply the simplicial profile theorem (see Theorem 5.4) and the robustness results (see Lemma 6.5, Lemma 6.6, and Lemma 6.6). In one direction, we have

$$\begin{aligned} \text{Ent}_R(h_R) &\leq \text{Ent}_K(h_K) + \theta(\varepsilon) && (\text{Lemma 6.5 and (44)}) \\ &\leq \text{Ent}_{K_n}(B(K_n, h_K, \varepsilon \ell)) && (\text{Theorem 5.4}) \\ &\quad + \theta(\varepsilon) + \theta_{\varepsilon, \ell}(\tfrac{1}{n}) \\ &\leq \text{Ent}_{K_n}(B(K_n, h_R, \tfrac{1}{2}\varepsilon \ell)) && (\text{Lemma 6.6 and (43)}) \\ &\quad + \theta(\varepsilon) + \theta_{\varepsilon, \ell}(\tfrac{1}{n}) \\ &\leq \text{Ent}_{R_n}(B(R_n, h_R, \tfrac{1}{2}\varepsilon \ell)) && (\text{Lemma 6.7 and (42)}) \\ &\quad + \theta(\varepsilon) + \theta_{\varepsilon, \ell}(\tfrac{1}{n}) \\ &= \text{Ent}_{R_n}(B(R_n, h_R, \tfrac{1}{2}\delta)) && (\text{choice of } \varepsilon \text{ and } \ell) \\ &\quad + \theta(\delta) + \theta_\delta(\tfrac{1}{n}). \end{aligned}$$

By taking $\delta' = \frac{1}{2}\delta$, this yields

$$\text{Ent}_R(h_R) \leq \text{Ent } R_n(B(R_n, h_R, \delta')) + \theta(\delta') + \theta_{\delta'}(\tfrac{1}{n}). \quad (45)$$

In the other direction,

$$\begin{aligned}
\text{Ent}_R(h_R) &\geq \text{Ent}_K(h_K) + \theta(\varepsilon) && (\text{Lemma 6.5 and (44)}) \\
&\geq \text{Ent}_{K_n}(B(K_n, h_K, \varepsilon\ell)) && (\text{Theorem 5.4}) \\
&\quad + \theta(\varepsilon) + \theta_{\varepsilon,\ell}(\tfrac{1}{n}) \\
&\geq \text{Ent}_{K_n}(B(K_n, h_R, 2\varepsilon\ell)) && (\text{Lemma 6.6 and (43)}) \\
&\quad + \theta(\varepsilon) + \theta_{\varepsilon,\ell}(\tfrac{1}{n}) \\
&\geq \text{Ent}_{R_n}(B(R_n, h_R, \tfrac{6}{c}(\varepsilon\ell)^{1/2})) && (\text{Lemma 6.7 and (42)}) \\
&\quad + \theta(\varepsilon) + \theta_{\varepsilon,\ell}(\tfrac{1}{n}) \\
&= \text{Ent}_{R_n}(B(R_n, h_R, \tfrac{6}{c}\delta^{1/2})) && (\text{choice of } \varepsilon \text{ and } \ell) \\
&\quad + \theta(\delta) + \theta_\delta(\tfrac{1}{n}).
\end{aligned}$$

By taking $\delta'' = \frac{6}{c}\delta^{1/2}$, this yields

$$\text{Ent}_R(h_R) \leq \text{Ent}_{R_n}(B(R_n, h_R, \delta'')) + \theta(\delta') + \theta_{\delta''}(\tfrac{1}{n}). \quad (46)$$

Combining (45) and (46) completes the proof of the profile theorem. \square

7. PROOF OF THE VARIATIONAL PRINCIPLE

Besides the profile theorem (Theorem 2.15), the proof of the variational principle (Theorem 2.16) relies on compactness of the space of asymptotic height functions. For robustness, we give a proof that does not assume that the macroscopic entropy functional admits a minimum. Note that the existence of such a minimizer is standard as soon as the local surface tension is convex and bounded below; see for example Section 2 of [CKP01] or [She05]. However, for greater generality we work with the infimum of the macroscopic entropy and we do not assume that a minimizer exists. At any rate, it will be necessary to deal with infima (rather than minima) later when proving the large deviations principle.

Proof of Theorem 2.16. First, we shall prove that

$$\inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) \geq \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) + \theta(\delta) + \theta_\delta(\tfrac{1}{n}) \quad (47)$$

via undercounting the number of height functions in $M(R_n, h_{\partial R_n}, \delta)$. The strategy is simple: we only count those height functions that are close to a “near-minimizer” of the macroscopic entropy. If we assume

that a minimizer exists, i.e. that there exists $h_R^{\min} \in M(R, h_{\partial R})$ such that

$$\text{Ent}_R(h_R^{\min}) = \inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R),$$

then the following proof suffices. For any $\delta > 0$ and $n \in \mathbb{N}$, the Definition 2.5 implies that

$$B(R_n, h_R^{\min}, \delta) \subseteq M(R_n, h_{\partial R_n}, 2\delta).$$

It follows immediately that

$$\text{Ent}_{R_n}(B(R_n, h_R^{\min}, \delta)) \geq \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, 2\delta)),$$

so after applying the profile theorem and replacing 2δ by δ ,

$$\text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) \leq \inf_{h \in M(R, h_{\partial R})} \text{Ent}_R(h) + \theta(\delta) + \theta_\delta(\frac{1}{n}).$$

However, as mentioned above we want to give a proof that does not rely on the existence of a minimizer. This idea is also important for proving the large deviation principle below (see the paragraphs following (56) below). The first step is to replace h_R^{\min} by a sequence of approximations, say $h_R^{(k)}$ satisfying

$$\text{Ent}_R(h_R^{(k)}) \leq \inf_{h_R \in M(R, h_{\partial R})} \text{Ent}(h_R) + \frac{1}{k} \quad \text{for } k \in \mathbb{N}.$$

Also let $\theta^{(k)}(\delta)$ and $\theta_\delta^{(k)}(\frac{1}{n})$ denote the θ terms from the profile theorem (Theorem 2.15) for the height function $h_R^{(k)}$. At this point one may be tempted to simply take the limit $k \rightarrow \infty$ for fixed δ and n . The problem is that the sequence $\theta^{(k)}(\delta)$ is not necessarily controlled as k goes to infinity, and could in general diverge for any fixed $\delta > 0$, and likewise for $\theta_\delta^{(k)}(\frac{1}{n})$. To correct this, we proceed as follows.

Let $\delta_0 = +\infty$ and $n_0 = 0$. For $k \geq 1$, choose δ_k such that

$$0 < \delta_k \leq \frac{1}{2}\delta_{k-1} \quad \text{and} \quad \theta^{(k)}(\delta_k) \leq \frac{1}{k}.$$

Now let $\delta > 0$ and $n \in \mathbb{N}$ be given. Fix k such that $\delta \in (\delta_k, \delta_{k-1}]$. Note that this is possible since $\delta_k \leq \frac{1}{2}\delta_{k-1}$ forces $\delta_k \rightarrow 0$ as $k \rightarrow \infty$. Since $\delta_k < \delta$, we have

$$\text{Ent}_{R_n}(B(R_n, h_{\partial R_n}, \delta)) \leq \text{Ent}_{R_n}(B(R_n, h_{\partial R_n}, \delta_k)).$$

By the profile theorem (Theorem 2.15) applied to $h_R^{(k)}$,

$$\text{Ent}_{R_n}(B(R_n, h_{\partial R_n}, \delta_k)) \leq \text{Ent}_R(h_R^{(k)}) + \theta^{(k)}(\delta_k) + \theta_{\delta_k}^{(k)}(\frac{1}{n}).$$

By choice of $h_R^{(k)}$, we have

$$\text{Ent}_R(h_R^{(k)}) \leq \inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) + \frac{1}{k}.$$

By choice of δ_k ,

$$\theta^{(k)}(\delta_k) \leq \frac{1}{k},$$

Finally, since k and δ_k are determined from δ ,

$$\theta_{\delta_k}^{(k)}\left(\frac{1}{n}\right) = \theta_\delta\left(\frac{1}{n}\right) \quad \text{and} \quad \frac{1}{k} = \theta(\delta).$$

Putting it all together, we have

$$\begin{aligned} \text{Ent}_{R_n}(B(R_n, h_{\partial R_n}, \delta)) &\leq \inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) + \frac{2}{k} + \theta_{\delta_k}^{(k)}\left(\frac{1}{n}\right) \\ &= \inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) + \theta(\delta) + \theta_\delta\left(\frac{1}{n}\right). \end{aligned}$$

Now we prove the reverse inequality, namely

$$\inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) \leq \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) + \theta(\delta) + \theta_\delta\left(\frac{1}{n}\right). \quad (48)$$

Let $\varepsilon > 0$. For each $h_R \in M(R, h_{\partial R}, 2\delta)$, by the profile theorem (Theorem 2.15)

$$\left| \text{Ent}_{R_n}(B(R_n, h_R, \delta)) - \text{Ent}_R(h_R) \right| < \theta(\delta) + \theta_\delta\left(\frac{1}{n}\right). \quad (49)$$

For each $h_R \in M(R, h_{\partial R})$, fix $\eta(h_R) > 0$ such that the $\theta(\delta)$ term in (49) satisfies

$$\theta(\eta(h_R)) \leq \frac{\varepsilon}{2}. \quad (50)$$

Recall from Definition 2.6 that

$$\begin{aligned} M(R, h_{\partial R}, 2\delta) \\ := \{h_R \in \text{Lip}(R) \mid \forall x \in \partial R, |h(x) - h_{\partial R}(x)| \leq 2\delta\}. \end{aligned}$$

This set is compact as an easy consequence of the Arzelà–Ascoli theorem. Choose $h_R^{(1)} \in M(R, h_{\partial R})$ such that

$$\text{Ent}_R(h_R^{(1)}) \leq \inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) + \varepsilon,$$

and pick $h_R^{(2)}, \dots, h_R^{(k)}$ so that the union $\bigcup_{i=1}^k B(R, h_R^{(i)}, \eta(h_R^{(i)}))$ covers $M(R, h_{\partial R}, 2\delta)$, where $B(R, h_R^{(i)}, \eta(h_R^{(i)}))$ is the set of asymptotic height functions $h_R \in M(R)$ that are uniformly within distance $\eta(h_R^{(i)})$ of $h_R^{(i)}$. Note that the number k of sets in this cover depends only on δ . We

abbreviate $\eta_i := \eta(h_R^{(i)})$. Moreover, we fix $n_i \in \mathbb{N}$ such that for all $n \geq n_i$, the $\theta_\delta(\frac{1}{n})$ from (49) satisfies

$$\theta_{\eta_i}(\frac{1}{n}) \leq \varepsilon. \quad (51)$$

We use the cover of the set $M(R, h_{\partial R}, 2\delta)$ to cover the set of height functions $M(R_n, h_{\partial R_n}, \delta)$. Indeed, consider an arbitrary height function $h_{R_n} \in M(R_n, h_{\partial R_n}, \delta)$. After rescaling and interpolating (via the classical Kirszbraun theorem), we identify h_{R_n} with a continuous function in $M(R, h_{\partial R}, 2\delta)$. Under this identification,

$$M(R_n, h_{\partial R_n}, \delta) \subseteq M(R, h_{\partial R}, 2\delta) \subseteq \bigcup_{i=1}^k B(R, h_R^{(i)}, \eta_i).$$

This means that for any discrete height function $h_{R_n} \in M(R_n, h_{\partial R_n}, \delta)$ with continuous (rescaled) interpolation $\tilde{h}_{R_n} \in M(R, h_{\partial R}, 2\delta)$ (note that δ increases to 2δ from discretization errors), there is $i \in \{1, \dots, k\}$ such that $\sup_{x \in R} |\tilde{h}_{R_n}(x) - h_R^{(i)}(x)| < \eta_i$. By Definition 2.5, it follows that

$$M(R_n, h_{\partial R_n}, \delta) \subseteq \bigcup_{i=1}^k B(R_n, h_R^{(i)}, \eta_i). \quad (52)$$

Hence,

$$\begin{aligned} \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) &\geq \text{Ent}_{R_n}\left(\bigcup_{i=1}^k B(R_n, h_R^{(i)}, \eta_i)\right) \\ &\geq -\frac{1}{|R_n|} \ln \left(\sum_{i=1}^k |B(R_n, h_R^{(i)}, \eta_i)| \right). \end{aligned} \quad (53)$$

Let us estimate $|B(R_n, h_R^{(i)}, \eta_i)|$. Assuming that n is larger than the constants n_1, \dots, n_k , then for all $i = 1, \dots, k$,

$$\begin{aligned} \text{Ent}_{R_n}(B(R_n, h_R^{(i)}, \eta_i)) &\geq \text{Ent}_R(h_R^{(i)}) - 2\varepsilon \quad (\text{By (49), (50), and (51)}) \\ &\geq \inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) - 2\varepsilon \\ &\geq \text{Ent}_R(h_R^{(1)}) - 3\varepsilon \\ &\geq \text{Ent}_{R_n}(B(R_n, h_R^{(1)}, \eta_1)) - 5\varepsilon \quad (\text{By (49), (50), and (51)}). \end{aligned}$$

In other words,

$$|B(R_n, h_R^{(i)}, \eta_i)| \leq |B(R_n, h_R^{(1)}, \eta_1)| e^{5\varepsilon |R_n|}.$$

We apply this last estimate in (53) to derive

$$\begin{aligned}
\text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) &\geq -\frac{1}{|R_n|} \ln \left(k |B(R_n, h_R^{(1)}, \eta_1)| e^{5\varepsilon |R_n|} \right) \\
&= -\frac{1}{|R_n|} \ln |B(R_n, h_R^{(1)}, \eta_1)| - \frac{\ln k}{|R_n|} - 5\varepsilon \\
&= -\frac{1}{|R_n|} \ln |B(R_n, h_R^{(1)}, \eta_1)| - \theta_\delta(\tfrac{1}{n}) - 5\varepsilon \\
&= -\inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R) - \theta_\delta(\tfrac{1}{n}) - 6\varepsilon.
\end{aligned} \tag{54}$$

Here, note that since k depends only on δ , $\frac{k}{|R_n|} = \theta_\delta(\frac{1}{n})$. Because $\varepsilon > 0$ was arbitrary, this yields the desired estimate (48). \square

8. LARGE DEVIATIONS PRINCIPLE

In this section we prove Theorem 2.17, the large deviations principle. For the reader's convenience, we recall the following definitions from the statement of the theorem in Section 2.5. For $\delta > 0$, $n \in \mathbb{N}$, and $h_R \in M(R)$:

$$\begin{aligned}
\mu_{\delta, n} &:= \frac{1}{|M(R_n, h_{\partial R_n}, \delta)|} |\{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\}|, \\
r_{\delta, n} &:= |R_n|, \\
I(h_R) &:= \begin{cases} \text{Ent}_R(h_R) - E & \text{if } h_R|_{\partial R} = h_{\partial R}, \\ \infty & \text{otherwise,} \end{cases}
\end{aligned}$$

where $E = \inf_{h_R \in M(R, h_{\partial R})} \text{Ent}_R(h_R)$.

The proof of the large deviations principle that we give here is based on the proof of the variational principle, Theorem 2.16, given in Section 7. We encourage the reader to read Section 7 first.

Proof of Theorem 2.17. First, we prove the LDP lower bound (14), i.e.

$$-\inf_{h_R \in A^\circ} I(h_R) \leq \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{r_{\delta, n}} \log \mu_{\delta, n}(A).$$

Without loss of generality we may assume that A is open. We may assume also that $\inf_{h_R \in A} I(h_R) < \infty$, or else (14) is trivial. By using these assumptions and replacing the symbols $\mu_{\delta, n}$, $r_{\delta, n}$, and $I(h_R)$ by

their definitions, (14) simplifies to

$$\begin{aligned} & - \inf_{h_R \in A} \left(\text{Ent}_R(h_R) - E \right) \\ & \leq \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \frac{|\{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\}|}{|M(R_n, h_{\partial R_n}, \delta)|}. \end{aligned}$$

Simplifying further by multiplying by -1 and using our definition of the microscopic entropy (9), it suffices to prove

$$\begin{aligned} & \inf_{h_R \in A} \text{Ent}_R(h_R) - E \\ & \geq \overline{\lim}_{\delta \rightarrow 0} \overline{\lim}_{n \rightarrow \infty} \text{Ent}_{R_n}(\{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\}) \quad (55) \\ & \quad - \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)). \end{aligned}$$

By the variational principle (Theorem 2.16),

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \text{Ent}_{R_n}(M(R_n, h_{\partial R_n}, \delta)) = E.$$

After cancelling the corresponding terms in (55), and after replacing $\overline{\lim}$ by our preferred θ asymptotics, it suffices to show that

$$\begin{aligned} \inf_{h_R \in A} \text{Ent}_R(h_R) & \geq \text{Ent}_{R_n}(\{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\}) \\ & \quad + \theta_A(\delta) + \theta_{A, \delta}(\tfrac{1}{n}). \end{aligned} \quad (56)$$

Note the analogy between (56) and inequality (47) from the proof of the variational principle. Indeed, we prove (56) in a similar manner to (47). We fix a sequence of asymptotic height function $h_R^{(k)} \in A$ that saturates the infimum; for concreteness, let us take

$$\text{Ent}_R(h_R^{(k)}) \leq \inf_{h_R \in A} \text{Ent}_R(h_R) + \frac{1}{k}.$$

Write $\theta^{(k)}(\delta)$ and $\theta_\delta^{(k)}(\frac{1}{n})$ for the error terms from the profile theorem for $h_R^{(k)}$. Choose a decreasing sequence δ_k such that $\delta_k \leq \frac{1}{2}\delta_{k-1}$ and such that $\theta^{(k)}(\delta_k) \leq \frac{1}{k}$. Given $\delta > 0$, choose k such that $\delta \in (\delta_k, \delta_{k-1}]$; then by the profile theorem applied to $h_R^{(k)}$,

$$\begin{aligned} & \text{Ent}_{R_n}(\{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\}) \\ & \leq \text{Ent}_{R_n}(B(R_n, h_R^{(k)}, \delta_k)) \\ & \leq \inf_{h_R \in A} \text{Ent}_R(h_R) + \frac{2}{k} + \theta_{\delta_k}^{(k)}(\tfrac{1}{n}) \\ & = \inf_{h_R \in A} \text{Ent}_R(h_R) + \theta(\delta) + \theta_\delta(\tfrac{1}{n}). \end{aligned}$$

Now, we turn to the LDP upper bound (15), i.e.

$$\overline{\lim}_{\delta \rightarrow 0} \overline{\lim}_{n \rightarrow \infty} \frac{1}{r_{\delta,n}} \log \mu_{\delta,n}(A) \leq - \inf_{h_R \in A} I(h_R),$$

We observe that $(\mu_{\delta,n})_{\delta,n}$ is exponentially tight, i.e. that for every $b \in (0, \infty)$, there exists $K_b \subset M(R)$ such that

$$\overline{\lim}_{\delta \rightarrow 0} \overline{\lim}_{n \rightarrow \infty} \frac{1}{r_{\delta,n}} \log \mu_{\delta,n}(K_b^c) \leq -b.$$

Indeed, we may take K_b to be the closure of $M(h_R, h_{\partial R}, 1)$, independent of b . For $\delta < \frac{1}{3}$ and n large enough that

$$\max_{z \in \partial R_n} \left| \frac{1}{n} h_{\partial R_n}(z) - h_{\partial R}\left(\frac{1}{n}z\right) \right| \leq \frac{1}{3},$$

any $h_{R_n} \in M(R_n, h_{\partial R}, \delta)$ satisfies $\tilde{h}_{R_n} \in M(h_R, h_{\partial R}, 1)$ by the triangle inequality, so $\mu_{\delta,n}(K_b^c) = 0$. By the general theory of large deviations, exponential tightness implies that it is sufficient to prove the upper bound (15) for compact sets $A \subset M(R)$.

If $\inf_{h_R \in A} I(h_R) = \infty$, then every height function in A differs from $h_{\partial R}$ at some point on the boundary. In fact by compactness, there exists δ_0 such that for every $h_R \in A$, $\sup_{x \in \partial R} |h_{\partial R}(x) - h_R(x)| \geq \delta_0$. Clearly, as in the proof of exponential tightness above, this implies that $\{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\}$ is empty once δ is small enough and n large enough. For all such δ, n we have $\mu_{\delta,n}(A) = 0$ and (15) follows.

It remains to prove the upper bound (15) when $\inf_{h_R \in A} I(h_R) < \infty$ and A is compact. Just like for the lower bound before, we reduce to proving the following inequality:

$$\begin{aligned} \inf_{h_R \in A} \text{Ent}_R(h_R) &\leq \text{Ent}_{R_n}(\{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\}) \\ &\quad + \theta_A(\delta) + \theta_{A,\delta}\left(\frac{1}{n}\right). \end{aligned}$$

We will closely follow the proof of (48) from Theorem 2.16. Let $\varepsilon > 0$, and choose $h_R^{(1)}$ such that

$$\text{Ent}_R(h_R^{(1)}) \leq \inf_{h_R \in A} \text{Ent}_R(h_R) + \varepsilon.$$

As in (52), choose $h_R^{(2)}, \dots, h_R^{(k)}$ such that

$$A \subset \bigcup_{i=1}^k B(R, h_R^{(i)}, \eta_i),$$

where η_1, \dots, η_k are chosen so that for each i , the $\theta(\delta)$ term from the profile theorem for $h_R^{(i)}$ satisfies $\theta(\eta_i) \leq \varepsilon$. Exactly as in the proof of

Theorem 2.16 (see (53)),

$$\begin{aligned} \text{Ent}_{R_n}(\{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\}) \\ \geq -\frac{1}{|R_n|} \ln \sum_{i=1}^k |B(R_n, h_R^{(i)}, \eta_i)|. \end{aligned}$$

From this we deduce the analogue of (54), namely

$$\begin{aligned} \text{Ent}_{R_n}(\{h_{R_n} \in M(R_n, h_{\partial R_n}, \delta) \mid \tilde{h}_{R_n} \in A\}) \\ \geq \text{Ent}_R(h_R^{(1)}) + \theta_{A,\delta}(\tfrac{1}{n}) - 5\varepsilon \\ \geq \inf_{h_R \in A} \text{Ent}_R(h_R) + \theta_{A,\delta}(\tfrac{1}{n}) - 6\varepsilon. \end{aligned}$$

□

ACKNOWLEDGMENT

The authors want to thank Tim Austin, Marek Biskup, Thomas Liggett, Igor Pak, Greta Panova for the fruitful discussions and helpful comments. This research has been partially supported by NSF grant DMS-1712632.

REFERENCES

- [BCG16] Alexei Borodin, Ivan Corwin, and Vadim Gorin. Stochastic six-vertex model. *Duke Math. J.*, 165(3):563–624, 2016.
- [CEP96] Henry Cohn, Noam Elkies, and James Propp. Local statistics for random domino tilings of the Aztec diamond. *Duke Math. J.*, 85(1):117–166, 1996.
- [Cer06] Raphaël Cerf. *The Wulff Crystal in Ising and Percolation Models*, volume 1878 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2006.
- [CKP01] Henry Cohn, Richard Kenyon, and James Propp. A variational principle for domino tilings. *J. Amer. Math. Soc.*, 14(2):297–346 (electronic), 2001.
- [CPST18] Nishant Chandgotia, Ron Peled, Scott Sheffield, and Martin Tassy. Delocalization of uniform graph homomorphisms from \mathbb{Z}^2 to \mathbb{Z} . *arXiv:1810.10124 [math.PR]*, 2018.
- [CPT18] Nishant Chandgotia, Igor Pak, and Martin Tassy. Kirszbraum-type theorems for graphs. *J. Comb. Theory Ser. B*, 2018.
- [CS16] Filippo Colomo and Andrea Sportiello. Arctic curves of the six-vertex model on generic domains: The tangent method. *J. Stat. Phys.*, 164(6):1488–1523, 2016.
- [Des98] Nicolas Destainville. Entropy and boundary conditions in random rhombus tilings. *J. Phys. A*, 31(29):6123–6139, 1998.
- [DGI00] Jean-Dominique Deuschel, Giambattista Giacomin, and Dmitry Ioffe. Large deviations and concentration properties for $\nabla\phi$ interface models. *Probab. Theory Relat. Fields*, 117(1):49–111, 2000.

- [DKS92] Roland Lvovich Dobrushin, Roman Kotecký, and Senya Shlosman. *Wulff Construction*, volume 104 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1992. A global shape from local interaction, Translated from the Russian by the authors.
- [DSS08] Daniela De Silva and Ovidiu Savin. Minimizers of convex functionals arising in random surfaces. *Duke Math. J.*, 151, 10 2008.
- [Dur10] Rick Durrett. *Probability: Theory and Examples*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, fourth edition, 2010.
- [DZ09] Amir Dembo and Ofer Zeitouni. *Large Deviations Techniques and Applications*. Stochastic Modelling and Applied Probability. Springer Berlin Heidelberg, 2009.
- [FO04] Tadahisa Funaki and Hirofumi Osada. *Stochastic Analysis on Large Scale Interacting Systems*. Advanced studies in pure mathematics. Mathematical Society of Japan, 2004.
- [FS06] Patrik L. Ferrari and Herbert Spohn. Scaling limit for the space-time covariance of the stationary totally asymmetric simple exclusion process. *Commun. Math. Phys.*, 265(1):1–44, 2006.
- [Kas63] Pieter W Kasteleyn. Dimer statistics and phase transitions. *J. Math. Phys.*, 4:287–293, 1963.
- [Kir34] Mojżesz D Kirszbraun. Über die zusammenziehende und Lipschitzsche Transformationen. *Fundam. Math.*, 22(1):77–108, 1934.
- [KK92] Claire Kenyon and Rick Kenyon. Tiling a polygon with rectangles. *Proc. of 33rd Fundamentals of Computer Science (FOCS)*, pages 610–619, 1992.
- [KMT17] Anndrew Krieger, Georg Menz, and Martin Tassy. A quenched variational principle for discrete random maps. *arXiv:1710.11330*, 2017.
- [KS99] Michael K.-H. Kiessling and Herbert Spohn. A note on the eigenvalue density of random matrices. *Commun. Math. Phys.*, 199(3):683–695, Jan 1999.
- [LP08] Lionel Levine and Yuval Peres. Strong spherical asymptotics for rotor-router aggregation and the divisible sandpile. *Potential Anal.*, 30(1):1, 2008.
- [LRS01] Michael Luby, Dana Randall, and Alistair Sinclair. Markov chain algorithms for planar lattice structures. *SIAM J. Comput.*, 31(1):167–192, 2001.
- [LS77] Benjamin F Logan and Larry A Shepp. A variational problem for random Young tableaux. *Adv. in Math.*, 26(2):206–222, 1977.
- [MT16] Georg Menz and Martin Tassy. A variational principle for a non-integrable model. *arXiv:1610.08103 [math.PR]*, 2016.
- [PR07] Boris Pittel and Dan Romik. Limit shapes for random square Young tableaux. *Adv. in Appl. Math.*, 38(2):164–209, 2007.
- [RAS15] Firas Rassoul-Agha and Timo Seppinen. *A Course on Large Deviations with an Introduction to Gibbs Measures*. American Mathematical Society, Providence, RI, 05 2015.
- [RS18] Nicolai Reshetikhin and Ananth Sridhar. Limit shapes of the stochastic six vertex model. *Commun. Math. Phys.*, 363(3):741–765, 2018.

- [Sch14] Jean Van Schaftingen. Approximation in Sobolev spaces by piecewise affine interpolation. *J. Math. Anal. Appl.*, 420(1):40–47, 2014.
- [She01] Scott Sheffield. Ribbon tilings and multidimensional height functions. *Trans. Am. Math. Soc.*, 354, 08 2001.
- [She05] Scott Sheffield. Random surfaces. *Astérisque*, (304):vi+175, 2005.
- [SS95] Lynn Arthur Steen and J Arthur Seebach. *Counterexamples in Topology*. Dover books on mathematics. Dover Publications, 1995.
- [Thu90] William P. Thurston. Conway’s tiling groups. *Am. Math. Monthly*, 97(8):757–756, September 1990.
- [vB77] Henk van Beijeren. Exactly solvable model for the roughening transition of a crystal surface. *Phys. Rev. Lett.*, 38(18), 1977.
- [VK77] Anatolii Moiseevich Vershik and Sergei V. Kerov. Asymptotic behavior of the Plancherel measure of the symmetric group and the limit form of Young tableaux. *Dokl. Akad. Nauk SSSR*, 233(6):1024–1027, 1977.
- [Wig59] Eugene Paul Wigner. Statistical properties of real symmetric matrices with many dimensions. In *4th Can. Math. Congress (Banff 1957)*, pages 174–184. Univ. Toronto Press, 1959.
- [Wil04] David Bruce Wilson. Mixing times of lozenge tiling and card shuffling Markov chains. *Ann. Appl. Probab.*, 14(1):274–325, 02 2004.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES
E-mail address: `akrieger@math.ucla.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES
E-mail address: `gmenz@math.ucla.edu`

DARTMOUTH COLLEGE, HANOVER
E-mail address: `mtassy@math.dartmouth.edu`