

SCHWINGER-BASED QCD FORMULATION'S DERIVATION OF ELASTIC PP SCATTERING

H.M. Fried, Y.M.Sheu, P.H.Tsang^a
Brown University, Department of Physics, RI, USA

Y. Gabellini, T.Grandou
Institut de Physique de Nice, UMR 7010 CNRS, Site Sophia 06560 Valbonne, France

Using previously described functional techniques for some non-perturbative, gauge invariant, renormalized QCD processes, a simplified version of the amplitudes - in which forms akin to Pomerons naturally appear - provides fits to ISR and LHC-TOTEM pp elastic scattering data. Those amplitudes rely on a specific function $\phi(b)$ which describes the fluctuations of the transverse position of quarks inside hadrons.

Presented at EDS Blois 2019
 The 18th Conference on Elastic and Diffractive Scattering
 Quy Nhon, Vietnam,
 June 2328, 2019

1 Introduction

This talk covers the work in ¹. Beginning with Schwinger's Generating Functional and the QCD Lagrangian, applying two procedures that were overlooked in the last four decades, an analytic, gauge-invariant correlation functions for Non-Perturbative QCD is obtained. This formulation produces the following results thus far:

- First-principled dynamical quark confining potential for quarks ²,
- A potential obtained from QCD that allows nucleons to be bounded, thus provided the first-principled model deuteron ³,
- New property of Effective Locality, provides gauge-invariant summation of all gluonic exchanges between quarks; more over, the interaction becomes a local interaction. ⁴.
- Obtained Chiral Symmetry Breaking for dynamical quarks out of the new property of Effective Locality ⁵.
- Extended Asymptotic freedom as supported by other non-perturbative approaches: Dyson Schwinger Equation ⁶.
- A qualitative description of the Hadron Confinement mass scale(s) ⁷.

^aspeaker

- The full SU(3) algebraic content of QCD amplitudes, both C_2 and C_3 casimir invariants are preserved⁸.
- First-principled calculation of elastic proton-proton scattering at ISR and LHC energies. This will be the focus of this talk¹.

1.1 Schwinger Generating Functional

The Schwinger Generating Functional can be rewritten into gaussian operations on gaussian fields.

$$\mathfrak{Z}_{QCD}[\bar{\eta}, \eta, j] = \mathfrak{N} e^{-\frac{i}{2} \int \frac{\partial}{\partial A} \cdot D_F^{(0)} \cdot \frac{\partial}{\partial A} \cdot e^{\frac{i}{4} \int \mathbf{F}^2 + \frac{i}{2} \int \mathbf{A} \cdot (-\partial^2) \cdot \mathbf{A}} \cdot e^{i \int \bar{\eta} \cdot \mathbf{G}_F[\mathbf{A}] \cdot \mathbf{J} + L[\mathbf{A}]} \Big|_{A=f D_F^{(0)} \cdot j} \quad (1)$$

where $G_F(x, y|A) = [m + \gamma \cdot (\delta - igA\tau)]^{-1}$, and $L[A] = \ln[1 - i\gamma A\tau[0]]$.
The \mathbf{F}^2 can be linearized with Halpern's half a century old expression:

$$e^{-\frac{i}{4} \int \mathbf{F}^2} = N \int d[\chi] e^{\frac{i}{4} \int \chi^2 + \frac{i}{2} \int F \cdot \chi} \quad (2)$$

where $\chi_{\mu\nu}^a = -\chi_{\nu\mu}^a$ is anti symmetric tensor. It is this added χ field that plays the central role in the summation of all gluons in the non-perturbative regime.

1.2 Gauge-Invariance

Schwinger's Generating Functional now becomes

$$\mathfrak{Z}_{QCD}[\bar{\eta}, \eta, j] = N \int d[\chi] e^{\frac{i}{4} \int \chi^2} e^{\mathfrak{D}_A^{(0)}} \cdot e^{\frac{i}{2} \int \chi \cdot \mathbf{F} + \frac{i}{2} \int \mathbf{A} \cdot (-\partial) \cdot \mathbf{A}} e^{i \int \bar{\eta} \cdot G_F[A] \cdot \eta + L[A]} \Big|_{A=f D_F^{(0)} \cdot j} \quad (3)$$

Calculating 2n-point Fermionic Green's functions (e.g. n=2), gives

$$= N \int d[\chi] e^{\frac{i}{4} \int \chi^2} e^{\mathfrak{D}_A^{(0)}} e^{\frac{i}{2} \int A \cdot (D_F^{(0)})^{-1} \cdot A} G_F(1|gA) G_F(2|gA) e^{L[A]} \Big|_{A=0} \quad (4)$$

$$e^{\mathfrak{D}_A} F_1[A] = \exp\left[\frac{i}{2} \int \bar{Q} \cdot D_F^{(0)} \cdot (1 - \bar{K} \cdot D_F^{(0)})^{-1} \cdot \bar{Q} - \frac{1}{2} \text{Tr} \ln(1 - D_F \cdot \tilde{K})\right] \cdot \exp\left[\frac{1}{2} \int A \cdot \bar{K} \cdot (1 - D_F^{(0)} \cdot \bar{K})^{-1} \cdot A\right] \quad (5)$$

where

$$D_F^{(0)} \cdot (1 - \bar{K} \cdot D_F^{(0)})^{-1} = D_F^{(0)} \cdot [1 - \hat{K} + (D_F^{(0)})^{-1}] \cdot D_F^{(0)} \cdot (1 - \hat{K})^{-1} = -(\tilde{K}_{\mu\nu}^{ab} + g f^{abc} \chi_{\mu\nu}^c)^{-1} = -\hat{\mathbf{K}}^{-1} \quad (6)$$

with $F_1[A] = e^{\frac{i}{2} \int A \cdot \bar{K} \cdot A + i \int \bar{Q} \cdot A}$, $F_2[A] = e^{L[A]}$ and $\langle z | \bar{K}_{\mu\nu}^{ab} | z' \rangle = [\tilde{\mathbf{K}}_{\mu\nu}^{ab}(z) + g f^{abc} \chi_{\mu\nu}^c(z)] \delta^{(4)}(z - z') + \langle z | (D_F^{(0)})^{-1} |_{\mu\nu}^{ab} | z' \rangle$.

We then have

$$\begin{aligned} e^{\mathfrak{D}_A} F_1[A] F_2[A] &= \exp\left[-\frac{i}{2} \int \bar{Q} \cdot \hat{\mathbf{K}}^{-1} \cdot \bar{Q}\right] + \frac{1}{2} \text{Tr} \ln \hat{\mathbf{K}} + \frac{1}{2} \text{Tr} \ln(-D_F^{(0)}) \\ &\cdot \exp\left[\frac{i}{2} \int \frac{\partial}{\partial A} \cdot D_F^{(0)} \cdot \frac{\partial}{\partial A'}\right] \\ &\cdot \exp\left[\frac{i}{2} \int \frac{\partial}{\partial A'} \cdot \hat{\mathbf{K}}^{-1} \cdot \frac{\partial}{\partial A'} - \int \bar{Q} \cdot \hat{\mathbf{K}}^{-1} \cdot \frac{\partial}{\partial A'}\right] \\ &\cdot (e^{\mathfrak{D}_A} F_2[A']) \end{aligned} \quad (7)$$

and

$$e^{\mathcal{D}A} F_1[A] F_2[A] = N \exp\left[-\frac{i}{2} \int \bar{Q} \cdot \hat{\mathbf{K}}^{-1} \cdot \bar{Q} + \frac{1}{2} \text{Tr} \ln \hat{\mathbf{K}}\right] \cdot \exp\left[\frac{i}{2} \int \frac{\partial}{\partial A} \cdot \hat{\mathbf{K}}^{-1} \cdot \frac{\partial}{\partial A} - \int \bar{Q} \cdot \hat{\mathbf{K}}^{-1} \cdot \frac{\partial}{\partial A}\right] \cdot \exp(L[A])|_{A \rightarrow 0} \quad (8)$$

From above, all explicit gauge dependencies are cancelled. That is, Gauge Invariance is explicitly preserved by means of Gauge Independence.

$-\hat{\mathbf{K}}^{-1} \equiv (f \cdot \chi)^{-1}$ represents all gluonic exchanges between quarks, a **Gluon Bundle**, ie, all gluonic exchanges summed in the non-perturbative regime.

In QCD, where confinement and chiral symmetry breaking hold, the impact parameter, b must fluctuate^{10 11}. At this stage, we choose a fluctuating b by a deformed gaussian (gaussian resulted in zero potentials).

$$\varphi(b) = \frac{\mu^2}{\pi} \frac{1 - \xi/2}{\Gamma(\frac{1}{1-\xi/2})} e^{-(\mu b)^{(2-\xi)}} \quad (9)$$

where $0 < \xi \ll 1$

With this, non-perturbative QCD processes becomes processes of gluon bundles, $(f \cdot \chi)^{-1}$'s and closed quark loops, $L[A]$'s. With the introduction of a particular renormalization scheme as described in⁹, functional integrals of Halpern's χ 's reduces to ordinary integrals, where $d\chi^4$ becomes space-time integrals of infinitesimal sizes, giving δ functions joining with closed quark loops, $L[A]$'s.

2 Comparing theory with experimental proton-proton elastic scattering differential cross section at ISR and LHC energies.

As shown in^{1, 9}, the choice for $\delta^2 \ell = \kappa/\bar{m}$ results explicit 0 for all graphs except Quark Loop Chain graphs.

The non-perturbative QCD scattering amplitude can thus be calculated as

$$T(s, \vec{q}) = \frac{is}{2M^2} \int d^2 b e^{i\vec{q} \cdot \vec{b}} [1 - e^{i\mathbb{X}^{pp}(s, \vec{b})}] \quad (10)$$

Where \mathbb{X}^{pp} is the Eikonal of the proton-proton process where $e^{i\mathbb{X}^{pp}} = e^{i\mathbb{X}^{(GluonBundle)}} e^{i\mathbb{X}^{(QuarkLoopChain)}}$

$$T(s, \vec{q}) = \frac{is}{2M^2} \int d^b e^{i\vec{q} \cdot \vec{b}} [1 - e^{\sqrt{i} g \delta_q^2 \varphi(\vec{b})/2} e^{g^2 \delta_q^2 (\kappa/\bar{m}^2) \Delta \bar{\varphi}/4}] \quad (11)$$

Expanding above we get for one gluon bundle plus one quark loop chain exchange:

$$T_1(s, \vec{q}) = \frac{s}{2M^2} \frac{g}{2} \left(\frac{\lambda}{m}\right)^2 \left(\frac{m}{E}\right)^{2p} \left[-\frac{1}{\sqrt{2}} e^{-q^2/4m^2} + i\left(\frac{1}{\sqrt{2}} e^{-q^2/4m^2} + \frac{g}{2} \kappa \frac{q^2}{\bar{m}^2} e^{-q^2/2\bar{m}^2}\right)\right] \quad (12)$$

giving a differential cross section, (green dotted curve in figures)

$$\frac{d\sigma_1}{dt}(s, q^2) = K \frac{27}{4\pi} \frac{g^2}{4} \left(\frac{\lambda}{4}\right)^4 \left(\frac{6m}{\sqrt{s}}\right)^{4p} \left[\frac{1}{2} e^{-q^2/2m^2} + \left(\frac{1}{\sqrt{2}} e^{-q^2/4m^2} + \frac{g}{2} \kappa \frac{q^2}{\bar{m}^2} e^{-q^2/2\bar{m}^2}\right)^2\right] \quad (13)$$

And for one gluon bundle plus two gluon bundles plus one quark loop chain exchange:

$$T_2(s, \vec{q}) = \frac{s}{2M^2} \frac{g}{2} \left(\frac{\lambda}{m}\right)^2 \left(\frac{m}{E}\right)^{2p} \left[\left(-\frac{1}{\sqrt{2}} e^{-q^2/4m^2} + \frac{1}{2} \frac{g}{2} \delta_q^2 \frac{m^2}{2\pi} e^{-q^2/8m^2}\right) + i\left(\frac{1}{\sqrt{2}} e^{-q^2/4m^2} + \frac{g}{2} \kappa \frac{q^2}{\bar{m}^2} e^{-q^2/2\bar{m}^2}\right) \right] \quad (14)$$




Figure 1 – ISR = 23.5 GeV




Figure 2 – ISR = 30.7 GeV




Figure 3 – ISR = 44.7 GeV




Figure 4 – ISR = 52.8 GeV




Figure 5 – ISR = 62.5 GeV

Figure 6 – ISR energies for elastic proton-proton scattering. Green dotted line for one Gluon Bundle and one Quark Loop Chain. Red solid line for One Gluon Bundle plus two Gluon Bundles and one Quark Loop Chain.

with differential cross section (red solid line in figures)

$$\begin{aligned} \frac{d\sigma_2}{dt}(s, q^2) = K \frac{27}{4\pi} \frac{g^2}{4} \left(\frac{\lambda}{m}\right)^4 \left(\frac{6m}{\sqrt{s}}\right)^{4p} & \left[\left(\frac{1}{2} e^{-q^2/2m^2} \right. \right. \\ & \left. \left. - \frac{g}{2} \frac{\lambda^2}{4\pi} \left(\frac{6m}{\sqrt{s}}\right)^{2p} e^{-q^2/8m^2} \right)^2 + \left(\frac{1}{\sqrt{2}} e^{-q^2/4m^2} + \frac{g}{2} \kappa \frac{q^2}{\bar{m}^2} e^{-q^2/2\bar{m}^2} \right)^2 \right] \end{aligned} \quad (15)$$

K is conversion factor from mb to GeV at $0.44mbGeV^{-2}$. For the ISR data, we obtained parameters: $g=7.0$, $p=0.13$, $\lambda = 0.5$, $\kappa = -6.810^{-4}$, $m = 0.23GeV \approx 1.5m_\pi$, $\bar{m} = 0.64GeV \approx 4.5m_\pi$.

For the LHC TOTEM data, we obtained $g = 7.0$, $p = 0.55$, $\lambda = 0.72$, $\kappa = -4.210^{-3}$, $m = 0.16GeV \approx m_\pi$, $\bar{m} = 0.41GeV \approx 3m_\pi$.

Acknowledgments

This work was made possible by a generous grant from the Julian Schwinger Foundation.

References

1. H.M.Fried, Y. Gabellini, T. Grandou, Y.M. Sheu, P.H. Tsang. arXiv:1904.11083
2. H.M.Fried et al. Ann. Phys. 327, 2666-2690 (2012) DOI: 10.1016/j.aop.2012.07.008
3. Fried et al. Ann. Phys. 338, 2013 Volume 338, November 2013, Pages 107-122. DOI:10.1016/j.aop.2013.07.006
4. T. Grandou et al. Ann. Phys. 327 (2012), Mod. Phys. Lett. A, Vol.32(2017); arXiv:1706.02264
5. T. Grandou, P.H.Tsang, arXiv:1905.05666 . to be published
6. H.M.Fried, T.Grandou, Y.M.Sheu, Annals of Physics 344, 78-96 (2014), DOI:10.1016/j.aop.2014.02.015
7. H.M.Fried, P.H. Tsang arXiv:1502.04378.
8. T.Grandou, EPL (Europhysics Letters), Volume 107, Number 1. Article 11001. DOI:doi.org/10.1209/0295-5075/107/11001
9. H.M.Fried, Y.Gabellini, T.Grandou, Y.M.Sheu, P.H.Tsang., Annals of Physics Volume 359, August 2015, Pages 1-19. DOI: 10.1016/j.aop.2015.03.024
10. A.Casher(1979) Chiral symmetry breaking in quark confining theories. Phys Lett B 83:395398.
11. S.J.Brodsky, R.Shrock. Proc. Nat. Acad. Sci. 108:45-50,2011. DOI: 10.1073/pnas.1010113107