

DIMENSION FORMULAE AND GENERALISED DEEP HOLES OF THE LEECH LATTICE VERTEX OPERATOR ALGEBRA

SVEN MÖLLER^{a,b} AND NILS R. SCHEITHAUER^c

Dedicated to the memory of John Horton Conway.

ABSTRACT. We prove a dimension formula for the weight-1 subspace of a vertex operator algebra $V^{\text{orb}(g)}$ obtained by orbifolding a strongly rational, holomorphic vertex operator algebra V of central charge 24 with a finite-order automorphism g . Based on an upper bound derived from this formula we introduce the notion of a generalised deep hole in $\text{Aut}(V)$.

Then we show that the orbifold construction defines a bijection between the generalised deep holes of the Leech lattice vertex operator algebra V_Λ with non-trivial fixed-point Lie subalgebra and the strongly rational, holomorphic vertex operator algebras of central charge 24 with non-vanishing weight-1 space. This provides a uniform construction of these vertex operator algebras and naturally generalises the correspondence between the deep holes of the Leech lattice Λ and the 23 Niemeier lattices with non-vanishing root system found by Conway, Parker, Sloane and Borcherds.

CONTENTS

1. Introduction	1
2. Lie Algebras	5
3. Vertex Operator Algebras	8
4. Modular Forms	13
5. Dimension Formulae	22
6. Generalised Deep Holes	30
Appendix. List of Generalised Deep Holes	42
References	48

1. INTRODUCTION

In 1968 Niemeier classified the positive-definite, even, unimodular lattices of rank 24 [Nie68, Nie73]. He showed that up to isomorphism there are exactly 24 such lattices and that the isomorphism class is uniquely determined by the root system. The Leech lattice Λ is the unique lattice in this genus without roots. There are several proofs of this result. Niemeier applied Kneser's neighbourhood method. Venkov found a proof based on harmonic theta series [Ven80]. It can also be derived from Conway, Parker and Sloane's classification of deep holes of the Leech lattice [CPS82, Bor85] and from the Smith-Minkowski-Siegel mass formula [CS82a, CS99]. Finally, it also follows from the classification of certain automorphic representations of O_{24} [CL19].

Conway, Parker and Sloane found a nice construction of the Niemeier lattices starting from the Leech lattice [CPS82, CS82b]. They showed that up to equivalence there are exactly 23 deep holes in the Leech lattice Λ , i.e. points in $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ which have maximal distance to the Leech lattice, and that they are in bijection with the Niemeier lattices different from the Leech lattice. The construction is as follows. Let d be a deep hole corresponding to the Niemeier lattice N . Then the $\mathbb{Z}$ -module in $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ generated by $\{x \in \Lambda \mid (x, d) \in \mathbb{Z}\}$ and d is isomorphic to N .

^aRutgers University, Piscataway, NJ, United States of America.

^bResearch Institute for Mathematical Sciences, Kyoto University, Kyoto, Japan.

^cTechnische Universität Darmstadt, Darmstadt, Germany.

The classification of strongly rational, holomorphic vertex operator algebras of central charge 24 bears similarities to the classification of positive-definite, even, unimodular lattices of rank 24. The weight-1 subspace V_1 of a strongly rational, holomorphic vertex operator algebra V of central charge 24 is a reductive Lie algebra. In 1993 Schellekens [Sch93] (see also [EMS20a]) showed that there are at most 71 possibilities for this Lie algebra using the theory of Jacobi forms. He conjectured that all potential Lie algebras are realised and that the V_1 -structure fixes the vertex operator algebra up to isomorphism. By the work of many authors over the past three decades the following result is now proved.

Theorem. *Up to isomorphism there are exactly 70 strongly rational, holomorphic vertex operator algebras V of central charge 24 with $V_1 \neq \{0\}$. Such a vertex operator algebra is uniquely determined by its V_1 -structure.*

The proof is based on a case-by-case analysis and uses a variety of methods. One of the main results of this paper is a uniform proof of the existence part which generalises Conway and Sloane's construction of the Niemeier lattices from the Leech lattice.

One method to construct vertex operator algebras is the orbifold construction [EMS20a]. Let V be a strongly rational, holomorphic vertex operator algebra and g an automorphism of V of finite order n and type 0. Then the fixed-point subalgebra V^g is a strongly rational vertex operator algebra. It has exactly n^2 non-isomorphic irreducible modules, which can be realised as the eigenspaces of g acting on the twisted modules $V(g^i)$ of V . If the twisted modules $V(g^i)$ have positive conformal weight for $i \neq 0 \bmod n$, then the sum $V^{\text{orb}(g)} := \bigoplus_{i \in \mathbb{Z}/n\mathbb{Z}} V(g^i)^g$ is a strongly rational, holomorphic vertex operator algebra. There is also an inverse orbifold construction, i.e. an automorphism h of $V^{\text{orb}(g)}$ such that $(V^{\text{orb}(g)})^{\text{orb}(h)} \cong V$.

We can use the deep holes of the Leech lattice to construct the vertex operator algebras corresponding to the Niemeier lattices.

Theorem (Proposition 6.2). *Let $d \in \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ be a deep hole in the Leech lattice Λ corresponding to the Niemeier lattice N . Then $g = e^{2\pi i d_0}$ is an automorphism of the vertex operator algebra V_{Λ} associated with Λ of order equal to the Coxeter number of N and type 0. The corresponding orbifold construction $V_{\Lambda}^{\text{orb}(g)}$ is isomorphic to the vertex operator algebra V_N associated with N .*

We shall see that the other elements on Schellekens' list can be constructed in a similar way.

Modular forms for the Weil representation of $\text{SL}_2(\mathbb{Z})$ play an important role in many areas of mathematics. The simplest examples are theta series. Let L be a positive-definite, even lattice of even rank $2k$ and $D = L'/L$ its discriminant form. Then $\theta(\tau) = \sum_{\gamma \in D} \theta_{\gamma}(\tau) e^{\gamma}$ with $\theta_{\gamma}(\tau) = \sum_{\alpha \in \gamma + L} q^{(\alpha, \alpha)/2}$ is a modular form of weight k for the Weil representation of D .

Another example comes from orbifold theory. Under the same conditions as above the n^2 characters of the irreducible modules of V^g combine to a modular form of weight 0 for the Weil representation of the hyperbolic lattice $II_{1,1}(n)$. Pairing this form with a certain Eisenstein series of weight 2 for the dual Weil representation we obtain:

Theorem (Theorem 5.3 and Corollary 5.7). *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and g an automorphism of V of finite order $n > 1$ and type 0. We assume that the twisted modules $V(g^i)$ have positive conformal weight for $i \neq 0 \bmod n$. Then*

$$\dim(V_1^{\text{orb}(g)}) = 24 + \sum_{m|n} c_n(m) \dim(V_1^{g^m}) - R(g)$$

where the $c_n(m)$ are defined by $\sum_{m|n} c_n(m)(t, m) = n/t$ for all $t \mid n$ and the rest term $R(g)$ is non-negative. In particular

$$\dim(V_1^{\text{orb}(g)}) \leq 24 + \sum_{m|n} c_n(m) \dim(V_1^{g^m}).$$

The rest term $R(g)$ is given explicitly. It depends on the dimensions of the weight spaces of the irreducible V^g -modules of weight less than 1.

The dimension formula was first proved by Montague for $n = 2, 3$ [Mon94], then generalised to $n = 5, 7, 13$ [Möl16] and finally to all $n > 1$ such that $\Gamma_0(n)$ has genus 0 in [EMS20b]. The previous proofs all used explicit expansions of the corresponding Hauptmoduln. We show here that the dimension formula is an obstruction coming from the Eisenstein space. Pairing the character of V^g with other modular forms for the dual Weil representation we can obtain further restrictions.

The above upper bound is our motivation for the following definition. Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and g an automorphism of V of finite order $n > 1$. Suppose g has type 0 and V^g satisfies the positivity condition. Then g is called a *generalised deep hole* of V if

- (1) the upper bound in the dimension formula is attained, i.e. $\dim(V_1^{\text{orb}(g)}) = 24 + \sum_{m|n} c_n(m) \dim(V_1^{g^m})$,
- (2) $\text{rk}(V_1^{\text{orb}(g)})$ is minimal, i.e. $\text{rk}(V_1^{\text{orb}(g)}) = \text{rk}(V_1^g)$.

By convention, we also call the identity a generalised deep hole.

We show that if $d \in \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ is a deep hole of the Leech lattice, then $g = e^{2\pi i d_0}$ is a generalised deep hole of the vertex operator algebra V_{Λ} .

The existence of generalised deep holes is restricted by Deligne's bound on the Fourier coefficients of cusp forms. Pairing the character of V^g with a certain cusp form we obtain:

Theorem (Theorem 5.10). *In the above situation suppose that the order of g is a prime p such that $\Gamma_0(p)$ has positive genus. Then $R(g) \geq 24$. In particular, g is not a generalised deep hole.*

Using inverse orbifolding and an averaged version of Kac's very strange formula [Kac90] we show (recall that algebraic conjugacy means conjugacy of cyclic subgroups):

Theorem (Theorem 6.6). *The cyclic orbifold construction $g \mapsto V_{\Lambda}^{\text{orb}(g)}$ defines a bijection between the algebraic conjugacy classes of generalised deep holes g in $\text{Aut}(V_{\Lambda})$ with $\text{rk}((V_{\Lambda}^g)_1) > 0$ and the isomorphism classes of strongly rational, holomorphic vertex operator algebras V of central charge 24 with $V_1 \neq \{0\}$.*

Then, we give a uniform construction of the 70 non-trivial Lie structures on Schellekens' list:

Theorem (Theorem 6.7). *Let $\mathfrak{g}$ be one of the 70 non-zero Lie algebras on Schellekens' list (Table 1 in [Sch93]). Then there exists a generalised deep hole $g \in \text{Aut}(V_{\Lambda})$ such that $(V_{\Lambda}^{\text{orb}(g)})_1 \cong \mathfrak{g}$.*

We explicitly give a generalised deep hole for each case. In order to determine the V_1 -structure of the corresponding orbifold construction we use Schellekens' classification of the possible V_1 -structures and Kac's theory of finite-order automorphisms of simple Lie algebras [Kac90].

Our results generalise the holy construction of the Niemeier lattices with non-trivial root system by Conway, Parker, Sloane and Borchers.

We remark that another uniform construction of the vertex operator algebras on Schellekens' list with non-trivial V_1 was given by Höhn (see [Höh17] and [Lam20]) using a different approach based on commuting pairs.

The classification of the strongly rational, holomorphic vertex operator algebras of central charge 24 with non-trivial V_1 described above and Theorem 6.6 yield the classification of the generalised deep holes of V_Λ with $\text{rk}((V_\Lambda^g)_1) > 0$. We observe:

Theorem (Theorem 6.10). *Under the natural projection $\text{Aut}(V_\Lambda) \rightarrow \text{O}(\Lambda)$ the 70 algebraic conjugacy classes of generalised deep holes g with $\text{rk}((V_\Lambda^g)_1) > 0$ map to the 11 algebraic conjugacy classes in $\text{O}(\Lambda)$ with cycle shapes 1^{24} , $1^8 2^8$, $1^6 3^6$, 2^{12} , $1^4 2^2 4^4$, $1^4 5^4$, $1^2 2^2 3^2 6^2$, $1^3 7^3$, $1^2 2^4 4^1 8^2$, $2^3 6^3$ and $2^2 10^2$.*

More precisely, we exactly recover the decomposition of the genus of the Moonshine module described by Höhn in [Höh17] (see Table 1).

Finally, we give a full classification of the generalised deep holes of the Leech lattice vertex operator algebra V_Λ , including those with $\text{rk}((V_\Lambda^g)_1) = 0$, based on Carnahan's constructions of the Moonshine module [Car18] (see Theorem 6.9).

Outline. In Section 2 we review finite-order automorphisms of simple Lie algebras and prove an averaged version of Kac's very strange formula.

In Section 3 we recall some results on lattice vertex operator algebras, automorphisms of the vertex operator algebra V_Λ associated with the Leech lattice and the cyclic orbifold construction.

In Section 4 we review modular forms of weight 2 for $\Gamma_0(n)$ and vector-valued modular forms for the Weil representation. Then we construct a certain Eisenstein series of weight 2 for the dual Weil representation of $\Pi_{1,1}(n)$.

In Section 5 we derive a formula for the dimension of the weight-1 space of the cyclic orbifold construction by pairing the character of V^g with the Eisenstein series of the preceding section and show that the dimension is bounded from above. We introduce the notion of extremal automorphisms, for which the upper bound is attained.

In Section 6 we define generalised deep holes, certain extremal automorphisms, and prove a bijection between the strongly rational, holomorphic vertex operator algebras of central charge 24 with non-vanishing weight-1 space and the generalised deep holes of V_Λ with non-vanishing fixed-point Lie subalgebra. Then we describe 70 generalised deep holes realising all non-zero Lie algebras on Schellekens' list.

Notation. Throughout this text we use the notation $e(x) := e^{2\pi i x}$ and $\mathbb{Z}_n := \mathbb{Z}/n\mathbb{Z}$ for $n \in \mathbb{Z}_{>0}$. The variable τ is always from the complex upper half-plane $H = \{z \in \mathbb{C} \mid \text{Im}(z) > 0\}$ and $q = e(\tau) = e^{2\pi i \tau}$. The Dedekind eta function is defined as $\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$. We set $\Gamma := \text{SL}_2(\mathbb{Z})$, and we denote by $\sigma = \sigma_1$ the usual sum-of-divisors function extended to the positive rational numbers by $\sigma(r) = 0$ if $r \notin \mathbb{Z}_{>0}$.

Acknowledgements. The authors thank Moritz Dittmann, Jethro van Ekeren, Gerald Höhn, Yi-Zhi Huang, Ching Hung Lam, Jim Lepowsky, Masahiko Miyamoto, Hiroki Shimakura, Jakob Stix and Don Zagier for helpful discussions, and the anonymous referees for their suggestions.

Sven Möller was supported by a JSPS *Postdoctoral Fellowship for Research in Japan* and by JSPS Grant-in-Aid KAKENHI 20F40018. Nils Scheithauer was supported by the DFG through the CRC *Geometry and Arithmetic of Uniformized Structures*, project number 444845124. Both authors also thank the LOEWE research unit *Uniformized Structures in Arithmetic and Geometry* for its support.

2. LIE ALGEBRAS

In this section we review the theory of finite-order automorphisms of finite-dimensional, simple Lie algebras and the very strange formula [Kac90]. Then we prove an averaged version of the very strange formula.

The Lie algebras in this paper are algebras over $\mathbb{C}$.

2.1. Finite-Order Automorphisms of Simple Lie Algebras. The finite-order automorphisms of simple Lie algebras were classified in [Kac90], Chapter 8. Since we shall only need inner automorphisms, we restrict to this case.

Let $\mathfrak{g}$ be a finite-dimensional, simple Lie algebra of type X_l . Fix a Cartan subalgebra $\mathfrak{h}$ with corresponding root system $\Phi \subseteq \mathfrak{h}^*$ and a set of simple roots $\{\alpha_1, \dots, \alpha_l\} \subseteq \Phi$. The height of a root $\alpha = \sum_{i=1}^l m_i \alpha_i \in \Phi$ is defined as $\text{ht}(\alpha) = \sum_{i=1}^l m_i$. There is a unique root $\theta \in \Phi$ of maximal height. We normalise the non-degenerate, symmetric, invariant bilinear form $(\cdot, \cdot)$ on $\mathfrak{g}$ such that $(\theta, \theta) = 2$. The Weyl vector ρ of $\mathfrak{g}$ is the half sum of the positive roots $\rho = (1/2) \sum_{\alpha \in \Phi^+} \alpha$ (or the sum of the fundamental weights). The relation $\theta = \sum_{i=1}^l a_i \alpha_i$ defines the Kac labels (or marks) $a_1, \dots, a_l$, complemented with $a_0 = 1$. The untwisted affine Dynkin diagram $X_l^{(1)}$ associated with X_l has $l + 1$ nodes labelled by the indices $0, \dots, l$ (see Table Aff 1 in [Kac90]).

By Theorem 8.6 in [Kac90] the conjugacy classes of inner automorphisms of $\mathfrak{g}$ of order $n \in \mathbb{Z}_{>0}$ are in bijection with the sequences $s = (s_0, \dots, s_l)$ of non-negative, coprime integers satisfying $\sum_{i=0}^l a_i s_i = n$, modulo automorphisms of the affine Dynkin diagram $X_l^{(1)}$. The conjugacy class associated with a sequence s is represented by the inner automorphism σ_s uniquely specified by the relations

$$\sigma_s(x_i) = e(s_i/n)x_i$$

for $i = 1, \dots, l$ where x_i spans the 1-dimensional root space $\mathfrak{g}_{\alpha_i}$ corresponding to the simple root α_i .

The automorphism σ_s is characterised by the vector $\lambda_s \in \mathfrak{h}^*$ defined by $(\alpha_i, \lambda_s) = s_i/n$ for all simple roots α_i . Identifying the Cartan subalgebra $\mathfrak{h}$ with its dual $\mathfrak{h}^*$ via $(\cdot, \cdot)$ we can write $\sigma_s = e^{2\pi i \text{ad}(\lambda_s)}$.

2.2. Very Strange Formula. The very strange formula, which generalises the strange formula of Freudenthal–de Vries, relates the dimensions of the eigenspaces of a finite-order automorphism of a simple Lie algebra to the distance of the corresponding vector to the Weyl vector. Here, we only state it for inner automorphisms.

Let $\mathfrak{g}$ be a finite-dimensional, simple Lie algebra with dual Coxeter number $h^\vee$. Given an automorphism σ of $\mathfrak{g}$ of order n , we define

$$B(\sigma) := \frac{\dim(\mathfrak{g})}{24} - \frac{1}{4n^2} \sum_{j=1}^{n-1} j(n-j) \dim(\mathfrak{g}_{(j)})$$

where $\mathfrak{g}_{(j)}$ is the eigenspace $\mathfrak{g}_{(j)} = \{x \in \mathfrak{g} \mid \sigma x = e(j/n)x\}$. We note that $B(\sigma)$ can be written as

$$B(\sigma) = \frac{1}{4} \sum_{j=0}^{n-1} B_2(j/n) \dim(\mathfrak{g}_{(j)})$$

where $B_2(x) = x^2 - x + 1/6$ is the second Bernoulli polynomial. In both formulae we may replace the (exact) order n of σ by any multiple of it.

Proposition 2.1 (Very Strange Formula, formula (12.3.6) in [Kac90]). *Let $\mathfrak{g}$ be a finite-dimensional, simple Lie algebra and σ an inner automorphism of $\mathfrak{g}$ of order n corresponding to the sequence $s = (s_0, \dots, s_l)$. Then*

$$\frac{h^\vee}{2} \left| \lambda_s - \frac{\rho}{h^\vee} \right|^2 = \frac{\dim(\mathfrak{g})}{24} - \frac{1}{4n^2} \sum_{j=1}^{n-1} j(n-j) \dim(\mathfrak{g}_{(j)}).$$

The squared norm on the left-hand side is formed with respect to the bilinear form on $\mathfrak{h}^*$ induced by $(\cdot, \cdot)$.

We define the automorphism $\sigma_{\rho/h^\vee} = e^{2\pi i \text{ad}(\rho/h^\vee)}$, whose conjugacy class is characterised by $\lambda_s = \rho/h^\vee$. The following corollary is immediate:

Corollary 2.2. *Let $\mathfrak{g}$ be a finite-dimensional, simple Lie algebra. Then $B(\sigma) \geq 0$ for all finite-order, inner automorphisms σ of $\mathfrak{g}$, and equality is attained if and only if σ is conjugate to $\sigma_{\rho/h^\vee}$.*

The automorphism $\sigma_{\rho/h^\vee}$ has order $n = lh^\vee$ where $l \in \{1, 2, 3\}$ denotes the lacing number of $\mathfrak{g}$, is regular, i.e. the fixed-point Lie subalgebra is abelian, and rational, i.e. $\sigma_{\rho/h^\vee}$ is conjugate to $\sigma_{\rho/h^\vee}^i$ for all $i \in \mathbb{Z}_n^*$. In particular, $\sigma_{\rho/h^\vee}$ is quasirational, i.e. the dimensions of the eigenspaces $\mathfrak{g}_{(j)}$ only depend on (j, n) .

If $\mathfrak{g}$ is simply-laced, then $\sigma_{\rho/h^\vee}$ is the up to conjugacy unique minimal-order regular automorphism of $\mathfrak{g}$ and $s = (1, \dots, 1)$ (see Exercise 8.11 in [Kac90]).

Finally, we note that if an automorphism σ of $\mathfrak{g}$ is conjugate to $\sigma_{\rho/h^\vee}$, then it must already be conjugate to $\sigma_{\rho/h^\vee}$ under an inner automorphism of $\mathfrak{g}$ because the sequence s characterising $\sigma_{\rho/h^\vee}$ is invariant under the diagram automorphisms of the untwisted affine Dynkin diagram of $\mathfrak{g}$.

2.3. Averaged Very Strange Formula. In the following we prove an averaged version of Kac's very strange formula. Remarkably, the coefficients of the eigenspaces that will appear in this formula are identical to those in the dimension formula in Theorem 5.3.

Given an automorphism σ of $\mathfrak{g}$ of order n (or dividing n), we define

$$A(\sigma) := \frac{1}{\phi(n)} \sum_{i \in \mathbb{Z}_n^*} B(\sigma^i).$$

Clearly, $A(\sigma) = B(\sigma)$ if σ is quasirational. This is the case that was studied in [ELMS21]. Here, we do not make this assumption.

The definition of $A(\sigma)$ and Corollary 2.2 imply:

Proposition 2.3. *Let $\mathfrak{g}$ be a finite-dimensional, simple Lie algebra. Then $A(\sigma) \geq 0$ for all finite-order, inner automorphisms σ of $\mathfrak{g}$, and equality is attained if and only if σ is conjugate to $\sigma_{\rho/h^\vee}$.*

Proof. Let σ be an inner automorphism of $\mathfrak{g}$ of order n . Then $B(\sigma^i) \geq 0$ for all $i \in \mathbb{Z}_n^*$ implies $A(\sigma) \geq 0$. If $A(\sigma) = 0$, then $B(\sigma) = 0$ so that σ is conjugate to $\sigma_{\rho/h^\vee}$. Conversely, if σ is conjugate to $\sigma_{\rho/h^\vee}$, then so is σ^i for all $i \in \mathbb{Z}_n^*$ by the rationality of $\sigma_{\rho/h^\vee}$. Hence, $B(\sigma^i) = 0$ for all $i \in \mathbb{Z}_n^*$, and therefore $A(\sigma) = 0$. $\square$

We now compute the coefficients of the eigenspace dimensions $\dim(\mathfrak{g}_{(j)})$ in $A(\sigma)$. Because of the averaging these coefficients only depend on (j, n) . Equivalently, we may rewrite $A(\sigma)$ as a linear combination of the fixed-point dimensions $\dim(\mathfrak{g}^{\sigma^m})$ for $m \mid n$.

Let $n \in \mathbb{Z}_{>0}$. We define the constants $c_n(m) \in \mathbb{Q}$ for $m \mid n$ by the linear system of equations

$$\sum_{m \mid n} c_n(m)(t, m) = n/t$$

for all $t \mid n$. An explicit formula will be given in Section 5.2.

Theorem 2.4 (Averaged Very Strange Formula). *Let $\mathfrak{g}$ be a finite-dimensional, simple Lie algebra and σ an inner automorphism of $\mathfrak{g}$ of order n . Then*

$$\frac{1}{\phi(n)} \sum_{i \in \mathbb{Z}_n^*} \frac{h^\vee}{2} \left| \lambda_{s^{[i]}} - \frac{\rho}{h^\vee} \right|^2 = \frac{1}{24n} \sum_{m|n} c_n(m) \dim(\mathfrak{g}^{\sigma^m})$$

where $s^{[i]}$ is the sequence characterising σ^i .

Again, we may replace the order n in the above formula by any multiple of it.

Proof. We assume that $n > 1$. By the very strange formula (see Proposition 2.1) and the definition of $A(\sigma)$, the left-hand side equals

$$\begin{aligned} A(\sigma) &= \frac{1}{\phi(n)} \sum_{i \in \mathbb{Z}_n^*} \frac{1}{4} \sum_{j=0}^{n-1} B_2(j/n) \dim(\mathfrak{g}_{(ij)}) \\ &= \sum_{j=0}^{n-1} \left(\frac{1}{\phi(n/(j,n))} \sum_{\substack{i=0 \\ (i,n)=(j,n)}}^{n-1} \frac{1}{4} B_2(i/n) \right) \dim(\mathfrak{g}_{(j)}). \end{aligned}$$

Let $k > 1$. Then

$$\sum_{\substack{i=1 \\ (i,k)=1}}^k 1 = \phi(k), \quad \sum_{\substack{i=1 \\ (i,k)=1}}^k i = \frac{1}{2} k \phi(k), \quad \sum_{\substack{i=1 \\ (i,k)=1}}^k i^2 = \frac{1}{3} \phi(k) (k^2 + \frac{1}{2} \lambda(k))$$

with

$$\lambda(k) := \prod_{\substack{p|k \\ p \text{ prime}}} (-p).$$

(For the third sum see, e.g., [PS71], Abschnitt VIII, Aufgabe 27.) Hence, summing $B_2(i/n) = (i/n)^2 - i/n + 1/6$ over i yields

$$\frac{1}{\phi(n/(j,n))} \sum_{\substack{i=0 \\ (i,n)=(j,n)}}^{n-1} \frac{1}{4} B_2(i/n) = \frac{1}{24} \frac{\lambda(n/(j,n))}{(n/(j,n))^2}.$$

Noting that

$$\sum_{\substack{j=0 \\ (j,n)=d}}^{n-1} \dim(\mathfrak{g}_{(j)}) = \sum_{m|(n/d)} \mu(n/dm) \dim(\mathfrak{g}^{\sigma^m})$$

for all $d | n$ with the Möbius function μ we obtain

$$A(\sigma) = \sum_{m|n} \left(\sum_{d|(n/m)} \frac{\lambda(n/d) \mu(n/dm)}{24(n/d)^2} \right) \dim(\mathfrak{g}^{\sigma^m}).$$

Finally we show that

$$\sum_{m|n} \left(\sum_{d|(n/m)} \frac{\lambda(n/d) \mu(n/dm)}{(n/d)^2} \right) (t, m) = \frac{1}{t}$$

for all $t | n$ using the multiplicativity of the left-hand side in n , which implies that

$$\sum_{d|(n/m)} \frac{\lambda(n/d) \mu(n/dm)}{(n/d)^2} = \frac{1}{n} c_n(m).$$

Now the assertion follows. $\square$

3. VERTEX OPERATOR ALGEBRAS

In this section we recall some results about lattice vertex operator algebras and their automorphism groups and describe the cyclic orbifold construction developed in [EMS20a, Möl16].

The vertex operator algebras in this paper are defined over $\mathbb{C}$.

3.1. Lattice Vertex Operator Algebras. To keep the exposition self-contained we recall a few well-known facts about lattice vertex operator algebras [Bor86, FLM88].

For a positive-definite, even lattice L with bilinear form $(\cdot, \cdot): L \times L \rightarrow \mathbb{Z}$ the associated vertex operator algebra of central charge $c = \text{rk}(L)$ is given by

$$V_L = M(1) \otimes \mathbb{C}[L]_\varepsilon$$

where $M(1)$ is the Heisenberg vertex operator algebra associated with $L_{\mathbb{C}} = L \otimes_{\mathbb{Z}} \mathbb{C}$ and $\mathbb{C}[L]_\varepsilon$ is the twisted group algebra, i.e. the algebra with basis $\{e^\alpha \mid \alpha \in L\}$ and products $e^\alpha e^\beta = \varepsilon(\alpha, \beta) e^{\alpha+\beta}$ for all $\alpha, \beta \in L$ where $\varepsilon: L \times L \rightarrow \{\pm 1\}$ is a 2-cocycle satisfying $\varepsilon(\alpha, \beta)/\varepsilon(\beta, \alpha) = (-1)^{(\alpha, \beta)}$. Note that ε defines a central extension $\hat{L}$ of L by $1 \rightarrow \{\pm 1\} \rightarrow \{\pm 1\} \times_\varepsilon L \rightarrow L \rightarrow 0$ and $\mathbb{C}[L]_\varepsilon = \mathbb{C}[\hat{L}]/J$ where J is the kernel of the linear homomorphism $\mathbb{C}[\hat{L}] \rightarrow \mathbb{C}[L]$ defined by $e^{(\pm 1, \alpha)} \mapsto \pm e^\alpha$.

Let $\nu \in O(L)$, the orthogonal group of the lattice L , and let $\eta: L \rightarrow \{\pm 1\}$ be a function. Then the map $\phi_\eta(\nu)$ acting on $\mathbb{C}[L]_\varepsilon$ as $\phi_\eta(\nu)(e^\alpha) = \eta(\alpha) e^{\nu\alpha}$ for $\alpha \in L$ defines an automorphism of $\mathbb{C}[L]_\varepsilon$ if and only if

$$\frac{\eta(\alpha)\eta(\beta)}{\eta(\alpha+\beta)} = \frac{\varepsilon(\alpha, \beta)}{\varepsilon(\nu\alpha, \nu\beta)}$$

for all $\alpha, \beta \in L$. In this case $\phi_\eta(\nu)$ is called a *lift* of ν . Since the lifts can be identified with the automorphisms of $\hat{L}$ which preserve the norm in L , we denote the group of lifts by $O(\hat{L})$. In contrast to $O(L)$ the group $O(\hat{L})$ acts naturally on the vertex operator algebra $V_L = M(1) \otimes \mathbb{C}[L]_\varepsilon$.

There is a short exact sequence

$$1 \rightarrow \text{Hom}(L, \{\pm 1\}) \rightarrow O(\hat{L}) \rightarrow O(L) \rightarrow 1$$

with the surjection $O(\hat{L}) \rightarrow O(L)$ given by $\phi_\eta(\nu) \mapsto \nu$. The image of $\text{Hom}(L, \{\pm 1\})$ in $O(\hat{L})$ are exactly the lifts of the identity in $O(L)$.

If the restriction of η to the fixed-point sublattice L^ν is trivial, we call $\phi_\eta(\nu)$ a *standard lift* of ν . It is always possible to choose η in this way (see [Lep85], Section 5).

Let $\phi_\eta(\nu)$ be a standard lift of ν , and suppose that ν has order m . If m is odd or if m is even and $(\alpha, \nu^{m/2}\alpha)$ is even for all $\alpha \in L$, then the order of $\phi_\eta(\nu)$ is also m . Otherwise the order of $\phi_\eta(\nu)$ is $2m$, in which case we say ν exhibits *order doubling*.

For any vertex operator algebra V of CFT-type $K := \langle e^{v_0} \mid v \in V_1 \rangle$ is a subgroup of $\text{Aut}(V)$, called the *inner automorphism group* of V . By [DN99], Theorem 2.1, the automorphism group of V_L can be written as

$$\text{Aut}(V_L) = O(\hat{L}) \cdot K.$$

Here, K is normal in $\text{Aut}(V_L)$, $\text{Hom}(L, \{\pm 1\})$ is a subgroup of $K \cap O(\hat{L})$ and $\text{Aut}(V_L)/K$ is isomorphic to a quotient of $O(L)$.

All standard lifts of a given lattice automorphism $\nu \in O(L)$ are conjugate in $\text{Aut}(V_L)$, in fact under elements of K (see [EMS20a], Proposition 7.1).

Choosing a section $O(L) \rightarrow O(\hat{L})$, $\nu \mapsto \phi_\eta(\nu)$, we can write every automorphism of V_L as $\phi_\eta(\nu)\sigma$ for some $\nu \in O(L)$, $\sigma \in K$ because any two lifts of ν only differ by a homomorphism $L \rightarrow \{\pm 1\}$, which can be absorbed into σ .

We also recall some results about the twisted modules of lattice vertex operator algebras. The irreducible $\phi_\eta(\nu)$ -twisted modules of a lattice vertex operator algebra V_L for standard lifts $\phi_\eta(\nu)$ are described in [DL96, BK04]. Together with the results in Section 5 of [Li96] this allows us to describe the irreducible g -twisted V_L -modules for all finite-order automorphisms $g \in \text{Aut}(V_L)$ (cf. [HM20]).

By construction, $(V_L)_1$ is a reductive Lie algebra, and the subspace $\mathfrak{h} := \{h(-1) \otimes e^0 \mid h \in L_{\mathbb{C}}\} \cong L_{\mathbb{C}}$ is a Cartan subalgebra of $(V_L)_1$. Now, let $g = \phi_\eta(\nu)\sigma_h$ where $\phi_\eta(\nu)$ is some lift of $\nu \in \text{O}(L)$ and $\sigma_h = e^{2\pi i h_0}$ for some $h \in L_{\mathbb{C}}$. Note that g has finite order if and only if $h \in L \otimes_{\mathbb{Z}} \mathbb{Q}$. Suppose ν has order m . In Lemma 8.3 of [EMS20b] (see also Lemma 3.4 in [LS19]) it is shown that $\phi_\eta(\nu)\sigma_h$ is conjugate to $\phi_\eta(\nu)\sigma_{\pi_\nu(h)}$ where $\pi_\nu = \frac{1}{m} \sum_{i=0}^{m-1} \nu^i$ is the projection of $L_{\mathbb{C}}$ onto the elements of $L_{\mathbb{C}}$ fixed by ν . Hence we may assume that $h \in \pi_\nu(L_{\mathbb{C}})$. Then σ_h and $\phi_\eta(\nu)$ commute, and we can apply the results in [Li96] to $g = \phi_\eta(\nu)\sigma_h$.

We shall be interested in the case that L is unimodular. Then V_L is holomorphic and there is a unique g -twisted V_L -module $V_L(g)$ for each $g \in \text{Aut}(V_L)$ of finite order. Let $g = \phi_\eta(\nu)\sigma_h$ where $\nu \in \text{O}(L)$ has order m , $\phi_\eta(\nu)$ is a standard lift of ν , $\sigma_h = e^{2\pi i h_0}$ and $h \in \pi_\nu(L \otimes_{\mathbb{Z}} \mathbb{Q})$. Suppose that ν has cycle shape $\prod_{t|m} t^{b_t}$, i.e. ν has characteristic polynomial $\prod_{t|m} (x^t - 1)^{b_t}$ as automorphism of $L_{\mathbb{C}}$. Note that $\sum_{t|m} t b_t = \text{rk}(L) = c$. Then the conformal weight of $V_L(g)$ is given by

$$\rho(V_L(g)) = \frac{1}{24} \sum_{t|m} b_t \left(t - \frac{1}{t} \right) + \min_{\alpha \in -h + \pi_\nu(L)} \frac{(\alpha, \alpha)}{2} \geq 0.$$

The number

$$\rho_\nu := \frac{1}{24} \sum_{t|m} b_t \left(t - \frac{1}{t} \right)$$

is called the *vacuum anomaly* of $V_L(g)$. It is positive for $\nu \neq \text{id}$.

3.2. Automorphisms of the Leech Lattice Vertex Operator Algebra. We now specialise the exposition to the Leech lattice.

The Leech lattice Λ is the up to isomorphism unique positive-definite, even, unimodular lattice of rank 24 without roots, i.e. vectors of squared norm 2. The automorphism group $\text{O}(\Lambda)$ of the Leech lattice is Conway's group Co_0 . Because $(V_\Lambda)_1 = \mathfrak{h} = \{h(-1) \otimes e^0 \mid h \in \Lambda_{\mathbb{C}}\}$, the inner automorphism group is given by

$$K = \{\sigma_h \mid h \in \Lambda_{\mathbb{C}}\}$$

with $\sigma_h = e^{2\pi i h_0}$ and is abelian. Moreover, $\text{Aut}(V_\Lambda)/K \cong \text{O}(\Lambda)$, i.e. there is a short exact sequence

$$1 \rightarrow K \rightarrow \text{Aut}(V_\Lambda) \rightarrow \text{O}(\Lambda) \rightarrow 1,$$

because $K \cap \text{O}(\hat{\Lambda}) = \text{Hom}(\Lambda, \{\pm 1\})$ in the special case of the Leech lattice. Hence, every automorphism of V_Λ is of the form

$$\phi_\eta(\nu)\sigma_h$$

for a lift $\phi_\eta(\nu)$ of some $\nu \in \text{O}(\Lambda)$ and some $h \in \Lambda_{\mathbb{C}}$. The surjective homomorphism $\text{Aut}(V_\Lambda) \rightarrow \text{O}(\Lambda)$ in the short exact sequence is given by $\phi_\eta(\nu)\sigma_h \mapsto \nu$.

Recall that it suffices to take the lift $\phi_\eta(\nu)$ from a fixed section. Moreover, since $\sigma_h = \text{id}$ if and only if $h \in \Lambda' = \Lambda$, we can take $h \in \Lambda_{\mathbb{C}}/\Lambda$.

In the following we describe the conjugacy classes of $\text{Aut}(V_\Lambda)$. As mentioned above, the automorphism $\phi_\eta(\nu)\sigma_h$ is conjugate to $\phi_\eta(\nu)\sigma_{\pi_\nu(h)}$ for any $h \in \Lambda_{\mathbb{C}}$, and $\phi_\eta(\nu)$ and $\sigma_{\pi_\nu(h)}$ commute.

Fix a section $\varphi: \text{O}(L) \rightarrow \text{O}(\hat{L})$, and for $\nu \in \text{O}(\Lambda)$ denote by $\eta_\nu: \Lambda \rightarrow \{\pm 1\}$ the corresponding function characterising the action of $\varphi(\nu)$ on $\mathbb{C}[\Lambda]_\varepsilon$.

Lemma 3.1. *Any automorphism in $\text{Aut}(V_\Lambda)$ is conjugate to an automorphism of the form $\varphi(\nu)\sigma_h$ where*

- (1) ν is in a fixed set N of representatives for the conjugacy classes in $O(\Lambda)$,
- (2) h is in a fixed set H_ν of representatives for the orbits of the action of $C_{O(\Lambda)}(\nu)$ on $\pi_\nu(\Lambda_\mathbb{C})/\pi_\nu(\Lambda)$.

Proof. Let $g \in \text{Aut}(V_\Lambda)$. We have already argued that g is conjugate to $\varphi(\nu)\sigma_h$ for some $\nu \in O(\Lambda)$ and $h \in \pi_\nu(\Lambda_\mathbb{C})/\pi_\nu(\Lambda)$. Now suppose that $\mu = \tau^{-1}\nu\tau$ for some $\tau, \mu \in O(\Lambda)$. Let $\phi_\eta(\tau)$ be any lift of τ . Then $\varphi(\mu)^{-1}\phi_\eta(\tau)^{-1}\varphi(\nu)\sigma_h\phi_\eta(\tau)$ acts as $\mu^{-1}\tau^{-1}\nu\tau = \text{id}$ on $(V_\Lambda)_1 \cong \Lambda_\mathbb{C}$. By Lemma 2.5 in [DN99] this implies that $\varphi(\mu)^{-1}\phi_\eta(\tau)^{-1}\varphi(\nu)\sigma_h\phi_\eta(\tau) = \sigma_{h'}$ for some $h' \in \Lambda_\mathbb{C}$, i.e. $\varphi(\nu)\sigma_h$ is conjugate to $\varphi(\mu)\sigma_{h'}$. As explained above we may even assume that $h' \in \pi_\mu(\Lambda_\mathbb{C})/\pi_\mu(\Lambda)$. This proves item (1).

To see item (2) we consider $\varphi(\nu)\sigma_h$ with $h \in \pi_\nu(\Lambda_\mathbb{C})$ and let $h' \in \pi_\nu(\Lambda_\mathbb{C})$ with $h' + \pi_\nu(\Lambda) = \tau h + \pi_\nu(\Lambda)$ for some $\tau \in C_{O(\Lambda)}(\nu)$ and prove that $\varphi(\nu)\sigma_h$ and $\varphi(\nu)\sigma_{h'}$ are conjugate. Indeed, $(\varphi(\nu)\sigma_{h'})^{-1}\varphi(\tau)\varphi(\nu)\sigma_h\varphi(\tau)^{-1}$ acts as identity on $(V_\Lambda)_1 \cong \Lambda_\mathbb{C}$ and on $e^\alpha \in \mathbb{C}[\Lambda]_\varepsilon$ as

$$\begin{aligned} & \sigma_{-h'}\varphi(\nu)^{-1}\varphi(\tau)\varphi(\nu)\sigma_h\varphi(\tau)^{-1}e^\alpha \\ &= \frac{e^{-2\pi i(h', \nu^{-1}\tau\nu\tau^{-1}\alpha)}\eta_\tau(\nu\tau^{-1}\alpha)\eta_\nu(\tau^{-1}\alpha)e^{2\pi i(h, \tau^{-1}\alpha)}}{\eta_\nu(\nu^{-1}\tau\nu\tau^{-1}\alpha)\eta_\tau(\tau^{-1}\alpha)}e^{\nu^{-1}\tau\nu\tau^{-1}\alpha} \\ &= e^{2\pi i(\tau h - h', \alpha)}\frac{\eta_\tau(\nu\tau^{-1}\alpha)\eta_\nu(\tau^{-1}\alpha)}{\eta_\nu(\alpha)\eta_\tau(\tau^{-1}\alpha)}e^\alpha \\ &=: \lambda(\alpha)e^\alpha. \end{aligned}$$

The map $\lambda: \Lambda \rightarrow \mathbb{C}^*$ is a homomorphism that is trivial on Λ^ν . This means that $(\varphi(\nu)\sigma_{h'})^{-1}\varphi(\tau)\varphi(\nu)\sigma_h\varphi(\tau)^{-1} = \sigma_f$ for some $f \in \Lambda_\mathbb{C}$ with $(f, \alpha) \in \mathbb{Z}$ for all $\alpha \in \Lambda^\nu$. We just showed that $\varphi(\nu)\sigma_h$ and $\varphi(\nu)\sigma_{h'+f}$ are conjugate. The latter is conjugate to $\varphi(\nu)\sigma_{h'+\pi_\nu(f)}$, but $\pi_\nu(f) \in (\Lambda^\nu)'$ and $(\Lambda^\nu)' = \pi_\nu(\Lambda)$ because Λ is unimodular. This proves the claim. $\square$

The above lemma provides the surjective map $(\nu, h) \mapsto \varphi(\nu)\sigma_h$ from the set $Q := \{(\nu, h) \mid \nu \in N, h \in H_\nu\}$ onto the conjugacy classes of $\text{Aut}(V_\Lambda)$. We shall now prove that this map is in fact a bijection.

Proposition 3.2. *The map $(\nu, h) \mapsto \varphi(\nu)\sigma_h$ is a bijection from the set Q to the conjugacy classes of $\text{Aut}(V_\Lambda)$.*

Proof. We saw in Lemma 3.1 that this map is surjective. To see the injectivity let (ν, h) and (μ, f) be in the set Q and assume that the corresponding automorphisms $\varphi(\nu)\sigma_h$ and $\varphi(\mu)\sigma_f$ are conjugate in $\text{Aut}(V_\Lambda)$. We need to show that $(\nu, h) = (\mu, f)$.

Applying the surjection $\text{Aut}(V_\Lambda) \rightarrow O(\Lambda)$ immediately yields that ν and μ are conjugate and hence $\nu = \mu$.

Now assume $\varphi(\nu)\sigma_h$ and $\varphi(\nu)\sigma_f$ are conjugate under some automorphism $\varphi(\tau)\sigma_s$. Then $\tau \in C_{O(\Lambda)}(\nu)$ and $(\varphi(\nu)\sigma_f)^{-1}(\varphi(\tau)\sigma_s)^{-1}\varphi(\nu)\sigma_h\varphi(\tau)\sigma_s = \text{id}$. The latter acts on $e^\alpha \in \mathbb{C}[\Lambda]_\varepsilon$ by multiplication with

$$\frac{\eta_\tau(\alpha)\eta_\nu(\tau\alpha)}{\eta_\tau(\nu\alpha)\eta_\nu(\alpha)}e^{2\pi i(s - \nu^{-1}s + \tau^{-1}h - f, \alpha)} = 1$$

for all $\alpha \in \Lambda$. Since ν and τ commute and η_ν is trivial on Λ^ν , this implies that $e^{2\pi i(\tau^{-1}h - f, \alpha)} = 1$ for all $\alpha \in \Lambda^\nu$. Hence $\tau^{-1}h - f \in (\Lambda^\nu)' = \pi_\nu(\Lambda)$. This means that $h + \pi_\nu(\Lambda)$ and $f + \pi_\nu(\Lambda)$ are in the same orbit under the action of $C_{O(\Lambda)}(\nu)$ and hence identical. $\square$

Recall that the automorphism $\varphi(\nu)\sigma_h$ in $\text{Aut}(V_\Lambda)$ has finite order if and only if h is in $\pi_\nu(\Lambda \otimes_\mathbb{Z} \mathbb{Q})$.

3.3. Orbifold Construction. In [EMS20a] (see also [Möl16]) we develop a cyclic orbifold theory for holomorphic vertex operator algebras. Below we give a short summary.

A vertex operator algebra V is called *strongly rational* if it is rational (as defined, e.g., in [DLM97]), C_2 -cofinite (or lisse), self-contragredient (or self-dual) and of CFT-type. Then V is also simple. Moreover, a rational vertex operator algebra V is said to be *holomorphic* if V itself is the only irreducible V -module. Note that the central charge c of a strongly rational, holomorphic vertex operator algebra V is necessarily a non-negative multiple of 8, a simple consequence of Zhu's modular invariance result [Zhu96]. Examples of strongly rational vertex operator algebras are the lattice vertex operator algebras discussed above.

Let V be a strongly rational, holomorphic vertex operator algebra and $G = \langle g \rangle$ a finite, cyclic group of automorphisms of V of order n .

By [DLM00], for each $i \in \mathbb{Z}_n$, there is an irreducible g^i -twisted V -module $V(g^i)$, which is unique up to isomorphism. The uniqueness of $V(g^i)$ implies that there is a representation $\phi_i: G \rightarrow \text{Aut}_{\mathbb{C}}(V(g^i))$ of G on the vector space $V(g^i)$ such that

$$\phi_i(g)Y_{V(g^i)}(v, x)\phi_i(g)^{-1} = Y_{V(g^i)}(gv, x)$$

for all $v \in V$. This representation is unique up to an n -th root of unity. Let $W^{(i,j)}$ denote the eigenspace of $\phi_i(g)$ in $V(g^i)$ corresponding to the eigenvalue $e(j/n)$. On $V(g^0) = V$ we choose $\phi_0(g) = g$.

The fixed-point vertex operator subalgebra $V^g = W^{(0,0)}$ is strongly rational by [DM97, Miy15, CM16, McR21] and has exactly n^2 irreducible modules, namely the $W^{(i,j)}$, $i, j \in \mathbb{Z}_n$ [MT04, DRX17]. We further show that the conformal weight $\rho(V(g))$ of $V(g)$ is in $(1/n^2)\mathbb{Z}$, and we define the type $t \in \mathbb{Z}_n$ of g by $t = n^2\rho(V(g)) \bmod n$.

Assume for simplicity that g has type $t = 0$, i.e. that $\rho(V(g)) \in (1/n)\mathbb{Z}$. We may choose the representations ϕ_i such that the conformal weights satisfy

$$\rho(W^{(i,j)}) = \frac{ij}{n} \bmod 1$$

and V^g has fusion rules

$$W^{(i,j)} \boxtimes W^{(l,k)} \cong W^{(i+l, j+k)}$$

for all $i, j, k, l \in \mathbb{Z}_n$ [EMS20a], i.e. the fusion algebra of V^g is the group ring $\mathbb{C}[D]$ where D is the discriminant form with group structure $\mathbb{Z}_n \times \mathbb{Z}_n$ and quadratic form defined by the conformal weights modulo 1. This is the discriminant form of the hyperbolic lattice $II_{1,1}(n)$, i.e. the lattice with Gram matrix $\begin{pmatrix} 0 & n \\ n & 0 \end{pmatrix}$. In particular, all V^g -modules are simple currents.

In essence, the results in [EMS20a] show that for a strongly rational, holomorphic vertex operator algebra V and cyclic $G \cong \mathbb{Z}_n$ the module category of V^G is the twisted group double $\mathcal{D}_\omega(G)$ where the 3-cocycle $[\omega] \in H^3(G, \mathbb{C}^*) \cong \mathbb{Z}_n$ is determined by the type $t \in \mathbb{Z}_n$. This proves a conjecture by Dijkgraaf, Vafa, Verlinde and Verlinde [DVVV89] who stated it for arbitrary finite G .

If $24 \mid c$, then the characters of the irreducible V^g -modules

$$\text{ch}_{W^{(i,j)}}(\tau) = \text{tr}|_{W^{(i,j)}} q^{L_0 - c/24},$$

$i, j \in \mathbb{Z}_n$, form a vector-valued modular form of weight 0 for the Weil representation ρ_D of $\Gamma = \text{SL}_2(\mathbb{Z})$ on $\mathbb{C}[D]$. Since D has level n , the individual characters $\text{ch}_{W^{(i,j)}}(\tau)$ are modular forms for the congruence subgroup $\Gamma(n)$. Moreover, $\text{ch}_{W^{(0,0)}}(\tau) = \text{ch}_{V^g}(\tau)$ is modular for $\Gamma_0(n)$. Similar results hold if $24 \nmid c$ or if $t \neq 0$. We shall explain these notions in more detail in Section 4.

In general, we say that a simple vertex operator algebra V satisfies the *positivity condition* if $\rho(W) > 0$ for any irreducible V -module $W \not\cong V$ and $\rho(V) = 0$.

Now, if V^g satisfies the positivity condition, then the direct sum of V^g -modules

$$V^{\text{orb}(g)} := \bigoplus_{i \in \mathbb{Z}_n} W^{(i,0)}$$

carries the structure of a strongly rational, holomorphic vertex operator algebra of the same central charge as V and is called the *orbifold construction* of V associated with g [EMS20a]. The vertex operator algebra structure of $V^{\text{orb}(g)}$ as extension of V^g is unique up to isomorphism. It depends only up to algebraic conjugacy on g , i.e. if $\langle g \rangle$ and $\langle h \rangle$ for some $h \in \text{Aut}(V)$ are conjugate in $\text{Aut}(V)$, then $V^{\text{orb}(g)} \cong V^{\text{orb}(h)}$. Note that the sum $\bigoplus_{j \in \mathbb{Z}_n} W^{(0,j)}$ is just the old vertex operator algebra V .

There is also an *inverse* (or *reverse*) orbifold construction. Given a vertex operator algebra $V^{\text{orb}(g)}$ obtained by the orbifold construction, we define an automorphism ζ of $V^{\text{orb}(g)}$ of order n via $\zeta v := e(i/n)v$ for $v \in W^{(i,0)}$, and the unique irreducible ζ^j -twisted $V^{\text{orb}(g)}$ -module is given by $V^{\text{orb}(g)}(\zeta^j) = \bigoplus_{i \in \mathbb{Z}_n} W^{(i,j)}$, $j \in \mathbb{Z}_n$ (see [Möl16], Theorem 4.9.6). Then

$$(V^{\text{orb}(g)})^{\text{orb}(\zeta)} = \bigoplus_{j \in \mathbb{Z}_n} W^{(0,j)} = V,$$

i.e. orbifolding with ζ is inverse to orbifolding with g .

If two elements $g, h \in \text{Aut}(V)$ of type 0 and satisfying the positivity condition are (algebraically) conjugate, the inverse-orbifold automorphisms corresponding to g and h are (algebraically) conjugate under some isomorphism $\varphi: V^{\text{orb}(g)} \rightarrow V^{\text{orb}(h)}$.

3.4. Orbifold Characters. We recall a formula for the character of the orbifold construction from [GK19], which we shall use in the proof of Theorem 6.7.

We continue in the setting of the previous section, i.e. we let V be a strongly rational, holomorphic vertex operator algebra and g an automorphism of V of order n such that V^g satisfies the positivity condition. Suppose that g has type 0 and $24 \mid c$. Then the character

$$\text{ch}_{V^{\text{orb}(g)}}(\tau) = \text{tr} |_{V^{\text{orb}(g)}} q^{L_0 - c/24}$$

is a modular form for $\Gamma = \text{SL}_2(\mathbb{Z})$ of weight 0, holomorphic on H with a possible pole at the cusp ∞ . For $k \mid n$ define

$$D_k(\tau) = \sum_{\substack{j \in \mathbb{Z}_n \\ (j,n)=k}} T(0,j)(\tau)$$

with $T(0,j)(\tau) = \text{tr} |_V g^j q^{L_0 - c/24}$. Then D_k is a modular form for $\Gamma_0(n/k)$ and

$$\text{ch}_{V^{\text{orb}(g)}}(\tau) = \frac{1}{n} \sum_{k \mid n} \sum_{M \in \Gamma_0(n/k) \backslash \Gamma} D_k|_M(\tau)$$

where $D_k|_M(\tau) = D_k(M\tau)$ (this notation is explained in Section 4).

Now, we specialise to the case where $V = V_L$ is the lattice vertex operator algebra associated with a positive-definite, even, unimodular lattice L such that $24 \mid \text{rk}(L)$. Let $g = \phi_\eta(\nu)\sigma_h$ where $\phi_\eta(\nu)$ is a standard lift of $\nu \in \text{O}(L)$ and $h \in L_\mathbb{C}$. The eta product associated with ν is defined as

$$\eta_\nu(\tau) = \prod_{t \mid m} \eta(t\tau)^{b_t}$$

if ν has order m and cycle shape $\prod_{t|m} t^{b_t}$. Define

$$w_k: L^{\nu^k} \rightarrow \mathbb{C}^*, \quad \alpha \mapsto e^{2\pi i k(\pi_\nu(h), \pi_\nu(\alpha))} \begin{cases} (-1)^{\langle \alpha, \nu^{k/2} \alpha \rangle} & \text{if } m, k \in 2\mathbb{Z}, \\ 1 & \text{else.} \end{cases}$$

Then w_k is a homomorphism and the kernel $M_{k,d}$ of w_k^d is a finite-index sublattice of L^{ν^k} . A slight generalisation of the argument given in [GK19] shows that

$$D_k(\tau) = \frac{1}{\eta_{\nu^k}(\tau)} \sum_{d|n/k} \frac{n}{kd} \mu(d) \theta_{M_{k,d}}(\tau)$$

where $\theta_{M_{k,d}}$ is the theta series of $M_{k,d}$. (Note that there is a typo in Theorem 3.9 of [GK19] and its proof.)

4. MODULAR FORMS

In this section we describe some results on modular forms of weight 2. In particular, we construct an Eisenstein series of weight 2 for the dual Weil representation of $\Pi_{1,1}(n)$.

4.1. Modular Forms of Weight 2 for $\Gamma_0(n)$. We construct a special Eisenstein series of weight 2 for $\Gamma_0(n)$ and recall the Deligne bound. For background on modular forms we refer to [DS05, Miy06].

The group $\Gamma = \mathrm{SL}_2(\mathbb{Z})$ acts on the upper half-plane H by Möbius transformations. Given a positive integer n the subgroup $\Gamma_0(n) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma \mid c \equiv 0 \pmod{n} \right\}$ has index $\psi(n)$ in Γ where $\psi(n) = n \prod_{p|n} (1 + 1/p)$ denotes the Dedekind ψ -function.

The quotient $X_0(n) = \Gamma_0(n) \backslash H^*$ where $H^* = H \cup P$ is the upper half-plane complemented by the cusps $P = \mathbb{Q} \cup \{\infty\}$ is a compact Riemann surface. The class of $a/c \in \mathbb{Q}$ with $(a, c) = 1$ in $\Gamma_0(n) \backslash P$ is characterised by the invariants (c, n) , a divisor of n , and $ac/(c, n) \in \mathbb{Z}_{(c, n/(c, n))}^*$.

It follows that $\Gamma_0(n)$ has $\varepsilon = \sum_{c|n} \phi((c, n/c))$ classes of cusps. The cusp $a/c \in \mathbb{Q}$ with $(a, c) = 1$ has width $t_{a/c} := n/(c^2, n)$. Note that $\sum_{s \in \Gamma_0(n) \backslash P} t_s = \psi(n)$. A complete system of representatives for the cusps $\Gamma_0(n) \backslash P$ is given by the rational numbers a/c where c ranges over the positive divisors of n and a over a complete set of representatives of $\mathbb{Z}_{(c, n/c)}^*$ such that $(a, c) = 1$.

We denote by $\mathcal{M}_k(\Gamma_0(n))$ the space of holomorphic modular forms for $\Gamma_0(n)$ of weight $k \in \mathbb{Z}_{>0}$ and trivial character. These are holomorphic functions on H which are invariant under modular transformations, i.e. $f|_M = f$ for all $M \in \Gamma_0(n)$ where $f|_M(\tau) := (c\tau + d)^{-k} f(M\tau)$ for $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{R})$, and which are holomorphic at the cusps.

Let $f \in \mathcal{M}_k(\Gamma_0(n))$ and $s \in \Gamma_0(n) \backslash P$ a cusp represented by a/c with $(a, c) = 1$. Choose $M_s = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$. Then $M_s \infty = s$. The Fourier expansion of $f|_{M_s}$ is called an expansion of f at s . It is an expansion in integral powers of $q^{1/t_s} = e(\tau/t_s)$. In general, it depends on the choice of M_s . However, its coefficients at integral powers of q do not.

We now specialise to weight 2. The Eisenstein series

$$E_2(\tau) = 1 - 24 \sum_{m=1}^{\infty} \sigma(m) q^m$$

transforms under $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ as

$$E_2|_M(\tau) = E_2(\tau) + \frac{12}{2\pi i} \frac{c}{c\tau + d}.$$

For $k \in \mathbb{Z}_{>0}$ we define $F_k := \frac{1}{\sqrt{k}} \begin{pmatrix} k & 0 \\ 0 & 1 \end{pmatrix}$ and

$$E_2^{(k)}(\tau) := E_2(k\tau) = \frac{1}{k} E_2|_{F_k}(\tau).$$

We describe the transformation behaviour of $E_2^{(k)}$ under Γ :

Proposition 4.1. *Let $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ and $k \in \mathbb{Z}_{>0}$. Define*

$$u = (k, c) > 0, \quad w = \frac{k}{(k, c)} > 0$$

and choose $v \in \mathbb{Z}$ such that

$$v = \left(\frac{c}{(k, c)} \right)^{-1} d \pmod{\frac{k}{(k, c)}}$$

where $(c/(k, c))^{-1}$ denotes the inverse of $c/(k, c)$ modulo $k/(k, c)$. Then

$$E_2^{(k)}|_M(\tau) = \frac{1}{w^2} E_2 \left(\frac{u\tau + v}{w} \right) + \frac{12}{2\pi i} \frac{c}{k(c\tau + d)}.$$

Proof. Let $N := \frac{1}{\sqrt{uw}} \begin{pmatrix} u & v \\ 0 & w \end{pmatrix}$. Then

$$F_k M N^{-1} = \begin{pmatrix} aw & bu - av \\ c/u & (du - cv)/k \end{pmatrix} \in \Gamma$$

and

$$\begin{aligned} E_2^{(k)}|_M(\tau) &= \frac{1}{k} E_2|_{F_k M}(\tau) = \frac{1}{k} E_2|_{F_k M N^{-1} N}(\tau) \\ &= \frac{1}{k} \frac{u}{w} \left(E_2(N\tau) + \frac{12}{2\pi i} \frac{c}{u(c\tau + d)/w} \right) \\ &= \frac{1}{w^2} E_2(N\tau) + \frac{12}{2\pi i} \frac{c}{k(c\tau + d)}. \end{aligned}$$

□

The Fourier expansion of the function $E_2((u\tau + v)/w)$ in the proposition is

$$E_2 \left(\frac{u\tau + v}{w} \right) = 1 - 24 \sum_{m=1}^{\infty} \sigma(m) e(mv/w) q^{mu/w},$$

which is an expansion in powers of $q^{k/(c^2, k)}$ because $u/w = \frac{u/(u, w)}{w/(u, w)} = \frac{u/(u, w)}{k/(c^2, k)}$.

The space of modular forms of weight 2 for $\Gamma_0(n)$ decomposes as

$$\mathcal{M}_2(\Gamma_0(n)) = \mathcal{S}_2(\Gamma_0(n)) \oplus \mathcal{E}_2(\Gamma_0(n))$$

where $\mathcal{S}_2(\Gamma_0(n))$ denotes the space of cusp forms and $\mathcal{E}_2(\Gamma_0(n))$ the Eisenstein space, the orthogonal complement of $\mathcal{S}_2(\Gamma_0(n))$ with respect to the Petersson inner product. The dimensions of the spaces are given by

$$\dim(\mathcal{S}_2(\Gamma_0(n))) = g(X_0(n)),$$

$$\dim(\mathcal{E}_2(\Gamma_0(n))) = \varepsilon - 1$$

where $g(X_0(n))$ is the genus of the modular curve $X_0(n) = \Gamma_0(n) \backslash H^*$ and ε the number of cusps of $\Gamma_0(n)$. An explicit basis of $\mathcal{E}_2(\Gamma_0(n))$ is given, for example, in [DS05], Theorem 4.6.2.

Let $n > 1$ and $k \mid n$, $k > 1$. Then

$$f_k(\tau) := E_2(\tau) - k E_2^{(k)}(\tau) = E_2(\tau) - k E_2(k\tau)$$

is in $\mathcal{E}_2(\Gamma_0(n))$. If n is square-free, the functions f_k already form a basis of $\mathcal{E}_2(\Gamma_0(n))$. The expansion of $E_2^{(k)}(\tau)$ at a cusp $s = a/c$ is given by

$$E_2^{(k)}|_{M_s}(\tau) = \frac{(k, c)^2}{k^2} \left(1 - 24 \sum_{m=1}^{\infty} \sigma(m) e(mv/w) q^{m(k, c)^2/k} \right) + \dots$$

where we omitted the non-modular term, which cancels out in f_k . This is an expansion in q^{1/t_s} where $t_s = n/(c^2, n)$ is the width of the cusp s since $k/(c^2, k) \mid t_s$. It follows that the constant term of f_k at the cusp $s = a/c$ is

$$[f_k|_{M_s}](0) = 1 - \frac{(k, c)^2}{k}.$$

We construct an Eisenstein series with certain constant terms at the cusps:

Proposition 4.2. *Suppose $n > 1$. Then there is a unique $f \in \mathcal{E}_2(\Gamma_0(n))$ satisfying*

$$[f|_{M_s}](0) = 1$$

for all cusps $s \neq \infty$. This function can be written as

$$f(\tau) = \frac{1}{\phi(n)} \sum_{\substack{k|n \\ k>1}} k\mu(n/k) f_k(\tau) = \frac{1}{\phi(n)} \sum_{\substack{k|n \\ k>1}} k\mu(n/k) (E_2(\tau) - kE_2(k\tau))$$

and the constant coefficient of the Fourier expansion of f at ∞ is given by

$$[f](0) = 1 - \psi(n).$$

Proof. Let S be the set of cusps of $\Gamma_0(n)$ without ∞ . Then $\mathcal{E}_2(\Gamma_0(n))$ has a basis consisting of Eisenstein series g_s , $s \in S$, which are 1 at s and vanish at the other cusps in S . This implies the first statement. Now let

$$f(\tau) = \frac{1}{\phi(n)} \sum_{\substack{k|n \\ k>1}} k\mu(n/k) f_k(\tau)$$

and $s = a/c$ with $(c, n) < n$. Then

$$\begin{aligned} [f|_{M_s}](0) &= \frac{1}{\phi(n)} \sum_{\substack{k|n \\ k>1}} k\mu(n/k) [f_k|_{M_s}](0) = \frac{1}{\phi(n)} \sum_{\substack{k|n \\ k>1}} k\mu(n/k) \left(1 - \frac{(k, c)^2}{k} \right) \\ &= \frac{1}{\phi(n)} \sum_{k|n} k\mu(n/k) - \frac{1}{\phi(n)} \sum_{k|n} \mu(n/k) (k, c)^2 \\ &= 1 - 0 = 1. \end{aligned}$$

This proves the second statement. The third follows from the residue theorem. $\square$

The Eisenstein series f in the previous proposition can also be written as

$$f(\tau) = E_2(\tau) - \frac{1}{\phi(n)} \sum_{k|n} k^2 \mu(n/k) E_2(k\tau)$$

and its Fourier expansion at a cusp $s = a/c$ is

$$\begin{aligned} f|_{M_s}(\tau) &= 1 - 24 \sum_{m=1}^{\infty} \sigma(m) q^m \\ &\quad - \frac{1}{\phi(n)} \sum_{k|n} \mu(n/k) (k, c)^2 \left(1 - 24 \sum_{m=1}^{\infty} \sigma(m) e(mv/w) q^{m(k, c)^2/k} \right) \end{aligned}$$

with $M_s = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ and v, w as in Proposition 4.1.

We now consider certain cusp forms. Let p be a prime and $W_p := \frac{1}{\sqrt{p}} \begin{pmatrix} 0 & -1 \\ p & 0 \end{pmatrix}$. Then the Fricke involution

$$W_p : \mathcal{S}_2(\Gamma_0(p)) \rightarrow \mathcal{S}_2(\Gamma_0(p)), \quad f \mapsto f|_{W_p}$$

is diagonalisable with eigenvalues ± 1 . The corresponding eigenspace decomposition

$$\mathcal{S}_2(\Gamma_0(p)) = \mathcal{S}_2(\Gamma_0(p))^+ \oplus \mathcal{S}_2(\Gamma_0(p))^-$$

is orthogonal with respect to the Petersson inner product. Let $\Gamma_0(p)^+ := \Gamma_0(p) \cup W_p \Gamma_0(p)$ and $X_0(p)^+ := \Gamma_0(p)^+ \backslash H^*$.

Proposition 4.3. *Let p be a prime such that $X_0(p)$ has genus $g(X_0(p)) > 0$. Then $\mathcal{S}_2^-(\Gamma_0(p))$ is non-trivial.*

Proof. By a classical result of Fricke (see [Fri11], p. 366)

$$g(X_0(p)^+) = \frac{1}{2}(g(X_0(p)) + 1) - \varepsilon_p h(p)$$

for all primes $p \geq 5$ where $h(p)$ denotes the class number of $\mathbb{Q}(\sqrt{-p})$ and

$$\varepsilon_p = \begin{cases} \frac{1}{4} & \text{if } p \equiv 1, 5 \pmod{8}, \\ \frac{1}{2} & \text{if } p \equiv 7 \pmod{8}, \\ 1 & \text{if } p \equiv 3 \pmod{8}. \end{cases}$$

Hence

$$\begin{aligned} \dim(\mathcal{S}_2(\Gamma_0(p))^-) &= \dim(\mathcal{S}_2(\Gamma_0(p))) - \dim(\mathcal{S}_2(\Gamma_0(p))^+) \\ &= g(X_0(p)) - g(X_0(p)^+) \\ &= \frac{1}{2}(g(X_0(p)) - 1) + \varepsilon_p h(p) > 0 \end{aligned}$$

if $g(X_0(p)) > 0$. □

The space $\mathcal{S}_2(\Gamma_0(p))^-$ has a basis consisting of simultaneous eigenforms for all Hecke operators T_m , $m \in \mathbb{Z}_{>0}$ (see [AL70] and Theorem 9.27 in [Kna92]).

We state *Deligne's bound* on the growth of the coefficients of cusp forms:

Theorem 4.4. *Let p be a prime such that $g(X_0(p)) > 0$. Let $f \in \mathcal{S}_2(\Gamma_0(p))^-$ be a normalised eigenform, i.e. $f(\tau) = \sum_{m=1}^{\infty} a(m)q^m$ with $a(1) = 1$ and $T_m f = a(m)f$ for all $m \in \mathbb{Z}_{>0}$. Then*

- (1) $a(p^k) = 1$ for all $k \in \mathbb{Z}_{>0}$,
- (2) $a(2), \dots, a(p-1) \in \mathbb{R}$,
- (3) $|a(m)| \leq \sigma_0(m)\sqrt{m}$ for all $m \in \mathbb{Z}_{>0}$.

Proof. By Theorem 9.27 in [Kna92], $a(p) = 1$ and $a(p^k) = a(p)^k = 1$ for all $k \in \mathbb{Z}_{>0}$. Since the Hecke operators T_m for $(m, p) = 1$ are self-adjoint with respect to the Petersson inner product, the coefficients $a(2), \dots, a(p-1)$ are real (see Theorem 9.18 in [Kna92]). Finally the L -function of f is of the form

$$L(f, s) = \frac{1}{1 - a(p)p^{-s}} \prod_{\substack{t \text{ prime} \\ t \neq p}} \frac{1}{1 - a(t)t^{-s} + t^{1-2s}}.$$

Deligne has shown (see [Del74], Théorème 8.2) that the reciprocal roots of the polynomial $1 - a(t)X + tX^2$ have absolute value $\sqrt{t}$, i.e. that

$$1 - a(t)X - tX^2 = (1 - \alpha_t X)(1 - \bar{\alpha}_t X)$$

with $|\alpha_t| = \sqrt{t}$. Writing $(1 - a(t)t^{-s} + t^{1-2s})^{-1} = (1 - \alpha_t t^{-s})^{-1}(1 - \bar{\alpha}_t t^{-s})^{-1}$ and expanding each factor in a geometric series we find that the coefficient of t^{-ms} is

given by $\alpha_t^m + \alpha_t^{m-1}\bar{\alpha}_t + \dots + \bar{\alpha}_t^m$. This means that if the positive integer m with $(m, p) = 1$ has prime factorisation $m = t_1^{e_1} \dots t_r^{e_r}$, then

$$a(m) = \prod_{j=1}^r (\alpha_{t_j}^{e_j} + \alpha_{t_j}^{e_j-1}\bar{\alpha}_{t_j} + \dots + \bar{\alpha}_{t_j}^{e_j}),$$

implying

$$|a(m)| \leq \prod_{j=1}^r (e_j + 1) t_j^{e_j/2} = \sigma_0(m) \sqrt{m}.$$

The general case follows from the multiplicativity of the $a(m)$ and the first statement. $\square$

4.2. Modular Forms for the Weil Representation. In this section we recall some properties of modular forms for the Weil representation of Γ from [Sch09].

Let D be a discriminant form with quadratic form $q: D \rightarrow \mathbb{Q}/\mathbb{Z}$ and associated bilinear form $(\cdot, \cdot): D \times D \rightarrow \mathbb{Q}/\mathbb{Z}$ (see, e.g., [Nik80, CS99]). The level of D is the smallest positive integer n such that $nq(\gamma) = 0 \pmod{1}$ for all $\gamma \in D$.

Let c be an integer. Then c acts by multiplication on D and there is an exact sequence $0 \rightarrow D_c \rightarrow D \rightarrow D^c \rightarrow 0$ where D_c is the kernel and D^c the image of this map. The group D^c is the orthogonal complement of D_c .

Let D^{c*} be the set of elements $\alpha \in D$ satisfying $cq(\gamma) + (\alpha, \gamma) = 0 \pmod{1}$ for all $\gamma \in D_c$. Then D^{c*} is a coset of D^c . Once we have chosen a Jordan decomposition of D , there is a canonical coset representative x_c of D^{c*} satisfying $2x_c = 0$. Write $\alpha \in D^{c*}$ as $\alpha = x_c + c\gamma$. Then $q_c(\alpha) := cq(\gamma) + (x_c, \gamma) \pmod{1}$ is independent of the choice of γ . This gives a well-defined map $q_c: D^{c*} \rightarrow \mathbb{Q}/\mathbb{Z}$.

If D has no odd 2-adic Jordan components, then $x_c = 0$ so that $D^c = D^{c*}$ and $q_c(\gamma) = cq(\mu) \pmod{1}$ for $\gamma = c\mu \in D^c$.

Suppose D has even signature. We define a scalar product $(\cdot, \cdot)$ on the group ring $\mathbb{C}[D]$ which is linear in the first and antilinear in the second variable by $(e^\gamma, e^\beta) := \delta_{\gamma, \beta}$. Then there is a unitary action ρ_D of the group Γ on $\mathbb{C}[D]$ satisfying

$$\begin{aligned} \rho_D(T)e^\gamma &= e(q(\gamma))e^\gamma, \\ \rho_D(S)e^\gamma &= \frac{e(\text{sign}(D)/8)}{\sqrt{|D|}} \sum_{\beta \in D} e(-(\gamma, \beta))e^\beta \end{aligned}$$

where $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ are the standard generators of Γ . The representation ρ_D is called the *Weil representation* of Γ on $\mathbb{C}[D]$. It commutes with $O(D)$. We also consider the *dual Weil representation* $\bar{\rho}_D$, the complex conjugate of ρ_D .

Suppose that the level of D divides n . We define a quadratic Dirichlet character $\chi_D: \Gamma_0(n) \rightarrow \{\pm 1\}$ by

$$\chi_D(M) := \left(\frac{a}{|D|} \right) e((a-1)\text{odddity}(D)/8)$$

for $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(n)$. Then $M \in \Gamma_0(n)$ acts in Weil representation ρ_D as

$$\rho_D(M)e^\gamma = \chi_D(M)e(bdq(\gamma))e^{d\gamma}.$$

A general formula for $M \in \Gamma$ is given in [Sch09], Theorem 4.7.

Let

$$F(\tau) = \sum_{\gamma \in D} F_\gamma(\tau)e^\gamma$$

be a holomorphic function on the upper half-plane H with values in $\mathbb{C}[D]$ and $k \in \mathbb{Z}$. Then F is a modular form for ρ_D of weight k if

$$F(M\tau) = (c\tau + d)^k \rho_D(M)F(\tau)$$

for all $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in Γ and F is meromorphic at the cusp ∞ . We call F *holomorphic* if it is also holomorphic at ∞ .

Classical examples of modular forms transforming under the Weil representation are theta series. Let L be a positive-definite, even lattice of even rank $2k$ with bilinear form $(\cdot, \cdot): L \times L \rightarrow \mathbb{Z}$ and let $D = L'/L$ be the associated discriminant form. Define

$$\theta = \sum_{\gamma \in D} \theta_{\gamma} e^{\gamma} \quad \text{with} \quad \theta_{\gamma}(\tau) := \sum_{\alpha \in \gamma + L} q^{(\alpha, \alpha)/2}$$

for all $\gamma \in D$. Then θ is a modular form for the Weil representation ρ_D of weight k which is holomorphic at ∞ .

Another example arises from orbifold theory (see Section 3.3). Let V be a strongly rational, holomorphic vertex operator algebra of central charge c and g an automorphism of V of finite order n and type 0 such that V^g satisfies the positivity condition. Suppose $24 \mid c$. Then the irreducible V^g -modules are given by W^{γ} , $\gamma \in D$, with conformal weights $\rho(W^{\gamma}) = q(\gamma) \bmod 1$ where the discriminant form $D = \mathbb{Z}_n \times \mathbb{Z}_n$ has quadratic form $q((i, j)) = ij/n \bmod 1$. Note that D has order n^2 and level n . The vector-valued character $\text{Ch}_{V^g}: H \rightarrow \mathbb{C}[D]$ defined by

$$\text{Ch}_{V^g}(\tau) = \sum_{\gamma \in D} \text{ch}_{W^{\gamma}}(\tau) e^{\gamma} \quad \text{with} \quad \text{ch}_{W^{\gamma}}(\tau) = \text{tr} |_{W^{\gamma}} q^{L_0 - c/24}$$

is a modular form of weight 0 for the Weil representation ρ_D of Γ , holomorphic on H with a possible pole at the cusp ∞ .

Let D be a discriminant form of even signature and n a positive integer such that the level of D divides n . Let $F = \sum_{\gamma \in D} F_{\gamma} e^{\gamma}$ be a modular form for ρ_D . Then the formula for ρ_D shows that F_0 is a modular form for $\Gamma_0(n)$ of character χ_D . Conversely:

Proposition 4.5. *Let D be a discriminant form of even signature and level dividing n . Let f be a scalar-valued modular form for $\Gamma_0(n)$ of weight k and character χ_D . Then*

$$F = \sum_{M \in \Gamma_0(n) \backslash \Gamma} f|_M \rho_D(M^{-1}) e^0$$

is a modular form for ρ_D of weight k which is invariant under $O(D)$.

This is Theorem 6.2 in [Sch06] in the special case of trivial support. An explicit description of the components of the lift F is given in [Sch09], Theorem 5.7 (and for n square-free in [Sch06], Theorem 6.5). An analogous statement holds if we replace ρ_D by the dual Weil representation $\bar{\rho}_D$.

The dimension formulae proved in this text are instances of the following pairing argument:

Proposition 4.6. *Let D be a discriminant form of even signature and G a modular form for ρ_D of weight $2 - k$, $k \in \mathbb{Z}$. Let F be a modular form for $\bar{\rho}_D$ of weight k . Then the constant coefficient of $(G, \bar{F}) = \sum_{\gamma \in D} G_{\gamma} \bar{F}_{\gamma}$ vanishes.*

Proof. The unitarity of ρ_D with respect to $(\cdot, \cdot)$ implies that $(G, \bar{F})$ is a scalar-valued modular form for Γ of weight 2 which is holomorphic on H and meromorphic at ∞ . Hence $(G, \bar{F})d\tau$ defines a 1-form on the Riemann sphere with a pole at ∞ . By the residue theorem its residue has to vanish. Since $d\tau = \frac{1}{2\pi i} \frac{dq}{q}$, the constant term in the Fourier expansion of $(G, \bar{F})$ must be 0. $\square$

4.3. Vector-Valued Eisenstein Series. In this section we construct an Eisenstein series F for the dual Weil representation of $II_{1,1}(n)$ by lifting the Eisenstein series $f \in \mathcal{E}_2(\Gamma_0(n))$ from Proposition 4.2 and calculate some of its Fourier coefficients. Pairing F with the character of V^g will give the first dimension formula.

Let n be a positive integer. Consider the discriminant form $D = (\mathbb{Z}_n \times \mathbb{Z}_n, q)$ with quadratic form $q((i, j)) = ij/n \pmod{1}$. This is the discriminant form of the hyperbolic lattice $II_{1,1}(n)$ and of the fusion algebra of the orbifold vertex operator algebra V^g described above. We note that the character χ_D is trivial because D has no odd 2-adic components and its order is n^2 .

Let f be a modular form for $\Gamma_0(n)$ of weight 2 with trivial character. Then Proposition 4.5 allows us to lift f to a vector-valued modular form F for the dual Weil representation $\bar{\rho}_D$. We describe the components of F . The expansion of f at a cusp $s \in \Gamma_0(n) \backslash P$ is an expansion in q^{1/t_s} where t_s denotes the width of s . We decompose

$$f|_{M_s}(\tau) = g_{t_s,0}(\tau) + \dots + g_{t_s,t_s-1}(\tau)$$

with $g_{t_s,j}|_T = e(j/t_s)g_{t_s,j}$ for $j \in \mathbb{Z}_{t_s}$. In general, the terms in this decomposition depend on the choice of M_s but $g_{t_s,0}$ does not. The special case of Theorem 5.7 in [Sch09] for trivial support applied to $D = (\mathbb{Z}_n \times \mathbb{Z}_n, q)$ gives:

Proposition 4.7. *Let $f \in \mathcal{M}_2(\Gamma_0(n))$ and F be the lift (from Proposition 4.5) of f to a vector-valued modular form for the dual Weil representation $\bar{\rho}_D$ where $D = (\mathbb{Z}_n \times \mathbb{Z}_n, q)$. Then*

$$F_\gamma(\tau) = \sum_{s \in \Gamma_0(n) \backslash P} \delta_{\gamma \in D^c} \frac{1}{((c, n), n/(c, n))} e(dq_c(\gamma)) g_{t_s, j_\gamma}(\tau)$$

for all $\gamma \in D$ where $M_s = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ is such that $M_s \infty = s$ and $j_\gamma \in \{0, \dots, t_s - 1\}$ such that $j_\gamma/t_s = -q(\gamma) \pmod{1}$.

Recall that since D has no odd 2-adic components, $q_c(\gamma) = cq(\mu) \pmod{1}$ for any $\mu \in D$ such that $\gamma = c\mu$.

For the remainder of this section, let $F = \sum_{\gamma \in D} F_\gamma e^\gamma$ be the lift of the Eisenstein series $f \in \mathcal{E}_2(\Gamma_0(n))$ from Proposition 4.2 to a modular form for $\bar{\rho}_D$. We calculate the lowest-order Fourier coefficients of the components F_γ and $[F_0](1)$.

For $k \in \mathbb{Z}_{>0}$ and $m \in \mathbb{Z}_k$ we define the *Ramanujan sum*

$$C_k(m) := \sum_{a \in \mathbb{Z}_k^*} e(am/k) = \sum_{c|(k,m)} c\mu(k/c).$$

Proposition 4.8. *Let $\gamma \in D$ with $q(\gamma) = 0 \pmod{1}$. Then the constant term in the Fourier expansion of F_γ is*

$$[F_\gamma](0) = -\delta_{\gamma,0}\psi(n) + \sum_{c|n} \delta_{\gamma \in D^c} \frac{1}{(c, n/c)} C_{(c, n/c)}((c, n/c)q_c(\gamma)).$$

Proof. Let $\gamma \in D$ with $q(\gamma) = 0 \pmod{1}$. Then Proposition 4.7 implies that

$$\begin{aligned} [F_\gamma](0) &= \sum_{s \in \Gamma_0(n) \backslash P} \delta_{\gamma \in D^c} \frac{1}{((c, n), n/(c, n))} e(dq_c(\gamma)) [g_{t_s,0}](0) \\ &= \sum_{s \in \Gamma_0(n) \backslash P} \delta_{\gamma \in D^c} \frac{1}{((c, n), n/(c, n))} e(dq_c(\gamma)) [f|_{M_s}](0) \\ &= \sum_{s \in \Gamma_0(n) \backslash P} \delta_{\gamma \in D^c} \frac{1}{((c, n), n/(c, n))} e(dq_c(\gamma)) - \delta_{\gamma,0}\psi(n) \end{aligned}$$

where for each cusp $s \in \Gamma_0(n) \backslash P$ we have chosen a matrix $M_s = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ such that $M_s \infty = s$. The sum can be partially carried out to remove the complex factor $e(dq_c(\gamma))$. Indeed, choosing the system of representatives for the cusps described in Section 4.1 we obtain

$$\begin{aligned} [F_\gamma](0) &= \sum_{c|n} \delta_{\gamma \in D^c} \frac{1}{(c, n/c)} \sum_{a \in \mathbb{Z}_{(c, n/c)}^*} e(dq_c(\gamma)) - \delta_{\gamma, 0} \psi(n) \\ &= \sum_{c|n} \delta_{\gamma \in D^c} \frac{1}{(c, n/c)} \sum_{a \in \mathbb{Z}_{(c, n/c)}^*} e(aq_c(\gamma)) - \delta_{\gamma, 0} \psi(n) \end{aligned}$$

since $q_c(\gamma) = 0 \pmod{1/c}$ and $ad = 1 \pmod{c}$. Note that $(c, n/c)q_c(\gamma) = 0 \pmod{1}$ if $\gamma \in D^c$ and $q(\gamma) = 0 \pmod{1}$. Hence

$$\begin{aligned} [F_\gamma](0) &= \sum_{c|n} \delta_{\gamma \in D^c} \frac{1}{(c, n/c)} \sum_{a \in \mathbb{Z}_{(c, n/c)}^*} e\left(\frac{a(c, n/c)q_c(\gamma)}{(c, n/c)}\right) - \delta_{\gamma, 0} \psi(n) \\ &= \sum_{c|n} \delta_{\gamma \in D^c} \frac{1}{(c, n/c)} C_{(c, n/c)}((c, n/c)q_c(\gamma)) - \delta_{\gamma, 0} \psi(n). \quad \square \end{aligned}$$

For $c \mid n$ we define the isotropic subgroup H_c of D by

$$H_c := \{(ci, nj/c) \mid i, j \in \mathbb{Z}_n\} \cong \mathbb{Z}_{n/c} \times \mathbb{Z}_c.$$

Then we can rewrite the Fourier coefficients $[F_\gamma](0)$ for isotropic $\gamma \in D$ as follows:

Proposition 4.9. *Let $\gamma \in D$ with $q(\gamma) = 0 \pmod{1}$. Then*

$$[F_\gamma](0) = -\delta_{\gamma, 0} \psi(n) + \sum_{c|n} \delta_{\gamma \in H_c} \frac{\phi((c, n/c))}{(c, n/c)}.$$

Recall that we extended the sum-of-divisors function σ to the positive rational numbers by $\sigma(r) = 0$ if $r \notin \mathbb{Z}_{>0}$.

Proposition 4.10. *The coefficient at q in the Fourier expansion of F_0 is given by*

$$[F_0](1) = 24 - 24 \sum_{c|n} \frac{\phi((c, n/c))}{(c, n/c)}.$$

Proof. Proposition 4.7 implies that

$$F_0(\tau) = \sum_{s \in \Gamma_0(n) \backslash P} \frac{1}{((c, n), n/(c, n))} g_{t_s, 0}(\tau).$$

From the Fourier expansion of f at s calculated above we obtain

$$\begin{aligned} [g_{t_s, 0}](1) &= [f|_{M_s}](1) \\ &= -24 + \frac{24}{\phi(n)} \sum_{k|n} \mu(n/k)(k, c)^2 \sigma\left(\frac{k}{(k, c)^2}\right) e\left(\frac{v}{(k, c)}\right) \end{aligned}$$

with $v \in \mathbb{Z}$ such that $v = (c/(k, c))^{-1}d \pmod{k/(k, c)}$ and the inverse taken modulo $k/(k, c)$ (see Proposition 4.1). It suffices to show that

$$\sum_{s \in \Gamma_0(n) \backslash P} \frac{1}{((c, n), n/(c, n))} \sum_{k|n} \mu(n/k)(k, c)^2 \sigma\left(\frac{k}{(k, c)^2}\right) e\left(\frac{v}{(k, c)}\right) = \phi(n).$$

As in the proof of Proposition 4.8, we partially carry out the sum over the cusps to remove the complex factors. Since $(c/(k, c))^{-1}/(k, c) = 0 \pmod{1/c}$ and $ad = 1 \pmod{c}$, we obtain

$$\begin{aligned} & \sum_{c|n} \frac{1}{(c, n/c)} \sum_{k|n} \mu(n/k)(k, c)^2 \sigma \left(\frac{k}{(k, c)^2} \right) \sum_{a \in \mathbb{Z}_{(c, n/c)}^*} e \left(\frac{d(c/(k, c))^{-1}}{(k, c)} \right) \\ &= \sum_{c|n} \frac{1}{(c, n/c)} \sum_{k|n} \mu(n/k)(k, c)^2 \sigma \left(\frac{k}{(k, c)^2} \right) \sum_{a \in \mathbb{Z}_{(c, n/c)}^*} e \left(\frac{a(c/(k, c))^{-1}}{(k, c)} \right). \end{aligned}$$

Now $c \mid n$, $k \mid n$, n/k square-free and $(k, c)^2 \mid k$ imply $(k, c) = (c, n/c)$. Then the last sum becomes

$$\sum_{a \in \mathbb{Z}_{(c, n/c)}^*} e \left(\frac{a(c/(k, c))^{-1}}{(c, n/c)} \right) = \sum_{a \in \mathbb{Z}_{(c, n/c)}^*} e \left(\frac{a}{(c, n/c)} \right) = C_{(c, n/c)}(1) = \mu((c, n/c))$$

because $(c, n/c)$ and $(c/(k, c))^{-1}$ are coprime. Overall, the expression of interest simplifies to

$$\sum_{c|n} \frac{1}{(c, n/c)} \sum_{k|n} \mu(n/k)(k, c)^2 \sigma \left(\frac{k}{(k, c)^2} \right) \mu((c, n/c)),$$

which is multiplicative in n , as is $\phi(n)$. We verify that both expressions coincide for prime powers, which concludes the proof. $\square$

For $\gamma \in D$ with $q(\gamma) \neq 0 \pmod{1}$ we introduce $r_\gamma \in (0, 1)$ such that $r_\gamma = -q(\gamma) \pmod{1}$.

Proposition 4.11. *Let $\gamma \in D$ with $q(\gamma) \neq 0 \pmod{1}$. Then*

$$\begin{aligned} [F_\gamma](r_\gamma) &= \frac{24}{\phi(n)} \sum_{c|n} \delta_{\gamma \in D^c} \frac{1}{(c, n/c)} \sum_{k|n} \mu(n/k)(k, c)^2 \times \\ &\quad \times \sigma \left(r_\gamma \frac{k}{(k, c)^2} \right) C_{(c, n/c)} \left((c, n/c) \left(q_c(\gamma) + r_\gamma \frac{(c/(k, c))^{-1}}{(k, c)} \right) \right) \end{aligned}$$

where the inverse is taken modulo $k/(k, c)$.

Proof. By Proposition 4.7

$$[F_\gamma](r_\gamma) = \sum_{s \in \Gamma_0(n) \setminus P} \delta_{\gamma \in D^c} \frac{1}{((c, n), n/(c, n))} e(dq_c(\gamma)) [f|_{M_s}](r_\gamma)$$

with $M_s = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ such that $M_s \infty = s$. The formula for the Fourier expansion of f at s gives

$$[f|_{M_s}](r_\gamma) = \frac{24}{\phi(n)} \sum_{k|n} \mu(n/k)(k, c)^2 \sigma \left(r_\gamma \frac{k}{(k, c)^2} \right) e \left(r_\gamma \frac{v}{(k, c)} \right)$$

where $v = d(c/(k, c))^{-1}$ with inverse taken modulo $k/(k, c)$. Hence,

$$\begin{aligned} [F_\gamma](r_\gamma) &= \frac{24}{\phi(n)} \sum_{s \in \Gamma_0(n) \setminus P} \delta_{\gamma \in D^c} \frac{1}{((c, n), n/(c, n))} \sum_{k|n} \mu(n/k)(k, c)^2 \times \\ &\quad \times \sigma \left(r_\gamma \frac{k}{(k, c)^2} \right) e \left(d \left(q_c(\gamma) + r_\gamma \frac{(c/(k, c))^{-1}}{(k, c)} \right) \right). \end{aligned}$$

Again, we take the standard representatives of the cusps and note that $q_c(\gamma) + r_\gamma(c/(k, c))^{-1}/(k, c) = 0 \pmod{1/c}$ and $ad = 1 \pmod{c}$. Then

$$\begin{aligned} [F_\gamma](r_\gamma) &= \frac{24}{\phi(n)} \sum_{c|n} \delta_{\gamma \in D^c} \frac{1}{(c, n/c)} \sum_{k|n} \mu(n/k)(k, c)^2 \times \\ &\quad \times \sigma \left(r_\gamma \frac{k}{(k, c)^2} \right) \sum_{a \in \mathbb{Z}_{(c, n/c)}^*} e(ay_{n, \gamma}(c, k)/(c, n/c)) \end{aligned}$$

with

$$y_{n, \gamma}(c, k) := (c, n/c) \left(q_c(\gamma) + r_\gamma \frac{(c/(k, c))^{-1}}{(k, c)} \right)$$

for $c | n$, $k | n$, $\gamma \in D^c$ and $r_\gamma k/(k, c)^2 \in \mathbb{Z}$. One can show that $y_{n, \gamma}(c, k) = 0 \pmod{1}$. Hence,

$$\sum_{a \in \mathbb{Z}_{(c, n/c)}^*} e(ay_{n, \gamma}(c, k)/(c, n/c)) = C_{(c, n/c)}(y_{n, \gamma}(c, k)),$$

which implies the assertion. $\square$

We shall see in Proposition 5.4 that the coefficients $[F_\gamma](r_\gamma)$ are positive.

5. DIMENSION FORMULAE

In this section we prove a formula for the dimension $\dim(V_1^{\text{orb}(g)})$ of the weight-1 space of the orbifold construction in central charge 24, depending on the dimensions of the fixed-point Lie subalgebras $\dim(V_1^{g^m})$, $m | n$, where n is the order of g , and terms of lower L_0 -weight.

This generalises previous results for $n = 2, 3$ in [Mon94] (see also [LS19]), for $n = 5, 7, 13$ in [Möl16] and for all $n > 1$ such that the modular curve $X_0(n)$ has genus 0 in [EMS20b]. However, our approach is fundamentally different.

5.1. First Dimension Formula. The first dimension formula will be proved using the pairing argument described in Proposition 4.6.

Let V be a strongly rational, holomorphic vertex operator algebra V of central charge 24 and $g \in \text{Aut}(V)$ of finite order n and type 0. Suppose that the fixed-point subalgebra V^g satisfies the positivity condition. Then V^g has fusion algebra $\mathbb{C}[D]$ with discriminant form $D = (\mathbb{Z}_n \times \mathbb{Z}_n, q)$ where $q((i, j)) = ij/n \pmod{1}$. In particular, the irreducible V^g -modules are given by W^γ , $\gamma \in D$, with conformal weights $\rho(W^\gamma) = q(\gamma) \pmod{1}$ (see Section 3.3). The character

$$\text{Ch}_{V^g}(\tau) = \sum_{\gamma \in D} \text{ch}_{W^\gamma}(\tau) e^\gamma$$

is a vector-valued modular form of weight 0 for the Weil representation ρ_D of Γ , holomorphic on H and with a pole at the cusp ∞ (see Section 4.2). The lowest-order terms in the Fourier expansion of Ch_{V^g} are

$$\text{ch}_{W^\gamma}(\tau) = \begin{cases} q^{-1} + \dim(V_1^g) + O(q) & \text{if } \gamma = 0, \\ \dim(W_1^\gamma) + O(q) & \text{if } \gamma \neq 0 \text{ and } q(\gamma) = 0 \pmod{1}, \\ \dim(W_{1-r_\gamma}^\gamma) q^{-r_\gamma} + O(q^{1-r_\gamma}) & \text{if } \gamma \neq 0 \text{ and } q(\gamma) \neq 0 \pmod{1} \end{cases}$$

because $W^0 = W^{(0,0)} = V^g$ is of CFT type and V^g satisfies the positivity condition. Recall that $r_\gamma \in (0, 1)$ is chosen such that $r_\gamma = -q(\gamma) \pmod{1}$.

Pairing Ch_{V^g} with a holomorphic vector-valued modular form F of weight 2 for the dual Weil representation $\bar{\rho}_D$ (see Proposition 4.6) we obtain:

Proposition 5.1. *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of finite order n and type 0 such that V^g satisfies the positivity condition. Suppose F is a holomorphic modular form of weight 2 for the dual Weil representation $\bar{\rho}_D$. Then*

$$[F_0](1) + \sum_{\substack{\gamma \in D \\ q(\gamma) \equiv 0 \pmod{1}}} [F_\gamma](0) \dim(W_1^\gamma) + \sum_{\substack{\gamma \in D \\ q(\gamma) \not\equiv 0 \pmod{1}}} [F_\gamma](r_\gamma) \dim(W_{1-r_\gamma}^\gamma) = 0.$$

We emphasise that any modular form $F \neq 0$ imposes a restriction on the character Ch_{V^g} . Our first dimension formula is obtained by choosing for F the lift of the Eisenstein series from Proposition 4.2,

$$f(\tau) = E_2(\tau) - \frac{1}{\phi(n)} \sum_{k|n} k^2 \mu(n/k) E_2(k\tau).$$

Theorem 5.2 (First Dimension Formula). *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of finite order $n > 1$ and type 0 such that V^g satisfies the positivity condition. Then*

$$\psi(n) \dim(V_1^g) - \sum_{c|n} \frac{\phi((c, n/c))}{(c, n/c)} \dim(V_1^{\text{orb}(g^c)}) = 24 - 24 \sum_{c|n} \frac{\phi((c, n/c))}{(c, n/c)} + \tilde{R}(g)$$

with rest term

$$\tilde{R}(g) = \sum_{\substack{\gamma \in D \\ q(\gamma) \not\equiv 0 \pmod{1}}} \tilde{d}_n(\gamma) \dim(W_{1-r_\gamma}^\gamma)$$

where $\tilde{d}_n(\gamma)$ equals the expression $[F_\gamma](r_\gamma)$ from Proposition 4.11.

Proof. Let F be the lift of f . Proposition 4.9 and $V^{\text{orb}(g^c)} = \bigoplus_{\gamma \in H_c} W^\gamma$ imply

$$\sum_{\substack{\gamma \in D \\ q(\gamma) \equiv 0 \pmod{1}}} [F_\gamma](0) \dim(W_1^\gamma) = \sum_{c|n} \frac{\phi((c, n/c))}{(c, n/c)} \dim(V_1^{\text{orb}(g^c)}) - \psi(n) \dim(V_1^g).$$

Finally, Proposition 5.1 together with Proposition 4.10 gives

$$\psi(n) \dim(V_1^g) - \sum_{c|n} \frac{\phi((c, n/c))}{(c, n/c)} \dim(V_1^{\text{orb}(g^c)}) = 24 - 24 \sum_{c|n} \frac{\phi((c, n/c))}{(c, n/c)} + \tilde{R}(g)$$

with $\tilde{R}(g)$ as stated. $\square$

The formula extends to $n = 1$ if we define $\tilde{R}(g) = 0$ in this case.

The rest term $\tilde{R}(g)$ contains exactly the terms involving the dimensions of the weight spaces of the irreducible V^g -modules of weight less than 1. We shall show that $\tilde{R}(g) \geq 0$ (see Proposition 5.4).

Specialising to $n = 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 16, 18, 25$, i.e. those $n > 1$ for which the modular curve $X_0(n)$ has genus 0, the result becomes Theorem 3.1 in [EMS20b], which was proved using expansions of Hauptmoduln.

In the case that the order of g is a prime p such that $X_0(p)$ has positive genus we shall see that $\tilde{R}(g) \geq 24$. This follows from pairing Ch_{V^g} with the lift of a Hecke eigenform in $\mathcal{S}_2(\Gamma_0(n))^-$.

5.2. Second Dimension Formula. Linearly combining the dimension formula in Theorem 5.2 applied to the powers $g^{n/m}$ of g with $m \mid n$ we obtain the second dimension formula, a closed formula for $\dim(V_1^{\text{orb}(g)})$. This will be the main result of this section.

We introduce the arithmetic function λ by

$$\lambda(n) := \prod_{\substack{p|n \\ p \text{ prime}}} (-p)$$

for $n \in \mathbb{Z}_{>0}$ and define

$$\begin{aligned} \xi_n(m) &:= \frac{\lambda(n/m)}{n/m} \frac{\phi((m, n/m))}{(m, n/m)}, \\ c_n(m) &:= \xi_n(n/m) \psi(n/m) = \frac{\lambda(m)}{m} \frac{\phi((m, n/m))}{(m, n/m)} \psi(n/m) \end{aligned}$$

for $n \in \mathbb{Z}_{>0}$ and $m \mid n$ (see also Section 2.3). Note that for a given $n \in \mathbb{Z}_{>0}$, the numbers $\xi_n(m)$ and $c_n(m)$ for $m \mid n$ can be equivalently specified by the systems of linear equations

$$\sum_{t|m|n} \xi_n(m) \frac{\phi((t, m/t))}{(t, m/t)} = \delta_{t,n}$$

(so that in particular $\sum_{m|n} \xi_n(m) = \delta_{1,n}$) and

$$\sum_{m|n} c_n(m)(t, m) = n/t$$

for all $t \mid n$.

Taking the linear combination of the dimension formulae in Theorem 5.2 for $g^{n/m}$ with coefficients $\xi_n(m)$ for $m \mid n$ yields:

Theorem 5.3 (Second Dimension Formula). *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of order $n > 1$ and type 0 such that V^g satisfies the positivity condition. Then*

$$\dim(V_1^{\text{orb}(g)}) = 24 + \sum_{m|n} c_n(m) \dim(V_1^{g^m}) - R(g)$$

with rest term

$$R(g) = \sum_{m|n} \xi_n(n/m) \tilde{R}(g^m) = \sum_{\substack{\gamma \in D \\ q(\gamma) \not\equiv 0 \pmod{1}}} d_n(\gamma) \dim(W_{1-r_\gamma}^\gamma)$$

where for $\gamma = (i, j) \in D$ such that $q(\gamma) \not\equiv 0 \pmod{1}$

$$d_n(\gamma) = \sum_{m|(i,n)} \xi_n(n/m) \tilde{d}_{n/m}(i/m, j).$$

Proof. The dimension formula in Theorem 5.2 applied to $g^{n/m}$ of order m for $m \mid n$ reads

$$\begin{aligned} \psi(m) \dim(V_1^{g^{n/m}}) - \sum_{c|m} \frac{\phi((c, m/c))}{(c, m/c)} \dim(V_1^{\text{orb}(g^{nc/m})}) \\ = 24 - 24 \sum_{c|m} \frac{\phi((c, m/c))}{(c, m/c)} + \tilde{R}(g^{n/m}). \end{aligned}$$

Summing this equation over $m \mid n$ with coefficients $\xi_n(m)$ yields

$$\begin{aligned} \sum_{m|n} c_n(n/m) \dim(V_1^{g^{n/m}}) - \sum_{m|n} \xi_n(m) \sum_{c|m} \frac{\phi((c, m/c))}{(c, m/c)} \dim(V_1^{\text{orb}(g^{nc/m})}) \\ = 24\delta_{1,n} - 24 \sum_{m|n} \xi_n(m) \sum_{c|m} \frac{\phi((c, m/c))}{(c, m/c)} + R(g). \end{aligned}$$

Replacing c by m/c and swapping the order of the two summations we obtain

$$\begin{aligned}
& \sum_{m|n} \xi_n(m) \sum_{c|m} \frac{\phi((c, m/c))}{(c, m/c)} \dim(V_1^{\text{orb}(g^{nc/m})}) \\
&= \sum_{m|n} \xi_n(m) \sum_{c|m} \frac{\phi((c, m/c))}{(c, m/c)} \dim(V_1^{\text{orb}(g^{n/c})}) \\
&= \sum_{c|n} \dim(V_1^{\text{orb}(g^{n/c})}) \sum_{c|m|n} \xi_n(m) \frac{\phi((c, m/c))}{(c, m/c)} \\
&= \sum_{c|n} \dim(V_1^{\text{orb}(g^{n/c})}) \delta_{c,n} = \dim(V_1^{\text{orb}(g)}).
\end{aligned}$$

Similarly,

$$\sum_{m|n} \xi_n(m) \sum_{c|m} \frac{\phi((c, m/c))}{(c, m/c)} = 1.$$

This proves

$$\dim(V_1^{\text{orb}(g)}) = 24 - 24\delta_{1,n} + \sum_{m|n} c_n(m) \dim(V_1^{g^m}) - R(g).$$

Recall that

$$\tilde{R}(g) = \sum_{\substack{i,j \in \mathbb{Z}_n \\ ij \not\equiv 0 \pmod n}} \tilde{d}_n(i, j) \dim_g(i, j)$$

with $\dim_g(i, j) := \dim(W_{1-r(i,j)}^{(i,j)})$. To compute $\tilde{R}(g^m)$ for $m | n$ we note that

$$\dim_{g^m}(i, j) = \sum_{l \in \mathbb{Z}_m} \dim_g(im, j + ln/m).$$

Hence

$$\begin{aligned}
R(g) &= \sum_{m|n} \xi_n(n/m) \tilde{R}(g^m) = \sum_{m|n} \xi_n(n/m) \sum_{\substack{i,j \in \mathbb{Z}_{n/m} \\ ij \not\equiv 0 \pmod{n/m}}} \tilde{d}_{n/m}(i, j) \dim_{g^m}(i, j) \\
&= \sum_{m|n} \xi_n(n/m) \sum_{\substack{i,j \in \mathbb{Z}_{n/m} \\ ij \not\equiv 0 \pmod{n/m}}} \tilde{d}_{n/m}(i, j) \sum_{l \in \mathbb{Z}_m} \dim_g(im, j + ln/m),
\end{aligned}$$

which may be rewritten as

$$\begin{aligned}
R(g) &= \sum_{\substack{i,j \in \mathbb{Z}_n \\ ij \not\equiv 0 \pmod n}} \left(\sum_{m|(i,n)} \xi_n(n/m) \tilde{d}_{n/m}(i/m, j) \right) \dim_g(i, j) \\
&= \sum_{\substack{i,j \in \mathbb{Z}_n \\ ij \not\equiv 0 \pmod n}} d_n(i, j) \dim_g(i, j)
\end{aligned}$$

with

$$d_n(i, j) := \sum_{m|(i,n)} \xi_n(n/m) \tilde{d}_{n/m}(i/m, j)$$

for all $i, j \in \mathbb{Z}_n$ with $ij \not\equiv 0 \pmod n$. This proves the theorem. $\square$

Again, the rest term $R(g)$ contains exactly the terms involving the dimensions of the weight spaces of the irreducible V^g -modules of weight less than 1. We shall prove in Proposition 5.5 that $R(g) \geq 0$. The theorem generalises Corollary 4.3 in [EMS20b].

We remark that if the order of g is a prime p , then the first and second dimension formulae both reduce to

$$\dim(V_1) + \dim(V_1^{\text{orb}(g)}) = 24 + (p+1)\dim(V_1^g) - R(g)$$

with rest term

$$R(g) = \tilde{R}(g) = \frac{24}{p-1} \sum_{k=1}^{p-1} \sigma(p-k) \sum_{\substack{\gamma \in D \\ q(\gamma) \equiv k/p \pmod{1}}} \dim(W_{k/p}^\gamma).$$

A similar formula was derived in [BM21] in the special case of $p = 11, 17, 19$, the primes with $g(X_0(p)) = 1$ using the theory of mock modular forms.

5.3. Dimension Bounds. In this section we show that the rest terms in Theorem 5.2 and Theorem 5.3 are non-negative. This implies that there is an upper bound for the dimension of the weight-1 subspace of the orbifold construction. Based on this observation we define the notion of an extremal orbifold.

Proposition 5.4. *In the rest term $\tilde{R}(g)$ in Theorem 5.2 the coefficients $\tilde{d}_n(\gamma)$ for all $\gamma \in D$ with $q(\gamma) \not\equiv 0 \pmod{1}$ satisfy*

$$\tilde{d}_n(\gamma) = [F_\gamma](r_\gamma) \in \frac{24}{\phi(n)} \mathbb{Z}_{>0}.$$

This implies

$$\tilde{R}(g) \geq 0.$$

Proof. Since the argument is tedious, we only sketch it. We write

$$\tilde{d}_n(\gamma) = [F_\gamma](r_\gamma) = \frac{24}{\phi(n)} \sigma(r_{\gamma,[n]}) \sum_{c|n} \tilde{f}_{n,\gamma}(c)$$

with

$$\tilde{f}_{n,\gamma}(c) := \delta_{\gamma \in D^c} \frac{1}{(c, n/c)} \sum_{k|n} \mu(n/k) (k, c)^2 \sigma \left(r_{\gamma,n} \frac{k}{(k, c)^2} \right) C_{(c, n/c)}(y_{n,\gamma}(c, k))$$

where we factored the rational number $r_\gamma = r_{\gamma,n} r_{\gamma,[n]}$ such that $r_{\gamma,n}$ contains all the prime divisors of n , i.e. $r_{\gamma,n} = \prod_{p|n} p^{\nu_p(r_\gamma)}$. Note that $r_{\gamma,[n]} \in \mathbb{Z}_{>0}$ because $nr_\gamma \in \mathbb{Z}_{>0}$. Then the functions $\tilde{f}_{n,\gamma}$ are multiplicative after normalisation. Writing $\sum_{c|n} \tilde{f}_{n,\gamma}(c)$ as a product over the primes dividing n we see that this sum is a positive integer. $\square$

For the rest term in Theorem 5.3 we show a similar statement. The results differ slightly, insofar as the coefficients in the rest term $R(g)$ are only non-negative rather than positive.

We denote by i_n for $i \in \mathbb{Z}_n$ the representative of i in $\{0, 1, \dots, n-1\}$.

Proposition 5.5. *In the rest term $R(g)$ in Theorem 5.3 the coefficients $d_n(\gamma)$ for all $\gamma = (i, j) \in D$ with $q(\gamma) \not\equiv 0 \pmod{1}$ satisfy*

$$d_n(\gamma) \in \frac{24}{\phi(n)} \mathbb{Z}_{\geq 0}.$$

This implies

$$R(g) \geq 0.$$

Moreover, $d_n(i, j) = 0$ if and only if $(i, j, n) \nmid [i_n j_n / n]$.

Proof. The proof is analogous to that of Proposition 5.4. Here we write

$$d_n(\gamma) = \frac{24}{\phi(n)} \sigma(r_{\gamma,[n]}) \sum_{c|n} f_{n,\gamma}(c)$$

with

$$\begin{aligned} f_{n,\gamma}(c) := & \delta_{\gamma \in D^c} \sum_{m|(i,n)/c} \frac{\lambda(m)}{m} \frac{\phi((m, n/m))}{(m, n/m)} \frac{\phi(n)}{\phi(n/m)} \frac{1}{(c, (n/m)/c)} \times \\ & \times \sum_{k|n/m} \mu((n/m)/k) (k, c)^2 \sigma \left(r_{\gamma,n} \frac{k}{(k, c)^2} \right) \times \\ & \times C_{(c, (n/m)/c)} \left((c, (n/m)/c) \left(q_c(\gamma) + r_\gamma \frac{(c/(k, c))^{-1}}{(k, c)} \right) \right) \end{aligned}$$

where the inverse is taken modulo $k/(k, c)$. Then $f_{n,\gamma}$ is multiplicative in c after normalisation. Writing $\sum_{c|n} f_{n,\gamma}(c)$ as a product over the prime divisors of n we can deduce the assertions. $\square$

Note that, while the coefficients $\tilde{d}_n(i, j)$ in Theorem 5.2 are symmetric in $i, j \in \mathbb{Z}_n$, the $d_n(i, j)$ in general are not. However, the condition that $d_n(i, j)$ vanishes is symmetric.

We now state two important corollaries of the above propositions.

Corollary 5.6 (First Dimension Bound). *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of finite order $n > 1$ and type 0 such that V^g satisfies the positivity condition. Then*

$$\psi(n) \dim(V_1^g) - \sum_{c|n} \frac{\phi((c, n/c))}{(c, n/c)} \dim(V_1^{\text{orb}(g^c)}) \geq 24 - 24 \sum_{c|n} \frac{\phi((c, n/c))}{(c, n/c)}.$$

The next corollary provides an upper bound for the dimension of the weight-1 Lie algebra $V_1^{\text{orb}(g)}$ purely in terms of the restriction of the automorphism g to the Lie algebra V_1 .

Corollary 5.7 (Second Dimension Bound). *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of finite order $n > 1$ and type 0 such that V^g satisfies the positivity condition. Then*

$$\dim(V_1^{\text{orb}(g)}) \leq 24 + \sum_{m|n} c_n(m) \dim(V_1^{g^m}).$$

We introduce the notion of extremal automorphisms that attain the upper bound for $\dim(V_1^{\text{orb}(g)})$.

Definition 5.8 (Extremality). *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of finite order $n > 1$. Suppose that g has type 0 and that V^g satisfies the positivity condition. Then g is called *extremal* if*

$$\dim(V_1^{\text{orb}(g)}) = 24 + \sum_{m|n} c_n(m) \dim(V_1^{g^m}),$$

i.e. if $R(g) = 0$. By convention, we also call the identity extremal.

It is in general not easy to identify extremal orbifolds besides explicitly computing $\dim(V_1^{\text{orb}(g)})$, either via the rest term $R(g)$ in the dimension formula or directly (as explained in, e.g., Section 5.6 of [Möl16]). Both approaches are in general computationally demanding. However, we can formulate some easily verifiable necessary and sufficient conditions for extremality:

Proposition 5.9. *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of order $n > 1$ and type 0 such that V^g satisfies the positivity condition. Then:*

- (1) *If the conformal weight $\rho(V(g^i)) \geq 1$ for all $i \in \mathbb{Z}_n \setminus \{0\}$, then g is extremal.*
- (2) *If g is extremal, then $\rho(V(g^i)) \geq 1$ for all $i \in \mathbb{Z}_n$ with $(i, n) = 1$.*
- (3) *If n is prime, then g is extremal if and only if $\rho(V(g^i)) \geq 1$ for all $i \in \mathbb{Z}_n \setminus \{0\}$.*

Proof. Item (1) is immediate since the condition implies that $\dim(W_{1-r_\gamma}^\gamma) = 0$ for all $\gamma \in D$ with $q(\gamma) \not\equiv 0 \pmod{1}$.

Suppose that $i \in \mathbb{Z}_n$ such that $(i, n) = 1$. Then Theorem 5.3 shows that for all $j \in \mathbb{Z}_n \setminus \{0\}$, $d_n(i, j) = \tilde{d}_n(i, j)$, which is positive by Proposition 5.4. Then, $R(g)$ can only vanish if $\dim(W_{1-r(i,j)}^{(i,j)}) = 0$ for all $j \in \mathbb{Z}_n \setminus \{0\}$, which implies that $\rho(V(g^i)) \geq 1$. This proves item (2).

Item (3) follows from items (1) and (2). $\square$

Extremal automorphism cannot exist for all orders n . This follows from the existence of certain Hecke eigenforms and the Deligne bound.

Theorem 5.10. *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of prime order p and type 0 such that V^g satisfies the positivity condition. If $g(X_0(p)) > 0$, then $R(g) \geq 24$ and hence g cannot be extremal.*

Proof. Let p be a prime such that $g(X_0(p)) > 0$. Then by Proposition 4.3 the space $\mathcal{S}_2(\Gamma_0(p))^-$ of cusp forms of weight 2 on which the Fricke involution W_p acts as -1 is non-trivial. Recall that $\mathcal{S}_2(\Gamma_0(p))^-$ has a basis consisting of Hecke eigenforms. Let $f \in \mathcal{S}_2(\Gamma_0(p))^-$ with $f(\tau) = \sum_{m=1}^{\infty} a(m)q^m$ be such an eigenform, normalised so that $a(1) = 1$. Then Theorem 4.4 shows that $a(2), \dots, a(p-1) \in \mathbb{R}$, $a(p) = 1$ and $|a(m)| \leq \sigma_0(m)\sqrt{m}$ for all $m \in \mathbb{Z}_{>0}$.

Since $S = W_p \frac{1}{\sqrt{p}} \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix}$, we obtain

$$f|_S(\tau) = \frac{1}{p} f|_{W_p}(\tau/p) = -\frac{1}{p} f(\tau/p).$$

Then we decompose $f(\tau/p)$, which has a Fourier expansion in $q^{1/p}$, as

$$f(\tau/p) = g_0(\tau) + \dots + g_{p-1}(\tau)$$

with $g_j|_T = e(j/p)g_j$ for $j \in \mathbb{Z}_p$.

Let $F(\tau) = \sum_{\gamma \in D} F_\gamma(\tau)e^\gamma$ be the lift (see Proposition 4.5) of f to a modular form for the dual Weil representation $\bar{\rho}_D$. Then Proposition 4.7 yields

$$F_\gamma(\tau) = \delta_{\gamma,0} f(\tau) - \frac{1}{p} g_{j_\gamma}(\tau)$$

where $j_\gamma \in \{0, \dots, p-1\}$ such that $j_\gamma/p = -q(\gamma) \pmod{1}$. We insert F into Proposition 5.1. Clearly, since f is a cusp form, so is F , i.e. $[F_\gamma](0)$ vanishes for all $\gamma \in D$. We compute

$$[F_0](1) = [f](1) - \frac{1}{p}[g_0](1) = [f](1) - \frac{1}{p}[f](p) = a(1) - a(p)/p = 1 - 1/p$$

and

$$[F_\gamma](r_\gamma) = -\frac{1}{p}[g_{j_\gamma}](r_\gamma) = -\frac{1}{p}[f](j_\gamma) = -\frac{1}{p}a(j_\gamma)$$

for $\gamma \in D$ with $q(\gamma) \not\equiv 0 \pmod{1}$. Recall that $r_\gamma \in (0, 1)$ such that $r_\gamma = -q(\gamma) \pmod{1}$. Hence, by Proposition 5.1,

$$p - 1 = \sum_{\substack{\gamma \in D \\ q(\gamma) \equiv 0 \pmod{1}}} a(j_\gamma) \dim(W_{1-r_\gamma}^\gamma) = \sum_{k=1}^{p-1} a(p-k) d_k$$

with

$$d_k := \sum_{\substack{\gamma \in D \\ q(\gamma) \equiv k/n \pmod{1}}} \dim(W_{k/n}^\gamma).$$

On the other hand, the rest term $R(g)$ reduces to

$$R(g) = \tilde{R}(g) = \frac{24}{p-1} \sum_{k=1}^{p-1} \sigma(p-k) d_k$$

since p is prime.

Finally, Deligne's bound $|a(m)| \leq \sigma_0(m) \sqrt{m} \leq \sigma(m)$ for $m \in \mathbb{Z}_{>0}$ implies

$$\begin{aligned} \sum_{k=1}^{p-1} \sigma(p-k) d_k &= \sum_{k=1}^{p-1} a(p-k) d_k + \sum_{k=1}^{p-2} (\sigma(p-k) - a(p-k)) d_k \\ &= p - 1 + \sum_{k=1}^{p-2} (\sigma(p-k) - a(p-k)) d_k \\ &\geq p - 1, \end{aligned}$$

proving the assertion. $\square$

We remark that the theorem does not extend to the non-prime case. For example, there are extremal orbifolds of order 14 (see Table 2).

Recall the inverse orbifold construction summarised at the end of Section 3.3. We show that it preserves extremality:

Proposition 5.11. *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of finite order $n > 1$ and type 0 such that V^g satisfies the positivity condition. Then g is extremal if and only if the inverse-orbifold automorphism $\zeta \in \text{Aut}(V^{\text{orb}(g)})$ is extremal.*

Proof. The rest term $R(g)$ in Theorem 5.3 is given by

$$R(g) = \sum_{\substack{i, j \in \mathbb{Z}_n \\ ij \not\equiv 0 \pmod{n}}} d_n(i, j) \dim(W_{1-r(i, j)}^{(i, j)}).$$

Let $\zeta \in \text{Aut}(V^{\text{orb}(g)})$ be the inverse-orbifold automorphism. By definition of ζ , $W^{(i, j)}$ is the eigenspace of ζ acting on $V^{\text{orb}(g)}(\zeta^j)$ with eigenvalue $e(i/n)$. Hence, the rest term in the corresponding dimension formula is given by

$$R(\zeta) = \sum_{\substack{i, j \in \mathbb{Z}_n \\ ij \not\equiv 0 \pmod{n}}} d_n(i, j) \dim(W_{1-r(i, j)}^{(j, i)}) = \sum_{\substack{i, j \in \mathbb{Z}_n \\ ij \not\equiv 0 \pmod{n}}} d_n(j, i) \dim(W_{1-r(i, j)}^{(i, j)}),$$

noting that $r(i, j) = r(j, i)$.

Since g is extremal if and only if $\dim(W_{1-r(i, j)}^{(i, j)}) = 0$ or $d_n(i, j) = 0$ for all $i, j \in \mathbb{Z}_n$ with $ij \not\equiv 0 \pmod{n}$, and similarly for ζ , the assertion follows from the fact that $d_n(i, j) = 0$ if and only if $d_n(j, i) = 0$ for all $i, j \in \mathbb{Z}_n$ with $ij \not\equiv 0 \pmod{n}$ (see Proposition 5.5). $\square$

Finally, as consequence of the dimension bounds, we give another proof of a result on Kneser neighbours of Niemeier lattices by Chenevier and Lannes [CL19]. For $n \in \mathbb{Z}_{>0}$, two positive-definite, even, unimodular lattices L_1 and L_2 are called (*Kneser*) n -neighbours if their intersection $K := L_1 \cap L_2$ satisfies $L_1/K \cong \mathbb{Z}_n \cong L_2/K$. Equivalently, K is a positive-definite, even lattice satisfying $K'/K \cong D$, the hyperbolic discriminant form with group structure $\mathbb{Z}_n \times \mathbb{Z}_n$ and quadratic form $q((i, j)) = ij/n \pmod{1}$, and L_1 and L_2 are unions of the cosets of K corresponding to two trivially intersecting, maximal isotropic subgroups of D .

Let L be a positive-definite, even lattice. For $d \in L \otimes_{\mathbb{Z}} \mathbb{Q}$ we define the finite-index sublattice $L^d = \{\alpha \in L \mid (\alpha, d) \in \mathbb{Z}\}$ and denote by $N_r(L)$ the number of lattice vectors $\alpha \in L$ of squared norm $(\alpha, \alpha) = r$. If L is also unimodular, then the n -neighbours of L are exactly the lattices M of the form $M = \text{span}_{\mathbb{Z}}\{L^d, d\}$ with $d \in L \otimes_{\mathbb{Z}} \mathbb{Q}$ such that $(d, d) \in 2\mathbb{Z}$ and $d + L$ has order n in $(L \otimes_{\mathbb{Z}} \mathbb{Q})/L$. Moreover, $N_2(L)$ is the number of roots of L .

Corollary 5.12. *Let L_1 and L_2 be Niemeier lattices. If $n > 1$ and L_1 and L_2 are n -neighbours, then $L_2 = \text{span}_{\mathbb{Z}}\{L_1^d, d\}$ for some $d \in L_1 \otimes_{\mathbb{Z}} \mathbb{Q}$ as above and*

$$N_2(L_2) \leq 24n + \sum_{m|n} c_n(m) N_2(L_1^{md})$$

where the $c_n(m)$ are defined by $\sum_{m|n} c_n(m)(t, m) = n/t$ for all $t \mid n$.

Proof. Let $V := V_{L_1}$ be the strongly rational, holomorphic vertex operator algebra of central charge 24 associated with the Niemeier lattice L_1 and $g := \sigma_d = e^{2\pi i d_0}$ the inner automorphism of V_{L_1} associated with d . Then g has order d and type 0 and it is not difficult to see that $V^{g^m} = V_{L_1^{md}}$ and $V^{\text{orb}(g)} \cong V_{L_2}$ (more details of this construction are given in Section 6.1). Finally, the assertion follows from Corollary 5.7 with the observation that $\dim((V_L)_1) = \text{rk}(L) + N_2(L)$ for any positive-definite, even lattice L . $\square$

Since the Leech lattice Λ has no roots, $N_2(L) = 0$ for all sublattices L of Λ . Then the above corollary simplifies to:

Corollary 5.13. *Let L be a Niemeier lattice. If $n > 1$ and L is an n -neighbour of the Leech lattice Λ , then $N_2(L) \leq 24n$.*

This is the statement of Proposition 3.4.1.1 in [CL19]. They even show that the converse of the statement holds (cf. Theorem B in [CL19]).

Using Theorem 5.10 we can replace the term $24p$ by $24(p-1)$ in both results if $n = p$ is a prime with $g(X_0(p)) > 0$.

We remark that it is not necessary to use vertex operator algebras to prove these results with the methods in this text. It suffices to consider the theta series of the lattices involved divided by the modular discriminant Δ , which are modular forms of weight 0 satisfying the assumptions of this section (and are precisely the characters of the corresponding lattice vertex operator algebras).

6. GENERALISED DEEP HOLES

In this section we introduce the notion of *generalised deep holes*, certain extremal automorphisms of holomorphic vertex operator algebras of central charge 24.

Then we show that there is a bijection

$$g \mapsto V_{\Lambda}^{\text{orb}(g)}$$

between the generalised deep holes g of the Leech lattice vertex operator algebra V_{Λ} with $\text{rk}((V_{\Lambda}^g)_1) > 0$ and the strongly rational, holomorphic vertex operator algebras V of central charge 24 with $V_1 \neq \{0\}$. This naturally generalises Conway, Parker

and Sloane's and Borcherds' correspondence between the Niemeier lattices and the deep holes of the Leech lattice Λ .

We also prove a uniform construction of the 70 non-vanishing Lie algebras on Schellekens' list by explicitly giving a generalised deep hole of V_Λ for each case. Finally, we classify all generalised deep holes of V_Λ up to algebraic conjugacy.

6.1. Deep Holes of the Leech Lattice. As a motivating example for the definition of a generalised deep hole we study the orbifold constructions associated with the deep holes of the Leech lattice Λ .

Recall that up to isomorphism there are exactly 24 positive-definite, even, unimodular lattices of rank 24 and that the Leech lattice Λ is the unique one amongst them without roots.

The weight-1 space of the Leech lattice vertex operator algebra V_Λ is the abelian Lie algebra $(V_\Lambda)_1 \cong \Lambda_{\mathbb{C}}$. Hence, the dimension formula simplifies.

Corollary 6.1. *Let g be an automorphism of V_Λ of finite order $n > 1$ and type 0 such that V_Λ^g satisfies the positivity condition. Suppose that g projects to the automorphism $\nu \in \text{O}(\Lambda)$ of cycle shape $\prod_{t|m} t^{b_t}$. Then*

$$\dim((V_\Lambda^{\text{orb}(g)})_1) = 24 + n \sum_{t|m} \frac{b_t}{t} - R(g) = 24 + 24n(1 - \rho_\nu) - R(g)$$

where ρ_ν is the vacuum anomaly introduced in Section 3.1.

Proof. The automorphism ν^k has cycle shape $\prod_{t|m} (t/(k, t))^{b_t(k, t)}$. Hence

$$\dim((V_\Lambda^{g^k})_1) = \text{rk}(\Lambda^{\nu^k}) = \sum_{t|m} b_t(k, t)$$

so that the relevant term in Theorem 5.3 becomes

$$\sum_{k|n} c_n(k) \dim((V_\Lambda^{g^k})_1) = \sum_{k|n} c_n(k) \sum_{t|m} b_t(k, t) = n \sum_{t|m} \frac{b_t}{t} = 24n(1 - \rho_\nu).$$

This proves the assertion. $\square$

The corollary shows that, given an automorphism $\nu \in \text{O}(\Lambda)$, the dimensions of extremal orbifold constructions associated with automorphisms of V_Λ projecting to ν lie in $24 + (m \sum_{t|m} b_t/t) \mathbb{Z}_{>0}$. Note that for automorphisms of the Leech lattice $(1/24) \sum_{t|m} b_t/t = 1 - \rho_\nu$ has a denominator dividing $m(12, m)$.

We now recall the bijection between the 23 classes of deep holes of the Leech lattice Λ and the 23 Niemeier lattices other than the Leech lattice described in [CPS82, CS82b].

A *hole* in a positive-definite lattice L is a point $d \in L \otimes_{\mathbb{Z}} \mathbb{R}$ in which the distance to the lattice L assumes a local maximum. If the distance is a global maximum, d is called a *deep hole*. In [CPS82] the authors show that the covering radius of the Leech lattice Λ is $\sqrt{2}$ and that there are exactly 23 orbits of deep holes in $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ under the affine automorphism group $\text{O}(\Lambda) \ltimes \Lambda$. The authors observe that these are in a natural bijection with the 23 isomorphism classes of Niemeier lattices other than the Leech lattice itself.

In [CS82b], for each deep hole, a simultaneous construction of the Leech lattice and of the corresponding Niemeier lattice is given. Let $d \in \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ be a deep hole of the Leech lattice. Let n be the order of $d + \Lambda \in (\Lambda \otimes_{\mathbb{Z}} \mathbb{R})/\Lambda$. We consider the sublattice $\Lambda^d := \{\alpha \in \Lambda \mid (\alpha, d) \in \mathbb{Z}\}$ of Λ of index n . Then $\text{span}_{\mathbb{Z}}\{\Lambda^d, d\}$ contains Λ^d as a sublattice of index n and is even, unimodular and in fact isomorphic to the Niemeier lattice N corresponding to d . Note that $n = h$ where h is the Coxeter number of (any irreducible component of) the root system of N .

The original proof of these results is based on a case-by-case analysis. Borcherds found a conceptual proof in [Bor85] by embedding all lattices into $\mathbb{H}_{25,1}$, the unique even, unimodular lattice of Lorentzian signature $(25, 1)$.

We now study the orbifold constructions of the Leech lattice vertex operator algebra V_Λ corresponding to deep holes of the Leech lattice.

Proposition 6.2. *Let $d \in \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ be a deep hole of the Leech lattice Λ corresponding to the Niemeier lattice N . Then $g = \sigma_d = e^{2\pi i d_0}$ is an automorphism of V_Λ of order equal to the Coxeter number of N and type 0. The corresponding orbifold construction $V_\Lambda^{\text{orb}(g)}$ is isomorphic to V_N .*

Proof. Let $d \in \Lambda \otimes_{\mathbb{Z}} \mathbb{Q}$ and consider the inner automorphism $g = \sigma_d = e^{2\pi i d_0} \in \text{Aut}(V_\Lambda)$. Then g is an automorphism of V_Λ of order n , the smallest positive integer such that $nd \in \Lambda' = \Lambda$. The conformal weight (see Section 3.1) of the unique irreducible g -twisted V_Λ -module $V_\Lambda(g)$ is $\rho(V_\Lambda(g)) = \min_{\alpha \in -d + \Lambda} (\alpha, \alpha)/2$, which is at most 1 because the covering radius of Λ is $\sqrt{2}$.

Assume that d is a deep hole of Λ , which is the case if and only if $\rho(V_\Lambda(g)) = 1$. Hence, g has type 0, and we may consider the orbifold construction $V_\Lambda^{\text{orb}(g)}$. The fixed-point vertex operator subalgebra V_Λ^g is the lattice vertex operator algebra V_{Λ^d} . Hence, rather than applying the general theory of cyclic orbifolds described in Section 3.3 it suffices to make use of the representation theory of lattice vertex operator algebras. The n^2 irreducible V_{Λ^d} -modules are given by $V_{\alpha + \Lambda^d}$ for $\alpha + \Lambda^d \in (\Lambda^d)' / \Lambda^d \cong \mathbb{Z}_n^2$. Note that $(\Lambda^d)' = \text{span}_{\mathbb{Z}}\{\Lambda, d\}$.

Now, Λ / Λ^d and $\text{span}_{\mathbb{Z}}\{\Lambda^d, d\} / \Lambda^d \cong N / \Lambda^d$ form a pair of trivially intersecting, maximal isotropic subgroups of $(\Lambda^d)' / \Lambda^d$. The sum of irreducible V_{Λ^d} -modules corresponding to the former gives back V_Λ while the one corresponding to the latter yields

$$V_\Lambda^{\text{orb}(g)} = \bigoplus_{j \in \mathbb{Z}_n} V_{jd + \Lambda^d} \cong V_N. \quad \square$$

The proposition shows that we can replicate the deep-hole construction of the 23 Niemeier lattices other than the Leech lattice on the level of strongly rational, holomorphic vertex operator algebras of central charge 24, simply by studying inner automorphisms.

We collect some properties of these 23 orbifold constructions, which will serve as defining properties of generalised deep holes.

Proposition 6.3. *Let d be a deep hole of Λ and $g = \sigma_d = e^{2\pi i d_0}$ the corresponding inner automorphism of V_Λ . Then g has type 0 and V_Λ^g satisfies the positivity condition. Furthermore*

- (1) g is extremal,
- (2) $\text{rk}((V_\Lambda^g)_1) = \text{rk}((V_\Lambda^{\text{orb}(g)})_1) = 24$.

Proof. It was already stated in Proposition 6.2 that g has type 0. By Corollary 6.1, the upper bound from the dimension formula for orbifold constructions of V_Λ with inner automorphisms of order n is given by

$$\dim((V_\Lambda^{\text{orb}(g)})_1) \leq 24 + 24n.$$

On the other hand, let N be the Niemeier lattice with $V_\Lambda^{\text{orb}(g)} \cong V_N$. Then

$$\dim((V_\Lambda^{\text{orb}(g)})_1) = \dim((V_N)_1) = 24 + 24h$$

because the number of roots of N is $24h$, h being the Coxeter number of N . Since $n = h$, the upper bound is attained and g is extremal.

The automorphism g acts trivially on the 24-dimensional abelian Lie algebra $(V_\Lambda)_1$ so that $\text{rk}((V_\Lambda^g)_1) = 24$. Because $(V_\Lambda^{\text{orb}(g)})_1 \cong (V_N)_1$ is a semisimple Lie algebra of rank 24, the assertion follows. $\square$

Note that $\rho(V_\Lambda(g)) = 1$ is necessary albeit in general not sufficient for g to be extremal (see Proposition 5.9). For many but not all of the 23 deep holes, $\rho(V_\Lambda(g^i)) = 1$ for all $i \in \mathbb{Z}_n \setminus \{0\}$, which also implies the extremality of g .

6.2. Definition of Generalised Deep Holes. We use the properties of the deep-hole orbifold constructions listed in Proposition 6.3 to define generalised deep holes. In order to obtain non-lattice cases on Schellekens' list as orbifold constructions associated with V_Λ , we allow $(V_\Lambda^{\text{orb}(g)})_1$ to have rank less than 24.

Definition 6.4 (Generalised Deep Hole). Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 and $g \in \text{Aut}(V)$ of finite order $n > 1$. Suppose g has type 0 and V^g satisfies the positivity condition. Then g is called a *generalised deep hole* of V if

- (1) g is extremal, i.e. $\dim(V_1^{\text{orb}(g)}) = 24 + \sum_{m|n} c_n(m) \dim(V_1^{g^m})$,
- (2) $\text{rk}(V_1^{\text{orb}(g)}) = \text{rk}(V_1^g)$.

Note that the Lie algebras V_1^g and $V_1^{\text{orb}(g)}$ are reductive [DM04b]. By Lemma 8.1 in [Kac90] the centraliser in $V_1^{\text{orb}(g)}$ of any choice of Cartan subalgebra of V_1^g is a Cartan subalgebra of $V_1^{\text{orb}(g)}$. The second condition is hence equivalent to requiring that any Cartan subalgebra of V_1^g also be a Cartan subalgebra of $V_1^{\text{orb}(g)}$. Since a finite-order automorphism of a reductive Lie algebra is inner if and only if it fixes a Cartan subalgebra pointwise (see the proof of Proposition 8.6 in [Kac90]), the second condition can also be replaced by the condition that the inverse-orbifold automorphism restricts to an inner automorphism on $V_1^{\text{orb}(g)}$.

If $V = V_\Lambda$, the vertex operator algebra associated with the Leech lattice Λ , then the rank condition is equivalent to demanding that $(V_\Lambda^g)_1$, which as subalgebra of $(V_\Lambda)_1$ is abelian, be a Cartan subalgebra of $(V_\Lambda^{\text{orb}(g)})_1$.

Again, we also call the identity a generalised deep hole.

6.3. Uniform Construction of Schellekens' List. In the following, we show that there is a bijection between the strongly rational, holomorphic vertex operator algebras of central charge 24 with non-trivial weight-1 subspace and the generalised deep holes of the Leech lattice vertex operator algebra V_Λ with non-trivial fixed-point Lie subalgebra. This is a natural generalisation of the bijection between the deep holes of the Leech lattice Λ and the 23 Niemeier lattices with non-trivial root system proved in [CPS82, CS82b, Bor85].

Then we prove a uniform construction of the 70 non-zero Lie algebra structures on Schellekens' list by explicitly giving a generalised deep hole of V_Λ for each case.

This is an improvement over the potpourri of constructions in [FLM88, DGM90, Don93, DGM96, Lam11, LS12, LS15, Miy13, EMS20a, Möl16, SS16, LS16b, LS16a, LL20]. We remark that other uniform constructions are given in [Höh17] (and [Lam20]), as well as in [HM20].

First, we combine the dimension formula in Theorem 5.3 with the averaged very strange formula in Theorem 2.4. This generalises Theorem 3.3 in [ELMS21] to non-quasirational automorphisms.

Proposition 6.5. *Let V be a strongly rational, holomorphic vertex operator algebra of central charge 24 whose weight-1 Lie algebra $V_1 = \bigoplus_{i=1}^r \mathfrak{g}_i$ is semisimple and g*

an inner automorphism of V of finite order $n > 1$ such that g is of type 0 and that V^g satisfies the positivity condition. Then

$$\dim(V_1^{\text{orb}(g)}) = 24 + \frac{12n}{\phi(n)} \sum_{i=1}^r \sum_{j \in \mathbb{Z}_n^*} h_i^\vee \left| \lambda_{s_i^{[j]}} - \frac{\rho_i}{h_i^\vee} \right|^2 - R(g).$$

Here, $R(g) \geq 0$ is as in Theorem 5.3, ρ_i is the Weyl vector and $h_i^\vee$ the dual Coxeter number of $\mathfrak{g}_i$ and $s_i^{[j]}$ the sequence characterising $g^j|_{\mathfrak{g}_i}$.

Proof. As inner automorphism, g restricts to each simple ideal of V_1 so that the relevant term in the dimension formula in Theorem 5.3 becomes

$$\begin{aligned} \sum_{m|n} c_n(m) \dim(V_1^{g^m}) &= \sum_{i=1}^r \sum_{m|n} c_n(m) \dim(\mathfrak{g}_i^{g^m}) \\ &= \frac{12n}{\phi(n)} \sum_{i=1}^r \sum_{j \in \mathbb{Z}_n^*} h_i^\vee \left| \lambda_{s_i^{[j]}} - \frac{\rho_i}{h_i^\vee} \right|^2 \end{aligned}$$

by Theorem 2.4, where n is in general a multiple of the order of $g|_{V_1}$ and $g|_{\mathfrak{g}_i}$. $\square$

Note that the assumption that V_1 is semisimple and g inner is merely made for convenience and could be easily removed.

Theorem 6.6 (Holy Correspondence). *The cyclic orbifold construction $g \mapsto V_\Lambda^{\text{orb}(g)}$ defines a bijection between the algebraic conjugacy classes of generalised deep holes $g \in \text{Aut}(V_\Lambda)$ with $\text{rk}((V_\Lambda^g)_1) > 0$ and the isomorphism classes of strongly rational, holomorphic vertex operator algebras V of central charge 24 with $V_1 \neq \{0\}$.*

Proof. Let V be a strongly rational, holomorphic vertex operator algebra V of central charge 24 with non-trivial weight-1 Lie algebra V_1 . Then V_1 is either abelian and of rank 24 or semisimple [DM04a]. In the first case V must be isomorphic to the Leech lattice vertex operator algebra V_Λ [DM04b]. Moreover, if g is a generalised deep hole such that $V_\Lambda^{\text{orb}(g)} \cong V_\Lambda$, then $24 = \text{rk}((V_\Lambda^{\text{orb}(g)})_1) = \text{rk}((V_\Lambda^g)_1) = \text{rk}(\Lambda^\nu)$ where ν is the projection of g to $\text{O}(\Lambda)$. Hence, ν is the identity and Corollary 6.1 implies that so is g . Therefore, in the following we assume that $\mathfrak{g} := V_1$ is semisimple and that g is non-trivial.

We sketch the proof of the surjectivity, which is given in [ELMS21]. We decompose $\mathfrak{g} = \mathfrak{g}_1 \oplus \dots \oplus \mathfrak{g}_r$ into simple components $\mathfrak{g}_i$. Then the vertex operator subalgebra of V generated by V_1 is isomorphic to the affine vertex operator algebra $L_{\hat{\mathfrak{g}}_1}(k_1, 0) \otimes \dots \otimes L_{\hat{\mathfrak{g}}_r}(k_r, 0)$ with levels $k_i \in \mathbb{Z}_{>0}$ [DM06]. Schellekens' lowest-order trace identity [Sch93] (see also [DM04a, EMS20a]) states that $h_i^\vee/k_i = (\dim(V_1) - 24)/24$ for all $i = 1, \dots, r$ where $h_i^\vee$ is the dual Coxeter number of $\mathfrak{g}_i$. Let ρ_i denote the Weyl vector of $\mathfrak{g}_i$. Define the inner automorphism $\sigma_u = e^{2\pi i u_0} \in \text{Aut}(V)$ with $u := \sum_{i=1}^r \rho_i/h_i^\vee$. It restricts to the automorphism $\sigma_i := \sigma_{\rho_i/h_i^\vee}$ introduced in Section 2.2 on each $\mathfrak{g}_i$. The order of σ_u is $\text{lcm}(\{l_i h_i^\vee\}_{i=1}^r)$ with lacing numbers $l_i \in \{1, 2, 3\}$. The fixed-point vertex operator subalgebra V^{σ_u} satisfies the positivity condition and the lowest-order trace identity implies that σ_u has type 0. Hence, the orbifold construction $V^{\text{orb}(\sigma_u)}$ exists. Inserting σ_u in Proposition 6.5 yields $\dim(V_1^{\text{orb}(\sigma_u)}) = 24 - R(\sigma_u) \leq 24$ since $\lambda_{s_i^{[j]}} = \frac{\rho_i}{h_i^\vee}$ for all $i = 1, \dots, r$ and $j \in \mathbb{Z}_n^*$, each $\sigma_u^j|_{\mathfrak{g}_i}$ being conjugate to σ_i by the rationality of σ_i . Again by the lowest-order trace identity, $V_1^{\text{orb}(\sigma_u)}$ cannot be semisimple (or else $\dim(V_1^{\text{orb}(\sigma_u)}) > 24$) so that it is abelian of rank 24. Hence, $V^{\text{orb}(\sigma_u)} \cong V_\Lambda$ by [DM04b]. Since $R(\sigma_u) = 0$, σ_u is extremal.

Finally, we consider the inverse-orbifold automorphism $g \in \text{Aut}(V_\Lambda)$ with respect to $\sigma_u \in \text{Aut}(V)$. The properties of σ_u imply that g is a generalised deep hole and

$V \cong V_\Lambda^{\text{orb}(g)}$. Indeed, g is extremal by Proposition 5.11 and the rank condition is satisfied because σ_u is inner. This proves the surjectivity.

We remark that we could have used the very strange formula rather than its averaged version since $\sigma_u|_{V_1}$ is quasirational (see Theorem 3.3 in [ELMS21]).

Next we prove the injectivity. Let $g \in \text{Aut}(V_\Lambda)$ be a generalised deep hole of order $n > 1$. Define $V := V_\Lambda^{\text{orb}(g)}$. Then the inverse-orbifold automorphism σ corresponding to g is of type 0, inner and extremal by Proposition 5.11. As above we decompose $\mathfrak{g} = \mathfrak{g}_1 \oplus \dots \oplus \mathfrak{g}_r$ into simple ideals. Then, Proposition 6.5 implies that

$$24 = \dim((V_\Lambda)_1) = \dim(V_1^{\text{orb}(\sigma)}) = 24 + \frac{12n}{\phi(n)} \sum_{i=1}^r \sum_{j \in \mathbb{Z}_n^*} h_i^\vee \left| \lambda_{s_i^{[j]}} - \frac{\rho_i}{h_i^\vee} \right|^2$$

where $s_i^{[j]}$ is the sequence characterising $\sigma^j|_{\mathfrak{g}_i}$. This equation can only be satisfied if $\sigma|_{V_1}$ is conjugate to $e^{2\pi i \text{ad}(u)}$ with $u = \sum_{i=1}^r \rho_i / h_i^\vee$. In fact, $\sigma|_{V_1}$ is conjugate to $e^{2\pi i \text{ad}(u)}$ under an inner automorphism of V_1 (see remark at the end of Section 2.2). Since inner automorphisms of V_1 can be extended to inner automorphisms of V , σ is conjugate in $\text{Aut}(V)$ to $\sigma_u = e^{2\pi i u_0}$. Hence, g is conjugate to the inverse-orbifold automorphism of σ_u . The same is true for any other generalised deep hole h of V_Λ with $V_\Lambda^{\text{orb}(h)} \cong V$. This proves the injectivity of the map. $\square$

As a consequence of the correspondence, it is possible to classify the strongly rational, holomorphic vertex operator algebras of central charge 24 with non-trivial weight-1 space by classifying the generalised deep holes $g \in \text{Aut}(V_\Lambda)$ satisfying $\text{rk}((V_\Lambda^g)_1) > 0$. This shall be addressed in a forthcoming publication [MS21] using geometric properties of generalised deep holes. (In the next section, we proceed in the opposite direction to obtain a classification of the generalised deep holes.)

The surjectivity of the map also follows (less conceptually) from Theorem 6.7 below together with Schellekens' classification of the possible V_1 -structures of the strongly rational, holomorphic vertex operator algebras of central charge 24 and the uniqueness of such a vertex operator algebra for non-trivial V_1 .

We now give a uniform construction of the 70 semisimple Lie algebra structures on Schellekens' list. The proof uses Schellekens' classification of the possible V_1 -structures and Kac's theory of finite-order automorphisms of simple Lie algebras [Kac90].

Theorem 6.7 (Uniform Construction). *Let $\mathfrak{g}$ be one of the 70 non-zero Lie algebras on Schellekens' list (Table 1 in [Sch93]). Then the corresponding automorphism $g \in \text{Aut}(V_\Lambda)$ listed in the appendix (Table 3 and Table 4) is a generalised deep hole and satisfies $(V_\Lambda^{\text{orb}(g)})_1 \cong \mathfrak{g}$.*

Some properties that only depend on the algebraic conjugacy class of the generalised deep hole g are listed in Table 1 and Table 2.

Proof. In Table 3 and Table 4 of the appendix (see also Table 1 and Table 2) we list representatives of 70 algebraic conjugacy classes of automorphisms $g \in \text{Aut}(V_\Lambda)$ with $\text{rk}((V_\Lambda^g)_1) > 0$. In each case, we shall identify the weight-1 Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$ of the corresponding orbifold construction as the one listed in Table 2 and prove that g is a generalised deep hole. Since each of the 70 non-zero Lie algebras on Schellekens' list appears exactly once, this will prove the assertion.

First, we describe these 70 automorphisms. A complete system of representatives of the conjugacy classes in $\text{Aut}(V_\Lambda)$ is given in Proposition 3.2. It is a special property of the Leech lattice Λ that the algebraic conjugacy classes in $\text{O}(\Lambda) \cong \text{Co}_0$ are uniquely determined by their cycle shape. The 70 automorphisms project to

TABLE 1. Summary of the generalised deep holes of the Leech lattice vertex operator algebra V_Λ and the corresponding orbifold constructions.

[Höh17]	Cycle shape	n	#	Rk.	Dim.
A	1^{24}	$1, 2, \dots, 25, 30, 46$	24	24	$24 + 24n - 24\delta_{n,1}$
B	$1^8 2^8$	$2, 4, \dots, 18, 22, 30$	17	16	$24 + 12n$
C	$1^6 3^6$	$3, 6, 9, 12, 18$	6	12	$24 + 8n$
D	2^{12}	$2, 6, 10, \dots, 22, 46$	9	12	$24 + 6n$
E	$1^4 2^2 4^4$	$4, 8, 12, 16$	5	10	$24 + 6n$
F	$1^4 5^4$	$5, 10$	2	8	$24 + (24/5)n$
G	$1^2 2^2 3^2 6^2$	$6, 12$	2	8	$24 + 4n$
H	$1^3 7^3$	7	1	6	$24 + (24/7)n$
I	$1^2 2^1 4^1 8^2$	8	1	6	$24 + 3n$
J	$2^3 6^3$	$6, 18$	2	6	$24 + 2n$
K	$2^2 10^2$	10	1	4	$24 + (6/5)n$
L	$1^{-24} 2^{24}, 1^{-12} 3^{12}, \dots$	$2, 3, \dots, 70, 78$	38	0	0

11 different cycle shapes in $O(\Lambda)$ (listed in Table 1) with non-trivial fixed-point sublattices. These are exactly the same conjugacy classes that appear in [Höh17], Table 4. Once a conjugacy class in $O(\Lambda)$ is specified, we choose suitable elements $h \in \pi_\nu(\Lambda_{\mathbb{C}})/\pi_\nu(\Lambda)$ (listed in Table 4) and consider the automorphism $g = \phi_\eta(\nu)\sigma_h$ where $\phi_\eta(\nu)$ is a standard lift of $\nu \in O(\Lambda)$. Up to conjugacy g does not depend on the choice of the standard lift of ν .

We explain the entries of Table 2. Each block corresponds to a cycle shape of ν and each row to a choice of h , determining the automorphism g of order n (up to conjugacy). We list the (yet to be proved) Lie algebra structure of $(V_\Lambda^{\text{orb}(g)})_1$ together with its dimension and rank, of which the latter only depends on ν . The first column corresponds to the labelling in Table 1 in [Sch93] and the second one to the labelling in Table 4 of [Höh17]. Then we list the conformal weights $\rho(V_\Lambda(g^c))$ of the unique irreducible g^c -twisted V_Λ -modules for $c \mid n$ in ascending order. (Note that for the 70 chosen automorphisms $\rho(V_\Lambda(g^i))$ only depends on (i, n) .) The last column will be explained below.

And now to the proof:

(1) First, we note that in all cases, $\rho(V_\Lambda(g)) = 1$ so that g has type 0, which implies that the corresponding cyclic orbifold construction $V_\Lambda^{\text{orb}(g)}$ exists and is a strongly rational, holomorphic vertex operator algebra of central charge 24. Moreover $\text{rk}((V_\Lambda^{\text{orb}(g)})_1) \geq \text{rk}((V_\Lambda^g)_1) = \text{rk}(\Lambda^\nu) > 0$ since ν has non-trivial fixed-point sublattice Λ^ν . Hence, the Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$ is one of the 70 non-zero (semisimple or abelian) Lie algebras on Schellekens' list.

(2) The 24 automorphisms with $\nu = \text{id}$ are exactly the inner automorphisms $g = \sigma_h$ corresponding to the 23 classes of deep holes h of the Leech lattice Λ and the identity. We proved in Proposition 6.3 that they are generalised deep holes and in Proposition 6.2 that the corresponding orbifold constructions realise the 24 Niemeier lattice vertex operator algebras, with weight-1 Lie algebras of rank 24.

(3) It remains to study the 46 classes of automorphisms $g = \phi_\eta(\nu)\sigma_h$ with non-trivial ν , i.e. with $\text{rk}((V_\Lambda^g)_1) = \text{rk}(\Lambda^\nu) < 24$. We begin by proving that $\text{rk}((V_\Lambda^{\text{orb}(g)})_1) = \text{rk}((V_\Lambda^g)_1)$ or equivalently that the abelian Lie algebra $(V_\Lambda^g)_1$ is

TABLE 2. The 70 generalised deep holes of V_Λ whose corresponding orbifold constructions realise all non-zero Lie algebras on Schellekens' list (continued on next page).

No.	No.	$(V_\Lambda^{\text{orb}(g)})_1$	Dim.	n	$\rho(V_\Lambda(g^c))$	$(V_\Lambda(g))_1$
Rk. 24, cycle shape 1^{24}						
70	A1	$D_{24,1}$	1128	46	$1, 22/23, 1, 0$	$\tilde{D}_{24}$
69	A2	$D_{16,1}E_{8,1}$	744	30	$1, 14/15, 1, 1, 1, 1, 0$	$\tilde{D}_{16}\tilde{E}_8$
68	A3	$E_{8,1}^3$	744	30	$1, 14/15, 9/10, 5/6, 1, 1, 1, 0$	$\tilde{E}_8^3$
67	A4	$A_{24,1}$	624	25	$1, 1, 0$	$\tilde{A}_{24}$
66	A5	$D_{12,1}^2$	552	22	$1, 10/11, 1, 0$	$\tilde{D}_{12}^2$
65	A6	$A_{17,1}E_{7,1}$	456	18	$1, 1, 1, 1, 1, 0$	$\tilde{A}_{17}\tilde{E}_7$
64	A7	$D_{10,1}E_{7,1}^2$	456	18	$1, 8/9, 1, 1, 1, 0$	$\tilde{D}_{10}\tilde{E}_7^2$
63	A8	$A_{15,1}D_{9,1}$	408	16	$1, 1, 1, 1, 0$	$\tilde{A}_{15}\tilde{D}_9$
61	A9	$D_{8,1}^3$	360	14	$1, 6/7, 1, 0$	$\tilde{D}_8^3$
60	A10	$A_{12,1}^2$	336	13	$1, 0$	$\tilde{A}_{12}^2$
59	A11	$A_{11,1}D_{7,1}E_{6,1}$	312	12	$1, 1, 1, 1, 1, 0$	$\tilde{A}_{11}\tilde{D}_7\tilde{E}_6$
58	A12	$E_{6,1}^4$	312	12	$1, 1, 3/4, 1, 1, 0$	$\tilde{E}_6^4$
55	A13	$A_{9,1}^2D_{6,1}$	264	10	$1, 1, 1, 0$	$\tilde{A}_9^2\tilde{D}_6$
54	A14	$D_{6,1}^4$	264	10	$1, 4/5, 1, 0$	$\tilde{D}_6^4$
51	A15	$A_{8,1}^3$	240	9	$1, 1, 0$	$\tilde{A}_8^3$
49	A16	$A_{7,1}^2D_{5,1}^2$	216	8	$1, 1, 1, 0$	$\tilde{A}_7^2\tilde{D}_5^2$
46	A17	$A_{6,1}^4$	192	7	$1, 0$	$\tilde{A}_6^4$
43	A18	$A_{5,1}^4D_{4,1}$	168	6	$1, 1, 1, 0$	$\tilde{A}_5^4\tilde{D}_4$
42	A19	$D_{4,1}^6$	168	6	$1, 2/3, 1, 0$	$\tilde{D}_4^6$
37	A20	$A_{4,1}^6$	144	5	$1, 0$	$\tilde{A}_4^6$
30	A21	$A_{3,1}^8$	120	4	$1, 1, 0$	$\tilde{A}_3^8$
24	A22	$A_{2,1}^{12}$	96	3	$1, 0$	$\tilde{A}_2^{12}$
15	A23	$A_{1,1}^{24}$	72	2	$1, 0$	$\tilde{A}_1^{24}$
1	A24	$\mathbb{C}^{24}$	24	1	0	$\emptyset$
Rk. 16, cycle shape $1^8 2^8$						
62	B1	$B_{8,1}E_{8,2}$	384	30	$1, 14/15, 1, 1, 1, 1, 1, 0$	$A_1\tilde{E}_8$
56	B2	$C_{10,1}B_{6,1}$	288	22	$1, 10/11, 1, 0$	A_1A_9
52	B3	$C_{8,1}F_{4,1}^2$	240	18	$1, 8/9, 1, 1, 1, 0$	$A_2^2A_7$
53	B4	$B_{5,1}E_{7,2}F_{4,1}$	240	18	$1, 8/9, 1, 1, 1, 0$	$A_1A_2\tilde{E}_7$
50	B5	$A_{7,1}D_{9,2}$	216	16	$1, 1, 1, 1, 0$	$\tilde{D}_9$
47	B6	$B_{4,1}^2D_{8,2}$	192	14	$1, 6/7, 1, 0$	$A_1^2\tilde{D}_8$
48	B7	$B_{4,1}C_{6,1}^2$	192	14	$1, 6/7, 1, 0$	$A_1A_5^2$
44	B8	$A_{5,1}C_{5,1}E_{6,2}$	168	12	$1, 1, 1, 1, 1, 0$	$A_4\tilde{E}_6$
40	B9	$A_{4,1}A_{9,2}B_{3,1}$	144	10	$1, 1, 1, 0$	$A_1\tilde{A}_9$
39	B10	$B_{3,1}^2C_{4,1}D_{6,2}$	144	10	$1, 4/5, 1, 0$	$A_1^2A_3\tilde{D}_6$
38	B11	$C_{4,1}^4$	144	10	$1, 4/5, 1, 0$	A_3^4
33	B12	$A_{3,1}A_{7,2}C_{3,1}^2$	120	8	$1, 1, 1, 0$	$A_2^2\tilde{A}_7$
31	B13	$A_{3,1}^2D_{5,2}^2$	120	8	$1, 1, 1, 0$	$\tilde{D}_5^2$
26	B14	$A_{2,1}^2A_{5,2}^2B_{2,1}$	96	6	$1, 1, 1, 0$	$A_1\tilde{A}_5^2$
25	B15	$B_{2,1}^4D_{4,2}^2$	96	6	$1, 2/3, 1, 0$	$A_1^4\tilde{D}_4^2$
16	B16	$A_{1,1}^4A_{3,2}^4$	72	4	$1, 1, 0$	$\tilde{A}_3^4$
5	B17	$A_{1,2}^{16}$	48	2	$1, 0$	$\tilde{A}_1^{16}$

TABLE 2. (continued)

No.	No.	$(V_{\Lambda}^{\text{orb}(g)})_1$	Dim.	n	$\rho(V_{\Lambda}(g^c))$	$(V_{\Lambda}(g))_1$
Rk. 12, cycle shape $1^6 3^6$						
45	C1	$A_{5,1}E_{7,1}$	168	18	1, 1, 1, 1, 1, 0	$\tilde{E}_7$
34	C2	$A_{3,1}D_{7,3}G_{2,1}$	120	12	1, 1, 1, 1, 1, 0	$A_1\tilde{D}_7$
32	C3	$E_{6,3}G_{2,1}^3$	120	12	1, 1, 3/4, 1, 1, 0	$A_1^3\tilde{E}_6$
27	C4	$A_{2,1}^2A_{8,3}$	96	9	1, 1, 0	$\tilde{A}_8$
17	C5	$A_{1,1}^3A_{5,3}D_{4,3}$	72	6	1, 1, 1, 0	$\tilde{A}_5\tilde{D}_4$
6	C6	$A_{2,3}^6$	48	3	1, 0	$\tilde{A}_2^6$
Rk. 12, cycle shape 2^{12} (order doubling)						
57	D1a	$B_{12,2}$	300	46	1, 22/23, 1, 0	A_1
41	D1b	$B_{6,2}^2$	156	22	1, 10/11, 1, 0	A_1^2
29	D1c	$B_{4,2}^3$	108	14	1, 6/7, 1, 0	A_1^3
23	D1d	$B_{3,2}^4$	84	10	1, 4/5, 1, 0	A_1^4
12	D1e	$B_{2,2}^6$	60	6	1, 2/3, 1, 0	A_1^6
2	D1f	$A_{1,4}^{12}$	36	2	1, 0	$\tilde{A}_1^{12}$
36	D2a	$A_{8,2}F_{4,2}$	132	18	1, 1, 1, 1, 1, 0	A_2
22	D2b	$A_{4,2}^2C_{4,2}$	84	10	1, 1, 1, 0	A_3
13	D2c	$A_{2,2}^4D_{4,4}$	60	6	1, 1, 1, 0	$\tilde{D}_4$
Rk. 10, cycle shape $1^4 2^2 4^4$						
35	E1	$A_{3,1}C_{7,2}$	120	16	1, 1, 1, 1, 1, 0	A_6
28	E2	$A_{2,1}B_{2,1}E_{6,4}$	96	12	1, 1, 1, 1, 1, 0	$\tilde{E}_6$
18	E3	$A_{1,1}^3A_{7,4}$	72	8	1, 1, 1, 0	$\tilde{A}_7$
19	E4	$A_{1,1}^2C_{3,2}D_{5,4}$	72	8	1, 1, 1, 0	$A_2\tilde{D}_5$
7	E5	$A_{1,2}A_{3,4}^3$	48	4	1, 1, 0	$\tilde{A}_3^3$
Rk. 8, cycle shape $1^4 5^4$						
20	F1	$A_{1,1}^2D_{6,5}$	72	10	1, 1, 1, 0	$\tilde{D}_6$
9	F2	$A_{4,5}^2$	48	5	1, 0	$\tilde{A}_4^2$
Rk. 8, cycle shape $1^2 2^2 3^2 6^2$						
21	G1	$A_{1,1}C_{5,3}G_{2,2}$	72	12	1, 1, 1, 1, 1, 0	A_1A_4
8	G2	$A_{1,2}A_{5,6}B_{2,3}$	48	6	1, 1, 1, 0	$A_1\tilde{A}_5$
Rk. 6, cycle shape $1^3 7^3$						
11	H1	$A_{6,7}$	48	7	1, 0	$\tilde{A}_6$
Rk. 6, cycle shape $1^2 2^1 4^1 8^2$						
10	I1	$A_{1,2}D_{5,8}$	48	8	1, 1, 1, 0	$\tilde{D}_5$
Rk. 6, cycle shape $2^3 6^3$ (order doubling)						
14	J1a	$A_{2,2}F_{4,6}$	60	18	1, 1, 1, 1, 1, 0	A_2
3	J1b	$A_{2,6}D_{4,12}$	36	6	1, 1, 1, 0	$\tilde{D}_4$
Rk. 4, cycle shape $2^2 10^2$ (order doubling)						
4	K1	$C_{4,10}$	36	10	1, 1, 1, 0	A_3

a Cartan subalgebra of $(V_\Lambda^{\text{orb}(g)})_1$, which is one of the two properties of a generalised deep hole and additionally determines the rank of the Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$. (In particular, we may then assume that $(V_\Lambda^{\text{orb}(g)})_1$ is semisimple as the only abelian case on Schellekens' list has rank 24.)

Indeed, by the inverse orbifold construction, $(V_\Lambda^{\text{orb}(g)})_1$ is a $\mathbb{Z}_n$ -graded Lie algebra with 0-component $(V_\Lambda^g)_1$. If the centraliser $C_{W_1^{(i,0)}}((V_\Lambda^g)_1) \subseteq C_{V_\Lambda(g^i)_1}((V_\Lambda^g)_1) = \{0\}$ for all $i \in \mathbb{Z}_n \setminus \{0\}$, then Lemma 8.1 in [Kac90] implies that $(V_\Lambda^g)_1$ is a Cartan subalgebra of $(V_\Lambda^{\text{orb}(g)})_1$. Now, by the definition of the twisted modules for lattice vertex operator algebras (see Section 3.1), the centraliser condition for $i = 1$ follows from $h \notin \pi_\nu(\Lambda)$ and similarly for the other powers of g (essentially, the condition becomes $ih \notin \pi_\nu(\Lambda)$ but an extra term appears if $\phi_\eta(\nu)^i$ is not a standard lift of ν^i). It is easy to check that all automorphisms $g = \phi_\eta(\nu)\sigma_h$ satisfy this (in fact, they were chosen this way), proving that $\text{rk}((V_\Lambda^{\text{orb}(g)})_1) = \text{rk}((V_\Lambda^g)_1)$.

(4) In 31 of the remaining 46 cases, we observe that $\rho(V_\Lambda(g^i)) \geq 1$ for all $i \in \mathbb{Z}_n \setminus \{0\}$, implying that g is extremal by Proposition 5.9. This shows that g is a generalised deep hole. Moreover, the dimension formula in Theorem 5.3 then yields the dimension of $(V_\Lambda^{\text{orb}(g)})_1$ as the one listed in Table 2.

For 16 of these generalised deep holes g , the rank and dimension of the Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$ uniquely determine it on Schellekens' list. These are the cases 2, 3, 4, 5, 6, 7, 14, 16, 17, 27, 28, 35, 36, 44, 45 and 50.

(5) For the other 15 generalised deep holes g , namely those we claim to correspond to cases 8, 9, 10, 11, 13, 18, 19, 20, 21, 22, 34, 26, 31, 33 and 40 on Schellekens' list, the Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$ is not uniquely determined by its rank and dimension. Here, we need an additional argument.

The inverse orbifold automorphism ζ on $V_\Lambda^{\text{orb}(g)}$ restricts to an inner automorphism of order dividing n of the semisimple Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$. The theory of finite-order automorphisms of simple Lie algebras was developed by Kac (see Section 2.1 and [Kac90]). The said automorphism is regular, i.e. the fixed-point Lie subalgebra $(V_\Lambda^g)_1$ is abelian, and in fact a Cartan subalgebra of $(V_\Lambda^{\text{orb}(g)})_1$.

Then, for a fixed $i \in \mathbb{Z}_n$ with $(i, n) = 1$ we consider the adjoint action of $(V_\Lambda^g)_1$ on $(V_\Lambda(g^i))_1 \subseteq (V_\Lambda^{\text{orb}(g)})_1$. This defines a subset of the root system of $(V_\Lambda^{\text{orb}(g)})_1$. Crucially, $(V_\Lambda(g^i))_1 = W_1^{(i,0)}$ is the eigenspace of the automorphism $\zeta^{i^{-1}}$ (where i^{-1} denotes the inverse of i modulo n) restricted to $(V_\Lambda^{\text{orb}(g)})_1$ with eigenvalue $e(1/n)$. By Proposition 8.1 c) in [Kac90] the roots of $(V_\Lambda(g^i))_1$ are the *simple* roots of a subdiagram of the (untwisted) affinisation of the Dynkin diagram associated with the semisimple Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$. (More precisely, it is the subdiagram consisting of those nodes in the affine Dynkin diagram for which the corresponding entry of the sequence $s = (s_0, \dots, s_l)$ characterising $\zeta^{i^{-1}}$ equals 1.)

For a given automorphism $g = \phi_\eta(\nu)\sigma_h$ this Dynkin diagram associated with the action of $(V_\Lambda^g)_1$ on $(V_\Lambda(g^i))_1$ can be easily determined using the definition of the twisted modules for lattice vertex operator algebras. We list the result in the last column of Table 2 and note that it is the same for all $i \in \mathbb{Z}_n$ with $(i, n) = 1$.

Then, for each of the 15 generalised deep holes we check if this Dynkin diagram of $(V_\Lambda(g))_1$ is compatible with the remaining potential Lie algebras for $(V_\Lambda^{\text{orb}(g)})_1$. In each case, this narrows down the list of possible Lie algebras on Schellekens' list to just one. For example, the generalised deep hole we list for case 40 on Schellekens' list has an orbifold Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$ of rank 16 and dimension 144. This leaves the cases 38, 39 and 40 on Schellekens' list. However, of the affine diagrams $\tilde{C}_4^4$,

$\tilde{B}_3^2 \tilde{C}_4 \tilde{D}_6$ and $\tilde{A}_4 \tilde{A}_9 \tilde{B}_3$ only the latter contains $A_1 \tilde{A}_9$ as a subdiagram, leaving only case 40, i.e. $A_4 A_9 B_3$, as weight-1 Lie algebra of $V_\Lambda^{\text{orb}(g)}$.

(6) It remains to study the $15 = 46 - 31$ automorphisms g of which we do not know the dimension of $(V_\Lambda^{\text{orb}(g)})_1$ a priori by using the dimension formula. (By Corollary 5.7 we do however know that the dimensions listed in Table 2 must be upper bounds.) For ten of these automorphisms, namely those we claim to correspond to cases 25, 32, 38, 39, 47, 48, 52, 53, 56 and 62 on Schellekens' list, the argument we just described involving the Dynkin diagram of $(V_\Lambda(g))_1$ suffices to reduce the list of possible Lie algebra structures of $(V_\Lambda^{\text{orb}(g)})_1$ to just one case. In particular, this determines the dimension of $(V_\Lambda^{\text{orb}(g)})_1$ and gives an a posteriori proof that these automorphisms are generalised deep holes.

For example, the automorphism we list for case 62 has an orbifold Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$ of rank 16 and dimension bounded from above by 384, which is compatible with all 17 rank-16 Lie algebras on Schellekens' list. However, the Dynkin diagram associated with the action of $(V_\Lambda^g)_1$ on $(V_\Lambda(g))_1$ is $A_1 \tilde{E}_8$. Noting that the diagram $\tilde{E}_8$ is not a subdiagram of any other irreducible affine Dynkin diagram, an inspection of the Lie algebras of rank 16 in Table 2 reveals that only case 62 on Schellekens' list is compatible. This Lie algebra $B_8 E_8$ has dimension 384 so that g is extremal and a generalised deep hole.

(7) We are left with the five automorphisms g of order 6, 10, 14, 22 and 46 that we postulate to yield cases 12, 23, 29, 41 and 57, i.e. the Lie algebras B_2^6 , B_3^4 , B_4^3 , B_6^2 and B_{12} , respectively, in the orbifold construction. These Lie algebras of rank 12 are too similar to be distinguished by the Dynkin diagram of $(V_\Lambda(g))_1$. We can however show that none of the rank-12 Lie algebras on Schellekens' list, except for cases 2, 12, 23, 29, 41 and 57, are compatible with the Dynkin diagram of $(V_\Lambda(g))_1$ in each case. This means that the Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$ is uniquely determined once we know its dimension.

This dimension can be determined using the formulae in Section 3.4. It is a standard exercise to compute the q -expansions of the theta series of the lattices $M_{k,d}$ appearing in the functions D_k (for instance, this takes less than a second using the function `ThetaSeriesModularForm` in `Magma` [BCP97]) to sufficiently high order so as to identify them as modular forms. Let $p = n/2$, half the order of g . Then we find

$$\begin{aligned} D_1(\tau) &= (p-1)T_{2p+p}(\tau), \\ D_2(\tau) &= (p-1)(T_{p+}(\tau) + 24), \\ D_p(\tau) &= T_{2-}(\tau), \\ D_{2p}(\tau) &= T_1(\tau) + 24 = j(\tau) - 720 \end{aligned}$$

in terms of certain McKay-Thompson series (see [CN79]). We can also write these functions as eta products. For example,

$$D_1(\tau) = (p-1) \left(\frac{\eta(\tau)\eta(p\tau)}{\eta(2\tau)\eta(2p\tau)} \right)^{24/(p+1)} + \frac{24(p-1)}{p+1}.$$

It is now easy to compute the expansions of the D_k in the different cusps (for instance by using the well-known modular transformation properties of the eta function). We obtain

$$\text{ch}_{V_\Lambda^{\text{orb}(g)}}(\tau) = j(\tau) - 720 + 12p = q^{-1} + 24 + 12p + O(q)$$

for the character of $V_\Lambda^{\text{orb}(g)}$. Hence, $\dim((V_\Lambda^{\text{orb}(g)})_1) = 24 + 12p = 24 + 6n$, which together with the previous remarks uniquely determines the Lie algebra $(V_\Lambda^{\text{orb}(g)})_1$

for the five automorphisms g in question as cases 12, 23, 29, 41 and 57, respectively, on Schellekens' list and in particular shows that the automorphisms are extremal and hence generalised deep holes. This completes the proof. $\square$

6.4. Classification of Generalised Deep Holes. Finally, we give a complete classification of the generalised deep holes of the Leech lattice vertex operator algebra V_Λ .

Theorem 6.6 and the classification of strongly rational, holomorphic vertex operator algebras of central charge 24 (see [DM04a, LS19, LS15, LL20, EMS20b, LS20, KLL18], a uniform proof is given in [HM20]) yield a classification of the generalised deep holes with $\text{rk}((V_\Lambda^g)_1) > 0$. It remains to study those with $\text{rk}((V_\Lambda^g)_1) = 0$.

Recall that the Moonshine module $V^\natural$ is a strongly rational, holomorphic vertex operator algebra V of central charge 24 with $V_1 = \{0\}$ and conjecturally the only such vertex operator algebra up to isomorphism. There are 38 orbifold constructions of the Moonshine module $V^\natural$ from the Leech lattice vertex operator algebra V_Λ [Car18], including the original construction in [FLM88] of order 2.

In [Car18], based on ideas in [Tui92, Tui95], the author proves that for all fixed-point free automorphisms $\nu \in O(\Lambda)$ with vacuum anomaly $\rho_\nu > 1$ (there are exactly 40 such conjugacy classes that give type 0, 38 up to algebraic conjugacy) the corresponding standard lift $g = \phi_\eta(\nu) \in \text{Aut}(V_\Lambda)$ (if a lattice automorphism is fixed-point free, all its lifts are standard lifts and conjugate) defines an orbifold construction that yields the Moonshine module $V^\natural$. (By generalising the orbifold construction to non-zero type, a total of 53 conjugacy classes, 51 up to algebraic conjugacy, are considered; hence the title of the paper.) No order doubling occurs. We list the 38 algebraic conjugacy classes in Table 5.

Proposition 6.8. *Let $g \in \text{Aut}(V_\Lambda)$ be one of the 38 automorphisms in [Car18] such that $V_\Lambda^{\text{orb}(g)} \cong V^\natural$. Then g is a generalised deep hole.*

Proof. All fixed-point free automorphisms $\nu \in O(\Lambda)$ with $\rho_\nu > 1$ satisfy (see Section 3.1)

$$\rho(V_\Lambda(g)) = \rho_\nu = 1 + 1/n$$

where n is the order of ν and of $g = \phi_\eta(\nu)$. Then the dimension bound in the special case of the Leech lattice vertex operator algebra V_Λ (see Corollary 6.1) yields

$$\dim((V_\Lambda^{\text{orb}(g)})_1) \leq 24 + 24n(1 - \rho_\nu) = 0,$$

which implies that $\dim((V_\Lambda^{\text{orb}(g)})_1) = 0$ and that g is extremal. Since $(V_\Lambda^g)_1$ is a subspace of $(V_\Lambda^{\text{orb}(g)})_1$, $\text{rk}((V_\Lambda^g)_1) = \text{rk}((V_\Lambda^{\text{orb}(g)})_1) = 0$. This shows that each $g = \phi_\eta(\nu)$ is a generalised deep hole with $\dim((V_\Lambda^{\text{orb}(g)})_1) = 0$ suggesting that $V_\Lambda^{\text{orb}(g)} \cong V^\natural$, which would immediately follow if it were known that $V^\natural$ is the unique strongly rational, holomorphic vertex operator algebra V of central charge 24 with $V_1 = \{0\}$. The main work of [Car18] is to prove that indeed $V_\Lambda^{\text{orb}(g)} \cong V^\natural$. $\square$

We shall see that these 38 automorphisms of V_Λ are in fact all the generalised deep holes of V_Λ with $\text{rk}(V_\Lambda^g) = 0$ up to algebraic conjugacy.

Note that one can show in a case-by-case study that the orbifold constructions associated with the exactly 42 conjugacy classes, 39 up to algebraic conjugacy, of fixed-point free automorphisms $\nu \in O(\Lambda)$ with $\rho_\nu \leq 1$ all yield the Leech lattice vertex operator algebra V_Λ again (see also Remark 7.5 in [Car18]).

Now we can prove:

Theorem 6.9 (Classification of Generalised Deep Holes). *There are exactly 108 algebraic conjugacy classes of generalised deep holes g of the Leech lattice vertex operator algebra V_Λ (see Table 1), namely*

- (1) 70 classes with $\text{rk}((V_\Lambda^g)_1) > 0$, whose corresponding orbifold constructions yield the 70 strongly rational, holomorphic vertex operator algebras V of central charge 24 with $V_1 \neq \{0\}$ (listed in Table 3 and Table 4 and summarised in Table 2), and
- (2) 38 classes with $\text{rk}((V_\Lambda^g)_1) = 0$, whose corresponding orbifold constructions all yield the Moonshine module $V^\natural$ (listed in Table 5).

Proof. By Theorem 6.7 there are at least 70 algebraic conjugacy classes of generalised deep holes g with $\text{rk}((V_\Lambda^g)_1) > 0$. Schellekens' classification and the uniqueness of a strongly rational, holomorphic vertex operator algebra of central charge 24 with a given non-zero weight-1 space together with Theorem 6.6 imply that there are exactly 70 classes.

We study the generalised deep holes $g \in \text{Aut}(V_\Lambda)$ with $\text{rk}((V_\Lambda^g)_1) = 0$. As explained above, the 38 automorphisms whose corresponding orbifold constructions yield the Moonshine module $V^\natural$ are generalised deep holes. Clearly, they must satisfy $\text{rk}((V_\Lambda^g)_1) = 0$. In fact, these are all the generalised deep holes of V_Λ with $\text{rk}((V_\Lambda^g)_1) = 0$.

Indeed, suppose that g is an automorphism in $\text{Aut}(V_\Lambda)$ with $\text{rk}((V_\Lambda^g)_1) = 0$, i.e. g projects to a fixed-point free automorphism $\nu \in \text{O}(\Lambda)$. By Proposition 3.2 the (algebraic) conjugacy classes of such automorphisms are in bijection with the (algebraic) conjugacy classes of fixed-point free automorphisms $\nu \in \text{O}(\Lambda)$, of which there are exactly 95 (90). Now suppose that g is a generalised deep hole. Then $\text{rk}((V_\Lambda^{\text{orb}(g)})_1) = 0$ and $\rho(V_\Lambda(g)) \geq 1$. We even have $\rho(V_\Lambda(g)) > 1$ because $(V_\Lambda(g))_1 \subseteq (V_\Lambda^{\text{orb}(g)})_1$. As the automorphism is fixed-point free on Λ , the conformal weight is simply $\rho(V_\Lambda(g)) = \rho_\nu$, which only depends on the cycle shape of ν . There are exactly 53 (51) automorphisms in $\text{O}(\Lambda)$ which are fixed-point free and satisfy $\rho_\nu > 1$ up to (algebraic) conjugacy and of those 40 (38) have type 0. These are exactly the 38 automorphisms listed in Table 5. $\square$

Finally, we record the following observation (see also Table 1):

Theorem 6.10. *Under the natural projection $\text{Aut}(V_\Lambda) \rightarrow \text{O}(\Lambda)$ the 70 algebraic conjugacy classes of generalised deep holes g with $\text{rk}((V_\Lambda^g)_1) > 0$ map to the 11 algebraic conjugacy classes in $\text{O}(\Lambda)$ with cycle shapes 1^{24} , $1^8 2^8$, $1^6 3^6$, 2^{12} , $1^4 2^4 4^4$, $1^4 5^4$, $1^2 2^2 3^2 6^2$, $1^3 7^3$, $1^2 2^1 4^1 8^2$, $2^3 6^3$ and $2^2 10^2$.*

This shows that we recover the decomposition of the genus of the Moonshine module described by Höhn in [Höh17]. There is probably an automorphic proof of this result (cf. [Sch17]).

APPENDIX. LIST OF GENERALISED DEEP HOLES

We give a complete list of the generalised deep holes of V_Λ .

First, we describe the 70 generalised deep holes $g = \phi_\eta(\nu)\sigma_h \in \text{Aut}(V_\Lambda)$ from Theorem 6.7 that give cyclic orbifold constructions of the 70 strongly rational, holomorphic vertex operator algebras of central charge 24 with $V_1 \neq \{0\}$. Recall that many of their properties are listed in Table 2.

We shall explain our notation. Let $\mathcal{G}_{24} \subseteq \mathbb{F}_2^{24}$ be the *extended binary Golay code*, a doubly-even, self-dual binary code in dimension 24. Let $\Omega := \{1, 2, \dots, 24\}$ denote the coordinates of $\mathbb{F}_2^{24}$ (and of $\mathbb{R}^{24}$). The automorphism group of $\mathcal{G}_{24}$, i.e. the group of permutations of Ω mapping $\mathcal{G}_{24}$ to itself, is the Mathieu group $M_{24} \subseteq \mathcal{S}_\Omega$. Following Curtis' *Miracle Octad Generator* we shall arrange vectors in $\mathbb{F}_2^{24}$ (and in $\mathbb{R}^{24}$) in (4×6) -arrays. For definiteness, let $\mathcal{G}_{24}$ be the binary code generated by the standard basis (see, e.g., (5.35) in [Gri98]).

Let $\mathbb{R}^{24}$ be equipped with the standard scalar product $(\cdot, \cdot)$ so that $(e_i, e_j) = \delta_{i,j}$ for all $i, j \in \Omega$. The *Leech lattice* $\Lambda \subseteq \mathbb{R}^{24}$ is a positive-definite, even, unimodular lattice of rank 24. It can be constructed from $\mathcal{G}_{24}$ as the $\mathbb{Z}$ -span of

$$\frac{1}{\sqrt{2}}x, x \in \mathcal{G}_{24}, \quad \sqrt{2}(e_i \pm e_j), i, j \in \Omega, \quad \frac{1}{\sqrt{8}}(-3, 1, \dots, 1)$$

where we consider the entries of $x \in \mathbb{F}_2^{24}$ as $0, 1 \in \mathbb{R}$. An explicit basis of Λ is given, for example, in [CS99]. The automorphism group of the Leech lattice is the Conway group Co_0 . M_{24} can be considered as a subgroup of Co_0 , acting on Λ by permuting the coordinates.

In Table 3 (see also Table 1) we list representatives ν of the 11 algebraic conjugacy classes of automorphisms in $O(\Lambda) \cong \text{Co}_0$ that the 70 generalised deep holes project to. For all cycle shapes but $2^3 6^3$ these classes have representatives in M_{24} , allowing us to write them as permutations of the coordinates Ω of $\mathbb{R}^{24}$. We list particularly nice representatives, partly taken from [Gri98], (5.38) and [CS99], Chapter 11. The cycle shape $2^3 6^3$ may be realised as a signed permutation. In Table 3 arrows denote permutations of the corresponding coordinates and a hollow node indicates a sign change.

In Table 4, for each automorphism $\nu \in O(\Lambda)$ in Table 3 (separated by a horizontal line), we list elements $h \in \Lambda_{\mathbb{C}}$ such that $g = \phi_{\eta}(\nu)\sigma_h \in \text{Aut}(V_{\Lambda})$ are generalised deep holes, amounting to a total of 70 generalised deep holes. To improve legibility the least common denominator of h is factored out and written under the vector. The first 23 entries are representatives of the classes of deep holes of Λ as they are given in [CPS82].

Finally, we list the 38 generalised deep holes of V_{Λ} whose orbifolds give the Moonshine module $V^{\natural}$.


TABLE 3. Representatives of 11 conjugacy classes in $O(\Lambda)$.

TABLE 4. Representatives of 70 orbits of the action of $C_{O(\Lambda)}(\nu)$ on $\pi_\nu(\Lambda_C)/\pi_\nu(\Lambda)$ with ν from Table 3 (continued on next page)

<table><tr><td>12</td><td>18</td><td>14</td><td>12</td><td>16</td><td>12</td></tr><tr><td>34</td><td>16</td><td>12</td><td>12</td><td>12</td><td>14</td></tr><tr><td>34</td><td>12</td><td>12</td><td>12</td><td>20</td><td>20</td></tr><tr><td>34</td><td>24</td><td>12</td><td>12</td><td>22</td><td>18</td></tr></table> A1: $23\sqrt{8}$ <table><tr><td>75</td><td>7</td><td>-25</td><td>9</td><td>25</td><td>11</td></tr><tr><td>3</td><td>1</td><td>13</td><td>17</td><td>13</td><td>17</td></tr><tr><td>5</td><td>1</td><td>13</td><td>15</td><td>23</td><td>13</td></tr><tr><td>5</td><td>3</td><td>11</td><td>15</td><td>15</td><td>15</td></tr></table> A4: $25\sqrt{8}$ <table><tr><td>5</td><td>7</td><td>5</td><td>5</td><td>5</td><td>5</td></tr><tr><td>13</td><td>7</td><td>5</td><td>5</td><td>5</td><td>5</td></tr><tr><td>13</td><td>5</td><td>5</td><td>5</td><td>9</td><td>7</td></tr><tr><td>13</td><td>9</td><td>5</td><td>5</td><td>9</td><td>7</td></tr></table> A7: $9\sqrt{8}$ <table><tr><td>39</td><td>3</td><td>-13</td><td>5</td><td>13</td><td>5</td></tr><tr><td>3</td><td>1</td><td>5</td><td>9</td><td>7</td><td>9</td></tr><tr><td>1</td><td>1</td><td>7</td><td>9</td><td>11</td><td>7</td></tr><tr><td>3</td><td>1</td><td>7</td><td>7</td><td>7</td><td>9</td></tr></table> A10: $13\sqrt{8}$ <table><tr><td>15</td><td>1</td><td>-5</td><td>2</td><td>5</td><td>2</td></tr><tr><td>1</td><td>0</td><td>3</td><td>3</td><td>3</td><td>3</td></tr><tr><td>1</td><td>0</td><td>2</td><td>4</td><td>3</td><td>3</td></tr><tr><td>1</td><td>1</td><td>2</td><td>3</td><td>3</td><td>4</td></tr></table> A13: $5\sqrt{8}$ <table><tr><td>12</td><td>1</td><td>-4</td><td>2</td><td>4</td><td>2</td></tr><tr><td>1</td><td>0</td><td>3</td><td>3</td><td>3</td><td>3</td></tr><tr><td>1</td><td>0</td><td>2</td><td>2</td><td>2</td><td>2</td></tr><tr><td>1</td><td>0</td><td>2</td><td>2</td><td>2</td><td>2</td></tr></table> A16: $4\sqrt{8}$ <table><tr><td>2</td><td>2</td><td>2</td><td>2</td><td>2</td><td>2</td></tr><tr><td>4</td><td>2</td><td>2</td><td>2</td><td>2</td><td>2</td></tr><tr><td>4</td><td>4</td><td>2</td><td>2</td><td>2</td><td>2</td></tr><tr><td>4</td><td>2</td><td>2</td><td>2</td><td>2</td><td>2</td></tr></table> A19: $3\sqrt{8}$ <table><tr><td>10</td><td>0</td><td>2</td><td>0</td><td>2</td><td>0</td></tr><tr><td>0</td><td>2</td><td>0</td><td>2</td><td>0</td><td>2</td></tr><tr><td>0</td><td>2</td><td>0</td><td>2</td><td>0</td><td>2</td></tr><tr><td>0</td><td>2</td><td>0</td><td>2</td><td>0</td><td>2</td></tr></table> A22: $3\sqrt{8}$	12	18	14	12	16	12	34	16	12	12	12	14	34	12	12	12	20	20	34	24	12	12	22	18	75	7	-25	9	25	11	3	1	13	17	13	17	5	1	13	15	23	13	5	3	11	15	15	15	5	7	5	5	5	5	13	7	5	5	5	5	13	5	5	5	9	7	13	9	5	5	9	7	39	3	-13	5	13	5	3	1	5	9	7	9	1	1	7	9	11	7	3	1	7	7	7	9	15	1	-5	2	5	2	1	0	3	3	3	3	1	0	2	4	3	3	1	1	2	3	3	4	12	1	-4	2	4	2	1	0	3	3	3	3	1	0	2	2	2	2	1	0	2	2	2	2	2	2	2	2	2	2	4	2	2	2	2	2	4	4	2	2	2	2	4	2	2	2	2	2	10	0	2	0	2	0	0	2	0	2	0	2	0	2	0	2	0	2	0	2	0	2	0	2	<table><tr><td>8</td><td>12</td><td>10</td><td>8</td><td>10</td><td>8</td></tr><tr><td>22</td><td>10</td><td>8</td><td>8</td><td>8</td><td>8</td></tr><tr><td>22</td><td>8</td><td>8</td><td>8</td><td>14</td><td>12</td></tr><tr><td>22</td><td>16</td><td>8</td><td>8</td><td>14</td><td>12</td></tr></table> A2: $15\sqrt{8}$ <table><tr><td>6</td><td>8</td><td>6</td><td>6</td><td>8</td><td>6</td></tr><tr><td>16</td><td>8</td><td>6</td><td>6</td><td>6</td><td>6</td></tr><tr><td>16</td><td>6</td><td>6</td><td>6</td><td>10</td><td>10</td></tr><tr><td>16</td><td>12</td><td>6</td><td>6</td><td>10</td><td>8</td></tr></table> A5: $11\sqrt{8}$ <table><tr><td>24</td><td>2</td><td>-8</td><td>3</td><td>8</td><td>3</td></tr><tr><td>1</td><td>1</td><td>4</td><td>7</td><td>4</td><td>5</td></tr><tr><td>1</td><td>0</td><td>4</td><td>6</td><td>5</td><td>5</td></tr><tr><td>2</td><td>1</td><td>4</td><td>4</td><td>5</td><td>5</td></tr></table> A8: $8\sqrt{8}$ <table><tr><td>18</td><td>2</td><td>-6</td><td>2</td><td>6</td><td>3</td></tr><tr><td>1</td><td>0</td><td>3</td><td>4</td><td>3</td><td>5</td></tr><tr><td>1</td><td>0</td><td>3</td><td>4</td><td>4</td><td>4</td></tr><tr><td>1</td><td>1</td><td>3</td><td>3</td><td>4</td><td>3</td></tr></table> A11: $6\sqrt{8}$ <table><tr><td>3</td><td>3</td><td>3</td><td>3</td><td>3</td><td>3</td></tr><tr><td>7</td><td>3</td><td>3</td><td>3</td><td>3</td><td>3</td></tr><tr><td>7</td><td>3</td><td>3</td><td>3</td><td>5</td><td>5</td></tr><tr><td>7</td><td>3</td><td>3</td><td>3</td><td>5</td><td>5</td></tr></table> A14: $5\sqrt{8}$ <table><tr><td>21</td><td>1</td><td>-7</td><td>5</td><td>7</td><td>3</td></tr><tr><td>1</td><td>1</td><td>5</td><td>5</td><td>3</td><td>3</td></tr><tr><td>1</td><td>1</td><td>5</td><td>5</td><td>3</td><td>3</td></tr><tr><td>1</td><td>1</td><td>5</td><td>5</td><td>3</td><td>3</td></tr></table> A17: $7\sqrt{8}$ <table><tr><td>4</td><td>4</td><td>2</td><td>6</td><td>6</td><td>6</td></tr><tr><td>8</td><td>4</td><td>2</td><td>2</td><td>2</td><td>2</td></tr><tr><td>4</td><td>4</td><td>6</td><td>2</td><td>2</td><td>2</td></tr><tr><td>4</td><td>4</td><td>6</td><td>2</td><td>2</td><td>2</td></tr></table> A20: $5\sqrt{8}$ <table><tr><td>4</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr></table> A23: $1\sqrt{8}$	8	12	10	8	10	8	22	10	8	8	8	8	22	8	8	8	14	12	22	16	8	8	14	12	6	8	6	6	8	6	16	8	6	6	6	6	16	6	6	6	10	10	16	12	6	6	10	8	24	2	-8	3	8	3	1	1	4	7	4	5	1	0	4	6	5	5	2	1	4	4	5	5	18	2	-6	2	6	3	1	0	3	4	3	5	1	0	3	4	4	4	1	1	3	3	4	3	3	3	3	3	3	3	7	3	3	3	3	3	7	3	3	3	5	5	7	3	3	3	5	5	21	1	-7	5	7	3	1	1	5	5	3	3	1	1	5	5	3	3	1	1	5	5	3	3	4	4	2	6	6	6	8	4	2	2	2	2	4	4	6	2	2	2	4	4	6	2	2	2	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	<table><tr><td>45</td><td>6</td><td>-15</td><td>6</td><td>15</td><td>6</td></tr><tr><td>1</td><td>2</td><td>8</td><td>7</td><td>10</td><td>7</td></tr><tr><td>4</td><td>2</td><td>8</td><td>10</td><td>10</td><td>10</td></tr><tr><td>1</td><td>-1</td><td>8</td><td>10</td><td>10</td><td>10</td></tr></table> A3: $15\sqrt{8}$ <table><tr><td>27</td><td>2</td><td>-9</td><td>3</td><td>9</td><td>4</td></tr><tr><td>1</td><td>1</td><td>4</td><td>7</td><td>4</td><td>6</td></tr><tr><td>2</td><td>0</td><td>5</td><td>6</td><td>7</td><td>5</td></tr><tr><td>2</td><td>1</td><td>4</td><td>6</td><td>6</td><td>5</td></tr></table> A6: $9\sqrt{8}$ <table><tr><td>4</td><td>6</td><td>4</td><td>4</td><td>4</td><td>4</td></tr><tr><td>10</td><td>4</td><td>4</td><td>4</td><td>4</td><td>4</td></tr><tr><td>10</td><td>4</td><td>4</td><td>4</td><td>6</td><td>6</td></tr><tr><td>10</td><td>8</td><td>4</td><td>4</td><td>6</td><td>6</td></tr></table> A9: $7\sqrt{8}$ <table><tr><td>18</td><td>3</td><td>-6</td><td>4</td><td>4</td><td>4</td></tr><tr><td>1</td><td>0</td><td>5</td><td>3</td><td>3</td><td>3</td></tr><tr><td>1</td><td>0</td><td>5</td><td>3</td><td>3</td><td>3</td></tr><tr><td>1</td><td>0</td><td>5</td><td>3</td><td>3</td><td>3</td></tr></table> A12: $6\sqrt{8}$ <table><tr><td>27</td><td>3</td><td>-9</td><td>3</td><td>9</td><td>3</td></tr><tr><td>1</td><td>1</td><td>5</td><td>5</td><td>7</td><td>5</td></tr><tr><td>1</td><td>1</td><td>5</td><td>5</td><td>7</td><td>5</td></tr><tr><td>1</td><td>1</td><td>5</td><td>5</td><td>7</td><td>5</td></tr></table> A15: $9\sqrt{8}$ <table><tr><td>5</td><td>3</td><td>5</td><td>3</td><td>3</td><td>3</td></tr><tr><td>9</td><td>3</td><td>5</td><td>3</td><td>3</td><td>3</td></tr><tr><td>9</td><td>3</td><td>3</td><td>5</td><td>5</td><td>5</td></tr><tr><td>9</td><td>3</td><td>3</td><td>5</td><td>5</td><td>5</td></tr></table> A18: $6\sqrt{8}$ <table><tr><td>-1</td><td>1</td><td>2</td><td>0</td><td>2</td><td>0</td></tr><tr><td>1</td><td>1</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>1</td><td>1</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>1</td><td>1</td><td>0</td><td>0</td><td>0</td><td>0</td></tr></table> A21: $1\sqrt{8}$ <table><tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr></table> A24: $1\sqrt{8}$	45	6	-15	6	15	6	1	2	8	7	10	7	4	2	8	10	10	10	1	-1	8	10	10	10	27	2	-9	3	9	4	1	1	4	7	4	6	2	0	5	6	7	5	2	1	4	6	6	5	4	6	4	4	4	4	10	4	4	4	4	4	10	4	4	4	6	6	10	8	4	4	6	6	18	3	-6	4	4	4	1	0	5	3	3	3	1	0	5	3	3	3	1	0	5	3	3	3	27	3	-9	3	9	3	1	1	5	5	7	5	1	1	5	5	7	5	1	1	5	5	7	5	5	3	5	3	3	3	9	3	5	3	3	3	9	3	3	5	5	5	9	3	3	5	5	5	-1	1	2	0	2	0	1	1	0	0	0	0	1	1	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
12	18	14	12	16	12																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
34	16	12	12	12	14																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
34	12	12	12	20	20																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
34	24	12	12	22	18																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
75	7	-25	9	25	11																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
3	1	13	17	13	17																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
5	1	13	15	23	13																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
5	3	11	15	15	15																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
5	7	5	5	5	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
13	7	5	5	5	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
13	5	5	5	9	7																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
13	9	5	5	9	7																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
39	3	-13	5	13	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
3	1	5	9	7	9																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	7	9	11	7																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
3	1	7	7	7	9																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
15	1	-5	2	5	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	3	3	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	2	4	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	2	3	3	4																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
12	1	-4	2	4	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	3	3	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	2	2	2	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	2	2	2	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
2	2	2	2	2	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	2	2	2	2	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	4	2	2	2	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	2	2	2	2	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
10	0	2	0	2	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	2	0	2	0	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	2	0	2	0	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	2	0	2	0	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
8	12	10	8	10	8																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
22	10	8	8	8	8																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
22	8	8	8	14	12																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
22	16	8	8	14	12																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
6	8	6	6	8	6																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
16	8	6	6	6	6																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
16	6	6	6	10	10																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
16	12	6	6	10	8																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
24	2	-8	3	8	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	4	7	4	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	4	6	5	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
2	1	4	4	5	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
18	2	-6	2	6	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	3	4	3	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	3	4	4	4																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	3	3	4	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
3	3	3	3	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
7	3	3	3	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
7	3	3	3	5	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
7	3	3	3	5	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
21	1	-7	5	7	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	5	5	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	5	5	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	5	5	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	4	2	6	6	6																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
8	4	2	2	2	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	4	6	2	2	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	4	6	2	2	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	0	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	0	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	0	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	0	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
45	6	-15	6	15	6																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	2	8	7	10	7																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	2	8	10	10	10																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	-1	8	10	10	10																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
27	2	-9	3	9	4																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	4	7	4	6																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
2	0	5	6	7	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
2	1	4	6	6	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	6	4	4	4	4																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
10	4	4	4	4	4																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
10	4	4	4	6	6																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
10	8	4	4	6	6																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
18	3	-6	4	4	4																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	5	3	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	5	3	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	0	5	3	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
27	3	-9	3	9	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	5	5	7	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	5	5	7	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	5	5	7	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
5	3	5	3	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
9	3	5	3	3	3																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
9	3	3	5	5	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
9	3	3	5	5	5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
-1	1	2	0	2	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	1	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	0	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	0	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	0	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
0	0	0	0	0	0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													

TABLE 4. (continued)

<table><tr><td>10</td><td>4</td><td>-1</td><td>-5</td><td>-7</td><td>7</td></tr><tr><td>0</td><td>-2</td><td>-5</td><td>-1</td><td>-5</td><td>-3</td></tr><tr><td>0</td><td>-2</td><td>-5</td><td>-1</td><td>-5</td><td>-3</td></tr><tr><td>0</td><td>-2</td><td>-5</td><td>-1</td><td>-5</td><td>-3</td></tr></table> C1: $9\sqrt{8}$	10	4	-1	-5	-7	7	0	-2	-5	-1	-5	-3	0	-2	-5	-1	-5	-3	0	-2	-5	-1	-5	-3	<table><tr><td>245</td><td>-142</td><td>100</td><td>47</td><td>56</td><td>-34</td></tr><tr><td>-95</td><td>28</td><td>-66</td><td>15</td><td>38</td><td>0</td></tr><tr><td>-95</td><td>28</td><td>-66</td><td>15</td><td>38</td><td>0</td></tr><tr><td>-95</td><td>28</td><td>-66</td><td>15</td><td>38</td><td>0</td></tr></table> C2: $6\sqrt{8}$	245	-142	100	47	56	-34	-95	28	-66	15	38	0	-95	28	-66	15	38	0	-95	28	-66	15	38	0	<table><tr><td>191</td><td>-1</td><td>-77</td><td>-21</td><td>111</td><td>-9</td></tr><tr><td>3</td><td>51</td><td>-45</td><td>-17</td><td>-9</td><td>-1</td></tr><tr><td>3</td><td>51</td><td>-45</td><td>-17</td><td>-9</td><td>-1</td></tr><tr><td>3</td><td>51</td><td>-45</td><td>-17</td><td>-9</td><td>-1</td></tr></table> C3: $12\sqrt{8}$	191	-1	-77	-21	111	-9	3	51	-45	-17	-9	-1	3	51	-45	-17	-9	-1	3	51	-45	-17	-9	-1
10	4	-1	-5	-7	7																																																																					
0	-2	-5	-1	-5	-3																																																																					
0	-2	-5	-1	-5	-3																																																																					
0	-2	-5	-1	-5	-3																																																																					
245	-142	100	47	56	-34																																																																					
-95	28	-66	15	38	0																																																																					
-95	28	-66	15	38	0																																																																					
-95	28	-66	15	38	0																																																																					
191	-1	-77	-21	111	-9																																																																					
3	51	-45	-17	-9	-1																																																																					
3	51	-45	-17	-9	-1																																																																					
3	51	-45	-17	-9	-1																																																																					
<table><tr><td>-5</td><td>-113</td><td>171</td><td>57</td><td>-69</td><td>-31</td></tr><tr><td>-93</td><td>-37</td><td>-5</td><td>29</td><td>43</td><td>5</td></tr><tr><td>-93</td><td>-37</td><td>-5</td><td>29</td><td>43</td><td>5</td></tr><tr><td>-93</td><td>-37</td><td>-5</td><td>29</td><td>43</td><td>5</td></tr></table> C4: $9\sqrt{8}$	-5	-113	171	57	-69	-31	-93	-37	-5	29	43	5	-93	-37	-5	29	43	5	-93	-37	-5	29	43	5	<table><tr><td>-5</td><td>-2</td><td>-13</td><td>4</td><td>2</td><td>-2</td></tr><tr><td>5</td><td>4</td><td>1</td><td>4</td><td>-2</td><td>0</td></tr><tr><td>5</td><td>4</td><td>1</td><td>4</td><td>-2</td><td>0</td></tr><tr><td>5</td><td>4</td><td>1</td><td>4</td><td>-2</td><td>0</td></tr></table> C5: $3\sqrt{8}$	-5	-2	-13	4	2	-2	5	4	1	4	-2	0	5	4	1	4	-2	0	5	4	1	4	-2	0	<table><tr><td>-2</td><td>0</td><td>-4</td><td>2</td><td>0</td><td>0</td></tr><tr><td>2</td><td>0</td><td>0</td><td>2</td><td>0</td><td>0</td></tr><tr><td>2</td><td>0</td><td>0</td><td>2</td><td>0</td><td>0</td></tr><tr><td>2</td><td>0</td><td>0</td><td>2</td><td>0</td><td>0</td></tr></table> C6: $3\sqrt{8}$	-2	0	-4	2	0	0	2	0	0	2	0	0	2	0	0	2	0	0	2	0	0	2	0	0
-5	-113	171	57	-69	-31																																																																					
-93	-37	-5	29	43	5																																																																					
-93	-37	-5	29	43	5																																																																					
-93	-37	-5	29	43	5																																																																					
-5	-2	-13	4	2	-2																																																																					
5	4	1	4	-2	0																																																																					
5	4	1	4	-2	0																																																																					
5	4	1	4	-2	0																																																																					
-2	0	-4	2	0	0																																																																					
2	0	0	2	0	0																																																																					
2	0	0	2	0	0																																																																					
2	0	0	2	0	0																																																																					
<table><tr><td>-14</td><td>31</td><td>-60</td><td>-35</td><td>-51</td><td>14</td></tr><tr><td>-14</td><td>31</td><td>-60</td><td>-35</td><td>-51</td><td>14</td></tr><tr><td>9</td><td>9</td><td>-22</td><td>-22</td><td>14</td><td>14</td></tr><tr><td>48</td><td>48</td><td>-9</td><td>-9</td><td>55</td><td>55</td></tr></table> D1a: $23\sqrt{8}$	-14	31	-60	-35	-51	14	-14	31	-60	-35	-51	14	9	9	-22	-22	14	14	48	48	-9	-9	55	55	<table><tr><td>14</td><td>8</td><td>-7</td><td>5</td><td>3</td><td>7</td></tr><tr><td>14</td><td>8</td><td>-7</td><td>5</td><td>3</td><td>7</td></tr><tr><td>19</td><td>19</td><td>8</td><td>8</td><td>2</td><td>2</td></tr><tr><td>17</td><td>17</td><td>14</td><td>14</td><td>16</td><td>16</td></tr></table> D1b: $11\sqrt{8}$	14	8	-7	5	3	7	14	8	-7	5	3	7	19	19	8	8	2	2	17	17	14	14	16	16	<table><tr><td>-7</td><td>5</td><td>-22</td><td>-6</td><td>-11</td><td>3</td></tr><tr><td>-7</td><td>5</td><td>-22</td><td>-6</td><td>-11</td><td>3</td></tr><tr><td>-2</td><td>-2</td><td>-11</td><td>-11</td><td>4</td><td>4</td></tr><tr><td>6</td><td>6</td><td>-3</td><td>-3</td><td>14</td><td>14</td></tr></table> D1c: $7\sqrt{8}$	-7	5	-22	-6	-11	3	-7	5	-22	-6	-11	3	-2	-2	-11	-11	4	4	6	6	-3	-3	14	14
-14	31	-60	-35	-51	14																																																																					
-14	31	-60	-35	-51	14																																																																					
9	9	-22	-22	14	14																																																																					
48	48	-9	-9	55	55																																																																					
14	8	-7	5	3	7																																																																					
14	8	-7	5	3	7																																																																					
19	19	8	8	2	2																																																																					
17	17	14	14	16	16																																																																					
-7	5	-22	-6	-11	3																																																																					
-7	5	-22	-6	-11	3																																																																					
-2	-2	-11	-11	4	4																																																																					
6	6	-3	-3	14	14																																																																					
<table><tr><td>-1</td><td>6</td><td>-7</td><td>-12</td><td>-5</td><td>-4</td></tr><tr><td>-1</td><td>6</td><td>-7</td><td>-12</td><td>-5</td><td>-4</td></tr><tr><td>-1</td><td>-1</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>8</td><td>8</td><td>-3</td><td>-3</td><td>5</td><td>5</td></tr></table> D1d: $5\sqrt{8}$	-1	6	-7	-12	-5	-4	-1	6	-7	-12	-5	-4	-1	-1	0	0	0	0	8	8	-3	-3	5	5	<table><tr><td>-2</td><td>2</td><td>-6</td><td>-10</td><td>-2</td><td>-2</td></tr><tr><td>-2</td><td>2</td><td>-6</td><td>-10</td><td>-2</td><td>-2</td></tr><tr><td>3</td><td>3</td><td>2</td><td>2</td><td>0</td><td>0</td></tr><tr><td>3</td><td>3</td><td>-6</td><td>-6</td><td>0</td><td>0</td></tr></table> D1e: $3\sqrt{8}$	-2	2	-6	-10	-2	-2	-2	2	-6	-10	-2	-2	3	3	2	2	0	0	3	3	-6	-6	0	0	<table><tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>1</td><td>1</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>1</td><td>1</td><td>0</td><td>0</td><td>0</td><td>0</td></tr></table> D1f: $1\sqrt{8}$	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	0	1	1	0	0	0	0
-1	6	-7	-12	-5	-4																																																																					
-1	6	-7	-12	-5	-4																																																																					
-1	-1	0	0	0	0																																																																					
8	8	-3	-3	5	5																																																																					
-2	2	-6	-10	-2	-2																																																																					
-2	2	-6	-10	-2	-2																																																																					
3	3	2	2	0	0																																																																					
3	3	-6	-6	0	0																																																																					
0	0	0	0	0	0																																																																					
0	0	0	0	0	0																																																																					
1	1	0	0	0	0																																																																					
1	1	0	0	0	0																																																																					
<table><tr><td>3</td><td>11</td><td>-16</td><td>-16</td><td>-20</td><td>0</td></tr><tr><td>3</td><td>11</td><td>-16</td><td>-16</td><td>-20</td><td>0</td></tr><tr><td>9</td><td>9</td><td>-8</td><td>-8</td><td>1</td><td>1</td></tr><tr><td>21</td><td>21</td><td>0</td><td>0</td><td>21</td><td>21</td></tr></table> D2a: $9\sqrt{8}$	3	11	-16	-16	-20	0	3	11	-16	-16	-20	0	9	9	-8	-8	1	1	21	21	0	0	21	21	<table><tr><td>-6</td><td>4</td><td>-22</td><td>4</td><td>-6</td><td>4</td></tr><tr><td>-6</td><td>4</td><td>-22</td><td>4</td><td>-6</td><td>4</td></tr><tr><td>-17</td><td>-17</td><td>-4</td><td>-4</td><td>6</td><td>6</td></tr><tr><td>1</td><td>1</td><td>-2</td><td>-2</td><td>20</td><td>20</td></tr></table> D2b: $5\sqrt{8}$	-6	4	-22	4	-6	4	-6	4	-22	4	-6	4	-17	-17	-4	-4	6	6	1	1	-2	-2	20	20	<table><tr><td>-2</td><td>4</td><td>-2</td><td>-4</td><td>-4</td><td>-2</td></tr><tr><td>-2</td><td>4</td><td>-2</td><td>-4</td><td>-4</td><td>-2</td></tr><tr><td>1</td><td>1</td><td>0</td><td>0</td><td>0</td><td>0</td></tr><tr><td>3</td><td>3</td><td>-2</td><td>-2</td><td>2</td><td>2</td></tr></table> D2c: $3\sqrt{8}$	-2	4	-2	-4	-4	-2	-2	4	-2	-4	-4	-2	1	1	0	0	0	0	3	3	-2	-2	2	2
3	11	-16	-16	-20	0																																																																					
3	11	-16	-16	-20	0																																																																					
9	9	-8	-8	1	1																																																																					
21	21	0	0	21	21																																																																					
-6	4	-22	4	-6	4																																																																					
-6	4	-22	4	-6	4																																																																					
-17	-17	-4	-4	6	6																																																																					
1	1	-2	-2	20	20																																																																					
-2	4	-2	-4	-4	-2																																																																					
-2	4	-2	-4	-4	-2																																																																					
1	1	0	0	0	0																																																																					
3	3	-2	-2	2	2																																																																					
<table><tr><td>-93</td><td>103</td><td>-64</td><td>-24</td><td>273</td><td>-19</td></tr><tr><td>-93</td><td>103</td><td>95</td><td>95</td><td>-62</td><td>-62</td></tr><tr><td>-93</td><td>103</td><td>95</td><td>95</td><td>-62</td><td>-62</td></tr><tr><td>103</td><td>-93</td><td>78</td><td>78</td><td>-197</td><td>-197</td></tr></table> E1: $8\sqrt{8}$	-93	103	-64	-24	273	-19	-93	103	95	95	-62	-62	-93	103	95	95	-62	-62	103	-93	78	78	-197	-197	<table><tr><td>30</td><td>-22</td><td>71</td><td>117</td><td>-219</td><td>-17</td></tr><tr><td>30</td><td>-22</td><td>-21</td><td>-21</td><td>-59</td><td>-59</td></tr><tr><td>30</td><td>-22</td><td>-21</td><td>-21</td><td>-59</td><td>-59</td></tr><tr><td>-22</td><td>30</td><td>-53</td><td>-53</td><td>39</td><td>39</td></tr></table> E2: $6\sqrt{8}$	30	-22	71	117	-219	-17	30	-22	-21	-21	-59	-59	30	-22	-21	-21	-59	-59	-22	30	-53	-53	39	39	<table><tr><td>-31</td><td>35</td><td>-19</td><td>5</td><td>73</td><td>-11</td></tr><tr><td>-31</td><td>35</td><td>31</td><td>31</td><td>-35</td><td>-35</td></tr><tr><td>-31</td><td>35</td><td>31</td><td>31</td><td>-35</td><td>-35</td></tr><tr><td>35</td><td>-31</td><td>27</td><td>27</td><td>-69</td><td>-69</td></tr></table> E3: $8\sqrt{8}$	-31	35	-19	5	73	-11	-31	35	31	31	-35	-35	-31	35	31	31	-35	-35	35	-31	27	27	-69	-69
-93	103	-64	-24	273	-19																																																																					
-93	103	95	95	-62	-62																																																																					
-93	103	95	95	-62	-62																																																																					
103	-93	78	78	-197	-197																																																																					
30	-22	71	117	-219	-17																																																																					
30	-22	-21	-21	-59	-59																																																																					
30	-22	-21	-21	-59	-59																																																																					
-22	30	-53	-53	39	39																																																																					
-31	35	-19	5	73	-11																																																																					
-31	35	31	31	-35	-35																																																																					
-31	35	31	31	-35	-35																																																																					
35	-31	27	27	-69	-69																																																																					
<table><tr><td>7</td><td>-7</td><td>5</td><td>-7</td><td>-22</td><td>-6</td></tr><tr><td>7</td><td>-7</td><td>-13</td><td>-13</td><td>12</td><td>12</td></tr><tr><td>7</td><td>-7</td><td>-13</td><td>-13</td><td>12</td><td>12</td></tr><tr><td>-7</td><td>7</td><td>-7</td><td>-7</td><td>20</td><td>20</td></tr></table> E4: $4\sqrt{8}$	7	-7	5	-7	-22	-6	7	-7	-13	-13	12	12	7	-7	-13	-13	12	12	-7	7	-7	-7	20	20	<table><tr><td>-5</td><td>7</td><td>-1</td><td>-5</td><td>15</td><td>-1</td></tr><tr><td>-5</td><td>7</td><td>5</td><td>5</td><td>1</td><td>1</td></tr><tr><td>-5</td><td>7</td><td>5</td><td>5</td><td>1</td><td>1</td></tr><tr><td>7</td><td>-5</td><td>3</td><td>3</td><td>-9</td><td>-9</td></tr></table> E5: $4\sqrt{8}$	-5	7	-1	-5	15	-1	-5	7	5	5	1	1	-5	7	5	5	1	1	7	-5	3	3	-9	-9																									
7	-7	5	-7	-22	-6																																																																					
7	-7	-13	-13	12	12																																																																					
7	-7	-13	-13	12	12																																																																					
-7	7	-7	-7	20	20																																																																					
-5	7	-1	-5	15	-1																																																																					
-5	7	5	5	1	1																																																																					
-5	7	5	5	1	1																																																																					
7	-5	3	3	-9	-9																																																																					
<table><tr><td>-3</td><td>-3</td><td>-3</td><td>-7</td><td>-1</td><td>-37</td></tr><tr><td>-3</td><td>-3</td><td>-3</td><td>13</td><td>-5</td><td>-9</td></tr><tr><td>-3</td><td>-3</td><td>-3</td><td>13</td><td>-5</td><td>-5</td></tr><tr><td>13</td><td>13</td><td>13</td><td>-3</td><td>-5</td><td>-5</td></tr></table> F1: $10\sqrt{8}$	-3	-3	-3	-7	-1	-37	-3	-3	-3	13	-5	-9	-3	-3	-3	13	-5	-5	13	13	13	-3	-5	-5	<table><tr><td>1</td><td>-1</td><td>-1</td><td>3</td><td>19</td><td>13</td></tr><tr><td>1</td><td>-1</td><td>1</td><td>1</td><td>-1</td><td>1</td></tr><tr><td>1</td><td>1</td><td>-1</td><td>1</td><td>-1</td><td>-1</td></tr><tr><td>1</td><td>1</td><td>1</td><td>-1</td><td>-1</td><td>-1</td></tr></table> F2: $5\sqrt{8}$	1	-1	-1	3	19	13	1	-1	1	1	-1	1	1	1	-1	1	-1	-1	1	1	1	-1	-1	-1																									
-3	-3	-3	-7	-1	-37																																																																					
-3	-3	-3	13	-5	-9																																																																					
-3	-3	-3	13	-5	-5																																																																					
13	13	13	-3	-5	-5																																																																					
1	-1	-1	3	19	13																																																																					
1	-1	1	1	-1	1																																																																					
1	1	-1	1	-1	-1																																																																					
1	1	1	-1	-1	-1																																																																					
<table><tr><td>36</td><td>46</td><td>-18</td><td>-18</td><td>15</td><td>15</td></tr><tr><td>36</td><td>14</td><td>-18</td><td>8</td><td>15</td><td>-7</td></tr><tr><td>14</td><td>36</td><td>-18</td><td>8</td><td>15</td><td>-7</td></tr><tr><td>14</td><td>-56</td><td>-18</td><td>-18</td><td>15</td><td>15</td></tr></table> G1: $6\sqrt{8}$	36	46	-18	-18	15	15	36	14	-18	8	15	-7	14	36	-18	8	15	-7	14	-56	-18	-18	15	15	<table><tr><td>-2</td><td>4</td><td>1</td><td>1</td><td>0</td><td>0</td></tr><tr><td>-2</td><td>2</td><td>1</td><td>-1</td><td>0</td><td>-4</td></tr><tr><td>2</td><td>-2</td><td>1</td><td>-1</td><td>0</td><td>-4</td></tr><tr><td>2</td><td>4</td><td>1</td><td>1</td><td>0</td><td>0</td></tr></table> G2: $3\sqrt{8}$	-2	4	1	1	0	0	-2	2	1	-1	0	-4	2	-2	1	-1	0	-4	2	4	1	1	0	0																									
36	46	-18	-18	15	15																																																																					
36	14	-18	8	15	-7																																																																					
14	36	-18	8	15	-7																																																																					
14	-56	-18	-18	15	15																																																																					
-2	4	1	1	0	0																																																																					
-2	2	1	-1	0	-4																																																																					
2	-2	1	-1	0	-4																																																																					
2	4	1	1	0	0																																																																					
<table><tr><td>-2</td><td>-2</td><td>-2</td><td>-2</td><td>14</td><td>14</td></tr><tr><td>-2</td><td>-2</td><td>-2</td><td>-2</td><td>14</td><td>14</td></tr><tr><td>-2</td><td>-2</td><td>-2</td><td>-2</td><td>14</td><td>14</td></tr><tr><td>-2</td><td>-2</td><td>-2</td><td>-10</td><td>14</td><td>-34</td></tr></table> H1: $7\sqrt{8}$	-2	-2	-2	-2	14	14	-2	-2	-2	-2	14	14	-2	-2	-2	-2	14	14	-2	-2	-2	-10	14	-34																																																		
-2	-2	-2	-2	14	14																																																																					
-2	-2	-2	-2	14	14																																																																					
-2	-2	-2	-2	14	14																																																																					
-2	-2	-2	-10	14	-34																																																																					
<table><tr><td>-6</td><td>-6</td><td>-6</td><td>0</td><td>3</td><td>9</td></tr><tr><td>-6</td><td>0</td><td>0</td><td>0</td><td>9</td><td>9</td></tr><tr><td>-6</td><td>0</td><td>0</td><td>0</td><td>-15</td><td>-15</td></tr><tr><td>-6</td><td>-6</td><td>-6</td><td>0</td><td>9</td><td>-9</td></tr></table> I1: $4\sqrt{8}$	-6	-6	-6	0	3	9	-6	0	0	0	9	9	-6	0	0	0	-15	-15	-6	-6	-6	0	9	-9																																																		
-6	-6	-6	0	3	9																																																																					
-6	0	0	0	9	9																																																																					
-6	0	0	0	-15	-15																																																																					
-6	-6	-6	0	9	-9																																																																					
<table><tr><td>10</td><td>0</td><td>50</td><td>0</td><td>124</td><td>0</td></tr><tr><td>0</td><td>-42</td><td>0</td><td>70</td><td>0</td><td>-14</td></tr><tr><td>0</td><td>-42</td><td>0</td><td>70</td><td>0</td><td>-14</td></tr><tr><td>0</td><td>-42</td><td>0</td><td>70</td><td>0</td><td>-14</td></tr></table> J1a: $9\sqrt{8}$	10	0	50	0	124	0	0	-42	0	70	0	-14	0	-42	0	70	0	-14	0	-42	0	70	0	-14	<table><tr><td>4</td><td>0</td><td>-16</td><td>0</td><td>-10</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>-4</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>-4</td><td>0</td><td>0</td></tr><tr><td>0</td><td>0</td><td>0</td><td>-4</td><td>0</td><td>0</td></tr></table> J1b: $3\sqrt{8}$	4	0	-16	0	-10	0	0	0	0	-4	0	0	0	0	0	-4	0	0	0	0	0	-4	0	0																									
10	0	50	0	124	0																																																																					
0	-42	0	70	0	-14																																																																					
0	-42	0	70	0	-14																																																																					
0	-42	0	70	0	-14																																																																					
4	0	-16	0	-10	0																																																																					
0	0	0	-4	0	0																																																																					
0	0	0	-4	0	0																																																																					
0	0	0	-4	0	0																																																																					
<table><tr><td>-2</td><td>-2</td><td>-2</td><td>6</td><td>6</td><td>6</td></tr><tr><td>6</td><td>6</td><td>6</td><td>-2</td><td>-2</td><td>-2</td></tr><tr><td>6</td><td>6</td><td>6</td><td>-2</td><td>-2</td><td>-2</td></tr><tr><td>6</td><td>-1</td><td>-1</td><td>7</td><td>7</td><td>-2</td></tr></table> K1: $5\sqrt{8}$	-2	-2	-2	6	6	6	6	6	6	-2	-2	-2	6	6	6	-2	-2	-2	6	-1	-1	7	7	-2																																																		
-2	-2	-2	6	6	6																																																																					
6	6	6	-2	-2	-2																																																																					
6	6	6	-2	-2	-2																																																																					
6	-1	-1	7	7	-2																																																																					

TABLE 5. The 38 generalised deep holes of V_Λ whose corresponding orbifold constructions are isomorphic to $V^\natural$ (and their class in $\text{Co}_1 = \text{O}(\Lambda)/\{\pm 1\}$ [CCN⁺85]).

n	Cycle shape	$\rho(V_\Lambda(g)) - 1$	Co_1
2	$1^{-24}2^{24}$	1/2	1A
3	$1^{-12}3^{12}$	1/3	3A
4	$1^{-8}4^8$	1/4	4A
5	$1^{-6}5^6$	1/5	5A
6	$1^{-6}2^63^{-6}6^6$	1/6	3B
6	$1^{-4}2^{-4}3^46^4$	1/6	6A
6	$1^{-5}2^13^{-1}6^5$	1/6	6D
7	$1^{-4}7^4$	1/7	7A
8	$1^{-4}2^24^{-2}8^4$	1/8	8C
9	$1^{-3}9^3$	1/9	9A
10	$1^{-4}2^45^{-4}10^4$	1/10	5B
10	$1^{-2}2^{-2}5^210^2$	1/10	10A
10	$1^{-3}2^15^{-1}10^3$	1/10	10E
12	$1^{-4}2^43^44^{-4}6^{-4}12^4$	1/12	12A
12	$1^{-2}3^{-2}4^212^2$	1/12	12E
12	$1^{-3}2^23^14^{-1}6^{-2}12^3$	1/12	12K
13	$1^{-2}13^2$	1/13	13A
14	$1^{-3}2^37^{-3}14^3$	1/14	7B
15	$1^{-2}3^25^{-2}15^2$	1/15	15B
16	$1^{-2}2^18^{-1}16^2$	1/16	16B
18	$1^{-3}2^33^26^{-2}9^{-3}18^3$	1/18	9C
18	$1^{-1}2^{-1}9^118^1$	1/18	18A
18	$1^{-2}2^13^16^{-1}9^{-1}18^2$	1/18	18B
20	$1^{-2}2^24^{-2}5^210^{-2}20^2$	1/20	20A
21	$1^{-1}3^{-1}7^121^1$	1/21	21B
22	$1^{-2}2^211^{-2}22^2$	1/22	11A
24	$1^{-2}2^13^24^16^{-1}8^{-2}12^{-1}24^2$	1/24	24B
28	$1^{-1}4^17^{-1}28^1$	1/28	28A
30	$1^{-3}2^33^35^36^{-3}10^{-3}15^{-3}30^3$	1/30	15A
30	$1^{-1}2^13^{-1}5^{-1}6^110^115^{-1}30^1$	1/30	15D
30	$1^{-2}2^23^15^16^{-1}10^{-1}15^{-2}30^2$	1/30	15E
33	$1^{-1}3^111^{-1}33^1$	1/33	33A
36	$1^{-1}2^14^{-1}9^118^{-1}36^1$	1/36	36A
42	$1^{-2}2^23^26^{-2}7^214^{-2}21^{-2}42^2$	1/42	21A
46	$1^{-1}2^123^{-1}46^1$	1/46	23A/B
60	$1^{-1}3^14^15^112^{-1}15^{-1}20^{-1}60^1$	1/60	60A
70	$1^{-1}2^15^17^110^{-1}14^{-1}35^{-1}70^1$	1/70	35A
78	$1^{-1}2^13^16^{-1}13^126^{-1}39^{-1}78^1$	1/78	39A/B

REFERENCES

- [AL70] A. O. L. Atkin and Joseph Lehner. Hecke operators on $\Gamma_0(m)$. *Math. Ann.*, 185:134–160, 1970.
- [BCP97] Wieb Bosma, John Cannon and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3–4):235–265, 1997. <http://magma.maths.usyd.edu.au>.
- [BK04] Bojko N. Bakalov and Victor G. Kac. Twisted modules over lattice vertex algebras. In Heinz-Dietrich Doebner and Vladimir K. Dobrev, editors, *Lie theory and its applications in physics V*, pages 3–26. World Scientific, 2004. (arXiv:math/0402315v3 [math.QA]).
- [BM21] Lea Beneish and Michael H. Mertens. On Weierstrass mock modular forms and a dimension formula for certain vertex operator algebras. *Math. Z.*, 297(1-2):59–80, 2021. (arXiv:1910.06942v3 [math.NT]).
- [Bor85] Richard E. Borcherds. The Leech lattice. *Proc. Roy. Soc. London Ser. A*, 398:365–376, 1985.
- [Bor86] Richard E. Borcherds. Vertex algebras, Kac-Moody algebras, and the Monster. *Proc. Nat. Acad. Sci. U.S.A.*, 83(10):3068–3071, 1986.
- [Car18] Scott Carnahan. 51 constructions of the Moonshine module. *Commun. Number Theory Phys.*, 12(2):305–334, 2018. (arXiv:1707.02954v2 [math.RT]).
- [CCN⁺85] John H. Conway, Robert T. Curtis, Simon P. Norton, Richard A. Parker and Robert A. Wilson. *Atlas of finite groups: Maximal subgroups and ordinary characters for simple groups*. Oxford University Press, 1985. With computational assistance from J. G. Thackray.
- [CL19] Gaëtan Chenevier and Jean Lannes. *Automorphic forms and even unimodular lattices. Kneser neighbors of Niemeier lattices*, volume 69 of *Ergeb. Math. Grenzgeb. (3)*. Springer, 2019.
- [CM16] Scott Carnahan and Masahiko Miyamoto. Regularity of fixed-point vertex operator subalgebras. (arXiv:1603.05645v4 [math.RT]), 2016.
- [CN79] John H. Conway and Simon P. Norton. Monstrous moonshine. *Bull. Lond. Math. Soc.*, 11(3):308–339, 1979.
- [CPS82] John H. Conway, Richard A. Parker and Neil J. A. Sloane. The covering radius of the Leech lattice. *Proc. Roy. Soc. London Ser. A*, 380(1779):261–290, 1982.
- [CS82a] John H. Conway and Neil J. A. Sloane. On the enumeration of lattices of determinant one. *J. Number Theory*, 15(1):83–94, 1982.
- [CS82b] John H. Conway and Neil J. A. Sloane. Twenty-three constructions for the Leech lattice. *Proc. Roy. Soc. London Ser. A*, 381(1781):275–283, 1982.
- [CS99] John H. Conway and Neil J. A. Sloane. *Sphere packings, lattices and groups*, volume 290 of *Grundlehren Math. Wiss.* Springer, 3rd edition, 1999.
- [Del74] Pierre Deligne. La conjecture de Weil. I. *Inst. Hautes Études Sci. Publ. Math.*, (43):273–307, 1974.
- [DGM90] Louise Dolan, Peter Goddard and Paul S. Montague. Conformal field theory, triality and the Monster group. *Phys. Lett. B*, 236(2):165–172, 1990.
- [DGM96] Louise Dolan, Peter Goddard and Paul S. Montague. Conformal field theories, representations and lattice constructions. *Comm. Math. Phys.*, 179(1):61–120, 1996. (arXiv:hep-th/9410029v1).
- [DL96] Chongying Dong and James I. Lepowsky. The algebraic structure of relative twisted vertex operators. *J. Pure Appl. Algebra*, 110(3):259–295, 1996. (arXiv:q-alg/9604022v1).
- [DLM97] Chongying Dong, Haisheng Li and Geoffrey Mason. Regularity of rational vertex operator algebras. *Adv. Math.*, 132(1):148–166, 1997. (arXiv:q-alg/9508018v1).
- [DLM00] Chongying Dong, Haisheng Li and Geoffrey Mason. Modular-invariance of trace functions in orbifold theory and generalized moonshine. *Commun. Math. Phys.*, 214:1–56, 2000. (arXiv:q-alg/9703016v2).
- [DM97] Chongying Dong and Geoffrey Mason. On quantum Galois theory. *Duke Math. J.*, 86(2):305–321, 1997. (arXiv:hep-th/9412037v1).
- [DM04a] Chongying Dong and Geoffrey Mason. Holomorphic vertex operator algebras of small central charge. *Pacific J. Math.*, 213(2):253–266, 2004. (arXiv:math/0203005v1 [math.QA]).
- [DM04b] Chongying Dong and Geoffrey Mason. Rational vertex operator algebras and the effective central charge. *Int. Math. Res. Not.*, 2004(56):2989–3008, 2004. (arXiv:math/0201318v1 [math.QA]).
- [DM06] Chongying Dong and Geoffrey Mason. Integrability of C_2 -cofinite vertex operator algebras. *Int. Math. Res. Not.*, 2006:80468, 2006. (arXiv:math/0601569v1 [math.QA]).

- [DN99] Chongying Dong and Kiyokazu Nagatomo. Automorphism groups and twisted modules for lattice vertex operator algebras. In Naihuan Jing and Kailash C. Misra, editors, *Recent developments in quantum affine algebras and related topics*, volume 248 of *Contemp. Math.*, pages 117–133. Amer. Math. Soc., 1999. (arXiv:math/9808088v1 [math.QA]).
- [Don93] Chongying Dong. Vertex algebras associated with even lattices. *J. Alg.*, 161(1):245–265, 1993.
- [DRX17] Chongying Dong, Li Ren and Feng Xu. On orbifold theory. *Adv. Math.*, 321:1–30, 2017. (arXiv:1507.03306v2 [math.QA]).
- [DS05] Fred Diamond and Jerry Shurman. *A first course in modular forms*, volume 228 of *Grad. Texts in Math.* Springer, 2005.
- [DVVV89] Robbert Dijkgraaf, Cumrun Vafa, Erik Verlinde and Herman Verlinde. The operator algebra of orbifold models. *Comm. Math. Phys.*, 123(3):485–526, 1989.
- [ELMS21] Jethro van Ekeren, Ching Hung Lam, Sven Möller and Hiroki Shimakura. Schellekens’ list and the very strange formula. *Adv. Math.*, 380:107567, 2021. (arXiv:2005.12248v2 [math.QA]).
- [EMS20a] Jethro van Ekeren, Sven Möller and Nils R. Scheithauer. Construction and classification of holomorphic vertex operator algebras. *J. Reine Angew. Math.*, 759:61–99, 2020. (arXiv:1507.08142v3 [math.RT]).
- [EMS20b] Jethro van Ekeren, Sven Möller and Nils R. Scheithauer. Dimension formulae in genus zero and uniqueness of vertex operator algebras. *Int. Math. Res. Not.*, 2020(7):2145–2204, 2020. (arXiv:1704.00478v3 [math.QA]).
- [FLM88] Igor B. Frenkel, James I. Lepowsky and Arne Meurman. *Vertex operator algebras and the Monster*, volume 134 of *Pure Appl. Math.* Academic Press, 1988.
- [Fri11] Robert Fricke. *Die elliptischen Funktionen und ihre Anwendungen. Zweiter Teil. Die algebraischen Ausführungen.* Springer, 2011. Reprint of the 1922 original.
- [GK19] Thomas Gemünden and Christoph A. Keller. Orbifolds of lattice vertex operator algebras at $d = 48$ and $d = 72$. *J. Algebra*, 523:93–118, 2019. (arXiv:1802.10581v1 [math.QA]).
- [Gri98] Robert L. Griess, Jr. *Twelve sporadic groups.* Springer Monogr. Math. Springer, 1998.
- [HM20] Gerald Höhn and Sven Möller. Systematic orbifold constructions of Schellekens’ vertex operator algebras from Niemeier lattices. *J. Lond. Math. Soc.*, to appear, 2020. (arXiv:2010.00849v2 [math.QA]).
- [Höh17] Gerald Höhn. On the genus of the Moonshine module. (arXiv:1708.05990v1 [math.QA]), 2017.
- [Kac90] Victor G. Kac. *Infinite-dimensional Lie algebras.* Cambridge University Press, 3rd edition, 1990.
- [KLL18] Kazuya Kawasetsu, Ching Hung Lam and Xingjun Lin. $\mathbb{Z}_2$ -orbifold construction associated with (-1) -isometry and uniqueness of holomorphic vertex operator algebras of central charge 24. *Proc. Amer. Math. Soc.*, 146(5):1937–1950, 2018. (arXiv:1611.07655v2 [math.QA]).
- [Kna92] Anthony W. Knap. *Elliptic Curves*, volume 40 of *Math. Notes.* Princeton University Press, 1992.
- [Lam11] Ching Hung Lam. On the constructions of holomorphic vertex operator algebras of central charge 24. *Comm. Math. Phys.*, 305(1):153–198, 2011.
- [Lam20] Ching Hung Lam. Cyclic orbifolds of lattice vertex operator algebras having group-like fusions. *Lett. Math. Phys.*, 110(5):1081–1112, 2020. (arXiv:1805.10778v2 [math.QA]).
- [Lep85] James I. Lepowsky. Calculus of twisted vertex operators. *Proc. Natl. Acad. Sci. USA*, 82(24):8295–8299, 1985.
- [Li96] Haisheng Li. Local systems of twisted vertex operators, vertex operator superalgebras and twisted modules. In Chongying Dong and Geoffrey Mason, editors, *Moonshine, the Monster, and Related Topics*, volume 193 of *Contemp. Math.*, pages 203–236. Amer. Math. Soc., 1996. (arXiv:q-alg/9504022v1).
- [LL20] Ching Hung Lam and Xingjun Lin. A holomorphic vertex operator algebra of central charge 24 with the weight one Lie algebra $F_{4,6}A_{2,2}$. *J. Pure Appl. Algebra*, 224(3):1241–1279, 2020. (arXiv:1612.08123v2 [math.QA]).
- [LS12] Ching Hung Lam and Hiroki Shimakura. Quadratic spaces and holomorphic framed vertex operator algebras of central charge 24. *Proc. Lond. Math. Soc. (3)*, 104(3):540–576, 2012. (arXiv:1010.5303v2 [math.QA]).
- [LS15] Ching Hung Lam and Hiroki Shimakura. Classification of holomorphic framed vertex operator algebras of central charge 24. *Amer. J. Math.*, 137(1):111–137, 2015. (arXiv:1209.4677v1 [math.QA]).

- [LS16a] Ching Hung Lam and Hiroki Shimakura. A holomorphic vertex operator algebra of central charge 24 whose weight one Lie algebra has type $A_{6,7}$. *Lett. Math. Phys.*, 106(11):1575–1585, 2016. (arXiv:1606.04688v2 [math.QA]).
- [LS16b] Ching Hung Lam and Hiroki Shimakura. Orbifold construction of holomorphic vertex operator algebras associated to inner automorphisms. *Comm. Math. Phys.*, 342(3):803–841, 2016. (arXiv:1501.05094v2 [math.QA]).
- [LS19] Ching Hung Lam and Hiroki Shimakura. Reverse orbifold construction and uniqueness of holomorphic vertex operator algebras. *Trans. Amer. Math. Soc.*, 372(10):7001–7024, 2019. (arXiv:1606.08979v3 [math.QA]).
- [LS20] Ching Hung Lam and Hiroki Shimakura. On orbifold constructions associated with the Leech lattice vertex operator algebra. *Math. Proc. Cambridge Philos. Soc.*, 168(2):261–285, 2020. (arXiv:1705.01281v2 [math.QA]).
- [McR21] Robert McRae. On rationality for C_2 -cofinite vertex operator algebras. (arXiv:2108.01898v2 [math.QA]), 2021.
- [Miy06] Toshitsune Miyake. *Modular forms*. Springer Monogr. Math. Springer, 2006. Translated from the Japanese by Yoshitaka Maeda, Reprint of the 1989 edition.
- [Miy13] Masahiko Miyamoto. A $\mathbb{Z}_3$ -orbifold theory of lattice vertex operator algebra and $\mathbb{Z}_3$ -orbifold constructions. In Kenji Iohara, Sophie Morier-Genoud and Bertrand Rémy, editors, *Symmetries, Integrable Systems and Representations*, volume 40 of *Springer Proc. Math. Stat.*, pages 319–344. Springer, 2013. (part of arXiv:1003.0237v2 [math.QA]).
- [Miy15] Masahiko Miyamoto. C_2 -cofiniteness of cyclic-orbifold models. *Comm. Math. Phys.*, 335(3):1279–1286, 2015. (arXiv:1306.5031v1 [math.QA]).
- [Möl16] Sven Möller. *A Cyclic Orbifold Theory for Holomorphic Vertex Operator Algebras and Applications*. Ph.D. thesis, Technische Universität Darmstadt, 2016. (arXiv:1611.09843v2 [math.QA]).
- [Mon94] Paul S. Montague. Orbifold constructions and the classification of self-dual $c = 24$ conformal field theories. *Nucl. Phys. B*, 428(1–2):233–258, 1994. (arXiv:hep-th/9403088v1).
- [MS21] Sven Möller and Nils R. Scheithauer. A geometric classification of the holomorphic vertex operator algebras of central charge 24. (arXiv:2112.12291v1 [math.QA]), 2021.
- [MT04] Masahiko Miyamoto and Kenichiro Tanabe. Uniform product of $A_{g,n}(V)$ for an orbifold model V and G -twisted Zhu algebra. *J. Algebra*, 274(1):80–96, 2004. (arXiv:math/0112054v3 [math.QA]).
- [Nie68] Hans-Volker Niemeier. *Definite quadratische Formen der Dimension 24 und Diskriminante 1*. Dissertation, Georg-August-Universität Göttingen, 1968.
- [Nie73] Hans-Volker Niemeier. Definite quadratische Formen der Dimension 24 und Diskriminante 1. *J. Number Theory*, 5:142–178, 1973.
- [Nik80] Viacheslav V. Nikulin. Integral symmetric bilinear forms and some of their applications. *Math. USSR Izv.*, 14(1):103–167, 1980.
- [PS71] George Pólya and Gábor Szegő. *Aufgaben und Lehrsätze aus der Analysis. Band II: Funktionentheorie, Nullstellen, Polynome Determinanten, Zahlentheorie*, volume 74 of *Heidelberger Taschenbücher*. Springer, 4th edition, 1971.
- [Sch93] A. N. Schellekens. Meromorphic $c = 24$ conformal field theories. *Comm. Math. Phys.*, 153(1):159–185, 1993. (arXiv:hep-th/9205072v1).
- [Sch06] Nils R. Scheithauer. On the classification of automorphic products and generalized Kac-Moody algebras. *Invent. Math.*, 164(3):641–678, 2006.
- [Sch09] Nils R. Scheithauer. The Weil representation of $SL_2(\mathbb{Z})$ and some applications. *Int. Math. Res. Not.*, 2009(8):1488–1545, 2009.
- [Sch17] Nils R. Scheithauer. Automorphic products of singular weight. *Compos. Math.*, 153(9):1855–1892, 2017.
- [SS16] Daisuke Sagaki and Hiroki Shimakura. Application of a $\mathbb{Z}_3$ -orbifold construction to the lattice vertex operator algebras associated to Niemeier lattices. *Trans. Amer. Math. Soc.*, 368(3):1621–1646, 2016. (arXiv:1302.4826v4 [math.QA]).
- [Tui92] Michael P. Tuite. Monstrous Moonshine from orbifolds. *Comm. Math. Phys.*, 146(2):277–309, 1992.
- [Tui95] Michael P. Tuite. On the relationship between Monstrous Moonshine and the uniqueness of the Moonshine module. *Comm. Math. Phys.*, 166(3):495–532, 1995. (arXiv:hep-th/9305057v1).
- [Ven80] Boris B. Venkov. On the classification of integral even unimodular 24-dimensional quadratic forms. *Proc. Steklov Inst. Math.*, 148:63–74, 1980.
- [Zhu96] Yongchang Zhu. Modular invariance of characters of vertex operator algebras. *J. Amer. Math. Soc.*, 9(1):237–302, 1996.