

Non-commutative graphs and quantum error correction for a two-mode quantum oscillator

G.G. Amosov^{*1}, A.S. Mokeev^{†1}, and A.N. Pechen^{‡1}

¹Steklov Mathematical Institute of Russian Academy of Sciences, Gubkina str., 8, Moscow 119991, Russia

January 3, 2022

Abstract

An important topic in quantum information is the theory of error correction codes. Practical situations often involve quantum systems with states in an infinite dimensional Hilbert space, for example coherent states. Motivated by these practical needs, we develop the theory of non-commutative graphs, which is a tool to analyze error correction codes, to infinite dimensional Hilbert spaces. As an explicit example, a family of non-commutative graphs associated with the Schrödinger equation describing the dynamics of a two-mode quantum oscillator is constructed and maximal quantum anticliques for these graphs are found.

Keywords: quantum error correction, non-commutative graphs, quantum anticliques, quantum oscillator, two-mode field, coherent states.

1 Introduction

An important topic in quantum information is the theory of error correction codes [1–6] used to protect quantum information from decoherence and other noises. The information is encoded in the density matrix ρ of the system.

^{*}gramos@mi-ras.ru

[†]alexandrmokeev@yandex.ru

[‡]apechen@gmail.com

Given a quantum channel Φ (i.e., a completely positive trace preserving map) which describes information transmission from sender to receiver and includes noise and other destructive for information transmission effects, an error correction code is a set of states in \mathcal{H} which can be exactly distinguished after a transmission via the channel Φ . Quantum error correcting codes are actively studied theoretically [7–12] and experimentally [13, 14]. They can rely on encoding the information in finite-dimensional states of a quantum system, as was originally considered for example in [2], or in infinite-dimensional states as for example using states of a quantum oscillator [7, 12]. Continuous variables are also used, e.g, for generating continuous variable quadripartite cluster and Greenberger-Horne-Zeilinger entangled states of electromagnetic fields [15], studying attacks and security analysis for protocols of quantum cryptography [16, 17], etc. Controlling a two-mode quantum oscillator already appears a nontrivial complex NP-complete problem [18].

Every quantum channel has a Kraus decomposition $\Phi(\rho) = \sum_j V_j \rho V_j^*$ with some operators V_j . The crucial role in finding an appropriate error correction code is played by the linear space \mathcal{V} spanned by the operators $V_j^* V_k$. The famous Knill–Laflamme condition claims that a zero error transmission via channel Φ is possible iff $PVP = CP$ for the orthogonal projection P on the subspace generated by an error correction code [3]. Throughout this paper we call \mathcal{V} a non-commutative graph.

In [19–21] the study of non-commutative graphs generated by covariant resolutions of identity was initiated for finite-dimensional Hilbert spaces. The cases of the commutative circle group [20] and the non-commutative Heisenberg–Weyl group [21] were considered. In this work we consider non-commutative graphs for an infinite dimensional space \mathcal{H} . An example of a non-commutative operator graph generated by the dynamics of diatomic molecule, or coupled oscillators, is explicitly constructed and its maximal anticlique is found.

The paper is organized as follows. Section 2 is devoted to the introduction of basic notions of the theory of non-commutative graphs for analyzing error correction codes in infinite dimensional Hilbert spaces. In Section 3.1 the action of the unitary group used for a generation of graph is obtained in the explicit form. In Section 3.2 a family of graphs generated by this group is constructed and in Section 3.3 their maximal quantum anticliques are found. In the Conclusion Section we give some conclusive remarks.

2 Non-commutative graphs generated by orbits of unitary groups

Definition 1. A **non-commutative graph** is a linear subspace \mathcal{V} of bounded operators in a Hilbert space \mathcal{H} possessing the properties

- $\mathbf{V} \in \mathcal{V}$ implies that $\mathbf{V}^* \in \mathcal{V}$;
- $\mathbf{I} \in \mathcal{V}$

Such objects were introduced in [22] as operator systems and recently re-defined in quantum information theory under the name of a non-commutative graph [23].

In finite-dimensional space, an operator graph \mathcal{V} can be generated by a unitary representation $g \rightarrow \mathbf{U}_g$ in a Hilbert space \mathcal{H} of a compact group G with the Haar measure μ as

$$\mathcal{V} = \text{span}\{\mathbf{U}_g \mathbf{Q} \mathbf{U}_g^*, g \in G\}, \quad (1)$$

Here \mathbf{Q} is an orthogonal projection such that

$$\int_G \mathbf{U}_g \mathbf{Q} \mathbf{U}_g^* d\mu(g) = \mathbf{I} \quad (2)$$

to guarantee that $\mathbf{I} \in \mathcal{V}$. At the moment, only finite dimensional case $\dim \mathcal{H} < +\infty$ has been studied in this framework. Below we consider the infinite dimensional case $\dim \mathcal{H} = +\infty$.

Suppose that the dynamics of the quantum system is determined by the Schrödinger equation

$$i\psi_t = \mathbf{H}\psi. \quad (3)$$

Then instead of (1) and (2), we shall search for a linear subspace \mathcal{V} and a set of orthogonal projections \mathbf{Q}_β with a parameter β belonging to some set \mathfrak{B} such that

$$\mathcal{V} = \overline{\text{span}\{\mathbf{U}_t \mathbf{Q}_\beta \mathbf{U}_t^*, t \in \mathbb{R}, \beta \in \mathfrak{B}\}}, \quad (4)$$

$$\mathbf{I} \in \mathcal{V}, \quad (5)$$

where $(\mathbf{U}_t = e^{-it\mathbf{H}}, t \in \mathbb{R})$ is a strong continuous unitary group. Since \mathcal{H} has infinite dimension, we should take a closure in (4) of the linear span of $\mathbf{U}_t \mathbf{Q}_\beta \mathbf{U}_t^*$ to guarantee that it generates a graph \mathcal{V} . In the finite dimensional case (1) taking closure is not necessary.

Following to [25], we give the following definition.

Definition 2. An orthogonal projection \mathbf{P} such that $\text{rank } \mathbf{P} \geq 2$ is a **quantum anticlique** for a non-commutative graph \mathcal{V} if it satisfies:

$$\dim \mathbf{P}\mathcal{V}\mathbf{P} = 1. \quad (6)$$

Using the spectral order on the set of Hermitian operators ($A > B$ iff $A - B > 0$) we can give the following notion for considering a maximal error correction code.

Definition 3. A quantum anticlique \mathbf{P} for a non-commutative graph \mathcal{V} is called **maximal** if there does not exist a quantum anticlique $\hat{\mathbf{P}}$ for \mathcal{V} such that $\mathbf{P} < \hat{\mathbf{P}}$.

In the next section we consider a two-mode quantum oscillator with the Hilbert space $\mathcal{H} = L^2(\mathbb{R}^2) \cong L^2(\mathbb{R}) \otimes L^2(\mathbb{R})$. We show that the inclusion (5) of the identity operator in \mathcal{V} can take place for a two-mode quantum oscillator described by the Hamiltonian

$$\mathbf{H} = \frac{\mathbf{p}_1^2}{2} + \frac{\mathbf{p}_2^2}{2} + \frac{(\mathbf{q}_2 - \mathbf{q}_1)^2}{2}, \quad (7)$$

where \mathbf{q}_j and \mathbf{p}_j are the position and momentum operators for j th oscillator, $j = 1, 2$. The Hamiltonian (7) is associated with the dynamics of a diatomic molecule. The corresponding operators \mathbf{Q} are found and the associated operator graphs (4) are described in detail. Maximal quantum anticliques for these operator graphs are found. For this system the infinite dimensional projection \mathbf{P} is generated by one-dimensional entangled projections, while \mathbf{Q} is a linear envelope of separable projections.

3 Operator graphs and maximal anticliques for a pair of coupled quantum oscillators

3.1 Explicit action of the group \mathbf{U}_t

We begin by deriving an explicit action of the unitary group \mathbf{U}_t .

Equation (3) in the coordinate representation has the form

$$i \frac{\partial \psi}{\partial t} = -\frac{1}{2} \frac{\partial^2 \psi}{\partial x^2} - \frac{1}{2} \frac{\partial^2 \psi}{\partial y^2} + \frac{(x - y)^2 \psi}{2}. \quad (8)$$

Making change of the variables $\tilde{x} = x + y$ and $\tilde{y} = x - y$ transforms this equation into

$$i \frac{\partial \psi}{\partial t} = -\frac{\partial^2 \psi}{\partial \tilde{x}^2} - \frac{\partial^2 \psi}{\partial \tilde{y}^2} + \frac{\tilde{y}^2}{2} \psi. \quad (9)$$

Here the operator

$$-\frac{\partial^2}{\partial \tilde{y}^2} + \frac{1}{2}\tilde{y}^2$$

is the Hamiltonian of a one-dimensional oscillator with mass $m = 1/2$ and frequency $\omega = \sqrt{2}$. Its eigenstates and the corresponding eigenvalues are

$$\psi_n(\tilde{y}) = \frac{1}{\sqrt{2^n n!} (\sqrt{2\pi})^{\frac{1}{4}}} H_n \left(\frac{\tilde{y}}{\sqrt{2}} \right) e^{-\frac{\tilde{y}^2}{2}}, \quad \lambda_n = \sqrt{2} \left(\frac{1}{2} + n \right),$$

where

$$H_n(\tilde{y}) = (-1)^n e^{\tilde{y}^2} \frac{d}{d\tilde{y}^n} \left(e^{-\tilde{y}^2} \right)$$

are Hermite polynomials, $n = 0, 1, 2, \dots$.

By representing a solution to (9) in the form $\psi(t, \tilde{x}, \tilde{y}) = \sum_{n=0}^{+\infty} c_n(t, \tilde{x}) \psi_n(\tilde{y})$ we obtain the following Cauchy problem for the coefficients $c_n(t, \tilde{x}) = \langle \psi_n(\cdot), \psi(t, \tilde{x}, \cdot) \rangle$:

$$\begin{aligned} i \frac{\partial c_n(t, \tilde{x})}{\partial t} &= -\frac{\partial^2 c_n(t, \tilde{x})}{\partial \tilde{x}^2} + \frac{2n+1}{\sqrt{2}} c_n(t, \tilde{x}) \\ c_n(0, \tilde{x}) &= \langle \psi_n(\cdot), \psi(0, \tilde{x}, \cdot) \rangle \end{aligned}$$

Its solution is given by the Fresnel integral

$$c_n(t, \tilde{x}) = \frac{1}{2\sqrt{i\pi t}} e^{-it\left(\frac{2n+1}{\sqrt{2}}\right)} \int_{-\infty}^{+\infty} e^{\frac{i(\tilde{x}-v)^2}{4t}} c_n(0, v) dv$$

Thus the solution to (8) is given by the formula

$$\begin{aligned} (\mathbf{U}_t \psi)(x, y) &= \sum_{n=0}^{\infty} \frac{1}{2^{\frac{9}{4}} \pi \sqrt{it} 2^n n!} e^{-it\left(\frac{2n+1}{\sqrt{2}}\right)} H_n \left(\frac{x-y}{2^{\frac{1}{4}}} \right) e^{-\frac{(x-y)^2}{2\sqrt{2}}} \\ &\times \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} e^{\frac{i(x+y-(v+u))^2}{4t}} H_n \left(\frac{u-v}{2^{\frac{1}{4}}} \right) e^{-\frac{(u-v)^2}{2\sqrt{2}}} \psi(u, v) du dv. \end{aligned} \quad (10)$$

Lemma 1. *The functions $f_n(y) = \frac{1}{\sqrt{n! 2^n}} H_n(y) e^{-\frac{y^2}{2}}$ satisfy the following equality*

$$\int_{-\infty}^{+\infty} e^{-\frac{ixy}{2t}} e^{i\frac{y^2}{4t}} f_n(y) dy = \frac{\sqrt{4\pi ti}}{\pi^{1/4} \sqrt{1+2ti} \sqrt{n! 2^n} e^{i\frac{x^2}{4t}}} H_n \left(\frac{x}{\sqrt{1+2ti}} \right) e^{-\frac{x^2}{2(1+2ti)}} \quad (11)$$

Proof. Consider the integral operator $K : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ determined by the formula

$$(Kf)(x) = \int_{-\infty}^{+\infty} K(x, y) f(y) dy$$

with the kernel

$$K(x, y) = \frac{1}{\sqrt{4\pi ti}} e^{i\frac{x^2}{4t}} e^{-\frac{ixy}{2t}} e^{i\frac{y^2}{4t}}.$$

Note that the integral operators with the kernels given below coincide

$$\begin{aligned} \frac{d}{dx} K(x, y) &= \frac{ix}{2t} K(x, y) + K(x, y) \left(-\frac{iy}{2t} \right) \\ K(x, y) \frac{d}{dy} &= \frac{ix}{2t} K(x, y) + K(x, y) \left(-\frac{iy}{2t} \right). \end{aligned}$$

It follows that

$$\frac{d}{dx} K = K \frac{d}{dx}.$$

On the other hand,

$$\begin{aligned} xK(x, y) &= -2tiK(x, y) \frac{d}{dy} + K(x, y)y, \\ K(x, y)y &= 2ti \frac{d}{dx} K(x, y) + xK(x, y). \end{aligned}$$

Hence, for the creation operator $a^+ = \frac{x - \frac{d}{dx}}{\sqrt{2}}$ we get

$$Ka^+ = \frac{1}{\sqrt{2}} \left(2ti \frac{d}{dx} K + xK - \frac{d}{dx} K \right) = b^+ K,$$

where

$$b^+ = \frac{x + (2ti - 1) \frac{d}{dx}}{\sqrt{2}}$$

Define $f_0(x) = \frac{1}{\pi^{1/4}} \exp(-x^2/2)$. Then

$$\begin{aligned} (Kf_0)(x) &= \frac{1}{\pi^{1/4}} \frac{1}{\sqrt{4\pi ti}} e^{i\frac{x^2}{4t}} \int_{-\infty}^{+\infty} e^{-\frac{ixy}{2t}} e^{i\frac{y^2}{4t}} e^{-\frac{y^2}{2}} dy \\ &= \frac{1}{\pi^{1/4}} \frac{1}{\sqrt{4\pi ti}} e^{i\frac{x^2}{4t}} \int_{-\infty}^{+\infty} \exp \left(-\frac{(\sqrt{2t-i}y + i\frac{x}{\sqrt{2t-i}})^2}{4t} \right) \exp \left(-\frac{x^2}{4t(2t-i)} \right) dy \\ &= \frac{1}{\pi^{1/4}} \frac{1}{\sqrt{4\pi ti(2t-i)}} e^{-\frac{x^2}{2(1+2ti)}} \int_{-\infty}^{+\infty} \exp \left(-\frac{z^2}{4t} \right) dz = \frac{1}{\pi^{1/4} \sqrt{1+2ti}} e^{-\frac{x^2}{2(1+2ti)}} \equiv g_0(x) \end{aligned}$$

Since

$$f_n(y) = \frac{1}{\sqrt{n!2^n}} H_n(y) e^{-\frac{y^2}{2}} = \frac{1}{\sqrt{n!}} (a^+)^n (f_0)(y)$$

we obtain

$$\begin{aligned} K(f_n)(x) &= K\left(\frac{(a^+)^n}{\sqrt{n!}} f_0\right)(x) = \frac{(b^+)^n}{\sqrt{n!}} K(f_0)(x) = \frac{(b^+)^n}{\sqrt{n!}} (g_0)(x) \\ &= \frac{1}{\pi^{1/4} \sqrt{1+2ti} \sqrt{n!2^n}} H_n\left(\frac{x}{\sqrt{1+2ti}}\right) e^{-\frac{x^2}{2(1+2ti)}}. \end{aligned}$$

□

Denote

$$\psi_{lm}(x, y) = \frac{1}{\sqrt{2}\sqrt{\pi}2^l l! 2^m m!} H_l\left(\frac{x-y}{\sqrt[4]{2}}\right) H_m\left(\frac{x+y}{\sqrt[4]{2}}\right) e^{-\frac{(x-y)^2 + (x+y)^2}{2\sqrt{2}}}. \quad (12)$$

Proposition 1. Unitary group \mathbf{U}_t acts on ψ_{lm} as

$$\begin{aligned} (\mathbf{U}_t \psi_{lm})(x, y) &= \frac{1}{(2\pi)^{\frac{3}{4}} \sqrt{2^l l! 2^m m!} (1 + \sqrt{2}ti)} e^{-it\left(\frac{2l+1}{\sqrt{2}}\right)} \\ &\quad \times H_l\left(\frac{x-y}{\sqrt[4]{2}}\right) e^{-\frac{(x-y)^2}{2\sqrt{2}}} H_m\left(\frac{x+y}{\sqrt[4]{2}\sqrt{1+\sqrt{2}ti}}\right) e^{-\frac{(x+y)^2}{2\sqrt{2}(1+\sqrt{2}ti)}} \end{aligned}$$

Proof. We prove the Proposition by computing the integral

$$\begin{aligned} I &= \frac{1}{\sqrt{it}} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} e^{\frac{i(x+y-(v+u))^2}{4t}} H_n\left(\frac{u-v}{\sqrt[4]{2}}\right) e^{-\frac{(u-v)^2}{2\sqrt{2}}} \\ &\quad H_l\left(\frac{u-v}{\sqrt[4]{2}}\right) H_m\left(\frac{u+v}{\sqrt[4]{2}}\right) e^{-\frac{(u-v)^2 + (u+v)^2}{2\sqrt{2}}} dudv = \\ &= \frac{\sqrt[4]{2}}{\sqrt{it}} \int_{-\infty}^{+\infty} e^{\frac{i(x+y-z\sqrt[4]{2})^2}{4t}} e^{-\frac{z^2}{2}} H_m(z) \int_{-\infty}^{+\infty} H_n(b) H_l(b) e^{-b^2} db dz = \\ &= \frac{\sqrt[4]{2} \delta_{nl} 2^n n! \sqrt{\pi}}{\sqrt{it}} \int_{-\infty}^{+\infty} e^{\frac{i(x+y-z\sqrt[4]{2})^2}{4t}} e^{-\frac{z^2}{2}} H_m(z) dz = \\ &= \frac{\sqrt[4]{2} \delta_{nl} 2^n n! \sqrt{\pi}}{\sqrt{it}} e^{\frac{i(x+y)^2}{4t}} \int_{-\infty}^{+\infty} e^{\frac{-i(x+y)z\sqrt[4]{2}}{2t}} e^{\frac{i\sqrt{2}}{4t} z^2} e^{-\frac{z^2}{2}} H_m(z) dz = \\ &= \frac{2\pi^{\frac{3}{4}} \sqrt{2^n n!} \delta_{nl}}{\sqrt{1+\sqrt{2}ti} \sqrt{2^m m!}} H_m\left(\frac{x+y}{\sqrt[4]{2}\sqrt{1+\sqrt{2}ti}}\right) e^{-\frac{(x+y)^2}{2\sqrt{2}(1+\sqrt{2}ti)}} \end{aligned}$$

□

Below we will use coherent states [26,27]. Given a complex number $\alpha \in \mathbb{C}$, the coherent state $\xi_\alpha \in L^2(\mathbb{R})$ is an eigenvector of the annihilation operator with eigenvalue α . Its explicit form is

$$\xi_\alpha(u) = \frac{1}{\pi^{\frac{1}{4}}} \exp\left(-\frac{|\alpha|^2}{2}\right) \exp\left(-\frac{u^2 - 2\sqrt{2}\alpha u + \alpha^2}{2}\right) \quad (13)$$

Consider in the tensor product of the Hilbert space $\mathcal{H} = L^2(\mathbb{R}^2) \cong L^2(\mathbb{R}) \otimes L^2(\mathbb{R})$ the product of two coherent states

$$\psi_{\alpha\beta}(x, y) = \frac{1}{\sqrt{2}} \xi_\alpha\left(\frac{x+y}{\sqrt{2}}\right) \xi_\beta\left(\frac{x-y}{\sqrt{2}}\right). \quad (14)$$

As a result we have

Corollary 1.

$$(\mathbf{U}_t \psi_{\alpha\beta})(x, y) = \frac{e^{-\frac{it}{\sqrt{2}}}}{\sqrt{2}\sqrt{1+\sqrt{2}ti}} \xi_\alpha\left(\frac{x+y}{\sqrt{2}\sqrt{1+\sqrt{2}ti}}\right) \xi_{e^{-i\sqrt{2}t}\beta}\left(\frac{x-y}{\sqrt{2}}\right).$$

Proof. Taking into account that

$$\psi_{\alpha\beta}(x, y) = e^{-\frac{|\alpha|^2+|\beta|^2}{2}} \sum_{l,m=0}^{+\infty} \frac{\alpha^l \beta^m}{\sqrt{l!} \sqrt{m!}} \psi_{lm}(x, y),$$

where ψ_{lm} are defined by (12), and applying Proposition 1 we get the result.

□

3.2 The graph generated by orbits of (\mathbf{U}_t)

It is straightforward to check that

$$\psi_{\alpha\beta}(x, y) = \frac{1}{\sqrt{2}} \xi_{\alpha+\beta}\left(\frac{x}{\sqrt{2}}\right) \xi_{\alpha-\beta}\left(\frac{y}{\sqrt{2}}\right). \quad (15)$$

Given $\gamma \in \mathbb{C}$ denote

$$\langle x|\gamma\rangle = \frac{1}{\sqrt{2}} \xi_\gamma\left(\frac{x}{\sqrt{2}}\right)$$

the corresponding squeezed coherent state. Taking an arbitrary $\beta \in \mathbb{C}$ let us define the operator

$$\mathbf{Q}_\beta = \frac{1}{\pi} \int_{\mathbb{C}} |\beta + \alpha, \alpha - \beta\rangle \langle \beta + \alpha, \alpha - \beta| d^2\alpha \quad (16)$$

Proposition 2. *The operator \mathbf{Q}_β is an orthogonal projection.*

Proof. The operator is clearly positive. It follows from (14) that

$$\mathbf{Q}_\beta \cong \frac{1}{\pi} \int_{\mathbb{C}} |\alpha\rangle \langle \alpha| d^2\alpha \otimes |\beta\rangle \langle \beta|. \quad (17)$$

Taking into account that

$$\frac{1}{\pi} \int_{\mathbb{C}} |\alpha\rangle \langle \alpha| d^2\alpha = \mathbf{I}_{L^2(\mathbb{R})} \quad (18)$$

we obtain that \mathbf{Q}_β is an orthogonal projection to the subspace

$$\mathcal{H}_\beta = \overline{\text{span}\{\psi_{\alpha\beta}, \alpha \in \mathbb{C}\}}. \quad (19)$$

□

Lemma 2. *Fix $\varphi \in [0, 2\pi)$, then the following inclusion holds*

$$\mathbf{I} \in \overline{\text{span}\{\mathbf{U}_t \mathbf{Q}_{re^{i\varphi}} \mathbf{U}_t^*, t \in \mathbb{R}, r \in \mathbb{R}_+\}}. \quad (20)$$

Moreover the right hand side of (20) doesn't depend on a choice of φ .

Proof. It follows from Corollary 1 that

$$(\mathbf{U}_t \psi_{\alpha\beta})(x, y) = \frac{e^{-\frac{it}{\sqrt{2}}}}{\sqrt{2}\sqrt{1+\sqrt{2}ti}} \xi_\alpha \left(\frac{x-y}{\sqrt[4]{2}\sqrt{1+\sqrt{2}ti}} \right) \xi_{e^{-i\sqrt{2}t}\beta} \left(\frac{x+y}{\sqrt[4]{2}} \right).$$

Taking into account (18) we obtain for (19)

$$\mathbf{U}_t \mathcal{H}_\beta = \mathcal{H}_{e^{-i\sqrt{2}t}\beta}$$

and

$$\mathbf{U}_t \mathbf{Q}_\beta \mathbf{U}_t^* = \mathbf{Q}_{e^{-i\sqrt{2}t}\beta}.$$

Hence a convex hull of the set

$$\{\mathbf{U}_t \mathbf{Q}_{re^{i\varphi}} \mathbf{U}_t^*, t \in \mathbb{R}, r \in \mathbb{R}_+\}$$

contains the integral over all squeezed coherent vectors. The result follows.

□

The next statement immediately follows from Lemma 2.

Theorem 1. *The operator space*

$$\mathcal{V} = \overline{\text{span}\{\mathbf{U}_t \mathbf{Q}_{re^{i\varphi}} \mathbf{U}_t^*, t \in \mathbb{R}, r \in \mathbb{R}_+\}} = \overline{\text{span}\{\mathbf{Q}_{re^{i(-\sqrt{2}t+\varphi)}}, t \in \mathbb{R}, r \in \mathbb{R}_+\}}$$

is a non-commutative graph.

3.3 Maximal anticlique

Fix in $L^2(\mathbb{R})$ a unit vector g_0 and an orthonormal basis $\{\varphi_k\}$, $k = 1, 2, 3, \dots$. Consider pairwise orthogonal unit vectors in $L^2(\mathbb{R}^2)$ defined by the formula

$$\eta_k(x, y) = \varphi_k\left(\frac{x-y}{\sqrt{2}}\right) g_0\left(\frac{x+y}{\sqrt{2}}\right), \quad k = 1, 2, \dots \quad (21)$$

Theorem 2. *The projection*

$$\mathbf{P} = \sum_{k=1}^{\infty} |\eta_k\rangle \langle \eta_k|$$

is a maximal quantum anticlique for \mathcal{V} .

Proof. Using the unitary equivalence like in (17) we get

$$\begin{aligned} \mathbf{P} \mathbf{Q}_{e^{-i\sqrt{2}t}\beta} \mathbf{P} &\cong \left(\sum_{k=1}^{\infty} |\varphi_k\rangle \langle \varphi_k| \otimes |g_0\rangle \langle g_0| \right) \left(\mathbf{I} \otimes |e^{-i\sqrt{2}t}\beta\rangle \langle e^{-i\sqrt{2}t}\beta| \right) \\ &\times \left(\sum_{k=1}^{\infty} |\varphi_k\rangle \langle \varphi_k| \otimes |g_0\rangle \langle g_0| \right) \\ &= |\langle e^{-i\sqrt{2}t}\beta | g_0 \rangle|^2 (\mathbf{I} \otimes |g_0\rangle \langle g_0|) \cong c_{t,\beta} \mathbf{P}. \end{aligned}$$

Hence

$$\dim \mathbf{P} \mathcal{V} \mathbf{P} = 1. \quad (22)$$

Suppose that there exists a quantum anticlique $\hat{\mathbf{P}}$ such that $\mathbf{P} < \hat{\mathbf{P}}$. Then, it should have the form

$$\hat{\mathbf{P}} \cong \mathbf{I} \otimes \tilde{\mathbf{P}},$$

where $\tilde{\mathbf{P}} > |g_0\rangle \langle g_0|$. To satisfy (22) we need

$$\tilde{\mathbf{P}} |e^{-i\sqrt{2}t}\beta\rangle \langle e^{-i\sqrt{2}t}\beta| \tilde{\mathbf{P}} = \tilde{c}_{t,\beta} \tilde{\mathbf{P}} \quad (23)$$

for all t and $\beta = re^{i\varphi}$. It immediately follows from (23) that $\tilde{\mathbf{P}}$ is one-dimensional, i.e. $\tilde{\mathbf{P}} = |g_0\rangle \langle g_0|$. \square

The non-commutative graph of Theorem 1 is generated by elementary errors which act on a density matrix ρ of the oscillators as follows

$$\rho \rightarrow \mathbf{Q}_{re^{i\varphi}} \mathbf{U}_t \rho \mathbf{U}_t^* \mathbf{Q}_{re^{i\varphi}}. \quad (24)$$

Theorem 2 guarantees that encoding the information into pure states defined by (21) implies that these errors can be corrected. The reason is that these errors affect only the state $g_0\left(\frac{x+y}{\sqrt{2}}\right)$ while images of φ_k remain pairwise orthogonal and thus can be distinguished from each other.

4 Conclusion

Motivated by the needs of quantum error correction, we extend the notion of operator graphs to infinite-dimensional Hilbert spaces. Our approach has allowed for the first time to construct operator graphs using the dynamical evolution in an infinite dimensional Hilbert space. As an explicit example, non-commutative operator graphs are constructed which are generated by the orbits of the unitary group given by the solution of the Schrödinger equation describing dynamics of a pair of oscillators. The quantum anticliques for this graph are found and their maximality is proved.

5 Acknowledgments

This work was supported by the Russian Science Foundation (Project No. 17-11-01388) and performed in Steklov Mathematical Institute of Russian Academy of Sciences.

References

- [1] Holevo A.S.: Quantum Systems, Channels, Information: A Mathematical Introduction. De Gruyter, Berlin/Boston (2012)
- [2] Shor P.W.: Scheme for reducing decoherence in quantum computer memory. *Physical Review A* 52, R2493 (1995).
- [3] Knill E. and Laflamme R.: Theory of error-correction codes. *Physical Review A* 55, 900–911 (1997).
- [4] Knill E., Laflamme R. and Viola L.: Theory of quantum error correction for general noise. *Physical Review Letters* 84, 2525 (2000).
- [5] Gaitan F.: Quantum Error Correction and Fault Tolerant Quantum Computing. Taylor & Francis (2008)
- [6] Lidar D. and Brun T. (eds): Quantum Error Correction. Cambridge University Press (2013)
- [7] Gottesman D., Kitaev A., and Preskill J.: Encoding a qubit in an oscillator. *Physical Review A* 64, 012310 (2001).
- [8] Kribs D.W., Laflamme R. and Poulin D.: Unified and generalized approach to quantum error correction. *Physical Review Letters* 94, 180501 (2005).

- [9] Kosut R.L. and Lidar D.A.: Quantum error correction via convex optimization. *Quantum Information Processing* 8, 441 (2009).
- [10] Beny C., Kempf A. and Kribs D.W.: Quantum error correction on infinite-dimensional Hilbert spaces. *Journal of Mathematical Physics* 50, 062108 (2009).
- [11] Kribs D., Mintah C., Nathanson M., Pereira R.: Operator structures and quantum one-way LOCC conditions. *Journal of Mathematical Physics* 58, 092201 (2017).
- [12] Albert V.V. et al: Performance and structure of single-mode bosonic codes. *Physical Review A* 97, 032346 (2018).
- [13] Schindler P. et al: Experimental repetitive quantum error correction. *Science* 332, 1059–1061 (2011).
- [14] Reed M.D. et al: Realization of three-qubit quantum error correction with superconducting circuits. *Nature* 482, 382–385 (2012).
- [15] Su X., Tan A., Jia X. et al: Experimental preparation of quadripartite cluster and Greenberger-Horne-Zeilinger entangled states for continuous variables. *Physical Review Letters* 98, 070502 (2007).
- [16] Avanesov A.S., Kronberg D.A. and Pechen A.N.: Active beam splitting attack applied to differential phase shift quantum key distribution protocol. *P-Adic Numbers Ultrametric Anal. Appl.* 10:3, 222-232 (2018).
- [17] Bochkov M.K. and Trushechkin A.S.: Security of quantum key distribution with detection-efficiency mismatch in the single-photon case: Tight bounds. *Phys. Rev. A* 99, 032308 (2019).
- [18] Bondar D.I. and Pechen A.N.: Uncomputability and complexity of quantum control. *Scientific Reports* (2019, in press). arXiv:1907.10082
- [19] Amosov G.G.: On general properties of non-commutative operator graphs. *Lobachevskii Journal of Mathematics* 39, 304–308 (2018).
- [20] Amosov G.G. and Mokeev A.S.: On non-commutative operator graphs generated by covariant resolutions of identity. *Quantum Information Processing* 17, 325 (2018).
- [21] Amosov G.G. and Mokeev A.S.: On non-commutative operator graphs generated by reducible unitary representation of the Heisenberg-Weyl group. *International Journal of Theoretical Physics* (2018). <https://doi.org/10.1007/s10773-018-3963-4>; arXiv:1812.02515.

- [22] Choi M.D. and Effros E.G.: Injectivity and operator spaces. *J. Funct. Anal.* 24, 156–209 (1977).
- [23] Duan R., Severini S., Winter A.: Zero-error communication via quantum channels, noncommutative graphs and a quantum Lovasz theta function. *IEEE Trans. Inf. Theory* 59, 1164–1174 (2013); arXiv:1002.2514.
- [24] Amosov G.G. and Mokeev A.S.: On construction of anticliques for non-commutative operator graphs. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 456, 5–15 (2017); *J. Math. Sci.* 234, 269–275 (2018).
- [25] Weaver N.: A "quantum" Ramsey theorem for operator systems. *Proc. Amer. Math. Soc.* 145, 4595–4605 (2017).
- [26] Glauber R.G.: Coherent and incoherent states of the radiation field. *Physical Review* 131, 2766–2788 (1963).
- [27] Sudarshan E.C.G.: Equivalence of semiclassical and quantum mechanical descriptions of statistical light beams. *Physical Review Letters*, 10:7, 277–279 (1963).