

New Early Dark Energy

Florian Niedermann^{1,*} and Martin S. Sloth^{1,†}

¹*CP³-Origins, Center for Cosmology and Particle Physics Phenomenology
University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark*

New measurements of the expansion rate of the universe have plunged the standard model of cosmology into a severe crisis. In this letter we propose a simple resolution to the problem. We propose that a first order phase transition in a dark sector in the early universe, before recombination, can resolve the problem. This will lead to a short phase of a New Early Dark Energy (New EDE) component and can explain the observations. Fitting our model to measurements of the Cosmic Microwave Background, Baryonic Acoustic Oscillations, and supernovae yields a significant improvement of the best fit compared with the standard cosmological model without EDE at the cost of only two extra parameters. We find the mean value of the present Hubble parameter in the New EDE model to be $H_0 = 70.4 \pm 1.0 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

PACS numbers: 98.80.Cq, 98.80.-k

INTRODUCTION

Recent measurements of the expansion of the universe have led to an apparent crisis for the standard model of cosmology, the Λ CDM model.

Within the Λ CDM model, we can calculate the evolution of the universe from the earliest times until today, and until recently all our measurements were consistent with the model. In particular, we can use the measurements of the Cosmic Microwave Background (CMB) radiation to infer the present value of the Hubble parameter, H_0 . If the Λ CDM model is correct, this value will have to agree with the value obtained by *directly* measuring the expansion rate today using supernovae redshift measurements. Now, the problem is that the measurements, direct and indirect, do not agree, and this puts the Λ CDM model in a crisis.

The most precise measurements we have of the temperature fluctuations, polarisation and lensing in the CMB radiation are from the European Space Agency satellite, *Planck*, which, assuming the Λ CDM model, infers the value of the expansion rate today to be $H_0 = 67.36 \pm 0.54 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [1]. Comparing that with the expansion rate measured from Cepheids-calibrated supernovae by the *SH₀ES* team [2], $H_0 = 74.03 \pm 1.42 \text{ km s}^{-1} \text{ Mpc}^{-1}$, there is a 4.4σ discrepancy. Other measurements of the current Hubble rate, such as the *H₀LiCoW* [3], are also significantly discrepant with the *Planck* measurement [4].

The *Planck* measurement of the CMB is a very clean experiment with the systematics well under control, and it is therefore unlikely that there is non-understood systematics in the CMB measurements which can explain the discrepancy. The local supernova observations, on the other hand, involves astronomical distance measurements, which are notoriously difficult, and have been plagued by non-understood systematic errors in the past. Various possible sources of systematics have been considered extensively in the literature already. In [5] it is shown that adding up different possible systematic er-

rors in the determination of the distance ladder cannot resolve the discrepancy. Also the effect of the local bulk flow was considered in [6–8] with the same negative conclusion. So far astronomers have no commonly accepted idea of possible systematic effects to explain the discrepancy, and an often echoed conclusion is that new physics beyond the Λ CDM model is required to resolve the tension (see f.ex. [9]). While it is important to continue to look for possible systematic effects, in the present letter, we will rather consider a simple solution in terms of new physics.

We will study the possibility that a *first order phase transition* in a dark sector at zero temperature happened shortly before recombination in the early universe. Such a phase transition will have the effect of lowering an initially high value of the cosmological constant in the early universe down to the value today, inferred from the measurement of H_0 . Effectively this means that there has been an extra component of dark energy in the early universe, providing a short burst of additional repulsion. Currently, an extra component of Early Dark Energy (EDE) is the best way to resolve the tension between the early and late measurements of H_0 [10–17]. So far people have typically considered a dynamical EDE component that disappears due to a 2nd order phase transition of a slowly rolling scalar field. Such scenarios have complications if monomial potentials are used due to the inevitable oscillations of the scalar field after the transition [18]. This problem can be overcome by using specific terms from the non-perturbative form of the axion potential [10–12], which however represents a non-generic choice [13].

On the other hand, we believe that a first order phase transition may be more natural, and also holds in it the potential to fully resolve the discrepancy between the early and late measurements of H_0 . Provided the string landscape [19] is the correct framework for understanding the cosmological constant problem, New EDE could be a manifestation of this. In addition, a first order phase

transition will lead to different experimental signatures, such as gravitational waves; so, there is good reason to believe it might be possible to settle the question by more precise measurements in the future.

THE MODEL

In order to have a change in the vacuum energy due to a field that undergoes a first order phase transition, we will consider a scalar field with two non-degenerate minima at zero temperature. However, if the tunneling probability from the false to the true vacuum is initially high, the field will tunnel immediately and EDE never makes a sizeable contribution. On the other hand, once tunneling commences, we need a large rate in order to produce enough bubbles of true vacuum that will quickly collide and decay. If the rate is too small, then part of the universe will be in the true and part of it in the false vacuum, which will lead to large inhomogeneities ruled out by observations. We therefore require an additional subdominant trigger field, which suddenly, at the right moment, makes the tunneling rate very high. Analogous to previously considered mechanisms for ending inflation [20, 21], we will therefore consider models with a general potential of the form,

$$V(\psi, \phi) = \frac{\lambda}{4}(M^4 + \psi^4) + \frac{1}{2}\beta M^2\psi^2 - \frac{1}{3}\alpha M\psi^3 + \frac{1}{2}m^2\phi^2 + \frac{1}{2}\tilde{\lambda}\phi^2\psi^2, \quad (1)$$

where ψ is the tunneling field and ϕ is the trigger field. The subdominant trigger field will be frozen as long as its mass is smaller than the Hubble rate, but as soon as the Hubble rate drops below its mass, it will start decaying and this will trigger the tunneling of the ψ field. For a second minimum to develop after the point of inflection, we need to impose $\alpha^2 > 4\beta\lambda$, $\beta > 0$. In figure 1, we have included a 3-D visualisation of the evolution of the potential as the trigger field, ϕ , starts evolving along the orange path opening up the new vacuum for ψ , to which it tunnels with high probability.

The decay rate per unit volume is $\Gamma = K \exp(-S_E)$, where K is a determinant factor which is generically set by the energy scale of the phase transition [22, 23] and S_E is the Euclidian action corresponding to a so-called bounce solution [24]. While it is possible to find an analytic expression in the thin wall limit for a single field, the general case requires a numerical approach. In [25] it is argued that a good approximation of the Euclidian action (approximating the potential as being effectively one-dimensional) can be written as

$$S_E \approx \frac{4\pi^2}{3\lambda} (2 - \delta)^{-3} (\alpha_1\delta + \alpha_2\delta^2 + \alpha_3\delta^3), \quad (2)$$




FIG. 1. Schematic plot of the two-field potential in (1). For $H \lesssim m$ the field rolls along the orange line corresponding to $\psi = 0$. At the inflection point (blue dot) the potential (in ψ direction) develops a second minimum which becomes degenerate shortly after (orange dot). The nucleation probability is highest when $\phi = 0$ (red dot). The true vacuum corresponds to the white dot.

with numerically determined coefficients $\alpha_1 = 13.832$, $\alpha_2 = -10.819$, $\alpha_3 = 2.0765$ and

$$\delta = 9 \frac{\lambda}{\alpha^2} \left(\beta + \tilde{\lambda} \frac{\phi^2}{M^2} \right). \quad (3)$$

The important message from this is that S_E diverges in the thin-wall limit ($\delta \rightarrow 2$) and vanishes in the second limit ($\delta \rightarrow 0$). One can check that for $\delta > 9/4 \sim 2$, there is no second minimum. In the present model, we have $K \sim M^4$. As a result, the tunneling rate diminishes in the thin-wall limit and $\Gamma \rightarrow M^4$ as $\delta \rightarrow 0$. Initially, ϕ is slowly rolling until its mass becomes of order the Hubble rate, at which point ϕ begins to oscillate. For a suitable choice of model parameters, δ becomes small when $\phi \rightarrow 0$ for the first time and the vacuum at $\psi = 0$ becomes unstable, initiating the decay of EDE. We demand $S_E \lesssim 10$ which according to (2) and (3) can be easily achieved when α , β and λ are $\mathcal{O}(1)$.¹

At early times, on the other hand, we require the transition rate to be vanishing, which is guaranteed if $\delta \gg 2$. This fixes the initial value of the trigger field, ϕ_{ini} , and can be satisfied consistently with the condition that $\phi_{ini}/M_{pl} \ll 1$, which is sufficient to ensure that the contribution of ϕ to the total energy density is subdominant.

Now, we also have to ensure that EDE, given by the potential energy in the ψ field, gives a sizeable contribution to the energy budget at the time t_* where bubble percolation of the ψ vacuum becomes efficient. If the transition occurs at a redshift of order $z \sim 10^4$, $\lambda \sim \alpha \sim \mathcal{O}(1)$ and there is a sizeable amount of EDE, say a fraction of the total energy density $f_{EDE} \sim 0.1$, we have $M \sim eV$, and a tiny mass scale of order $m \sim 10^{-27} eV$. A microphysical model, explaining the mass hierarchy between

¹ For simplicity we assume to be in a regime where a semi-classical treatment is at least marginally applicable, i.e. $S_E/\hbar > 1$.

the M and the m scale, would be a model of axion monodromy with two axion fields (see [27] for a field theory version). Here, the masses are protected by softly broken shift symmetries.

We also have to make sure that the nucleation itself happens sufficiently quickly. To that end, we define the percolation parameter $p = \Gamma/H^4$. Provided $p \gg 1$, a large number of bubbles is nucleated within one Hubble patch and one Hubble time. At leading order $p \sim M^4/H_*^4 e^{-S_E} \gg 1$, corresponding to a violent percolation process. In fact, for a typical choice of parameters, where the phase transition occurs at $z \sim 10^4$ and $S_E \approx 10$, we find $p \sim 10^{104}$. This means that percolation is extremely efficient and will cover the entire space with bubbles of true vacuum in a tiny fraction of a Hubble time. Therefore, we can treat it as an instantaneous process on cosmological time scales, which takes place at time t_* .

After the space has been filled with bubbles of true vacuum, they expand and start to collide. In our case this happens almost instantly as there is little space left between the bubbles. This coalescence phase is governed by complicated dynamics, which can be studied analytically only in simplified two-bubble scenarios as in [28]. In particular, it leads to the production of density perturbations that are in general not scale invariant and therefore appear to be dangerous. However, this is not a problem, because all bubbles are of microscopical size $\sim 1/M \sim \mu m$ when they collide, whereas the CMB is only sensitive to structures with comoving size greater than about $10h^{-1}\text{Mpc}$ [29]. As part of the collision process the complicated ψ condensate starts to decay. Microscopically, this is due to interactions of ψ within the dark sector, which gets “reheated” during the coalescence phase.

MATCHING CONDITIONS

We use a simple background model describing the instantaneous transition from a background fluid with an equation of state (e.o.s.) parameter that changes from -1 to w_{EDE}^* ,

$$w_{EDE}(t) = \begin{cases} -1 & \text{for } t < t_*, \\ w_{EDE}^* & \text{for } t > t_*, \end{cases} \quad (4)$$

where the transition happens at time t_* . In terms of our field theory model in (1) this corresponds to a situation where all of the liberated vacuum energy is transferred to a fluid with e.o.s. parameter w_{EDE}^* . Note that we are interested in a regime where the energy stored in ϕ can be neglected.

Background Matching

The above condition fixes the evolution of the background energy density uniquely,

$$\bar{\rho}_{EDE}(t) = \bar{\rho}_{EDE}^* \left(\frac{a(t_*)}{a(t)} \right)^{3[1+w_{EDE}(t)]}, \quad (5)$$

where $\bar{\rho}_{EDE}^* = \text{const.}$ The energy density of EDE is normalized with respect to the true vacuum and continuous across the transition. In order to denote the discontinuity of a time dependent function $f(t)$ across the transition surface at time t_* , we introduce the notation

$$[f]_{\pm} = \lim_{\epsilon \rightarrow 0} [f(t_* - \epsilon) - f(t_* + \epsilon)] \equiv f^{(-)} - f^{(+)} . \quad (6)$$

Applying this operation to the Friedmann equations, we then find

$$[H]_{\pm} = 0, \quad (7a)$$

$$\left[\dot{H} \right]_{\pm} = 4\pi G(1 + w_{EDE}^*)\bar{\rho}_{EDE}^*, \quad (7b)$$

where we used the continuity of the background energy density, $[\bar{\rho}]_{\pm} = 0$, which holds due to (5) and the instantaneous character of the transition. The derivation of (7b) also assumes that the e.o.s. of all other fluid components (except for EDE) is preserved during the transition. Besides the EDE component, we also track the evolution of the sub-dominant field ϕ to turn on the phase transition.

Perturbation Matching

It is not enough to implement the modifications on the background level. In fact, neglecting the fluctuations of the EDE sector would be inconsistent after the decay has occurred, making it mandatory to track their evolution. Before the decay, on the other hand, we can set them to zero as EDE simply behaves as a (non-fluctuating) cosmological constant. This raises the issue of how to initialize them at time t_* . Since the transition is allowed to happen at a relatively late stage in the evolution of the primordial plasma (in the extreme case right before recombination), we cannot assume that all relevant modes are outside the horizon. This makes it quite different to the standard problem of choosing adiabatic initial conditions for super-horizon modes. In the specific case of our two-field model, we use the field value of the trigger field to determine the transition. This is motivated by the ϕ dependence of the parameter δ in (3) which controls the exponential in the tunneling rate through (2). The transition surface Σ for both sub- and super-horizon modes is therefore defined by a constant value of the trigger field

$$\phi(t_*, \mathbf{x})|_{\Sigma} = \text{const} . \quad (8)$$

As a consequence, fluctuations in ϕ lead to spatial variations of the time t^* at which the decay takes place. These variations, $\delta\phi(t_*, \mathbf{x}) = \phi(t_*, \mathbf{x}) - \bar{\phi}(t_*)$, then provide the initial conditions for the fluctuations in the EDE fluid after the phase transition.

In order to match the conventions used in the Boltzmann code community, we work in synchronous gauge,

$$ds^2 = -dt^2 + a(t)^2 (\delta_{ij} + h_{ij}) dx^i dx^j, \quad (9)$$

where in momentum space

$$h_{ij} = \frac{k_i k_j}{k^2} h + \left(\frac{k_i k_j}{k^2} - \frac{1}{3} \delta_{ij} \right) \eta, \quad (10)$$

and $h = \delta^{ij} h_{ij}$. In the following we will make use of the equations for the metric perturbations that are first order in time derivatives [30],

$$\frac{1}{2} H \dot{h} - \frac{k^2}{a^2} \eta = 4\pi G \delta\rho, \quad (11a)$$

$$\frac{k^2}{a^2} \dot{\eta} = 4\pi G (\bar{\rho} + \bar{p}) \frac{\theta}{a}, \quad (11b)$$

where $(\bar{\rho} + \bar{p})\theta = \sum_i (\bar{\rho}_i + \bar{p}_i)\theta_i$ and $\delta\rho = \sum_i \delta\rho_i$ are the total divergence of the fluid velocity and the total energy density perturbation, respectively. The dynamical equations have to be supplemented with Israel's matching conditions [26, 31]. They relate the time derivatives of η and h before and after the transition,

$$\left[\dot{h} \right]_{\pm} = -6 [\dot{\eta}]_{\pm} = 6 \left[\dot{H} \right]_{\pm} \frac{\delta\phi(t_*, \mathbf{k})}{\dot{\phi}(t_*)}, \quad (12)$$

where $\left[\dot{H} \right]_{\pm}$ is specified in (7b), and we used the residual gauge freedom in the synchronous gauge to bring the matching conditions on this simple form. So, we see that the discontinuities of $\dot{\eta}$ and \dot{h} are proportional to the spatial variations of the trigger field. We further find that all perturbations without a derivative, including the fluid sector, are continuous, i.e. $[h]_{\pm} = [\eta]_{\pm} = [\delta_i]_{\pm} = [\theta_i]_{\pm} = 0$, where $\delta_i = \delta\rho_i/\bar{\rho}_i$. This does not apply to EDE perturbations because the derivation assumed that the e.o.s. of a particular matter component i is not changing during the transition, in contrast with (4). In fact, as early dark energy behaves as a perfect cosmological constant before the transition, we can consistently set

$$\delta_{EDE} = \theta_{EDE} = \sigma_{EDE} = F_{EDEl} = 0 \quad \text{for } t < t_*, \quad (13)$$

because these fluctuations are not sourced by the gravitational potential in contrast to ordinary fluids. Here, σ_{EDE} and F_{EDEl} (with $l \geq 3$) are the higher multipoles in the Boltzmann hierarchy [30]. We further introduce the notation $\delta_{EDE}^{(+)} \equiv \delta_{EDE}^*$ and $\theta_{EDE}^{(+)} \equiv \theta_{EDE}^*$ to denote

the fluctuations right after the transition. We can now evaluate the discontinuity of Einstein's equations (11) in order to fix δ_{EDE}^* and θ_{EDE}^* , providing the initial conditions for the EDE perturbations after the transition. Using (12) and (7b), we have

$$\delta_{EDE}^* = -3(1 + w_{EDE}^*) H(t_*) \frac{\delta\phi(t_*, \mathbf{k})}{\dot{\phi}(t_*)}, \quad (14a)$$

$$\theta_{EDE}^* = \frac{k^2}{a(t_*)} \frac{\delta\phi(t_*, \mathbf{k})}{\dot{\phi}(t_*)}. \quad (14b)$$

These two equations together with the junction conditions (12) are the main result of this section. They will allow us to consistently implement our model in a Boltzmann code. In order to close the differential system, we assumed the vanishing of the higher multipoles after the transition. Also, we did not need to make any assumption about the spatial momentum k , so the junction conditions apply equally to sub- and super-horizon modes.

DATA ANALYSIS AND RESULTS

In order to fit the New EDE model to the CMB data, we have incorporated the New EDE model into the Boltzmann code *CLASS* [32, 33]. To that end, we make two simplifying assumptions about the underlying field theory model (we intend to relax both assumptions in later work):

- All liberated vacuum energy is ultimately converted to radiation in some dark sector, i.e. $w_{EDE}^* = 1/3$. This requires the presence of a corresponding channel the ψ condensate can decay into.²
- The fluctuations in the descendent EDE fluid are free streaming, i.e. they fulfil the collisionless Boltzmann equations.

The cosmological parameters are then extracted with the Monte Carlo Markov Chain code *MontePython* [35, 36], employing a Metropolis-Hastings algorithm. Compared to Λ CDM we introduce two new parameters: the fraction of EDE before the decay, $f_{EDE} = \bar{\rho}_{EDE}^*/\bar{\rho}(t_*)$, and the logarithm of the mass of the trigger field $\log_{10} m$, which defines the redshift at decay time, z_* , via $H(z_*) = 0.2m$ (corresponding approximately to the first zero crossing of the trigger field). In total we vary 8 parameters $\{\omega_b, \omega_{cdm}, h, \ln 10^{10} A_s, n_s, \tau_{reio}, f_{EDE}, \log_{10} m\}$, on

² This scenario also assumes that there are no sizeable oscillations in ϕ after the transition, which would give rise to a pressureless fluid component (see a similar study in [34]). We checked that this can be realized with a mild tuning of parameters, $\delta_* - 9\beta\lambda/\alpha^2 \ll 1$, where $\delta_* \lesssim 1$. Relaxing this assumption offers the prospect of a unified DM/EDE scenario.




FIG. 2. 2D posterior distributions of H_0 , f_{EDE} , z_* and ω_{cdm} . The SH_0ES measurement is plotted as a dashed line.

which we impose flat priors. The neutrino sector is modelled in terms of two massless and one massive species with $M_\nu = 0.06$ eV. We impose the initial value $\phi_{ini}/M_{pl} = 10^{-5}$ to make sure that the trigger field is always sub-dominant. We will use the following datasets: the most recent SH_0ES measurement, which is $H_0 = 74.03 \pm 1.42 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [2]; the *Pantheon* dataset [38] comprised of 1048 SNe Ia in a range $0.01 < z < 2.3$; the *BOSS DR 12* anisotropic BAO and growth function measurements at redshift $z = 0.38, 0.51$ and 0.61 based on the *CMASS* and *LOWZ* galaxy samples [39], as well as small- z , isotropic BAO measurements of the 6dF Galaxy Survey [40] and the *SDSS DR7* main Galaxy sample [41] at $z = 0.106$ and $z = 0.15$, respectively; the *Planck 2015 TT, TE, EE* and lensing likelihood [42] with all nuisance parameters.

Performing a likelihood analysis shows that the best fit improves by $\Delta\chi^2 = -9.7$ compared to ΛCDM .³ The extracted mean values can be found in Tab. I; in particular, $H_0 = 70.4 \pm 1.0 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (bestfit $H_0 = 71.2 \text{ km s}^{-1} \text{ Mpc}^{-1}$). The contours for a subset of parameters are depicted in Fig. 2, with the first column showing an overlap of the 95% contours between New EDE and SH_0ES .

³ We assume that chains are converged if the Gelman-Rubin criterion [37] yields $R - 1 < 0.01$.

CONCLUSIONS

We have studied a New EDE model where the decay of EDE happens through a first order phase transition instead of a second order phase transition, as has been studied previously in the literature. This model holds in it the potential to fully resolve the discrepancy in H_0 as inferred from early CMB and BAO measurements and late time distance ladder measurements. In our first most simplified implementation of the model (fixing as many free parameters as possible by making simple assumptions) already yields a significant improvement in the fit over the ΛCDM model of $\Delta\chi^2 = -9.7$. It also yields a significant improvement compared to models with extra dark radiation [43–45] of $\Delta\chi^2 = -4.3$. We expect that the model will fit the data even better when the simplifying assumptions made in the present short letter are dropped in future work.

We thank Edmund Copeland, José Espinosa, Nemanja Kaloper, Antonio Padilla and Kari Rummukainen for discussions and/or useful comments on the manuscript. This work is supported by Villum Fonden grant 13384. CP3-Origins is partially funded by the Danish National Research Foundation, grant number DNRF90.

* niedermann@cp3.sdu.dk

† sloth@cp3.sdu.dk

- [1] N. Aghanim *et al.* [Planck Collaboration], arXiv:1807.06209 [astro-ph.CO].
- [2] A. G. Riess, S. Casertano, W. Yuan, L. M. Macri and D. Scolnic, *Astrophys. J.* **876** (2019) no.1, 85 doi:10.3847/1538-4357/ab1422 [arXiv:1903.07603 [astro-ph.CO]].
- [3] G. C.-F. Chen *et al.*, doi:10.1093/mnras/stz2547 arXiv:1907.02533 [astro-ph.CO].
- [4] L. Verde, T. Treu and A. G. Riess, *Nature Astronomy* 2019 doi:10.1038/s41550-019-0902-0 [arXiv:1907.10625 [astro-ph.CO]].
- [5] T. M. Davis, S. R. Hinton, C. Howlett and J. Calcino, arXiv:1907.12639 [astro-ph.CO].
- [6] R. Wojtak, A. Knebe, W. A. Watson, I. T. Iliev, S. He, D. Rapetti, G. Yepes and S. Gottlber, *Mon. Not. Roy. Astron. Soc.* **438** (2014) no.2, 1805 doi:10.1093/mnras/stt2321 [arXiv:1312.0276 [astro-ph.CO]].
- [7] I. Odderskov, S. Hannestad and J. Brandbyge, *JCAP* **1703** (2017) 022 doi:10.1088/1475-7516/2017/03/022 [arXiv:1701.05391 [astro-ph.CO]].
- [8] H. Y. Wu and D. Huterer, *Mon. Not. Roy. Astron. Soc.* **471** (2017) no.4, 4946 doi:10.1093/mnras/stx1967 [arXiv:1706.09723 [astro-ph.CO]].
- [9] J. L. Bernal, L. Verde and A. G. Riess, *JCAP* **1610** (2016) 019 doi:10.1088/1475-7516/2016/10/019 [arXiv:1607.05617 [astro-ph.CO]].
- [10] V. Poulin, T. L. Smith, D. Grin, T. Karwal and M. Kamionkowski, *Phys. Rev. D* **98** (2018) no.8, 083525 doi:10.1103/PhysRevD.98.083525 [arXiv:1806.10608]

	$100\omega_b$	ω_{cdm}	H_0 [km s ⁻¹ Mpc ⁻¹]	$\ln 10^{10} A_s$	n_s	τ_{reio}	f_{EDE}	z_*
Λ CDM	$2.242^{+0.014}_{-0.015}$	0.1175 ± 0.0012	$68.39^{+0.52}_{-0.55}$	$3.084^{+0.023}_{-0.025}$	$0.9696^{+0.0043}_{-0.0045}$	0.077 ± 0.013	–	–
New EDE	$2.252^{+0.017}_{-0.018}$	$0.1242^{+0.0028}_{-0.003}$	$70.43^{+0.95}_{-1.03}$	$3.1^{+0.025}_{-0.027}$	$0.9862^{+0.0083}_{-0.0086}$	$0.075^{+0.014}_{-0.013}$	$0.047^{+0.018}_{-0.017}$	20900^{+1700}_{-8200}

TABLE I. The mean and 1σ errors obtained from the cosmological parameter extraction.

- [astro-ph.CO]].
- [11] V. Poulin, T. L. Smith, T. Karwal and M. Kamionkowski, Phys. Rev. Lett. **122** (2019) no.22, 221301 doi:10.1103/PhysRevLett.122.221301 [arXiv:1811.04083 [astro-ph.CO]].
- [12] T. L. Smith, V. Poulin and M. A. Amin, arXiv:1908.06995 [astro-ph.CO].
- [13] N. Kaloper, arXiv:1903.11676 [hep-th].
- [14] S. Alexander and E. McDonough, Phys. Lett. B **797** (2019) 134830 doi:10.1016/j.physletb.2019.134830 [arXiv:1904.08912 [astro-ph.CO]].
- [15] M. X. Lin, G. Benevento, W. Hu and M. Raveri, Phys. Rev. D **100** (2019) no.6, 063542 doi:10.1103/PhysRevD.100.063542 [arXiv:1905.12618 [astro-ph.CO]].
- [16] E. Hardy and S. Parameswaran, arXiv:1907.10141 [hep-th].
- [17] L. Knox and M. Millea, arXiv:1908.03663 [astro-ph.CO].
- [18] P. Agrawal, F. Y. Cyr-Racine, D. Pinner and L. Randall, arXiv:1904.01016 [astro-ph.CO].
- [19] L. Susskind, In *Carr, Bernard (ed.): Universe or multi-universe?* 247-266 [hep-th/0302219].
- [20] E. J. Copeland, A. R. Liddle, D. H. Lyth, E. D. Stewart and D. Wands, Phys. Rev. D **49** (1994) 6410 doi:10.1103/PhysRevD.49.6410 [astro-ph/9401011].
- [21] M. Cortes and A. R. Liddle, Phys. Rev. D **80** (2009) 083524 doi:10.1103/PhysRevD.80.083524 [arXiv:0905.0289 [astro-ph.CO]].
- [22] C. G. Callan, Jr. and S. R. Coleman, Phys. Rev. D **16** (1977) 1762. doi:10.1103/PhysRevD.16.1762
- [23] A. D. Linde, Nucl. Phys. B **216** (1983) 421 Erratum: [Nucl. Phys. B **223** (1983) 544]. doi:10.1016/0550-3213(83)90293-6, 10.1016/0550-3213(83)90072-X
- [24] S. R. Coleman, Phys. Rev. D **15** (1977) 2929 Erratum: [Phys. Rev. D **16** (1977) 1248]. doi:10.1103/PhysRevD.15.2929, 10.1103/PhysRevD.16.1248
- [25] F. C. Adams, Phys. Rev. D **48** (1993) 2800 doi:10.1103/PhysRevD.48.2800 [hep-ph/9302321].
- [26] N. Deruelle and V. F. Mukhanov, Phys. Rev. D **52** (1995) 5549 doi:10.1103/PhysRevD.52.5549
- [27] N. Kaloper and L. Sorbo, Phys. Rev. Lett. **102** (2009) 121301 doi:10.1103/PhysRevLett.102.121301 [arXiv:0811.1989 [hep-th]].
- [28] S. W. Hawking, I. G. Moss and J. M. Stewart, Phys. Rev. D **26** (1982) 2681. doi:10.1103/PhysRevD.26.2681
- [29] A. R. Liddle and D. Wands, Mon. Not. Roy. Astron. Soc. **253** (1991) 637.
- [30] C. P. Ma and E. Bertschinger, Astrophys. J. **455** (1995) 7 doi:10.1086/176550 [astro-ph/9506072].
- [31] W. Israel, Nuovo Cim. B **44S10** (1966) 1 [Nuovo Cim. B **44** (1966) 1] Erratum: [Nuovo Cim. B **48** (1967) 463]. doi:10.1007/BF02710419, 10.1007/BF02712210
- [32] J. Lesgourgues, arXiv:1104.2932 [astro-ph.IM].
- [33] D. Blas, J. Lesgourgues and T. Tram, JCAP **1107** (2011) 034 doi:10.1088/1475-7516/2011/07/034 [arXiv:1104.2933 [astro-ph.CO]].
- [34] L. Amendola and R. Barbieri, Phys. Lett. B **642** (2006) 192 doi:10.1016/j.physletb.2006.08.069 [hep-ph/0509257].
- [35] B. Audren, J. Lesgourgues, K. Benabed and S. Prunet, JCAP **1302** (2013) 001 doi:10.1088/1475-7516/2013/02/001 [arXiv:1210.7183 [astro-ph.CO]].
- [36] T. Brinckmann and J. Lesgourgues, arXiv:1804.07261 [astro-ph.CO].
- [37] A. Gelman and D. B. Rubin, Statist. Sci. **7** (1992) 457. doi:10.1214/ss/1177011136
- [38] D. M. Scolnic *et al.*, Astrophys. J. **859** (2018) no.2, 101 doi:10.3847/1538-4357/aab9bb [arXiv:1710.00845 [astro-ph.CO]].
- [39] S. Alam *et al.* [BOSS Collaboration], Mon. Not. Roy. Astron. Soc. **470** (2017) no.3, 2617 doi:10.1093/mnras/stx721 [arXiv:1607.03155 [astro-ph.CO]].
- [40] F. Beutler *et al.*, Mon. Not. Roy. Astron. Soc. **416** (2011) 3017 doi:10.1111/j.1365-2966.2011.19250.x [arXiv:1106.3366 [astro-ph.CO]].
- [41] A. J. Ross, L. Samushia, C. Howlett, W. J. Percival, A. Burden and M. Manera, Mon. Not. Roy. Astron. Soc. **449** (2015) no.1, 835 doi:10.1093/mnras/stv154 [arXiv:1409.3242 [astro-ph.CO]].
- [42] N. Aghanim *et al.* [Planck Collaboration], Astron. Astrophys. **594** (2016) A11 doi:10.1051/0004-6361/201526926 [arXiv:1507.02704 [astro-ph.CO]].
- [43] M. Wyman, D. H. Rudd, R. A. Vanderveld and W. Hu, Phys. Rev. Lett. **112** (2014) no.5, 051302 doi:10.1103/PhysRevLett.112.051302 [arXiv:1307.7715 [astro-ph.CO]].
- [44] C. Dvorkin, M. Wyman, D. H. Rudd and W. Hu, Phys. Rev. D **90** (2014) no.8, 083503 doi:10.1103/PhysRevD.90.083503 [arXiv:1403.8049 [astro-ph.CO]].
- [45] G. Rossi, C. Yeche, N. Palanque-Delabrouille and J. Lesgourgues, Phys. Rev. D **92** (2015) no.6, 063505 doi:10.1103/PhysRevD.92.063505 [arXiv:1412.6763 [astro-ph.CO]].