

Algebraic assignments of truth values to experimental quantum propositions

Arkady Bolotin*

Ben-Gurion University of the Negev, Beersheba (Israel)

March 18, 2022

Abstract

Of what are experimental quantum propositions primary bearers? As it is widely accepted in the modern literature, rather than being bearers of truth and falsity, these entities are bearers of probability values. Consequently, their truth values can be regarded as no more than degenerate probabilities (i.e., ones that have only the values 0 and 1). The mathematical motivation for precedence of probabilistic semantics over propositional semantic for the logic of experimental quantum propositions is Gleason's theorem. It proves that the theory of probability measures on closed linear subspaces of a Hilbert space (which represent experimental quantum propositions) does not admit any probability measure having only the values 0 and 1.

By contrast, in the present paper, it is proclaimed that experimental propositions about quantum systems are primary bearers of truth values. As this paper demonstrates, algebraic properties of separable Hilbert spaces of finite dimension equal or greater than 2 do not allow in valuations (that is, truth assignments) which are dispersion-free, i.e., total functions from the set of atomic propositions to the set of two objects, true and false. Providing a probability function can be interpreted as a measure of the (un)certainly in the assignment of truth values, the fact that valuations cannot be dispersion-free gives rise to probabilistic semantics for the logic of experimental quantum propositions.

Keywords: Truth value assignment; Hilbert space; Experimental quantum propositions; Propositional semantic; Probabilistic semantics; Gleason's theorem.

1 Introduction

The relations between logic, probability theory and quantum mechanics are still a matter for investigation. One of the open questions is this: Among *truth* and *probability* pertaining to quantum systems, what is prior to what?

In the literature (see [1, 2, 3, 4] to name but a few), it has been mainly accepted that *quantum probability precedes quantum truth*. That is, rather than being bearers of truth values, experimental quantum propositions – i.e., meaningful declarative sentences that are (or make) statements about a quantum system – are primary bearers of probability values. This means that the logic of the

*Email : arkadyv@bgu.ac.il

experimental quantum propositions presupposes the probability theory of these entities. In other words, probabilistic semantics underlies propositional semantic for the logic of the experimental quantum propositions. As a result, valuations v denoted by

$$v: \mathbb{P} \rightarrow \mathbb{B}_2 \quad , \quad (1)$$

where \mathbb{P} is the set of atomic propositions and \mathbb{B}_2 denotes either the set $\{\text{true}, \text{false}\}$ or the set $\{1, 0\}$, are replaced with probability functions Pr mapping elements of \mathbb{P} to real numbers in the interval $[0, 1]$

$$\text{Pr}: \mathbb{P} \rightarrow \mathbb{R} \quad . \quad (2)$$

Unlike axiomatic probability theory, in which probability functions are usually defined on σ -algebra of subsets of a given sample space Ω , in the formula (2), functions Pr are defined on elements of the set \mathbb{P} (the exposition of the notion of a probability function for the propositional language, which has a countable set of atomic propositions and the usual truth-functional connectives, see in [5, 6]). Therefore, the image of every proposition $P \in \mathbb{P}$ under any function Pr , denoted $\text{Pr}(P)$, is the probability of the proposition P having the truth value of 1 in the given pure state of the system. Along these lines, valuations v are merely degenerate probability functions Pr , namely,

$$[[P]]_v = \begin{cases} 1, & \text{Pr}(P) = 1 \\ 0, & \text{Pr}(P) = 0 \end{cases} \quad , \quad (3)$$

where the double-bracket notation $[[\cdot]]_v$ is used to express $v(\cdot)$.

Since, in accordance with Birkhoff and von Neumann's proposal [7], the mathematical representative of any experimental proposition about the quantum system is a closed linear subspace of a Hilbert space \mathcal{H} associated with the system, it follows that probability functions Pr are defined on closed linear subspaces of \mathcal{H} . Providing the space \mathcal{H} is separable (meaning that it admits an orthonormal basis consisting of a denumerable family of vectors), one can prove that Pr are continuous functions. Because no continuous function in the interval $[0, 1]$ can take only the two values 0 and 1, the continuity of Pr implies that experimental quantum propositions cannot admit any Pr having only the values 0 and 1. This constitutes Gleason's theorem [8].

However, the question of precedence between quantum truth and quantum probability might have a different answer.

Indeed, imagine that v , a function from \mathbb{P} to \mathbb{B}_2 , is not total but *partial*. In that case, for any proposition $P \in \mathbb{P}$, either $[[P]]_v$ belongs to \mathbb{B}_2 or $[[P]]_v$ is undefined (i.e., the proposition P has something called *truth-value gap* [9], meaning that P is neither true nor false). In addition, assume that there exists a probability function Pr from \mathbb{P} to \mathbb{R} mapping each proposition P to $\text{Pr}(P) \in [0, 1]$ in a way that $[[P]]_v \neq [[P']]_v$ implies $\text{Pr}(P) \neq \text{Pr}(P')$, and also

$$\text{Pr}(P) = \begin{cases} 1, & [[P]]_v = 1 \\ 0, & [[P]]_v = 0 \end{cases} \quad . \quad (4)$$

Then, in case $\llbracket P \rrbracket_v$ is neither 1 nor 0, it must be that $\Pr(P) \notin \{0, 1\}$.

Now, consider the following mathematical statement (which can be called the lemma of partial valuation): Without any additional supposition, algebraic properties of separable Hilbert spaces \mathcal{H} cannot allow in truth assignments v which are total functions from \mathbb{P} to \mathbb{B}_2 . If this statement is true, it will enable the same inference as Gleason's theorem does – that experimental quantum propositions do not admit probabilities $\Pr(P)$ which are only $\{0, 1\}$ -valued.

The present paper provides the proof of the above statement.

2 Truth assignments in agreement with quantum theory

Recall that any closed linear subspace of \mathcal{H} is *the range* of some projection operator \hat{P} acting on \mathcal{H} [10], explicitly,

$$\text{ran}(\hat{P}) = \left\{ |\psi\rangle \in \mathcal{H} : \hat{P}|\psi\rangle = |\psi\rangle \right\} \quad . \quad (5)$$

In view of that, the zero subspace $\{0\}$ and the identity subspace \mathcal{H} are *trivial closed linear subspaces* of \mathcal{H} , namely, $\{0\} = \text{ran}(\hat{0})$ and $\mathcal{H} = \text{ran}(\hat{1})$, where $\hat{0}$ and $\hat{1}$ are the zero and identity operators, correspondingly.

Because the set of the eigenvalues of each projection operator \hat{P} is contained in $\{0, 1\}$, one can assume correspondence between an experimental proposition P and a projection operator \hat{P} , which is another way of stating that the mathematical representative of an experimental proposition P is a nontrivial closed linear subspace $\text{ran}(\hat{P})$.

Let the system be in the pure state described by the unit vector $|\Psi\rangle \in \mathcal{H}$, i.e., one that has unit norm, $\langle \Psi | \Psi \rangle = 1$. Then, the truth value of the experimental proposition P in the state $|\Psi\rangle$ can be determined by the formula

$$\llbracket P \rrbracket_v = \mathfrak{P}_{\in}(|\Psi\rangle, \text{ran}(\hat{P})) \quad , \quad (6)$$

where $\mathfrak{P}_{\in}(|\Psi\rangle, \text{ran}(\hat{P}))$ is the image of a couple $(|\Psi\rangle, \text{ran}(\hat{P}))$ under the propositional function (in other words, *predicate*) \mathfrak{P}_{\in} denoted by the mapping

$$\mathfrak{P}_{\in} : \mathfrak{H} \times \wp(\mathcal{H}) \rightarrow \mathbb{B}_2 \quad , \quad (7)$$

in which \mathfrak{H} is the set of all unit vectors in \mathcal{H} and $\wp(\mathcal{H})$ is the set of all closed linear subspaces of \mathcal{H} . The predicate \mathfrak{P}_{\in} is used to indicate set membership: $\mathfrak{P}_{\in}(|\Psi\rangle, \text{ran}(\hat{P})) = 1$ if the vector $|\Psi\rangle$ belongs to $\text{ran}(\hat{P})$; contrastively, $\mathfrak{P}_{\in}(|\Psi\rangle, \text{ran}(\hat{P})) = 0$ if $|\Psi\rangle$ does not belong to $\text{ran}(\hat{P})$.

Consider *the kernel* of the projection operator \hat{P} : It is the closed linear subspace of \mathcal{H} that corresponds the set of vectors $|\phi\rangle$ in \mathcal{H} which are mapped to zero by \hat{P} , i.e.,

$$\ker(\hat{P}) = \text{ran}(\hat{1} - \hat{P}) = \left\{ |\phi\rangle \in \mathcal{H}: \hat{P}|\phi\rangle = 0 \right\} \quad . \quad (8)$$

As every unit vector $|\Psi\rangle \in \mathcal{H}$ can be decomposed uniquely as $|\Psi\rangle = |\psi\rangle + |\phi\rangle$ with $|\psi\rangle = \hat{P}|\Psi\rangle$ and $|\phi\rangle = |\Psi\rangle - \hat{P}|\Psi\rangle = (\hat{1} - \hat{P})|\Psi\rangle$, where $|\psi\rangle \in \text{ran}(\hat{P})$ and $|\phi\rangle \in \ker(\hat{P})$, the subspaces $\text{ran}(\hat{P})$ and $\ker(\hat{P})$ decompose the Hilbert space \mathcal{H} into the direct sum:

$$\mathcal{H} = \text{ran}(\hat{P}) \oplus \ker(\hat{P}) = \text{ran}(\hat{P}) \oplus \text{ran}(\hat{1} - \hat{P}) \quad . \quad (9)$$

Since $|\Psi\rangle \notin \{0\}$, one can infer from here that

$$|\Psi\rangle \in \ker(\hat{P}) \implies |\Psi\rangle \notin \text{ran}(\hat{P}) \quad , \quad (10)$$

i.e., if $|\Psi\rangle$ belongs to $\ker(\hat{P})$, then $|\Psi\rangle$ does not belong to $\text{ran}(\hat{P})$ and so $\mathfrak{P}_{\in}(|\Psi\rangle, \text{ran}(\hat{P})) = 0$.

But suppose $|\Psi\rangle$ belongs to neither $\text{ran}(\hat{P})$ nor $\ker(\hat{P})$, i.e., both $|\Psi\rangle \notin \text{ran}(\hat{P})$ and $|\Psi\rangle \notin \ker(\hat{P})$ are true. This is logically equivalent to truth of *the joint denial* $|\Psi\rangle \in \text{ran}(\hat{P}) \downarrow |\Psi\rangle \in \ker(\hat{P})$ (recall that the joint denial or *logical nor* is a truth-functional operator which produces a result that is the negation of *logical or*, \sqcup).

To take into consideration this case, one can put forward the hypothesis of indistinguishability:

Hypothesis: *The statements $|\Psi\rangle \in \ker(\hat{P})$ and $|\Psi\rangle \in \text{ran}(\hat{P}) \downarrow |\Psi\rangle \in \ker(\hat{P})$ are indistinguishable from one another under the propositional function \mathfrak{P}_{\in} .*

In a semantics defined upon this hypothesis, \mathfrak{P}_{\in} is a total function and presented by

$$\mathfrak{P}_{\in}(|\Psi\rangle, \text{ran}(\hat{P})) = \begin{cases} 1, & |\Psi\rangle \in \text{ran}(\hat{P}) \\ 0, & |\Psi\rangle \in \ker(\hat{P}) \\ 0, & |\Psi\rangle \in \text{ran}(\hat{P}) \downarrow |\Psi\rangle \in \ker(\hat{P}) \end{cases} \quad . \quad (11)$$

Despite being bivalent, such a semantics is not classical since it does not hold the distributive law of classical logic. Let us show this.

Assume that $\neg P$, the negation of the proposition P , corresponds to the projection operator $\hat{1} - \hat{P}$, and the disjunction $P \sqcup \neg P$ is represented by the lattice-theoretic join $\text{ran}(\hat{P}) \vee \text{ran}(\hat{1} - \hat{P})$. This gives

$$\text{ran}(\hat{P}) \vee \text{ran}(\hat{1} - \hat{P}) = \text{ran}(\hat{P}) \oplus \ker(\hat{P}) = \mathcal{H} \quad . \quad (12)$$

Let the unit vector $|\Phi\rangle \in \mathcal{H}$ be such that it belongs to the subspace $\text{ran}(\hat{Q})$, which represents the proposition Q , but with this, $|\Phi\rangle$ does not belong to neither $\text{ran}(\hat{P})$ nor $\text{ran}(\hat{1} - \hat{P})$. Then, according to (11),

$$\mathfrak{P}_{\in}(|\Phi\rangle, \text{ran}(\hat{Q})) = 1 \quad , \quad (13)$$

$$\mathfrak{P}_{\in}(|\Phi\rangle, \text{ran}(\hat{P})) = 0 \quad , \quad (14)$$

$$\mathfrak{P}_{\in}(|\Psi\rangle, \ker(\hat{P})) = 0 \quad . \quad (15)$$

Let us also assume that the lattice-theoretic meet $\text{ran}(\hat{Q}) \wedge \text{ran}(\hat{P})$ is the set-theoretic intersection $\text{ran}(\hat{Q}) \cap \text{ran}(\hat{P})$. In the set builder notation, the latter is written down as:

$$\text{ran}(\hat{Q}) \cap \text{ran}(\hat{P}) = \left\{ |\psi\rangle \in \mathcal{H}: \mathfrak{F}(|\psi\rangle) \right\} \quad , \quad (16)$$

where the rule $\mathfrak{F}(|\psi\rangle)$ is the logical conjunction of two predicates, namely,

$$\mathfrak{F}(|\psi\rangle) = \mathfrak{P}_{\in}(|\psi\rangle, \text{ran}(\hat{Q})) \sqcap \mathfrak{P}_{\in}(|\psi\rangle, \text{ran}(\hat{P})) \quad . \quad (17)$$

If $|\psi\rangle = |\Phi\rangle$, then $\mathfrak{F}(|\Phi\rangle) = 0$, and so the non-zero vector $|\Phi\rangle$ is not an element of $\text{ran}(\hat{Q}) \wedge \text{ran}(\hat{P})$. This indicates that $\text{ran}(\hat{Q}) \wedge \text{ran}(\hat{P}) = \{0\}$ and, likewise, $\text{ran}(\hat{Q}) \wedge \text{ran}(\hat{1} - \hat{P}) = \{0\}$.

Providing $(Q \sqcap P) \sqcup (Q \sqcap \neg P)$ is represented by $(\text{ran}(\hat{Q}) \wedge \text{ran}(\hat{P})) \vee (\text{ran}(\hat{Q}) \wedge \text{ran}(\hat{1} - \hat{P})) = \{0\} \vee \{0\} = \{0\}$, and $Q \sqcap (P \sqcup \neg P)$ is represented by $\text{ran}(\hat{Q}) \wedge \mathcal{H} = \text{ran}(\hat{Q})$, the failure of the distributive law $Q \sqcap (P \sqcup \neg P) = (Q \sqcap P) \sqcup (Q \sqcap \neg P)$ ensues:

$$\llbracket Q \sqcap (P \sqcup \neg P) \rrbracket_v = \mathfrak{P}_{\in}(|\Phi\rangle, \text{ran}(\hat{Q})) = 1 \quad , \quad (18)$$

however

$$\llbracket (Q \sqcap P) \sqcup (Q \sqcap \neg P) \rrbracket_v = \mathfrak{P}_{\in}(|\Phi\rangle, \{0\}) = 0 \quad . \quad (19)$$

The described semantics is identified with quantum logic (of Birkhoff and von Neumann [7]).

It is important to lay stress on the fact that the hypothesis of indistinguishability stated above is neither intuitive nor plausible nor justifiable by experimental evidence. This hypothesis appears to be added just for the purpose of maintaining valuations v as total functions from \mathbb{P} to \mathbb{B}_2 .

Hence, *if a semantics is presumed bivalent but no further hypothesis is assumed*, the predicate \mathfrak{P}_{\in} can only be a partial function, that is,

$$\mathfrak{P}_{\in}(|\Psi\rangle, \text{ran}(\hat{P})) = \begin{cases} 1, & |\Psi\rangle \in \text{ran}(\hat{P}) \\ 0, & |\Psi\rangle \in \ker(\hat{P}) \\ 0/0, & |\Psi\rangle \in \text{ran}(\hat{P}) \downarrow |\Psi\rangle \in \ker(\hat{P}) \end{cases}, \quad (20)$$

where $0/0$ symbolizes a truth-value gap.

In a semantics of this kind, $\mathfrak{P}_{\in}(|\Phi\rangle, \text{ran}(\hat{P})) = 0/0$; therefore, the rule $\mathfrak{F}(|\Phi\rangle)$ cannot be determined, which causes the meet of two subspaces $\text{ran}(\hat{Q})$ and $\text{ran}(\hat{P})$ to be *undecidable*.

What is more, in this semantics, the disjunction $P \sqcup \neg P$ is true even in the case where neither P nor $\neg P$ has a truth value: To be sure, $\llbracket P \sqcup \neg P \rrbracket_v = \mathfrak{P}_{\in}(|\Phi\rangle, \mathcal{H}) = 1$ even though $\llbracket P \rrbracket_v = \mathfrak{P}_{\in}(|\Phi\rangle, \text{ran}(\hat{P})) = 0/0$ and $\llbracket \neg P \rrbracket_v = \mathfrak{P}_{\in}(|\Phi\rangle, \text{ran}(\hat{1} - \hat{P})) = 0/0$.

The above semantics is identified with *supervaluationism*, i.e., the form of *partial logic* (a deeper study of truth-value gaps and logics that allow for truth-value gaps can be found, for example, in [11, 12]).

As one can see from (20), the logic of experimental quantum propositions could have been classical (i.e., total and bivalent), if the joint denial $|\Psi\rangle \in \text{ran}(\hat{P}) \downarrow |\Psi\rangle \in \ker(\hat{P})$ would have been *always false*, i.e., false for all couples $(|\Psi\rangle, \text{ran}(\hat{P}))$.

Let us clarify the reason why this condition was not fulfilled.

3 Set membership from the algebraic perspective

Consider a separable Hilbert space \mathcal{H} of finite dimension n . Let the projection operator \hat{P} acting on $\mathcal{H} = \mathbb{C}^n$ be expressed in terms of the complex $n \times n$ matrix \mathbf{P}

$$\mathbf{P} = \begin{bmatrix} P_{11} & \cdots & P_{1n} \\ \vdots & \ddots & \vdots \\ P_{n1} & \cdots & P_{nn} \end{bmatrix} = (P_{ij})_{i=1, j=1}^{n,n} \in \mathbf{Mat}_{\mathbf{n} \times \mathbf{n}}(\mathbb{C}) \quad . \quad (21)$$

Then, the range of \hat{P} is the same as the span of the column vectors \mathbf{P}_j of the matrix \mathbf{P} , i.e.,

$$\text{ran}(\hat{P}) = \text{Span}(\mathbf{P}_1, \dots, \mathbf{P}_n) \quad , \quad (22)$$

where either $(\mathbf{P}_1, \dots, \mathbf{P}_n)$ is a basis of $\text{ran}(\hat{P})$ or some \mathbf{P}_j can be removed to obtain a basis of $\text{ran}(\hat{P})$; explicitly,

$$\text{ran}(\hat{P}) = \left\{ c_1, \dots, c_n \in \mathbb{C} : c_1 \begin{bmatrix} P_{11} \\ \vdots \\ P_{n1} \end{bmatrix} + \cdots + c_n \begin{bmatrix} P_{1n} \\ \vdots \\ P_{nn} \end{bmatrix} \right\} \quad . \quad (23)$$

Provided a basis set of vectors $\{|e_1\rangle, \dots, |e_n\rangle\}$ in the Hilbert space \mathbb{C}^n , the unit vector $|\Psi\rangle$ describing the pure state of the quantum system can be expressed as the column vector $\Psi \in \mathbf{Mat}_{n \times 1}(\mathbb{C})$ whose i^{th} row has the entry $\langle e_i | \Psi \rangle$.

One can make here the following observation: From the algebraic perspective, the truth of the statement $|\Psi\rangle \in \text{ran}(\hat{P})$ entails the existence of at least one solution to the system of linear equations

$$\mathbf{R}\mathbf{X} = \Psi \quad , \quad (24)$$

where $\mathbf{X} \in \mathbf{Mat}_{m \times 1}(\mathbb{C})$ is the column vector with $m \leq n$ unknowns x_1, \dots, x_m which are put in the place of weights c_1, \dots, c_m for the linearly independent column vectors $\mathbf{P}_1, \dots, \mathbf{P}_m$ of the matrix \mathbf{P} , so that

$$\mathbf{R}\mathbf{X} = \begin{bmatrix} P_{11} \\ \vdots \\ P_{n1} \end{bmatrix} x_1 + \dots + \begin{bmatrix} P_{1m} \\ \vdots \\ P_{nm} \end{bmatrix} x_m \quad . \quad (25)$$

Denoting $(\delta_{ij})_{i=1}^n \in \mathbf{Mat}_{n \times 1}(\mathbb{C})$ by \mathbf{I}_j , the kernel of \hat{P} can be presented as the span of the column vectors $\mathbf{I}_j - \mathbf{P}_j$ of the matrix $\mathbf{I} - \mathbf{P}$, that is,

$$\ker(\hat{P}) = \text{Span}(\mathbf{I}_1 - \mathbf{P}_1, \dots, \mathbf{I}_n - \mathbf{P}_n) \quad , \quad (26)$$

where either $(\mathbf{I}_1 - \mathbf{P}_1, \dots, \mathbf{I}_n - \mathbf{P}_n)$ is a basis of $\ker(\hat{P})$ or some $\mathbf{I}_j - \mathbf{P}_j$ can be removed to obtain a basis of $\ker(\hat{P})$; explicitly,

$$\ker(\hat{P}) = \left\{ c_1, \dots, c_n \in \mathbb{C} : c_1 \begin{bmatrix} 1 - P_{11} \\ \vdots \\ -P_{n1} \end{bmatrix} + \dots + c_j \begin{bmatrix} \vdots \\ \delta_{ij} - P_{ij} \\ \vdots \end{bmatrix} + \dots + c_n \begin{bmatrix} -P_{1n} \\ \vdots \\ 1 - P_{nn} \end{bmatrix} \right\} \quad . \quad (27)$$

Accordingly, to decide whether $|\Psi\rangle$ belongs to $\ker(\hat{P})$ means to answer the question whether the following system of linear equations has at least one solution:

$$\mathbf{K}\mathbf{X} = \Psi \quad , \quad (28)$$

where $\mathbf{X} \in \mathbf{Mat}_{k \times 1}(\mathbb{C})$ is the column vector with $k \leq n$ unknowns x_1, \dots, x_k which substitute weights c_1, \dots, c_k for the linearly independent column vectors of the matrix $\mathbf{I} - \mathbf{P}$ so that

$$\mathbf{K}\mathbf{X} = \begin{bmatrix} 1 - P_{11} \\ \vdots \\ -P_{n1} \end{bmatrix} x_1 + \dots + \begin{bmatrix} \vdots \\ \delta_{ij} - P_{ij} \\ \vdots \end{bmatrix} x_j + \dots + \begin{bmatrix} -P_{1n} \\ \vdots \\ 1 - P_{nk} \end{bmatrix} x_k \quad . \quad (29)$$

To illustrate this observation, consider the Hilbert space \mathbb{C}^4 characterizing the spin $\frac{3}{2}$ system. In terms of the complex 4×4 matrix, the projection operator $\hat{Y}_{+3/2}$ corresponding to the experimental atomic proposition “The spin of the system along the Y axis is $+\frac{3}{2}\hbar$ ”, denoted $Y_{+3/2}$, takes the form

$$\mathbf{Y}_{+3/2} = \frac{1}{8} \begin{bmatrix} 1 & -i\sqrt{3} & -\sqrt{3} & i \\ i\sqrt{3} & 3 & -i3 & -\sqrt{3} \\ -\sqrt{3} & i3 & 3 & -i\sqrt{3} \\ -i & -\sqrt{3} & i\sqrt{3} & 1 \end{bmatrix} ; \quad (30)$$

its range and kernel are

$$\text{ran}(\hat{Y}_{+3/2}) = \left\{ a \in \mathbb{C} : a \begin{bmatrix} 1 \\ i\sqrt{3} \\ -\sqrt{3} \\ -i \end{bmatrix} \right\} , \quad (31)$$

$$\text{ker}(\hat{Y}_{+3/2}) = \left\{ a, b, c \in \mathbb{C} : a \begin{bmatrix} 7 \\ -i\sqrt{3} \\ \sqrt{3} \\ i \end{bmatrix} + b \begin{bmatrix} i\sqrt{3} \\ 5 \\ -i3 \\ \sqrt{3} \end{bmatrix} + c \begin{bmatrix} \sqrt{3} \\ i3 \\ 5 \\ -i\sqrt{3} \end{bmatrix} \right\} . \quad (32)$$

Suppose that the spin $\frac{3}{2}$ system is in the pure state that is described by the ket $|Y_{+3/2}\rangle$ expressed as the column vector written out in the coordinates over the z -basis, namely,

$$|Y_{+3/2}\rangle = \frac{1}{2\sqrt{2}} \begin{bmatrix} i \\ -\sqrt{3} \\ -i\sqrt{3} \\ 1 \end{bmatrix} . \quad (33)$$

To decide whether the statement $|Y_{+3/2}\rangle \in \text{ran}(\hat{Y}_{+3/2})$ is true, let us present it as the system of linear equations

$$\begin{bmatrix} 1 \\ i\sqrt{3} \\ -\sqrt{3} \\ -i \end{bmatrix} x = \frac{1}{2\sqrt{2}} \begin{bmatrix} i \\ -\sqrt{3} \\ -i\sqrt{3} \\ 1 \end{bmatrix} . \quad (34)$$

Even though this linear system is overdetermined, it contains 3 linearly dependent equations; hence, it has the solution, $x = i\frac{\sqrt{2}}{4}$, which means that the ket $|Y_{+3/2}\rangle$ belongs to $\text{ran}(\hat{Y}_{+3/2})$, and so in the state described by $|Y_{+3/2}\rangle$, the proposition $Y_{+3/2}$ is true:

$$\llbracket Y_{+3/2} \rrbracket_v = \mathfrak{P}_{\in}(|Y_{+3/2}\rangle, \text{ran}(\hat{Y}_{+3/2})) = 1 . \quad (35)$$

Yet, the linear system with the same left-hand side as (34) but the different right-hand side, namely,

$$|Y_{+1/2}\rangle = \frac{1}{2\sqrt{2}} \begin{bmatrix} -i\sqrt{3} \\ 1 \\ -i \\ \sqrt{3} \end{bmatrix} , \quad (36)$$

has no solution. To confirm that the ket $|Y_{+1/2}\rangle$ does not belong to $\text{ran}(\hat{Y}_{+3/2})$, consider the statement $|Y_{+1/2}\rangle \in \ker(\hat{Y}_{+3/2})$ and present it as the system of linear equations, that is,

$$\begin{bmatrix} 7 & i\sqrt{3} & \sqrt{3} \\ -i\sqrt{3} & 5 & i3 \\ \sqrt{3} & -i3 & 5 \\ -i & \sqrt{3} & -i\sqrt{3} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \frac{1}{2\sqrt{2}} \begin{bmatrix} -i\sqrt{3} \\ 1 \\ -i \\ \sqrt{3} \end{bmatrix} . \quad (37)$$

Although this linear system is overdetermined (i.e., $\dim(\mathbf{X}) = 3 < 4$), it has the solution

$$\mathbf{X} = \frac{1}{8\sqrt{2}} \begin{bmatrix} -i\sqrt{3} \\ 2 \\ i \end{bmatrix} , \quad (38)$$

therefore, the statement $|Y_{+1/2}\rangle \in \ker(\hat{Y}_{+3/2})$ is true. This implies that ket $|Y_{+1/2}\rangle$ does not belong to $\text{ran}(\hat{Y}_{+3/2})$, and so in the state $|Y_{+1/2}\rangle$ the proposition $Y_{+3/2}$ is false:

$$[[Y_{+3/2}]]_v = \mathfrak{P}_{\in}(|Y_{+1/2}\rangle, \text{ran}(\hat{Y}_{+3/2})) = 0 \quad . \quad (39)$$

By contrast, in case the state of the spin $3/2$ system is described by the ket $|X_{+3/2}\rangle$ identified with the column vector

$$|X_{+3/2}\rangle = \frac{1}{2\sqrt{2}} \begin{bmatrix} 1 \\ \sqrt{3} \\ \sqrt{3} \\ 1 \end{bmatrix} , \quad (40)$$

neither the equations

$$\begin{bmatrix} 1 \\ i\sqrt{3} \\ -\sqrt{3} \\ -i \end{bmatrix} x = \frac{1}{2\sqrt{2}} \begin{bmatrix} 1 \\ \sqrt{3} \\ \sqrt{3} \\ 1 \end{bmatrix} , \quad (41)$$

nor the equations

$$\begin{bmatrix} 7 & i\sqrt{3} & \sqrt{3} \\ -i\sqrt{3} & 5 & i3 \\ \sqrt{3} & -i3 & 5 \\ -i & \sqrt{3} & -i\sqrt{3} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \frac{1}{2\sqrt{2}} \begin{bmatrix} 1 \\ \sqrt{3} \\ \sqrt{3} \\ 1 \end{bmatrix} \quad (42)$$

have a solution. As a result, the ket $|X_{+3/2}\rangle$ belongs to neither $\text{ran}(\hat{Y}_{+3/2})$ nor $\text{ker}(\hat{Y}_{+3/2})$. In a bivalent semantics with no extra hypothesis, this implies that in the state described by $|X_{+3/2}\rangle$, the proposition $Y_{+3/2}$ is neither true nor false:

$$\llbracket Y_{+3/2} \rrbracket_v = \mathfrak{P}_{\in}(|X_{+3/2}\rangle, \text{ran}(\hat{Y}_{+3/2})) = 0/0 \quad . \quad (43)$$

4 Lemma of partial valuation

Suppose that the unit vector $|\Psi\rangle$ lies in the range of the projection operator \hat{P} , so that $\hat{P}|\Psi\rangle = |\Psi\rangle$. This equation corresponds to the matrix equation

$$\mathbf{P}\Psi = \Psi \quad , \quad (44)$$

which indicates that $\mathbf{P}^2 = \mathbf{P}$. Similarly, the bra equation $\langle\Psi| = \langle\Psi|\hat{P}$ can be written as

$$\Psi^\dagger = \Psi^\dagger \mathbf{P} \quad , \quad (45)$$

where the row vector $\Psi^\dagger \in \mathbf{Mat}_{1 \times n}(\mathbb{C})$, whose i^{th} column has the entry $\langle\Psi|e_i\rangle$, is the adjoint matrix of Ψ . From here, it follows that

$$\mathbf{P}\Psi\Psi^\dagger = \Psi\Psi^\dagger\mathbf{P} = \Psi\Psi^\dagger \quad , \quad (46)$$

which can be if

$$\mathbf{P} = \Psi\Psi^\dagger \quad . \quad (47)$$

As Ψ is a $n \times 1$ matrix and Ψ^\dagger is a $1 \times n$ matrix, the above factorization means (see, for example [13]) that the rank of the matrix \mathbf{P} is $\text{Rank}(\mathbf{P}) = 1$. This implies that for any $n \geq 2$, the number of linearly independent column vectors of the matrix \mathbf{P} is less than n , i.e.

$$\text{Rank}(\mathbf{P}) = \text{Span}(\mathbf{P}_1, \dots, \mathbf{P}_n) = m = 1 \quad . \quad (48)$$

By the rank-nullity theorem [14], the nullity of the matrix \mathbf{P} is $\text{Nullity}(\mathbf{P}) = n - 1$; so, for any $n \geq 2$, the number of linearly independent column vectors of the matrix $\mathbf{I} - \mathbf{P}$ is also less than n , namely,

$$\text{Nullity}(\mathbf{P}) = \text{Span}(\mathbf{I}_1 - \mathbf{P}_1, \dots, \mathbf{I}_n - \mathbf{P}_n) = k = n - 1 \quad . \quad (49)$$

Consequently, both linear systems, $\mathbf{R}\mathbf{X} = \mathbf{\Psi}$ and $\mathbf{K}\mathbf{X} = \mathbf{\Psi}$, are overdetermined.

Let's prove the lemma of partial valuation:

Lemma: *With no extra hypothesis, algebraic properties of the separable Hilbert space \mathcal{H} of finite dimension $n \geq 2$ cannot allow in valuations v which are total functions from \mathbb{P} to \mathbb{B}_2 .*

Suppose, the opposite is true, namely, valuations v that are total functions from \mathbb{P} to \mathbb{B}_2 are admitted. Then, for every nonzero vector $|\Psi\rangle$ in \mathcal{H} and an arbitrary projection operator \hat{P} such that $\hat{P} \neq \hat{0}$ and $\hat{P} \neq \hat{1}$, it holds true that $|\Psi\rangle$ belongs to either $\text{ran}(\hat{P})$ or $\text{ker}(\hat{P})$. Then again, this can happen only if at least one solution to either system of equations, $\mathbf{R}\mathbf{X} = \mathbf{\Psi}$ or $\mathbf{K}\mathbf{X} = \mathbf{\Psi}$, is guaranteed regardless of either \mathbf{R} and $\mathbf{\Psi}$, or \mathbf{K} and $\mathbf{\Psi}$, that is, in case either m or k is equal to n .

But, because both systems are overdetermined, this cannot be guaranteed: $\mathbf{R}\mathbf{X} = \mathbf{\Psi}$ and $\mathbf{K}\mathbf{X} = \mathbf{\Psi}$ will necessarily be unsolvable for some choice of values for the left-hand and right-hand sides of their equations, i.e., for some couples $(|\Psi\rangle, \text{ran}(\hat{P}))$.

Stipulating that $U_{\mathbf{R}}$ is the solution set for the linear system $\mathbf{R}\mathbf{X} = \mathbf{\Psi}$

$$U_{\mathbf{R}} = \{\mathbf{X} \in \text{Mat}_{m \times 1}(\mathbb{C}) : \mathbf{R}\mathbf{X} = \mathbf{\Psi}\} \quad , \quad (50)$$

and $U_{\mathbf{K}}$ is the solution set for the linear system $\mathbf{K}\mathbf{X} = \mathbf{\Psi}$

$$U_{\mathbf{K}} = \{\mathbf{X} \in \text{Mat}_{k \times 1}(\mathbb{C}) : \mathbf{K}\mathbf{X} = \mathbf{\Psi}\} \quad , \quad (51)$$

this inference can be written by the series of the sentences:

$$\begin{aligned} |\Psi\rangle \in \text{ran}(\hat{P}) &\iff U_{\mathbf{R}} \neq \emptyset \\ |\Psi\rangle \in \text{ker}(\hat{P}) &\iff U_{\mathbf{K}} \neq \emptyset \\ |\Psi\rangle \in \text{ran}(\hat{P}) \downarrow |\Psi\rangle \in \text{ker}(\hat{P}) &\iff U_{\mathbf{R}} \neq \emptyset \downarrow U_{\mathbf{K}} \neq \emptyset \end{aligned} \quad , \quad (52)$$

where the symbol \iff stands for *the logical biconditional* (which is true when its antecedent and consequent are either true or false at the same time).

According to the rank-nullity theorem, $U_{\mathbf{R}} \neq \emptyset \downarrow U_{\mathbf{K}} \neq \emptyset$ may be true (meaning that both solution sets $U_{\mathbf{R}}$ and $U_{\mathbf{K}}$ may be empty) for any $n \geq 2$. Thus, without additional hypotheses, the valuations v , which are total functions mapping experimental quantum propositions to elements of \mathbb{B}_2 , cannot

be admitted for any separable Hilbert space \mathcal{H} of finite dimension $n \geq 2$.

5 Closing remarks

Consider the statement (which can be called experimental quantum proposition) “A physical quantity of a quantum system has a certain value”. *Of what is this statement a primary bearer?*

The answer, that is ordinarily being given in the literature, is that this sentence is *a primary bearer of probability values*. This answer suggests that truth values of experimental quantum propositions can be regarded as degenerate probabilities. Consequently, rather than truth preservation, the logic of experimental quantum propositions is about probability preservation.

Given that the mathematical representative of any experimental quantum proposition is a closed linear subspace of a Hilbert space \mathcal{H} , this implies that quantum mechanics can be reduced to the theory of probability measures on closed linear subspaces of \mathcal{H} . According to Gleason’s theorem, if the Hilbert space has a finite dimension $n \geq 3$, this theory does not admit probability measures having only the values 0 and 1. The above result can be interpreted as evidence ruling out the possibility of hidden variables in quantum mechanics.

In contrast, in the present paper, another answer is offered asserting that an experimental quantum proposition is a primary bearer of truth values. Upon that, the paper demonstrates that without additional suppositions, algebraic properties of separable Hilbert spaces \mathcal{H} of finite dimension $n \geq 2$ do not allow in truth assignments v which are dispersion-free, i.e., which have being total functions from the set of atomic (i.e., elementary) experimental quantum propositions to the set of two objects, true and false (or 1 and 0, correspondingly).

As long as a probability is interpreted as a measure of the (un)certainly in the assignment of truth values to an atomic proposition, the fact that the valuations v cannot be dispersion-free indicates that probabilities cannot be only $\{0, 1\}$ -valued. In this way, one can say that gaps in truth assignments entail the emergence of probabilistic semantics for the logic of experimental quantum propositions.

It follows – independently of Gleason’s theorem – that non-contextual hidden variables identified as globally defined $\{0, 1\}$ -valued observables (resembling experimental propositions) must be excluded from the interpretation of quantum mechanics that is based on the Hilbert space formalism.

References

- [1] Alexander Wilce. Quantum Logic and Probability Theory. In Edward N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, 2017.
- [2] Enrico Beltrametti and Gianni Cassinelli. The Logic of Quantum Mechanics. In Gian-Carlo Rota, editor, *Encyclopedia of Mathematics and its Applications*, volume 15. Cambridge University Press, 1984.

- [3] M.L. Dalla Chiara and R. Giuntini. Quantum Logic. arXiv:quant-ph/0101028, Jan 2001.
- [4] Rafael D. Sorkin. An exercise in “anhomomorphic logic”. *J. Phys. Conf. Ser.*, 67(012018), 2007.
- [5] Alan Hájek. Probability, Logic, and Probability Logic. In Lou Goble, editor, *The Blackwell Guide to Philosophical Logic*, pages 362–384. John Wiley and Sons Ltd., 2001.
- [6] Jon Williamson. Probability Logic. In D. M. Gabbay, R. H. Johnson, H. J. Ohlbach, and J. Woods, editors, *Handbook of the Logic of Argument and Inference: The Turn Towards the Practical*, pages 397–424. Elsevier, Amsterdam, 2002.
- [7] G. Birkhoff and J. von Neumann. The logic of quantum mechanics. *Annals of Mathematics*, 37:823–843, 1936.
- [8] A. Gleason. Measures on the closed subspaces of a Hilbert space. *J. Math. Mech.*, 6(6):885–893, 1957.
- [9] J.-Y. Bèziau. Bivalence, Excluded Middle and Non Contradiction. In L. Behounek, editor, *The Logica Yearbook 2003*, pages 73–84. Academy of Sciences, Prague, 2003.
- [10] G. Kalmbach. *Orthomodular Lattices*. Academic Press, London, 1983.
- [11] Stephen Blamey. Partial logic. In Dov M. Gabbay and Franz Guenther, editors, *Handbook of Philosophical Logic, Volume III: Alternatives in Classical Logic*, pages 1–70. Springer Science and Business Media Dordrecht, 1986.
- [12] Tore Langholm. *Partiality, Truth and Persistence. CSLI (Center for the Study of Language and Information) Lecture Notes*. CSLI Publications, Stanford University, 1988.
- [13] L. Mirsky. *An Introduction to Linear Algebra*. Dover Publications, 2011.
- [14] Stephen H. Friedberg, Arnold J. Insel, and Lawrence E. Spence. *Linear Algebra, 5th Edition*. Pearson, 2019.