

# Energy Stablensess for Schrödinger Operators with Time-Dependent Potentials

by **Masaki Kawamoto**

Department of Engineering for Production, Graduate School of Science and Engineering, Ehime University, 3 Bunkyo-cho Matsuyama, Ehime, 790-8577. Japan.

Email: kawamoto.masaki.zs@ehime-u.ac.jp

**Abstract.** In this paper, we prove the energy stable property for time-dependent (generalized) Schrödinger operators by using Hardy inequality. Such property acts very important roles in quantum scattering theory and nonlinear problem. As an application, we prove Sobolev type inequality.

**Keywords:** Time-dependent potentials; Time-dependent Schrödinger equations; Energy Stablensess; Sobolev inequality;

## 1 Introduction

We consider the (generalized) Schrödinger equations with time-dependent potential;

$$\begin{aligned}i\partial_t\phi(t, x) &= ((-\Delta)^\theta/(2m) + V(t))\phi(t, x), \\ \phi(s, x) &= \phi(s) \in H^1(\mathbf{R}^n),\end{aligned}$$

where  $\theta \geq 1/2$ ,  $x = (x_1, x_2, \dots, x_n) \in \mathbf{R}^n$ ,  $\Delta = \partial_1^2 + \partial_2^2 + \dots + \partial_n^2$  is the Laplacian,  $m > 0$ ,  $s \in \mathbf{R}$  is a fixed constant, and  $V(t)$  is a potential; a multiplication operator of real valued function  $V(t, x)$  defined later. We use notations  $p = -i\nabla = -i(\partial_1, \partial_2, \dots, \partial_n)$  i.e.,  $-\Delta = p^2$ . Let  $H(t) = p^{2\theta}/(2m) + V(t)$  and call *energy* in  $t$ . We say a family of unitary operators  $\{U(t, s)\}_{(t,s) \in \mathbf{R}^2}$  a propagator for  $H(t)$  if each component satisfies

$$\begin{aligned}i\frac{\partial}{\partial t}U(t, s) &= H(t)U(t, s), & i\frac{\partial}{\partial s}U(t, s) &= -U(t, s)H(s), \\ U(t, \tau)U(\tau, s) &= U(t, s), & U(s, s) &= \text{Id}_{L^2(\mathbf{R}^n)}.\end{aligned}$$

In this paper, we state the following assumptions on the potential;

**Assumption 1.** Let  $V(t, x) : \mathbf{R} \times \mathbf{R}^n \rightarrow \mathbf{R}$  satisfies  $V \in C^1(\mathbf{R}; C^1(\mathbf{R}^n))$ . Suppose that for all  $u \in H^\theta(\mathbf{R}^n)$ , there exist positive constants  $C_H > 0$  and  $C > 0$  such that

$$(p^{2\theta}u, u)_{L^2(\mathbf{R}^n)} \geq C_H(V(t)u, u)_{L^2(\mathbf{R}^n)} \quad (1)$$

and

$$\int_s^\infty \left\| \frac{\partial}{\partial t} V(t, \cdot) \right\|_{L^\infty(\mathbf{R}^n)} dt \leq C. \quad (2)$$

holds. Moreover, assume that the propagator  $U(t, s)$  uniquely exists and that for all  $t \in \mathbf{R}$ ,  $U(t, s)$  satisfies

$$U(t, s)H^\theta(\mathbf{R}^n) \subset H^\theta(\mathbf{R}^n).$$

In order to obtain energy stableness, we further state the following two conditions;

**Assumption 2.** For all  $u \in H^\theta(\mathbf{R}^n)$ , potential  $V(t)$  satisfies the following inequality

$$\left( \left( p^{2\theta}/4m + V(s) - \int_s^t \left\| \frac{\partial}{\partial \tau} V(\tau, \cdot) \right\|_{L^\infty(\mathbf{R}^n)} d\tau \right) u, u \right)_{L^2(\mathbf{R}^n)} \geq 0.$$

**Assumption 3.** Energy  $H(t)$  is strictly positive, that is, for all  $t \in \mathbf{R}$ ,  $x \in \mathbf{R}^n$  and  $u \in H^\theta(\mathbf{R}^n)$ , there exists  $\delta > 0$  such that

$$(H(t)u, u)_{L^2(\mathbf{R}^n)} = \left( \left( p^{2\theta}/(2m) + V(t, x) \right) u, u \right)_{L^2(\mathbf{R}^n)} \geq \delta \left( p^{2\theta} u, u \right)_{L^2(\mathbf{R}^n)} \quad (3)$$

**Remark 4.** If the potential is independent of the time, the assumption 1 admits Coulomb type potential, that is,  $V = C_{H,n}|x|^{-2\theta_0}$ , where  $0 < \theta_0 < \theta$  and  $C_{H,n} > 0$  depends only on the dimension  $n \in \mathbf{N}$ , see e.g. Secchi-Smets-Willem [3] since we can obtain the unique existence of the propagator  $U(t, s) = e^{-i(t-s)H}$  with  $H = p^\theta/(2m) + V$  by using the Stone's theorem and self-adjointness of  $H$ . On the other hand, in the case where potential depends on time, it seems difficult to include singular potentials such as Coulomb type potential in Assumption 1 because of (2). For  $\theta = 1$  the unique existence of the propagator is guaranteed by Yajima [4] even if the potential depends on time and has singularities. If  $V$  is bounded, then we can easily prove the unique existence of  $U(t, s)$ .

**Remark 5.** As an example, we consider the potential written as the form

$$V(t, x) = V_0(x + c(t)) \in \mathbf{R},$$

where  $V_0 \in \mathcal{B}^1(\mathbf{R})$ , and  $c = (c_1, \dots, c_n)$  satisfies that for all  $j \in \{1, \dots, n\}$ ,  $c_j \in \mathcal{B}^1(\mathbf{R})$  and integrable condition

$$\int_s^\infty |c'(t)| dt < \infty.$$

Additionally if  $V(t)$  satisfies

$$\sup_{x \in \mathbf{R}^n} \left| |x|^{2\theta_0} V(t, x) \right| \leq C_H$$

for some constant  $C > 0$  and some  $\theta_0 > \theta$  then  $V$  satisfies Assumption 1. For  $\theta = 1$ , such a potential appears for Schrödinger equations with time-decaying electric fields, see e.g., Adachi-Fujiwara-Ishida [1]. What we emphasize here is we do not need to assume that  $|c(t)| \rightarrow 0$  as  $|t| \rightarrow \infty$ .

Under the assumption 1, we have the following Theorem;

**Theorem 6** (Stableness for Laplacian). *Let  $\phi(s) \in H^\theta(\mathbf{R}^n)$  satisfies that for some  $0 < a < R$ ,  $\text{supp}(\hat{\phi}(s)) \subset \{\xi \in \mathbf{R}^n \mid a \leq |\xi| \leq R\}$ , where  $\hat{\cdot}$  stands for the Fourier transform. If we assume Assumption 1 and 2. Then for any  $t \in \mathbf{R}$  there exists a  $t$ -independent constant  $\tilde{a} > 0$  such that*

$$\left( p^{2\theta} U(t, s) \phi(s), U(t, s) \phi(s) \right)_{L^2(\mathbf{R}^n)} \geq \tilde{a} \|\phi(s)\|_{L^2(\mathbf{R}^n)}^2 \quad (4)$$

holds. On the other hand, if we assume Assumption 1 and 3. Then for any  $t \in \mathbf{R}$  there exists a  $t$ -independent constant  $\tilde{R} > 0$  such that

$$\left( p^{2\theta} U(t, s) \phi(s), U(t, s) \phi(s) \right)_{L^2(\mathbf{R}^n)} \leq \tilde{R} \|\phi(s)\|_{L^2(\mathbf{R}^n)}^2 \quad (5)$$

holds. Hence if we assume Assumption 1, 2 and 3, then for any  $t \in \mathbf{R}$  there exist  $t$ -independent constants  $\tilde{a} > 0$  and  $\tilde{R} > 0$  such that

$$\tilde{a} \|\phi(s)\|_{L^2(\mathbf{R}^n)}^2 \leq \left( p^{2\theta} U(t, s) \phi(s), U(t, s) \phi(s) \right)_{L^2(\mathbf{R}^n)} \leq \tilde{R} \|\phi(s)\|_{L^2(\mathbf{R}^n)}^2$$

holds.

As the corollary, we can obtain the following energy stableness property;

**Theorem 7** (Energy stableness). *Let  $\phi(s) \in H^\theta(\mathbf{R}^n)$  satisfies*

$$a_1 \|\phi(s)\|_{L^2(\mathbf{R}^n)}^2 \leq (H(s)\phi(s), \phi(s))_{L^2(\mathbf{R}^n)} \leq R_1 \|\phi(s)\|_{L^2(\mathbf{R}^n)}^2$$

for some constants  $0 < a_1 < R_1$ . Assume Assumption 1, 2 and 3. Then for any  $t \in \mathbf{R}$ , there exist  $t$ -independent constants  $0 < \tilde{a}_1 < \tilde{R}_1$  such that

$$\tilde{a}_1 \|\phi(s)\|_{L^2(\mathbf{R}^n)}^2 \leq (H(t)U(t, s)\phi(s), U(t, s)\phi(s))_{L^2(\mathbf{R}^n)} \leq \tilde{R}_1 \|\phi(s)\|_{L^2(\mathbf{R}^n)}^2.$$

**Corollary 8** (Uniformly stableness in  $H^\theta(\mathbf{R}^n)$ ). *Suppose  $\phi(s) \in H^\theta(\mathbf{R}^n)$ . If we assume Assumption 1 and 2. Then for any  $t \in \mathbf{R}$  there exists a  $t$ -independent constant  $c_m > 0$  such that*

$$\|U(t, s)\phi(s)\|_{H^\theta(\mathbf{R}^n)}^2 \geq c_m \|\phi(s)\|_{H^\theta(\mathbf{R}^n)}^2 \quad (6)$$

holds. If we assume Assumption 1 and 3. Then for any  $t \in \mathbf{R}$  there exists a  $t$ -independent constant  $c_M > 0$  such that

$$\|U(t, s)\phi(s)\|_{H^\theta(\mathbf{R}^n)}^2 \leq c_M \|\phi(s)\|_{H^\theta(\mathbf{R}^n)}^2 \quad (7)$$

holds

As an application of this theorem, we shall introduce the Sobolev type inequality;

**Corollary 9.** Under the assumption 1 and 3, for all  $0 \leq \gamma \leq \theta$  and  $\phi(s) \in H^\gamma(\mathbf{R}^n)$ , there exists a constant  $c > 0$  such that

$$\|U(t, s)\phi(s)\|_{L^p(\mathbf{R}^n)} \leq c \|\phi(s)\|_{H^\gamma(\mathbf{R}^n)}$$

holds, where  $p$  satisfies

$$\frac{1}{p} + \frac{\gamma}{n} = \frac{1}{2}.$$

and  $c$  does not depend on  $t$ .

For  $\theta = 1$ , energy stableness property can be proven for the case where  $\|V(t, \cdot)\|_{L^\infty(\mathbf{R}^n)}$  decays in  $t$  or  $\|V(t, \cdot)\|_{L^\infty(\mathbf{R}^n)}$  is sufficiently small or  $V(t, \cdot)$  is periodic in time, respectively. However, in the assumption 1, we do not assume these conditions, and the energy stableness under Assumption 1 – 3 has not been seen yet, as far as we know. For such potentials, linear scattering theory (in particular asymptotic completeness), Strchartz estimates (see, e.g., Naibo-Stefanov [2]) and global well-posedness in  $L^\infty$  for NLS, also have not been proven yet, as far as we know. The energy stableness and Sobolev inequality may be applicable to such studies.

## 2 Proof of Theorem

In this section, we shall prove Theorem 6, Corollary 8 and 9. For simplicity, we assume that  $\|\cdot\|_2$  denotes  $\|\cdot\|_{L^2(\mathbf{R}^n)}$  and  $(\cdot, \cdot)$  denotes  $(\cdot, \cdot)_{L^2(\mathbf{R}^n)}$ .

Define  $F(t)$  as

$$F(t) = \| |p|^\theta u(t) \|_2^2 = (p^{2\theta} u(t), u(t)),$$

where  $u(t) = U(t, s)\phi(s)$ . Then, by the assumption (1) (Hardy type inequality), there exists  $C_H > 0$  such that

$$F(t) \geq C_H (V(t)u(t), u(t)) =: G(t).$$

Here, by the simple calculation, we have

$$\frac{d}{dt} F(t) = \left( i[V(t), p^{2\theta}]u(t), u(t) \right),$$

where  $[\cdot, \cdot]$  stands for the commutator of operators. On the other hand,

$$\begin{aligned}\frac{d}{dt}G(t) &= C_H \left( i[p^{2\theta}/2m, V(t)]u(t), u(t) \right) + C_H(V'(t)u(t), u(t)) \\ &= -(C_H/2m)\frac{d}{dt}F(t) + C_H(V'(t)u(t), u(t))\end{aligned}$$

Hence we have

$$G(t) = C_H \int_s^t (V'(\tau)u(\tau), u(\tau))d\tau - \frac{C_H}{2m}(F(t) - F(s)) + G(s) \quad (8)$$

Condition  $F(t) \geq G(t)$  implies

$$\begin{aligned}\frac{2m + C_H}{2m}F(t) &\geq \frac{C_H}{2m}F(s) + G(s) + C_H \int_s^t (V'(\tau)u(\tau), u(\tau)) d\tau \\ &\geq \frac{C_H}{4m}F(s) + C_H \left( \left( \frac{1}{4m}p^{2\theta} + V(s) - \int_s^t \|V'(\tau, \cdot)\|_{L^\infty(\mathbf{R}^n)} d\tau \right) \phi(s), \phi(s) \right).\end{aligned}$$

If we assume Assumption 2, we notice that there exists a positive constant  $C > 0$  such that

$$F(t) \geq CF(s),$$

which implies (4) holds. On the other hand, if we assume Assumption 3, then we have

$$\begin{aligned}\frac{C_H}{2m}F(t) &= \frac{C_H}{2m\delta} \left( \delta p^{2\theta} u(t), u(t) \right) \leq \frac{C_H}{2m\delta} \left( \left( \frac{1}{2m}p^{2\theta} + V(t) \right) u(t), u(t) \right) \\ &= \frac{1}{2m\delta} \left( \frac{C_H}{2m}F(t) + G(t) \right) \\ &\leq \frac{1}{2m\delta} \left( G(s) + \frac{C_H}{2m}F(s) + C_H \int_s^t \|V'(\tau, \cdot)\|_{L^\infty(\mathbf{R}^n)} d\tau \|\phi(s)\|_2^2 \right),\end{aligned} \quad (9)$$

which implies (5) holds. Inequalities (6) and (7) also hold by (9) and (1). Theorem 7 can be immediately proven by using following inequalities

$$\begin{aligned}(H(t)u(t), u(t)) &\leq \left( \left( p^{2\theta}/(2m) + V(t) \right) u(t), u(t) \right) \\ &\leq (1/2m + 1/C_H)(p^{2\theta}u(t), u(t))\end{aligned}$$

and (3).

Finally, we prove Corollary 9. By interpolating the followings

$$\left\| |p|^0 U(t, s) \langle p \rangle^{-0} \right\|_{\mathcal{B}(L^2(\mathbf{R}^n))} \leq C_0$$

and

$$\left\| |p|^\theta U(t, s) \langle p \rangle^{-\theta} \right\|_{\mathcal{B}(L^2(\mathbf{R}^n))} \leq C_1,$$

we get for some  $0 \leq \gamma_0 \leq \theta$  and  $(t, s)$ -independent constant  $C_{\gamma_0} > 0$ ,

$$\left\| |p|^{\gamma_0} U(t, s) \langle p \rangle^{-\gamma_0} \right\|_{\mathcal{B}(L^2(\mathbf{R}^n))} \leq C_{\gamma_0},$$

where  $\langle \cdot \rangle = (1 + \cdot^2)^{1/2}$ ,  $C_0 > 0$  and  $C_1 > 0$  are  $(t, s)$ -independent constants and  $\mathcal{B}(L^2(\mathbf{R}^n))$  stands for the operator norm on  $L^2(\mathbf{R}^n)$ . That provides for a pair  $(p, \gamma)$  with  $1/p + \gamma/n = 1/2$ ,

$$\begin{aligned} \|U(t, s)\phi(s)\|_{L^p(\mathbf{R}^n)} &\leq C \| |p|^\gamma U(t, s)\phi(s) \|_2 \\ &\leq C \| |p|^\gamma U(t, s) \langle p \rangle^{-\gamma} \|_{\mathcal{B}(L^2(\mathbf{R}^n))} \| \langle p \rangle^\gamma \phi(s) \|_2, \end{aligned}$$

and which is the desired result, where we use the Gagliardo-Nirenberg inequality.

## References

- [1] Adachi, T., Fujiwara, Y., Ishida, A.: On multidimensional inverse scattering in time-dependent electric fields, *Inverse Problems*, **29**, 085012 (2013).
- [2] Naibo, V., Stefanov, A.: On some Schrödinger and wave equations with time-dependent potentials, *Math., Ann.*, **334**, 325–338 (2006).
- [3] Secchi, S., Smets, D., Willem, M.: Remarks on a Hardy-Sobolev inequality, *Comptes Rendus Math.*, **336**, 811–815 (2003).
- [4] Yajima, K.: Existence of solutions for Schrödinger evolution equations, *Comm. Math. Phys.*, **110**, 415–426 (1987)