

RANDOM PLANE PARTITIONS AND CORNER DISTRIBUTIONS

DAMIR YELIUSSIZOV

ABSTRACT. We explore some probabilistic applications arising in connections with K -theoretic symmetric functions. For instance, we determine certain corner distributions of random lozenge tilings and plane partitions. We also introduce some distributions that are naturally related to the corner growth model. Our main tools are dual symmetric Grothendieck polynomials and normalized Schur functions.

1. INTRODUCTION

Combinatorics arising in connection with K -theoretic Schubert calculus is quite rich. Accompanied by certain families of symmetric functions, it usually presents some inhomogeneous deformations of objects beyond classical Schur (or Schubert) case. While the subject is intensively studied from combinatorial, algebraic and geometric aspects, see [Len00, Buc02, Buc05, Vak06, LP07, IN13, Yel17] and many references therein, much less is known about probabilistic connections (unlike interactions between probability and representation theory). Some work in this direction was done in [TY11, MS14] and related problems were addressed in [Yel20].¹

In this paper, we give several probabilistic applications obtained with tools from combinatorial K -theory. We mostly focus on one deformation of Schur functions, the *dual Grothendieck polynomials*, whose associated combinatorics is fairly neat. In particular, we show that these functions are naturally related to the *corner growth model* [Joh00, Joh01, Sep09, Rom15] (which can also be viewed as a totally asymmetric simple exclusion process or a directed last passage percolation), see subsec. 1.6 and Sec. 8.

We are now going to discuss our results across few related models.

1.1. \mathbb{N} -matrices with bounded last passage time. A lattice path Π with vertices indexed by \mathbb{N}^2 (where $\mathbb{N} = \{0, 1, 2, \dots\}$) is called a *monotone path* if it uses only steps of the form $(i, j) \rightarrow (i+1, j), (i, j+1)$. An \mathbb{N} -matrix is a matrix of nonnegative integers with only finitely many nonzero entries. Let a, b, c be positive integers. Given an \mathbb{N} -matrix $D = (d_{ij})$ with b rows and c columns, the *last passage time* $G(b, c)$ is defined as

$$G(b, c) = \max_{\Pi} \sum_{(i,j) \in \Pi} d_{ij},$$

where the maximum is taken over monotone paths Π from $(1, 1)$ to (b, c) . Let $\text{BM}(a, b, c)$ be the set of $b \times c$ \mathbb{N} -matrices whose last passage time is *bounded* by a , i.e.

$$\text{BM}(a, b, c) := \left\{ D = (d_{ij})_{i,j=1}^{b,c} : G(b, c) \leq a, \quad d_{ij} \in \mathbb{N} \right\}$$

¹See also Remark 8.6 on recent works.

This set is in fact equinumerous with *boxed plane partitions* (see Theorem 2.2).

Consider the uniform probability measure on $\text{BM}(a, b, c)$. For a random matrix $D \in \text{BM}(a, b, c)$, define the *column marginals*

$$C_\ell = \sum_{i=1}^b d_{i\ell}$$

i.e. as the sum of entries in ℓ -th column of D where $\ell \in [1, c]$. It is natural to ask what are distributions of C_ℓ . We determine these distributions. The results about C_ℓ are presented via lozenge tilings (see Theorem 1.3) below. We show that the random variables (C_ℓ) are exchangeable (see Theorem 1.2; in particular, C_i and C_j have the same distribution for all $i, j \in [1, c]$) and obtain limiting (joint) distributions with different asymptotic regimes for the parameters a, b, c . In some regimes, the variables C_ℓ become *asymptotically independent*. By symmetry, we also get similar results for *row marginals*.

1.2. Boxed plane partitions. A *plane partition* is an \mathbb{N} -matrix $\pi = (\pi_{ij})_{i,j \geq 1}$ satisfying

$$\pi_{ij} \geq \pi_{i+1,j}, \quad \pi_{ij} \geq \pi_{i,j+1}, \quad i, j \geq 1.$$

By default we ignore zeros of π . We denote by $\text{sh}(\pi) := \{(i, j) : \pi_{ij} > 0\}$ the *shape* of π .

Let $\text{PP}(a, b, c)$ be the set of *boxed plane partitions* that fit inside the box $a \times b \times c$, i.e. the first row of the shape is at most a , the first column is at most b , and the first entry is at most c . We show that $|\text{PP}(a, b, c)| = |\text{BM}(a, b, c)|$ (Theorem 2.2).

Consider the uniform probability measure on the set $\text{PP}(a, b, c)$. Recall that classical MacMahon's theorem on boxed plane partitions gives the explicit product formula




$$Z_{abc} := |\text{PP}(a, b, c)| = \prod_{i=1}^a \prod_{j=1}^b \prod_{k=1}^c \frac{i+j+k-1}{i+j+k-2}$$

Note that we also have


$$Z_{abc} = s_{(a^b)}(1^{b+c}),$$

where s_λ is the *Schur polynomial* and $1^n := (1, \dots, 1)$ repeated n times. (See e.g. [Sta99, Ch. 7].)

For a uniformly random plane partition $\pi \in \text{PP}(a, b, c)$, let X_ℓ be the number of columns of π that contain entry $\ell \in [1, c]$. In fact, the variables X_ℓ become image of the marginals C_ℓ described previously under the bijection which we describe in Sec. 2. Similarly, (X_ℓ) are exchangeable, and X_i and X_j have the same distribution for all $i, j \in [1, c]$. Theorem 1.3 below (stated in terms of lozenge tilings) gives limiting distributions in several asymptotic regimes. By symmetry, we also get similar results for the random variables Y_ℓ , the number of rows containing ℓ in a random plane partition. Hence the results also imply some bounds (deviation) for the *area* containing ℓ .

1.3. Lozenge tilings of a hexagon. A *lozenge tiling* is a tiling of a planar domain with three types of *lozenges*    which we refer to as *left*, *top* and *right* tiles.

Let $\text{LT}(a, b, c)$ be the set of lozenge tilings of a *hexagon* with sides (a, b, c, a, b, c) as in Fig. 1. For a lozenge tiling $T \in \text{LT}(a, b, c)$, define:

- *left corners* (or simply *corners*) of T , that are local tile configurations of the form 
- for each corner α , its *height* $h(\alpha) \in [1, c]$ is the z -coordinate of the top tile of α , when T is viewed in \mathbb{R}^3 as a pile of cubes (boxed plane partition), see Fig. 1.




FIGURE 1. A plane partition and the corresponding lozenge tiling of a hexagon with sides $a = 5, b = 4, c = 3$. It has 11 left corners 5 of which have height 2.

Consider the uniform probability measure on the set $\text{LT}(a, b, c)$. We clearly have $|\text{LT}(a, b, c)| = |\text{PP}(a, b, c)|$, where the bijection is given by writing the height of each top tile. The random variables C_ℓ and X_ℓ defined for previous models translate into the following random variables for lozenge tilings.

Definition 1.1 (Corner distributions). Let Γ_ℓ be the number of left corners of height ℓ .

We show the following symmetry.

Theorem 1.2 (Exchangeability of levels, cf. Corollary 5.2). *The random variables (Γ_ℓ) are exchangeable. In particular, Γ_i and Γ_j have the same distribution for all $i, j \in [1, c]$.*

We find limits of corner distributions Γ in several asymptotic regimes.

Theorem 1.3. *We have convergence in distribution in the following regimes:*

(i) *Poisson: for $a, b, c \rightarrow \infty$ with $ab/c \rightarrow t > 0$ we have*

$$\Gamma_\ell \rightarrow \text{Poisson}(t)$$

(ii) *Negative binomial: for b fixed and $a, c \rightarrow \infty$ with $a/(a+c) \rightarrow q \in (0, 1)$ we have*

$$\Gamma_\ell \rightarrow \text{NB}(b, q)$$

(iii) *Gaussian: for $a, b, c \rightarrow \infty$ with $a/(b+c) \rightarrow u > 0$ and $b/(b+c) \rightarrow q \in (0, 1)$ we have*

$$\frac{\Gamma_\ell - \mu N}{\sigma \sqrt{N}} \rightarrow \mathcal{N}(0, 1),$$

where $N = b + c$, $\mu = uq$, and $\sigma = \sqrt{u(1+u)q(1-q)}$.

Moreover, the joint distribution of any collection of random variables Γ_ℓ weakly converges to the distribution of independent random variables given in corresponding regimes (i), (ii) or (iii).

Consider the Gaussian regime, say for $a = b = c = n \rightarrow \infty$. It is known that (with a probability going to 1) random lozenge tilings have the *arctic circle* phenomenon: outside of the inscribed circle of the hexagon, there are *frozen* regions, and inside there is a *liquid* region [CLP98, CKP01, Joh02, Ken09], see Fig. 2. Note that left corners on the topmost level at height c form a profile for the upper frozen region. Then the number of such corners is about $n/2$ with fluctuations of order \sqrt{n} , which is compatible with known edge behavior. However what may seem surprising is the behavior across height levels. There is a kind of invariance

following combinatorial presentation

$$g_\lambda(x_1, x_2, \dots) = \sum_{T \in \text{PP}(\lambda)} \prod_{i \geq 1} x_i^{c_i(T)},$$

where the sum runs over plane partitions T of shape λ and $c_i(T)$ is the number of columns of T containing i . These polynomials were explicitly introduced and studied in [LP07] (and earlier implicitly in [Len00, Buc02]) and they are related to K -homology of Grassmannians. More properties of these functions can be found in [Yel17, Yel19]. The polynomials g_λ can be considered as an inhomogeneous K -theoretic deformation of Schur polynomials. By definition, it is easy to see that the top degree homogeneous component of g_λ is the Schur polynomial s_λ , i.e. $g_\lambda = s_\lambda + \text{lower degree terms}$.

1.5. g -measure on plane partitions. Let $q_1, \dots, q_c \in (0, 1)$. For a plane partition π define the *descent set*

$$\text{Des}(\pi) := \{(i, j) : \pi_{ij} > \pi_{i+1, j}\}$$

and let $\text{des}(\pi) = |\text{Des}(\pi)|$.

Consider the probability distribution $P_{g,b,c}$, which we call *g -measure* or *g -distribution*, on the (infinite) set $\text{PP}(\infty, b, c)$ of plane partitions with at most b rows and maximal entry at most c , defined as follows:

$$P_{g,b,c}(\pi) := \frac{1}{Z_{b,c}} \prod_{(i,j) \in \text{Des}(\pi)} q_{\pi_{ij}},$$

where the associated partition function (normalization) can be computed as (see Corollary 7.2)

$$Z_{b,c} = \prod_{i=1}^c (1 - q_i)^{-b}.$$

Proposition 1.5 (cf. Proposition 7.1). *Let $\pi \in \text{PP}(\infty, b, c)$ and $\lambda = (\lambda_1 \geq \dots \geq \lambda_b \geq 0)$. We have*

$$P_{g,b,c}(\text{sh}(\pi) = \lambda) = \frac{1}{Z_{b,c}} g_\lambda(q_1, \dots, q_c).$$

By this formula it is convenient to view $P_{g,b,c}$ as a distribution on integer partitions λ with at most b parts. We write $P_{g,b,c}(\lambda)$ to mean this distribution. This measure is somewhat similar to the *Schur measure* [Oko01, OR03] as it relies on the Cauchy-type identity for dual Grothendieck polynomials (which is a special case of a more general Cauchy identity involving two families of symmetric Grothendieck functions [Yel19]).

Theorem 1.6 (cf. Theorem 7.3). *For $\pi \in \text{PP}(\infty, b, c)$, let $\lambda = (\lambda_1 \geq \dots \geq \lambda_b \geq 0)$ be the shape of π . The distribution of the first part λ_1 can be expressed as*

$$P_{g,b,c}(\lambda_1 \leq a) = \frac{1}{Z_{b,c}} s_{(a^b)}(1^b, q_1, \dots, q_c).$$

In particular, this formula implies Toeplitz (and Fredholm) determinantal expressions for this distribution, see Sec. 7.

1.6. Corner growth model. Let $W = (w_{ij})_{i,j \geq 1}$ be a random matrix with i.i.d. entries w_{ij} that have geometric distribution with parameter $q \in (0, 1)$, i.e.

$$P(w_{ij} = k) = (1 - q)q^k, \quad k \in \mathbb{N}.$$

Consider the last passage times for all $m, n \geq 1$

$$G(m, n) = \max_{\Pi} \sum_{(i,j) \in \Pi} w_{ij},$$

where the maximum is over monotone paths Π from $(1, 1)$ to (m, n) .

The *corner growth model* can be viewed as evolution of a random Young diagram $Y(t)$ given at time t by the region

$$Y(t) = \{(i, j) \in \mathbb{N}^2 : G(i, j) \leq t\}.$$

The matrix $G = (G(i, j))_{i,j \geq 1}$ is called the *performance table*. This model was studied in [Joh00, Bar01], see also [Joh01, Sep09, Rom15] and references therein for more on the topic. In Sec. 8 we review and use some known results.

The main relationship between this stochastic process and the g -distribution defined above is the following.

Theorem 1.7 (cf. Theorem 8.1). *Let $\lambda = (\lambda_1, \dots, \lambda_b)$ be a partition with at most b parts. We have*

$$P(G(b, c) = \lambda_1, \dots, G(1, c) = \lambda_b) = P_{g,b,c}(\lambda),$$

where $q_i = q$ for all $i \in [1, c]$ in the g -measure.

This formula is useful for both objects that it relates, i.e. as a formula for row distributions of the performance table as well as a source for new properties of dual Grothendieck polynomials.

1.7. On methods and tools. Random plane partitions were studied intensively and there is large literature on the subject, see [CLP98, Joh01, OR03, JN06, Ken09, BP14, Pet14, GP15] and many references therein. Yet, exploiting tools from combinatorial K -theory turns out to be beneficial for establishing new properties. Let us summarize the key ingredients crucial to our results. One of them is the bijection between plane partitions and \mathbb{N} -matrices, described in the following section. This map is different from the well-known Robinson-Schensted-Knuth (RSK) correspondence; its description is somewhat simpler but has analogous properties. Another key tool is in dual Grothendieck polynomials whose combinatorics is tied to plane partitions. As we see, these functions are useful for uncovering new properties and symmetries for some natural statistics on plane partitions, which seem to be difficult to obtain otherwise. The determinantal formulas in certain cases allow us to make a transition to Schur polynomials, which is non-trivial at the combinatorial level. Combined with this interesting coincidence (see Lemma 3.4), we then use known asymptotics for normalized Schur polynomials from [BP14, GP15].

2. A BIJECTION BETWEEN PLANE PARTITIONS AND \mathbb{N} -MATRICES

Given a plane partition π , define the *descent level sets*

$$(1) \quad D_{i\ell} := \{j : \pi_{ij} = \ell > \pi_{i+1,j}\},$$




FIGURE 3. A geometric view of the map Φ . The dots \bullet correspond to left corners.

i.e. $D_{i\ell}$ is the set of column indices of the entry ℓ in i th row of π that are strictly larger than the entry below.

Before proceeding, let us mention some important properties of these sets:


- The elements in $D_{i\ell}$ form a consecutive segment.
- For $(i, \ell) \neq (i_1, \ell_1)$ such that $i \geq i_1, \ell \geq \ell_1$, the sets $D_{i\ell}$ and $D_{i_1\ell_1}$ are *disjoint* (more precisely, every element in $D_{i_1\ell_1}$ is larger than any element of $D_{i\ell}$).

Let now $d_{i\ell} := |D_{i\ell}|$ and $D := (d_{i\ell})_{i,\ell \geq 1}$. Define the map $\Phi : \{\text{plane partitions}\} \rightarrow \{\mathbb{N}\text{-matrices}\}$ by setting

$$(2) \quad \Phi(\pi) = D.$$

Example 2.1. Consider $\Phi : \pi \mapsto D$ as follows

$$\Phi : \begin{array}{|c|c|c|} \hline 4 & 4 & 2 \\ \hline 4 & 2 & 1 \\ \hline 2 & 2 & \\ \hline \end{array} \mapsto \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 2 & 0 & 0 \end{pmatrix}$$

It is convenient to view the map Φ geometrically: present π as a pile of cubes in \mathbb{R}^3 ; then mark corners  on the surface of π with \bullet as in Fig. 3. Then $d_{i\ell}$ is the number of such marks of height $z = \ell$ and $x = i$. Alternatively, we may view this as a projection of these marks on the $y = 0$ plane, see Fig. 3.

Theorem 2.2. *The map Φ defines a bijection between the set $\text{PP}(a, b, c)$ of boxed plane partitions and $\text{BM}(a, b, c)$ of \mathbb{N} -matrices with bounded last passage time.*


Proof. Suppose $\pi \in \text{PP}(a, b, c)$. Let us show that $D = \Phi(\pi) \in \text{BM}(a, b, c)$. By construction, it is clear that D has at most b rows and c columns. Take any monotone path $(1, 1) = (i_1, \ell_1) \rightarrow \dots \rightarrow (i_k, \ell_k) \leq (b, c)$ in D . Then the descent level sets $D_{i_s \ell_s}$ for $s = 1, \dots, k$ are pairwise disjoint by the property mentioned above. Hence

$$\sum_{s=1}^k d_{i_s \ell_s} = \sum_{s=1}^k |\{j : \pi_{i_s j} = \ell_s > \pi_{i_{s+1}, j}\}| \leq a$$

as needed.

Let us now describe the inverse map Φ^{-1} . Given an \mathbb{N} -matrix $D \in \text{BM}(a, b, c)$, we show how to uniquely reconstruct a plane partition $\pi \in \text{PP}(a, b, c)$ such that $\Phi(\pi) = D$. We will build π sequentially by scanning the columns of D starting from the last one.

Let us show how to *add* an element ℓ in i -th row of some plane partition π' . To do so, find the first available column of π' whose length is less than i , then add the elements ℓ in that column so that its length becomes i . For example, suppose π' already had the following shape and we want to add ℓ in the 3rd row:



Let us initially set $\pi = \emptyset$. To *add* the ℓ -th column $(d_{1\ell}, \dots, d_{b\ell})^T$ of D for $\ell = c, c-1, \dots, 1$ we do the following. For each $i = b, b-1, \dots, 1$ add ℓ in i -th row of π exactly $d_{i\ell}$ times.

Let us check that after this procedure we have $\pi \in \text{PP}(a, b, c)$. By construction, it is clear that π has at most b rows and the largest entry at most c . We know that in D the maximal weight of a monotone path from $(1, 1)$ to (b, c) is at most a . Suppose to the contrary that the length of the first row of π is larger than a . Consider the elements of the first row of π , say $(\ell_1 \geq \dots \geq \ell_k)$ for $k > a$ and suppose they were added in rows i_1, \dots, i_k during the procedure. Then we must have $i_1 \geq \dots \geq i_k$ and hence there is a monotone path in D that passes through all points $(i_k, \ell_k) \rightarrow \dots \rightarrow (i_1, \ell_1)$ (here if we have the same point (i_j, ℓ_j) repeated several times we may use it just once as its multiplicity is recorded in $d_{i_j \ell_j}$). The weight of any such monotone path is at least $\sum_j d_{i_j \ell_j} \geq k > a$ which is a contradiction. Finally, it is easy to see that π has the desired property $\Phi(\pi) = D$. \square

Example 2.3. Let us show how Φ^{-1} applies to the matrix in Example 2.1. We have:

$$\Phi^{-1} : D = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 2 & 0 & 0 \end{pmatrix} \mapsto \begin{array}{|c|c|c|} \hline 4 & 4 & 2 \\ \hline 4 & 2 & 1 \\ \hline 2 & 2 & \\ \hline \end{array}$$

which is obtained as follows:

add column 4: add 4 in row 2:

4
4

 add 4 in row 1:

4	4
4	

add column 3: no addition

add column 2: add 2 in row 3 two times:

4	4
4	2
2	2

 add 2 in row 1:

4	4	2
4	2	
2	2	

add column 1: add 1 in row 2:

4	4	2
4	2	1
2	2	

Corollary 2.4. $|\text{PP}(a, b, c)| = |\text{BM}(a, b, c)|$.

Note that by letting $a, b, c \rightarrow \infty$ one can view Φ as a bijection between plane partitions and \mathbb{N} -matrices.

We discuss few more properties of the bijection Φ that will be needed later.

Lemma 2.5. *Let π be a plane partition and $\Phi(\pi) = D = (d_{i\ell})$.*

(i) *Let $c_\ell(\pi)$ be the number of columns of π containing entry ℓ , and $c_\ell(D) = \sum_i d_{i\ell}$ be column sums of D . Then we have $c_\ell(\pi) = c_\ell(D)$ for all $\ell \geq 1$.*

(ii) *Let $\lambda = \text{sh}(\pi)$ be the shape of π . We have for all $k \geq 1$*

$$(3) \quad \lambda_k = \max_{\Pi: (k,1) \rightarrow (b,c)} \sum_{(i,\ell) \in \Pi} d_{i\ell},$$

where the maximum is taken over monotone paths Π from $(k, 1)$ to (b, c) , if D has b rows and c columns.

Proof. The part (i) is clear by construction. Indeed, by definition of Φ we have

$$c_\ell(D) = \sum_i d_{i\ell} = \sum_i |\{j : \pi_{ij} = \ell > \pi_{i+1,j}\}| = c_\ell(\pi).$$

(ii) Take any monotone path Π from $(k, 1)$ to (b, c) . Then the descent level sets $D_{i\ell}$ for $(i, \ell) \in \Pi$ are pairwise disjoint. Using this and since $i \geq k$ for all $(i, \ell) \in \Pi$, we obtain

$$(4) \quad \sum_{(i,\ell) \in \Pi} d_{i\ell} = \sum_{(i,\ell)} |\{j : \pi_{ij} = \ell > \pi_{i+1,j}\}| \leq \lambda_k.$$

On the other hand, suppose the k -th row of π has entries $(\ell_1 \geq \dots \geq \ell_m)$ where $m = \lambda_k$. Assume the entries ℓ_1, \dots, ℓ_m end in rows $i_1 \geq \dots \geq i_m$ of π . Then there is a monotone path Π from $(k, 1)$ to (b, c) that contains all points $(i_m, \ell_m), \dots, (i_1, \ell_1)$. The weight of any such path is at least $\sum_j d_{i_j \ell_j} \geq m = \lambda_k$. Combining this with the inequality (4) we obtain (3). \square

Remark 2.6. The RSK correspondence also gives a bijection between plane partitions and \mathbb{N} -matrices, see e.g. [Sta99, Ch. 7]. The bijection Φ has another (simpler) description and different yet related properties. We explore more properties and applications of this bijection in [Yel21].

Remark 2.7. Many RSK-type dynamics were described in [BP16] and it would be interesting to see how the map Φ can be related to that classification.

3. DUAL GROTHENDIECK POLYNOMIALS

Recall that a *partition* is a sequence $\lambda = (\lambda_1, \dots, \lambda_\ell)$ of nonincreasing positive integers. Here $\ell = \ell(\lambda)$ is the *length* of λ . The *Young diagram* of λ is the set $\{(i, j) : i \in [1, \ell], j \in [1, \lambda_i]\}$. We denote by λ' the *conjugate* partition of λ , which has transposed Young diagram.

Definition 3.1 ([LP07]). Let $\text{PP}_n(\lambda)$ be the set of plane partitions of shape λ and largest entry at most n . The *dual symmetric Grothendieck polynomials* $\{g_\lambda\}$ are defined by the following combinatorial formula

$$g_\lambda(x_1, \dots, x_n) = \sum_{T \in \text{PP}_n(\lambda)} x^T,$$

where $x^T = \prod_{i=1}^n x_i^{c_i}$ and c_i is the number of columns of T containing i .

The following properties hold: g_λ is an inhomogeneous symmetric polynomial whose top degree component is the Schur polynomial, i.e. $g_\lambda = s_\lambda + \text{lower degree elements}$.

Besides combinatorial definition we need the following determinantal formula. We denote $\mathbf{x} = (x_1, x_2, \dots)$ and $\{e_n\}$ are the elementary symmetric functions.

Lemma 3.2 (Jacobi-Trudi identity, see [Yel17]). *We have*

$$(5) \quad g_\lambda(\mathbf{x}) = \det \left[e_{\lambda'_i - i + j}(1^{\lambda'_i - 1}, \mathbf{x}) \right]_{1 \leq i, j \leq \lambda_1}.$$

Remark 3.3. Jacobi-Trudi formulas for g_λ were first obtained in [SZ03]. A proof was given in [Yel17]. More properties of these polynomials can be found in [LP07, Yel17, Yel19].

The following properties will be important for our results.

Lemma 3.4 (Coincidence lemma). *Let $\rho := (a^b)$. We have*

$$g_\rho(\mathbf{x}) = s_\rho(1^{b-1}, \mathbf{x}).$$

Proof. We show that the Jacobi-Trudi formulas for both polynomials coincide under these conditions. Note firstly that by the classical Jacobi-Trudi formula we have

$$(6) \quad s_\rho(1^{b-1}, \mathbf{x}) = \det \left[e_{b-i+j}(1^{b-1}, \mathbf{x}) \right]_{1 \leq i, j \leq a}$$

For the polynomials g , the Jacobi-Trudi identity (5) for $\lambda = \rho$ gives (6). \square

Remark 3.5. While this coincidence formula is evident from determinantal identities, we should point out that it is *not* at all obvious combinatorially. Here one would use an argument as in [Yel17] on combinatorial proof of the identity (5).

Corollary 3.6. *We have the following formula*

$$(7) \quad \sum_{\lambda \subset (a^b)} g_\lambda(\mathbf{x}) = s_{(a^b)}(1^b, \mathbf{x})$$

Proof. Let $\rho = (a^b)$. By combinatorial definition of g , we have the branching formula

$$g_\rho(1, \mathbf{x}) = \sum_{\lambda \subset \rho} g_\lambda(\mathbf{x}).$$

On the other hand, $g_\rho(\mathbf{x}, 1) = s_\rho(1^b, \mathbf{x})$ by Lemma 3.4. \square

Lemma 3.7. *The following identity holds*

$$(8) \quad \sum_{\ell(\lambda) \leq b} g_\lambda(\mathbf{x}) = \prod_n (1 - x_n)^{-b}$$

Proof. This identity follows from the Cauchy identity for the pair G_λ, g_λ of Grothendieck polynomials and the following property of the symmetric Grothendieck polynomials $G_\lambda(1^b) = 1$ if $\ell(\lambda) \leq b$ and 0 otherwise, see [Yel20]. In fact, one can also derive it from the bijection Φ , we address this in [Yel21]. \square

Remark 3.8. The formulas (7), (8) can perhaps be compared to the following identities for Schur functions, see [Mac98]

$$\begin{aligned} \sum_{\lambda \subset (a^b)} s_\lambda(x_1, \dots, x_b) &= \frac{\det \left[x_j^{a+2b-i} - x_j^{i-1} \right]_{1 \leq i, j \leq b}}{\det \left[x_j^{2b-i} - x_j^{i-1} \right]_{1 \leq i, j \leq b}}, \\ \sum_{\ell(\lambda) \leq b} s_\lambda(x_1, \dots, x_b) &= \prod_{1 \leq n \leq b} \frac{1}{1 - x_n} \prod_{1 \leq i < j \leq b} \frac{1}{1 - x_i x_j}. \end{aligned}$$

4. NORMALIZED SCHUR POLYNOMIALS

The *normalized Schur polynomial* $S_\lambda(\mathbf{x}; N)$ is defined for $N \geq \max(k, \ell(\lambda))$ as follows

$$(9) \quad S_\lambda(x_1, \dots, x_k; N) := \frac{s_\lambda(x_1, \dots, x_k, 1^{N-k})}{s_\lambda(1^N)}$$

We are interested in a special case when $\lambda = (a^b)$ has rectangular shape. We present some known asymptotics for normalized Schur polynomials.

Lemma 4.1 (see [BP14]). *Let $N = b + c$, $\rho = (a^b)$ and $x_i \in \mathbb{C}$. We have:*

For $a, b, c \rightarrow \infty$ with $ab/c \rightarrow t > 0$

$$S_\rho(x_1, \dots, x_k; N) \rightarrow e^{-kt} \prod_{i=1}^k e^{tx_i}$$

For b fixed and $a, c \rightarrow \infty$ with $a/(a+c) \rightarrow q \in (0, 1)$

$$S_\rho(x_1, \dots, x_k; N) \rightarrow (1-q)^{bk} \prod_{i=1}^k (1-qx_i)^{-b}$$

Consider now the following *Gaussian* asymptotic regime. Let $u > 0, q \in (0, 1)$ be real parameters. Suppose $a = \lfloor uN \rfloor$, $b = \lfloor qN \rfloor$, $b + c = N$ and let $N \rightarrow \infty$. Then the partitions $\rho = (a^b)$ (when each row is rescaled by N) have the following simple limit shape $f : [0, 1] \rightarrow \mathbb{R}$ given by

$$f(x) = \begin{cases} u, & \text{if } x \leq q, \\ 0, & \text{if } x > q. \end{cases}$$

Define two more parameters

$$(10) \quad m = \int_0^1 f(x) dx = uq, \quad v = \int_0^1 f(x)^2 dx - m^2 + \int_0^1 f(x)(1-2x) dx = u(u+1)q(1-q).$$

Lemma 4.2 ([GP15]). *Let a, b, c be in the Gaussian regime as above. As $N \rightarrow \infty$, we have*

$$S_\rho(e^{x_1/\sqrt{vN}}, \dots, e^{x_k/\sqrt{vN}}; N) \cdot \prod_{i=1}^k e^{-x_i m \sqrt{N/v}} \rightarrow \prod_{i=1}^k e^{x_i^2/2}$$

uniformly on compact subsets of $(\mathbb{R} \setminus \{0\})^k$.

Remark 4.3. In fact, this is a special case of more general results proved in [GP15].

5. MULTIVARIATE GENERATING FUNCTION FOR CORNER DISTRIBUTIONS

5.1. Distributions of Γ_ℓ in lozenge tilings. Recall that we consider the uniform probability measure on the set of lozenge tilings $\text{LT}(a, b, c)$, and Γ_ℓ is the number of left corners of height $\ell \in [1, c]$ in a random lozenge tiling. Throughout the section (and later) we set $\rho = (a^b)$, $N = b + c$, and $\Gamma^{(c)} = (\Gamma_1, \dots, \Gamma_c)$.

Theorem 5.1. *We have the generating function*

$$(11) \quad \sum_{\gamma \in \mathbb{N}^c} \mathbb{P}(\Gamma^{(c)} = \gamma) \mathbf{x}^\gamma = S_\rho(x_1, \dots, x_c; N).$$

Proof. By definition of g_λ and the bijection between lozenge tilings and plane partitions so that $\Gamma_i \mapsto X_i$ we have

$$(12) \quad \sum_{\gamma \in \mathbb{N}^c} P(\Gamma^{(c)} = \gamma) \mathbf{x}^\gamma = \sum_{\gamma \in \mathbb{N}^c} P(X_1 = \gamma_1, \dots, X_c = \gamma_c) x^\gamma = \frac{1}{Z_{abc}} \sum_{\lambda \subset \rho} g_\lambda(x_1, \dots, x_c).$$

Using (7) and the fact that $Z_{abc} = s_\rho(1^{b+c})$ we obtain

$$\frac{1}{Z_{abc}} \sum_{\lambda \subset \rho} g_\lambda(x_1, \dots, x_c) = \frac{s_\rho(1^b, x_1, \dots, x_c)}{s_\rho(1^{b+c})} = S_\rho(x_1, \dots, x_c; N).$$

as needed. \square

Corollary 5.2 (Symmetry, exchangeability). $P(\Gamma^{(c)} = \gamma)$ is invariant under permutations of the coordinates $(\gamma_1, \dots, \gamma_c)$.

Corollary 5.3. Let $k \in [1, c]$. Setting $x_{k+1} = \dots = x_c = 1$ we have

$$\sum_{\gamma \in \mathbb{N}^k} P(\Gamma^{(k)} = \gamma) \mathbf{x}^\gamma = S_\rho(x_1, \dots, x_k; N).$$

More generally, for $\{\ell_1, \dots, \ell_k\} \subset [1, c]$ we also have

$$(13) \quad \sum_{\gamma \in \mathbb{N}^k} P(\Gamma_{\ell_1} = \gamma_1, \dots, \Gamma_{\ell_k} = \gamma_k) \mathbf{x}^\gamma = S_\rho(x_{\ell_1}, \dots, x_{\ell_k}; N).$$

Corollary 5.4. For a single variable we have

$$(14) \quad \sum_n P(\Gamma_\ell = n) x^n = S_\rho(x; N).$$

In particular, $P(\Gamma_i = n) = P(\Gamma_j = n)$ for all $i, j \in [1, c]$.

We are now going to prove the limiting distributions for Γ stated in Theorem 1.3.

Proof of Theorem 1.3. Suppose we have collection of random variables $\Gamma := (\Gamma_{\ell_1}, \dots, \Gamma_{\ell_k})$.

In the first Poisson regime $a, b \rightarrow \infty$ with $ab/c \rightarrow t$ by Lemma 4.1 we have

$$S_\rho(x_{\ell_1}, \dots, x_{\ell_k}; N) \rightarrow e^{-kt} \prod_{i=1}^k e^{x_{\ell_i}}.$$

Hence by (13), Γ weakly converges to k i.i.d. Poisson random variables with parameter t .

In the second negative binomial regime for fixed b and $a, c \rightarrow \infty$ with $a/(a+c) \rightarrow q \in (0, 1)$ again by Lemma 4.1 we have

$$S_\rho(x_{\ell_1}, \dots, x_{\ell_k}; N) \rightarrow (1-q)^{bk} \prod_{i=1}^k (1-x_{\ell_i}q)^{-b}$$

Hence by (13), Γ weakly converges to k i.i.d. negative binomial random variables with parameters b, q .

Finally, in the third Gaussian regime by (14) and Lemma 4.2 for $k = 1$ we obtain that $\frac{\Gamma_\ell - mN}{\sqrt{vN}}$ weakly converges to $N(0, 1)$ as $N \rightarrow \infty$, where $v = \sigma^2$. By Lemma 4.2 and (13), we get factorization of the characteristic functions in the limit which is equivalent to asymptotic




FIGURE 4. An example of k -th slice parallel to a and the corresponding right tiles marked with \bullet . Here $Y^{(3)} = (4, 2, 0)$, the y -coordinates of marked tiles.

independence. Hence the collection of variables $\frac{\Gamma_{\ell_i - mN}}{\sqrt{vN}}$ become asymptotically independent as $N \rightarrow \infty$. \square

5.2. Distributions of X_ℓ in plane partitions and marginals C_ℓ in \mathbb{N} -matrices. The results on Γ_ℓ easily translate to results on X_ℓ on random boxed plane partitions and C_ℓ on \mathbb{N} -matrices with bounded last passage time, discussed in the introduction.

Take a uniformly random plane lozenge tilings $T \in \text{LT}(a, b, c)$ and let Ψ be the map that records heights of all top tiles, the result is clearly a plane partition $\pi = \Psi(T) \in \text{PP}(a, b, c)$ and Ψ is a bijection. We have $\Gamma_\ell(D) = X_\ell(\pi)$.

Now take a uniformly random matrix $D \in \text{BM}(a, b, c)$, then by Lemma 2.5 (i), we have $C_\ell(D) = X_\ell(\pi)$ for $\pi = \Phi^{-1}(D)$.

6. JOINT DISTRIBUTIONS OF LEFT CORNERS AND RIGHT LOZENGES

Definition 6.1 (Right lozenge distributions). Let $Y^{(k)}$ be the vector of positions i.e. y -coordinates of the right tiles \blacklozenge along k -th slice parallel to the side a of the hexagon, see Fig. 4. For $k \leq \min(b, c)$, these positions have the form

$$\mu_1 \geq \dots \geq \mu_k \geq 0.$$

Let $\mathcal{P}_{n,k} := \{\mu : \mu \vdash n, \ell(\mu) \leq k\}$ be the set of partitions of n with length at most k . We have the following Schur expansion (see [BP14]) of normalized Schur polynomials

$$(15) \quad S_\rho(x_1, \dots, x_k; N) = \sum_{\mu \in \mathcal{P}_{n,k}} \mathbb{P}(Y^{(k)} = \mu) \frac{s_\mu(x_1, \dots, x_k)}{s_\mu(1^k)}$$

where $\rho = (a^b)$, $N = b + c$, and $k \leq \min(b, c)$. The distribution of Y has the formula (e.g. [BP14])

$$\mathbb{P}(Y^{(k)} = \mu) = \text{const}_{a,b,c,k} \prod_{1 \leq i < j \leq k} (\mu_i - i - \mu_j + j)^2 \prod_{i=1}^k w(\mu_i + k - i),$$

where

$$w(y) = \frac{(a + c - 1 - y)!(b - k + y)!}{y!(a + k - 1 - y)!}$$

which is a case of more general *orthogonal polynomial ensembles*. Alternatively, there is also another formula (e.g. [GP15, Prop. 5.3])

$$(16) \quad \mathbb{P}(Y^{(k)} = \mu) = \frac{1}{Z_{abc}} s_\mu(1^k) s_{\mu^c}(1^{b+c-k}),$$

where $\mu^c = (a - \mu_b, \dots, a - \mu_1)$ is the complement to μ in ρ .

It is proved in [JN06, GP15] that for every k , as $N \rightarrow \infty$ we have convergence in distribution

$$\frac{1}{\sqrt{vN}}(Y^{(k)} - (mN, \dots, mN)) \rightarrow \text{GUE}_k,$$

where GUE_k is the distribution of the spectrum of a random $k \times k$ Hermitian matrix from the *Gaussian unitary ensemble* (GUE), and m, v are defined in (10) (cf. Lemma 4.2).

Now we would like to relate the two distributions Γ and Y .

Recall that the *Kostka numbers* $K_{\lambda\gamma} \in \mathbb{N}$ (see e.g. [Mac98, Sta99]) are coefficients defined from the monomial expansion of Schur polynomials

$$s_\lambda = \sum_{\gamma} K_{\lambda\gamma} \mathbf{x}^\gamma.$$

Consider the random vectors

$$\Gamma^{(k)} = (\Gamma_1, \dots, \Gamma_k)$$

Theorem 6.2 (Joint distribution of left corners and right tiles). *Let $\gamma = (\gamma_1, \dots, \gamma_k) \in \mathbb{N}^k$ with $|\gamma| := \gamma_1 + \dots + \gamma_k = n$ for $k \leq \min(b, c)$ and $\mu \in \mathcal{P}_{n,k}$. We have:*

(i) *The joint distribution of $\Gamma^{(k)}$ and $Y^{(k)}$ is given by*

$$\mathbb{P}(\Gamma^{(k)} = \gamma, Y^{(k)} = \mu) = \frac{1}{Z_{abc}} K_{\mu\gamma} s_{\mu^c}(1^{b+c-k}),$$

where $K_{\mu\gamma}$ is the Kostka number and $\mu^c = (a - \mu_b, \dots, a - \mu_1)$ is complement to μ in (a^b) .

(ii) *The distribution of $\Gamma^{(k)}$ can be computed as follows*

$$\mathbb{P}(\Gamma^{(k)} = \gamma) = \sum_{\mu \in \mathcal{P}_{n,k}} p_\mu(\gamma) \mathbb{P}(Y^{(k)} = \mu),$$

where the conditional probability is determined as

$$p_\mu(\gamma) = \mathbb{P}(\Gamma^{(k)} = \gamma \mid Y^{(k)} = \mu) = \frac{K_{\mu\gamma}}{s_\mu(1^k)}.$$

Proof. By Corollary 5.3 and the Schur polynomial expansion (15) we have

$$\begin{aligned} \sum_{\gamma \in \mathbb{N}^k} \mathbb{P}(\Gamma^{(k)} = \gamma) x^\gamma &= S_\rho(x_1, \dots, x_k; N) \\ &= \sum_{\mu \in \mathcal{P}_{n,k}} \mathbb{P}(Y^{(k)} = \mu) \frac{s_\mu(x_1, \dots, x_k)}{s_\mu(1^k)} \\ &= \sum_{\mu \in \mathcal{P}_{n,k}} \mathbb{P}(Y^{(k)} = \mu) \sum_{\alpha} \frac{K_{\mu\alpha}}{s_\mu(1^k)} x^\alpha. \end{aligned}$$

Therefore,

$$P(\Gamma^{(k)} = \gamma) = \sum_{\mu \in \mathcal{P}_{n,k}} \frac{K_{\mu\gamma}}{s_\mu(1^k)} P(Y^{(k)} = \mu), \quad \frac{K_{\mu\gamma}}{s_\mu(1^k)} = P(\Gamma^{(k)} = \gamma | Y^{(k)} = \mu).$$

To find the joint distribution we have

$$\begin{aligned} P(\Gamma^{(k)} = \gamma, Y^{(k)} = \mu) &= P(\Gamma^{(k)} = \gamma | Y^{(k)} = \mu) \cdot P(Y^{(k)} = \mu) \\ &= \frac{K_{\mu\gamma}}{s_\mu(1^k)} \cdot \frac{s_\mu(1^k) s_{\mu^c}(1^{b+c-k})}{Z_{abc}} \\ &= \frac{K_{\mu\gamma} s_{\mu^c}(1^{b+c-k})}{Z_{abc}} \end{aligned}$$

where we used the formula (16). \square

Now using some properties of Kostka numbers we obtain the following inequalities.

Corollary 6.3. *Since $K_{\mu\mu} = 1$ we get the following inequality for $\mu \in \mathcal{P}_{n,k}$*

$$s_\mu(1^k) P(\Gamma^{(k)} = \mu) \geq P(Y^{(k)} = \mu).$$

For $\alpha, \beta \in \mathbb{N}^k$, say that β dominates α and write $\beta \succeq \alpha$ if $\sum_{i=1}^\ell \beta_i \geq \sum_{i=1}^\ell \alpha_i$ for all $\ell \in [1, k]$ and $\sum_{i=1}^k \alpha_i = \sum_{i=1}^k \beta_i$.

Corollary 6.4 (Monotonicity). *The following inequality holds*

$$\forall \beta \succeq \alpha : \quad P(\Gamma^{(k)} = \alpha) \geq P(\Gamma^{(k)} = \beta).$$

Proof. It is known that $K_{\mu\alpha} \geq K_{\mu\beta}$ for $\beta \succeq \alpha$ (see [Mac98]). Hence $p_\mu(\alpha) \geq p_\mu(\beta)$ and subsequently $P(\Gamma^{(k)} = \alpha) \geq P(\Gamma^{(k)} = \beta)$. \square

7. g -MEASURE

Let $q_1, \dots, q_c \in (0, 1)$. Recall that we define the probability distribution, called g -measure or g -distribution, on the (infinite) set $\text{PP}(\infty, b, c)$ of plane partitions with at most b rows and maximal entry at most c

$$P_{g,b,c}(\pi) = \frac{1}{Z_{b,c}} \prod_{(i,j) \in \text{Des}(\pi)} q_{\pi_{ij}},$$

where $Z_{b,c}$ is the normalization constant.

Proposition 7.1. *Let $\pi \in \text{PP}(\infty, b, c)$ and $\lambda = (\lambda_1 \geq \dots \geq \lambda_b \geq 0)$. We have*

$$P_{g,b,c}(\text{sh}(\pi) = \lambda) = \frac{1}{Z_{b,c}} g_\lambda(q_1, \dots, q_c).$$

Proof. Using the definition of g_λ , it is not difficult to see that

$$P_{g,b,c}(\lambda) = \sum_{\text{sh}(\pi)=\lambda} P_{b,c}(\pi) = \frac{1}{Z_{b,c}} \sum_{\text{sh}(\pi)=\lambda} \prod_{(i,j) \in \text{Des}(\pi)} q_{\pi_{ij}} = \frac{1}{Z_{b,c}} g_\lambda(q_1, \dots, q_c),$$

where the sum runs over $\pi \in \text{PP}(\infty, b, c)$. \square

Corollary 7.2. *We have*

$$Z_{b,c} = \prod_{i=1}^c (1 - q_i)^{-b}.$$

Proof. Follows from the proposition and the identity (8). \square

Theorem 7.3. *For $\pi \in \text{PP}(\infty, b, c)$, let $\lambda = (\lambda_1 \geq \dots \geq \lambda_b \geq 0)$ be the shape of π . The distribution of the first part λ_1 can be expressed as*

$$(17) \quad P_{g,b,c}(\lambda_1 \leq a) = \frac{1}{Z_{b,c}} s_{(a^b)}(1^b, q_1, \dots, q_c).$$

Proof. For the distribution of λ_1 we now have

$$\begin{aligned} P_{g,b,c}(\lambda_1 \leq a) &= \sum_{\lambda \subset (a^b)} P_{g,b,c}(\lambda) \\ &= \frac{1}{Z_{b,c}} \sum_{\lambda \subset (a^b)} g_{\lambda}(q_1, \dots, q_c) \\ &= \frac{1}{Z_{b,c}} g_{(a^b)}(1, q_1, \dots, q_c) \\ &= \frac{1}{Z_{b,c}} s_{(a^b)}(1^b, q_1, \dots, q_c), \end{aligned}$$

where we used Lemma 3.4 in the last identity. \square

Corollary 7.4. *The distribution of λ_1 can be expressed via the Toeplitz determinant*

$$(18) \quad P_{b,c}(\lambda_1 \leq a) = \frac{1}{Z_{b,c}} \det [\hat{\varphi}_{i-j}]_{i,j=1}^a,$$

where

$$\hat{\varphi}_k = \sum_{\ell=0}^{\infty} \binom{b}{\ell+k} e_{\ell}(q_1, \dots, q_c) \quad \text{or} \quad \varphi(z) := \sum_{k \in \mathbb{Z}} \hat{\varphi}_k z^k = (1+z)^b \prod_{i=1}^c (1+q_i z^{-1}).$$

Proof. Using the Jacobi-Trudi identity for Schur polynomials we obtain

$$s_{(a^b)}(1^b, q_1, \dots, q_c) = \det[e_{b-i+j}(1^b, q_1, \dots, q_c)]_{i,j=1}^a$$

Note that

$$e_{b-i+j}(1^b, q_1, \dots, q_c) = \sum_{\ell \geq 0} \binom{b}{b-i+j-\ell} e_{\ell}(q_1, \dots, q_c) = \sum_{\ell \geq 0} \binom{b}{\ell+i-j} e_{\ell}(q_1, \dots, q_c) = \hat{\varphi}_{i-j}$$

as needed. \square

Remark 7.5. From the Toeplitz determinant we can also obtain *Fredholm determinant* formula of the form $\det[I - K]$ for some kernel K using the Borodin-Okounkov formula [BO00].

Remark 7.6. Let Λ be the ring of symmetric functions. A specialization (ring homomorphism) $\rho : \Lambda \rightarrow \mathbb{R}$ is called *g-positive* if $\rho(g_{\lambda}) \geq 0$ for all λ . In light of the *g-measure* one can be interested in characterizations of *g-positive* specializations. See [Yel20] for a class of *g-positive* specializations and related questions.

8. CORNER GROWTH MODEL

Consider a random matrix $W = (w_{ij})$ with i.i.d. entries w_{ij} that are geometrically distributed with parameter q , i.e.

$$P(W_{ij} = k) = (1 - q)q^k, \quad k \in \mathbb{N}.$$

For $m, n \geq 1$, recall that the last passage times $G(m, n)$ are defined as

$$G(m, n) = \max_{\Pi} \sum_{(i,j) \in \Pi} w_{ij},$$

where the maximum is taken over monotone paths Π from $(1,1)$ to (m,n) . Note that we have the recurrence relation

$$G(i, j) = \max(G(i-1, j), G(i, j-1)) + w_{ij}, \quad i, j \geq 1$$

(one needs to set the boundary conditions $G(i, 0) = G(0, i) = 0$).

The *corner growth model* can be viewed as evolution of a random Young diagram $Y(t)$ given at time t by the region

$$Y(t) = \{(i, j) \in \mathbb{N}^2 : G(i, j) \leq t\}.$$

One can see that $Y(t+1)$ adds boxes to some corners of $Y(t)$. We refer to [Sep09, Rom15] for more on the topic.

The main result of this section is the following.

Theorem 8.1. *Let $\lambda = (\lambda_1, \dots, \lambda_b)$ be a partition with at most b parts. We have*

$$P(G(b, c) = \lambda_1, \dots, G(1, c) = \lambda_b) = P_{g,b,c}(\lambda) = (1 - q)^{bc} g_{\lambda}(\underbrace{q, \dots, q}_{c \text{ times}}),$$

where $q_i = q$ for all $i \in [1, c]$ in the g -measure.

To prove this theorem we need the following lemma.

Lemma 8.2. *Let $D = (D_{ij})_{i,j=1}^{b,c}$ be a random $b \times c$ matrix, where D_{ij} are i.i.d. geometrically distributed random variables with parameter q . Then*

$$P(\Phi^{-1}(D) = \pi) = P_{g,b,c}(\pi),$$

where $q_i = q$ for all $i \in [1, c]$ in the g -measure.

Proof. Note firstly that for $q_i = q$ we have

$$P_{g,b,c}(\pi) = (1 - q)^{bc} q^{\text{des}(\pi)}.$$

Let $d_{i\ell} = |\{j : \pi_{ij} = \ell > \pi_{i+1j}\}|$, i.e. $(d_{i\ell}) = \Phi(\pi)$. Then

$$P(D = \Phi(\pi)) = \prod_{i=1}^b \prod_{\ell=1}^c P(D_{i\ell} = d_{i\ell}) = \prod_{i=1}^b \prod_{\ell=1}^c (1 - q) q^{d_{i\ell}} = P_{b,c}(\pi)$$

as needed. □

Proof of Theorem 8.1. Follows now from Lemma 2.5 (ii), Lemma 8.2 and Proposition 7.1. □

We now list some implications of Theorem 8.1.

Corollary 8.3 (Marginal distributions). *Let $k \in [1, b]$. We have*

$$P(G(b - k + 1, c) \leq n) = P_{g,b,c}(\lambda_k \leq n).$$

In particular, $\lambda_b \sim G(1, c)$ has binomial distribution with parameters c, q . For $\lambda_1 \sim G(b, c)$ combining with Theorem 7.3 we obtain.

Corollary 8.4. *We have*

$$P(G(b, c) \leq a) = (1 - q)^{bc} s_{(a^b)}(1^b, \underbrace{q, \dots, q}_{c \text{ times}})$$

as well as Toeplitz determinantal expressions (18).

More generally using the Jacobi-Trudi-type determinantal formula (5) we get

Corollary 8.5. *We have the following formula*

$$P(G(b, c) = \lambda_1, \dots, G(1, c) = \lambda_b) = (1 - q)^{bc} \det \left[e_{\lambda'_i - i + j}(1^{\lambda'_i - 1}, \underbrace{q, \dots, q}_{c \text{ times}}) \right]_{i,j=1}^{\lambda_1}$$

Let us now discuss several results about the corner growth model that imply results on the shape λ from the g -distribution.

It was proved in [Joh00] that for $x > 0$ we have almost sure convergence to deterministic limit shape

$$\lim_{b \rightarrow \infty} \frac{1}{b} G(\lfloor xb \rfloor, b) \rightarrow \psi(x) = (1 - q)^{-1}(qx + q + 2\sqrt{qx})$$

and *fluctuations* around this limit shape are of order $b^{1/3}$ given by

$$\lim_{b \rightarrow \infty} P \left(\frac{G(\lfloor xb \rfloor, b) - \psi(x)b}{\sigma(x)b^{1/3}} \leq t \right) = F_{\text{TW}}(t),$$

where $\sigma(x) = (1 - q)^{-1}(qx)^{1/6}(\sqrt{x} + \sqrt{q})^{2/3}(1 + \sqrt{qx})^{2/3}$ and $F_{\text{TW}}(t)$ is the *Tracy-Widom distribution*, the limiting distribution of the properly scaled largest eigenvalue in GUE [TW94]. Therefore, combining these with Corollary 8.3, λ from the g -distribution has limit shape

$$\lim_{b \rightarrow \infty} \frac{\lambda_{\lfloor xb \rfloor}}{b} \rightarrow \psi(1 - x)$$

where $x \in (0, 1]$, $c = b$; and similar fluctuations.

It was proved in [Bar01] that for fixed b , as $c \rightarrow \infty$, the variables

$$(19) \quad \frac{G(k, c) - \frac{q}{1-q}c}{\frac{\sqrt{q}}{1-q}\sqrt{c}}, \quad k = 1, \dots, b$$

jointly converge in distribution to largest eigenvalues of b principal minors in $b \times b$ GUE matrix, which is known as *GUE corners process* (or GUE minors process [JN06]). Therefore, for λ from the g -distribution we have

$$(20) \quad \frac{\lambda_{b-k+1} - \frac{q}{1-q}c}{\frac{\sqrt{q}}{1-q}\sqrt{c}}, \quad k = 1, \dots, b$$

jointly converge in distribution to largest eigenvalues in $b \times b$ GUE corners process. It is also known [GW91] that as $c \rightarrow \infty$ the random variables (19) converge in distribution to the process

$$\sup_{0=t_0 < \dots < t_k=1} \sum_{i=0}^{k-1} (B_i(t_{i+1}) - B_i(t_i)), \quad k = 1, 2, \dots$$

where B_i are independent standard Brownian motions. Hence the same holds for the random variables (20).

Remark 8.6. After this paper was written, more related works on probabilistic models for dual Grothendieck polynomials appeared in [Yel20a, MS20].

Remark 8.7. The g -measure and Theorem 8.1 have higher-dimensional generalizations obtained recently in [AY20].

ACKNOWLEDGEMENTS

I am grateful to Pavlo Pylyavskyy for many stimulating conversations on K -theoretic combinatorics. I am also grateful to Leonid Petrov for helpful comments and some references. I am also thankful to the referees for careful reading of the text and many useful comments.

REFERENCES

- [AY20] A. Amanov and D. Yeliussizov, MacMahon's statistics on higher-dimensional partitions, arXiv:2009.00592 (2020).
- [Bar01] Y. Baryshnikov, GUEs and queues, Probability Theory and Related Fields **119** (2001), 256–274.
- [BF14] A. Borodin and P. Ferrari, Anisotropic growth of random surfaces in 2+1 dimensions, Commun. Math. Phys. **325** (2014), 603–684.
- [BO00] A. Borodin and A. Okounkov, A Fredholm determinant formula for Toeplitz determinants, Integral Equations and Operator Theory **37** (2000), 386–396.
- [BP14] A. Borodin and L. Petrov, Integrable probability: From representation theory to Macdonald processes, Probability Surveys **11** (2014), 1–58.
- [BP16] A. Borodin and L. Petrov, Nearest neighbor Markov dynamics on Macdonald processes, Adv. Math. **300** (2016), 71–155.
- [Buc02] A. Buch, A Littlewood Richardson rule for the K-theory of Grassmannians, Acta Math. **189** (2002), 37–78.
- [Buc05] A. Buch, Combinatorial K -theory, Topics in cohomological studies of algebraic varieties, Birkhäuser Basel, 2005, 87–103.
- [CKP01] H. Cohn, R. Kenyon, and J. Propp, A variational principle for domino tilings. J. Amer. Math. Soc. **14** (2001), 297–346.
- [CLP98] H. Cohn, M. Larsen, and J. Propp, The shape of a typical boxed plane partition, New York J. Math. **4** (1998), 137–165.
- [GW91] P. Glynn and W. Whitt, Departures from many queues in series, Ann. Appl. Probability, **1** (1991), 546–572.
- [GP15] V. Gorin and G. Panova, Asymptotics of symmetric polynomials with applications to statistical mechanics and representation theory, Ann. Probability **43** (2015), 3052–3132.
- [Joh00] K. Johansson, Shape fluctuations and random matrices, Commun. Math. Phys. **209** (2000), 437–476.
- [Joh01] K. Johansson, Random growth and random matrices, European Congress of Mathematics, Birkhäuser, Basel, 2001.
- [Joh02] K. Johansson, Non-intersecting paths, random tilings and random matrices, Probability theory and related fields **123** (2002), 225–280.

- [JN06] K. Johansson and E. Nordenstam, Eigenvalues of GUE minors, *Electronic J. Probability* **11** (2006), 1342–1371.
- [Ken09] R. Kenyon, Lectures on dimers, in *Statistical mechanics*, AMS, Providence, RI, 2009, 191–230.
- [IN13] T. Ikeda, and H. Naruse, K -theoretic analogues of factorial Schur P - and Q -functions, *Adv. Math.* **243** (2013), 22–66.
- [LP07] T. Lam and P. Pylyavskyy, Combinatorial Hopf algebras and K -homology of Grassmannians, *Int. Math. Res. Not. Vol. 2007*, (2007), rnm 125.
- [Len00] C. Lenart, Combinatorial aspects of the K -theory of Grassmannians, *Ann. Comb.* **4** (2000), 67–82.
- [Mac98] I. G. Macdonald, *Symmetric functions and Hall-Littlewood polynomials*, Oxford Univ. Press, Oxford (1998).
- [MS14] K. Motegi and K. Sakai, Vertex models, TASEP and Grothendieck polynomials, *Journal of Physics A: Mathematical and Theoretical* **46** (2013): 355201.
- [MS20] K. Motegi and T. Scrimshaw, Refined dual Grothendieck polynomials, integrability, and the Schur measure, *arXiv:2012.15011* (2020).
- [Oko01] A. Okounkov, Infinite wedge and measures on partitions, *Selecta Math.* **7** (2001), 57–81.
- [OR03] A. Okounkov and N. Reshetikhin, Correlation function of Schur process with application to local geometry of a random 3-dimensional Young diagram, *J. Amer. Math. Soc.* **16** (2003), 581–603.
- [Pet14] L. Petrov, Asymptotics of random lozenge tilings via Gelfand–Tsetlin schemes, *Probability theory and related fields* **160** (2014), 429–487.
- [Rom15] D. Romik, *The surprising mathematics of longest increasing subsequences*, Cambridge Univ. Press, New York, 2015.
- [SZ03] M. Schimozono and M. Zabrocki, Stable Grothendieck symmetric functions and Ω -calculus, preprint, 2003.
- [Sep09] T. Seppäläinen, Lecture notes on the corner growth model, Unpublished notes (2009).
- [Sos00] A. Soshnikov, Determinantal random point fields, *Russian Math. Surveys* **55** (2000), 923–975.
- [Sta99] R. Stanley, *Enumerative Combinatorics*, Vol 2, Cambridge, 1999.
- [TY11] H. Thomas and A. Yong, Longest increasing subsequences, Plancherel-type measure and the Hecke insertion algorithm, *Adv. Appl. Math.* **46** (2011), 610–642.
- [TW94] C. Tracy and H. Widom, Level Spacing Distributions and the Airy Kernel, *Commun. Math. Phys.* **159** (1994), 151–174.
- [Vak06] R. Vakil, A geometric Littlewood–Richardson rule, *Ann. Math.* (2006), 371–422.
- [Yel17] D. Yeliussizov, Duality and deformations of stable Grothendieck polynomials, *J. Algebraic Combin.* **45** (2017), 295–344.
- [Yel19] D. Yeliussizov, Symmetric Grothendieck polynomials, skew Cauchy identities, and dual filtered Young graphs, *J. Combin. Theory Ser. A* **161** (2019), 453–485.
- [Yel20] D. Yeliussizov, Positive specializations of symmetric Grothendieck polynomials, *Adv. Math.* **363** (2020), Article 107000.
- [Yel20a] D. Yeliussizov, Dual Grothendieck polynomials via last-passage percolation, *C. R. Math. Acad. Sci. Paris* **358** (2020), 497–503.
- [Yel21] D. Yeliussizov, Enumeration of plane partitions by descents, *J. Combin. Theory Ser. A* **178** (2021), Article 105367.

KBTU, ALMATY, KAZAKHSTAN

Email address: yeldamir@gmail.com