

Strictly linear light cones in long-range interacting systems of arbitrary dimensions

Tomotaka Kuwahara^{1,2*}

¹ *Mathematical Science Team, RIKEN Center for Advanced Intelligence Project (AIP),
1-4-1 Nihonbashi, Chuo-ku, Tokyo 103-0027, Japan and*

² *Interdisciplinary Theoretical & Mathematical Sciences Program
(iTHEMS) RIKEN 2-1, Hirosawa, Wako, Saitama 351-0198, Japan*

Keiji Saito^{3†}

³ *Department of Physics, Keio University, Yokohama 223-8522, Japan*

In locally interacting quantum many-body systems, the velocity of information propagation is finitely bounded and a linear light cone can be defined. Outside the light cone, the amount of information rapidly decays with distance. When systems have long-range interactions, it is highly nontrivial whether such a linear light cone exists. Herein, we consider generic long-range interacting systems with decaying interactions, such as $R^{-\alpha}$ with distance R . We prove the existence of the linear light cone for $\alpha > 2D + 1$ (D : the spatial dimension), where we obtain the Lieb–Robinson bound as $\|[O_i(t), O_j]\| \lesssim t^{2D+1}(R - \bar{v}t)^{-\alpha}$ with $\bar{v} = \mathcal{O}(1)$ for two arbitrary operators O_i and O_j separated by a distance R . Moreover, we provide an explicit quantum-state transfer protocol that achieves the above bound up to a constant coefficient and violates the linear light cone for $\alpha < 2D + 1$. In the regime of $\alpha > 2D + 1$, our result characterizes the best general constraints on the information spreading.

II. INTRODUCTION

In deep understanding of many-body physics, we necessarily encounter the question on how fast information propagates in the dynamics. In this context, the most fundamental principle is the causality; that is, in relativistic systems, information propagation is completely prohibited outside the light cone. On the other hand, in non-relativistic quantum many-body systems, a rigorous light cone is not defined. Lieb and Robinson proved in 1972 [1] that an effective light cone can be defined, outside which information propagation exponentially decreases with distance. In their study, the effective light cone is characterized by the so-called Lieb–Robinson velocity.

The Lieb–Robinson bound imposes one of the most fundamental restrictions to the dynamics [2–13] and has been improved in various ways [9, 14–20]. Moreover, after Hastings’ work on the Lieb–Schultz–Mattis theorem [21], the Lieb–Robinson bound has been recognized as a crucial ingredient for analyzing the universal physics in many-body systems, such as quasi-adiabatic continuation [22], the area law of entanglement [23–26], thermalization [27–30], quantization of Hall conductance [31, 32], stability of the topological order [33–35], clustering theorem for correlation functions [14, 15, 36–38], effective Hamiltonian theory [39, 40], classical simulation of many-body systems [41–47]. More recently, the Lieb–Robinson bound has been further applied to the digital quantum simulation of many-body systems [48–50] and quantum information scrambling [51–63], where the Lieb–Robinson velocity gives an upper bound of butterfly speed [51]. In addition, experimental advancement enables direct observation of the Lieb–Robinson bounds [64–70].

In the case of short-range interacting spin systems, an effective light cone is characterized by a finite velocity, and information propagation is restricted inside the

“linear light cone.” However, when we consider long-range interacting systems, the existence of a linear light cone is quite subtle because long-range interactions enable immediate communication between two arbitrarily distant parties. Here, long-range interaction implies that the interaction strength between separated sites shows a power-law decay of $R^{-\alpha}$ with the distance R . Depending on the exponent α , both the linear and non-linear light cones can appear. Recent experiments have realized long-range interacting systems with various values of α [71–80], and hence, exploring the universal aspects of long-range interacting systems is attracting increasing attention [81–87]. From these backgrounds, one of the most important and intriguing open problems is the so-called *linear light cone problem*, which clarifies whether linear light cones can exist in long-range interacting systems, and what is the general criterion for it.

So far, various studies have clarified dynamical properties in the long-range interacting systems [88–104]. As one of the generic aspects of the Lieb–Robinson bound, Refs. [15, 17, 105–107] showed that the amount of information propagation is suppressed at least outside the effective light cone exponentially growing in time, irrespective of α . Later, a more detailed universal upper bound was provided by Foss–Feig et al. [108]. They proved that the effective light cone is at most polynomial with respect to time; in more detail, the shape of the light-cone was given by $t^{(\alpha-D+1)/(\alpha-2D)}$ ($\alpha > 2D$) with D as the spatial dimensions. However, despite significant efforts [49, 109–111], the critical value of α to obtain a linear light cone is still unclear even in the numerical level.

In this work, we rigorously prove that a linear light cone is obtained in generic long-range interacting systems under the condition of $\alpha > 2D + 1$. As related work, in one-dimensional two-body interacting systems, the long-range Lieb–Robinson bound has been proved very recently in the form of $\|[O_i(t), O_j]\| \lesssim t/R$ for $\alpha > 3$ [112], which gives a non-trivial upper bound up to the time $t = \mathcal{O}(R)$. In our analyses, the Hamiltonian is not restricted to few-body interactions, and is applica-

* tomotaka.kuwahara@riken.jp

† saitoh@rk.phys.keio.ac.jp




FIG. 1. When we consider time-evolution $O_i(t)$ of a local operator O_i , the quasi-locality of the interaction ensures that $O_i(t)$ is well-approximated by an operator defined on a ball region $i[r]$ having the maximum distance of r from i . The dynamics define the linear light cone if we can achieve an arbitrary approximation error for $r = O(t)$ as in (5).

ble to arbitrary spatial dimensions. Our Lieb–Robinson bound is given in the stronger form of $\|[O_i(t), O_j]\| \lesssim t^{2D+1}(R - \bar{v}t)^{-\alpha}$. However, only the above commutation relation is not sufficient to upper-bound the entire information propagation outside the light cone. To obtain a linear light cone in a strict sense (see (5) and Fig. 1), we also prove that the error of the local approximation of $O_i(t)$ decays as $t^{D+1}R^{-\alpha+D}$ outside the light cone (see (7) below). Our result can improve various existing analyses that depend on a polynomial light cone [49, 108].

We also discuss whether $\alpha = 2D + 1$ is the critical value to ensure a linear light cone in general setups. In other words, we investigate the achievability of our Lieb–Robinson bound and the possibility to violate the linear-light cone for $\alpha < 2D + 1$, by which we show that our bound is the best general upper bound. We consider a quantum state-transfer protocol through the dynamics on a spin network and give an explicit example that achieves our Lieb–Robinson bound for $\alpha > 2D + 1$ and violates the linear light cone for $\alpha < 2D + 1$. Our protocol is applied to $(1/2)$ -spin systems and comprises only the Ising-type long-range interactions and simple short-range interactions that generate the controlled NOT gate operation. This example ensures the optimality of our results.

The rest of this paper is organized as follows. In Sec. II, we formulate the precise setting and state the main results and their implications. In Sec. III, we sketch the intuitive explanation for the condition that linear light cone appears. We also show the brief strategy for the proof. In Sec. IV, we discuss the optimality of the present Lieb–Robinson bound. In more detail, we explicitly show a quantum state transfer protocol which achieves our theoretical upper bound up to a coefficient. Finally, in Sec. V, we summarize the paper making a brief discussion.

III. MAIN RESULTS

We consider a quantum many-body system with n sites, where each site is located on a D -dimensional lattice with the total set Λ ($|\Lambda| = n$). We assume that a

finite-dimensional Hilbert space is assigned to each of the sites.

For simplicity, we restrict ourselves to two-body interactions, but our results are extended to a more general setup, as shown in Appendix VII. We focus on the following Hamiltonian H with power-law decaying interactions:

$$H = \sum_{i,j \in \Lambda} h_{i,j} + \sum_{i=1}^n h_i \quad \text{with} \quad \|h_{i,j}\| \leq \frac{g_0}{d_{i,j}^\alpha}, \quad (1)$$

where $d_{i,j}$ is the distance between the sites i and j , namely the minimum path length from the site i to j , $\{h_{i,j}\}_{i < j}$ are the bi-partite interaction operators, $\{h_i\}_{i=1}^n$ are the local potentials, g_0 is a positive constant of $\mathcal{O}(1)$, and $\|\cdots\|$ is the operator norm. Although in some literatures [113, 114] the long-range interaction often implies the power-law decaying interactions with $\alpha \leq D$, we refer to systems with *arbitrary power-law decaying interactions* as long-range interacting systems in distinction from the finite-range (or, exponentially decaying) interactions.

Our analysis can be also applied to a time-dependent Hamiltonian, but for simplicity, we present the analysis only for the time-independent case. One typical example is the following one-dimensional long-range Ising model:

$$H = \sum_{i < j} \frac{g_0}{d_{i,j}^\alpha} \sigma_i^x \sigma_j^x + B \sum_i \sigma_i^z \quad (D = 1), \quad (2)$$

where $\{\sigma^x, \sigma^y, \sigma^z\}$ are the Pauli matrices. This class of Hamiltonians is experimentally realized for the power-law exponent in $\alpha \leq 3$ [69, 79].

We are now interested in the time-evolution using the Hamiltonian H . For simplicity, we consider an operator O_i that is locally defined on the site i and analyze

$$O_i(t) := e^{iHt} O_i e^{-iHt}.$$

Mainly, we focus on the following two quantities:

$$\|[O_i(t), O_j]\| \quad \text{for} \quad \forall j \in \Lambda, \quad (3)$$

and

$$\begin{aligned} & \|[O_i(t) - O_i(t, i[r])]\| \quad \text{with} \\ & O_i(t, i[r]) := \frac{1}{\text{tr}_{i[r]^c}(\hat{1})} \text{tr}_{i[r]^c} [O_i(t)] \otimes \hat{1}_{i[r]^c}, \end{aligned} \quad (4)$$

where $i[r]$ denotes the set of sites having a maximum distance of r from the site i and $i[r]^c$ is its complement set. The quantity (4) characterizes the error of the local approximation for the time-evolved operator $O_i(t)$ into the region $i[r]$ (see Fig. 1). We note that the decay of (4) is not necessarily derived only from the decay of (3).

Here, we define the linear light cone in the following sense. We say that the Hamiltonian dynamics e^{-iHt} has a linear light cone if the following inequality holds for an arbitrary error $\delta \in \mathbb{R}$ and t :

$$\|[O_i(t) - O_i(t, i[r])]\| \leq \delta \quad \text{for} \quad r \geq v_{t,\delta}|t|, \quad (5)$$

where $v_{t,\delta}$ decreases in time and eventually converges to a finite value (i.e., $v_{\infty,\delta} = \text{const.}$). From the definition,




FIG. 2. Effect of the operator spreading. When a bipartite interaction $h_{i,j}$ evolves using the Hamiltonian H_1 , interactions between the sites i' and j' such that $d_{i,i'} \lesssim vt$ and $d_{j,j'} \lesssim vt$ are effectively induced. The operator $h_{i,j}(H_1, t)$ now acts on a subsystem with roughly $\mathcal{O}(t^D)$ sites. Hence, one-site energy of $H_2(H_1, t)$ defined as in Eq. (9) becomes $\mathcal{O}(t^D)$ times the original one.

the amount of information propagation is smaller than δ outside the region separated by the distance $v_{t,\delta}|t|$.

Here, we show our main results. For $\alpha > 2D + 1$, the Hamiltonian in Eq. (1) satisfies the Lieb–Robinson bound for (3) as

$$\|[O_i(t), O_j]\| \leq \mathcal{C}_H |t|^{2D+1} (R - \bar{v}|t|)^{-\alpha} \quad (6)$$

with $R = d_{i,j}$. In addition, the Lieb–Robinson bound for (4) is given by

$$\|[O_i(t) - O_i(t, i[R])]\| \leq \mathcal{C}'_H |t|^{D+1} (R - \bar{v}|t|)^{-\alpha+D}, \quad (7)$$

where $R > \bar{v}|t|$ is considered and $\mathcal{C}_H, \mathcal{C}'_H$, and \bar{v} are constants that depend only on the parameters $\{D, g_0, \alpha\}$ and a geometric constant defined by the lattice structure. We emphasize that the same upper bound is obtained for generic operators O_X and O_Y (see Appendix VII).

As long as we consider the commutator for local observables, the first inequality (6) is stronger than the second one (7) in the sense that asymptotic decay is as small as $\mathcal{O}(R^{-\alpha})$. From the inequality (7), we can calculate the Lieb–Robinson velocity $v_{\delta,t}$ defined in Eq. (5):

$$v_{\delta,t} = \bar{v} + c\delta^{-\frac{1}{\alpha-D}} |t|^{\frac{-\alpha+2D+1}{\alpha-D}} \xrightarrow{t \rightarrow \infty} \bar{v}$$

where c is a constant of $\mathcal{O}(1)$, and where we use the condition $\alpha > 2D + 1$.

IV. INTUITIVE EXPLANATION OF THE CONDITION $\alpha > 2D + 1$

We here show an intuitive explanation of why the condition $\alpha > 2D + 1$ appears. The point to obtain the linear light cone is that the contribution to the Lieb–Robinson velocity from the long-range interaction with very large distance becomes asymptotically negligible. To get better insights into this condition, we consider the simplest setup as follows.

Let us consider the Hamiltonian given by $H = H_1 + H_2$, where H_1 has only nearest-neighbor interactions, while H_2 consists of the long-range interactions with the length scales from ℓ to 2ℓ as $H_2 = \sum_{\ell \leq d_{i,j} \leq 2\ell} h_{i,j}$. Note that the condition (1) implies $\|h_{i,j}\| \leq g_0 d_{i,j}^{-\alpha}$. We

eventually take the large ℓ limit to consider asymptotic behavior arising from the interactions of the large distance. Note that the Hamiltonian H_1 consists of short-range interactions; hence, the unitary operator $e^{-iH_1 t}$ satisfies the standard Lieb–Robinson bound [1, 14, 15] giving a finite Lieb–Robinson velocity. Here, we denote it by v_1 . We focus on the time range of $t \lesssim \ell/v_1$ and then consider the condition for which the Lieb–Robinson velocity for the Hamiltonian H is given by a finite velocity related to v_1 in the large ℓ limit.

As a simple exercise, we first consider the product of the unitary operators $e^{-iH_1 t} e^{-iH_2 t}$. Then, to estimate the contribution from the long-range interactions, we only have to consider the Lieb–Robinson bound for $e^{-iH_2 t}$. The Lieb–Robinson bound for H_2 is given by $e^{-c(x/\ell - v_\ell t)}$, and v_ℓ is proportional to the one-site energy:

$$g = \max_{i \in \Lambda} \sum_{j \in \Lambda} \|h_{i,j}\| = \mathcal{O}(\ell^{-\alpha+D}),$$

where $\sum_{j \in \Lambda} \|h_{i,j}\| \leq g_0 \sum_{j: \ell \leq d_{i,j} \leq 2\ell} d_{i,j}^{-\alpha}$ is a summation of all the interaction terms that act on the site i . Hence, the Lieb–Robinson velocity is proportional to $\ell^{-\alpha+D+1}$, which vanishes in the limit of $\ell \rightarrow \infty$ for $\alpha > D + 1$. Therefore, the unitary operator $e^{-iH_1 t} e^{-iH_2 t}$ has finite Lieb–Robinson velocity v_1 as long as $\alpha > D + 1$.

Now, let us discuss the unitary operator $e^{-i(H_1+H_2)t}$. For this unitary operator, we use the following representation to decompose the contributions from H_1 and H_2 :

$$e^{-i(H_1+H_2)t} = e^{-iH_1 t} \mathcal{T} e^{-i \int_0^t H_2(H_1, \tau) d\tau}, \quad (8)$$

where $H_2(H_1, \tau) := e^{iH_1 \tau} H_2 e^{-iH_1 \tau}$ and \mathcal{T} denotes the time-ordering operator. Because the one-site operator spreads up to a distance $\mathcal{O}(\tau)$ owing to the time-evolution $e^{-iH_1 \tau}$ (Fig. 2), the one-site energy is now given by

$$g(\tau) = \max_{i \in \Lambda} \sum_{Z: Z \ni i} \|h_{\tau, Z}\| = \tau^D \mathcal{O}(\ell^{-\alpha+D}), \quad (9)$$

where $h_{\tau, Z}$ is an interaction term on the subset Z that constitutes $H_2(H_1, \tau)$, i.e., $H_2(H_1, \tau) = \sum_Z h_{\tau, Z}$.

Therefore, the time-evolution $\mathcal{T} e^{-i \int_0^t H_2(H_1, \tau) d\tau}$ gives the Lieb–Robinson bound as $e^{-c(x/\ell - v_{t,\ell} t)}$ with the velocity $v_{t,\ell} \propto g(t) = t^D \mathcal{O}(\ell^{-\alpha+D})$. For $t \lesssim \ell/v_1$, this estimation provides the Lieb–Robinson velocity as $t^D \ell^{-\alpha+D+1} \lesssim \mathcal{O}(\ell^{-\alpha+2D+1})$. Hence, the contribution to the Lieb–Robinson velocity from H_2 vanishes in the limit of $\ell \rightarrow \infty$ for $\alpha > 2D + 1$. This leads to the finite Lieb–Robinson velocity v_1 for the unitary operator $e^{-i(H_1+H_2)t}$.

In summary, the spread of the operator changes the effective one-site energy by t^D times (see Eq. (9)), which yields the condition of $\alpha > 2D + 1$ for the linear light cone. In our proof for the general Hamiltonian (1), we decompose the total length scale into pieces and consider the multi-unitary decomposition by generalizing Eq. (8) (see also Appendix VIII). We then obtain the Lieb–Robinson bound for each of the decomposed unitary operators and connect them into a single Lieb–Robinson bound. The technical difficulties lie in that we




FIG. 3. Schematic of our state-transfer protocol. We show many copies of the input state here and transform the receiver into the GHZ state. The protocol effectively amplifies the long-range interaction strength between the sender and the receiver by t^{2D} times ($D = 1$). We can see that this simple protocol gives an example that realizes the intuitive picture in Fig. 2. We here consider 1D systems, but the generalization to high-dimensional systems is straightforward.

need to connect infinitely many Lieb–Robinson bounds; in the step-by-step connections, a simple estimation makes the Lieb–Robinson velocity diverge rapidly, and hence, highly refined analyses are required to obtain a finite velocity.

V. OPTIMALITY OF THE PRESENT LIEB–ROBINSON BOUND

We have proved that the condition $\alpha > 2D+1$ is a sufficient condition for arbitrary Hamiltonians to have the linear-light cone. We here discuss whether this condition can be further improved. It has been conjectured in previous studies [49, 94] that the best general condition may be given by $\alpha > D+1$ from numerical and theoretical analyses of specific models. Against the conventional expectation, we show that any improvement from $\alpha > 2D+1$ is impossible as long as we consider the general long-range interacting systems (1). In the following, we explicitly provide a quantum-state transfer protocol that achieves a nonlinear light cone for $\alpha < 2D+1$.

We follow a similar setup as in Ref. [2] and consider the quantum-state transfer between two separated spins A and B through a spin network, where we define R as the distance between the spins A and B (Fig. 3). We start from the initial state $|\psi\rangle$ such that all spins are given by $|0\rangle$, namely $|\psi\rangle := |0\rangle^{\otimes n}$. We then apply the unitary operation U_0 or U_1 to the spin A , where U_0 is the identity operator and U_1 is the spin flip operator (i.e., $U_1 = |0\rangle\langle 1| + |1\rangle\langle 0|$). After applying U_0 or U_1 , the

initial quantum state is transformed into $|\psi_s\rangle = |s\rangle \otimes |0\rangle^{\otimes n-1}$ with $s = 0, 1$. After time t , the quantum state evolves according to the unitary time operation $U(t) = \mathcal{T}[e^{-i \int_0^t H(\tau) d\tau}]$. We then define the output state for spin B as

$$\rho_B^{(s)} := \text{tr}_{B^c} [U(t)|\psi_s\rangle\langle\psi_s|U(t)^\dagger]. \quad (10)$$

If we can distinguish between the state $\rho_B^{(0)}$ from $\rho_B^{(1)}$ with the probability 1, we can achieve the perfect quantum state transfer.

By utilizing the Lieb–Robinson bound (6), we have [2]

$$\|\rho_B^{(1)} - \rho_B^{(0)}\|_1 \lesssim |t|^{2D+1} R^{-\alpha}, \quad (11)$$

where $\|\cdot\|_1$ is the trace norm. When $\|\rho_B^{(1)} - \rho_B^{(0)}\|_1 = 2$, the two states $\rho_B^{(0)}$ and $\rho_B^{(1)}$ are orthogonal to each other and completely distinguishable. We prove that the upper bound of (11) is achievable by combining controlled-NOT-type short-range interactions and the Ising-type long-range interactions.

We here consider a one-dimensional (1/2)-spin system and decompose the time-evolution into three steps (see Fig. 3). In each of the steps, we take a time of $\mathcal{O}(t)$ (e.g., $t/3$). In the first step, we copy the state of the spin A using the controlled NOT (CNOT) gate operation, which is generated by a simple bipartite interaction [94]. We define the subset L_A as the spins that are the same state as that of spin A (see Fig. 3). We consider the dynamics by short-range interactions, and the number of spins in the subset L_A is thus upper-bounded by $\mathcal{O}(t)$. At the same time, we prepare the Greenberger-Horne-Zeilinger (GHZ) state that includes the spin B . The GHZ state is also generated by combining the rotation of the spin B and the CNOT gate operation. We define the subset L_B as the spins that are involved in the GHZ state. Owing to the short-rangeness of the interactions, the number of spins in L_B is also of $\mathcal{O}(t)$.

In the second step, we apply long-range Ising interactions by the Hamiltonian

$$H_{\text{Ising}} = \frac{g}{R^\alpha} \sum_{i \in L_A} \sum_{j \in L_B} \sigma_i^z \otimes \sigma_j^z, \quad (12)$$

with $g \leq g_0$. Because the states of spins L_A are given by $|0\cdots 0\rangle$ or $|1\cdots 1\rangle$, the unitary time evolution by H_{Ising} only changes the phase factor of the GHZ state of spins L_B ; that is, the GHZ state $(|00\cdots 0\rangle + |11\cdots 1\rangle)/\sqrt{2}$ is rotated as $(|00\cdots 0\rangle + e^{\pm 2i\theta}|11\cdots 1\rangle)/\sqrt{2}$ with \pm depending on the states of spins L_A . Simple and straightforward calculations can give the phase shift θ as $\theta = g\mathcal{O}(t)|L_A| \cdot |L_B|R^{-\alpha}$. Then, if θ is taken as $\theta = \pi/4$, the rotated GHZ states $(|00\cdots 0\rangle + e^{\pm 2i\theta}|11\cdots 1\rangle)/\sqrt{2}$ are mutually orthogonal.

In the final step, we untangle the rotated GHZ state and concentrate the phase term on the spin B , which transforms this state to $|00\cdots 0\rangle \otimes (|0\rangle + e^{\pm 2i\theta}|1\rangle)/\sqrt{2}$. This procedure is also performed by the CNOT gate operation; hence, we only need the short-range interactions. In this protocol, we obtain the lower bound of

$$\|\rho_B^{(1)} - \rho_B^{(0)}\|_1 = 2 \sin(2\theta) \gtrsim t|L_A| \cdot |L_B|R^{-\alpha}. \quad (13)$$

Considering that $|L_A| = |L_B| = \mathcal{O}(t)$, we can achieve the theoretical upper bound of (6) with $D = 1$. Thus, as long as $\alpha < 3$, the information can reach a distance of $R = \mathcal{O}(t^{3/\alpha})$. This protocol can be generalized to high-dimensional setups, and we obtain the same lower bound as (13), where we have $|L_A| = |L_B| = \mathcal{O}(t^D)$. Then, the shape of the light cone becomes $t^{(2D+1)/\alpha}$.

This simple quantum model (i.e., two-body interaction and $(1/2)$ -spin systems) already saturates the Lieb–Robinson bound of (6); hence, our condition $\alpha > 2D + 1$ for the linear light cone cannot be improved unless we consider a special class of Hamiltonians. At the same time of our submission, a similar protocol to achieve the Lieb–Robinson bound (6) was given [115] with a more explicit lower bound on the commutator $\|[O_i(t), O_j]\|$.

VI. SUMMARY AND DISCUSSION

In this study, we proved the existence of the linear light cone (see (5) for the definition) in general long-range interacting systems, where the interaction decays as according to the relation $R^{-\alpha}$ ($\alpha > 2D + 1$) with respect to the distance R . Our Lieb–Robinson bound in (6) provided an approximate commutation relation as $\|[O_i(t), O_j]\| \lesssim t^{2D+1}(R - \bar{v}t)^{-\alpha}$, with rapid decay beyond $r \gtrsim \bar{v}t$. Moreover, the error of the local approximation for $O_i(t)$ was estimated as in (7), namely $\|O_i(t) - O_i(t, i[R])\| \lesssim t^{D+1}(R - \bar{v}t)^{-\alpha+D}$. Our result was obtained for Hamiltonians with two-body interactions but can be extended to a more general setup (see Eq. (14) with (15) in Appendix VII), where even the few-body interactions are not assumed. We also show an explicit example that our Lieb–Robinson bound is saturated for $\alpha > 2D + 1$, and the linearity of the light cone deteriorates for $\alpha < 2D + 1$. Therefore, our condition for the linear light cone is optimal as long as we consider the general class of Hamiltonians. Although we consider Hamiltonian dynamics throughout this work, we expect that the same analysis can be applied to generic Markovian dynamics using the procedures in Refs. [18, 116].

We finally present an open question. In the present work, although we provided the optimal Lieb–Robinson bound for $\alpha > 2D + 1$, it is still unclear what can be obtained in the $\alpha \leq 2D + 1$ regimes. The most important problem is to identify the regime of the exponent α that ensures the polynomial (or superlinear) light cone. A state-of-the-art analysis [49] has defined the polynomial light cone in the form $r = t^{(\alpha-D)/(\alpha-2D)}$ for $\alpha > 2D$. In contrast, the super-polynomial light cone has been explicitly shown for $\alpha \leq D$ [94]. In tackling this problem, the simplest case with only two length scales as in Eq. (8) may be a good starting point. We expect that our present analysis will provide a better polynomial light cone for high-dimensional systems.

ACKNOWLEDGMENTS

T.K. was supported by the RIKEN Center for AIP and JSPS KAKENHI grant no. 18K13475. TK gives thanks to God for his wisdom. K.S. was supported by

JSPS Grants-in-Aid for Scientific Research (grant nos. JP16H02211 and JP19H05603).

VII. FORMAL EXPRESSION OF THE THEOREM

We demonstrate our theorem here in a general manner. First, some necessary notations are provided; for arbitrary subsystems $X, Y \subset \Lambda$, we define $d_{X,Y}$ as the shortest path length on the lattice that connects X and Y . If $X \cap Y \neq \emptyset$, $d_{X,Y} = 0$. For a subset $X \subseteq \Lambda$, we define $\text{diam}(X) := \max_{i,j \in X} (d_{i,j}) + 1$, the cardinality $|X|$ as the number of vertices contained in X , and the complementary subset of X as $X^c := \Lambda \setminus X$.

We consider a general class of the Hamiltonian beyond the two-body interaction (1) as

$$H = \sum_{Z \subseteq \Lambda} h_Z, \quad (14)$$

where each of the interaction terms $\{h_Z\}_{Z \subseteq \Lambda}$ acts on the sites in $Z \subseteq \Lambda$. Notably, we *do not assume* the few-body interaction here, i.e., $|Z|$ can be arbitrarily large up to $|Z| = n$. Therefore, the Hamiltonian H includes macroscopic interactions such as $\sigma_1^z \otimes \sigma_2^z \otimes \cdots \otimes \sigma_n^z$. However, the assumption (15) below restricts the amplitude of such interactions as $\text{poly}(1/n)$. Here, the only assumption is the following power-law decay of the interactions:

$$\begin{aligned} \sup_{i \in \Lambda} \sum_{Z: Z \ni i, \text{diam}(Z) \geq r} \|h_Z\| &\leq gr^{-\alpha+D}, \\ \sup_{i,j \in \Lambda} \sum_{Z: Z \ni \{i,j\}} \|h_Z\| &\leq g_0(d_{i,j} + 1)^{-\alpha} \end{aligned} \quad (15)$$

with

$$\alpha > 2D + 1 \quad (16)$$

for an arbitrary site pair of $\{i, j\} \subset \Lambda$. Here, $\sum_{Z: Z \ni \{i,j\}}$ denotes the summation that encompasses all the interaction terms $\{h_Z\}_{Z \subseteq \Lambda}$, including the sites i and j , and $\|\cdots\|$ is the operator norm. By considering an appropriate energy unit, we set $g = 1$.

To formulate our main theorem, we first define the coarse-grained subsets (see Fig. 6). For a subset $X \subseteq \Lambda$, we first define $X[r]$ as the extended subset:

$$X[r] := \{i \in \Lambda | d_{X,i} \leq r\}, \quad (17)$$

where $X[0] = X$ and r is an arbitrary positive number. We also define the coarse-grained total set $\Lambda^{(\xi)}$ as the minimum subset such that $\Lambda^{(\xi)}[\xi] = \Lambda$, namely

$$\Lambda^{(\xi)} := \arg \min_{Z \subseteq \Lambda | Z[\xi] = \Lambda} |Z|, \quad (18)$$

where $\Lambda^{(0)} = \Lambda$. Similarly, for an arbitrary subset $X \subseteq \Lambda$, we define $X^{(\xi)} \subseteq \Lambda^{(\xi)}$ as follows:

$$X^{(\xi)} := \arg \min_{Z \subseteq \Lambda^{(\xi)} | Z[\xi] \supset X} |Z|, \quad (19)$$

where $X^{(0)} = X$. From the definition, the cardinality of the subset $X^{(\xi)}$ is roughly $(1/\xi)^D$ times that of the original, namely $|X^{(\xi)}| \approx (1/\xi)^D |X|$.




FIG. 4. (a) Extended subset. The subset $X[r]$ is defined by extending the original subset X by a distance r . (b) The subset $\Lambda^{(\xi)} \subseteq \Lambda$ is the coarse grained lattice (orange sites) which is defined as the minimum subset such that $\Lambda^{(\xi)}[\xi] = \Lambda$. (c) For an arbitrary subset $X \subseteq \Lambda$ (pink region), $X^{(\xi)} \subseteq \Lambda^{(\xi)}$ is the coarse grained subset (red sites) which is defined as the minimum subset such that $X^{(\xi)}[\xi] \supseteq X$. The cardinality of the subset $X^{(\xi)}$ is roughly $(1/\xi)^D$ times as the original one, namely $|X^{(\xi)}| \approx (1/\xi)^D |X|$.

Using the notation $X^{(\xi)}$, we state our main theorem as follows:

Main theorem. *Let us consider the long-range interacting Hamiltonian H of the form (14) with the assumption (15). For $|t| \geq 1$, this Hamiltonian H satisfies the Lieb–Robinson bound for arbitrary operators O_X and O_Y that are supported on $X \subseteq \Lambda$ and $Y \subseteq \Lambda$, respectively:*

$$\frac{\|[O_X(t), O_Y]\|}{\|O_X\| \cdot \|O_Y\|} \leq C_H |X^{(\bar{v}|t)}| \cdot |Y^{(\bar{v}|t)}| \frac{|t|^{2D+1} \log^{2D}(x+1)}{(x - \bar{v}|t|)^\alpha}$$

and

$$\frac{\|[O_X(t), O_Y]\|}{\|O_X\| \cdot \|O_Y\|} \leq C'_H |X^{(\bar{v}|t)}|^2 \frac{|t|^{D+1} \log^{2D}(x+1)}{(x - \bar{v}|t|)^{\alpha-D}},$$

where C_H , C'_H , and \bar{v} are constants that depend only on parameters $\{D, g_0, \alpha\}$ and a geometric constant that is determined by the lattice structure. Note that we use the notation from Eq. (19) with $\xi = \bar{v}|t|$.

Here, the coefficient $\log^{2D}(x+1)$ exists because of a technical reason, which results from the macroscopic interactions in Eq. (14) (e.g., $\mathcal{O}(n)$ -body interactions). If we restrict ourselves to the few-body (or k -local) Hamiltonians with $k = \mathcal{O}(1)$, namely

$$H = \sum_{Z \subseteq \Lambda, |Z| \leq k} h_Z, \quad (20)$$

the Lieb–Robinson bound is slightly improved as follows:

Main theorem (k -local Hamiltonians). *Let us consider the long-range interacting Hamiltonian H of the form (20) with the assumption (15). For $|t| \geq 1$, this Hamiltonian H satisfies the Lieb–Robinson bound as follows:*

$$\frac{\|[O_X(t), O_Y]\|}{\|O_X\| \cdot \|O_Y\|} \leq C_H^{(k)} |X^{(\bar{v}^{(k)}|t)}| \cdot |Y^{(\bar{v}^{(k)}|t)}| \frac{|t|^{2D+1}}{(x - \bar{v}^{(k)}|t|)^\alpha}$$

and

$$\frac{\|[O_X(t), O_Y]\|}{\|O_X\| \cdot \|O_Y\|} \leq C_H^{(k)'} |X^{(\bar{v}^{(k)}|t)}|^2 \frac{|t|^{D+1}}{(x - \bar{v}^{(k)}|t|)^{\alpha-D}},$$

where $C_H^{(k)}$, $C_H^{(k)'}$ and $\bar{v}^{(k)}$ are constants that depend only on the parameters $\{D, g_0, \alpha, k\}$ and a geometric constant that is determined by the lattice structure.

The first and second inequalities in the above theorem reduce to the inequalities (6) and (7) in the main part, respectively; in deriving the inequality (7), we use the discussion in Ref. [2] to relate the commutator inequality to the local approximation.

VIII. SKETCH OF THE PROOF

We herein show the essential ideas, and further details are provided in the supplementary materials [117]. For the proof, we first decompose the length scale into $\leq \ell_t$ and $> \ell_t$ for a fixed t and consider the following decomposition for the total Hamiltonian: $H = H_{\leq \ell_t} + H_{> \ell_t}$, where we define $H_{\leq \ell}$ for arbitrary $\ell \in \mathbb{N}$ as the operator that includes all the interaction terms whose length scales are less than ℓ : $H_{\leq \ell} = \sum_{\text{diam}(Z) \leq \ell} h_Z$. In the case where the length ℓ is short range or $\ell = \mathcal{O}(1)$, the Hamiltonian $H_{\leq \ell}$ provides the Lieb–Robinson bound with a finite velocity. However, we consider the case of $\ell = \ell_t$ with ℓ_t depending on time t here. In the following computations, we choose $\ell_t = |t|^{\tilde{\eta}}$ with $\tilde{\eta} := 1 - \frac{\alpha-2D-1}{2(\alpha-D)} < 1$. When the length scale ℓ is in the middle-range between $\ell = \mathcal{O}(t^0)$ and $\ell = \infty$, it is no longer considered trivial if the light-cone for the dynamics by $H_{\leq \ell_t}$ is retained.

To obtain the Lieb–Robinson bound for $H_{\leq \ell}$ with a generic ℓ , we further decompose $H_{\leq \ell}$ as follows (Fig. 5):

$$H_{\leq \ell} = \sum_{q=1}^{q^*} H_q, \quad H_q := \sum_{\ell_{q-1} \leq \text{diam}(Z) < \ell_q} h_Z.$$

We define a set of length scales $\{\ell_q\}_{q=1}^{q^*}$ as $\ell_q = \ell_{q-1}^{1+\eta_q}$ with $\eta_q > 0$, where $\{\eta_q\}_{q=2}^{q^*}$ is appropriately chosen such that $\{\ell_q\}_{q=1}^{q^*}$ is an integer and there exists an integer $q^* \in \mathbb{N}$ satisfying $\ell_{q^*} = \ell_1^{(1+\eta_2)(1+\eta_3)\cdots(1+\eta_{q^*})} = \ell$. In this case, ℓ_q increases by the double exponential function with respect to q .




FIG. 5. Decomposition of the length scale. We first decompose the total Hamiltonian into two regimes: $H_{\leq \ell_t}$ and $H_{> \ell_t}$. The Hamiltonian $H_{\leq \ell_t}$ includes all the interactions up to the length scale ℓ_t that is dependent on time, and $H_{> \ell_t}$ includes the other interactions. To obtain the middle-range Lieb–Robinson bound, we further decompose the range $[1, \ell_t]$ into q^* segments. We start from the Hamiltonian that only includes the length scale ℓ_1 and iteratively consider the increasing length scales.

In deriving the Lieb–Robinson bound for $e^{-iH_{\leq \ell_t} t}$, we iteratively take in the increasing length scales. We begin with the unitary operator $e^{-iH_1 t}$; as long as ℓ_1 is independent of t , the Lieb–Robinson bound for $e^{-iH_1 t}$ is in the shorter range with a velocity of $v_1 = \mathcal{O}(1)$. Then, using this Lieb–Robinson bound, we derive a new Lieb–Robinson bound for $e^{-i(H_1+H_2)t} = e^{-iH_{1:2}t}$, where we define $H_{1:q} := \sum_{s=1}^q H_s$ with $1 \leq q \leq q^*$. This process is repeated to extend the length scales for $\ell_1 \rightarrow \ell_2 \rightarrow \dots \rightarrow \ell_{q^*} = \ell$. In each of these steps, based on the Lieb–Robinson bound for $e^{-iH_{1:q-1}t}$, we update the Lieb–Robinson bound for $e^{-iH_{1:q}t}$.

In the first update, we start from the following decomposition of the unitary operator $e^{-iH_{1:2}t}$:

$$U_{2,t} := e^{-i(H_1+H_2)t} = e^{-iH_1 t} \mathcal{T} e^{-\int_0^t H_2(H_1, \tau) d\tau}.$$

The operator spreading by $e^{-iH_1 t}$ is of the order of $\mathcal{O}(v_1 t)$; hence, as long as $t \leq \Delta t_2 \approx \ell_2/v_1$, $H_2(H_1, \tau)$ ($\tau \leq \Delta t_2$) has the same interaction length as the original, namely ℓ_2 . Thus, we can obtain the Lieb–Robinson bound with a linear light cone for $\mathcal{T} e^{-\int_0^{\Delta t_2} H_2(H_1, \tau) d\tau}$. For any unspecified time t , we consider $U_{2,t} = U_{2, \Delta t_2}^m U_{2, \Delta t'_2}$, with $t = m\Delta t_2 + \Delta t'_2$, where $\Delta t_2 \propto \ell_2/v_1$ and $\Delta t'_2 < \Delta t_2$. By appropriately connecting all the Lieb–Robinson bounds, this results in a Lieb–Robinson bound of the following form:

$$\|[O_X(H_{\leq \ell_2}, t), O_Y]\| \lesssim (1 + x/\ell_2)^{D-1} e^{-2(x-v_2|t|)/\ell_2}$$

with $x = d_{X,Y}$, where the velocity $v_2 > v_1$ is upper-bounded using the Lieb–Robinson velocity v_1 for $e^{-iH_1 t}$.

For general $q-1$, we define v_{q-1} as the Lieb–Robinson velocity for $e^{-iH_{1:q-1}t}$ and analyze the unitary operator $U_{q,t} := e^{-iH_{1:q}t}$, which we decompose as $U_{q,t} = U_{q, \Delta t_q}^m U_{q, \Delta t'_q}$ with

$$U_{q, \Delta t_q} = e^{-iH_{1:q-1} \Delta t_q} \mathcal{T} e^{-\int_0^{\Delta t_q} H_q(H_{1:q-1}, \tau) d\tau}$$

and $t = m\Delta t_q + \Delta t'_q$; here, $\Delta t_q \propto \ell_q/v_{q-1}$ and $\Delta t'_q < \Delta t_q$. From the choice of Δt_q , we can ensure that $H_q(H_{1:q-1}, \tau)$ has the same interaction length as that of H_q , namely ℓ_q . We then obtain the Lieb–Robinson bound for $H_{1:q}$ as follows:

$$\|[O_X(H_{1:q}, t), O_Y]\| \lesssim (1 + x/\ell_q)^{D-1} e^{-2(x-v_q|t|)/\ell_q}$$

where v_q depends on the v_{q-1} . Thus, we iteratively estimate the Lieb–Robinson velocity v_q using v_{q-1} . Further, we can derive the following recursion relation:

$$v_q = v_{q-1} \left(1 + \frac{c \log(\ell_q)}{\ell_{q-1}^\eta} + \frac{c'}{\log(\ell_{q-1})} \right),$$

where $\eta = \sqrt{1 + \frac{\alpha - (2D+1)}{D+2}} - 1$ and c, c' are constants. We provide the explicit form in [117]. The length scale ℓ_q is now lower-bounded by a double exponential function with respect to q . Therefore, $\lim_{q \rightarrow \infty} v_q$ converges to a constant v^* , and we obtain the following “middle-range Lieb–Robinson bound” for $e^{-iH_{\leq \ell} t}$:

$$\|[O_X(H_{\leq \ell}, t), O_Y]\| \lesssim (1 + x/\ell)^{D-1} e^{-2(x-v^*|t|)/\ell}. \quad (21)$$

In this manner, we can ensure that $H_{\leq \ell}$ retains the linear light cone for $\ell \lesssim t$ from (21), whereas no such confirmation is possible for $\ell \gtrsim t$. Thus, we consider the case of $\ell = \ell_t$ and decompose the total time evolution as

$$e^{-iHt} = e^{-iH_{\leq \ell_t} t} U_{> \ell_t},$$

where $U_{> \ell_t} := \mathcal{T} e^{-i \int_0^t H_{> \ell_t}(H_{\leq \ell_t}, \tau) d\tau}$. In order to estimate the quasi-locality of the interaction for $H_{> \ell_t}(H_{\leq \ell_t}, \tau)$, we apply the middle-range Lieb–Robinson bound (21). Based on the quasi-locality, we utilize the standard recursion approach [14, 15] to obtain the Lieb–Robinson bound for $U_{> \ell_t}$. After intricate calculations, we obtain the Lieb–Robinson bound for $U_{> \ell_t}$ as

$$\|[U_{> \ell_t}^\dagger O_X U_{> \ell_t}, O_Y]\| \lesssim \frac{|t|^{2D+1} \log^{2D}(x+1)}{(x - \kappa_0 v^* |t|)^\alpha}, \quad (22)$$

where κ_0 is a constant. We then connect the two Lieb–Robinson bounds for $e^{-iH_{\leq \ell_t} t}$ (21) and $U_{> \ell_t}$ (22) to derive the total Lieb–Robinson bound in the main theorem.

-
- [1] Elliott H. Lieb and Derek W. Robinson, “*The finite group velocity of quantum spin systems*,” *Communications in Mathematical Physics* **28**, 251–257 (1972).
 - [2] S. Bravyi, M. B. Hastings, and F. Verstraete, “*Lieb-Robinson Bounds and the Generation of Correlations and Topological Quantum Order*,” *Phys. Rev. Lett.* **97**, 050401 (2006).
 - [3] Jens Eisert and Tobias J. Osborne, “*General Entanglement Scaling Laws from Time Evolution*,” *Phys. Rev. Lett.* **97**, 150404 (2006).
 - [4] Maurizio Fagotti and Pasquale Calabrese, “*Evolution of entanglement entropy following a quantum quench: Analytic results for the XY chain in a transverse magnetic field*,” *Phys. Rev. A* **78**, 010306 (2008).
 - [5] Alioscia Hamma, Fotini Markopoulou, Isabeau Prémont-Schwarz, and Simone Severini, “*Lieb-Robinson Bounds and the Speed of Light from Topological Order*,” *Phys. Rev. Lett.* **102**, 017204 (2009).
 - [6] Lars Bonnes, Fabian H. L. Essler, and Andreas M. Läuchli, “*“Light-Cone” Dynamics After Quantum Quenches in Spin Chains*,” *Phys. Rev. Lett.* **113**, 187203 (2014).
 - [7] Andreas M. Läuchli and Corinna Kollath, “*Spreading of correlations and entanglement after a quench in the one-dimensional Bose-Hubbard model*,” *Journal of Statistical Mechanics: Theory and Experiment* **2008**, P05018 (2008).
 - [8] Christian K. Burrell, Jens Eisert, and Tobias J. Osborne, “*Information propagation through quantum chains with fluctuating disorder*,” *Phys. Rev. A* **80**, 052319 (2009).
 - [9] J. Jünemann, A. Cadarso, D. Pérez-García, A. Bermudez, and J. J. García-Ripoll, “*Lieb-Robinson Bounds for Spin-Boson Lattice Models and Trapped Ions*,” *Phys. Rev. Lett.* **111**, 230404 (2013).
 - [10] Jean-Sébastien Bernier, Ryan Tan, Lars Bonnes, Chu Guo, Dario Poletti, and Corinna Kollath, “*Light-Cone and Diffusive Propagation of Correlations in a Many-Body Dissipative System*,” *Phys. Rev. Lett.* **120**, 020401 (2018).
 - [11] David Damanik, Marius Lemm, Milivoje Lukic, and William Yessen, “*New Anomalous Lieb-Robinson Bounds in Quasiperiodic XY Chains*,” *Phys. Rev. Lett.* **113**, 127202 (2014).
 - [12] J. Eisert and D. Gross, “*Supersonic Quantum Communication*,” *Phys. Rev. Lett.* **102**, 240501 (2009).
 - [13] Isabeau Prémont-Schwarz, Alioscia Hamma, Israel Klich, and Fotini Markopoulou-Kalamara, “*Lieb-Robinson bounds for commutator-bounded operators*,” *Phys. Rev. A* **81**, 040102 (2010).
 - [14] Bruno Nachtergaele and Robert Sims, “*Lieb-Robinson Bounds and the Exponential Clustering Theorem*,” *Communications in Mathematical Physics* **265**, 119–130 (2006).
 - [15] Matthew B. Hastings and Tohru Koma, “*Spectral Gap and Exponential Decay of Correlations*,” *Communications in Mathematical Physics* **265**, 781–804 (2006).
 - [16] Christian K. Burrell and Tobias J. Osborne, “*Bounds on the Speed of Information Propagation in Disordered Quantum Spin Chains*,” *Phys. Rev. Lett.* **99**, 167201 (2007).
 - [17] Bruno Nachtergaele and Robert Sims, “*Lieb-Robinson bounds in quantum many-body physics*,” *Contemp. Math* **529**, 141 (2010), [arXiv:1004.2086](https://arxiv.org/abs/1004.2086).
 - [18] David Poulin, “*Lieb-Robinson Bound and Locality for General Markovian Quantum Dynamics*,” *Phys. Rev. Lett.* **104**, 190401 (2010).
 - [19] Norbert Schuch, Sarah K. Harrison, Tobias J. Osborne, and Jens Eisert, “*Information propagation for interacting-particle systems*,” *Phys. Rev. A* **84**, 032309 (2011).
 - [20] Isabeau Prémont-Schwarz and Jeff Hnybida, “*Lieb-Robinson bounds on the speed of information propagation*,” *Phys. Rev. A* **81**, 062107 (2010).
 - [21] M. B. Hastings, “*Lieb-Schultz-Mattis in higher dimensions*,” *Phys. Rev. B* **69**, 104431 (2004).
 - [22] M. B. Hastings and Xiao-Gang Wen, “*Quasiadiabatic continuation of quantum states: The stability of topological ground-state degeneracy and emergent gauge invariance*,” *Phys. Rev. B* **72**, 045141 (2005).
 - [23] M. B. Hastings, “*An area law for one-dimensional quantum systems*,” *Journal of Statistical Mechanics: Theory and Experiment* **2007**, P08024 (2007), [arXiv:0705.2024](https://arxiv.org/abs/0705.2024).
 - [24] Lluís Masanes, “*Area law for the entropy of low-energy states*,” *Phys. Rev. A* **80**, 052104 (2009).
 - [25] Fernando G. S. L. Brandão, Toby S. Cubitt, Angelo Lucia, Spyridon Michalakis, and David Perez-Garcia, “*Area law for fixed points of rapidly mixing dissipative quantum systems*,” *Journal of Mathematical Physics* **56**, 102202 (2015).
 - [26] Karel Van Acoleyen, Michaël Mariën, and Frank Verstraete, “*Entanglement Rates and Area Laws*,” *Phys. Rev. Lett.* **111**, 170501 (2013).
 - [27] Markus P. Müller, Emily Adlam, Lluís Masanes, and Nathan Wiebe, “*Thermalization and Canonical Typicality in Translation-Invariant Quantum Lattice Systems*,” *Communications in Mathematical Physics* **340**, 499–561 (2015).
 - [28] Leandro Aolita, Fernando de Melo, and Luiz Davidovich, “*Open-system dynamics of entanglement: a key issues review*,” *Reports on Progress in Physics* **78**, 042001 (2015).
 - [29] Christian Gogolin and Jens Eisert, “*Equilibration, thermalisation, and the emergence of statistical mechanics in closed quantum systems*,” *Reports on Progress in Physics* **79**, 056001 (2016).
 - [30] Eiki Iyoda, Kazuya Kaneko, and Takahiro Sagawa, “*Fluctuation Theorem for Many-Body Pure Quantum States*,” *Phys. Rev. Lett.* **119**, 100601 (2017).
 - [31] Matthew B. Hastings and Spyridon Michalakis, “*Quantization of Hall Conductance for Interacting Electrons on a Torus*,” *Communications in Mathematical Physics* **334**, 433–471 (2014).
 - [32] M. B. Hastings, “*Quasi-adiabatic continuation for disordered systems: Applications to correlations, Lieb-Schultz-Mattis, and Hall conductance*,” *arXiv preprint arXiv:1001.5280* (2010), [arXiv:1001.5280](https://arxiv.org/abs/1001.5280).
 - [33] Sergey Bravyi, Matthew B. Hastings, and Spyridon Michalakis, “*Topological quantum order: Stability under local perturbations*,” *Journal of Mathematical Physics* **51**, 093512 (2010), [http://dx.doi.org/10.1063/1.3490195](https://doi.org/10.1063/1.3490195).
 - [34] Sergey Bravyi and Matthew B. Hastings, “*A Short Proof of Stability of Topological Order under Local Perturbations*,” *Communications in Mathematical Physics* **307**, 609–627 (2011).
 - [35] Spyridon Michalakis and Justyna P. Zwolak, “*Stability of Frustration-Free Hamiltonians*,” *Communications in Mathematical Physics* **322**, 277–302 (2013).
 - [36] M. B. Hastings, “*Locality in Quantum and Markov Dynamics on Lattices and Networks*,” *Phys. Rev. Lett.* **93**, 140402 (2004).
 - [37] Senaida Hernández-Santana, Christian Gogolin, J. Ig-

- nacio Cirac, and Antonio Acín, “Correlation Decay in Fermionic Lattice Systems with Power-Law Interactions at Nonzero Temperature,” *Phys. Rev. Lett.* **119**, 110601 (2017).
- [38] M. B. Hastings, “Decay of Correlations in Fermi Systems at Nonzero Temperature,” *Phys. Rev. Lett.* **93**, 126402 (2004).
- [39] Tomotaka Kuwahara, Takashi Mori, and Keiji Saito, “Floquet-Magnus theory and generic transient dynamics in periodically driven many-body quantum systems,” *Annals of Physics* **367**, 96 – 124 (2016).
- [40] Francisco Machado, Dominic V. Else, Gregory D. Kahanamoku-Meyer, Chetan Nayak, and Norman Y. Yao, “Long-Range Prethermal Phases of Nonequilibrium Matter,” *Phys. Rev. X* **10**, 011043 (2020).
- [41] Tobias J Osborne, “Hamiltonian complexity,” *Reports on Progress in Physics* **75**, 022001 (2012).
- [42] Tobias J. Osborne, “Efficient Approximation of the Dynamics of One-Dimensional Quantum Spin Systems,” *Phys. Rev. Lett.* **97**, 157202 (2006).
- [43] Tobias J. Osborne, “Simulating adiabatic evolution of gapped spin systems,” *Phys. Rev. A* **75**, 032321 (2007).
- [44] Sergey Bravyi, David P. DiVincenzo, Daniel Loss, and Barbara M. Terhal, “Quantum Simulation of Many-Body Hamiltonians Using Perturbation Theory with Bounded-Strength Interactions,” *Phys. Rev. Lett.* **101**, 070503 (2008).
- [45] M. B. Hastings, “Quantum belief propagation: An algorithm for thermal quantum systems,” *Phys. Rev. B* **76**, 201102 (2007).
- [46] M. B. Hastings, “Light-cone matrix product,” *Journal of Mathematical Physics* **50**, 095207 (2009).
- [47] M. B. Hastings, “Observations outside the light cone: Algorithms for nonequilibrium and thermal states,” *Phys. Rev. B* **77**, 144302 (2008).
- [48] Jeongwan Haah, Matthew Hastings, Robin Kothari, and Guang Hao Low, “Quantum Algorithm for Simulating Real Time Evolution of Lattice Hamiltonians,” in *2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS)* (2018) pp. 350–360.
- [49] Minh C. Tran, Andrew Y. Guo, Yuan Su, James R. Garrison, Zachary Eldredge, Michael Foss-Feig, Andrew M. Childs, and Alexey V. Gorshkov, “Locality and Digital Quantum Simulation of Power-Law Interactions,” *Phys. Rev. X* **9**, 031006 (2019).
- [50] Wen Wei Ho, Cheryne Jonay, and Timothy H. Hsieh, “Ultrafast variational simulation of nontrivial quantum states with long-range interactions,” *Phys. Rev. A* **99**, 052332 (2019).
- [51] Daniel A. Roberts and Brian Swingle, “Lieb-Robinson Bound and the Butterfly Effect in Quantum Field Theories,” *Phys. Rev. Lett.* **117**, 091602 (2016).
- [52] Juan Maldacena, Stephen H. Shenker, and Douglas Stanford, “A bound on chaos,” *Journal of High Energy Physics* **2016**, 106 (2016).
- [53] Yingfei Gu, Xiao-Liang Qi, and Douglas Stanford, “Local criticality, diffusion and chaos in generalized Sachdev-Ye-Kitaev models,” *Journal of High Energy Physics* **2017**, 125 (2017).
- [54] Xiao Chen and Tianci Zhou, “Quantum chaos dynamics in long-range power law interaction systems,” *Phys. Rev. B* **100**, 064305 (2019).
- [55] Ivan Kukuljan, Sašo Grozdanov, and Tomaž Prosen, “Weak quantum chaos,” *Phys. Rev. B* **96**, 060301 (2017).
- [56] Adam Nahum, Jonathan Ruhman, Sagar Vijay, and Jeongwan Haah, “Quantum Entanglement Growth under Random Unitary Dynamics,” *Phys. Rev. X* **7**, 031016 (2017).
- [57] Vedika Khemani, David A. Huse, and Adam Nahum, “Velocity-dependent Lyapunov exponents in many-body quantum, semiclassical, and classical chaos,” *Phys. Rev. B* **98**, 144304 (2018).
- [58] Yong-Liang Zhang, Yichen Huang, and Xie Chen, “Information scrambling in chaotic systems with dissipation,” *Phys. Rev. B* **99**, 014303 (2019).
- [59] David J. Luitz and Yevgeny Bar Lev, “Information propagation in isolated quantum systems,” *Phys. Rev. B* **96**, 020406 (2017).
- [60] Jun Li, Ruihua Fan, Hengyan Wang, Bingtian Ye, Bei Zeng, Hui Zhai, Xinhua Peng, and Jiangfeng Du, “Measuring Out-of-Time-Order Correlators on a Nuclear Magnetic Resonance Quantum Simulator,” *Phys. Rev. X* **7**, 031011 (2017).
- [61] Silvia Pappalardi, Angelo Russomanno, Bojan Žunkovič, Fernando Iemini, Alessandro Silva, and Rosario Fazio, “Scrambling and entanglement spreading in long-range spin chains,” *Phys. Rev. B* **98**, 134303 (2018).
- [62] Daniel E. Parker, Xiangyu Cao, Alexander Avdoshkin, Thomas Scaffidi, and Ehud Altman, “A Universal Operator Growth Hypothesis,” *Phys. Rev. X* **9**, 041017 (2019).
- [63] Tianci Zhou, Shenglong Xu, Xiao Chen, Andrew Guo, and Brian Swingle, “Operator Lévy Flight: Light Cones in Chaotic Long-Range Interacting Systems,” *Phys. Rev. Lett.* **124**, 180601 (2020).
- [64] Marc Cheneau, Peter Barmettler, Dario Poletti, Manuel Endres, Peter Schauß, Takeshi Fukuhara, Christian Gross, Immanuel Bloch, Corinna Kollath, and Stefan Kuhr, “Light-cone-like spreading of correlations in a quantum many-body system,” *Nature* **481**, 484–487 (2012).
- [65] Tim Langen, Remi Geiger, Maximilian Kuhnert, Bernhard Rauer, and Joerg Schmiedmayer, “Local emergence of thermal correlations in an isolated quantum many-body system,” *Nature Physics* **9**, 640–643 (2013).
- [66] Florian Meinert, Manfred J. Mark, Emil Kirilov, Katharina Lauber, Philipp Weinmann, Michael Gröbner, Andrew J. Daley, and Hanns-Christoph Nägerl, “Observation of many-body dynamics in long-range tunneling after a quantum quench,” *Science* **344**, 1259–1262 (2014).
- [67] Sebastian Hild, Takeshi Fukuhara, Peter Schauß, Johannes Zeiher, Michael Knap, Eugene Demler, Immanuel Bloch, and Christian Gross, “Far-from-Equilibrium Spin Transport in Heisenberg Quantum Magnets,” *Phys. Rev. Lett.* **113**, 147205 (2014).
- [68] Philip Richerme, Zhe-Xuan Gong, Aaron Lee, Crystal Senko, Jacob Smith, Michael Foss-Feig, Spyridon Michalakis, Alexey V. Gorshkov, and Christopher Monroe, “Non-local propagation of correlations in quantum systems with long-range interactions,” *Nature* **511**, 198–201 (2014).
- [69] Petar Jurcevic, Ben P. Lanyon, Philipp Hauke, Cornelius Hempel, Peter Zoller, Rainer Blatt, and Christian F. Roos, “Quasiparticle engineering and entanglement propagation in a quantum many-body system,” *Nature* **511**, 202–205 (2014).
- [70] Vincent Lienhard, Sylvain de Léséleuc, Daniel Barredo, Thierry Lahaye, Antoine Browaeys, Michael Schuler, Louis-Paul Henry, and Andreas M. Läuchli, “Observing the Space- and Time-Dependent Growth of Correlations in Dynamically Tuned Synthetic Ising Models with Antiferromagnetic Interactions,” *Phys. Rev. X* **8**, 021070 (2018).
- [71] Vera Bendkowsky, Björn Butscher, Johannes Nipper, James P. Shaffer, Robert Löw, and Tilman Pfau, “Observation of ultralong-range Rydberg molecules,” *Nature* **458**, 1005 (2009).

- [72] Immanuel Bloch, Jean Dalibard, and Wilhelm Zwerger, “*Many-body physics with ultracold gases*,” *Rev. Mod. Phys.* **80**, 885–964 (2008).
- [73] M. Saffman, T. G. Walker, and K. Mølmer, “*Quantum information with Rydberg atoms*,” *Rev. Mod. Phys.* **82**, 2313–2363 (2010).
- [74] Bo Yan, Steven A Moses, Bryce Gadway, Jacob P Covey, Kaden RA Hazzard, Ana Maria Rey, Deborah S Jin, and Jun Ye, “*Observation of dipolar spin-exchange interactions with lattice-confined polar molecules*,” *Nature* **501**, 521 (2013).
- [75] K. Aikawa, A. Frisch, M. Mark, S. Baier, A. Rietzler, R. Grimm, and F. Ferlaino, “*Bose-Einstein Condensation of Erbium*,” *Phys. Rev. Lett.* **108**, 210401 (2012).
- [76] Joseph W Britton, Brian C Sawyer, Adam C Keith, C-C Joseph Wang, James K Freericks, Hermann Uys, Michael J Biercuk, and John J Bollinger, “*Engineered two-dimensional Ising interactions in a trapped-ion quantum simulator with hundreds of spins*,” *Nature* **484**, 489 (2012).
- [77] R. Islam, C. Senko, W. C. Campbell, S. Korenblit, J. Smith, A. Lee, E. E. Edwards, C.-C. J. Wang, J. K. Freericks, and C. Monroe, “*Emergence and Frustration of Magnetism with Variable-Range Interactions in a Quantum Simulator*,” *Science* **340**, 583–587 (2013).
- [78] Hannes Bernien, Sylvain Schwartz, Alexander Keesling, Harry Levine, Ahmed Omran, Hannes Pichler, Soonwon Choi, Alexander S Zibrov, Manuel Endres, Markus Greiner, *et al.*, “*Probing many-body dynamics on a 51-atom quantum simulator*,” *Nature* **551**, 579 (2017), article.
- [79] Jiehang Zhang, Guido Pagano, Paul W Hess, Antonis Kyprianidis, Patrick Becker, Harvey Kaplan, Alexey V Gorshkov, Z-X Gong, and Christopher Monroe, “*Observation of a many-body dynamical phase transition with a 53-qubit quantum simulator*,” *Nature* **551**, 601 (2017).
- [80] Brian Neyenhuis, Jiehang Zhang, Paul W. Hess, Jacob Smith, Aaron C. Lee, Phil Richerme, Zhe-Xuan Gong, Alexey V. Gorshkov, and Christopher Monroe, “*Observation of prethermalization in long-range interacting spin chains*,” *Science Advances* **3** (2017), 10.1126/sciadv.1700672.
- [81] Davide Vodola, Luca Lepori, Elisa Ercolessi, Alexey V. Gorshkov, and Guido Pupillo, “*Kitaev Chains with Long-Range Pairing*,” *Phys. Rev. Lett.* **113**, 156402 (2014).
- [82] Thomas Koffel, M. Lewenstein, and Luca Tagliacozzo, “*Entanglement Entropy for the Long-Range Ising Chain in a Transverse Field*,” *Phys. Rev. Lett.* **109**, 267203 (2012).
- [83] Mohammad F. Maghrebi, Zhe-Xuan Gong, and Alexey V. Gorshkov, “*Continuous Symmetry Breaking in 1D Long-Range Interacting Quantum Systems*,” *Phys. Rev. Lett.* **119**, 023001 (2017).
- [84] Tomotaka Kuwahara, Itai Arad, Luigi Amico, and Vlatko Vedral, “*Local reversibility and entanglement structure of many-body ground states*,” *Quantum Science and Technology* **2**, 015005 (2017).
- [85] Tomotaka Kuwahara, “*Asymptotic behavior of macroscopic observables in generic spin systems*,” *Journal of Statistical Mechanics: Theory and Experiment* **2016**, 053103 (2016).
- [86] Tomotaka Kuwahara and Keiji Saito, “*Gaussian concentration bound and Ensemble equivalence in generic quantum many-body systems including long-range interaction*,” arXiv preprint arXiv:1906.10872 (2019), arXiv:1906.10872.
- [87] Tomotaka Kuwahara and Keiji Saito, “*Area law of non-critical ground states in 1D long-range interacting systems*,” arXiv preprint arXiv:1908.11547 (2019), arXiv:1908.11547.
- [88] Jens Eisert, Mauritz van den Worm, Salvatore R. Manmana, and Michael Kastner, “*Breakdown of Quasilocality in Long-Range Quantum Lattice Models*,” *Phys. Rev. Lett.* **111**, 260401 (2013).
- [89] Zhe-Xuan Gong, Michael Foss-Feig, Spyridon Michalakis, and Alexey V. Gorshkov, “*Persistence of Locality in Systems with Power-Law Interactions*,” *Phys. Rev. Lett.* **113**, 030602 (2014).
- [90] David Métivier, Romain Bachelard, and Michael Kastner, “*Spreading of Perturbations in Long-Range Interacting Classical Lattice Models*,” *Phys. Rev. Lett.* **112**, 210601 (2014).
- [91] P. Hauke and L. Tagliacozzo, “*Spread of Correlations in Long-Range Interacting Quantum Systems*,” *Phys. Rev. Lett.* **111**, 207202 (2013).
- [92] M. Pino, “*Entanglement growth in many-body localized systems with long-range interactions*,” *Phys. Rev. B* **90**, 174204 (2014).
- [93] M. Ghasemi Nezhadhighi and M. A. Rajabpour, “*Entanglement dynamics in short- and long-range harmonic oscillators*,” *Phys. Rev. B* **90**, 205438 (2014).
- [94] Zachary Eldredge, Zhe-Xuan Gong, Jeremy T. Young, Ali Hamed Moosavian, Michael Foss-Feig, and Alexey V. Gorshkov, “*Fast Quantum State Transfer and Entanglement Renormalization Using Long-Range Interactions*,” *Phys. Rev. Lett.* **119**, 170503 (2017).
- [95] Lorenzo Cevolani, Giuseppe Carleo, and Laurent Sanchez-Palencia, “*Spreading of correlations in exactly solvable quantum models with long-range interactions in arbitrary dimensions*,” *New Journal of Physics* **18**, 093002 (2016).
- [96] Lea F. Santos, Fausto Borgonovi, and Giuseppe Luca Celardo, “*Cooperative Shielding in Many-Body Systems with Long-Range Interaction*,” *Phys. Rev. Lett.* **116**, 250402 (2016).
- [97] G. L. Celardo, R. Kaiser, and F. Borgonovi, “*Shielding and localization in the presence of long-range hopping*,” *Phys. Rev. B* **94**, 144206 (2016).
- [98] L Lepori, A Trombettoni, and D Vodola, “*Singular dynamics and emergence of nonlocality in long-range quantum models*,” *Journal of Statistical Mechanics: Theory and Experiment* **2017**, 033102 (2017).
- [99] Lorenzo Cevolani, Julien Despres, Giuseppe Carleo, Luca Tagliacozzo, and Laurent Sanchez-Palencia, “*Universal scaling laws for correlation spreading in quantum systems with short- and long-range interactions*,” *Phys. Rev. B* **98**, 024302 (2018).
- [100] Rajeev Singh, Roderich Moessner, and Dibyendu Roy, “*Effect of long-range hopping and interactions on entanglement dynamics and many-body localization*,” *Phys. Rev. B* **95**, 094205 (2017).
- [101] X. Deng, V. E. Kravtsov, G. V. Shlyapnikov, and L. Santos, “*Duality in Power-Law Localization in Disordered One-Dimensional Systems*,” *Phys. Rev. Lett.* **120**, 110602 (2018).
- [102] David J. Luitz and Yevgeny Bar Lev, “*Emergent locality in systems with power-law interactions*,” *Phys. Rev. A* **99**, 010105 (2019).
- [103] Benedikt Kloss and Yevgeny Bar Lev, “*Spin transport in a long-range-interacting spin chain*,” *Phys. Rev. A* **99**, 032114 (2019).
- [104] Shuji Tamaki and Keiji Saito, “*Energy current correlation in solvable long-range interacting systems*,” *Phys. Rev. E* **101**, 042118 (2020).
- [105] Tomotaka Kuwahara, “*Exponential bound on information spreading induced by quantum many-body dynamics with long-range interactions*,” *New Journal of Physics* **18**, 053034 (2016).

- [106] Ryan Sweke, Jens Eisert, and Michael Kastner, “*Lieb–Robinson bounds for open quantum systems with long-ranged interactions*,” *Journal of Physics A: Mathematical and Theoretical* **52**, 424003 (2019).
- [107] Andrew Y. Guo, Minh C. Tran, Andrew M. Childs, Alexey V. Gorshkov, and Zhe-Xuan Gong, “*Signaling and scrambling with strongly long-range interactions*,” *Phys. Rev. A* **102**, 010401 (2020).
- [108] Michael Foss-Feig, Zhe-Xuan Gong, Charles W. Clark, and Alexey V. Gorshkov, “*Nearly Linear Light Cones in Long-Range Interacting Quantum Systems*,” *Phys. Rev. Lett.* **114**, 157201 (2015).
- [109] Takuro Matsuta, Tohru Koma, and Shu Nakamura, “*Improving the Lieb–Robinson Bound for Long-Range Interactions*,” *Annales Henri Poincaré* **18**, 519–528 (2017).
- [110] Dominic V. Else, Francisco Machado, Chetan Nayak, and Norman Y. Yao, “*Improved Lieb–Robinson bound for many-body Hamiltonians with power-law interactions*,” *Phys. Rev. A* **101**, 022333 (2020).
- [111] Minh C. Tran, Adam Ehrenberg, Andrew Y. Guo, Paraj Titum, Dmitry A. Abanin, and Alexey V. Gorshkov, “*Locality and heating in periodically driven, power-law-interacting systems*,” *Phys. Rev. A* **100**, 052103 (2019).
- [112] Chi-Fang Chen and Andrew Lucas, “*Finite Speed of Quantum Scrambling with Long Range Interactions*,” *Phys. Rev. Lett.* **123**, 250605 (2019).
- [113] Thierry Dauxois, Stefano Ruffo, Ennio Arimondo, and Martin Wilkens, “*Dynamics and thermodynamics of systems with long-range interactions: An introduction*,” in *Dynamics and Thermodynamics of Systems with Long-Range Interactions* (2002) pp. 1–19.
- [114] Alessandro Campa, Thierry Dauxois, and Stefano Ruffo, “*Statistical mechanics and dynamics of solvable models with long-range interactions*,” *Physics Reports* **480**, 57 – 159 (2009).
- [115] Minh C. Tran, Chi-Fang Chen, Adam Ehrenberg, Andrew Y. Guo, Abhinav Deshpande, Yifan Hong, Zhe-Xuan Gong, Alexey V. Gorshkov, and Andrew Lucas, “*Hierarchy of Linear Light Cones with Long-Range Interactions*,” *Phys. Rev. X* **10**, 031009 (2020).
- [116] Thomas Barthel and Martin Kliesch, “*Quasilocality and Efficient Simulation of Markovian Quantum Dynamics*,” *Phys. Rev. Lett.* **108**, 230504 (2012).
- [117] Supplementary material.
- [118] Tomotaka Kuwahara, “*Fundamental inequalities in quantum many-body systems*,” PhD thesis, University of Tokyo (2015).

Supplementary Material for “Strictly linear light cone long-range interacting systems of arbitrary dimensions”

Tomotaka Kuwahara^{1,2} and Keiji Saito³

¹*Mathematical Science Team, RIKEN Center for Advanced Intelligence Project (AIP), 1-4-1 Nihonbashi, Chuo-ku, Tokyo 103-0027, Japan*

²*Interdisciplinary Theoretical & Mathematical Sciences Program (iTHEMS) RIKEN 2-1, Hirosawa, Wako, Saitama 351-0198, Japan*

³*Department of Physics, Keio University, Yokohama 223-8522, Japan*

CONTENTS

II. Introduction	1
III. Main results	2
IV. Intuitive explanation of the condition $\alpha > 2D + 1$	3
V. Optimality of the present Lieb–Robinson bound	4
VI. Summary and discussion	5
acknowledgments	5
VII. Formal expression of the theorem	5
VIII. Sketch of the Proof	6
References	8
IX. Setup	13
A. Definition of the lattice	13
B. Definition of the long-range interacting Hamiltonian	13
C. Coarse grained set (Fig. 6)	14
X. Main results	15
A. Definitions	15
B. Main theorem	16
C. Proof of Theorem 1 (see also Fig. 7 for the basic outline)	17
XI. Extending the Lieb–Robinson bound for local operators to that for generic operators	20
A. Proof of Theorem 4	22
B. Proof of Proposition 5	25
1. Proof of Lemma 2	26
XII. Proof of Theorem 2: Middle-range Lieb–Robinson bound	27
A. statement	27
B. Decomposition of the length scale	29
C. Conditions for ℓ_1	29
D. Proof of Theorem 2 (see also Figs. 8 and 9)	31
E. Proof of the inequality (S.193)	36
XIII. Detailed proof of the complementary statements for the middle-range Lieb–Robinson bound	38
A. Proof of Proposition 6: Quasi-locality of operator after time-evolution	38
1. Statement	38
2. Proof of Proposition 6	38
3. Proof of Lemma 5	39
B. Proof of Proposition 7: Lieb–Robinson bound by time-evolved Hamiltonian	40
1. Statement	40
2. Proof of Proposition 7	41
C. Proof of Lemma 3: Lieb–Robinson bound for Hamiltonian with a finite interaction length	44
1. Statement	44
2. Proof of Lemma 3	44
D. Proof of Proposition 8: connection of two Lieb–Robinson bounds from different unitary operators	46

1. Statement	46
2. Proof of Proposition 8	46
XIV. Contribution of the Long-range interactions to Lieb-Robinson bound: Proof of Theorem 3	48
A. Statement	48
B. Proof of the theorem (see Fig. 10 for the outline)	49
C. Proofs of Propositions 9 and 10	53
1. Statement	53
2. Proofs of Propositions 9 and 10.	54
D. Proof of Proposition 11	59
1. Statement	59
2. Proof of Proposition 11.	60
3. Derivation of the inequality (S.442)	63

IX. SETUP

A. Definition of the lattice

We here recall the setup in the main theorem. We consider a quantum spin system with n spins, where each of the spin sits on a vertex of the D -dimensional graph (or D -dimensional lattice) with Λ the total spin set, namely $|\Lambda| = n$. We assume that a finite dimensional Hilbert space is assigned to each of the spins. We note that our Lieb-Robinson bound does not depend on the spin dimensions. For a partial set $X \subseteq \Lambda$, we denote the cardinality, that is, the number of vertices contained in X , by $|X|$ (e.g. $X = \{i_1, i_2, \dots, i_{|X|}\}$). We also denote the complementary subset of X by $X^c := \Lambda \setminus X$.

For arbitrary subsets $X, Y \subseteq \Lambda$, we define $d_{X,Y}$ as the shortest path length on the graph that connects X and Y ; that is, if $X \cap Y \neq \emptyset$, $d_{X,Y} = 0$. When X is composed of only one element (i.e., $X = \{i\}$), we denote $d_{\{i\},Y}$ by $d_{i,Y}$ for the simplicity. We also define $\text{diam}(X)$ as follows:

$$\text{diam}(X) := 1 + \max_{i,j \in X} (d_{i,j}). \quad (\text{S.1})$$

B. Definition of the long-range interacting Hamiltonian

We consider a Hamiltonian as

$$H = \sum_{Z \subseteq \Lambda} h_Z, \quad (\text{S.2})$$

where each of the interaction terms $\{h_Z\}_{Z \subseteq \Lambda}$ acts on the sites of $Z \subseteq \Lambda$. Note that we here *do not* need to assume the few-body interaction (i.e., $|Z| = \mathcal{O}(1)$). We do not explicitly consider the time-dependence of the Hamiltonians, but all the analyses can be generalized to the time-dependent Hamiltonians.

When we discuss the few-body interactions, we impose the following additional assumption:

$$H = \sum_{Z \subseteq \Lambda, |Z| \leq k} h_Z. \quad (\text{S.3})$$

This Hamiltonian includes at most k -body interactions, and we refer to such a Hamiltonian as “ k -local Hamiltonian.” For example, in the case of $|Z| \leq 2$ (i.e., including up to two-body interactions), the Hamiltonian is given in the form of

$$H = \sum_{Z: |Z|=2} h_Z + \sum_{Z: |Z|=1} h_Z = \sum_{i < j} h_{i,j} + \sum_{i=1}^n h_i. \quad (\text{S.4})$$

In the main text, we considered the above Hamiltonian for the simplicity of the notation. If we consider the k -local Hamiltonian instead of the generic Hamiltonian (S.2), our main results are slightly modified as shown in Table I.

TABLE I. Dependence of the k -locality of our main theorems. Theorem 2 does not depend on the k -locality.

Hamiltonian	Theorem 1	Theorem 2	Theorem 3
Generic Hamiltonians in Eq. (S.2)	(S.27) and (S.28)	(S.32)	(S.38)
k -local Hamiltonians in Eq. (S.3)	(S.29) and (S.30)	(S.32)	(S.39)

Throughout the paper, for arbitrary operators A and O , we define

$$O(A, t) := e^{iAt} O e^{-iAt}. \quad (\text{S.5})$$

In particular, for $A = H$, we denote

$$O(t) := e^{iHt} O e^{-iHt} \quad (\text{S.6})$$

for the simplicity of the notation.

In order to characterize the long-range interaction of the Hamiltonian, we impose the following assumption for the Hamiltonian:

Assumption 1 (Power-law decaying interactions). *We assume the power-law decay of the interaction in the following senses:*

$$\sup_{i \in \Lambda} \sum_{Z: Z \ni i, \text{diam}(Z) \geq r} \|h_Z\| \leq g r^{-\alpha+D}, \quad (\text{S.7})$$

$$\sup_{i, j \in \Lambda} \sum_{Z: Z \ni \{i, j\}} \|h_Z\| \leq g_0 (d_{i, j} + 1)^{-\alpha} \quad (\text{S.8})$$

with

$$\alpha > 2D + 1, \quad (\text{S.9})$$

where $\|\cdots\|$ denotes the operator norm and the parameters g, g_0 are $\mathcal{O}(1)$ constants which do not depend on the system size n . By taking the energy unit appropriately, we set

$$g = 1. \quad (\text{S.10})$$

Here, $\sum_{Z: Z \ni i, \text{diam}(Z) \geq r}$ means the summation which picks up all the subsets $Z \subseteq \Lambda$ such that $Z \ni i$ and $\text{diam}(Z) \geq r$. In the similar way, $\sum_{Z: Z \ni \{i, j\}}$ means the summation which picks up all the subsets $Z \subseteq \Lambda$ which include $\{i, j\}$. We notice that we actually need only the condition (S.8) since it implies the condition (S.7).

From the assumption (S.8), we can also derive

$$\sum_{Z: Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset} \|h_Z\| \leq \sum_{i \in X} \sum_{j \in Y} \sum_{Z: Z \ni \{i, j\}} \|h_Z\| \leq g_0 \sum_{i \in X} \sum_{j \in Y} (d_{i, j} + 1)^{-\alpha} \leq g_0 |X| \cdot |Y| (d_{X, Y} + 1)^{-\alpha}, \quad (\text{S.11})$$

where we use $d_{X, Y} \leq d_{i, j}$ for $i \in X$ and $j \in Y$.

C. Coarse grained set (Fig. 6)

For a subset $X \subseteq \Lambda$, we define $X[r]$ as

$$X[r] := \{i \in \Lambda | d_{X, i} \leq r\}, \quad (\text{S.12})$$

where $X[0] = X$ and r is an arbitrary positive number (i.e., $r \in \mathbb{R}^+$). We also define the coarse grained total set $\Lambda^{(\xi)}$ as the minimum subset such that $\Lambda^{(\xi)}[\xi] = \Lambda$, namely

$$\Lambda^{(\xi)} := \arg \min_{Z \subseteq \Lambda | Z[\xi] = \Lambda} |Z|, \quad (\text{S.13})$$

where $\Lambda^{(0)} = \Lambda$. Similarly, for an arbitrary subset $X \subseteq \Lambda$, we define $X^{(\xi)} (\subseteq \Lambda^{(\xi)})$ as

$$X^{(\xi)} := \arg \min_{Z \subseteq \Lambda^{(\xi)} | Z[\xi] \supseteq X} |Z|, \quad (\text{S.14})$$

where $X^{(0)} = X$. From the definition, because of $X^{(\xi_2)}[\xi_1 + \xi_2] = (X^{(\xi_2)}[\xi_2])[\xi_1] \supseteq X[\xi_1]$, we notice that

$$X[\xi_1] \subseteq X^{(\xi_2)}[\xi_1 + \xi_2] \quad \text{and} \quad |X[\xi_1]| \leq |X^{(\xi_2)}[\xi_1 + \xi_2]|, \quad (\text{S.15})$$

where we use the fact that $|X| \leq |Y|$ for $X \subseteq Y$.

We introduce a geometric parameter γ which is determined only by the lattice structure. We define $\gamma \geq 1$ as a constant of $\mathcal{O}(1)$ which satisfies all the following inequalities for arbitrary $\xi \in \mathbb{R}^+$ and $X \subseteq \Lambda$:

$$|X| \leq \gamma [\text{diam}(X)]^D, \quad (\text{S.16})$$

$$|i[r]| \leq \gamma (2r)^D \quad (r \geq 1), \quad (\text{S.17})$$

$$|X^{(\xi)}| \leq \max(1, \gamma [\text{diam}(X)/\xi]^D), \quad (\text{S.18})$$

$$\max_{i \in \Lambda} \left(\#\{j \in \Lambda^{(\xi)} | r \leq d_{i, j} < r + \xi\} \right) \leq 2\gamma D (2r/\xi)^{D-1} \quad \text{for } r \geq \xi \geq 1, \quad (\text{S.19})$$




FIG. 6. (a) Extended subset. The subset $X[r]$ is defined by extending the original subset X by a distance r . (b) The subset $\Lambda^{(\xi)} \subseteq \Lambda$ is the coarse grained lattice (orange sites) which is defined as the minimum subset such that $\Lambda^{(\xi)}[\xi] = \Lambda$. (c) For an arbitrary subset $X \subseteq \Lambda$ (pink region), $X^{(\xi)} \subseteq \Lambda^{(\xi)}$ is the coarse grained subset (red sites) which is defined as the minimum subset such that $X^{(\xi)}[\xi] \supseteq X$. As shown in (S.18), the cardinality of the subset $X^{(\xi)}$ is roughly $(1/\xi)^D$ times as the original one, namely $|X^{(\xi)}| \approx (1/\xi)^D |X|$.

where we have defined $\text{diam}(X) = \max_{i,j \in X} (d_{i,j}) + 1 \geq 1$. Note that the inequality (S.17) implies

$$|X[r]| = \left| \bigcup_{i \in X} i[r] \right| \leq \sum_{i \in X} |i[r]| \leq \gamma(2r)^D |X|. \quad (\text{S.20})$$

Furthermore, we compare two subset $X^{(\xi)}$ and $X^{(c\xi)}$ with $c \geq 1$. Because of $X^{(c\xi)}[c\xi] \supseteq X$, we have $X^{(\xi)} \leq (X^{(c\xi)}[c\xi])^{(\xi)}$, and hence

$$|X^{(\xi)}| \leq |(X^{(c\xi)}[c\xi])^{(\xi)}| \leq \sum_{i \in X^{(c\xi)}} |(i[c\xi])^{(\xi)}| \leq \sum_{i \in X^{(c\xi)}} \gamma(2c)^D \leq \gamma(2c)^D |X^{(c\xi)}|, \quad (\text{S.21})$$

where in the third inequality we utilize (S.17).

X. MAIN RESULTS

A. Definitions

We first define $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound and $\mathcal{G}(x, t, \mathcal{X})$ -Lieb-Robinson bound as follows:

Definition 1. Let H_0 be an arbitrary Hamiltonian. Then, the Hamiltonian H_0 satisfies $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound if it satisfies

$$\frac{\|[O_{\mathcal{X}}(H_0, t), O_{\mathcal{Y}}]\|}{\|O_{\mathcal{X}}\| \cdot \|O_{\mathcal{Y}}\|} \leq \mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}), \quad x = d_{\mathcal{X}, \mathcal{Y}} \quad (\text{S.22})$$

for arbitrary operators $O_{\mathcal{X}}$ and $O_{\mathcal{Y}}$ which are supported on \mathcal{X} and \mathcal{Y} , respectively, where we define $O_{\mathcal{X}}(H_0, t)$ as in Eq. (S.5). If the function $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ does not depend on the subset \mathcal{Y} , we simply denote it by $\mathcal{G}(x, t, \mathcal{X})$.

The definition clearly implies $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) \leq 2$ from $\|[O_{\mathcal{X}}(H_0, t), O_{\mathcal{Y}}]\| \leq 2\|O_{\mathcal{X}}\| \cdot \|O_{\mathcal{Y}}\|$ and we assume that the cut off is automatically included in the definition. For example, when $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ is given in the form of

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = \min(2, |\mathcal{X}| \cdot |\mathcal{Y}| e^{-x+vt}), \quad (\text{S.23})$$

we denote it by omitting the $\min(2, \dots)$ for the simplicity of the notation:

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = |\mathcal{X}| \cdot |\mathcal{Y}| e^{-x+vt}. \quad (\text{S.24})$$

The function $\mathcal{G}(x, t, \mathcal{X})$ is connected to an error in approximating time-evolved operator on a local region. We show the following lemma which has been given in Ref. [2]:

Lemma 1 (Bravyi, Hastings and Verstraete [2]). *Let H_0 be an arbitrary Hamiltonian satisfying $\mathcal{G}(x, t, \mathcal{X})$ -Lieb-Robinson bound. For arbitrary subsets $X, \tilde{X} \subseteq \Lambda$, we define $O_X(H_0, t, \tilde{X})$ as the local approximation of $O_X(H_0, t)$ onto the subset \tilde{X} :*

$$O_X(H_0, t, \tilde{X}) := \frac{1}{\text{tr}_{\tilde{X}^c}(\hat{1})} \text{tr}_{\tilde{X}^c} [O_X(t)] \otimes \hat{1}_{\tilde{X}^c}, \quad (\text{S.25})$$

where $\text{tr}_{\tilde{X}^c}(\cdots)$ is the partial trace with respect to the subset \tilde{X}^c . The definition implies that the operator $O_X(H_0, t, \tilde{X})$ is supported on the subset \tilde{X} and it also satisfies $\|O_X(H_0, t, \tilde{X})\| \leq \|O_X\|$. Then, by choosing $\tilde{X} = X[r]$ ($r \in \mathbb{N}$), we obtain

$$\|O_X(H_0, t) - O_X(H_0, t, X[r])\| \leq \mathcal{G}(r, t, X), \quad (\text{S.26})$$

where $\|O_X\| = 1$ and $X[r]$ was defined in Eq. (S.12). Recall that we mean by $\mathcal{G}(x, t, \mathcal{X})$ the function $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ which does not depend on the subset \mathcal{Y} .

B. Main theorem

We here show our main theorems:

Theorem 1 (Lieb-Robinson bound for long-range interacting systems). *Let us consider the long-range interacting Hamiltonian H satisfying the assumption 1. For $|t| \geq 1$, the Hamiltonian H satisfies the $\mathcal{G}_H(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound as*

$$\mathcal{G}_H(x, t, \mathcal{X}, \mathcal{Y}) = C_H |\mathcal{X}^{(\bar{v}|t)}| \cdot |\mathcal{Y}^{(\bar{v}|t)}| \frac{|t|^{2D+1} \log^{2D}(x+1)}{(x - \bar{v}|t|)^\alpha} \quad (|t| \leq x/\bar{v}) \quad (\text{S.27})$$

and

$$\mathcal{G}_H(x, t, \mathcal{X}) = C'_H |\mathcal{X}^{(\bar{v}|t)}|^2 \frac{|t|^{D+1} \log^{2D}(x+1)}{(x - \bar{v}|t|)^{\alpha-D}} \quad (|t| \leq x/\bar{v}), \quad (\text{S.28})$$

where C_H, C'_H and \bar{v} are constants which depend only on the parameters $\{D, g_0, \alpha, \gamma\}$. Note that we use the notation in Eq. (S.14) for $\mathcal{X}^{(\bar{v}|t)}$ and $\mathcal{Y}^{(\bar{v}|t)}$.

Theorem 1' (k -local Hamiltonian). *Let us impose an additional assumption of the k -locality (S.3) with $k = \mathcal{O}(1)$. Then, the Hamiltonian H satisfies the $\mathcal{G}_H^{(k)}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound as*

$$\mathcal{G}_H^{(k)}(x, t, \mathcal{X}, \mathcal{Y}) = C_H^{(k)} |\mathcal{X}^{(\bar{v}^{(k)}|t)}| \cdot |\mathcal{Y}^{(\bar{v}^{(k)}|t)}| \frac{|t|^{2D+1}}{(x - \bar{v}^{(k)}|t|)^\alpha} \quad (|t| \leq x/\bar{v}^{(k)}) \quad (\text{S.29})$$

and

$$\mathcal{G}_H^{(k)}(x, t, \mathcal{X}) = C_H^{(k)'} |\mathcal{X}^{(\bar{v}^{(k)}|t)}|^2 \frac{|t|^{D+1}}{(x - \bar{v}^{(k)}|t|)^{\alpha-D}} \quad (|t| \leq x/\bar{v}^{(k)}), \quad (\text{S.30})$$

where $C_H^{(k)}, C_H^{(k)'}$ and $\bar{v}^{(k)}$ are constants which depend only on the parameters $\{D, g_0, \alpha, \gamma, k\}$.

Remark. From the theorem, outside the light-cone of $R \geq \bar{v}|t|$, the commutator is as small as $\mathcal{O}(R^{-\alpha})$. The first inequality (S.27) is stronger than the second one (S.28) when subset \mathcal{Y} is small. However, in the case where \mathcal{Y} is infinitely large, the inequality (S.27) is useless. For example, in order to obtain the local approximation (S.26) for time-evolved operators, we need the second inequality (S.28).

In the case of generic Hamiltonians, we need the logarithmic correction $\log^{2D}(x+1)$. This condition arises from Theorem 3 below, which characterizes a contribution from interactions with sufficiently long length scales. This logarithmic correction can be deleted when we assume the k -locality of the Hamiltonian (see Theorem 3').




FIG. 7. Outline of the proof. In the proof, we decompose the Hamiltonian as in Eq. (S.34). The unitary operator e^{-iHt} also consists of $e^{-iH_{\leq \ell_t}t}$ and $U_{> \ell_t} = \mathcal{T}e^{-i \int_0^t H_{> \ell_t}(H_{\leq \ell_t}, \tau) d\tau}$ as in Eq. (S.36). The former one characterizes the middle-range interactions, while the latter one gives the contribution from the long-range interactions. Then, the central tasks for the proof of Theorem 1 are the derivations of the Lieb-Robinson bounds for $e^{-iH_{\leq \ell_t}t}$ and $U_{> \ell_t}$, which are given by Theorems 2 and 3 (or Theorem 3'), respectively. The remaining task is to connect these two bounds, which are given by Eqs. (S.40)–(S.55). For a technical reason (see Sec. XI), we first restrict the subset Y such that $\text{diam}(Y) \leq 2v^*|t| + 1$ as in Eq. (S.41). Afterward, we remove the restriction by using Proposition 5, which completes the proof of Theorem 1.

C. Proof of Theorem 1 (see also Fig. 7 for the basic outline)

Before discussing the long-range interaction, we consider the Lieb-Robinson bound which is obtained from the following Hamiltonian with an interaction truncation:

$$H_{\leq \ell} = \sum_{Z \subseteq \Lambda: \text{diam}(Z) \leq \ell} h_Z. \quad (\text{S.31})$$

As long as $\ell = \mathcal{O}(1)$, the Hamiltonian gives the Lieb-Robinson bound with a finite velocity (see Lemma 3). However, when we choose the length ℓ depending on the time t , it is highly nontrivial whether the unitary operator $e^{-iH_{\leq \ell}t}$ satisfies the Lieb-Robinson bound with a finite velocity or not. This kind of “middle-range” Lieb-Robinson bound is crucial in discussing the long-range Lieb-Robinson bound. We here show the following theorem which we will prove in Sec. XII:

Theorem 2 (Middle-range Lieb-Robinson bound). *We consider a Hamiltonian H which satisfies Assumption (S.7) with $\alpha > 2D + 1$. Then, for arbitrary ℓ , the Hamiltonian $H_{\leq \ell}$ satisfies the $\mathcal{G}_{\leq \ell}(x, t, \mathcal{X})$ -Lieb-Robinson bound with*

$$\mathcal{G}_{\leq \ell}(x, t, \mathcal{X}) = \tilde{C}_0 |\mathcal{X}|^2 (1 + x/\ell)^{D-1} e^{-2(x-v^*|t|)/\ell}, \quad (\text{S.32})$$

where v^* is a constant which depends only on the parameters $\{D, g_0, \alpha, \gamma\}$ and

$$\tilde{C}_0 := \frac{4}{3} e^{10/3} 15^D D! \gamma \zeta_2, \quad \zeta_2 := 16e^3 \gamma^2 \cdot 45^D. \quad (\text{S.33})$$

From Theorem 2, we ensure that the middle-range Lieb-Robinson bound (S.32) retains the linear light cone for $\ell \leq \mathcal{O}(t)$. Afterward, we prove the long-range Lieb-Robinson bound by using this middle-range Lieb-Robinson bound. For the purpose, we consider the following decomposition of the total Hamiltonian into

$$H = H_{\leq \ell_t} + H_{> \ell_t} \quad (\text{S.34})$$

with

$$\ell_t = |t|^{\tilde{\eta}}, \quad \tilde{\eta} := 1 - \frac{\alpha - 2D - 1}{2(\alpha - D)}, \quad (\text{S.35})$$

where we defined $H_{\leq \ell_t}$ by Eq. (S.31). Note that under the condition of $\alpha > 2D + 1$ we have $\tilde{\eta} < 1$ and the dynamics by $H_{\leq \ell_t}$ retains the linear light cone. We then decompose the unitary operator e^{-iHt} into the form of

$$e^{-iHt} = e^{-iH_{\leq \ell_t}t} \mathcal{T} e^{-i \int_0^t H_{> \ell_t}(H_{\leq \ell_t}, \tau) d\tau}, \quad (\text{S.36})$$

where \mathcal{T} is the time-ordering operator and we use the notation (S.5), namely $H_{> \ell_t}(H_{\leq \ell_t}, \tau) = e^{iH_{\leq \ell_t}\tau} H_{> \ell_t} e^{-iH_{\leq \ell_t}\tau}$. We have already obtained the Lieb-Robinson bound for $e^{-iH_{\leq \ell_t}t}$ from Theorem 2. Hence, we need to consider the Lieb-Robinson bound for the unitary operator of

$$U_{> \ell_t} := \mathcal{T} e^{-i \int_0^t H_{> \ell_t}(H_{\leq \ell_t}, \tau) d\tau}. \quad (\text{S.37})$$

We can prove the following theorem on the Lieb-Robinson bound for the unitary operator $U_{> \ell_t}$ (see Sec. XIV for the proof):

Theorem 3 (Contribution by the long-range interacting terms). *Under the choice of ℓ_t by Eq. (S.35), the unitary operator (S.37) satisfies the $\mathcal{G}_{> \ell_t}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound as*

$$\mathcal{G}_{> \ell_t}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{J}_0 |\mathcal{X}^{(v^*|t)}| \cdot |\mathcal{Y}^{(v^*|t)}| \frac{|t|^{2D+1} \log^{2D}(x+1)}{(x - \kappa_0 v^*|t|)^\alpha}, \quad (\text{S.38})$$

where \mathcal{J}_0 and κ_0 are constants which depend only on the parameters $\{D, g_0, \alpha, \gamma\}$.

The logarithmic term $\log^{2D}(x+1)$ appears from the contribution by the macroscopic interactions (e.g., $\mathcal{O}(n)$ -body interactions). It vanishes under the assumption of the k -locality (i.e., few-body interactions) as follows:

Theorem 3' (k -local Hamiltonian). *Let us assume the k -locality of the Hamiltonian [see Eq. (S.3) for the definition]. Then, under the same setup as that of Theorem 3, we obtain the Lieb-Robinson bound $\mathcal{G}_{> \ell_t}^{(k)}(x, t, \mathcal{X}, \mathcal{Y})$*

$$\mathcal{G}_{> \ell_t}^{(k)}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{J}_0^{(k)} |\mathcal{X}^{(v^*|t)}| \cdot |\mathcal{Y}^{(v^*|t)}| \frac{|t|^{2D+1}}{(x - \kappa_0 v^*|t|)^\alpha}, \quad (\text{S.39})$$

where $\mathcal{J}_0^{(k)}$ and κ_0 are constants which depend only on the parameters $\{D, g_0, \alpha, \gamma, k\}$.

By using Theorems 2 and 3, we prove the main theorem as follows. We here estimate the norm of the commutator

$$\|[O_X(t), O_Y]\| = \|[U_{> \ell_t}^\dagger O_X U_{> \ell_t}, O_Y(H_{\leq \ell_t}, -t)]\|, \quad R = d_{X,Y} \quad (\text{S.40})$$

for operators O_X and O_Y with $\|O_X\| = \|O_Y\| = 1$, where we impose the restriction of

$$\text{diam}(Y) \leq 2v^*|t| + 1. \quad (\text{S.41})$$

This restriction is removed afterward. Note that we impose no restrictions on X . By using the notation of Eq. (S.25), we first decompose $O_Y(H_{\leq \ell_t}, -t)$ as

$$\begin{aligned} O_Y(H_{\leq \ell_t}, -t) &= O_Y(H_{\leq \ell_t}, -t, Y[\xi_1]) + O_{t,Y[\xi_{j_*}]} + \sum_{j=2}^{j_*} O_{t,Y[\xi_j]}, \\ O_{t,Y[\xi_j]} &:= O_Y(H_{\leq \ell_t}, -t, Y[\xi_j]) - O_Y(H_{\leq \ell_t}, -t, Y[\xi_{j-1}]), \\ O_{t,Y[\xi_{j_*}]} &:= O_Y(H_{\leq \ell_t}, -t) - O_Y(H_{\leq \ell_t}, -t, Y[\xi_{j_*}]), \quad \xi_{j_*} := \lfloor R/2 \rfloor, \end{aligned} \quad (\text{S.42})$$

where $\xi_1 < \xi_2 < \dots < \xi_{j_*} = \lfloor R/2 \rfloor$ and we will appropriately determine $\{\xi_j\}_{j=1}^{j_*-1}$ afterwards. From the inequality (S.26) in Lemma 1, we have

$$\begin{aligned} \|O_Y(H_{\leq \ell_t}, -t, Y[\xi_1])\| &\leq 1, \\ \|O_{t,Y[\xi_{j_*}]} \| &\leq \mathcal{G}_{\leq \ell_t}(\xi_{j_*}, t, Y) = \mathcal{G}_{\leq \ell_t}(\lfloor R/2 \rfloor, t, Y), \\ \|O_{t,Y[\xi_j]} \| &= \|O_Y(H_{\leq \ell_t}, -t, Y[\xi_j]) - O_Y(H_{\leq \ell_t}, -t, Y[\xi_{j-1}])\| \\ &\leq \|O_Y(H_{\leq \ell_t}, -t, Y[\xi_j]) - O_Y(H_{\leq \ell_t}, -t)\| + \|O_Y(H_{\leq \ell_t}, -t) - O_Y(H_{\leq \ell_t}, -t, Y[\xi_{j-1}])\| \\ &\leq 2\mathcal{G}_{\leq \ell_t}(\xi_{j-1}, t, Y), \end{aligned} \quad (\text{S.43})$$

where we use $\xi_{j_*} := \lfloor R/2 \rfloor$ and $\mathcal{G}_{\leq \ell_t}(\xi_j, t, Y) \leq \mathcal{G}_{\leq \ell_t}(\xi_{j-1}, t, Y)$. By using the above decomposition, we obtain the upper bound of the commutator (S.40) as follows:

$$\begin{aligned}
& \| [U_{>\ell_t}^\dagger O_X U_{>\ell_t}, O_Y(H_{\leq \ell_t}, -t)] \| \\
& \leq \| [U_{>\ell_t}^\dagger O_X U_{>\ell_t}, O_Y(H_{\leq \ell_t}, -t, Y[\xi_1])] \| + \| [U_{>\ell_t}^\dagger O_X U_{>\ell_t}, O_{t,Y[\xi_{j_*}]}] \| + \sum_{j=2}^{\infty} \| [U_{>\ell_t}^\dagger O_X U_{>\ell_t}, O_{t,Y[\xi_j]}] \| \\
& \leq \mathcal{G}_{>\ell_t}(R - \xi_1, t, X, Y[\xi_1]) \cdot \| O_Y(H_{\leq \ell_t}, -t, Y[\xi_1]) \| + 2\| O_X \| \cdot \| O_{t,Y[\xi_{j_*}]} \| + \sum_{j=2}^{j_*} \mathcal{G}_{>\ell_t}(R - \xi_j, t, X, Y[\xi_j]) \cdot \| O_{t,Y[\xi_j]} \| \\
& \leq \mathcal{G}_{>\ell_t}(R - \xi_1, t, X, Y[\xi_1]) + 2\mathcal{G}_{\leq \ell_t}(\lfloor R/2 \rfloor, t, Y) + 2 \sum_{j=2}^{j_*} \mathcal{G}_{>\ell_t}(R - \xi_j, t, X, Y[\xi_j]) \cdot \mathcal{G}_{\leq \ell_t}(\xi_{j-1}, t, Y), \tag{S.44}
\end{aligned}$$

where we use Theorems 3 from the second line to the third line, and use the inequalities in (S.43) and $\| O_X \| = 1$ from the third line to the fourth line.

We, in the following, determine $\{\xi_j\}_{j=1}^{j_*-1}$ such that

$$\mathcal{G}_{\leq \ell_t}(\xi_j, t, Y) = \tilde{C}_0 |Y|^2 (1 + \xi_j/\ell_t)^{D-1} e^{-2(\xi_j - v^*|t|)/\ell_t} \leq e^{-j} \tag{S.45}$$

for each of $j \geq 1$, where we use the form of Eq. (S.32) for $\mathcal{G}_{\leq \ell_t}(\xi_j, t, Y)$. First, because of $\text{diam}(Y) \leq 2v^*|t| + 1$, we obtain

$$|Y|^2 \leq \gamma^2 (2v^*|t| + 1)^{2D}, \tag{S.46}$$

where we use the inequality (S.16). Thus, we need to choose ξ_j as

$$\xi_j = c_1 v^*|t| + c_2 \ell_t j = c_1 v^*|t| + c_2 |t|^{\tilde{\eta}} j, \tag{S.47}$$

where c_1 and c_2 are constants which depend on v^* , \tilde{C}_0 and $\tilde{\eta}$; that is, they depend on $\{D, g_0, \alpha, \gamma\}$. Remember that we have chosen ℓ_t as in Eq. (S.35). By applying the inequality (S.45) to (S.44), we obtain

$$\| [U_{>\ell_t}^\dagger O_X U_{>\ell_t}, O_Y(H_{\leq \ell_t}, -t)] \| \leq 2\mathcal{G}_{\leq \ell_t}(\lfloor R/2 \rfloor, t, Y) + 2 \sum_{j=1}^{j_*} e^{-j+1} \cdot \mathcal{G}_{>\ell_t}(R - \xi_j, t, X, Y[\xi_j]), \tag{S.48}$$

where the first term in (S.44) is now involved in the summation.

We then consider the values of $\mathcal{G}_{>\ell_t}(R - \xi_j, t, X, Y[\xi_j])$ for $j \leq j_*$. Because of $\xi_j \leq \lfloor R/2 \rfloor \leq R/2$, we obtain

$$\begin{aligned}
\mathcal{G}_{>\ell_t}(R - \xi_j, t, X, Y[\xi_j]) &= \mathcal{J}_0 |X^{(v^*|t|)}| \cdot |Y[\xi_j]^{(v^*|t|)}| \frac{|t|^{2D+1} \log^{2D}(R - \xi_j + 1)}{(R - \xi_j - \kappa_0 v^*|t|)^\alpha} \\
&\leq \mathcal{J}_0 |X^{(v^*|t|)}| \gamma (3 + 2c_1 + 2c_2 j |t|^{-1+\tilde{\eta}}/v^*)^D \frac{|t|^{2D+1} \log^{2D}(R + 1)}{(R/2 - \kappa_0 v^*|t|)^\alpha}, \tag{S.49}
\end{aligned}$$

where we estimate $|Y[\xi_j]^{(v^*|t|)}|$ by using the inequality (S.18) as follows:

$$|Y[\xi_j]^{(v^*|t|)}| \leq \gamma \left(\frac{\text{diam}(Y) + 2\xi_j}{v^*|t|} \right)^D \leq \gamma (3 + 2c_1 + 2c_2 j |t|^{-1+\tilde{\eta}}/v^*)^D, \tag{S.50}$$

where in the second inequality we use $(v^*|t|)^{-1} \text{diam}(Y) \leq 2 + (v^*|t|)^{-1} \leq 3$ and the form of Eq. (S.47). By applying the inequality (S.49) to (S.48), we obtain

$$\begin{aligned}
& \| [U_{>\ell_t}^\dagger O_X U_{>\ell_t}, O_Y(H_{\leq \ell_t}, -t)] \| \\
& \leq 2\mathcal{G}_{\leq \ell_t}(\lfloor R/2 \rfloor, t, Y) + 2\mathcal{J}_0 \gamma |X^{(v^*|t|)}| \frac{|t|^{2D+1} \log^{2D}(R + 1)}{(R/2 - \kappa_0 v^*|t|)^\alpha} \sum_{j=1}^{j_*} e^{-j+1} (3 + 2c_1 + 2c_2 j |t|^{-1+\tilde{\eta}}/v^*)^D \\
& \leq 2\mathcal{G}_{\leq \ell_t}(\lfloor R/2 \rfloor, t, Y) + 2^{\alpha+1} c_3 \mathcal{J}_0 \gamma |X^{(v^*|t|)}| \frac{|t|^{2D+1} \log^{2D}(R + 1)}{(R - 2\kappa_0 v^*|t|)^\alpha}. \tag{S.51}
\end{aligned}$$

We here define c_3 as

$$c_3 := \sum_{j=1}^{\infty} e^{-j+1} (3 + 2c_1 + 2c_2 j |t|^{-1+\tilde{\eta}}/v^*)^D \geq \sum_{j=1}^{j_*} e^{-j+1} (3 + 2c_1 + 2c_2 j |t|^{-1+\tilde{\eta}}/v^*)^D, \tag{S.52}$$

where the inequality $c_2 j/v^* \geq c_2 j|t|^{-1+\tilde{\eta}}/v^*$ is satisfied from the condition of $|t| \geq 1$ in the theorem.

We define the parameter $\tilde{\kappa}_0$ ($> \kappa_0$) so that it satisfies the following inequality for $R > \tilde{\kappa}_0 v^*|t|$:

$$\mathcal{G}_{\leq \ell_t}(\lfloor R/2 \rfloor, t, Y) \leq 2^\alpha c_3 \mathcal{J}_0 \gamma |X^{(v^*|t|)}| \frac{|t|^{2D+1} \log^{2D}(R+1)}{(R - 2\tilde{\kappa}_0 v^*|t|)^\alpha}. \quad (\text{S.53})$$

More specifically, from the inequality (S.32), $\tilde{\kappa}_0$ is defined by

$$\tilde{\mathcal{C}}_0 |Y|^2 (1 + \lfloor R/2 \rfloor / |t|^{\tilde{\eta}})^{D-1} e^{-2|t|^{-\tilde{\eta}}(\lfloor R/2 \rfloor - v^*|t|)} \leq 2^\alpha c_3 \mathcal{J}_0 \gamma |X^{(v^*|t|)}| \frac{|t|^{2D+1} \log^{2D}(R+1)}{(R - 2\tilde{\kappa}_0 v^*|t|)^\alpha}. \quad (\text{S.54})$$

Because of $\tilde{\eta} < 1$ and $\text{diam}(Y) \leq 2v^*|t| + 1$, we can always find such a parameter $\tilde{\kappa}_0$ which depends only on $\{D, g_0, \alpha, \gamma\}$. The inequality (S.53) reduces the inequality (S.51) to

$$\|[O_X(t), O_Y]\| = \|[U_{>\ell_t}^\dagger O_X U_{>\ell_t}, O_Y(H_{\leq \ell_t}, -t)]\| \leq 2^{\alpha+2} c_3 \mathcal{J}_0 \gamma |X^{(v^*|t|)}| \frac{|t|^{2D+1} \log^{2D}(R+1)}{(R - 2\tilde{\kappa}_0 v^*|t|)^\alpha}. \quad (\text{S.55})$$

We note that the subset Y is now restricted by the condition (S.41), namely $\text{diam}(Y) \leq 2v^*|t| + 1$.

In order to remove the restriction, we use Proposition 5 in the subsequent section. Here, the inequality (S.55) is given in the form of (S.69) by choosing

$$\mathcal{F}(t, \mathcal{X}, \mathcal{Y}) = 2^{\alpha+2} c_3 \mathcal{J}_0 \gamma |\mathcal{X}^{(v^*|t|)}| \cdot |t|^{2D+1}, \quad p = 2D, \quad \xi = v^*|t|, \quad \alpha_0 = \alpha, \quad \kappa = 2\tilde{\kappa}_0. \quad (\text{S.56})$$

Then, from the inequalities (S.70) and (S.72), we obtain

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = 2^{\alpha+3} c_3 \mathcal{J}_0 \gamma |\mathcal{X}^{(v^*|t|)}| \cdot |\mathcal{Y}^{(v^*|t|)}| \frac{|t|^{2D+1} \log^{2D}(x+1)}{(x - 2\tilde{\kappa}_0 v^*|t|)^\alpha} \quad (\text{S.57})$$

and

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = 2^{\alpha+2} c_3 \mathcal{J}_0 \gamma \mathcal{C}'_{2\tilde{\kappa}_0, 2D, \alpha} (v^*)^{-D} |\mathcal{X}^{(v^*|t|)}|^2 \frac{|t|^{D+1} \log^{2D}(x+1)}{(x - 2\tilde{\kappa}_0 v^*|t| - 4v^*|t|)^{\alpha-D}} \quad (\text{S.58})$$

for $x \geq (2\tilde{\kappa}_0 + 5)v^*|t|$, where $\mathcal{C}'_{2\tilde{\kappa}_0, 2D, \alpha}$ is defined by Eq. (S.73). Therefore, the two inequalities (S.57) and (S.58) reduce to the inequalities (S.27) and (S.28) respectively by choosing $\bar{v} = (2\tilde{\kappa}_0 + 5)v^*$; in order to upper-bound $\mathcal{X}^{(v^*|t|)}$ by $\mathcal{X}^{(\bar{v}|t|)}$, we utilize the inequality (S.21). We note that all the constants depend only on $\{D, g_0, \alpha, \gamma\}$.

In deriving the Lieb-Robinson bound for k -local Hamiltonians, we follow the same analytical steps. The only difference is that we utilize Theorem 3' instead of Theorem 3. After straightforward calculations, we can derive the inequalities (S.29) and (S.30). This completes the proof of the main theorem. \square

XI. EXTENDING THE LIEB-ROBINSON BOUND FOR LOCAL OPERATORS TO THAT FOR GENERIC OPERATORS

We first consider the Lieb-Robinson function $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ under the constraint that the local subsets \mathcal{X}, \mathcal{Y} satisfy $\text{diam}(\mathcal{X}), \text{diam}(\mathcal{Y}) \leq 2\xi + 1$ for a given ξ . In our analyses, it is crucial to relate the Lieb-Robinson bound for local operators to that for generic operators, including global operators that have no restrictions on $\text{diam}(\mathcal{X})$ and $\text{diam}(\mathcal{Y})$.

To make the motivation clearer, we consider a connection of the Lieb-Robinson bounds from different time-evolutions $e^{-iH_1 t}$ and $e^{-iH_2 t}$. We are now interested in the commutator of $\|[e^{iH_2 t} e^{iH_1 t} O_X e^{-iH_1 t} e^{-iH_2 t}, O_Y]\|$. The difficulty in the connection lies in the subset dependence (i.e., $|X|$ or $|Y|$) of the Lieb-Robinson bound. We often obtain the Lieb-Robinson bounds for $e^{-iH_1 t}$ and $e^{-iH_2 t}$ in the form of

$$\|[O_X(H_j, t), O_Y]\| \leq |X| \mathcal{F}_j(x, t) \quad (\|O_X\| = \|O_Y\| = 1) \quad (\text{S.59})$$

for $j = 1, 2$ and $x = d_{X,Y}$ (see [15, 17] for example), where $\mathcal{F}_j(x, t)$ is determined by details of H_j . In order to connect the Lieb-Robinson bounds for $e^{-iH_1 t}$ and $e^{-iH_2 t}$ to obtain that for $e^{-iH_1 t} e^{-iH_2 t}$, we consider a similar decomposition to Eq. (S.42):

$$O_X(H_1, t) = \sum_{s=1}^{\infty} \tilde{O}_{X[s\xi]} \quad (\text{S.60})$$

with

$$\tilde{O}_{X[\xi]} = O_X(H_1, t, X[\xi]), \quad \tilde{O}_{X[s\xi]} = O_X(H_1, t, X[s\xi]) - O_X(H_1, t, X[(s-1)\xi]) \quad (s \geq 2), \quad (\text{S.61})$$

where ξ is appropriately chosen and we use the definition (S.25). Note that $\lim_{s \rightarrow \infty} O_X(H_1, t, X[s\xi]) = O_X(H_1, t)$ and the operator $\tilde{O}_{X[s\xi]}$ is supported on the subset $X[s\xi] \subseteq \Lambda$. By using the decomposition, we obtain

$$\| [e^{iH_2t} e^{iH_1t} O_X e^{-iH_1t} e^{-iH_2t}, O_Y] \| \leq \sum_{s=1}^{\infty} \| [\tilde{O}_{X[s\xi]}(H_2, t), O_Y] \| \leq \sum_{s=1}^{\infty} |X[s\xi]| \mathcal{F}_2(x - s\xi, t) \cdot \| \tilde{O}_{X[s\xi]} \|, \quad (\text{S.62})$$

where we use $d_{X[s\xi], Y} \geq d_{X, Y} - s\xi = x - s\xi$ and the Lieb-Robinson bound (S.59) with $j = 2$.

From the inequality (S.26), we can obtain $\| \tilde{O}_{X[s\xi]} \| \leq 2|X| \mathcal{F}_1((s-1)\xi, t)$, where we use the Lieb-Robinson bound (S.59) with $j = 1$. Hence, the upper bound of (S.62) reduces to

$$\| [e^{iH_2t} e^{iH_1t} O_X e^{-iH_1t} e^{-iH_2t}, O_Y] \| \leq 2|X| \sum_{s=1}^{\infty} |X[s\xi]| \mathcal{F}_2(x - s\xi, t) \mathcal{F}_1((s-1)\xi, t). \quad (\text{S.63})$$

Then, after connecting two unitary operators e^{-iH_1t} and e^{-iH_2t} , the new Lieb-Robinson bound depends on the support X in the form of $|X|^2$. If we connect Lieb-Robinson bounds for m unitary operators, $|X|$ -dependence of the coefficient grows as $|X|^m$ and leads to a meaningless Lieb-Robinson bound in the process of the connections. In order to prevent it, we consider only the operators O_X and O_Y which are supported on local subsystems with $\text{diam}(X), \text{diam}(Y) \leq 2\xi + 1$ with appropriate ξ . After the connection process, we remove the constraints on the subset sizes to obtain the Lieb-Robinson bound for generic operators. The following theorem relates the Lieb-Robinson bound for local operators to that for generic operators:

Theorem 4. *We consider a Hamiltonian H_0 which satisfies the $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound in the form of*

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{F}(t, \mathcal{X}, \mathcal{Y}) \mathcal{L}(x) \quad \text{with} \quad \mathcal{L}(x) := e^{-x/\xi_0} \quad (\text{S.64})$$

for $\forall \mathcal{X}, \mathcal{Y} \subseteq \Lambda$ such that $\text{diam}(\mathcal{X}), \text{diam}(\mathcal{Y}) \leq 2\xi + 1$, where we assume $\mathcal{F}(t, \mathcal{X}, \mathcal{Y}) = \mathcal{F}(-t, \mathcal{X}, \mathcal{Y})$. Then, for arbitrary subsets $X \subset \Lambda$ with $\text{diam}(X) \leq 2\xi + 1$ and $L \subseteq \Lambda$, we obtain the Lieb-Robinson function $\mathcal{G}(x, t, X, L)$ which is given by

$$\mathcal{G}(x, t, X, L) = 2|L^{(\xi)}| \tilde{\mathcal{F}}(t) \mathcal{L}(x), \quad x = d_{X, L}, \quad (\text{S.65})$$

$$\tilde{\mathcal{F}}(t) := \sup_{\substack{\mathcal{X}, \mathcal{Y} \subseteq \Lambda \\ \text{diam}(X), \text{diam}(Y) \leq 2\xi + 1}} [\mathcal{F}(t, \mathcal{X}, \mathcal{Y})]. \quad (\text{S.66})$$

Furthermore, for arbitrary two subsets $L, L' \subseteq \Lambda$, we obtain the Lieb-Robinson function $\mathcal{G}(x, t, L, L')$ which is given by

$$\mathcal{G}(x, t, L, L') = \mathcal{C}_{\xi, \xi_0} |L^{(\xi)}|^2 \tilde{\mathcal{F}}(t) (1 + x/\xi)^{D-1} e^{-x/\xi_0}, \quad x = d_{L, L'}, \quad (\text{S.67})$$

where we define

$$\mathcal{C}_{\xi, \xi_0} := 2^{D+2} D! \gamma (1 + \xi_0/\xi)^D e^{5\xi/\xi_0}. \quad (\text{S.68})$$

Note that we here do not impose any restrictions on L and L' .

We obtain a similar statement for the case where $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ decays polynomially with respect to x . We have utilized the following proposition in deriving the inequalities (S.57) and (S.58) in the proof of Theorem 1:

Proposition 5. *We consider a Hamiltonian H_0 which satisfies the $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound in the form of*

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{F}(t, \mathcal{X}, \mathcal{Y}) \mathcal{L}(x) \quad \text{with} \quad \mathcal{L}(x) := \frac{\log^p(x+1)}{(x - \kappa\xi)^{-\alpha_0}} \quad (\alpha_0 > D+1) \quad (\text{S.69})$$

for $\forall \mathcal{X} \subseteq \Lambda$ without any restrictions and $\forall \mathcal{Y} \subseteq \Lambda$ with $\text{diam}(\mathcal{Y}) \leq 2\xi + 1$, where we assume $\mathcal{F}(t, \mathcal{X}, \mathcal{Y}) = \mathcal{F}(-t, \mathcal{X}, \mathcal{Y})$. Then, for arbitrary two subsets $L, L' \subseteq \Lambda$, we obtain the Lieb-Robinson function $\mathcal{G}(x, t, L, L')$ which is given by

$$\mathcal{G}(x, t, L, L') = 2|L^{(\xi)}| \tilde{\mathcal{F}}(t, L) \mathcal{L}(x), \quad x = d_{L, L'}, \quad (\text{S.70})$$

or

$$\mathcal{G}(x, t, L, L') = \mathcal{C}'_{\kappa, p, \alpha_0} |L^{(\xi)}| \tilde{\mathcal{F}}(t, L) \xi^{-D} \frac{\log^p(x+1)}{[x - (4 + \kappa)\xi]^{\alpha_0 - D}}, \quad x = d_{L, L'}, \quad (\text{S.71})$$

where we assume $x \geq (\kappa + 5)\xi$, and define $\tilde{\mathcal{F}}(t, L)$ and $\mathcal{C}'_{\kappa, p, \alpha_0}$ as

$$\tilde{\mathcal{F}}(t, L) := \sup_{\substack{\mathcal{Y} \subseteq \Lambda \\ \text{diam}(\mathcal{Y}) \leq 2\xi + 1}} [\mathcal{F}(t, L, \mathcal{Y})], \quad (\text{S.72})$$

$$\mathcal{C}'_{\kappa, p, \alpha_0} := \tilde{C}_{\kappa, p, \alpha_0} 2^{D+2} \gamma D(\kappa + 3)^{\alpha_0 - 1}, \quad \tilde{C}_{\kappa, p, \alpha_0} := \sup_{z \in \mathbb{R} | z \geq \kappa + 3} \left(\frac{z^{D+1} [2 \log(z) + 1]^p}{(z - \kappa - 2)^{\alpha_0}} \right). \quad (\text{S.73})$$

Note that we here do not impose any restrictions on L and L' .

A. Proof of Theorem 4

We first prove the inequality (S.65). For the proof, we need to consider the norm of

$$\|[O_X(H_0, t), O_L]\| \quad (\text{diam}(X) \leq 2\xi + 1), \quad (\text{S.74})$$

where O_X and O_L are arbitrary operators with the unit norm (i.e., $\|O_X\| = \|O_L\| = 1$) supported on X and L , respectively. Now, we do not assume any constraints on the subset L . Because $O_X(H_0, t, L^c)$ is supported on L^c from the definition (S.25), we have $[O_X(H_0, t, L^c), O_L] = 0$, and hence we have

$$\begin{aligned} \|[O_X(H_0, t), O_L]\| &= \|[O_X(H_0, t) - O_X(H_0, t, L^c), O_L] + [O_X(H_0, t, L^c), O_L]\| \\ &= \|[O_X(H_0, t) - O_X(H_0, t, L^c), O_L]\| \\ &\leq 2\|O_X(H_0, t) - O_X(H_0, t, L^c)\|, \end{aligned} \quad (\text{S.75})$$

where we use $\|O_L\| = 1$ in the last inequality. The partial trace with respect to an arbitrary subset $X_0 \subseteq \Lambda$ is given by

$$\frac{1}{\text{tr}_{X_0}(\hat{1}_{X_0})} \text{tr}_{X_0}(O) \otimes \hat{1}_{X_0} = \int d\mu(U_{X_0}) U_{X_0}^\dagger O U_{X_0} \quad (\text{S.76})$$

for an arbitrary operator O , where U_{X_0} is a unitary operator acting on X_0 and $\mu(U_{X_0})$ be the Haar measure for U_{X_0} .

We here estimate an upper bound of the norm of $\|O_X(H_0, t) - O_X(H_0, t, L^c)\|$, which immediately gives the upper bound of $\|[O_X(H_0, t), O_L]\|$ by Ineq. (S.75). For the purpose, we consider the coarse grained subset of L [see Eq. (S.14)]. We here denote $L^{(\xi)}$ by $\{j_s\}_{s=1}^{n_L} \subseteq \Lambda^{(\xi)}$, where we define $n_L := |L^{(\xi)}|$. We then define the following subsets $\{L_s\}_{s=1}^{n_L}$ recursively:

$$\begin{aligned} L_1 &:= j_1[\xi] \cap L \quad \text{for } s = 1, \\ L_s &:= (j_s[\xi] \cap L) \setminus (L_1 \cup L_2 \cup \dots \cup L_{s-1}) \quad \text{for } s \geq 2. \end{aligned} \quad (\text{S.77})$$

Note that for arbitrary $s \neq s'$ we have $L_s \cap L_{s'} = \emptyset$. From the definition (S.14), we have $L^{(\xi)}[\xi] \supseteq L$ and hence

$$\bigcup_{s=1}^{n_L} L_s = L^{(\xi)}[\xi] \cap L = L. \quad (\text{S.78})$$

Also, we notice that

$$d_{X, L_s} \leq d_{X, L}, \quad \text{diam}(L_s) \leq \text{diam}(j_s[\xi]) \leq 2\xi + 1. \quad (\text{S.79})$$

By using the notations of (S.77), we obtain

$$O_X(H_0, t, L^c) = \frac{1}{\text{tr}_L(\hat{1}_L)} \text{tr}_L[O_X(H_0, t)] \otimes \hat{1}_L = \int d\mu(U_{L_1}) \int d\mu(U_{L_2}) \dots \int d\mu(U_{L_{n_L}}) U_L^\dagger O_X(H_0, t) U_L, \quad (\text{S.80})$$

where we define $U_L := \prod_{s=1}^{n_L} U_{L_s}$ with $U_{L_s} = \hat{1}$ for $L_s = \emptyset$. From the assumption of $\text{diam}(X) \leq 2\xi + 1$ and the inequality (S.79), we can apply the Lieb-Robinson bound (S.64) to $\|[O_X(H_0, t), U_{L_s}]\|$ as follows:

$$\|[O_X(H_0, t), U_{L_s}]\| \leq \mathcal{F}(t, X, L_s) \mathcal{L}(d_{X, L_s}), \quad (\text{S.81})$$

where U_{L_s} can be an arbitrary unitary operator. Therefore, we calculate

$$\begin{aligned} \|O_X(H_0, t) - O_X(H_0, t, L^c)\| &\leq \int d\mu(U_{L_1}) \int d\mu(U_{L_2}) \dots \int d\mu(U_{L_{n_L}}) \|[O_X(H_0, t), U_L]\| \\ &\leq \sum_{s=1}^{n_L} \sup_{U_{L_s}} (\|[O_X(H_0, t), U_{L_s}]\|) \\ &\leq \sum_{s=1}^{n_L} \mathcal{F}(t, X, L_s) \mathcal{L}(d_{X, L_s}) \\ &\leq n_L \tilde{\mathcal{F}}(t) \mathcal{L}(d_{X, L}) = |L^{(\xi)}| \tilde{\mathcal{F}}(t) \mathcal{L}(d_{X, L}), \end{aligned} \quad (\text{S.82})$$

where we use the definition (S.66) of $\tilde{\mathcal{F}}(t)$ and $\mathcal{L}(d_{X, L_s}) \leq \mathcal{L}(d_{X, L})$ for $d_{X, L_s} \geq d_{X, L}$ in the fourth inequality. By applying the above inequality to (S.75), we prove the inequality (S.65).

We then prove the second inequality (S.67). For the proof, we need to estimate the norm of

$$\|[O_L(H_0, t), O_{L'}]\|, \quad (\text{S.83})$$

where O_L and $O_{L'}$ are arbitrary operators supported on $L, L' \subseteq \Lambda$ with $\|O_L\| = \|O_{L'}\| = 1$. In order to estimate the commutator norm, we calculate

$$\|O_L(H_0, t) - O_L(H_0, t, L[d_{L, L'} - 1])\|, \quad (\text{S.84})$$

which gives the upper bound of

$$\begin{aligned} \|[O_L(H_0, t), O_{L'}]\| &= \|[O_L(H_0, t) - O_L(H_0, t, L[d_{L, L'} - 1]), O_{L'}] + [O_L(H_0, t, L[d_{L, L'} - 1]), O_{L'}]\| \\ &= \|[O_L(H_0, t) - O_L(H_0, t, L[d_{L, L'} - 1]), O_{L'}]\| \\ &\leq 2\|O_L(H_0, t) - O_L(H_0, t, L[d_{L, L'} - 1])\|. \end{aligned} \quad (\text{S.85})$$

In the following, we estimate an upper bound of the norm of $\|O_L(H_0, t) - O_L(H_0, t, L[R])\|$ for an arbitrary R . We will set $R = d_{L, L'} - 1$ (or $R + 1 = d_{L, L'}$) afterward. Similar to the definition (S.77), we consider a set of spins $(L[R]^c)^{(\xi)} := \{j_s\}_{s=1}^{\tilde{n}} \subseteq \Lambda^{(\xi)}$ with $\tilde{n} = |(L[R]^c)^{(\xi)}|$, and define a set of $\{X_s\}_{s=1}^{\tilde{n}}$ as follows:

$$X_s := (j_s[\xi] \cap L[R]^c) \setminus (X_1 \cup X_2 \cup \dots \cup X_{s-1}). \quad (\text{S.86})$$

We obtain the similar relations to (S.78) and (S.79) as follows:

$$\bigcup_{s=1}^{\tilde{n}} X_s = L[R]^c, \quad d_{X_s, L} \geq R + 1, \quad \text{and} \quad \text{diam}(X_s) \leq \text{diam}(j_s[\xi]) \leq 2\xi + 1, \quad (\text{S.87})$$

where we use $d_{X_s, L} \geq d_{L[R]^c, L} = R + 1$ in the second inequality. By following the derivation of Eq. (S.80), the partial trace with respect to $L[R]^c$ is given by

$$O_L(H_0, t, L[R]) = \int d\mu(U_{L_1}) \int d\mu(U_{L_2}) \cdots \int d\mu(U_{L_{\tilde{n}}}) U_{L[R]^c}^\dagger O_L(H_0, t) U_{L[R]^c}, \quad (\text{S.88})$$

where we define $U_{L[R]^c} := \prod_{s=1}^{\tilde{n}} U_{X_s}$. Because of $\text{diam}(X_s) \leq 2\xi + 1$, we can apply the inequality (S.65) to $\|[O_L(H_0, t), U_{X_s}]\|$, which yields

$$\|[O_L(H_0, t), U_{X_s}]\| = \|[U_{X_s}(H_0, -t), O_L]\| \leq 2|L^{(\xi)}| \tilde{\mathcal{F}}(t) \mathcal{L}(d_{X_s, L}), \quad (\text{S.89})$$

and hence

$$\begin{aligned} \|O_L(H_0, t) - O_L(H_0, t, L[R])\| &\leq \int d\mu(U_{L_1}) \int d\mu(U_{L_2}) \cdots \int d\mu(U_{L_{\tilde{n}}}) \|[O_L(H_0, t), U_{L[R]^c}]\| \\ &\leq \sum_{s=1}^{\tilde{n}} \sup_{U_{X_s}} (\|[O_L(H_0, t), U_{X_s}]\|) \\ &\leq 2\tilde{\mathcal{F}}(t) \sum_{s=1}^{\tilde{n}} |L^{(\xi)}| \mathcal{L}(d_{L, X_s}). \end{aligned} \quad (\text{S.90})$$

Thus, the remaining task is to estimate the summation with respect to s .

By using the relation $L^{(\xi)}[\xi] \supseteq L$ and the definition of $\mathcal{L}(x) := e^{-x/\xi_0}$, we obtain

$$d_{L, X_s} \geq d_{L^{(\xi)}[\xi], X_s} \quad \text{and} \quad \mathcal{L}(d_{L^{(\xi)}[\xi], X_s}) \leq \max_{i \in L^{(\xi)}} [\mathcal{L}(d_{i[\xi], X_s})] \leq \sum_{i \in L^{(\xi)}} \mathcal{L}(d_{i[\xi], X_s}), \quad (\text{S.91})$$

which yields

$$\sum_{s=1}^{\tilde{n}} \mathcal{L}(d_{L, X_s}) \leq \sum_{s=1}^{\tilde{n}} \sum_{i \in L^{(\xi)}} \mathcal{L}(d_{i[\xi], X_s}). \quad (\text{S.92})$$

Moreover, the definition (S.86) implies $X_s \subseteq j_s[\xi]$ and we obtain for $i \in L^{(\xi)}$

$$d_{i[\xi], X_s} \geq d_{i[\xi], j_s[\xi]} \geq d_{i, j_s} - 2\xi. \quad (\text{S.93})$$

We therefore obtain

$$\sum_{s=1}^{\tilde{n}} \sum_{i \in L^{(\xi)}} \mathcal{L}(d_{i[\xi], X_s}) \leq \sum_{s=1}^{\tilde{n}} \sum_{i \in L^{(\xi)}} \mathcal{L}(d_{i, j_s} - 2\xi) = \sum_{i \in L^{(\xi)}} \sum_{j \in (L[R]^c)^{(\xi)}} \mathcal{L}(d_{i, j} - 2\xi), \quad (\text{S.94})$$

where in the equation we use the definition of $\{j_s\}_{s=1}^{\tilde{n}} := (L[R]^c)^{(\xi)}$. The conditions $i \in L^{(\xi)}$, $j \in (L[R]^c)^{(\xi)}$ and $d_{L,L[R]^c} = R+1$ imply

$$d_{i,j} \geq R+1-2\xi, \quad (\text{S.95})$$

which yields

$$\text{for } \forall i \in L^{(\xi)}, \quad \sum_{j \in (L[R]^c)^{(\xi)}} \mathcal{L}(d_{i,j} - 2\xi) \leq \sum_{j \in \Lambda^{(\xi)}: d_{i,j} \geq R+1-2\xi} \mathcal{L}(d_{i,j} - 2\xi). \quad (\text{S.96})$$

By applying the above inequality to (S.94), we obtain

$$\sum_{s=1}^{\tilde{n}} \sum_{i \in L^{(\xi)}} \mathcal{L}(d_{i[\xi], X_s}) \leq \sum_{i \in L^{(\xi)}} \sum_{j \in \Lambda^{(\xi)}: d_{i,j} \geq R+1-2\xi} \mathcal{L}(d_{i,j} - 2\xi). \quad (\text{S.97})$$

By using the inequality (S.19), the summation with respect to j is bounded from above by

$$\begin{aligned} \sum_{j \in \Lambda^{(\xi)}: d_{i,j} \geq R+1-2\xi} \mathcal{L}(d_{i,j} - 2\xi) &\leq \sum_{s=0}^{\infty} \sum_{\substack{j \in \Lambda^{(\xi)} \\ R+1-2\xi+s\xi \leq d_{i,j} < R+1-2\xi+(s+1)\xi}} \mathcal{L}(d_{i,j} - 2\xi) \\ &\leq 2\gamma D \sum_{s=0}^{\infty} \mathcal{L}(R+1-4\xi+s\xi) [2(R+1-2\xi+s\xi)/\xi]^{D-1}. \end{aligned} \quad (\text{S.98})$$

The form of $\mathcal{L}(x)$ is now given by

$$\mathcal{L}(x) = e^{-x/\xi_0}, \quad (\text{S.99})$$

and hence we obtain

$$\begin{aligned} &2\gamma D \sum_{s=0}^{\infty} \mathcal{L}(R+1-4\xi+s\xi) [2(R+1-2\xi+s\xi)/\xi]^{D-1} \\ &\leq 2^D \gamma D e^{-(R+1-4\xi)/\xi_0} \int_0^{\infty} e^{-(x-1)\xi/\xi_0} [(R+1-2\xi)/\xi + x]^{D-1} dx \\ &\leq 2^D \gamma D e^{5\xi/\xi_0} (D-1)! (\xi_0/\xi) [(R+1-2\xi+\xi_0)/\xi]^{D-1} e^{-(R+1)/\xi_0} \\ &\leq 2^D D! \gamma (1+\xi_0/\xi)^D e^{5\xi/\xi_0} [1+(R+1)/\xi]^{D-1} e^{-(R+1)/\xi_0} = (\mathcal{C}_{\xi, \xi_0}/4) [1+(R+1)/\xi]^{D-1} e^{-(R+1)/\xi_0}, \end{aligned} \quad (\text{S.100})$$

where we use the inequalities

$$\int_0^{\infty} (x+x_0)^{D-1} e^{-x/x_1} dx \leq (D-1)! x_1 (x_1+x_0)^{D-1} \quad (\text{S.101})$$

and

$$(R+1-2\xi+\xi_0)/\xi = [(R+1+\xi)/\xi] \cdot [1+(-3\xi+\xi_0)/(R+1+\xi)] \leq [1+(R+1)/\xi] \cdot (1+\xi_0/\xi). \quad (\text{S.102})$$

Also, the definition of \mathcal{C}_{ξ, ξ_0} has been given in Eq. (S.68). We thus arrive at the inequality of

$$\sum_{j \in \Lambda^{(\xi)}: d_{i,j} \geq R+1-2\xi} \mathcal{L}(d_{i,j} - 2\xi) \leq (\mathcal{C}_{\xi, \xi_0}/4) [1+(R+1)/\xi]^{D-1} e^{-(R+1)/\xi_0}. \quad (\text{S.103})$$

By combining the inequalities (S.92), (S.97) and (S.103), we obtain

$$\begin{aligned} \sum_{s=1}^{\tilde{n}} \mathcal{L}(d_{L, X_s}) &\leq \sum_{i \in L^{(\xi)}} (\mathcal{C}_{\xi, \xi_0}/4) [1+(R+1)/\xi]^{D-1} e^{-(R+1)/\xi_0} \\ &= |L^{(\xi)}| (\mathcal{C}_{\xi, \xi_0}/4) [1+(R+1)/\xi]^{D-1} e^{-(R+1)/\xi_0}. \end{aligned} \quad (\text{S.104})$$

By applying the above inequality to (S.90), we obtain the upper bound of $\|O_L(H_0, t) - O_L(H_0, t, L[R])\|$. We combine the upper bound of $\|O_L(H_0, t) - O_L(H_0, t, L[R])\|$ with the inequality (S.85) by choosing $R = d_{L, L'} - 1$. Then, we arrive at the inequality (S.67). This completes the proof. \square

B. Proof of Proposition 5

First of all, because the proof of (S.65) does not rely on $\text{diam}(\mathcal{X})$ and the form of $\mathcal{L}(x)$, we obtain the inequality (S.70) in exactly the same way.

Our task is to prove the inequality (S.71). We start from the inequality (S.85):

$$\|[O_L(H_0, t), O_{L'}]\| \leq 2\|O_L(H_0, t) - O_L(H_0, t, L[d_{L, L'} - 1])\|. \quad (\text{S.105})$$

We obtain the same inequality as (S.90) for an arbitrary integer R as follows:

$$\begin{aligned} \|O_L(H_0, t) - O_L(H_0, t, L[R])\| &\leq \int d\mu(U_{X_1}) \int d\mu(U_{X_2}) \cdots \int d\mu(U_{X_{\tilde{n}}}) \|[O_L(H_0, t), U_{L[R]^c}]\| \\ &\leq \sum_{s=1}^{\tilde{n}} \sup_{U_{X_s}} (\|[O_L(H_0, t), U_{X_s}]\|) \\ &\leq \sum_{s=1}^{\tilde{n}} \mathcal{F}(t, L, X_s) \mathcal{L}(d_{L, X_s}) \leq \tilde{\mathcal{F}}(t, L) \sum_{s=1}^{\tilde{n}} \mathcal{L}(d_{L, X_s}), \end{aligned} \quad (\text{S.106})$$

where $\tilde{n} = |(L[R]^c)^{(\xi)}|$, the third inequality is derived from the Lieb-Robinson bound (S.69), and the last inequality is derived from the definition of $\tilde{\mathcal{F}}(t, L)$ in Eq. (S.72).

For the summation of $\mathcal{L}(d_{L, X_s})$, we can utilize the same inequality as (S.92) and (S.97). We then obtain

$$\sum_{s=1}^{\tilde{n}} \mathcal{L}(d_{L, X_s}) \leq \sum_{i \in L^{(\xi)}} \sum_{j \in \Lambda^{(\xi)} : d_{i,j} \geq R+1-2\xi} \mathcal{L}(d_{i,j} - 2\xi). \quad (\text{S.107})$$

In order to estimate the summation with respect to j , we utilize the following lemma (see Sec. XIB1):

Lemma 2. *Let $f(z)$ be an arbitrary function and C_{f, x_0} ($x_0 \geq \xi$) be a constant such that*

$$C_{f, x_0} = \sup_{z \in \mathbb{R} | z \geq x_0} \left(\frac{z^{D+1} f(z)}{x_0^{D+1} f(x_0)} \right). \quad (\text{S.108})$$

Then, for an arbitrary $i \in \Lambda$, we obtain the following upper bound:

$$\sum_{j \in \Lambda^{(\xi)} : d(i,j) \geq x_0} f(d_{i,j}) \leq 2^{D+1} C_{f, x_0} \gamma D \xi^{-D} x_0^D f(x_0). \quad (\text{S.109})$$

In the following, we set

$$\begin{aligned} x_0 &= R + 1 - 2\xi, \quad f(z) = \mathcal{L}(z - 2\xi) = \frac{\log^p(z - 2\xi + 1)}{[z - (2 + \kappa)\xi]^{\alpha_0}}, \\ C_{f, x_0} &= \sup_{z \in \mathbb{R} | z \geq R+1-2\xi} \left(\frac{z^{D+1}}{(R+1-2\xi)^{D+1}} \cdot \frac{\mathcal{L}(z - 2\xi)}{\mathcal{L}(R+1-4\xi)} \right) \end{aligned} \quad (\text{S.110})$$

in the above lemma. Note that $x_0 \geq \xi$ is satisfied from the condition of $R \geq (\kappa + 5)\xi$. We then obtain from the inequality (S.109)

$$\begin{aligned} \sum_{j \in \Lambda^{(\xi)} : d_{i,j} \geq R+1-2\xi} \mathcal{L}(d_{i,j} - 2\xi) &\leq 2^{D+1} \xi^{-D} C_{f, x_0} \gamma D (R+1-2\xi)^D \frac{\log^p(R-4\xi+2)}{[R+1-(4+\kappa)\xi]^{\alpha_0}} \\ &\leq 2^{D+1} \xi^{-D} C_{f, x_0} C'_{R, \kappa, D} \gamma D \frac{\log^p(R+2)}{[R+1-(4+\kappa)\xi]^{\alpha_0-D}}, \end{aligned} \quad (\text{S.111})$$

where we define

$$C'_{R, \kappa, D} := \frac{(R+1-2\xi)^D}{[R+1-(4+\kappa)\xi]^D}. \quad (\text{S.112})$$

We then discuss the values of $C'_{R, \kappa, D}$ and C_{f, x_0} in more details. First, from the condition of $R \geq (\kappa + 5)\xi$ in this proposition, we have

$$C'_{R, \kappa, D} \leq \frac{(\kappa\xi + 3\xi + 1)^D}{(\xi + 1)^D} \leq (\kappa + 3)^D. \quad (\text{S.113})$$

Note that for $x \leq y$ we have $(y/x)^D \leq [(y-\delta)/(x-\delta)]^D$ ($\delta > 0$). Second, we notice that $x_0 = R+1-2\xi \geq (\kappa+3)\xi$ due to $R \geq (\kappa+5)\xi$. Also, because C_{f,x_0} monotonically decreases with x_0 , we have

$$\begin{aligned} C_{f,x_0} &\leq C_{f,(\kappa+3)\xi} = \sup_{z \in \mathbb{R} | z \geq (\kappa+3)\xi} \left(\frac{z^{D+1}}{[(\kappa+3)\xi]^{D+1}} \cdot \frac{\mathcal{L}(z-2\xi)}{\mathcal{L}(\kappa\xi+\xi)} \right) \\ &= \sup_{z \in \mathbb{R} | z \geq (\kappa+3)\xi} \left(\frac{z^{D+1}}{[(\kappa+3)\xi]^{D+1}} \cdot \frac{\log^p(z-2\xi+1)}{[z-(\kappa+2)\xi]^{\alpha_0}} \cdot \frac{[(\kappa+1)\xi]^{\alpha_0}}{\log^p(\kappa\xi+\xi+1)} \right) \\ &\leq (\kappa+3)^{\alpha_0-D-1} \sup_{z \in \mathbb{R} | z \geq (\kappa+3)\xi} \left(\frac{(z/\xi)^{D+1} \log^p(z-2\xi+1)}{[z/\xi-\kappa-2]^{\alpha_0} \log^p(\kappa\xi+\xi+1)} \right). \end{aligned} \quad (\text{S.114})$$

Then, by upper-bounding

$$\frac{\log(z-2\xi+1)}{\log(\kappa\xi+\xi+1)} \leq \frac{\log(z/\xi-2+1/\xi)+\log(\xi)}{\log(\xi+1)} \leq \frac{\log(z/\xi)}{\log(2)}+1 \leq 2\log(z/\xi)+1, \quad (\text{S.115})$$

we have

$$C_{f,x_0} \leq C_{f,(\kappa+3)\xi} \leq (\kappa+3)^{\alpha_0-D-1} \sup_{z \in \mathbb{R} | z \geq \kappa+3} \left(\frac{z^{D+1}[2\log(z)+1]^p}{(z-\kappa-2)^{\alpha_0}} \right) =: (\kappa+3)^{\alpha_0-D-1} \tilde{C}_{\kappa,p,\alpha_0}. \quad (\text{S.116})$$

By applying the inequalities (S.113) and (S.116) to the inequality (S.111), we finally obtain

$$\sum_{j \in \Lambda^{(\xi)} : d_{i,j} \geq R+1-2\xi} \mathcal{L}(d_{i,j}-2\xi) \leq \tilde{C}_{\kappa,p,\alpha_0} 2^{D+1} \xi^{-D} \gamma D (\kappa+3)^{\alpha_0-1} \frac{\log^p(R+2)}{[R+1-(4+\kappa)\xi]^{\alpha_0-D}}. \quad (\text{S.117})$$

By combining the above inequality with (S.107), we have

$$\sum_{s=1}^{\tilde{n}} \mathcal{L}(d_{L,X_s}) \leq |L^{(\xi)}| \tilde{C}_{\kappa,p,\alpha_0} 2^{D+1} \xi^{-D} \gamma D (\kappa+3)^{\alpha_0-1} \frac{\log^p(R+2)}{[R+1-(4+\kappa)\xi]^{\alpha_0-D}}. \quad (\text{S.118})$$

The above inequality with (S.106) reduces the inequality (S.105) to the main inequality (S.71) by choosing $R = d_{L,L'} - 1$. This completes the proof. \square

1. Proof of Lemma 2

From the definition of C_{f,x_0} , the following summation with respect to j is bounded from above by

$$\sum_{j \in \Lambda^{(\xi)} : d_{i,j} \geq x_0} f(d_{i,j}) = f(x_0) x_0^{D+1} \sum_{j \in \Lambda^{(\xi)} : d_{i,j} \geq x_0} \frac{f(d_{i,j}) d_{i,j}^{D+1}}{f(x_0) x_0^{D+1}} d_{i,j}^{-D-1} \leq C_{f,x_0} f(x_0) x_0^{D+1} \sum_{j \in \Lambda^{(\xi)} : d_{i,j} \geq x_0} d_{i,j}^{-D-1}, \quad (\text{S.119})$$

where i is an arbitrary site ($i \in \Lambda$). Then, by using the inequality (S.19), we obtain

$$\begin{aligned} \sum_{j \in \Lambda^{(\xi)} : d_{i,j} \geq x_0} d_{i,j}^{-D-1} &\leq \sum_{s=0}^{\infty} \sum_{\substack{j \in \Lambda^{(\xi)} \\ x_0+s\xi \leq d_{i,j} < x_0+(s+1)\xi}} d_{i,j}^{-D-1} \\ &\leq 2\gamma D \sum_{s=0}^{\infty} (x_0+s\xi)^{-D-1} [2(x_0+s\xi)/\xi]^{D-1} \\ &\leq 2^D \gamma D \xi^{-D+1} \sum_{s=0}^{\infty} (x_0+s\xi)^{-2} \leq 2^{D+1} \gamma D \xi^{-D} x_0^{-1}, \end{aligned} \quad (\text{S.120})$$

where the last inequality is derived from

$$\sum_{s=0}^{\infty} (x_0+s\xi)^{-2} = x_0^{-2} + \sum_{s=1}^{\infty} (x_0+s\xi)^{-2} \leq x_0^{-2} + \int_0^{\infty} (x_0+x\xi)^{-2} dx = x_0^{-2} + (\xi x_0)^{-1} \leq 2(\xi x_0)^{-1}. \quad (\text{S.121})$$

Note that $x_0 \geq \xi$ from the assumption. By combining the inequalities (S.119) and (S.120), we obtain the inequality (S.109). This completes the proof. \square

XII. PROOF OF THEOREM 2: MIDDLE-RANGE LIEB-ROBINSON BOUND

A. statement

Theorem 2 (Middle-range Lieb-Robinson bound) *We consider the Hamiltonian H which satisfies Assumption (S.7) with $\alpha > 2D + 1$. Then, for arbitrary ℓ , the Hamiltonian $H_{\leq \ell}$ satisfies the $\mathcal{G}_{\leq \ell}(x, t, \mathcal{X})$ -Lieb-Robinson bound with*

$$\mathcal{G}_{\leq \ell}(x, t, \mathcal{X}) = \tilde{C}_0 |\mathcal{X}|^2 (1 + x/\ell)^{D-1} e^{-2(x-v^*|t|)/\ell}, \quad (\text{S.122})$$

where v^* is a constant which depends only on the parameters $\{D, g_0, \alpha, \gamma\}$ and

$$\tilde{C}_0 := \frac{4}{3} e^{10/3} 15^D D! \gamma \zeta_2, \quad \zeta_2 := 16e^3 \gamma^2 \cdot 45^D. \quad (\text{S.123})$$




FIG. 8. Outline of the proof on the middle-range Lieb-Robinson bound 1. For the proof, we first decompose the length ℓ into q^* pieces as $\{\ell_q\}_{q=1}^{q^*}$ with $\ell_{q^*} = \ell$. We choose them so that the length scale ℓ_q increases double exponentially with respect to q [see Ineq. (S.130)]. Our purpose is to estimate the Lieb-Robinson bound for $e^{-iH_{1:q^*}t}$, which is equal to $e^{-iH_{\leq \ell}t}$ from the definition (S.150). The main task is to derive the Lieb-Robinson bound for $e^{-iH_{1:q}t}$, which we denote by $\mathcal{G}_q(x, t, \mathcal{X})$, in the form of Eq. (S.152) with (S.153). If we can derive the bound, Theorem 2 is immediately derived by choosing $q = q^*$. In order to prove Eq. (S.152), we use the induction method. The case of $q = 1$ is derived by using the standard technique to obtain the short-range Lieb-Robinson bound (see Lemma 3 with its proof in Sec. XIII C). The derivation of the case q from the assumption of the case $q - 1$ is a bit intricate and we give the outline in Fig. 9.

[Derivation of the case q under the assumption of the case $q-1$]




FIG. 9. Outline of the proof on the middle-range Lieb-Robinson bound 2. In order to derive the case q from the assumption (S.156) for the case of $q-1$, we decompose the unitary operator $e^{-iH_{1:q}t}$ as in Eq. (S.166): $e^{-iH_{1:q-1}t}$ and $\mathcal{T}e^{-\int_0^t H_q(H_{1:q-1}, \tau) d\tau}$. From the assumption, the unitary operator $e^{-iH_{1:q-1}t}$ satisfies the $\mathcal{G}_{q-1}(x, t, \mathcal{X})$ -Lieb-Robinson bound in the form of Eq. (S.152). Then, the time-evolved Hamiltonian $H_q(H_{1:q-1}, t)$ has the interaction length of $\ell_q + v_{q-1}\mathcal{O}(t)$, which is of order of $\mathcal{O}(\ell_q)$ for $t \lesssim \ell_q/v_{q-1}$. We thus consider the time range $t \leq \Delta t_q \approx \ell_q/v_{q-1}$ [see Eq. (S.157) in detail] for the first, and extend the time scale afterward [see Eq. (S.187)]. In order to obtain the Lieb-Robinson bound for $\mathcal{T}e^{-\int_0^t H_q(H_{1:q-1}, \tau) d\tau}$ for $t \leq \Delta t_q$, we take two steps: i) in the first step, we estimate the quasi-locality of $H_q(H_{1:q-1}, \tau)$ by using Proposition 6 (the proof is given in Sec. XIII A), ii) in the second step, based on the quasi-locality, we derive the Lieb-Robinson bound (S.183) for $\mathcal{T}e^{-\int_0^t H_q(H_{1:q-1}, \tau) d\tau}$ by using Proposition 7 (the proof is given in Sec. XIII B). After obtaining the Lieb-Robinson bounds for $e^{-iH_{1:q-1}t}$ and $\mathcal{T}e^{-\int_0^t H_q(H_{1:q-1}, \tau) d\tau}$ with $t \leq \Delta t_q$, we extend the time scale to generic t . In order to connect all the Lieb-Robinson bounds appropriately, we first restrict the subsets X and Y such that $\text{diam}(\mathcal{X}) \leq \ell_q$ and $\text{diam}(\mathcal{Y}) \leq \ell_q$, and remove the restrictions afterward by using Theorem 4. The connections rely on Proposition 8, which yields the inequality (S.193) (see Sec. XII E for the proof). By combining all the ingredients, we derive the Lieb-Robinson bound for the case of q .

B. Decomposition of the length scale

We define a set of the length scales $\{\ell_q\}_{q=1}^{q^*}$. First, ℓ_1 is defined as a constant which depends only on the parameters $\{D, g_0, \alpha, \gamma\}$. We show the conditions for ℓ_1 in Sec. XII C. Other length scales $\{\ell_q\}_{q=2}^{q^*}$ are defined as

$$\ell_q := (\ell_{q-1})^{1+\eta_q} \quad (\eta \leq \eta_q \leq \bar{\eta}), \quad (\text{S.124})$$

$$\bar{\eta} := \frac{\alpha - (2D+1)}{D+2}, \quad \eta := \sqrt{1+\bar{\eta}} - 1, \quad (\text{S.125})$$

where we choose $\{\eta_q\}_{q=2}^{q^*}$ appropriately such that $\{\ell_q\}_{q=2}^{q^*}$ become integer and for $q^* \geq 2$ the length ℓ_{q^*} satisfies

$$\ell_{q^*} = \ell_1^{(1+\eta_2)(1+\eta_3)\cdots(1+\eta_{q^*})} = \ell := \ell_1^{p_\ell}. \quad (\text{S.126})$$

From the condition $\eta \leq \eta_q \leq \bar{\eta}$ ($1 \leq q \leq q^*$), we have

$$\frac{\log(p_\ell)}{\log(1+\bar{\eta})} \leq q^* - 1 \leq \frac{\log(p_\ell)}{\log(1+\eta)} = \frac{2\log(p_\ell)}{\log(1+\bar{\eta})}. \quad (\text{S.127})$$

Thus, as long as

$$p_\ell \geq 1 + \bar{\eta} \quad \text{or} \quad \ell \geq \ell_1^{1+\bar{\eta}}, \quad (\text{S.128})$$

there exists parameters $\{\eta_q\}_{q=2}^{q^*}$ such that $\ell_{q^*} = \ell$. From the definition, we note that

$$\ell_{q-1}^\eta \leq \frac{\ell_q}{\ell_{q-1}} = \ell_{q-1}^{\eta_q} \leq \ell_{q-1}^{\bar{\eta}} \quad (\text{S.129})$$

and

$$\ell_q \geq \ell_1^{(1+\eta)^{q-1}}. \quad (\text{S.130})$$

Therefore, the length scale ℓ_q increases double exponentially with respect to q .

We then define Hamiltonian H_q which picks up all the interaction terms with interaction length from ℓ_{q-1} to ℓ_q :

$$\begin{aligned} H_q &:= \sum_{Z: \ell_{q-1} \leq \text{diam}(Z) < \ell_q} h_Z \quad (q = 1, 2, \dots, q^* - 1), \\ H_{q^*} &:= \sum_{Z: \ell_{q^*-1} \leq \text{diam}(Z) \leq \ell_{q^*}} h_Z, \end{aligned} \quad (\text{S.131})$$

where we set $\ell_0 = 1$. From the assumption (S.7), we obtain

$$\sup_{i \in \Lambda} \sum_{\substack{Z: Z \ni i \\ \ell_{q-1} \leq \text{diam}(Z) < \ell_q}} \|h_Z\| \leq \sup_{i \in \Lambda} \sum_{\substack{Z: Z \ni i \\ \text{diam}(Z) \geq \ell_{q-1}}} \|h_Z\| \leq g_q \quad (\text{S.132})$$

with

$$g_q := g \ell_{q-1}^{-\alpha+D}. \quad (\text{S.133})$$

From the definition (S.125), we have

$$\ell_q^{D+1} g_q \leq g \ell_{q-1}^{(D+1)(1+\eta_q)-\alpha+D} = g \ell_{q-1}^{(D+2)\eta_q - \eta_q - \alpha + 2D+1} \leq g \ell_{q-1}^{-\eta_q} \leq g \ell_{q-1}^{-\eta}, \quad (\text{S.134})$$

where in the second inequality we use $\eta_q \leq \bar{\eta} = \frac{\alpha-(2D+1)}{D+2}$.

C. Conditions for ℓ_1

In the proof, we adopt various kinds of conditions for ℓ_1 . We choose ℓ_1 so that the following conditions for ℓ_q ($q \geq 1$) are satisfied. Some of the conditions include ℓ_q , but they reduce to lower bounds for ℓ_1 because of Eq. (S.124). We note that all the conditions depend only on the parameters $\{D, g_0, \alpha, \gamma\}$. We summarize all the conditions in the following:

1. Used in the inequality (S.159):

$$\frac{1}{1 - \delta_{q-1}} \leq 2, \quad \text{with} \quad \delta_{q-1} = \frac{6 \log[\tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D}] + 6(D-1) \log(1 + \ell_{q-1}^{\eta_q}/12)}{\ell_{q-1}^{\eta_q}}, \quad (\text{S.135})$$

where $\tilde{\mathcal{C}}_0$ is defined in Eq. (S.123).

2. Used in the inequalities (S.171) and (S.180):

$$\ell_{q-1}^{\eta_q} \geq \ell_{q-1}^\eta \geq \ell_1^\eta \geq \max(24 \log(2\gamma^2), 48). \quad (\text{S.136})$$

3. Used in the inequality (S.182):

$$\frac{e^2 g \gamma^2 (20D)^D}{3\ell_1 v_0} \leq 1, \quad (\text{S.137})$$

where $v_0 = \frac{2e^3 g \gamma (\alpha - 2D)}{\alpha - 2D - 1}$ as defined in Eq. (S.148).

4. Used in the inequality (S.196):

$$\frac{\ell_{q-1} \log(\zeta_1)}{\xi_q} \leq 1 \quad \text{or} \quad \frac{\ell_{q-1}^\eta}{\log(\ell_{q-1})} \geq \frac{\eta \log(\zeta_1)}{16}, \quad (\text{S.138})$$

where $\zeta_1 = 2^{D+1} e^2 \gamma^2 D!$ and $\xi_q := \frac{16\ell_q}{\eta \log(\ell_{q-1})}$ as defined in Eq. (S.192) and Eq. (S.194), respectively.

5. Used in the inequalities (S.206) and (S.216):

$$\frac{1}{18\ell_{q-1}} \geq \frac{\eta \log(\ell_{q-1})}{8\ell_q} = \frac{2}{\xi_q} \quad \text{or} \quad \frac{\ell_{q-1}^\eta}{\log(\ell_{q-1})} \geq \frac{9\eta}{4}, \quad (\text{S.139})$$

where $\xi_q := \frac{16\ell_q}{\eta \log(\ell_{q-1})}$ as defined in Eq. (S.194).

6. Used in the inequality (S.208):

$$\frac{\ell_q}{\xi_q} = \frac{\eta \log(\ell_{q-1})}{16} \geq 3 \quad \text{or} \quad \log(\ell_{q-1}) \geq \log(\ell_1) \geq \frac{48}{\eta}, \quad (\text{S.140})$$

where $\xi_q := \frac{16\ell_q}{\eta \log(\ell_{q-1})}$ as defined in Eq. (S.194).

7. Used in the inequality (S.212):

$$\frac{2\ell_{q-1} \log(\zeta_1)}{\ell_q} = \frac{2 \log(\zeta_1)}{\ell_{q-1}^\eta} \leq \frac{2 \log(\zeta_1)}{\ell_1^\eta} \leq 1, \quad (\text{S.141})$$

where $\zeta_1 = 2^{D+1} e^2 \gamma^2 D!$ as defined in Eq. (S.192).

8. Used in the inequality (S.199):

$$\ell_1 \geq 3. \quad (\text{S.142})$$

9. Used in the inequality (S.221):

$$\frac{2}{e} \ell_{q-1}^{-\eta/2} \leq \frac{1}{\zeta_1} \quad \text{or} \quad \ell_{q-1} \geq \ell_1 \geq \left(\frac{4\zeta_1}{e} \right)^{2/\eta}, \quad (\text{S.143})$$

where $\zeta_1 = 2^{D+1} e^2 \gamma^2 D!$ as defined in Eq. (S.192).

10. Used in the inequality (S.326):

$$\frac{\log(\tilde{\mathcal{C}}_0) + (D-1) \log(v_0 \ell_1 + 1)}{2v_0 \ell_1} \leq \frac{1}{2}, \quad (\text{S.144})$$

where $v_0 = \frac{2e^3 g \gamma (\alpha - 2D)}{\alpha - 2D - 1}$ as defined in Eq. (S.148).

D. Proof of Theorem 2 (see also Figs. 8 and 9)

We first consider the case of $\ell \leq \ell_1^{1+\bar{\eta}}$, where $\bar{\eta}$ was defined in Eq. (S.125). In this case, we cannot define $q^* (\geq 2)$ such that $\ell_{q^*} = \ell$ in Eq. (S.125), but the Hamiltonian $H_{\leq \ell}$ has a finite interaction length which is independent of t . As a general setup, we consider the Lieb-Robinson bound by a Hamiltonian which has a length scale at most ξ . As long as $\xi = \mathcal{O}(1)$, we trivially obtain the Lieb-Robinson velocity of order $\mathcal{O}(1)$, whereas the ξ -dependence of the velocity may be non-linear with respect to ξ . Our purpose here is to prove that the Lieb-Robinson velocity is at most linear to ξ as long as $\alpha > 2D + 1$. We prove the following lemma (the proof is given in Sec. XIII C):

Lemma 3. *Let \tilde{H} be a Hamiltonian such that*

$$\tilde{H} = \sum_{Z \subset \Lambda: \text{diam}(Z) \leq \xi} h_Z \quad (\text{S.145})$$

with the same condition as (S.7), namely

$$\sup_{i \in \Lambda} \sum_{\substack{Z: Z \ni i \\ r \leq \text{diam}(Z) \leq \xi}} \|h_Z\| \leq gr^{-\alpha+D} \quad \text{for } r \leq \xi. \quad (\text{S.146})$$

Then, the Hamiltonian \tilde{H} satisfies $\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X})$ -Lieb-Robinson bound with

$$\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X}) = \min \left(2|\mathcal{X}| \left(\frac{v_0|t|}{e^2 \lceil x/\xi \rceil} \right)^{\lceil x/\xi \rceil}, 2 \right) \leq 2|\mathcal{X}| e^{-2(x-\xi v_0|t|)/\xi} \quad (\text{S.147})$$

with

$$v_0 := \frac{2e^3 g \gamma (\alpha - 2D)}{\alpha - 2D - 1}. \quad (\text{S.148})$$

We note that the same inequality holds for time-dependent Hamiltonians.

From Lemma 3 or the inequality (S.147), we obtain for $\ell \leq \ell_1^{1+\bar{\eta}}$

$$\mathcal{G}_{\leq \ell}(x, t, \mathcal{X}) = 2|\mathcal{X}| e^{-(x-\ell v_0|t|)/\ell} \leq 2|\mathcal{X}| e^{-2(x-\ell_1^{1+\bar{\eta}} v_0|t|)/\ell}. \quad (\text{S.149})$$

Because $\ell_1^{1+\bar{\eta}} v_0$ depends only on the parameters $\{D, g_0, \alpha, \gamma\}$, we obtain the inequality (S.122) by choosing $v^* = \ell_1^{1+\bar{\eta}} v_0$ for the case of $\ell \leq \ell_1^{1+\bar{\eta}}$.

We next focus on the case of $\ell > \ell_1^{1+\bar{\eta}}$, where we can define $q^* \geq 2$ such that $\ell_{q^*} = \ell$. We here define the Hamiltonian $H_{1:q}$ as follows:

$$H_{1:q} := \sum_{s=1}^q H_s. \quad (\text{S.150})$$

For the proof, we need to estimate the Lieb-Robinson bound for the Hamiltonian $H_{1:q^*}$, which is equal to $H_{\leq \ell}$. We define that $\{H_{1:q}\}_{q=1}^{q^*}$ satisfy the $\mathcal{G}_q(x, t, \mathcal{X})$ -Lieb-Robinson bound for $q = 1, 2, \dots, q^*$, namely,

$$\frac{\| [e^{iH_{1:q}t} O_L e^{-iH_{1:q}t}, O_{L'}] \|}{\|O_L\| \cdot \|O_{L'}\|} \leq \mathcal{G}_q(x, t, L), \quad x = d_{L, L'} \quad (\text{S.151})$$

for arbitrary operators O_L and $O_{L'}$. We aim to prove the Lieb-Robinson bound for $H_{1:q}$ in the form of

$$\mathcal{G}_q(x, t, \mathcal{X}) = \tilde{\mathcal{C}}_0 |\mathcal{X}|^2 (1 + x/\ell_q)^{D-1} e^{-2(x-v_q|t|)/\ell_q}, \quad (\text{S.152})$$

where $\tilde{\mathcal{C}}_0$ is defined in Eq. (S.123), and v_q is given by the recurrence relation as follows:

$$\begin{aligned} v_1 &= \ell_1 v_0, \quad v_q = v_{q-1} \left(\frac{1}{1 - \delta_{q-1}} + \frac{192 \log(\zeta_2)}{\eta \log(\ell_{q-1})} \right), \\ \delta_{q-1} &= \frac{6 \log[\tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D}] + 6(D-1) \log(1 + \ell_{q-1}^{\eta_q}/12)}{\ell_{q-1}^{\eta_q}}, \quad \zeta_2 = 16e^3 \gamma^2 \cdot 45^D \end{aligned} \quad (\text{S.153})$$

with $v_0 = \frac{2e^3 g \gamma (\alpha - 2D)}{\alpha - 2D - 1}$ as defined in Eq. (S.148). By remembering that the ℓ_q is lower-bounded by a double exponential function with respect to q as in (S.130), we can ensure

$$v_\infty = \lim_{q \rightarrow \infty} v_q < \infty, \quad (\text{S.154})$$

and v_∞ depends only on the parameters $\{D, g_0, \alpha, \gamma\}$. From the inequality (S.149) and Eq. (S.154), we can choose $v^* = \max(\ell_1^{1+\eta} v_0, v_\infty)$. We thus prove the theorem.

In the following, we prove the inequality (S.152) by induction method. We first consider the case of $q = 1$. From the inequality (S.147), we obtain

$$\mathcal{G}_1(x, t, \mathcal{X}) = 2|\mathcal{X}|e^{-2(x-\ell_1 v_0|t|)/\ell_1}, \quad (\text{S.155})$$

which clearly reduces to the form of (S.152). We then adopt the assumption that the Hamiltonian $H_{1:q-1}$ ($q \geq 2$) satisfies the Lieb-Robinson bound of

$$\mathcal{G}_{q-1}(x, t, \mathcal{X}) = \tilde{\mathcal{C}}_0 |\mathcal{X}|^2 (1 + x/\ell_{q-1})^{D-1} e^{-2(x-v_{q-1}|t|)/\ell_{q-1}}. \quad (\text{S.156})$$

By using the assumption (S.156), we aim to derive the Lieb-Robinson bound for $H_{1:q}$ in the form of (S.152). We first restrict ourselves to

$$t \leq \Delta t_q := \frac{\ell_q}{12v_{q-1}}, \quad (\text{S.157})$$

and consider the generic t afterward [see Eq. (S.187)]. Before going to the proof, we derive the following lemma:

Lemma 4. *For arbitrary $q \geq 2$, the Lieb-Robinson function $\mathcal{G}_{q-1}(x, t, \mathcal{X})$ in (S.156) is reduced to the form of*

$$\mathcal{G}_{q-1}(x, t, \mathcal{X}) = |\mathcal{X}^{(\ell_q)}|^2 e^{-(x-\bar{v}_{q-1}\Delta t_q)/\ell_{q-1}} \quad (t \leq \Delta t_q), \quad (\text{S.158})$$

where we define \bar{v}_{q-1} as

$$\bar{v}_{q-1} := \frac{v_{q-1}}{1 - \delta_{q-1}} \leq 2v_{q-1} \quad (\text{S.159})$$

with

$$\delta_{q-1} = \frac{6 \log[\tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D}] + 6(D-1) \log(1 + \ell_{q-1}^{\eta_q}/12)}{\ell_{q-1}^{\eta_q}}. \quad (\text{S.160})$$

Note that $\bar{v}_{q-1} \leq 2v_{q-1}$ is a consequence from the condition (S.135).

Proof of Lemma 4. From Eq. (S.23) and the definition of δ_{q-1} , the Lieb-Robinson bound (S.158) gives a trivial bound for $x \leq v_{q-1}\Delta t_q$. Hence, we only have to consider the range of $x \geq v_{q-1}\Delta t_q$.

We start from the form of $\mathcal{G}_{q-1}(x, t, \mathcal{X})$ as in Eq. (S.156):

$$\mathcal{G}_{q-1}(x, t, \mathcal{X}) = \tilde{\mathcal{C}}_0 |\mathcal{X}|^2 (1 + x/\ell_{q-1})^{D-1} e^{-2(x-v_{q-1}|t|)/\ell_{q-1}}. \quad (\text{S.161})$$

First, from the definition (S.14), we have $X \subseteq (X^{(\ell_q)})[\ell_q]$ for an arbitrary subset $X \subseteq \Lambda$, and hence $|X|$ is upper-bounded as follows:

$$|X| \leq |(X^{(\ell_q)})[\ell_q]| \leq \sum_{i \in X^{(\ell_q)}} |i[\ell_q]| \leq \gamma (2\ell_q)^D |X^{(\ell_q)}|, \quad (\text{S.162})$$

where we use the inequality (S.17). Then, we have

$$\mathcal{G}_{q-1}(x, t, \mathcal{X}) \leq \mathcal{G}_{q-1}(x, \Delta t_q, \mathcal{X}) \leq \tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D} |X^{(\ell_q)}|^2 (1 + x/\ell_{q-1})^{D-1} e^{-2(x-v_{q-1}\Delta t_q)/\ell_{q-1}}, \quad (\text{S.163})$$

where we use the condition $t \leq \Delta t_q$ in the first inequality.

Second, by using $x \geq v_{q-1}\Delta t_q$, we obtain the upper bound of

$$\begin{aligned} & \tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D} |X^{(\ell_q)}|^2 (1 + x/\ell_{q-1})^{D-1} e^{-2(x-v_{q-1}\Delta t_q)/\ell_{q-1}} \\ &= \exp \left\{ \frac{-2x}{\ell_{q-1}} \left[1 - \frac{\ell_{q-1} \log[\tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D} (1 + x/\ell_{q-1})^{D-1}]}{2x} \right] \right\} e^{2v_{q-1}\Delta t_q/\ell_{q-1}} \\ &\leq \exp \left\{ \frac{-2x}{\ell_{q-1}} \left[1 - \frac{\ell_{q-1} \log[\tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D} (1 + v_{q-1}\Delta t_q/\ell_{q-1})^{D-1}]}{2v_{q-1}\Delta t_q} \right] \right\} e^{2v_{q-1}\Delta t_q/\ell_{q-1}} \\ &= \exp \left\{ \frac{-2(1-\delta_{q-1})}{\ell_{q-1}} \left[x - (1-\delta_{q-1})^{-1} v_{q-1} \right] \right\}, \end{aligned} \quad (\text{S.164})$$

where in the last equation we use the definitions $\Delta t_q = \ell_q/(12v_{q-1})$ and $\ell_q = \ell_{q-1}^{1+\eta_q}$ to derive

$$\begin{aligned} \frac{\ell_{q-1} \log[\tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D} (1 + v_{q-1} \Delta t_q / \ell_{q-1})^{D-1}]}{2v_{q-1} \Delta t_q} &= \frac{6 \log[\tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D}] + 6(D-1) \log[1 + \ell_q / (12\ell_{q-1})]}{\ell_q / \ell_{q-1}} \\ &= \frac{6 \log[\tilde{\mathcal{C}}_0 \gamma^2 (2\ell_q)^{2D}] + 6(D-1) \log(1 + \ell_{q-1}^{\eta_q} / 12)}{\ell_{q-1}^{\eta_q}} = \delta_{q-1}. \end{aligned} \quad (\text{S.165})$$

From the condition (S.135), we have $1/(1 - \delta_{q-1}) \leq 2$, which yields $2(1 - \delta_{q-1}) \geq 1$. Therefore, the inequality (S.164) reduces to (S.158). This completes the proof. \square

We now consider the unitary operator $e^{-iH_{1:q}t}$ for $t \leq \Delta t_q$ and start from the decomposition of

$$e^{-iH_{1:q}t} = e^{-iH_{1:q-1}t} \mathcal{T} e^{-\int_0^t e^{iH_{1:q-1}\tau} H_q e^{iH_{1:q-1}-\tau} d\tau} = e^{-iH_{1:q-1}t} \mathcal{T} e^{-\int_0^t H_q(H_{1:q-1}, \tau) d\tau}. \quad (\text{S.166})$$

We aim to derive the Lieb-Robinson bound for $\mathcal{T} e^{-\int_0^t H_q(H_{1:q-1}, \tau) d\tau}$, which we characterize by the function $\tilde{\mathcal{G}}_q(x, t, \mathcal{X})$. Note that the Lieb-Robinson bound for $e^{-iH_{1:q-1}t}$ has been already given by the function $\mathcal{G}_{q-1}(x, t, \mathcal{X})$ as in Eq. (S.158). In order to estimate $\tilde{\mathcal{G}}_q(x, t, \mathcal{X})$, we first estimate the quasi-locality^{*1} of $H_q(H_{1:q-1}, \tau)$. We prove the following proposition (see Sec. XIII A for the proof):

Proposition 6. *Let $\tilde{\ell}_q$ be a length scale such that*

$$2\gamma^2 e^{\bar{v}_{q-1}\tau/\ell_{q-1}} e^{-\tilde{\ell}_q/(2\ell_{q-1})} \leq 1 \quad (\text{S.167})$$

for $\tau \leq \Delta t_q$. Then, under the assumption of (S.158), we can give a decomposition of the time-evolved Hamiltonian $H_q(H_{1:q-1}, \tau)$ in the form of

$$H_q(H_{1:q-1}, \tau) = \sum_{Z: \text{diam}(Z) \leq \ell_q} \sum_{s=1}^{\infty} h_{\tau, Z[s\tilde{\ell}_q]} \quad (\text{S.168})$$

with

$$\sum_{\substack{Z: \text{diam}(Z) \leq \ell_q \\ Z[s\tilde{\ell}_q] \cap X \neq \emptyset}} \|h_{\tau, Z[s\tilde{\ell}_q]}\| \leq (2\tilde{\ell}_q)^D g_q \gamma |X^{(s_0\tilde{\ell}_q)}|(s + s_0)^D e^{-(s-1)\tilde{\ell}_q/(2\ell_{q-1})} \quad (\text{S.169})$$

for an arbitrary subset $X \subseteq \Lambda$, where s_0 can be arbitrarily chosen and $h_{\tau, Z[s\tilde{\ell}_q]}$ is an operator which acts on the subset $Z[s\tilde{\ell}_q]$.

In order that the condition (S.167) is satisfied for $\tau \leq \Delta t_q = \ell_q/(12v_{q-1})$, we choose

$$\tilde{\ell}_q = 5\ell_q/12. \quad (\text{S.170})$$

This choice implies the condition (S.167) as follows:

$$2\gamma^2 e^{\bar{v}_{q-1}\tau/\ell_{q-1}} e^{-\tilde{\ell}_q/(2\ell_{q-1})} \leq 2\gamma^2 e^{2v_{q-1}[\ell_q/(12v_{q-1})]/\ell_{q-1}} e^{-(5\ell_q/12)/(2\ell_{q-1})} = e^{-\ell_{q-1}^{\eta_q}/24 + \log(2\gamma^2)} < 1, \quad (\text{S.171})$$

where in the first inequality we use $\bar{v}_{q-1} \leq 2v_{q-1}$ which is given in (S.159), and the second inequality $\ell_{q-1}^{\eta_q}/24 \geq \log(2\gamma^2)$ is derived from the condition (S.136). From the choice of $\tilde{\ell}_q = 5\ell_q/12$, we have

$$\text{diam}(Z[s\tilde{\ell}_q]) = \text{diam}(Z) + 2s\tilde{\ell}_q \leq 11s\ell_q/6 \quad \text{for} \quad \text{diam}(Z) \leq \ell_q. \quad (\text{S.172})$$

Also, by replacing $s_0 \rightarrow 22s_0/5$, we have

$$(2\tilde{\ell}_q)^D g_q \gamma |X^{(s_0\tilde{\ell}_q)}|(s + s_0)^D \rightarrow (2\tilde{\ell}_q)^D g_q \gamma |X^{(22s_0\tilde{\ell}_q/5)}|(s + 22s_0/5)^D \leq (10\tilde{\ell}_q)^D g_q \gamma |X^{(11s_0\ell_q/6)}|(s + s_0)^D. \quad (\text{S.173})$$

Note that s_0 can be arbitrarily chosen.

Moreover, we can prove the following proposition on the Lieb-Robinson bound for quasi-local operators in the form of Eq. (S.168) with (S.169) (see Sec. XIII B for the proof):

^{*1} An interaction is said to be quasi-local if the interaction decays rapidly (exponentially or polynomially) with the interaction length.

Proposition 7. Let \tilde{H} be a Hamiltonian such that

$$\tilde{H} = \sum_{Z: Z \subseteq \Lambda} h_Z = \sum_{s=1}^{\infty} \sum_{\text{diam}(Z_s) \leq s\xi} h_{Z_s}. \quad (\text{S.174})$$

For an arbitrary subset X and an arbitrary positive s_0 , we assume

$$\sum_{\substack{\text{diam}(Z_s) \leq s\xi \\ Z_s \cap X \neq \emptyset}} \|h_{Z_s}\| \leq \tilde{g} |X^{(s_0\xi)}| (s + s_0)^D e^{-\mu(s-1)} \quad (\mu > 1). \quad (\text{S.175})$$

Then, the Hamiltonian \tilde{H} satisfies $\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X})$ -Lieb-Robinson bound with

$$\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X}) = e |\mathcal{X}^{(\xi)}| \left[\tilde{v} |t| \exp \left(-\frac{\mu-1}{2} \frac{x}{\xi} + \tilde{v} |t| \right) + \left(\frac{e^2 \tilde{v} |t|}{m_x^*} \right)^{m_x^*} \right] \quad (\text{S.176})$$

with

$$\tilde{v} := 4\tilde{g}\gamma(4D)^D, \quad m_x^* := \left\lfloor \frac{\mu-1}{2\mu} \frac{x}{\xi} \right\rfloor + 1. \quad (\text{S.177})$$

We note that the same inequality holds for time-dependent Hamiltonians.

From Proposition 6, in order to apply Proposition 7 to $H_q(H_{1:q-1}, \tau)$ ($\tau \leq \Delta t_q$), we choose $\{\xi, \mu, \tilde{g}\}$ as

$$\xi = 11\ell_q/6 \leq 2\ell_q, \quad \mu = \frac{\tilde{\ell}_q}{2\ell_{q-1}} \geq \frac{\ell_q}{4\ell_{q-1}} \quad \text{and} \quad \tilde{g} = (10\tilde{\ell}_q)^D g_q \gamma \leq (5\ell_q)^D g_q \gamma, \quad (\text{S.178})$$

where we use the inequalities (S.172) and (S.173) in determining the parameters ξ and \tilde{g} as in (S.175). Note that $\mu \geq \ell_q/(4\ell_{q-1}) = \ell_{q-1}^{\eta_q}/4 > 1$ is ensured from the condition (S.136). From Eq. (S.176), we obtain the Lieb-Robinson function $\tilde{\mathcal{G}}_q(x, t, \mathcal{X})$ for dynamics by the time-evolved Hamiltonian $H_q(H_{1:q-1}, \tau)$ ($0 \leq \tau \leq t$ with $t \leq \Delta t_q$) as follows:

$$\tilde{\mathcal{G}}_q(x, t, \mathcal{X}) \leq e |\mathcal{X}^{(11\ell_q/6)}| \left[\tilde{v}_q |t| \exp \left(-\frac{1-4\ell_{q-1}/\ell_q}{16\ell_{q-1}} x + \tilde{v}_q |t| \right) + \left(\frac{e^2 \tilde{v}_q |t|}{m_x^*} \right)^{m_x^*} \right] \quad (\text{S.179})$$

with

$$\tilde{v}_q := 4\gamma^2(20D\ell_q)^D g_q, \quad m_x^* := \left\lfloor \frac{(1-4\ell_{q-1}/\ell_q)x}{11\ell_q/3} \right\rfloor + 1 \geq \frac{x}{4\ell_q}, \quad (\text{S.180})$$

where we use the condition (S.136) to obtain $\ell_q/\ell_{q-1} = \ell_{q-1}^{\eta_q} \geq 48$, which yields $1 - 4\ell_{q-1}/\ell_q \geq 11/12$.

We further simplify the bound (S.179) as

$$\begin{aligned} \tilde{\mathcal{G}}_q(x, t, \mathcal{X}) &\leq e |\mathcal{X}^{(\ell_q)}| \left[\tilde{v}_q |t| \exp \left(-\frac{11}{192\ell_{q-1}} x + \tilde{v}_q |t| \right) + (e^2 \tilde{v}_q |t|)^{m_x^*/2} \cdot (e^2 \tilde{v}_q |t|)^{m_x^*/2} \right] \\ &\leq e |\mathcal{X}^{(\ell_q)}| \left[\tilde{v}_q |t| \exp \left(-\frac{1}{18\ell_{q-1}} x + \tilde{v}_q |t| \right) + (e^2 \tilde{v}_q |t|)^{1/2} \cdot \exp \left(\frac{\log(e^2 \tilde{v}_q |t|)}{8\ell_q} x \right) \right] \end{aligned} \quad (\text{S.181})$$

for $t \leq \Delta t_q$, where we use $|\mathcal{X}^{(11\ell_q/6)}| \leq |\mathcal{X}^{(\ell_q)}|$. Note that from $\Delta t_q = \ell_q/(12v_{q-1})$ and Def. (S.180) for \tilde{v}_q , we have

$$e^2 \tilde{v}_q \Delta t_q = 4e^2 \gamma^2 (20D\ell_q)^D g_q \cdot \ell_q / (12v_{q-1}) = \frac{e^2 \gamma^2 (20D)^D}{3v_{q-1}} \ell_q^{D+1} g_q \leq \frac{e^2 g \gamma^2 (20D)^D}{3\ell_1 v_0} \ell_{q-1}^{-\eta} \leq \ell_{q-1}^{-\eta}, \quad (\text{S.182})$$

where we use $v_{q-1} \geq v_1 := \ell_1 v_0$ [see Eq. (S.153) for the definition of v_1] and the inequality (S.134) for $\ell_q^{D+1} g_q$ in the first inequality and the condition (S.137) in the second inequality. This reduces the inequality (S.181) to

$$\tilde{\mathcal{G}}_q(x, t, \mathcal{X}) \leq e |\mathcal{X}^{(\ell_q)}| \left[\ell_{q-1}^{-\eta} \exp \left(-\frac{1}{18\ell_{q-1}} x \right) + \ell_{q-1}^{-\eta/2} \cdot \exp \left(\frac{-\eta \log(\ell_{q-1})}{8\ell_q} x \right) \right] \quad \text{for } t \leq \Delta t_q, \quad (\text{S.183})$$

where we use the following bound in deriving the first term:

$$\tilde{v}_q |t| \exp(\tilde{v}_q |t|) \leq \tilde{v}_q \Delta t_q \exp(\tilde{v}_q \Delta t_q) \leq \ell_{q-1}^{-\eta} \frac{\exp(\ell_{q-1}^{-\eta}/e^2)}{e^2} \leq \ell_{q-1}^{-\eta} \frac{\exp(1/e^2)}{e^2} \leq \ell_{q-1}^{-\eta}. \quad (\text{S.184})$$

In the following, by combining $\mathcal{G}_{q-1}(x, t, \mathcal{X})$ and $\tilde{\mathcal{G}}_q(x, t, \mathcal{X})$, we obtain the Lieb-Robinson function $\mathcal{G}_q(x, t, \mathcal{X}, \mathcal{Y})$ for generic time t . For the purpose, we first estimate $\mathcal{G}_q(x, t, \mathcal{X}, \mathcal{Y})$ under the constraints of

$$\text{diam}(\mathcal{X}) \leq \ell_q, \quad \text{diam}(\mathcal{Y}) \leq \ell_q. \quad (\text{S.185})$$

We then remove the constraints by using Theorem 4.

We first decompose the unitary operator $e^{-iH_{1:q}t}$ into a sequence of the small time unitary operators. For the purpose, for a fixed t , we define the sequence of times $\{t_0, t_1, t_2, \dots, t_{m_0}\}$ as

$$\begin{aligned} t_0 &:= 0, \quad t_m := (m-1)\Delta t_q + \Delta t'_q \quad (m \geq 1), \\ m_0 &= \lceil t/\Delta t_q \rceil, \quad \Delta t'_q = t - (m_0-1)\Delta t_q, \end{aligned} \quad (\text{S.186})$$

where the definition implies $\Delta t'_q \leq \Delta t_q$. We aim to derive the Lieb-Robinson bound for the unitary time evolution $e^{-iH_{1:q}t_{m_0}}$ which is decomposed as

$$\begin{aligned} e^{-iH_{1:q}t_{m_0}} &= e^{-iH_{1:q}\Delta t'_q} [e^{-iH_{1:q}\Delta t_q}]^{m_0} \\ &= e^{-iH_{1:q-1}\Delta t'_q} \mathcal{T} e^{-\int_0^{\Delta t'_q} H_q(H_{1:q-1}, \tau) d\tau} \left[e^{-iH_{1:q-1}\Delta t_q} \mathcal{T} e^{-\int_0^{\Delta t_q} H_q(H_{1:q-1}, \tau) d\tau} \right]^{m_0-1}. \end{aligned} \quad (\text{S.187})$$

We note that $e^{-iH_{1:q-1}\Delta t_q}$ and $e^{-iH_{1:q-1}\Delta t'_q}$ satisfies the $\mathcal{G}_{q-1}(x, t, \mathcal{X})$ -Lieb-Robinson bound, while $\mathcal{T} e^{-\int_0^{\Delta t_q} H_q(H_{1:q-1}, \tau) d\tau}$ and $\mathcal{T} e^{-\int_0^{\Delta t'_q} H_q(H_{1:q-1}, \tau) d\tau}$ satisfies the $\tilde{\mathcal{G}}_q(x, t, \mathcal{X})$ -Lieb-Robinson bound (S.183). Our task is to connect all the Lieb-Robinson bounds appropriately. We here utilize the following proposition (see Sec. XIID for the proof):

Proposition 8. *We consider a Hamiltonian H_0 which satisfies the Lieb-Robinson bound for a fixed t as*

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) \leq \mathcal{C} e^{-(x-v|t|)/\xi} \quad (\text{S.188})$$

for $\text{diam}(\mathcal{X}), \text{diam}(\mathcal{Y}) \leq \nu\xi$ ($\nu \geq 3$). Also, let H'_0 be a Hamiltonian which satisfies the Lieb-Robinson bound for a fixed t' as

$$\mathcal{G}'(x, t', \mathcal{X}) = |\mathcal{X}^{(\nu\xi)}|^2 \mathcal{F}'(x, t'), \quad (\text{S.189})$$

where the function $\mathcal{F}'(x, t')$ is assumed to satisfy

$$\mathcal{F}'(x, t') \leq \mathcal{F}'(x_0, t') e^{-2(x-x_0)/\xi} \quad \text{for } x \geq 0, \quad 0 \leq x_0 \leq x. \quad (\text{S.190})$$

Note that we impose no assumptions on $\text{diam}(\mathcal{X})$ for $\mathcal{G}'(x, t', \mathcal{X})$. We then obtain the Lieb-Robinson function $\mathcal{G}''(x, t, t', \mathcal{X}, \mathcal{Y})$ for $e^{-iH_0 t} e^{-iH'_0 t'}$ as

$$\mathcal{G}''(x, t, t', \mathcal{X}, \mathcal{Y}) = 4\mathcal{C}\gamma(3/\nu)^D (2\kappa^* + \nu)^D e^{-(x-\kappa^*\xi-v|t|)/\xi} \quad (\text{S.191})$$

for $\text{diam}(\mathcal{X}), \text{diam}(\mathcal{Y}) \leq \nu\xi$, where κ^* is defined as an integer ($\kappa^* \in \mathbb{N}$) which satisfies

$$\mathcal{F}'(\kappa^*\xi, t') \leq \frac{1}{2^{D+1}e^{2\gamma^2 D}} =: \zeta_1^{-1}. \quad (\text{S.192})$$

Note that t' -dependence of $\mathcal{G}''(x, t, t', \mathcal{X}, \mathcal{Y})$ is included in κ^* .

By using Proposition 8, we prove the form of

$$\mathcal{G}_q(x, t_m, \mathcal{X}, \mathcal{Y}) = \mathcal{R}_q^m e^{-(x-\bar{v}_{q-1}t_m)/\xi_q} \quad (m = 0, 1, 2, \dots, m_0) \quad (\text{S.193})$$

under the constraint of $\text{diam}(\mathcal{X}) \leq \ell_q$ and $\text{diam}(\mathcal{Y}) \leq \ell_q$ [see Sec. XIIE for the proof of Ineq. (S.193)], where

$$\xi_q := \frac{16\ell_q}{\eta \log(\ell_{q-1})} \quad (\text{S.194})$$

and

$$\mathcal{R}_q := 4e\gamma 9^D e^{\ell_{q-1} \log(\zeta_1)/\xi_q} \cdot 4e\gamma 5^D = 16e^2\gamma^2 \cdot 45^D \cdot e^{\ell_{q-1} \log(\zeta_1)/\xi_q}. \quad (\text{S.195})$$

For the parameters ξ_q and \mathcal{R}_q , we can derive

$$\xi_q \leq \frac{\ell_q}{3}, \quad \mathcal{R}_q \leq 16e^3\gamma^2 \cdot 45^D := \zeta_2, \quad (\text{S.196})$$

where in the first inequality we use the condition (S.140), and in the second inequality we use $e^{\ell_{q-1} \log(\zeta_1)/\xi_q} \leq e$ which is a direct consequence from the condition (S.138).

From the inequality (S.193), we can reduce $\mathcal{G}_q(x, t, \mathcal{X}, \mathcal{Y})$ ($= \mathcal{G}_q(x, t_{m_0}, \mathcal{X}, \mathcal{Y})$) to

$$\begin{aligned} \mathcal{G}_q(x, t, \mathcal{X}, \mathcal{Y}) &= \mathcal{R}_q \exp \left[- \left(\frac{x - \bar{v}_{q-1} t_{m_0} - \xi_q (m_0 - 1) \Delta t_q \Delta t_q^{-1} \log(\mathcal{R}_q)}{\xi_q} \right) \right] \\ &\leq \zeta_2 \exp \left[- \left(\frac{x - t[\bar{v}_{q-1} + \xi_q \Delta t_q^{-1} \log(\zeta_2)]}{\ell_q/2} \right) \right] = \zeta_2 e^{-2(x-v_q t)/\ell_q}, \end{aligned} \quad (\text{S.197})$$

where we use $(m_0 - 1) \Delta t_q \leq t_{m_0} = t$ and $\xi_q \leq \ell_q/3 \leq \ell_q/2$ from (S.196) for the inequality and define v_q as

$$v_q = \bar{v}_{q-1} + \frac{\xi_q \log(\zeta_2)}{\Delta t_q} = \bar{v}_{q-1} + v_{q-1} \frac{192 \log(\zeta_2)}{\eta \log(\ell_{q-1})}, \quad (\text{S.198})$$

where we have defined Δt_q and ξ_q in Eqs. (S.157) and (S.194), respectively. By remembering that $\bar{v}_{q-1} = (1 - \delta_{q-1})^{-1} v_{q-1}$ from Eq. (S.159), we obtain the recurrence equation (S.153) for v_q . In this way, we obtain the $\mathcal{G}_q(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound under the constraints of $\text{diam}(\mathcal{X})$ and $\text{diam}(\mathcal{Y}) \leq \ell_q$.

Finally, in order to remove the constraints of $\text{diam}(\mathcal{X}), \text{diam}(\mathcal{Y}) \leq \ell_q$, we apply Theorem 4 to $\mathcal{G}_q(x, t, \mathcal{X}, \mathcal{Y})$ in (S.197) by setting

$$\xi_0 = \ell_q/2, \quad \xi = (\ell_q - 1)/2 \geq \ell_q/3, \quad (\text{S.199})$$

where we use $\ell_q \geq \ell_1 \geq 3$ from the condition (S.142). From the inequality (S.67), we obtain the Lieb-Robinson bound for generic operators without the constraints as follows:

$$\mathcal{G}_q(x, t, \mathcal{X}) \leq \mathcal{C}_{\ell_q/3, \ell_q/2} |\mathcal{X}^{(\ell_q/3)}|^2 (1 + 3x/\ell_q)^{D-1} \zeta_2 e^{-2(x-v_q t)/\ell_q} \leq \tilde{\mathcal{C}}_0 |\mathcal{X}|^2 (1 + x/\ell_q)^{D-1} e^{-2(x-v_q t)/\ell_q}, \quad (\text{S.200})$$

where $\mathcal{C}_{\ell_q/3, \ell_q/2}$ is defined by Eq. (S.68), and $\tilde{\mathcal{C}}_0$ was defined as

$$\tilde{\mathcal{C}}_0 := \frac{4}{3} e^{10/3} 15^D D! \gamma \zeta_2. \quad (\text{S.201})$$

We thus prove the inequality (S.152). This completes the proof of Theorem 2. \square

E. Proof of the inequality (S.193)

We prove the inequality (S.193) by induction. For $m = 0$, we have $\mathcal{G}_q(x, t_0, \mathcal{X}, \mathcal{Y}) = 0$ because of $t_0 = 0$ and the inequality is trivially obtained.

Then, we assume the form of (S.193) for a fixed m and prove the case of $m+1$. We here consider the Lieb-Robinson bound for

$$e^{-iH_{1:q} t_{m+1}} = e^{-iH_{1:q} t_{m+1}} e^{-iH_{1:q} \Delta t_q} = e^{-iH_{1:q} t_m} e^{-iH_{1:q-1} \Delta t_q} \mathcal{T} e^{-\int_0^{\Delta t_q} H_q(H_{1:q-1}, \tau) d\tau}, \quad (\text{S.202})$$

where we use Eq. (S.166) for the decomposition of $e^{-iH_{1:q} \Delta t_q}$. In order to obtain the Lieb-Robinson bound for $e^{-iH_{1:q} t_{m+1}}$, we first connect the two Lieb-Robinson bounds of $e^{-iH_{1:q} t_m}$ and $e^{-iH_{1:q-1} \Delta t_q}$. We then connect the two Lieb-Robinson bounds of $e^{-iH_{1:q} t_m} e^{-iH_{1:q-1} \Delta t_q}$ and $\mathcal{T} e^{-\int_0^{\Delta t_q} H_q(H_{1:q-1}, \tau) d\tau}$. In each of the connections, we utilize Proposition 8.

For the first, we consider the Lieb-Robinson bound for $e^{-iH_{1:q} t_m} e^{-iH_{1:q-1} \Delta t_q}$, which we characterize by the function of

$$\mathcal{G}'_q(x, t_m, \Delta t_q, \mathcal{X}, \mathcal{Y}). \quad (\text{S.203})$$

The Lieb-Robinson bound for $e^{-iH_{1:q} t_m}$ is given by the assumption (S.193), namely

$$\mathcal{G}_q(x, t_m, \mathcal{X}, \mathcal{Y}) = \mathcal{R}_q^m e^{-(x-\bar{v}_{q-1} t_m)/\xi_q}, \quad (\text{S.204})$$

where we have defined ξ_q as in Eq. (S.194). Also, the Lieb-Robinson bound for $e^{-iH_{1:q-1} \Delta t_q}$ is given by $\mathcal{G}_{q-1}(x, t, \mathcal{X})$ in Eq. (S.158), which has the form of

$$\mathcal{G}_{q-1}(x, t, \mathcal{X}) = |\mathcal{X}^{(\ell_q)}|^2 \mathcal{F}_{q-1}(x, t), \quad \mathcal{F}_{q-1}(x, t) := e^{-(x-\bar{v}_{q-1} t)/\ell_{q-1}}. \quad (\text{S.205})$$

We notice that $\mathcal{F}_{q-1}(x, t)$ satisfies the condition (S.190) in Proposition 8 as follows:

$$\begin{aligned} \mathcal{F}_{q-1}(x, t) &= e^{-(x-\bar{v}_{q-1} t)/\ell_{q-1}} \leq e^{-(x_0-\bar{v}_{q-1} t)/\ell_{q-1}} e^{-(x-x_0)/\ell_{q-1}} \\ &\leq \mathcal{F}_{q-1}(x_0, t) e^{-2(x-x_0)/\xi_q} \end{aligned} \quad (\text{S.206})$$

for $0 \leq x_0 \leq x$, where $1/\ell_{q-1} \geq 2/\xi_q$ is a consequence from the condition (S.139), namely $1/(18\ell_{q-1}) \geq 2/\xi_q$.

In Proposition 8, we set

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{G}_q(x, t_m, \mathcal{X}, \mathcal{Y}), \quad \mathcal{G}'(x, t', \mathcal{X}) = \mathcal{G}_{q-1}(x, \Delta t_q, \mathcal{X}), \quad \mathcal{F}'(x, t') = \mathcal{F}_{q-1}(x, \Delta t_q), \quad (\text{S.207})$$

and the parameters therein are given by

$$t = t_m, \quad t' = \Delta t_q, \quad \mathcal{C} = \mathcal{R}_q^m, \quad v = \bar{v}_{q-1}, \quad \xi = \xi_q, \quad \nu = \nu_q := \frac{\ell_q}{\xi_q} \geq 3, \quad (\text{S.208})$$

where the inequality $\nu_q \geq 3$ is given by the condition (S.140). Then, from Eq. (S.191), the Lieb-Robinson function $\mathcal{G}'_q(x, t_m, \Delta t_q, \mathcal{X}, \mathcal{Y})$ is given by

$$\mathcal{G}'_q(x, t_m, \Delta t_q, \mathcal{X}, \mathcal{Y}) = 4\mathcal{R}_q^m \gamma(3/\nu_q)^D (2\kappa_{q,1}^* + \nu_q)^D e^{-(x - \kappa_{q,1}^* \xi_q - \bar{v}_{q-1} |t_m|)/\xi_q}, \quad (\text{S.209})$$

where $\kappa_{q,1}^*$ is chosen so that the following inequality holds

$$\mathcal{F}_{q-1}(\kappa_{q,1}^* \xi_q, \Delta t_q) = e^{-(\kappa_{q,1}^* \xi_q - \bar{v}_{q-1} \Delta t_q)/\ell_{q-1}} \leq \zeta_1^{-1}, \quad (\text{S.210})$$

where we have defined ζ_1 in (S.192). The parameter $\kappa_{q,1}^*$ is given by

$$\kappa_{q,1}^* = \left\lceil \frac{\bar{v}_{q-1} \Delta t_q + \ell_{q-1} \log(\zeta_1)}{\xi_q} \right\rceil \leq \frac{\bar{v}_{q-1} \Delta t_q + \ell_{q-1} \log(\zeta_1)}{\xi_q} + 1. \quad (\text{S.211})$$

We also obtain

$$(3/\nu_q)(2\kappa_{q,1}^* + \nu_q) \leq 3 \left(1 + \frac{2}{\nu_q} + \frac{2\bar{v}_{q-1} \ell_q / (12v_{q-1}) + 2\ell_{q-1} \log(\zeta_1)}{\ell_q} \right) \leq 9 \quad (\text{S.212})$$

with $\Delta t_q = \ell_q / (12v_{q-1})$, where we use $\xi_q \nu_q = \ell_q$, the condition (S.141), $\bar{v}_{q-1} \leq 2v_{q-1}$ [see Eq. (S.159)] and $\nu_q \geq 3$. The inequalities (S.211) and (S.212) reduce the inequality (S.209) to

$$\begin{aligned} \mathcal{G}'_q(x, t_m, \Delta t_q, \mathcal{X}, \mathcal{Y}) &\leq \mathcal{R}_q^m \cdot 4\gamma e^{-1/\xi_q} 9^D e^{-(x - \bar{v}_{q-1} \Delta t_q - \ell_{q-1} \log(\zeta_1) - \bar{v}_{q-1} |t_m|)/\xi_q} \\ &=: \mathcal{R}_q^m \mathcal{R}_{q,1} e^{-(x - \bar{v}_{q-1} |t_m+1|)/\xi_q}, \end{aligned} \quad (\text{S.213})$$

where we define

$$\mathcal{R}_{q,1} := 4\gamma e^{-1/\xi_q} 9^D e^{\ell_{q-1} \log(\zeta_1)/\xi_q} \leq 4e\gamma 9^D e^{\ell_{q-1} \log(\zeta_1)/\xi_q}. \quad (\text{S.214})$$

In the next step, we connect the Lieb-Robinson bounds for the unitary operators

$$e^{-iH_{1:q} t_m} e^{-iH_{1:q-1} \Delta t_q} \quad \text{and} \quad \mathcal{T} e^{-\int_0^{\Delta t_q} H_q(H_{1:q-1}, \tau) d\tau}, \quad (\text{S.215})$$

which gives the Lieb-Robinson bound for $e^{-iH_{1:q} t_{m+1}}$, namely $\mathcal{G}_q(x, t_{m+1}, \mathcal{X}, \mathcal{Y})$. In order to apply Proposition 8 to this case, we first simplify the Lieb-Robinson bound (S.183) for $\mathcal{T} e^{-\int_0^{\Delta t_q} H_q(H_{1:q-1}, \tau) d\tau}$ as follows:

$$\begin{aligned} \tilde{\mathcal{G}}_q(x, \Delta t_q, \mathcal{X}) &\leq e|\mathcal{X}^{(\ell_q)}| \left[\ell_{q-1}^{-\eta} \exp\left(-\frac{1}{18\ell_{q-1}} x\right) + \ell_{q-1}^{-\eta/2} \cdot \exp\left(\frac{-\eta \log(\ell_{q-1})}{8\ell_q} x\right) \right] \\ &\leq 2e|\mathcal{X}^{(\ell_q)}|^2 \ell_{q-1}^{-\eta/2} \cdot e^{-2x/\xi_q} = |\mathcal{X}^{(\ell_q)}|^2 \tilde{\mathcal{F}}_q(x, \Delta t_q) \end{aligned} \quad (\text{S.216})$$

with

$$\tilde{\mathcal{F}}_q(x, \Delta t_q) = 2e\ell_{q-1}^{-\eta/2} \cdot e^{-2x/\xi_q}, \quad (\text{S.217})$$

where we use the condition (S.139) and the trivial inequality of $|\mathcal{X}^{(\ell_q)}| \leq |\mathcal{X}^{(\ell_q)}|^2$. We notice that $\tilde{\mathcal{F}}_q(x, \Delta t_q)$ satisfies the condition (S.190) in Proposition 8.

In order to connect two unitary operators (S.215), we set in Proposition 8

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{G}'_q(x, t_m, \Delta t_q, \mathcal{X}, \mathcal{Y}), \quad \mathcal{G}'(x, t', \mathcal{X}) = \tilde{\mathcal{G}}_q(x, \Delta t_q, \mathcal{X}), \quad \mathcal{F}'(x, t') = \tilde{\mathcal{F}}_q(x, \Delta t_q), \quad (\text{S.218})$$

and the parameters therein are given by

$$t = t_m + \Delta t_q = t_{m+1}, \quad t' = \Delta t_q, \quad \mathcal{C} = \mathcal{R}_q^m \mathcal{R}_{q,1}, \quad v = \bar{v}_{q-1}, \quad \xi = \xi_q, \quad \nu = \nu_q := \frac{\ell_q}{\xi_q} \geq 3. \quad (\text{S.219})$$

We then obtain from Eq. (S.191)

$$\mathcal{G}_q(x, t_{m+1}, \mathcal{X}, \mathcal{Y}) = 4\mathcal{R}_q^m \mathcal{R}_{q,1} \gamma(3/\nu_q)^D (2\kappa_{q,2}^* + \nu_q)^D e^{-(x - \kappa_{q,2}^* \xi_q - \bar{v}_{q-1}|t_{m+1}|)/\xi_q}, \quad (\text{S.220})$$

where $\kappa_{q,2}^*$ is chosen such that it satisfies

$$\tilde{\mathcal{F}}_q(\kappa_{q,2}^* \xi_q, \Delta t_q) = 2e\ell_{q-1}^{-\eta/2} \cdot e^{-2\kappa_{q,2}^*} \leq \zeta_1^{-1}. \quad (\text{S.221})$$

From the condition (S.143), the above condition is satisfied by choosing $\kappa_{q,2}^* = 1$. Hence, we have

$$\mathcal{G}_q(x, t_{m+1}, \mathcal{X}, \mathcal{Y}) = 4\mathcal{R}_q^m \mathcal{R}_{q,1} \gamma(6/\nu_q + 3)^D e^{-(x - \xi_q - \bar{v}_{q-1}|t_{m+1}|)/\xi_q} = \mathcal{R}_q^m \mathcal{R}_{q,1} \mathcal{R}_{q,2} e^{-(x - \bar{v}_{q-1}|t_{m+1}|)/\xi_q} \quad (\text{S.222})$$

with

$$\mathcal{R}_{q,2} = 4e\gamma(6/\nu_q + 3)^D \leq 4e\gamma 5^D, \quad (\text{S.223})$$

where we use $\nu_q \geq 3$. Because of $\mathcal{R}_{q,1} \mathcal{R}_{q,2} \leq \mathcal{R}_q$ from the definition (S.195), we prove the inequality (S.193) for the case of $m + 1$. This completes the proof. \square

XIII. DETAILED PROOF OF THE COMPLEMENTARY STATEMENTS FOR THE MIDDLE-RANGE LIEB-ROBINSON BOUND

A. Proof of Proposition 6: Quasi-locality of operator after time-evolution

1. Statement

Proposition 6. Let $\tilde{\ell}_q$ be a length scale such that

$$2\gamma^2 e^{\bar{v}_{q-1}\tau/\ell_{q-1}} e^{-\tilde{\ell}_q/(2\ell_{q-1})} \leq 1 \quad (\text{S.224})$$

for $\tau \leq \Delta t_q$. Then, under the assumption of (S.158), we obtain a decomposition the time-evolved Hamiltonian $H_q(H_{1:q-1}, \tau)$ in the form of

$$H_q(H_{1:q-1}, \tau) = \sum_{Z: \text{diam}(Z) \leq \ell_q} \sum_{s=1}^{\infty} h_{\tau, Z[s\tilde{\ell}_q]} \quad (\text{S.225})$$

with

$$\sum_{\substack{Z: \text{diam}(Z) \leq \ell_q \\ Z[s\tilde{\ell}_q] \cap X \neq \emptyset}} \|h_{\tau, Z[s\tilde{\ell}_q]}\| \leq (2\tilde{\ell}_q)^D g_q \gamma |X^{(s_0 \tilde{\ell}_q)}| (s + s_0)^D e^{-(s-1)\tilde{\ell}_q/(2\ell_{q-1})} \quad (\text{S.226})$$

for an arbitrary subset $X \subseteq \Lambda$, where s_0 can be arbitrarily chosen and $h_{\tau, Z[s\tilde{\ell}_q]}$ is an operator which acts on the subset $Z[s\tilde{\ell}_q]$.

2. Proof of Proposition 6

For the proof, we consider a more general setup. We define two operators A_1 and A_2 . We assume that the operator A_1 satisfies $\mathcal{G}_{A_1}(x, t, \mathcal{X})$ -Lieb-Robinson bound with

$$\mathcal{G}_{A_1}(x, t, \mathcal{X}) = \mathcal{C} |\mathcal{X}^{(\xi_2)}|^2 e^{-(x - v|t|)/\xi_1}. \quad (\text{S.227})$$

Also, the operator A_2 consists of terms with interaction length at most of ξ_2 , namely

$$A_2 = \sum_{Z: \text{diam}(Z) \leq \xi_2} a_Z, \quad \sum_{Z: Z \ni i} \|a_Z\| \leq g_2. \quad (\text{S.228})$$

We here consider

$$A_2(A_1, t) = e^{iA_1 t} A_2 e^{-iA_1 t} = \sum_{Z: \text{diam}(Z) \leq \xi_2} a_Z(A_1, t), \quad (\text{S.229})$$

and denote $A_2(A_1, t)$ by

$$A_2(A_1, t) = \sum_{Z: \text{diam}(Z) \leq \xi_2} \sum_{s=1}^{\infty} a_{t, Z[s\tilde{\xi}_2]}, \quad (\text{S.230})$$

where $\tilde{\xi}_2$ is an arbitrary positive integer such that $\tilde{\xi}_2 \geq \xi_2$. We here define

$$\begin{aligned} a_{t, Z[\tilde{\xi}_2]} &:= a_Z(A_1, t, Z[\tilde{\xi}_2]), \\ a_{t, Z[s\tilde{\xi}_2]} &:= a_Z(A_1, t, Z[s\tilde{\xi}_2]) - a_Z(A_1, t, Z[\tilde{\xi}_2(s-1)]) \quad \text{for } s \geq 2, \end{aligned} \quad (\text{S.231})$$

where $a_Z(A_1, t, Z[s\tilde{\xi}_2])$ is defined by using the notation of (S.25). Notice that $\lim_{s \rightarrow \infty} a_Z(A_1, t, Z[s\tilde{\xi}_2]) = a_Z(A_1, t)$.

We then derive the following lemma for the time-evolved operator $A_2(A_1, t)$ which satisfies the quasi-locality:

Lemma 5. *Let s_0 be an arbitrary positive number. Then, we have*

$$\sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z[s\tilde{\xi}_2] \cap X \neq \emptyset}} \|a_{t, Z[s\tilde{\xi}_2]}\| \leq 2(2\tilde{\xi}_2)^D \mathcal{C} \gamma^3 g_2 e^{v|t|/\xi_1} |X^{(s_0\tilde{\xi}_2)}| (s + s_0)^D e^{-(s-1)\tilde{\xi}_2/\xi_1} \quad (\text{S.232})$$

for $s \geq 2$, where $\text{diam}(Z[s\tilde{\xi}_2]) = \text{diam}(Z) + 2s\tilde{\xi}_2 \leq \xi_2 + 2s\tilde{\xi}_2$ from $\text{diam}(Z) \leq \xi_2$. In particular, for $s = 1$, we have,

$$\sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z[\tilde{\xi}_2] \cap X \neq \emptyset}} \|a_{t, Z[\tilde{\xi}_2]}\| \leq (2\tilde{\xi}_2)^D g_2 \gamma |X^{(s_0\tilde{\xi}_2)}| (s_0 + 1)^D. \quad (\text{S.233})$$

In this lemma, we further simplify the upper bound (S.232) as follows. By choosing $\tilde{\xi}_2$ as

$$2\gamma^2 \mathcal{C} e^{v|t|/\xi_1} e^{-\tilde{\xi}_2/(2\xi_1)} \leq 1, \quad (\text{S.234})$$

we obtain for $s \geq 2$

$$\begin{aligned} 2(2\tilde{\xi}_2)^D \mathcal{C} \gamma^3 g_2 e^{v|t|/\xi_1} (s + s_0)^D e^{-(s-1)\tilde{\xi}_2/\xi_1} &= 2\gamma^2 \mathcal{C} e^{v|t|/\xi_1} e^{-(s-1)\tilde{\xi}_2/(2\xi_1)} \cdot (2\tilde{\xi}_2)^D g_2 \gamma (s + s_0)^D e^{-(s-1)\tilde{\xi}_2/(2\xi_1)} \\ &\leq 2\gamma^2 \mathcal{C} e^{v|t|/\xi_1} e^{-\tilde{\xi}_2/(2\xi_1)} \cdot (2\tilde{\xi}_2)^D g_2 \gamma (s + s_0)^D e^{-(s-1)\tilde{\xi}_2/(2\xi_1)} \\ &\leq (2\tilde{\xi}_2)^D g_2 \gamma (s + s_0)^D e^{-(s-1)\tilde{\xi}_2/(2\xi_1)}, \end{aligned} \quad (\text{S.235})$$

which reduces the inequality (S.232) to

$$\sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z[s\tilde{\xi}_2] \cap X \neq \emptyset}} \|a_{t, Z[s\tilde{\xi}_2]}\| \leq (2\tilde{\xi}_2)^D g_2 \gamma |X^{(s_0\tilde{\xi}_2)}| (s + s_0)^D e^{-(s-1)\tilde{\xi}_2/(2\xi_1)}. \quad (\text{S.236})$$

We note that the inequality (S.233) also reduces to (S.236) for $s = 1$.

Proposition 6 is immediately given by choosing $|t| = \tau$, $A_1 = H_{1:q-1}$, $A_2 = H_q$. Note the Lieb-Robinson bound $\mathcal{G}_{A_1}(x, t, \mathcal{X})$ in (S.227) is replaced by $\mathcal{G}_{q-1}(x, t, \mathcal{X})$ which has been given in (S.158), namely $\mathcal{G}_{q-1}(x, t, \mathcal{X}) = |\mathcal{X}^{(\ell_q)}|^2 e^{-(x - \bar{v}_{q-1}\Delta t_q)/\ell_{q-1}}$. Now, the parameters $\{g_2, \xi_1, \xi_2, \tilde{\xi}_2, \mathcal{C}, v\}$ in Lemma 5 are given by

$$g_2 = g_q, \quad \xi_1 = \ell_{q-1}, \quad \xi_2 = \ell_q, \quad \tilde{\xi}_2 = \tilde{\ell}_q, \quad \mathcal{C} = 1, \quad v = \bar{v}_{q-1}. \quad (\text{S.237})$$

Then, the condition (S.234) and the inequality (S.236) reduces to the condition (S.224) and the inequality (S.226), respectively. This completes the proof. \square

3. Proof of Lemma 5

In the proof, we aim to obtain the upper bound of

$$\sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z[s\tilde{\xi}_2] \cap X \neq \emptyset}} \|a_{t, Z[s\tilde{\xi}_2]}\|. \quad (\text{S.238})$$

First, from the Lieb-Robinson bound in the form of of Eq. (S.227) and Lemma 1, we obtain

$$\|a_Z(A_1, t) - a_Z(A_1, t, Z[s\tilde{\xi}_2])\| \leq \mathcal{G}_{A_1}(s\tilde{\xi}_2, t, Z) = \mathcal{C} |\mathcal{X}^{(\xi_2)}|^2 e^{-(s\tilde{\xi}_2 - v|t|)/\xi_1}, \quad (\text{S.239})$$

which upper-bounds the norm of $a_{t,Z[s\tilde{\xi}_2]}$ in Eq. (S.231) as follows:

$$\begin{aligned} \|a_{t,Z[s\tilde{\xi}_2]}\| &\leq \|a_Z(A_1, t) - a_Z(A_1, t, Z[s\tilde{\xi}_2])\| + \|a_Z(A_1, t) - a_Z(A_1, t, Z[s\tilde{\xi}_2 - \tilde{\xi}_2])\| \\ &\leq \|a_Z\| \cdot \mathcal{C} e^{v|t|/\xi_1} |Z^{(\xi_2)}|^2 e^{-s\tilde{\xi}_2/\xi_1} + \|a_Z\| \cdot \mathcal{C} e^{v|t|/\xi_1} |Z^{(\xi_2)}|^2 e^{-(s-1)\tilde{\xi}_2/\xi_1} \\ &\leq \|a_Z\| \cdot 2\gamma^2 \mathcal{C} e^{v|t|/\xi_1} e^{-(s-1)\tilde{\xi}_2/\xi_1}, \end{aligned} \quad (\text{S.240})$$

where we use the inequality (S.18) with the condition $\text{diam}(Z) \leq \xi_2$ to obtain $|Z^{(\xi_2)}| \leq \gamma[\text{diam}(Z)/\xi_2]^D = \gamma$. For $s = 1$, from the definitions (S.25) and (S.231), we have

$$\|a_{t,Z[\tilde{\xi}_2]}\| \leq \|a_Z\|. \quad (\text{S.241})$$

By applying the inequality (S.240) to the summation (S.238), we have

$$\sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z[s\tilde{\xi}_2] \cap X \neq \emptyset}} \|a_{t,Z[s\tilde{\xi}_2]}\| \leq 2\gamma^2 \mathcal{C} e^{v|t|/\xi_1} \sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z[s\tilde{\xi}_2] \cap X \neq \emptyset}} \|a_Z\| e^{-(s-1)\tilde{\xi}_2/\xi_1} \quad \text{for } s \geq 2. \quad (\text{S.242})$$

On the summation with respect to Z , we obtain

$$\begin{aligned} \sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z[s\tilde{\xi}_2] \cap X \neq \emptyset}} \|a_Z\| &= \sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z \cap X[s\tilde{\xi}_2] \neq \emptyset}} \|a_Z\| \leq \sum_{j \in X[s\tilde{\xi}_2]} \sum_{Z: Z \ni j} \|a_Z\| \leq \sum_{j \in X[s\tilde{\xi}_2]} g_2 \leq g_2 |X[s\tilde{\xi}_2]| \\ &\leq g_2 |X^{(s_0\tilde{\xi}_2)}[(s+s_0)\tilde{\xi}_2]| \leq g_2 \gamma |X^{(s_0\tilde{\xi}_2)}| \cdot [2(s+s_0)\tilde{\xi}_2]^D \end{aligned} \quad (\text{S.243})$$

for arbitrary $s_0 \geq 0$, where we use the inequalities (S.228), (S.15) and (S.20) in the second, fourth and fifth inequalities, respectively. Note that the condition $Z[s\tilde{\xi}_2] \cap X \neq \emptyset$ for $Z \subseteq \Lambda$ is equivalent to $Z \cap X[s\tilde{\xi}_2] \neq \emptyset$.

By combining the inequalities (S.242) and (S.243), we have

$$\sum_{Z: Z[s\tilde{\xi}_2] \cap X \neq \emptyset} \|a_{t,Z[s\tilde{\xi}_2]}\| \leq 2(2\tilde{\xi}_2)^D \mathcal{C} \gamma^3 g_2 e^{v|t|/\xi_1} |X^{(s_0\tilde{\xi}_2)}| (s+s_0)^D e^{-(s-1)\tilde{\xi}_2/\xi_1}. \quad (\text{S.244})$$

We thus prove the inequality (S.232). In the same way, we can prove the case of $s = 1$. By using the inequality (S.241), we have

$$\sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z[\tilde{\xi}_2] \cap X \neq \emptyset}} \|a_{t,Z[\tilde{\xi}_2]}\| \leq \sum_{\substack{Z: \text{diam}(Z) \leq \xi_2 \\ Z[\tilde{\xi}_2] \cap X \neq \emptyset}} \|a_Z\| \leq (2\tilde{\xi}_2)^D g_2 \gamma |X^{(s_0\tilde{\xi}_2)}| (s_0+1)^D, \quad (\text{S.245})$$

where in the last inequality we use (S.243) with $s = 1$. This completes the proof of Lemma 6. \square

B. Proof of Proposition 7: Lieb-Robinson bound by time-evolved Hamiltonian

1. Statement

Proposition 7. *Let \tilde{H} be a Hamiltonian such that*

$$\tilde{H} = \sum_{Z: Z \subseteq \Lambda} h_Z = \sum_{s=1}^{\infty} \sum_{\text{diam}(Z_s) \leq s\xi} h_{Z_s}. \quad (\text{S.246})$$

For an arbitrary subset X and an arbitrary positive s_0 , we assume

$$\sum_{\substack{\text{diam}(Z_s) \leq s\xi \\ Z_s \cap X \neq \emptyset}} \|h_{Z_s}\| \leq \tilde{g} |X^{(s_0\xi)}| (s+s_0)^D e^{-\mu(s-1)} \quad (\mu > 1). \quad (\text{S.247})$$

Then, the Hamiltonian \tilde{H} satisfies $\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X})$ -Lieb-Robinson bound with

$$\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X}) = e |\mathcal{X}^{(\xi)}| \left[\tilde{v} |t| \exp \left(-\frac{\mu-1}{2} \frac{x}{\xi} + \tilde{v} |t| \right) + \left(\frac{e^2 \tilde{v} |t|}{m_x^*} \right)^{m_x^*} \right] \quad (\text{S.248})$$

with

$$\tilde{v} := 4\tilde{g}\gamma(4D)^D, \quad m_x^* := \left\lceil \frac{\mu-1}{2\mu} \frac{x}{\xi} \right\rceil + 1. \quad (\text{S.249})$$

We note that the same inequality holds for time-dependent Hamiltonians.

2. Proof of Proposition 7

We estimate the norm of $\|[O_L(\tilde{H}, t), O_{L'}]\|$, where O_L and $O_{L'}$ are arbitrary operators with the unit norm (i.e., $\|O_L\| = \|O_{L'}\| = 1$) which are supported on L and L' , respectively. Following Ref. [15], we start from the inequality as

$$\frac{d}{dt}\|[O_L(\tilde{H}, t), O_{L'}]\| = \frac{d}{dt}\|[e^{i\tilde{H}t}O_L e^{-i\tilde{H}t}, O_{L'}]\| \leq 2\|O_L\| \cdot \|\tilde{H}_L(\tilde{H}, \tau), O_{L'}\| = 2\|\tilde{H}_L(\tilde{H}, \tau), O_{L'}\|, \quad (\text{S.250})$$

where we define \tilde{H}_L as follows:

$$\tilde{H}_L := \sum_{s=1}^{\infty} \sum_{\substack{\text{diam}(Z_s) \leq s\xi \\ Z_s \cap L \neq \emptyset}} h_{Z_s}. \quad (\text{S.251})$$

The operator \tilde{H}_L picks up all the interaction terms $\{h_Z\}_{Z \subseteq \Lambda}$ in (S.246) that act on the subset L .

By using the inequality (S.250), we first obtain

$$\begin{aligned} \|[O_L(\tilde{H}, t), O_{L'}]\| &\leq \|[O_L, O_{L'}]\| + 2 \int_0^t \|\tilde{H}_L(\tilde{H}, \tau), O_{L'}\| d\tau \\ &\leq 2 \sum_{s_1=1}^{\infty} \sum_{\substack{\text{diam}(Z_{s_1}) \leq s_1\xi \\ Z_{s_1} \cap L \neq \emptyset}} \int_0^t \|h_{Z_{s_1}}(\tilde{H}, \tau_1), O_{L'}\| d\tau_1, \end{aligned} \quad (\text{S.252})$$

where we use $d_{L, L'} \geq 1$ which implies $[O_L, O_{L'}] = 0$. By applying the inequality (S.250) to $\|h_{Z_{s_1}}(\tilde{H}, \tau_1), O_{L'}\|$, we have

$$\|h_{Z_{s_1}}(\tilde{H}, \tau_1), O_{L'}\| \leq \|h_{Z_{s_1}}, O_{L'}\| + 2\|h_{Z_{s_1}}\| \sum_{s_2=1}^{\infty} \sum_{\substack{\text{diam}(Z_{s_2}) \leq s_2\xi \\ Z_{s_2} \cap Z_{s_1} \neq \emptyset}} \int_0^{\tau_1} \|h_{Z_{s_2}}(\tilde{H}, \tau_2), O_{L'}\| d\tau_2. \quad (\text{S.253})$$

In the following, for the simplicity of the notation, we simply describe as

$$\sum_{\substack{\text{diam}(Z_{s_2}) \leq s_2\xi \\ Z_{s_2} \cap Z_{s_1} \neq \emptyset}} = \sum_{Z_{s_2} \cap Z_{s_1} \neq \emptyset} \quad (\text{S.254})$$

by omitting the constraint of $\text{diam}(Z_{s_2}) \leq s_2\xi$. By combining the inequalities (S.252) and (S.253), we have

$$\begin{aligned} \|[O_L(\tilde{H}, t), O_{L'}]\| &\leq 2|t| \sum_{s_1=1}^{\infty} \sum_{\substack{Z_{s_1} \cap L \neq \emptyset \\ Z_{s_1} \cap L' \neq \emptyset}} \|h_{Z_{s_1}}, O_{L'}\| \\ &\quad + 2^2 \sum_{s_1=1}^{\infty} \sum_{Z_{s_1} \cap L \neq \emptyset} \|h_{Z_{s_1}}\| \sum_{s_2=1}^{\infty} \sum_{Z_{s_2} \cap Z_{s_1} \neq \emptyset} \int_0^t \int_0^{\tau_1} \|h_{Z_{s_2}}(\tilde{H}, \tau_2), O_{L'}\| d\tau_2 d\tau_1, \end{aligned} \quad (\text{S.255})$$

where in the summation of the first term we add the restriction of $Z_{s_1} \cap L' \neq \emptyset$ because of $\|h_{Z_{s_1}}, O_{L'}\| = 0$ for $Z_{s_1} \cap L' = \emptyset$.

By repeating this process $(m^* - 1)$ times, we obtain

$$\begin{aligned} &\|[O_L(\tilde{H}, t), O_{L'}]\| \\ &\leq 2|t| \sum_{s_1=1}^{\infty} \sum_{\substack{Z_{s_1} \cap L \neq \emptyset \\ Z_{s_1} \cap L' \neq \emptyset}} \|h_{Z_{s_1}}, O_{L'}\| + \frac{(2|t|)^2}{2!} \sum_{s_1=1}^{\infty} \sum_{\substack{Z_{s_1} \cap L \neq \emptyset \\ Z_{s_1} \cap L' \neq \emptyset}} \|h_{Z_{s_1}}\| \sum_{s_2=1}^{\infty} \sum_{\substack{Z_{s_2} \cap Z_{s_1} \neq \emptyset \\ Z_{s_2} \cap L' \neq \emptyset}} \|h_{Z_{s_2}}, O_{L'}\| \\ &\quad + \frac{(2|t|)^3}{3!} \sum_{s_1=1}^{\infty} \sum_{\substack{Z_{s_1} \cap L \neq \emptyset \\ Z_{s_1} \cap L' \neq \emptyset}} \|h_{Z_{s_1}}\| \sum_{s_2=1}^{\infty} \sum_{\substack{Z_{s_2} \cap Z_{s_1} \neq \emptyset \\ Z_{s_2} \cap L' \neq \emptyset}} \|h_{Z_{s_2}}\| \sum_{s_3=1}^{\infty} \sum_{\substack{Z_{s_3} \cap Z_{s_2} \neq \emptyset \\ Z_{s_3} \cap L' \neq \emptyset}} \|h_{Z_{s_3}}, O_{L'}\| \\ &\quad + \cdots + 2^{m^*} \sum_{s_1=1}^{\infty} \sum_{\substack{Z_{s_1} \cap L \neq \emptyset \\ Z_{s_1} \cap L' \neq \emptyset}} \|h_{Z_{s_1}}\| \sum_{s_2=1}^{\infty} \sum_{\substack{Z_{s_2} \cap Z_{s_1} \neq \emptyset \\ Z_{s_2} \cap L' \neq \emptyset}} \|h_{Z_{s_2}}\| \cdots \sum_{s_m^*=1}^{\infty} \sum_{\substack{Z_{s_m^*} \cap Z_{s_{m^*-1}} \neq \emptyset \\ Z_{s_m^*} \cap L' \neq \emptyset}} \|h_{Z_{s_m^*}}(\tilde{H}, \tau_{m^*}), O_{L'}\| \\ &\quad \int_0^t \int_0^{\tau_1} \cdots \int_0^{\tau_{m^*-1}} d\tau_1 d\tau_2 \cdots d\tau_{m^*} \\ &\leq \sum_{m=1}^{m^*} \mathcal{L}_m. \end{aligned} \quad (\text{S.256})$$

In the above inequality, we define $\{\mathcal{L}_m\}_{m=1}^{m^*}$ as

$$\mathcal{L}_m := \frac{2(2|t|)^m}{m!} \sum_{s_1=1}^{\infty} \sum_{Z_{s_1} \cap L \neq \emptyset} \|h_{Z_{s_1}}\| \sum_{s_2=1}^{\infty} \sum_{Z_{s_2} \cap Z_{s_1} \neq \emptyset} \|h_{Z_{s_2}}\| \cdots \sum_{s_m=1}^{\infty} \sum_{\substack{Z_{s_m} \cap Z_{s_{m-1}} \neq \emptyset \\ Z_{s_m} \cap L' \neq \emptyset}} \|h_{Z_{s_m}}\| \quad (\text{S.257})$$

for $m = 1, 2, \dots, m^* - 1$, where we use $\|h_{Z_{s_m}}, O_{L'}\| \leq 2\|h_{Z_{s_m}}\|$ from $\|O_{L'}\| = 1$. For $m = m^*$, we define \mathcal{L}_{m^*} as

$$\mathcal{L}_{m^*} := \frac{2(2|t|)^{m^*}}{m^*!} \sum_{s_1=1}^{\infty} \sum_{Z_{s_1} \cap L \neq \emptyset} \|h_{Z_{s_1}}\| \sum_{s_2=1}^{\infty} \sum_{Z_{s_2} \cap Z_{s_1} \neq \emptyset} \|h_{Z_{s_2}}\| \cdots \sum_{s_m^*=1}^{\infty} \sum_{Z_{s_m^*} \cap Z_{s_{m^*-1}} \neq \emptyset} \|h_{Z_{s_m^*}}\|. \quad (\text{S.258})$$

We notice that for \mathcal{L}_{m^*} the summation with respect to $Z_{s_m^*}$ is not restricted to $\{Z_{s_m^*} | Z_{s_m^*} \cap L' \neq \emptyset\}$.

We, in the following, aim to upper-bound the m th term \mathcal{L}_m in Eq. (S.257). By using the condition (S.247) with $s_0 = s_{j-1}$, we obtain the upper bound of

$$\sum_{Z_{s_j} \cap Z_{s_{j-1}} \neq \emptyset} \|h_{Z_{s_j}}\| \leq \tilde{g} |Z_{s_{j-1}}^{(s_{j-1}\xi)}| (s_j + s_{j-1})^D e^{-\mu(s_{j-1})} \leq \tilde{g} \gamma (s_j + s_{j-1})^D e^{-\mu(s_{j-1})} \quad (\text{S.259})$$

for $j \geq 2$, where we use the inequality (S.18) with the condition $\text{diam}(Z_{s_{j-1}}) \leq s_{j-1}\xi$ to obtain

$$|Z_{s_{j-1}}^{(s_{j-1}\xi)}| \leq \gamma [\text{diam}(Z_{s_{j-1}})/(s_{j-1}\xi)]^D = \gamma. \quad (\text{S.260})$$

Also, from the condition (S.247) with $s_0 = 1$, we obtain

$$\sum_{Z_{s_1} \cap L \neq \emptyset} \|h_{Z_{s_1}}\| \leq \tilde{g} |L^{(\xi)}| (1 + s_1)^D e^{-\mu(s_1-1)} \leq \tilde{g} \gamma |L^{(\xi)}| (1 + s_1)^D e^{-\mu(s_1-1)}, \quad (\text{S.261})$$

where we use $\gamma \geq 1$. By combining the inequalities (S.259) and (S.261), we have

$$\sum_{Z_{s_1} \cap L \neq \emptyset} \|h_{Z_{s_1}}\| \sum_{Z_{s_2} \cap Z_{s_1} \neq \emptyset} \|h_{Z_{s_2}}\| \cdots \sum_{Z_{s_m} \cap Z_{s_{m-1}} \neq \emptyset} \|h_{Z_{s_m}}\| \leq \prod_{j=1}^m g_{s_{j-1}, s_j}, \quad (\text{S.262})$$

where $g_{s_{j-1}, s_j} := \tilde{g} \gamma (s_j + s_{j-1})^D e^{-\mu(s_{j-1})}$ for $j \geq 2$ and $g_{s_0, s_1} = \tilde{g} \gamma |L^{(\xi)}| (1 + s_1)^D e^{-\mu(s_1-1)}$. For the product of $\{g_{s_{j-1}, s_j}\}_{j=1}^m$, the following inequality holds as

$$\prod_{j=1}^m g_{s_{j-1}, s_j} = (\tilde{g} \gamma)^m e^{-\mu(S_m-m)} |L^{(\xi)}| (1 + s_1)^D \prod_{j=2}^m (s_j + s_{j-1})^D \leq |L^{(\xi)}| (\tilde{g} \gamma)^m \left(\frac{2S_m}{m}\right)^{mD} e^{-\mu(S_m-m)}, \quad (\text{S.263})$$

where $S_m := \sum_{j=1}^m s_j \geq m$ and we use

$$(1 + s_1)^D \prod_{j=2}^m (s_j + s_{j-1})^D \leq \left(\frac{2S_m}{m}\right)^{mD}. \quad (\text{S.264})$$

Under the constraints of $\{Z_{s_1} \cap L \neq \emptyset, Z_{s_2} \cap Z_{s_1} \neq \emptyset, \dots, Z_{s_m} \cap Z_{s_{m-1}} \neq \emptyset\}$, a necessary condition for $Z_{s_m} \cap L' \neq \emptyset$ is given by

$$S_m \xi \geq d_{L, L'}, \quad (\text{S.265})$$

and hence by combining (S.257), (S.262) and (S.263), the m th term \mathcal{L}_m is bounded from above by

$$\mathcal{L}_m \leq \frac{2(2\tilde{g}\gamma|t|)^m}{m!} |L^{(\xi)}| \sum_{\substack{s_1 \geq 1, s_2 \geq 1, \dots, s_m \geq 1 \\ S_m \xi \geq d_{L, L'}}} \left(\frac{2S_m}{m}\right)^{mD} e^{-\mu(S_m-m)}. \quad (\text{S.266})$$

We upper-bound the summation as follows:

$$\begin{aligned} \sum_{\substack{s_1 \geq 1, s_2 \geq 1, \dots, s_m \geq 1 \\ S_m \xi \geq d_{L, L'}}} \left(\frac{2S_m}{m}\right)^{mD} e^{-\mu(S_m-m)} &= \sum_{S \geq \max(m, d_{L, L'}/\xi)} \sum_{\substack{s_1 \geq 1, s_2 \geq 1, \dots, s_m \geq 1 \\ s_1 + s_2 + \dots + s_m = S}} \left(\frac{2S}{m}\right)^{mD} e^{-\mu(S-m)} \\ &= \sum_{S \geq \max(m, d_{L, L'}/\xi)} \binom{m}{S-m} \left(\frac{2S}{m}\right)^{mD} e^{-\mu(S-m)} \\ &= \sum_{S \geq \max(m, d_{L, L'}/\xi)} \binom{S-1}{m-1} \left(\frac{2S}{m}\right)^{mD} e^{-\mu(S-m)} \\ &\leq \frac{2^{mD} e^{\mu m}}{m^{mD} (m-1)!} \sum_{S \geq d_{L, L'}/\xi} S^{mD+m-1} e^{-\mu S}, \end{aligned} \quad (\text{S.267})$$

where the summation with respect to $\{s_1, s_2, \dots, s_m\}$ such that $s_1 + s_2 + \dots + s_m = S$ is equal to the $(S - m)$ -multicombination from a set of m elements, and we use $\binom{S-1}{m-1} \leq S^{m-1}/(m-1)!$ in the inequality. For an arbitrary constant $S_0 \geq 0$ and $z \in \mathbb{N}$, we obtain

$$\begin{aligned} \sum_{S \geq S_0} S^z e^{-\mu S} &= \sum_{S \geq S_0} S^z e^{-(\mu-1)S} e^{-S} \leq e^{-(\mu-1)S_0} \sum_{S \geq S_0} S^z e^{-S} \\ &\leq e^{-(\mu-1)S_0} \int_{S_0}^{\infty} x^z e^{-x+1} dx \leq e^{-(\mu-1)S_0+1} z!, \end{aligned} \quad (\text{S.268})$$

where in the first inequality we use the condition $\mu > 1$. By using the above inequality with $S_0 = d_{L,L'}/\xi$ and $z = mD + m - 1$, the inequality (S.267) reduces to

$$\begin{aligned} \sum_{\substack{s_1 \geq 1, s_2 \geq 1, \dots, s_m \geq 1 \\ S_m \xi \geq d_{L,L'}}} \left(\frac{2S_m}{m} \right)^{mD} e^{-\mu(S_m-m)} &\leq \frac{2^{mD} e^{1+\mu m - (\mu-1)d_{L,L'}/\xi}}{m^{mD}(m-1)!} (mD + m - 1)! \\ &= \frac{2^{mD} e^{1+\mu m - (\mu-1)d_{L,L'}/\xi}}{m^{mD}} (mD)! \binom{mD + m - 1}{m-1} \\ &\leq e(2D)^{mD} e^{\mu m - (\mu-1)d_{L,L'}/\xi} \binom{mD + m - 1}{m-1} \\ &\leq e(4D)^{mD} 2^{m-1} e^{\mu m - (\mu-1)d_{L,L'}/\xi}, \end{aligned} \quad (\text{S.269})$$

where in the second inequality we use $(mD)! \leq (mD)^{mD}$ and in the last inequality we use $\binom{mD+m-1}{m-1} \leq 2^{mD+m-1}$.

Therefore, by combining the inequalities (S.266) and (S.269), we finally arrive at the inequality for \mathcal{L}_m as

$$\mathcal{L}_m \leq \frac{e[4\tilde{g}\gamma(4D)^D |t|]^m}{m!} |L^{(\xi)}| e^{\mu m - (\mu-1)d_{L,L'}/\xi} = \frac{e(\tilde{v}|t|)^m}{m!} |L^{(\xi)}| e^{\mu m - (\mu-1)d_{L,L'}/\xi}, \quad (\text{S.270})$$

where we use the definition of \tilde{v} in Eq. (S.249). In the same way, we can derive the similar upper bound for \mathcal{L}_{m^*} . For \mathcal{L}_{m^*} , only the difference is that the constraint of $S_m \xi \geq d_{L,L'}$ in (S.266) is removed. Hence, the same calculations as (S.267) and (S.269) give

$$\begin{aligned} \sum_{s_1 \geq 1, s_2 \geq 1, \dots, s_m \geq 1} \left(\frac{2S_m}{m} \right)^{mD} e^{-\mu(S_m-m)} &= \sum_{S \geq m} \binom{S-1}{m-1} \left(\frac{2S}{m} \right)^{mD} e^{-\mu(S-m)} \\ &\leq \frac{2^{mD} e^{\mu m}}{m^{mD}(m-1)!} \sum_{S \geq m} S^{mD+m-1} e^{-\mu S} \\ &\leq \frac{2^{mD} e^{\mu m}}{m^{mD}(m-1)!} e^{-(\mu-1)m+1} (mD + m - 1)! \leq e(4D)^{mD} 2^{m-1} e^m. \end{aligned} \quad (\text{S.271})$$

We thus obtain from Eq. (S.258)

$$\mathcal{L}_{m^*} \leq \frac{e[4e\tilde{g}\gamma(4D)^D |t|]^{m^*}}{m^*!} |L^{(\xi)}| = \frac{e(e\tilde{v}|t|)^{m^*}}{m^*!} |L^{(\xi)}| \leq e \left(\frac{e^2 \tilde{v}|t|}{m^*} \right)^{m^*} |L^{(\xi)}|, \quad (\text{S.272})$$

where we use $m^*! \geq (m^*/e)^{m^*}$.

We here choose m^* such that

$$\begin{aligned} \mu(m^* - 1) - (\mu - 1) \frac{d_{L,L'}}{\xi} &\leq -\frac{\mu - 1}{2} \frac{d_{L,L'}}{\xi} \\ \text{or } m^* - 1 &= \left\lfloor \frac{\mu - 1}{2\mu} \frac{d_{L,L'}}{\xi} \right\rfloor. \end{aligned} \quad (\text{S.273})$$

Then, by using the upper-bounds for $\{\mathcal{L}_m\}_{m=1}^{m^*}$, we arrive at the following inequality:

$$\begin{aligned} \|[O_L(t), O_{L'}]\| &\leq \mathcal{L}_{m^*} + \sum_{m=1}^{m^*-1} \mathcal{L}_m \leq e |L^{(\xi)}| \left[\tilde{v}|t| \exp \left(\mu(m^* - 1) - (\mu - 1) \frac{d_{L,L'}}{\xi} + \tilde{v}|t| \right) + \left(\frac{e^2 \tilde{v}|t|}{m^*} \right)^{m^*} \right] \\ &\leq e |L^{(\xi)}| \left[\tilde{v}|t| \exp \left(-\frac{\mu - 1}{2} \frac{d_{L,L'}}{\xi} + \tilde{v}|t| \right) + \left(\frac{e^2 \tilde{v}|t|}{m^*} \right)^{m^*} \right], \end{aligned} \quad (\text{S.274})$$

where we use $\sum_{m=1}^{\infty} x^m/m! \leq x \sum_{m=0}^{\infty} x^m/(m+1)! \leq x \sum_{m=0}^{\infty} x^m/m! \leq x e^x$. We thus prove the inequality (S.248). This completes the proof. \square

C. Proof of Lemma 3: Lieb-Robinson bound for Hamiltonian with a finite interaction length

1. Statement

Lemma 3. Let \tilde{H} be a Hamiltonian such that

$$\tilde{H} = \sum_{Z \subset \Lambda: \text{diam}(Z) \leq \xi} h_Z \quad (\text{S.275})$$

with the same condition as (S.7), namely

$$\sup_{i \in \Lambda} \sum_{\substack{Z: Z \ni i \\ r \leq \text{diam}(Z) \leq \xi}} \|h_Z\| \leq gr^{-\alpha+D} \quad \text{for } r \leq \xi. \quad (\text{S.276})$$

Then, the Hamiltonian \tilde{H} satisfies $\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X})$ -Lieb-Robinson bound with

$$\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X}) = \min \left(2|\mathcal{X}| \left(\frac{v_0|t|}{e^{2\lceil x/\xi \rceil}} \right)^{\lceil x/\xi \rceil}, 2 \right) \leq 2|\mathcal{X}| e^{-2(x-\xi v_0|t|)/\xi} \quad (\text{S.277})$$

with

$$v_0 := \frac{2e^3 g \gamma (\alpha - 2D)}{\alpha - 2D - 1}. \quad (\text{S.278})$$

We note that the same inequality holds for time-dependent Hamiltonians.

Remark. When we consider a Hamiltonian with k -body interactions ($k = \mathcal{O}(1)$) (i.e., k -local Hamiltonian), the inequality (S.277) is trivially obtained for $\alpha > D$; for example, please see Sec. 2.3 in Ref. [118] or Appendix C in Ref. [24]. However, the k -dependence of the Lieb-Robinson bound is roughly given by

$$\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X}) \sim \left(\frac{\mathcal{O}(gk)|t|}{x/\xi} \right)^{x/\xi}. \quad (\text{S.279})$$

This estimation gives the Lieb-Robinson velocity of order $\mathcal{O}(1/(k\xi))$. The Hamiltonian (S.275) includes up to $\mathcal{O}(\xi^D)$ -body interactions, namely $k = \mathcal{O}(\xi^D)$. Hence, the Lieb-Robinson velocity is not given by $\mathcal{O}(\xi)$, but given by $\mathcal{O}(\xi^{D+1})$. Therefore, in order to obtain the Lieb-Robinson velocity of $\mathcal{O}(\xi)$, we need to utilize the power-law decay as in (S.276) with $\alpha > 2D + 1$.

2. Proof of Lemma 3

The proof is almost the same as Proposition 7. We note that $\min(\cdot, 2)$ is given by the trivial upper bound of $\mathcal{G}_{\tilde{H}}(x, t, \mathcal{X}) \leq 2$. We start from the same inequality as (S.256):

$$\| [O_L(\tilde{H}, t), O_{L'}] \| \leq \sum_{m=1}^{m^*} \mathcal{L}_m \quad (\text{S.280})$$

with $\|O_L\| = \|O_{L'}\| = 1$, where \mathcal{L}_m ($m < m^*$) and \mathcal{L}_{m^*} are defined as

$$\mathcal{L}_m = \frac{2(2|t|)^m}{m!} \sum_{Z_1 \cap L \neq \emptyset} \|h_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset} \|h_{Z_2}\| \cdots \sum_{\substack{Z_m \cap Z_{m-1} \neq \emptyset \\ Z_m \cap L' \neq \emptyset}} \|h_{Z_m}\| \quad (\text{S.281})$$

and

$$\mathcal{L}_{m^*} = \frac{2(2|t|)^{m^*}}{m^*!} \sum_{Z_1 \cap L \neq \emptyset} \|h_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset} \|h_{Z_2}\| \cdots \sum_{Z_{m^*} \cap Z_{m^*-1} \neq \emptyset} \|h_{Z_{m^*}}\|. \quad (\text{S.282})$$

Now, due to the finite interaction length of ξ , we have $\mathcal{L}_m = 0$ as long as $\xi m < d_{L, L'}$. Hence, if we choose

$$m^* = \lceil d_{L, L'} / \xi \rceil, \quad (\text{S.283})$$

we obtain $\mathcal{L}_m = 0$ for $m \leq m^* - 1 < d_{L,L'}/\xi$. This choice of m^* gives the inequality of

$$\|[O_L(\tilde{H}, t), O_{L'}]\| \leq \mathcal{L}_{m^*}. \quad (\text{S.284})$$

Our task is to estimate the upper bound of \mathcal{L}_{m^*} .

In order to estimate \mathcal{L}_{m^*} , we first decompose the Hamiltonian as

$$\tilde{H} = \sum_{\text{diam}(Z) \leq \xi} h_Z = \sum_{r=1}^{\xi} \sum_{Z_r: \text{diam}(Z_r)=r} h_{Z_r}. \quad (\text{S.285})$$

We notice that the condition (S.276) gives

$$\sum_{Z_r: \text{diam}(Z_r)=r, Z_r \ni i} \|h_{Z_r}\| \leq \sum_{Z: \text{diam}(Z) \geq r, Z \ni i} \|h_Z\| \leq gr^{-\alpha+D} \quad \text{for } \forall i \in \Lambda. \quad (\text{S.286})$$

Therefore, we have for arbitrary subset $L \subseteq \Lambda$

$$\sum_{Z_r: \text{diam}(Z_r)=r, Z_r \cap L \neq \emptyset} \|h_{Z_r}\| \leq \sum_{i \in L} \sum_{Z_r: \text{diam}(Z_r)=r, Z_r \ni i} \|h_{Z_r}\| \leq g|L|r^{-\alpha+D}, \quad (\text{S.287})$$

which gives the following upper bound:

$$\sum_{Z_{m^*} \cap Z_{m^*-1} \neq \emptyset} \|h_{Z_{m^*}}\| \leq g|Z_{m^*-1}| \sum_{r_{m^*}=1}^{\xi} r_{m^*}^{-\alpha+D}. \quad (\text{S.288})$$

From the inequality (S.16), we have $|Z| \leq \gamma[\text{diam}(Z)]^D$, which yields the inequality of

$$\sum_{Z_{m^*-1} \cap Z_{m^*-2} \neq \emptyset} \|h_{Z_{m^*-1}}\| \cdot |Z_{m^*-1}| \leq g|Z_{m^*-2}| \sum_{r_{m^*-1}=1}^{\xi} (\gamma r_{m^*-1}^D) r_{m^*-1}^{-\alpha+D}. \quad (\text{S.289})$$

By repeating the same process, we obtain

$$\begin{aligned} \sum_{Z_1 \cap L \neq \emptyset} \|h_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset} \|h_{Z_2}\| \cdots \sum_{Z_{m^*} \cap Z_{m^*-1} \neq \emptyset} \|h_{Z_{m^*}}\| &\leq |L| \sum_{r_1, r_2, \dots, r_{m^*}=1}^{\xi} gr_{m^*}^{-\alpha+D} \prod_{s=1}^{m^*-1} g(\gamma r_s^D) r_s^{-\alpha+D} \\ &\leq |L| \left(\sum_{r=1}^{\xi} g\gamma r^{-\alpha+2D} \right)^{m^*}, \end{aligned} \quad (\text{S.290})$$

where we use $r_{m^*}^{-\alpha+D} \leq \gamma r_{m^*}^{-\alpha+2D}$ for $r_{m^*} \geq 1$ from $\gamma \geq 1$. For $\alpha > 2D + 1$, we obtain

$$\sum_{r=1}^{\xi} r^{-\alpha+2D} \leq 1 + \sum_{r=2}^{\infty} r^{-\alpha+2D} \leq 1 + \int_1^{\infty} x^{-\alpha+2D} dx = \frac{\alpha - 2D}{\alpha - 2D - 1}. \quad (\text{S.291})$$

By applying the inequalities (S.290) and (S.291) to Eq. (S.282), we obtain the upper bound of \mathcal{L}_{m^*} by

$$\mathcal{L}_{m^*} \leq \frac{2(2|t|)^{m^*}}{m^*!} |L| \left(\frac{g\gamma(\alpha - 2D)}{\alpha - 2D - 1} \right)^{m^*} \leq 2|L| \left(\frac{2e^3 g\gamma(\alpha - 2D)}{\alpha - 2D - 1} \frac{|t|}{e^2 m^*} \right)^{m^*} = 2|L| \left(\frac{v_0 |t|}{e^2 m^*} \right)^{m^*}, \quad (\text{S.292})$$

where we use $m^*! \geq (m^*/e)^{m^*}$ and v_0 was defined in Eq. (S.278). We thus obtain the first inequality in (S.277) from the choice of m^* as in Eq. (S.283).

Finally, we need to prove

$$\min \left(2|L| \left(\frac{v_0 |t|}{e^2 \lceil x/\xi \rceil} \right)^{\lceil x/\xi \rceil}, 2 \right) \leq 2|L| e^{-2(x - \xi v_0 |t|)/\xi}. \quad (\text{S.293})$$

For $x < \xi v_0 |t|$, the inequality trivially holds, and hence we only need to consider $x \geq \xi v_0 |t|$. For $x \geq \xi v_0 |t|$, we use the following upper bounds:

$$\left(\frac{v_0 |t|}{e^2 \lceil x/\xi \rceil} \right)^{\lceil x/\xi \rceil} \leq \left(\frac{v_0 |t|}{e^2 (x/\xi)} \right)^{x/\xi} \leq e^{-2x/\xi} \leq e^{-2(x - \xi v_0 |t|)/\xi}, \quad (\text{S.294})$$

where the last inequality is a trivial inequality. We thus obtain the inequality (S.293). This completes the proof. \square

D. Proof of Proposition 8: connection of two Lieb-Robinson bounds from different unitary operators

1. Statement

Proposition 8. We consider a Hamiltonian H_0 which satisfies the Lieb-Robinson bound for a fixed t as

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) \leq \mathcal{C}e^{-(x-v|t|)/\xi} \quad (\text{S.295})$$

for $\text{diam}(\mathcal{X}), \text{diam}(\mathcal{Y}) \leq \nu\xi$ ($\nu \geq 3$). Also, let H'_0 be a Hamiltonian which satisfies the Lieb-Robinson bound for a fixed t' as

$$\mathcal{G}'(x, t', \mathcal{X}) = |\mathcal{X}^{(\nu\xi)}|^2 \mathcal{F}'(x, t'), \quad (\text{S.296})$$

where the function $\mathcal{F}'(x, t')$ is assumed to satisfy

$$\mathcal{F}'(x, t') \leq \mathcal{F}'(x_0, t')e^{-2(x-x_0)/\xi} \quad \text{for } x \geq 0, \quad 0 \leq x_0 \leq x. \quad (\text{S.297})$$

Note that we impose no assumptions on $\text{diam}(\mathcal{X})$ for $\mathcal{G}'(x, t', \mathcal{X})$. We then obtain the Lieb-Robinson function $\mathcal{G}''(x, t, t', \mathcal{X}, \mathcal{Y})$ for $e^{-iH_0 t}e^{-iH'_0 t'}$ as

$$\mathcal{G}''(x, t, t', \mathcal{X}, \mathcal{Y}) = 4\mathcal{C}\gamma(3/\nu)^D(2\kappa^* + \nu)^D e^{-(x-\kappa^*\xi-v|t|)/\xi} \quad (\text{S.298})$$

for $\text{diam}(\mathcal{X}), \text{diam}(\mathcal{Y}) \leq \nu\xi$, where κ^* is defined as an integer ($\kappa^* \in \mathbb{N}$) which satisfies

$$\mathcal{F}'(\kappa^*\xi, t') \leq \frac{1}{2^{D+1}e^{2\gamma^2 D!}} =: \zeta_1^{-1}. \quad (\text{S.299})$$

Note that t' -dependence of $\mathcal{G}''(x, t, t', \mathcal{X}, \mathcal{Y})$ is included in κ^* .

2. Proof of Proposition 8

We need to consider the norm of the commutator

$$[e^{iH'_0 t'} e^{iH_0 t} O_X e^{-iH_0 t} e^{-iH'_0 t'}, O_Y] \quad (\text{S.300})$$

for arbitrary operators O_X and O_Y with $\text{diam}(X), \text{diam}(Y) \leq \nu\xi$ ($\nu \geq 1$). In the following, we aim to estimate the upper bound of

$$\|[e^{iH'_0 t'} e^{iH_0 t} O_X e^{-iH_0 t} e^{-iH'_0 t'}, O_Y]\| = \|[O_X(H_0, t), O_Y(H'_0, -t')]\|. \quad (\text{S.301})$$

We first decompose $O_Y(H'_0, -t')$ as

$$\begin{aligned} O_Y(H'_0, -t') &= O_Y(H'_0, -t', Y[\kappa^*\xi]) + \sum_{s=\kappa^*+1}^{\infty} O_{Y[s\xi]}, \\ O_{Y[s\xi]} &:= O_Y(H'_0, -t', Y[s\xi]) - O_Y(H'_0, -t', Y[(s-1)\xi]), \end{aligned} \quad (\text{S.302})$$

where we determine κ^* afterward. From the Lieb-Robinson bound (S.296) and Lemma 1, the norm of $O_{Y[s\xi]}$ is bounded from above by

$$\begin{aligned} \|O_{Y[s\xi]}\| &\leq \|O_Y(H'_0, -t') - O_Y(H'_0, -t', Y[s\xi])\| + \|O_Y(H'_0, -t') - O_Y(H'_0, -t', Y[(s-1)\xi])\| \\ &\leq \mathcal{G}'(s\xi, t', Y) + \mathcal{G}'((s-1)\xi, t', Y) \\ &\leq 2|Y^{(\nu\xi)}|^2 \mathcal{F}'((s-1)\xi, t') \leq 2\gamma^2 \mathcal{F}'((s-1)\xi, t'), \end{aligned} \quad (\text{S.303})$$

where we use $|Y^{(\nu\xi)}| \leq \gamma[\text{diam}(Y)/(\nu\xi)]^D \leq \gamma$ which is derived from the inequality (S.18). On the other hand, we have the trivial upper bound of

$$\|O_Y(H'_0, -t', Y[\kappa^*\xi])\| \leq \|O_Y\| = 1. \quad (\text{S.304})$$

By using the decomposition (S.302), we obtain

$$\|[O_X(H_0, t), O_Y(H'_0, -t')]\| \leq \|[O_X(H_0, t), O_Y(H'_0, -t', Y[\kappa^*\xi])]\| + \sum_{s=\kappa^*+1}^{\infty} \|[O_X(H_0, t), O_{Y[s\xi]}\|]. \quad (\text{S.305})$$

In the estimation of the commutators $[O_X(H_0, t), O_{Y[s\xi]}]$, we cannot directly use the Lieb-Robinson bound (S.188) because the subset $Y[s\xi]$ may no longer satisfy the condition $\text{diam}(Y[s\xi]) \leq \nu\xi$. We thus utilize Theorem 4. From the definition of $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ in Eq. (S.295), we set

$$\mathcal{F}(t, \mathcal{X}, \mathcal{Y}) = \mathcal{C}e^{v|t|/\xi}, \quad \mathcal{L}(x) = e^{-x/\xi}, \quad (\text{S.306})$$

in Eq. (S.64). Because the assumption $\text{diam}(Y) \leq \nu\xi$ ($\nu \geq 3$) implies $\text{diam}(Y) \leq 2(\nu\xi/3) + 1$, the inequality (S.65) in the theorem gives

$$\begin{aligned} \|[O_X(H_0, t), O_Y(H'_0, -t', Y[\kappa^*\xi])]\| &\leq 2\mathcal{C}|Y[\kappa^*\xi]^{(\nu\xi/3)}|e^{-(d_{X,Y}-\kappa^*\xi-v|t|)/\xi}, \\ \|[O_X(H_0, t), O_{Y[s\xi]}]\| &\leq \|O_{Y[s\xi]}\| \cdot 2\mathcal{C}|Y[s\xi]^{(\nu\xi/3)}|e^{-(d_{X,Y}-s\xi-v|t|)/\xi}, \end{aligned} \quad (\text{S.307})$$

where we use (S.304) in the first inequality. Note that $d_{X,Y[s\xi]} = d_{X,Y} - s\xi$. Because of $\text{diam}(Y[s\xi]) \leq 2s\xi + \text{diam}(Y) \leq 2s\xi + \nu\xi$, we have

$$|Y[s\xi]^{(\nu\xi/3)}| \leq \gamma 3^D (1 + 2s/\nu)^D \quad (\text{S.308})$$

from the inequality (S.18). We thus reduce the inequality (S.305) to

$$\begin{aligned} &\|[O_X(H_0, t), O_Y(H'_0, -t')]\| \\ &\leq 2\mathcal{C}\gamma 3^D (1 + 2\kappa^*/\nu)^D e^{-(d_{X,Y}-\kappa^*\xi-v|t|)/\xi} + 2\mathcal{C}\gamma 3^D \sum_{s=\kappa^*+1}^{\infty} \|O_{Y[s\xi]}\| \cdot (1 + 2s/\nu)^D e^{-(d_{X,Y}-s\xi-v|t|)/\xi} \\ &\leq 2\mathcal{C}\gamma 3^D e^{-(d_{X,Y}-\kappa^*\xi-v|t|)/\xi} \left[(1 + 2\kappa^*/\nu)^D + 2\gamma^2 \sum_{s=\kappa^*+1}^{\infty} (1 + 2s/\nu)^D e^{s-\kappa^*} \mathcal{F}'((s-1)\xi, t') \right], \end{aligned} \quad (\text{S.309})$$

where we use the upper bound (S.303) for $\|O_{Y[s\xi]}\|$.

We then calculate an upper bound of

$$\sum_{s=\kappa^*+1}^{\infty} (1 + 2s/\nu)^D e^{s-\kappa^*} \mathcal{F}'((s-1)\xi, t') = \sum_{s=0}^{\infty} [1 + 2(\kappa^* + 1 + s)/\nu]^D e^{s+1} \mathcal{F}'((\kappa^* + s)\xi, t'). \quad (\text{S.310})$$

By using the inequality (S.297) with $x_0 = \kappa^*\xi$, we obtain

$$\mathcal{F}'((\kappa^* + s)\xi, t') \leq \mathcal{F}'(\kappa^*\xi, t') e^{-2s}, \quad (\text{S.311})$$

which upper-bounds (S.310) as

$$\begin{aligned} \sum_{s=\kappa^*+1}^{\infty} (1 + 2s/\nu)^D e^{s-\kappa^*} \mathcal{F}'((s-1)\xi, t') &\leq \mathcal{F}'(\kappa^*\xi, t') \sum_{s=0}^{\infty} [1 + 2(\kappa^* + 1 + s)/\nu]^D e^{-s+1} \\ &\leq \mathcal{F}'(\kappa^*\xi, t') e^2 (2/\nu)^D \int_0^{\infty} (x + \kappa^* + 1 + \nu/2)^D e^{-x} dx \\ &\leq \mathcal{F}'(\kappa^*\xi, t') \cdot e^2 (2/\nu)^D D! (\kappa^* + 2 + \nu/2)^D, \end{aligned} \quad (\text{S.312})$$

where in the last inequality we use (S.101). By applying the inequality (S.312) to (S.309), we finally obtain

$$\begin{aligned} &\|[O_X(H_0, t), O_Y(H'_0, -t')]\| \\ &\leq 2\mathcal{C}\gamma (3/\nu)^D e^{-(d_{X,Y}-\kappa^*\xi-v|t|)/\xi} [(2\kappa^* + \nu)^D + 2e^2\gamma^2 D! (2\kappa^* + 4 + \nu)^D \mathcal{F}'(\kappa^*\xi, t')]. \end{aligned} \quad (\text{S.313})$$

By choosing κ^* in the above inequality such that

$$\mathcal{F}'(\kappa^*\xi, t') \leq \frac{(2\kappa^* + \nu)^D}{2e^2\gamma^2 D! (2\kappa^* + 4 + \nu)^D}, \quad (\text{S.314})$$

we obtain the inequality (S.298). Because of $\kappa^* \geq 1$ and $\nu \geq 3$, we have

$$\frac{(2\kappa^* + \nu)^D}{(2\kappa^* + 4 + \nu)^D} = \left(1 + \frac{4}{2\kappa^* + \nu}\right)^{-D} \geq (9/5)^{-D} > 2^{-D}, \quad (\text{S.315})$$

and hence the condition (S.314) is simplified as

$$\mathcal{F}'(\kappa^*\xi, t') \leq \frac{1}{2^{D+1} e^2 \gamma^2 D!}. \quad (\text{S.316})$$

This completes the proof. \square

XIV. CONTRIBUTION OF THE LONG-RANGE INTERACTIONS TO LIEB-ROBINSON BOUND: PROOF OF THEOREM 3

A. Statement

Theorem 3 (Contribution by the long-range interacting terms). *Under the choice of ℓ_t by Eq. (S.35), the unitary operator (S.37) satisfies the $\mathcal{G}_{>\ell_t}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound as*

$$\mathcal{G}_{>\ell_t}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{J}_0 |\mathcal{X}^{(v^*|t|)}| \cdot |\mathcal{Y}^{(v^*|t|)}| \frac{|t|^{2D+1} \log^{2D}(x+1)}{(x - \kappa_0 v^* |t|)^\alpha}, \quad (\text{S.317})$$

where \mathcal{J}_0 and κ_0 are constants which depend only on the parameters $\{D, g_0, \alpha, \gamma\}$.

Theorem 3' (k -local Hamiltonian). *Let us assume the k -locality of the Hamiltonian [see Eq. (S.3) for the definition]. Then, under the same setup as that of Theorem 3, we obtain the Lieb-Robinson bound $\mathcal{G}_{>\ell_t}^{(k)}(x, t, \mathcal{X}, \mathcal{Y})$*

$$\mathcal{G}_{>\ell_t}^{(k)}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{J}_0^{(k)} |\mathcal{X}^{(v^*|t|)}| \cdot |\mathcal{Y}^{(v^*|t|)}| \frac{|t|^{2D+1}}{(x - \kappa_0 v^* |t|)^\alpha}, \quad (\text{S.318})$$

where $\mathcal{J}_0^{(k)}$ and κ_0 are constants which depend only on the parameters $\{D, g_0, \alpha, \gamma, k\}$.

Contribution by the long-range interactions




FIG. 10. Outline of the proof. In the proof of Theorem 3, we need to derive the Lieb-Robinson bound for $U_{>\ell_t}$ in Eq. (S.322). The proof consists of the following two steps: i) estimation of the quasi-locality of $H_{>\ell_t}(H_{\leq \ell_t}, \tau)$, ii) derivation of the Lieb-Robinson bound under the estimated quasi-locality. In the first step, we can utilize Theorem 2 for the time evolution by $H_{\leq \ell_t}$, and aim to prove Proposition 9 (the proof is given in Sec. XIV C). By using the proposition, we can derive the quasi-locality of $H_{>\ell_t}(H_{\leq \ell_t}, \tau)$ in the form of Eq. (S.347) with (S.348). Based on this estimation of the quasi-locality, we derive Proposition 11 (the proof is given in Sec. XIV D), which yields the Lieb-Robinson bound for $U_{>\ell_t}$ as in Ineq. (S.359). By appropriately transform the inequality (S.359), we arrive at the main inequality (S.317). In the case where the k -locality is assumed, the only difference arises from the point that Proposition 9 is replaced by Proposition 10.

B. Proof of the theorem (see Fig. 10 for the outline)

In the following, we define $\tilde{\alpha}$ as

$$\tilde{\alpha} := \alpha - 2D - 1, \quad (\text{S.319})$$

which gives $\tilde{\eta}$ in Eq. (S.35) as

$$\tilde{\eta} = 1 - \frac{\tilde{\alpha}}{2(\alpha - D)}. \quad (\text{S.320})$$

From the assumption of $\alpha > 2D + 1$, we have $\tilde{\alpha} > 0$. Also, the assumption (S.7) implies

$$\sum_{\substack{Z \subseteq \Lambda, Z \ni i \\ \text{diam}(Z) > \ell_t}} \|h_Z\| \leq g \ell_t^{-\alpha+D} = g |t|^{-\alpha+D+\tilde{\alpha}/2}, \quad (\text{S.321})$$

where we use the definition of $\ell_t := |t|^{\tilde{\eta}}$ [see Eq. (S.35)].

Here, the primary task is to derive the Lieb-Robinson bound for the unitary operator of

$$U_{>\ell_t} := \mathcal{T}e^{-i \int_0^t H_{>\ell_t}(H_{\leq \ell_t}, \tau) d\tau}, \quad (\text{S.322})$$

which appears from the decomposition of the unitary operator e^{-iHt} :

$$e^{-iHt} = e^{-iH_{\leq \ell_t} t} \mathcal{T} e^{-i \int_0^t H_{> \ell_t}(H_{\leq \ell_t}, \tau) d\tau}, \quad (\text{S.323})$$

Notice that we have already obtained the Lieb-Robinson bound for $e^{-iH_{\leq \ell_t} t}$ from Theorem 2.

From Theorem 2, $H_{\leq \ell_t}$ satisfies the $\mathcal{G}_{\leq \ell_t}(x, t, \mathcal{X})$ -Lieb-Robinson bound as

$$\mathcal{G}_{\leq \ell_t}(x, t, \mathcal{X}) = \tilde{\mathcal{C}}_0 |\mathcal{X}|^2 (1 + x/\ell_t)^{D-1} e^{-2(x-v^*|t|)/\ell_t}. \quad (\text{S.324})$$

We first rewrite the function $\mathcal{G}_{\leq \ell_t}(x, t, \mathcal{X})$ in the form of

$$\mathcal{G}_{\leq \ell_t}(x, t, \mathcal{X}) = |\mathcal{X}|^2 e^{-2(1-\delta_t)[x-(1-\delta_t)^{-1}v^*|t|]/\ell_t}. \quad (\text{S.325})$$

By following the same calculations in the proof of Lemma 4, we obtain δ_t as follows:

$$\delta_t = \frac{\log(\tilde{\mathcal{C}}_0) + (D-1) \log(v^*|t|/\ell_t + 1)}{2(v^*|t|/\ell_t)} \leq \frac{\log(\tilde{\mathcal{C}}_0) + (D-1) \log(v_0 \ell_1 + 1)}{2v_0 \ell_1} \leq \frac{1}{2}, \quad (\text{S.326})$$

where in the first inequality we use $v^*|t|/\ell_t = v^*|t|^{\tilde{\eta}} \geq v^* \geq v_0 \ell_1$ [see Eq. (S.153)], and in the second inequality we use the condition (S.144) for ℓ_1 . We thus obtain

$$1 - \delta_t \geq 1/2, \quad (\text{S.327})$$

which reduces (S.325) to

$$\mathcal{G}_{\leq \ell_t}(x, t, \mathcal{X}) \leq |\mathcal{X}|^2 e^{-(x-2v^*|t|)/\ell_t}. \quad (\text{S.328})$$

For the derivation of (S.317), we first show the quasi-locality of $H_{> \ell_t}(H_{\leq \ell_t}, \tau)$ for $\tau < t$. We prove the following propositions (see Sec. XIV C for the proof):

Proposition 9. *Let us consider an operator A as follows:*

$$A = \sum_{Z \subseteq \Lambda} a_Z, \quad (\text{S.329})$$

where A satisfies the assumptions (S.7) and (S.8) with $g = g_a$ and $g_0 = g_{a,0}$. We also consider an operator A_0 which satisfies $\mathcal{G}_0(x, t, \mathcal{X})$ -Lieb-Robinson bound as follows:

$$\mathcal{G}_0(x, t, \mathcal{X}) = \mathcal{C} |\mathcal{X}|^2 e^{-(x-v|t|)/\xi} \quad (\mathcal{C} \geq 1). \quad (\text{S.330})$$

For arbitrary $\tau \leq |t|$ such that $v|t|/\xi \geq 1$, we obtain the decomposition of

$$A(A_0, \tau) = \sum_{Z \subseteq \Lambda} a_{\tau, Z} \quad (\text{S.331})$$

such that

$$\begin{aligned} \sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|a_{\tau, Z}\| &\leq \min \left[J_1 g_{a,0} |X^{(v|t|)}| \cdot |Y^{(v|t|)}| (\tilde{f}_R \xi)^{2D} \left(R^{-\alpha} + e^{-(R-4\tilde{f}_R \xi)/(8\xi)} \right), J_2 g_a |X^{(v|t|)}| (\tilde{f}_R \xi)^D \right] \\ &\quad + g_a J_3 |X^{(v|t|)}| (v|t|)^D R^{-\alpha} \quad \text{with} \quad \tilde{f}_R := \log(2^{2D+1} e \gamma^2 \mathcal{C} R^{2\alpha}) + v|t|/\xi, \end{aligned} \quad (\text{S.332})$$

where $R = d_{X,Y}$ and we define $J_1 := 2^{6D+\alpha} e(2D)! \gamma^2$, $J_2 := e 6^D \gamma D!$ and $J_3 = e \gamma D! [8 \log(2e \gamma^2 \mathcal{C})]^D c_{\alpha,D}$ with $c_{\alpha,D} := D^{-1} \sup_{r \geq 1} ([1 + 2D \log(r)]^D / r^{\alpha-2D-1})$.

Because of $\tilde{f}_R \propto \log(R)$, this proposition implies that the interaction strength algebraically decays as $R^{-\alpha} \log^{2D}(R)$. In particular, if we assume the k -locality for the Hamiltonian, we can obtain a stronger statement as follows:

Proposition 10 (k -local operator). *Let us consider an operator A which includes at most k -body interactions (i.e., k -local operator), namely*

$$A = \sum_{Z \subseteq \Lambda, |Z| \leq k} a_Z. \quad (\text{S.333})$$

Then, under the same setup as that of Proposition 9, we obtain the decomposition of

$$A(A_0, \tau) = \sum_{Z \subseteq \Lambda} a_{\tau, Z} \quad (\text{S.334})$$

such that

$$\sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|a_{\tau, Z}\| \leq \min \left[J_1^{(k)} g_{a,0} |X^{(v|t|)}| \cdot |Y^{(v|t|)}| (v|t|)^{2D} \left(R^{-\alpha} + e^{-(R-4v|t|)/(8\xi)} \right), J_2^{(k)} g_a |X^{(v|t|)}| (v|t|)^D \right], \quad (\text{S.335})$$

where $R = d_{X,Y}$ and we define $J_1^{(k)} := (2e\mathcal{C}k^2)2^{6D+\alpha}e(2D)!\gamma^2$ and $J_2^{(k)} := (2e\mathcal{C}k^2)e6^D\gamma D!$.

In Proposition 9, we choose $A = H_{>\ell_t}$ and $A_0 = H_{\leq\ell_t}$, which gives the parameters

$$\xi = \ell_t, \quad \mathcal{C} = 1, \quad g_{a,0} = g_0, \quad g_a = g|t|^{-\alpha+D+\tilde{\alpha}/2}, \quad v = 2v^*. \quad (\text{S.336})$$

We note that these choices ensure the condition $v|t| \geq \xi$ because of $v^* > 1$ and $\ell_t = |t|^{\tilde{\eta}} \leq |t|$ ($|t| \geq 1$). Then, for $\tau \leq |t|$, we have

$$H_{>\ell_t}(H_{\leq\ell_t}, \tau) = \sum_{Z \subseteq \Lambda} h_{\tau, \ell_t, Z} \quad (\text{S.337})$$

with

$$\begin{aligned} \sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|h_{\tau, \ell_t, Z}\| &\leq g|t|^{-\alpha+D+\tilde{\alpha}/2} \tilde{J}_3 |X^{(2v^*|t|)}| (2v^*|t|)^D R^{-\alpha} \\ &+ \min \left[\tilde{J}_1 g_0 |X^{(2v^*|t|)}| \cdot |Y^{(2v^*|t|)}| (\tilde{f}_{t,R} \ell_t)^{2D} \left(R^{-\alpha} + e^{-(R-4\tilde{f}_{t,R} \ell_t)/(8\ell_t)} \right), \tilde{J}_2 g |t|^{-\alpha+D+\tilde{\alpha}/2} |X^{(2v^*|t|)}| (\tilde{f}_{t,R} \ell_t)^D \right] \end{aligned} \quad (\text{S.338})$$

and

$$\tilde{f}_{t,R} := \log(2^{2D+1} e \gamma^2 R^{2\alpha}) + 2v^*|t|/\ell_t, \quad (\text{S.339})$$

where $\{\tilde{J}_1, \tilde{J}_2, \tilde{J}_3\}$ are constants which are defined by $\{J_1, J_2, J_3\}$ in Proposition 9. Note that these parameters depend only on $\{D, g_0, \alpha, \gamma\}$. Furthermore, from $\tilde{f}_{t,R} \ell_t \geq 2v^*|t|$, we can reduce the inequality (S.338) to

$$\begin{aligned} \sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|h_{\tau, \ell_t, Z}\| &\leq \tilde{c}_3 \min \left[\tilde{J}_1 g_0 |X^{(v^*|t|)}| \cdot |Y^{(v^*|t|)}| (\tilde{f}_{t,R} \ell_t)^{2D} \left(R^{-\alpha} + e^{-(R-4\tilde{f}_{t,R} \ell_t)/(8\ell_t)} \right), \tilde{J}_2 g |t|^{-\alpha+D+\tilde{\alpha}/2} |X^{(v^*|t|)}| (\tilde{f}_{t,R} \ell_t)^D \right], \end{aligned} \quad (\text{S.340})$$

where the first term in (S.338) is absorbed by taking an appropriate constant \tilde{c}_3 which depends on g, g_0 and $\{\tilde{J}_1, \tilde{J}_2, \tilde{J}_3\}$. Note that we have $|X^{(2v^*|t|)}| \leq |X^{(v^*|t|)}|$.

From $\ell_t = |t|^{\tilde{\eta}}$, we have $\tilde{f}_{t,R} \ell_t$ in the form of

$$\tilde{f}_{t,R} \ell_t = |t|^{\tilde{\eta}} \log(2^{2D+1} e \gamma^2 R^{2\alpha}) + 2v^*|t|. \quad (\text{S.341})$$

From the above expression, there always exists a constant $\tilde{\kappa} (\geq 2)$ such that it satisfies the following inequalities

$$R - 4\tilde{f}_{t,R} \ell_t \geq \frac{R}{2} \quad \text{and} \quad e^{-R/(16\ell_t)} \leq R^{-\alpha} \quad \text{for} \quad R \geq \tilde{\kappa} v^*|t|, \quad (\text{S.342})$$

$$\frac{d}{dR} [R^{-\alpha} \log^{2D}(R+1)] = -\alpha R^{-\alpha+1} \log^{2D}(R+1) + \frac{2DR^{-\alpha} \log^{2D-1}(R+1)}{R+1} \leq 0 \quad \text{for} \quad R \geq \tilde{\kappa} v^*|t|, \quad (\text{S.343})$$

and

$$\tilde{f}_{t,R} \ell_t = 2v^*|t| \left(\frac{\log(2^{2D+1} e \gamma^2 R^{2\alpha})}{2v^*|t|^{1-\tilde{\eta}}} + 1 \right) \leq \begin{cases} c^* v^*|t| & \text{for } R < \tilde{\kappa} v^*|t|, \\ c^* v^*|t| \log(R+1) & \text{for } R \geq \tilde{\kappa} v^*|t|, \end{cases} \quad (\text{S.344})$$

where c_* depends on the parameters $\{D, g_0, \alpha, \gamma\}$. Note that the inequality (S.342) implies $R \geq 8\tilde{f}_{t,R} \ell_t \geq 16v^*|t|$ and hence $\tilde{\kappa}$ is at least larger than 16, namely $\tilde{\kappa} \geq 16$.

Under the above definition of $\tilde{\kappa}$, for $R \geq \tilde{\kappa} v^*|t|$, the inequalities (S.342) and (S.344) give

$$(\tilde{f}_{t,R} \ell_t)^{2D} \left(R^{-\alpha} + e^{-(R-4\tilde{f}_{t,R} \ell_t)/(8\ell_t)} \right) \leq 2(c^* v^*|t|)^{2D} R^{-\alpha} \log^{2D}(R+1). \quad (\text{S.345})$$

Also, for $R < \tilde{\kappa}v^*|t|$, the inequality (S.344) gives

$$|t|^{-\alpha+D+\tilde{\alpha}/2}(\tilde{f}_{t,R}\ell_t)^D \leq (c^*v^*)^D|t|^{-\alpha+2D+\tilde{\alpha}/2} \leq (c^*v^*)^D|t|^{-1+\tilde{\alpha}/2}, \quad (\text{S.346})$$

where the last inequality is given by the condition $\alpha > 2D + 1$. By using the inequalities (S.345) and (S.346), the inequality (S.340) reduces to

$$\sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|h_{\tau, \ell_t, Z}\| \leq g_t(R) |X^{(v^*|t)}| \cdot |Y^{(v^*|t)}| \quad (\text{S.347})$$

with

$$g_t(R) := \begin{cases} \tilde{c}_3 g \tilde{J}_2 (c^*v^*)^D |t|^{-1-\tilde{\alpha}/2} =: \tilde{J}'_2 |t|^{-1-\tilde{\alpha}/2} & \text{for } R < \tilde{\kappa}v^*|t|, \\ 2\tilde{c}_3 g_0 \tilde{J}_1 (c^*v^*)^{2D} \frac{|t|^{2D} \log^{2D}(R+1)}{R^\alpha} =: \tilde{J}'_1 |t|^{2D} R^{-\alpha} \log^{2D}(R+1) & \text{for } R \geq \tilde{\kappa}v^*|t|, \end{cases} \quad (\text{S.348})$$

where \tilde{J}'_1 and \tilde{J}'_2 depend on the parameters $\{D, g_0, \alpha, \gamma\}$. Note that the function $g_t(R)$ monotonically decreases with R because of the condition (S.343).

In the case where the Hamiltonian is k -local, we can apply the same analyses. Only the difference is that $\tilde{f}_{t,R}\ell_t$ is replaced with $2v^*|t|$ from Proposition 10. Hence, the logarithmic dependence $\log^{2D}(R+1)$, which results from the inequality (S.344), does not appear in this case. Then, the inequality (S.347) is replaced by

$$\sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|h_{\tau, \ell_t, Z}\| \leq g_t^{(k)}(R) |X^{(v^*|t)}| \cdot |Y^{(v^*|t)}| \quad (\text{S.349})$$

with

$$g_t^{(k)}(R) := \begin{cases} \tilde{J}_2^{(k)} |t|^{-1-\tilde{\alpha}/2} & \text{for } R < \tilde{\kappa}v^*|t|, \\ \tilde{J}_1^{(k)} |t|^{2D} R^{-\alpha} & \text{for } R \geq \tilde{\kappa}v^*|t|, \end{cases} \quad (\text{S.350})$$

where $\tilde{J}_1^{(k)}$ and $\tilde{J}_2^{(k)}$ are constants which depend on the parameters $\{D, g_0, \alpha, \gamma, k\}$.

We here show the Lieb-Robinson bound for time-evolved long-range Hamiltonians $H_{>\ell_t}(H_{\leq \ell_t}, \tau)$. We can prove the following proposition (see Sec. XIV D for the proof):

Proposition 11. *Let us consider a Hamiltonian \tilde{H} as follows:*

$$\tilde{H} = \sum_{Z \subseteq \Lambda} \tilde{h}_Z \quad (\text{S.351})$$

with

$$\sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|\tilde{h}_Z\| \leq |X^{(\xi_0)}| \cdot |Y^{(\xi_0)}| g(R), \quad R = d_{X,Y},$$

$$g(R) = \begin{cases} \tilde{g} & \text{for } R < \kappa \xi_0, \\ \tilde{g}_R R^{-\alpha} & \text{for } R \geq \kappa \xi_0, \end{cases} \quad \tilde{g}_R := \tilde{g}_0 \log^p(R+1) \quad (p \in \mathbb{N}), \quad (\text{S.352})$$

where $\kappa \geq 2$, $\xi_0 \geq 1$, and $g(R)$ monotonically decreases with R . Then, the Hamiltonian \tilde{H} satisfies $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound with

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = |\mathcal{X}^{(\xi_0)}| \cdot |\mathcal{Y}^{(\xi_0)}| \frac{2^{\alpha+1} e \Gamma(\alpha+2)}{\tilde{\lambda}} \left(e^{2e\tilde{\lambda}\gamma^2 9^D |t|} - 1 \right) g(R) + 2 |X^{(\xi_0)}| \left(\frac{4e\kappa\xi_0 \tilde{\lambda} \gamma^2 9^D |t|}{R} \right)^{R/(2\kappa\xi_0)}, \quad (\text{S.353})$$

where $\Gamma(x)$ is the gamma function and we define $\tilde{\lambda}$ as

$$\tilde{\lambda} := \tilde{g} \gamma [2(\kappa+3)]^D + \frac{c_{p,\kappa} \tilde{g}_0 \gamma D 2^{D+1} \log^p(\kappa\xi_0+1)}{\kappa+2} \frac{1}{\xi_0^\alpha},$$

$$c_{p,\kappa} := \sup_{z \in \mathbb{R} | z \geq \kappa+2} \left[\frac{(\log(z)+1)^p}{(z-2)^{\alpha-D-1}} \right]. \quad (\text{S.354})$$

In the following, we apply Proposition 11 to the unitary operator of

$$\mathcal{T} e^{-i \int_0^t H_{>\ell_t}(H_{\leq \ell_t}, \tau) d\tau}. \quad (\text{S.355})$$

From the inequality (S.347) with Eq. (S.348), the parameters in Proposition 11 is now given by

$$\xi_0 = v^*|t|, \quad \kappa = \tilde{\kappa}, \quad p = 2D, \quad \tilde{g} = \tilde{J}'_2|t|^{-\tilde{\alpha}/2}, \quad \tilde{g}_0 = \tilde{J}'_1|t|^{2D}. \quad (\text{S.356})$$

Then, the definition (S.354) gives $\tilde{\lambda}|t|$ as follows:

$$\begin{aligned} \tilde{\lambda}|t| &= \tilde{J}'_2\gamma[2(\tilde{\kappa}+3)]^D|t|^{-\tilde{\alpha}/2} + \frac{c_{2D,\tilde{\kappa}}\tilde{J}'_1\gamma D 2^{D+1}}{\tilde{\kappa}+2}|t|^{2D+1} \frac{\log^{2D}(\tilde{\kappa}v^*|t|+1)}{(v^*|t|)^\alpha} \\ &= \tilde{J}'_2\gamma[2(\tilde{\kappa}+3)]^D|t|^{-\tilde{\alpha}/2} + \frac{c_{2D,\tilde{\kappa}}\tilde{J}'_1\gamma D 2^{D+1}}{[v^*]^\alpha(\tilde{\kappa}+2)}|t|^{-\tilde{\alpha}} \log^{2D}(\tilde{\kappa}v^*|t|+1), \end{aligned} \quad (\text{S.357})$$

where $\tilde{\alpha} := \alpha - 2D - 1$ as defined in Eq. (S.319). Because of $\tilde{\alpha} > 0$ and $|t| \geq 1$, there exists a constant λ_0 which depends only on the parameters $\{D, g_0, \alpha, \gamma\}$ such that

$$\tilde{\lambda}|t| \leq \lambda_0|t|^{-\tilde{\alpha}/2} \leq \lambda_0. \quad (\text{S.358})$$

Then, from the inequality (S.353) in Proposition 11, the unitary operator (S.355) satisfies $\mathcal{G}_{>\ell_t}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound with

$$\begin{aligned} \mathcal{G}_{>\ell_t}(x, t, \mathcal{X}, \mathcal{Y}) &= |\mathcal{X}^{(v^*|t|)}| \cdot |\mathcal{Y}^{(v^*|t|)}| \frac{2^{\alpha+1}e\Gamma(\alpha+2) \left(e^{2e\lambda_0\gamma^2 9^D} - 1 \right)}{\lambda_0} |t| g_t(R) \\ &\quad + 2|X^{(v^*|t|)}| \left(\frac{4e\tilde{\kappa}v^*\lambda_0\gamma^2 9^D |t|^{1-\tilde{\alpha}/2}}{R} \right)^{R/(2\tilde{\kappa}v^*|t|)}. \end{aligned} \quad (\text{S.359})$$

From the definition of $g_t(R)$ in Eq. (S.348), the first term in (S.359) decays as $|t|^{2D+1}R^{-\alpha} \log^{2D}(R+1)$ for $R \geq \tilde{\kappa}v^*|t|$. On the other hand, the second term exponentially decays as $(R^{-1}|t|^{1-\tilde{\alpha}/2})^{R/|t|}$. Hence, there exists a constant $\kappa_0 \geq \tilde{\kappa}$ such that

$$\text{First term in (S.359)} \geq \text{Second term in (S.359)} \quad \text{for } R \geq \kappa_0 v^*|t|. \quad (\text{S.360})$$

We then obtain

$$\mathcal{G}_{>\ell_t}(x, t, \mathcal{X}, \mathcal{Y}) = 2|\mathcal{X}^{(v^*|t|)}| \cdot |\mathcal{Y}^{(v^*|t|)}| \frac{2^{\alpha+1}e\Gamma(\alpha+2) \left(e^{2e\lambda_0\gamma^2 9^D} - 1 \right)}{\lambda_0} \tilde{J}'_1|t|^{2D+1} R^{-\alpha} \log^{2D}(R+1) \quad (\text{S.361})$$

for $R \geq \kappa_0 v^*|t|$. Therefore, we finally obtain

$$\mathcal{G}_{>\ell_t}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{J}_0 |\mathcal{X}^{(v^*|t|)}| \cdot |\mathcal{Y}^{(v^*|t|)}| \frac{|t|^{2D+1} \log^{2D}(R+1)}{R^\alpha} \quad \text{for } R \geq \kappa_0 v^*|t| \quad (\text{S.362})$$

by choosing

$$\mathcal{J}_0 = \frac{2^{\alpha+2}e\Gamma(\alpha+2) \left(e^{2e\lambda_0\gamma^2 9^D} - 1 \right)}{\lambda_0} \tilde{J}'_1. \quad (\text{S.363})$$

The inequality (S.362) reduces to the main inequality (S.317).

In the case where the Hamiltonian is k -local, from the inequality (S.349) with Eq. (S.350), we set the parameters in Proposition 11 as follows:

$$\xi_0 = v^*|t|, \quad \kappa = \tilde{\kappa}, \quad p = 0, \quad \tilde{g} = \tilde{J}_2^{(k)}|t|^{-1-\tilde{\alpha}/2}, \quad \tilde{g}_0 = \tilde{J}_1^{(k)}|t|^{2D}. \quad (\text{S.364})$$

By following the same steps for the derivation of (S.362), we obtain for the k -local Hamiltonians

$$\mathcal{G}_{>\ell_t}(x, t, \mathcal{X}, \mathcal{Y}) = \mathcal{J}_0^{(k)} |\mathcal{X}^{(v^*|t|)}| \cdot |\mathcal{Y}^{(v^*|t|)}| \frac{|t|^{2D+1}}{R^\alpha} \quad \text{for } R \geq \kappa_0 v^*|t|, \quad (\text{S.365})$$

where $\mathcal{J}_0^{(k)}$ is a constant which depends on $\{D, g_0, \alpha, \gamma, k\}$. This completes the proof of the theorem. \square

C. Proofs of Propositions 9 and 10

1. Statement

Proposition 9. *Let us consider an operator A as follows:*

$$A = \sum_{Z \subseteq \Lambda} a_Z, \quad (\text{S.366})$$

where A satisfies the assumptions (S.7) and (S.8) with $g = g_a$ and $g_0 = g_{a,0}$. We also consider an operator A_0 which satisfies $\mathcal{G}_0(x, t, \mathcal{X})$ -Lieb-Robinson bound as follows:

$$\mathcal{G}_0(x, t, \mathcal{X}) = \mathcal{C}|\mathcal{X}|^2 e^{-(x-v|t|)/\xi} \quad (\mathcal{C} \geq 1). \quad (\text{S.367})$$

For arbitrary $\tau \leq |t|$ such that $v|t|/\xi \geq 1$, we obtain the decomposition of

$$A(A_0, \tau) = \sum_{Z \subseteq \Lambda} a_{\tau, Z} \quad (\text{S.368})$$

such that

$$\begin{aligned} \sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|a_{\tau, Z}\| &\leq \min \left[J_1 g_{a,0} |X^{(v|t|)}| \cdot |Y^{(v|t|)}| (\tilde{f}_R \xi)^{2D} \left(R^{-\alpha} + e^{-(R-4\tilde{f}_R \xi)/(8\xi)} \right), J_2 g_a |X^{(v|t|)}| (\tilde{f}_R \xi)^D \right] \\ &\quad + g_a J_3 |X^{(v|t|)}| (v|t|)^D R^{-\alpha} \quad \text{with} \quad \tilde{f}_R := \log(2^{2D+1} e \gamma^2 \mathcal{C} R^{2\alpha}) + v|t|/\xi, \end{aligned} \quad (\text{S.369})$$

where $R = d_{X,Y}$ and we define $J_1 := 2^{6D+\alpha} e(2D)! \gamma^2$, $J_2 := e 6^D \gamma D!$ and $J_3 = e \gamma D! [8 \log(2e \gamma^2 \mathcal{C})]^D c_{\alpha,D}$ with $c_{\alpha,D} := D^{-1} \sup_{r \geq 1} ([1 + 2D \log(r)]^D / r^{\alpha-2D-1})$.

Proposition 10 (k -local operator). *Let us consider an operator A which includes at most k -body interactions (i.e., k -local operator), namely*

$$A = \sum_{Z \subset \Lambda, |Z| \leq k} a_Z. \quad (\text{S.370})$$

Then, under the same setup as that of Proposition 9, we obtain the decomposition of

$$A(A_0, \tau) = \sum_{Z \subseteq \Lambda} a_{\tau, Z} \quad (\text{S.371})$$

such that

$$\sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|a_{\tau, Z}\| \leq \min \left[J_1^{(k)} g_{a,0} |X^{(v|t|)}| \cdot |Y^{(v|t|)}| (v|t|)^{2D} \left(R^{-\alpha} + e^{-(R-4v|t|)/(8\xi)} \right), J_2^{(k)} g_a |X^{(v|t|)}| (v|t|)^D \right], \quad (\text{S.372})$$

where $R = d_{X,Y}$ and we define $J_1^{(k)} := (2e \mathcal{C} k^2) 2^{6D+\alpha} e(2D)! \gamma^2$ and $J_2^{(k)} := (2e \mathcal{C} k^2) e 6^D \gamma D!$.

2. Proofs of Propositions 9 and 10.

[Few-body (k -local) operator]

We first consider the case where the operator A includes up to k -body interacting terms:

$$A = \sum_{Z \subset \Lambda, |Z| \leq k} a_Z. \quad (\text{S.373})$$

In this case, the proof is much simpler than that for generic A . We here denote

$$d_{X,Y} = R \quad (\text{S.374})$$

for the simplicity.

For the proof, we decompose A as

$$A = \sum_{r=1}^{\infty} \sum_{\substack{Z_r \subset \Lambda \\ |Z_r| \leq k, \text{diam}(Z_r)=r}} a_{Z_r}. \quad (\text{S.375})$$

We then obtain $A(A_0, \tau)$ in the following form:

$$A(A_0, \tau) = \sum_{r=1}^{\infty} \sum_{\substack{Z_r \subset \Lambda \\ |Z_r| \leq k, \text{diam}(Z_r)=r}} a_{Z_r}(A_0, \tau). \quad (\text{S.376})$$

We further decompose each of $\{a_{Z_r}(A_0, \tau)\}_{Z_r \subset \Lambda}$ as follows:

$$a_{Z_r}(A_0, \tau) = \sum_{s=0}^{\infty} a_{\tau, Z_r[s\xi + \xi_r]} \quad \text{with} \quad \text{diam}(Z_r) = r, \quad (\text{S.377})$$

where we define

$$a_{\tau, Z_r[s\xi + \xi_r]} := \begin{cases} a_{Z_r}(A_0, \tau, Z_r[\xi_r]) & \text{for } s = 0, \\ a_{Z_r}(A_0, \tau, Z_r[s\xi + \xi_r]) - a_{Z_r}(A_0, \tau, Z_r[(s-1)\xi + \xi_r]) & \text{for } s \geq 1. \end{cases} \quad (\text{S.378})$$

The parameter ξ_r is appropriately chosen afterward. By combining Eqs. (S.376) and (S.377), we have

$$A(A_0, \tau) = \sum_{r=1}^{\infty} \sum_{\substack{Z_r \subset \Lambda \\ |Z_r| \leq k, \text{diam}(Z_r) = r}} \sum_{s=0}^{\infty} a_{\tau, Z_r[s\xi + \xi_r]}. \quad (\text{S.379})$$

In order to obtain the inequality (S.369), we need to estimate the following quantity:

$$\sum_{r=1}^{\infty} \sum_{s=0}^{\infty} \sum_{\substack{Z_r \subset \Lambda, |Z_r| \leq k, \text{diam}(Z_r) = r \\ Z_r[s\xi + \xi_r] \cap X \neq \emptyset, Z_r[s\xi + \xi_r] \cap Y \neq \emptyset}} \|a_{\tau, Z_r[s\xi + \xi_r]}\|. \quad (\text{S.380})$$

We notice that the condition $Z_r[s\xi + \xi_r] \cap X \neq \emptyset$ for $Z_r \subset \Lambda$ is equivalent to $Z_r \cap X[s\xi + \xi_r] \neq \emptyset$, and hence

$$\sum_{r=1}^{\infty} \sum_{s=0}^{\infty} \sum_{\substack{Z_r \subset \Lambda, |Z_r| \leq k, \text{diam}(Z_r) = r \\ Z_r[s\xi + \xi_r] \cap X \neq \emptyset, Z_r[s\xi + \xi_r] \cap Y \neq \emptyset}} \|a_{\tau, Z_r[s\xi + \xi_r]}\| = \sum_{r=1}^{\infty} \sum_{s=0}^{\infty} \sum_{\substack{Z_r \subset \Lambda, |Z_r| \leq k, \text{diam}(Z_r) = r \\ Z_r \cap X[s\xi + \xi_r] \neq \emptyset, Z_r \cap Y[s\xi + \xi_r] \neq \emptyset}} \|a_{\tau, Z_r[s\xi + \xi_r]}\|. \quad (\text{S.381})$$

By using the $\mathcal{G}_0(x, t, \mathcal{X})$ -Lieb-Robinson bound of A_0 and Lemma 1, we obtain

$$\begin{aligned} \|a_{\tau, Z_r[s\xi + \xi_r]}\| &= \|a_{Z_r}(A_0, \tau, Z_r[s\xi + \xi_r]) - a_{Z_r}(A_0, \tau, Z_r[(s-1)\xi + \xi_r])\| \\ &\leq \|a_{Z_r}\| \mathcal{G}_0(s\xi + \xi_r, \tau, Z_r) + \|a_{Z_r}\| \mathcal{G}_0((s-1)\xi + \xi_r, \tau, Z_r) \\ &\leq 2\|a_{Z_r}\| \mathcal{G}_0((s-1)\xi + \xi_r, |t|, Z_r), \end{aligned} \quad (\text{S.382})$$

where we use $\mathcal{G}_0(x, \tau, \mathcal{X}) \leq \mathcal{G}_0(x, |t|, \mathcal{X})$ for $\tau \leq |t|$. By using the k -locality of the operator A , namely $|Z_r| \leq k$ for arbitrary r and choosing ξ_r as

$$\xi_r = v|t| \quad \text{for } \forall r \in \mathbb{N}, \quad (\text{S.383})$$

we reduce the inequality (S.382) to

$$\begin{aligned} \|a_{\tau, Z_r[s\xi + \xi_r]}\| &\leq 2\mathcal{C}|Z_r|^2 \|a_{Z_r}\| e^{-(s-1) - (\xi_r - v|t|)/\xi} \\ &\leq 2e\mathcal{C}k^2 e^{-(\xi_r - v|t|)/\xi} \|a_{Z_r}\| e^{-s} = 2e\mathcal{C}k^2 \|a_{Z_r}\| e^{-s}. \end{aligned} \quad (\text{S.384})$$

For $\|a_{\tau, Z_r[\xi_r]}\|$, we obtain the following trivial inequality:

$$\|a_{\tau, Z_r[\xi_r]}\| = \|a_{Z_r}(A_0, \tau, Z_r[\xi_r])\| \leq \|a_{Z_r}\| \leq 2e\mathcal{C}k^2 \|a_{Z_r}\|. \quad (\text{S.385})$$

Thus, we obtain the following upper bound for the summation (S.381):

$$\begin{aligned} \sum_{r=1}^{\infty} \sum_{s=0}^{\infty} \sum_{\substack{Z_r \subset \Lambda, |Z_r| \leq k, \text{diam}(Z_r) = r \\ Z_r \cap X[s\xi + \xi_r] \neq \emptyset, Z_r \cap Y[s\xi + \xi_r] \neq \emptyset}} \|a_{\tau, Z_r[s\xi + \xi_r]}\| &\leq 2e\mathcal{C}k^2 \sum_{s=0}^{\infty} e^{-s} \sum_{r=1}^{\infty} \sum_{\substack{Z_r \subset \Lambda, |Z_r| \leq k, \text{diam}(Z_r) = r \\ Z_r \cap X[s\xi + \xi_r] \neq \emptyset, Z_r \cap Y[s\xi + \xi_r] \neq \emptyset}} \|a_{Z_r}\| \\ &= 2e\mathcal{C}k^2 \sum_{s=0}^{\infty} e^{-s} \sum_{\substack{Z \subset \Lambda, |Z| \leq k \\ Z \cap X[s\xi + v|t|] \neq \emptyset, Z \cap Y[s\xi + v|t|] \neq \emptyset}} \|a_Z\| \\ &\leq 2e\mathcal{C}k^2 \sum_{s=0}^{\infty} e^{-s} \frac{g_{a,0} |X[s\xi + v|t|]| \cdot |Y[s\xi + v|t|]|}{[\max(R - 2s\xi - 2v|t|, 0) + 1]^\alpha} \end{aligned} \quad (\text{S.386})$$

with $R = d_{X,Y}$, where the last inequality is derived from the assumption (S.11) with $g_0 = g_{a,0}$ for the operator A . Note that we have $d_{X[s\xi + v|t|], Y[s\xi + v|t|]} \geq d_{X,Y} - 2(s\xi + v|t|) = R - 2(s\xi + v|t|)$.

We define s_R as $s_R := (R - 4v|t|)/(4\xi)$, which satisfies $R - 2s_R\xi - 2v|t| = R/2$. We first consider the case of $s_R > 0$ (i.e., $R > 4v|t|$). From the inequalities (S.15) and (S.20), we obtain

$$|X[s\xi + v|t|]| \leq |X^{(v|t|)}[s\xi + 2v|t|]| \leq \gamma |X^{(v|t|)}| \cdot [2(s\xi + 2v|t|)]^D. \quad (\text{S.387})$$

Then, for $s < s_R$, we obtain the following upper bound for the summation of

$$\begin{aligned} \sum_{s < s_R} e^{-s} \frac{g_{a,0} |X[s\xi + v|t|]| \cdot |Y[s\xi + v|t|]|}{[\max(R - 2s\xi - 2v|t|, 0) + 1]^\alpha} &\leq \frac{g_{a,0}\gamma^2 |X^{(v|t|)}| \cdot |Y^{(v|t|)}|}{(R/2)^\alpha} \sum_{s < s_R} [2(s\xi + 2v|t|)]^{2D} e^{-s} \\ &\leq \frac{e^{2D+\alpha} (2D)! g_{a,0}\gamma^2 |X^{(v|t|)}| \cdot |Y^{(v|t|)}|}{R^\alpha} (\xi + 2v|t|)^{2D}, \end{aligned} \quad (\text{S.388})$$

where we use $R - 2s\xi - 2v|t| \geq R/2$ for $s < s_R$ in the first inequality. Also, the second inequality is derived from

$$\begin{aligned} \sum_{s < s_R} [2(s\xi + 2v|t|)]^{2D} e^{-s} &\leq (2\xi)^{2D} \sum_{s=0}^{\infty} (s + 2v|t|/\xi)^{2D} e^{-s} \\ &\leq (2\xi)^{2D} \int_0^{\infty} (x + 2v|t|/\xi)^{2D} e^{-x+1} dx \leq e(2\xi)^{2D} (2D)! (1 + 2v|t|/\xi)^{2D}, \end{aligned} \quad (\text{S.389})$$

where we utilize the inequality (S.101). For $s \geq s_R$, we obtain

$$\begin{aligned} \sum_{s \geq s_R} e^{-s} \frac{g_{a,0} |X[s\xi + v|t|]| \cdot |Y[s\xi + v|t|]|}{[\max(R - 2s\xi - 2v|t|, 0) + 1]^\alpha} &\leq g_{a,0}\gamma^2 |X^{(v|t|)}| \cdot |Y^{(v|t|)}| \sum_{s \geq s_R} [2(s\xi + 2v|t|)]^{2D} e^{-s} \\ &\leq g_{a,0}\gamma^2 |X^{(v|t|)}| \cdot |Y^{(v|t|)}| e^{-s_R/2} \sum_{s \geq s_R} [2(s\xi + 2v|t|)]^{2D} e^{-s/2} \\ &\leq e^{2D+1} (2D)! g_{a,0}\gamma^2 |X^{(v|t|)}| \cdot |Y^{(v|t|)}| (2\xi + 2v|t|)^{2D} e^{-s_R/2}, \end{aligned} \quad (\text{S.390})$$

where we use the same inequality as (S.389) in estimating the summation in the third inequality. By applying the above two upper bounds to (S.386), we arrive at

$$\begin{aligned} &\sum_{r=1}^{\infty} \sum_{s=0}^{\infty} \sum_{\substack{Z_r \subset \Lambda, |Z_r| \leq k, \text{diam}(Z_r)=r \\ Z_r \cap X[s\xi + \xi_r] \neq \emptyset, Z_r \cap Y[s\xi + \xi_r] \neq \emptyset}} \|a_{\tau, Z_r[s\xi + \xi_r]}\| \\ &\leq 2e\mathcal{C}k^2 \left(\frac{e^{2D+\alpha} (2D)! g_{a,0}\gamma^2 |X^{(v|t|)}| \cdot |Y^{(v|t|)}|}{R^\alpha} (\xi + 2v|t|)^{2D} + e^{2D+1} (2D)! g_{a,0}\gamma^2 |X^{(v|t|)}| \cdot |Y^{(v|t|)}| (2\xi + 2v|t|)^{2D} e^{-s_R/2} \right) \\ &\leq (2e\mathcal{C}k^2) 2^{2D+\alpha} e(2D)! g_{a,0}\gamma^2 |X^{(v|t|)}| \cdot |Y^{(v|t|)}| (2\xi + 2v|t|)^{2D} \left(R^{-\alpha} + e^{-(R-4v|t|)/(8\xi)} \right) \\ &\leq (2e\mathcal{C}k^2) 2^{6D+\alpha} e(2D)! g_{a,0}\gamma^2 |X^{(v|t|)}| \cdot |Y^{(v|t|)}| (v|t|)^{2D} \left(R^{-\alpha} + e^{-(R-4v|t|)/(8\xi)} \right), \end{aligned} \quad (\text{S.391})$$

where in the last inequality we use $2v|t| + 2\xi \leq 4v|t|$ from the assumption of $v|t| \geq \xi$.

We second consider the case of $s_R \leq 0$ (i.e., $R \leq 4v|t|$). We take a different approach to estimate the summation (S.386) as follows:

$$\sum_{r=1}^{\infty} \sum_{s=0}^{\infty} \sum_{\substack{Z_r \subset \Lambda, |Z_r| \leq k, \text{diam}(Z_r)=r \\ Z_r \cap X[s\xi + \xi_r] \neq \emptyset, Z_r \cap Y[s\xi + \xi_r] \neq \emptyset}} \|a_{\tau, Z_r[s\xi + \xi_r]}\| \leq 2e\mathcal{C}k^2 \sum_{s=0}^{\infty} e^{-s} \sum_{\substack{Z \subset \Lambda, |Z| \leq k \\ Z \cap X[s\xi + v|t|] \neq \emptyset}} \|a_Z\|. \quad (\text{S.392})$$

By applying the assumption (S.7) with $g = g_a$ and the inequality (S.387), we have

$$\sum_{\substack{Z \subset \Lambda, |Z| \leq k \\ Z \cap X[s\xi + v|t|] \neq \emptyset}} \|a_Z\| \leq g_a |X[s\xi + v|t|]| \leq g_a \gamma |X^{(v|t|)}| \cdot [2(s\xi + 2v|t|)]^D, \quad (\text{S.393})$$

which gives the upper bound of

$$\sum_{s=0}^{\infty} e^{-s} \sum_{\substack{Z \subset \Lambda, |Z| \leq k \\ Z \cap X[s\xi + v|t|] \neq \emptyset}} \|a_Z\| \leq g_a \gamma |X^{(v|t|)}| \sum_{s=0}^{\infty} e^{-s} [2(s\xi + 2v|t|)]^D \leq e^{2D} g_a \gamma |X^{(v|t|)}| D! (\xi + 2v|t|)^D, \quad (\text{S.394})$$

where we use the same upper bound as (S.389). By combining the inequalities (S.392) and (S.394), we obtain

$$\begin{aligned} \sum_{r=1}^{\infty} \sum_{s=0}^{\infty} \sum_{\substack{Z_r \subset \Lambda, |Z_r| \leq k, \text{diam}(Z_r)=r \\ Z_r \cap X[s\xi + \xi_r] \neq \emptyset, Z_r \cap Y[s\xi + \xi_r] \neq \emptyset}} \|a_{\tau, Z_r[s\xi + \xi_r]}\| &\leq (2e\mathcal{C}k^2)e2^D g_a \gamma |X^{(v|t)}| D!(\xi + 2v|t|)^D \\ &\leq (2e\mathcal{C}k^2)e6^D g_a \gamma |X^{(v|t)}| D!(v|t|)^D, \end{aligned} \quad (\text{S.395})$$

where in the last inequality we use $v|t| + \xi \leq 3v|t|$ from the assumption of $v|t| \geq \xi$.

From the upper bounds of (S.391) and (S.395), we finally obtain

$$\begin{aligned} &\sum_{r=1}^{\infty} \sum_{s=0}^{\infty} \sum_{\substack{Z_r \subset \Lambda, |Z_r| \leq k, \text{diam}(Z_r)=r \\ Z_r \cap X[s\xi + \xi_r] \neq \emptyset, Z_r \cap Y[s\xi + \xi_r] \neq \emptyset}} \|a_{\tau, Z_r[s\xi + \xi_r]}\| \\ &\leq \min \left[(2e\mathcal{C}k^2)2^{6D+\alpha} e(2D)! g_{a,0} \gamma^2 |X^{(v|t)}| \cdot |Y^{(v|t)}| (v|t|)^{2D} \left(R^{-\alpha} + e^{-(R-4v|t|)/(8\xi)} \right), (2e\mathcal{C}k^2)e6^D g_a \gamma |X^{(v|t)}| D!(v|t|)^D \right] \\ &= \min \left[J_1^{(k)} g_{a,0} |X^{(v|t)}| \cdot |Y^{(v|t)}| (v|t|)^{2D} \left(R^{-\alpha} + e^{-(R-4v|t|)/(8\xi)} \right), J_2^{(k)} g_a |X^{(v|t)}| (v|t|)^D \right] \end{aligned} \quad (\text{S.396})$$

with

$$J_1^{(k)} := (2e\mathcal{C}k^2)2^{6D+\alpha} e(2D)! \gamma^2, \quad J_2^{(k)} := (2e\mathcal{C}k^2)e6^D \gamma D!. \quad (\text{S.397})$$

This completes the proof of Proposition 10. \square

[generic cases]

We here consider the general case without the assumption of the k -locality:

$$A = \sum_{Z \subseteq \Lambda} a_Z. \quad (\text{S.398})$$

In this case, we also start from the equation (S.381). Here, in estimating the norm of $\|a_{\tau, Z_r[s\xi + \xi_r]}\|$, we cannot utilize the upper bound of $|Z_r|^2 \leq k^2$ as in (S.384). Instead, we utilize the following upper bound which is derived from the inequality (S.16):

$$|Z_r| \leq \gamma [\text{diam}(Z_r)]^D = \gamma r^D. \quad (\text{S.399})$$

This replaces the inequality (S.384) by

$$\|a_{\tau, Z_r[s\xi + \xi_r]}\| \leq 2e\gamma^2 \mathcal{C} \|a_{Z_r}\| r^{2D} e^{-(\xi_r - v|t|)/\xi} e^{-s} \quad (s \geq 1). \quad (\text{S.400})$$

We here choose ξ_r as

$$\xi_r = f_r \xi, \quad f_r := \log(2e\gamma^2 \mathcal{C} r^{2D}) + v|t|/\xi. \quad (\text{S.401})$$

By applying the above choice of ξ_r to (S.400), we obtain

$$\|a_{\tau, Z_r[s\xi + \xi_r]}\| \leq \|a_{Z_r}\| e^{-s}. \quad (\text{S.402})$$

We note that this inequality also holds for $s = 0$ since $\|a_{\tau, Z_r[\xi_r]}\| = \|a_{Z_r}\|$ from the definition (S.378). We then estimate the same quantity as (S.386):

$$\sum_{r=1}^{\infty} \sum_{s=0}^{\infty} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[s\xi + \xi_r] \neq \emptyset, Z_r \cap Y[s\xi + \xi_r] \neq \emptyset}} \|a_{\tau, Z_r[s\xi + \xi_r]}\| \leq \sum_{s=0}^{\infty} e^{-s} \sum_{r=1}^{\infty} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[(s+f_r)\xi] \neq \emptyset, Z_r \cap Y[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\|. \quad (\text{S.403})$$

In order to estimate the summation with respect to r , we analyze the cases of $r \leq R_0$ and $r > R_0$ in different ways, where we will choose R_0 afterward. For $r \leq R_0$, we use the assumption (S.11) with $g_0 = g_{a,0}$, which implies

$$\begin{aligned} \sum_{r=1}^{R_0} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[(s+f_r)\xi] \neq \emptyset, Z_r \cap Y[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\| &\leq \sum_{\substack{Z \subseteq \Lambda \\ Z \cap X[(s+f_{R_0})\xi] \neq \emptyset, Z \cap Y[(s+f_{R_0})\xi] \neq \emptyset}} \|a_Z\| \\ &\leq \frac{g_{a,0} |X[(s+f_{R_0})\xi]| \cdot |Y[(s+f_{R_0})\xi]|}{[\max(R - 2s\xi - 2f_{R_0}\xi, 0) + 1]^\alpha}, \end{aligned} \quad (\text{S.404})$$

where we use $f_r \leq f_{R_0}$ for $r \leq R_0$ from the definition (S.401). From this inequality, we obtain

$$\sum_{s=0}^{\infty} e^{-s} \sum_{r=1}^{R_0} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[(s+f_r)\xi] \neq \emptyset, Z_r \cap Y[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\| \leq g_{a,0} \sum_{s=0}^{\infty} e^{-s} \frac{|X[(s+f_{R_0})\xi]| \cdot |Y[(s+f_{R_0})\xi]|}{[\max(R-2s\xi-2f_{R_0}\xi, 0) + 1]^\alpha}, \quad (\text{S.405})$$

and the remaining task is to estimate the summation with respect to s . This summation has been already estimated in (S.386), where we only need to replace $v|t|$ with $f_{R_0}\xi$. Hence, we can obtain the similar upper bound to (S.396):

$$\begin{aligned} & g_{a,0} \sum_{s=0}^{\infty} e^{-s} \frac{|X[(s+f_{R_0})\xi]| \cdot |Y[(s+f_{R_0})\xi]|}{[\max(R-2s\xi-2f_{R_0}\xi, 0) + 1]^\alpha} \\ & \leq \min \left[2^{2D+\alpha} e(2D)! g_{a,0} \gamma^2 |X^{(v|t)}| \cdot |Y^{(v|t)}| (f_{R_0}\xi + 2\xi + v|t|)^{2D} \left(R^{-\alpha} + e^{-(R-4f_{R_0}\xi)/(8\xi)} \right), \right. \\ & \quad \left. e 2^D g_a \gamma |X^{(v|t)}| D! (f_{R_0}\xi + \xi + v|t|)^D \right] \\ & \leq \min \left[2^{6D+\alpha} e(2D)! g_{a,0} \gamma^2 |X^{(v|t)}| \cdot |Y^{(v|t)}| (f_{R_0}\xi)^{2D} \left(R^{-\alpha} + e^{-(R-4f_{R_0}\xi)/(8\xi)} \right), e 6^D g_a \gamma |X^{(v|t)}| D! (f_{R_0}\xi)^D \right], \end{aligned} \quad (\text{S.406})$$

where we use $f_{R_0}\xi \geq v|t| \geq \xi$ from the definition (S.401) and the assumption $v|t| \geq \xi$. By applying the inequality (S.406) to (S.405), we arrive at the inequality of

$$\begin{aligned} & \sum_{s=0}^{\infty} e^{-s} \sum_{r=1}^{R_0} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[(s+f_r)\xi] \neq \emptyset, Z_r \cap Y[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\| \\ & \leq \min \left[J_1 g_{a,0} |X^{(v|t)}| \cdot |Y^{(v|t)}| (f_{R_0}\xi)^{2D} \left(R^{-\alpha} + e^{-(R-4f_{R_0}\xi)/(8\xi)} \right), J_2 g_a |X^{(v|t)}| (f_{R_0}\xi)^D \right] \end{aligned} \quad (\text{S.407})$$

with

$$J_1 := 2^{6D+\alpha} e(2D)! \gamma^2, \quad J_2 := e 6^D \gamma D!. \quad (\text{S.408})$$

We next consider the summation for $r > R_0$. We here use the assumption (S.7), which yields

$$\begin{aligned} & \sum_{r=R_0+1}^{\infty} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[(s+f_r)\xi] \neq \emptyset, Z_r \cap Y[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\| \leq \sum_{r=R_0+1}^{\infty} \sum_{\substack{Z \subseteq \Lambda, \text{diam}(Z) \geq r \\ Z \cap X[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\| \\ & \leq \sum_{r=R_0+1}^{\infty} \frac{g_a |X[(s+f_r)\xi]|}{r^{\alpha-D}} \\ & \leq \sum_{r=R_0+1}^{\infty} \frac{g_a \gamma |X^{(v|t)}| [2(s\xi + f_r\xi + v|t|)]^D}{r^{\alpha-D}} \\ & \leq g_a \gamma |X^{(v|t)}| (2\xi)^D \sum_{r=R_0+1}^{\infty} \frac{(s + f_r + v|t|/\xi)^D}{r^{\alpha-D}}, \end{aligned} \quad (\text{S.409})$$

where we use (S.387) in the third inequality. We thus obtain

$$\begin{aligned} & \sum_{s=0}^{\infty} e^{-s} \sum_{r=R_0+1}^{\infty} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[(s+f_r)\xi] \neq \emptyset, Z_r \cap Y[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\| \leq g_a \gamma |X^{(v|t)}| (2\xi)^D \sum_{r=R_0+1}^{\infty} \sum_{s=0}^{\infty} e^{-s} \frac{(s + f_r + v|t|/\xi)^D}{r^{\alpha-D}} \\ & \leq g_a \gamma |X^{(v|t)}| (2\xi)^D e D! \sum_{r=R_0+1}^{\infty} \frac{(1 + f_r + v|t|/\xi)^D}{r^{\alpha-D}}, \end{aligned} \quad (\text{S.410})$$

where we use the inequality (S.101) to obtain the following upper bound:

$$\sum_{s=0}^{\infty} e^{-s} (s + f_r + v|t|/\xi)^D \leq \int_0^{\infty} e^{-x+1} (x + f_r + v|t|/\xi)^D dx \leq e D! (1 + f_r + v|t|/\xi)^D. \quad (\text{S.411})$$

From the definition of f_r in Eq. (S.401), we have $f_r = f_1 + 2D \log(r)$, and hence

$$\begin{aligned}
\sum_{r=R_0+1}^{\infty} \frac{(1 + f_r + v|t|/\xi)^D}{r^{\alpha-D}} &= \sum_{r=R_0+1}^{\infty} \frac{[1 + f_1 + v|t|/\xi + 2D \log(r)]^D}{r^{\alpha-D}} \\
&\leq (1 + f_1 + v|t|/\xi)^D \sum_{r=R_0+1}^{\infty} \frac{[1 + 2D \log(r)]^D}{r^{\alpha-D}} \\
&\leq (1 + f_1 + v|t|/\xi)^D \sup_{r \geq 1} \left(\frac{[1 + 2D \log(r)]^D}{r^{\alpha-2D-1}} \right) \sum_{r=R_0+1}^{\infty} \frac{1}{r^{D+1}} \\
&\leq \frac{(1 + f_1 + v|t|/\xi)^D / D}{R_0^D} \sup_{r \geq 1} \left(\frac{[1 + 2D \log(r)]^D}{r^{\alpha-2D-1}} \right) =: c_{\alpha,D} \frac{(1 + f_1 + v|t|/\xi)^D}{R_0^D}, \quad (\text{S.412})
\end{aligned}$$

where we define $c_{\alpha,D} := D^{-1} \sup_{r \geq 1} ([1 + 2D \log(r)]^D / r^{\alpha-2D-1})$. Note that from $\alpha > 2D + 1$ the constant $c_{\alpha,D}$ has a finite value. By applying the above inequality to (S.410), we arrive at the following upper bound:

$$\begin{aligned}
\sum_{s=0}^{\infty} e^{-s} \sum_{r=R_0+1}^{\infty} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[(s+f_r)\xi] \neq \emptyset, Z_r \cap Y[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\| &\leq g_a \gamma |X^{(v|t)}| (2\xi)^D e D! c_{\alpha,D} \frac{(1 + f_1 + v|t|/\xi)^D}{R_0^D} \\
&\leq g_a |X^{(v|t)}| e \gamma D! [8 \log(2e\gamma^2 \mathcal{C})]^D c_{\alpha,D} \frac{(v|t|)^D}{R_0^D} = g_a J_3 |X^{(v|t)}| (v|t|)^D R_0^{-D}, \quad (\text{S.413})
\end{aligned}$$

where we use $2\xi(1 + f_1 + v|t|/\xi) = 2\xi + 2\xi \log(2e\gamma^2 \mathcal{C}) + 4v|t| \leq 8v|t| \log(2e\gamma^2 \mathcal{C})$ from the assumption of $v|t| \geq \xi$. Note that J_3 is defined as

$$J_3 = e \gamma D! [8 \log(2e\gamma^2 \mathcal{C})]^D c_{\alpha,D}. \quad (\text{S.414})$$

Finally, we choose R_0 as $R_0 = \lceil R^{\alpha/D} \rceil \leq 2R^{\alpha/D}$. The upper bounds in (S.407) and (S.413) reduce to

$$\begin{aligned}
\sum_{s=0}^{\infty} e^{-s} \sum_{r=1}^{R_0} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[(s+f_r)\xi] \neq \emptyset, Z_r \cap Y[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\| \\
\leq \min \left[J_1 g_{a,0} |X^{(v|t)}| \cdot |Y^{(v|t)}| (\tilde{f}_R \xi)^{2D} \left(R^{-\alpha} + e^{-(R-4\tilde{f}_R \xi)/(8\xi)} \right), J_2 g_a |X^{(v|t)}| (\tilde{f}_R \xi)^D \right] \quad (\text{S.415})
\end{aligned}$$

and

$$\sum_{s=0}^{\infty} e^{-s} \sum_{r=R_0+1}^{\infty} \sum_{\substack{Z_r \subseteq \Lambda, \text{diam}(Z_r)=r \\ Z_r \cap X[(s+f_r)\xi] \neq \emptyset, Z_r \cap Y[(s+f_r)\xi] \neq \emptyset}} \|a_{Z_r}\| \leq g_a J_3 |X^{(v|t)}| (v|t|)^D R^{-\alpha}, \quad (\text{S.416})$$

respectively, where in the first inequality we use $f_{R_0} \leq f_{2R^{\alpha/D}} =: \tilde{f}_R$. By combining the above two bounds with (S.403), we obtain the main inequality (S.369). This completes the proof of Proposition 9. \square

D. Proof of Proposition 11

1. Statement

Proposition 11. *Let us consider a Hamiltonian \tilde{H} as follows:*

$$\tilde{H} = \sum_{Z \subseteq \Lambda} \tilde{h}_Z \quad (\text{S.417})$$

with

$$\begin{aligned}
\sum_{\substack{Z \subseteq \Lambda \\ Z \cap X \neq \emptyset, Z \cap Y \neq \emptyset}} \|\tilde{h}_Z\| &\leq |X^{(\xi_0)}| \cdot |Y^{(\xi_0)}| g(R), \quad R = d_{X,Y}, \\
g(R) &= \begin{cases} \tilde{g} & \text{for } R < \kappa \xi_0, \\ \tilde{g}_R R^{-\alpha} & \text{for } R \geq \kappa \xi_0, \end{cases} \quad \tilde{g}_R := \tilde{g}_0 \log^p(R+1) \quad (p \in \mathbb{N}), \quad (\text{S.418})
\end{aligned}$$

where $\kappa \geq 2$, $\xi_0 \geq 1$, and $g(R)$ monotonically decreases with R . Then, the Hamiltonian \tilde{H} satisfies $\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y})$ -Lieb-Robinson bound with

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = |\mathcal{X}^{(\xi_0)}| \cdot |\mathcal{Y}^{(\xi_0)}| \frac{2^{\alpha+1} e \Gamma(\alpha+2)}{\tilde{\lambda}} \left(e^{2e\tilde{\lambda}\gamma^2 9^D |t|} - 1 \right) g(R) + 2 |X^{(\xi_0)}| \left(\frac{4e\kappa\xi_0\tilde{\lambda}\gamma^2 9^D |t|}{R} \right)^{R/(2\kappa\xi_0)}, \quad (\text{S.419})$$

where $\Gamma(x)$ is the gamma function and we define $\tilde{\lambda}$ as

$$\begin{aligned} \tilde{\lambda} &:= \tilde{g}\gamma[2(\kappa+3)]^D + \frac{c_{p,\kappa}\tilde{g}_0\gamma D 2^{D+1} \log^p(\kappa\xi_0+1)}{\kappa+2} \frac{1}{\xi_0^\alpha}, \\ c_{p,\kappa} &:= \sup_{z \in \mathbb{R} | z \geq \kappa+2} \left[\frac{(\log(z)+1)^p}{(z-2)^{\alpha-D-1}} \right]. \end{aligned} \quad (\text{S.420})$$

Remark. The theorem is similar to the ones in [15, 17]. Indeed, we will utilize their proof technique, but the theorems therein cannot be directly applied. The main reason is the following. The theorems in Ref. [15, 17] roughly give the following bound:

$$\mathcal{G}(x, t, \mathcal{X}, \mathcal{Y}) = |\mathcal{X}| \cdot |\mathcal{Y}| \left(e^{\text{const.} \lambda |t|} - 1 \right) g(R), \quad (\text{S.421})$$

where the parameter λ is defined by the inequality of

$$\max_{x,y \in \Lambda} \sum_{z \in \Lambda} g(d_{x,z}) g(d_{z,y}) \leq \lambda g(d_{x,y}). \quad (\text{S.422})$$

In our case, from the definition (S.418), the parameter λ is roughly upper-bounded by $\tilde{g}\xi_0^D$, and hence the exponential term is given by $e^{\mathcal{O}(\tilde{g}\xi_0^D |t|)}$. On the other hand, the inequality (S.420) gives it by

$$e^{\mathcal{O}(\tilde{g}|t|) + \mathcal{O}(\tilde{g}_0\xi_0^{-\alpha}|t|)},$$

which is much better than $e^{\mathcal{O}(\tilde{g}\xi_0^D |t|)}$ when \tilde{g} , \tilde{g}_0 and ξ_0 depend on the time as in Eq. (S.356). This improvement is crucial in proving the linear light cone for $\alpha > 2D + 1$.

2. Proof of Proposition 11.

We denote $\Lambda^{(\xi_0)} = \{i_s\}_{s=1}^{\tilde{n}}$ with $\tilde{n} = |\Lambda^{(\xi_0)}|$. We also define $\{L_s\}_{s=1}^{\tilde{n}}$ as

$$L_s := i_s[\xi_0] \quad \text{for } i_s \in \Lambda^{(\xi_0)}. \quad (\text{S.423})$$

Note that $L_1 \cup L_2 \cdots L_{\tilde{n}} = \Lambda$ and

$$\text{diam}(L_s) = \text{diam}(i_s[\xi_0]) \leq 2\xi_0 + 1. \quad (\text{S.424})$$

The essence of the proof is similar to the ones in [15, 17]. The key idea to improve the original results is to treat the coarse grained set $\Lambda^{(\xi_0)}$ instead of the original total set Λ . For example, this point is reflected in the inequality (S.437). In order to simplify the inequality to a convenient form, we will utilize the similar inequality (S.441) to (S.422). The parameter $\tilde{\lambda}$ defined in Eq. (S.420) will appear in order to upper-bound the summation (S.442).

We start from the same inequality as (S.256):

$$\|[O_X(\tilde{H}, t), O_Y]\| \leq \sum_{m=1}^{m^*} \mathcal{L}_m \quad (\text{S.425})$$

with $\|O_X\| = \|O_Y\| = 1$ and $R = d_{X,Y}$, where $\{\mathcal{L}_m\}_{m=1}^{m^*-1}$ and \mathcal{L}_{m^*} are defined as

$$\mathcal{L}_m = \frac{2(2|t|)^m}{m!} \sum_{Z_1 \cap X \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset} \|\tilde{h}_{Z_2}\| \cdots \sum_{\substack{Z_m \cap Z_{m-1} \neq \emptyset \\ Z_m \cap Y \neq \emptyset}} \|\tilde{h}_{Z_m}\| \quad (\text{S.426})$$

and

$$\mathcal{L}_{m^*} = \frac{2(2|t|)^{m^*}}{m^*!} \sum_{Z_1 \cap X \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset} \|\tilde{h}_{Z_2}\| \cdots \sum_{Z_{m^*} \cap Z_{m^*-1} \neq \emptyset} \|\tilde{h}_{Z_{m^*}}\|. \quad (\text{S.427})$$

In the proof, we choose m^* such that for $m \leq m^* - 1$ the following inequalities are satisfied:

$$\frac{R - 2(m+1)\xi_0}{m} \geq \frac{R}{2m} \quad \text{and} \quad \frac{R}{2m} \geq \kappa\xi_0. \quad (\text{S.428})$$

We note that the above inequalities hold for $m \leq R/(4\xi_0) - 1$ and $m \leq R/(2\kappa\xi_0)$, respectively. Because of $\kappa \geq 2$, we need to choose so that the parameter m may satisfy $m \leq R/(2\kappa\xi_0)$, and hence we adopt

$$m^* = \left\lfloor \frac{R}{2\kappa\xi_0} \right\rfloor + 1 \geq \frac{R}{2\kappa\xi_0}. \quad (\text{S.429})$$

We first consider \mathcal{L}_1 :

$$\mathcal{L}_1 = 2(2|t|) \sum_{Z_1 \cap X \neq \emptyset, Z_1 \cap Y \neq \emptyset} \|\tilde{h}_{Z_1}\| \leq 2(2|t|) |X^{(\xi_0)}| \cdot |Y^{(\xi_0)}| g(R), \quad (\text{S.430})$$

where the inequality is given by the definition of the interaction (S.418). We second consider \mathcal{L}_2 :

$$\mathcal{L}_2 = \frac{2(2|t|)^2}{2!} \sum_{Z_1 \cap X \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset, Z_2 \cap Y \neq \emptyset} \|\tilde{h}_{Z_2}\|. \quad (\text{S.431})$$

From $X^{(\xi_0)}[\xi_0] \supseteq X$, we obtain

$$\sum_{Z_1 \cap X \neq \emptyset} \|\tilde{h}_{Z_1}\| \leq \sum_{Z_1 \cap X^{(\xi_0)}[\xi_0] \neq \emptyset} \|\tilde{h}_{Z_1}\| \leq \sum_{i_s \in X^{(\xi_0)}} \sum_{Z_1 \cap i_s[\xi_0] \neq \emptyset} \|\tilde{h}_{Z_1}\| = \sum_{i_s \in X^{(\xi_0)}} \sum_{Z_1 \cap L_s \neq \emptyset} \|\tilde{h}_{Z_1}\|, \quad (\text{S.432})$$

where we use the notation of L_s in (S.423). By using the above inequality, we have

$$\sum_{Z_1 \cap X \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset, Z_2 \cap Y \neq \emptyset} \|\tilde{h}_{Z_2}\| \leq \sum_{i_s \in X^{(\xi_0)}} \sum_{i_{s'} \in Y^{(\xi_0)}} \sum_{Z_1 \cap L_s \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset, Z_2 \cap L_{s'} \neq \emptyset} \|\tilde{h}_{Z_2}\|. \quad (\text{S.433})$$

Furthermore, from $L_1 \cup L_2 \cup \dots \cup L_{\tilde{n}} = \Lambda$, we obtain

$$\sum_{Z_1 \cap L_s \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset, Z_2 \cap L_{s'} \neq \emptyset} \|\tilde{h}_{Z_2}\| \leq \sum_{s_1=1}^{\tilde{n}} \sum_{Z_1 \cap L_{s_1} \neq \emptyset, Z_1 \cap L_{s_1} \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap L_{s_1} \neq \emptyset, Z_2 \cap L_{s_1} \neq \emptyset} \|\tilde{h}_{Z_2}\|. \quad (\text{S.434})$$

The definition (S.418) gives the upper bound of

$$\sum_{Z_1 \cap L_s \neq \emptyset, Z_1 \cap L_{s_1} \neq \emptyset} \|\tilde{h}_{Z_1}\| \leq |L_s^{(\xi_0)}| \cdot |L_{s_1}^{(\xi_0)}| g(d_{L_s, L_{s_1}}) \leq \gamma^2 9^D g(d_{L_s, L_{s_1}}), \quad (\text{S.435})$$

where we use the inequality (S.18) to obtain $|L_s^{(\xi_0)}| \leq \gamma[\text{diam}(L_s)/\xi_0]^D \leq \gamma[(2\xi_0 + 1)/\xi_0]^D \leq \gamma 3^D$ ($\xi_0 \geq 1$). This upper bound reduces the inequality (S.434) to

$$\sum_{Z_1 \cap L_s \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset, Z_2 \cap L_{s'} \neq \emptyset} \|\tilde{h}_{Z_2}\| \leq (\gamma^2 9^D)^2 \sum_{s_1=1}^{\tilde{n}} g(d_{L_s, L_{s_1}}) g(d_{L_{s_1}, L_{s'}}). \quad (\text{S.436})$$

By combining (S.433) and (S.436), we have

$$\sum_{Z_1 \cap X \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset, Z_2 \cap Y \neq \emptyset} \|\tilde{h}_{Z_2}\| \leq (\gamma^2 9^D)^2 \sum_{i_s \in X^{(\xi_0)}} \sum_{i_{s'} \in Y^{(\xi_0)}} \sum_{s_1=1}^{\tilde{n}} g(d_{L_s, L_{s_1}}) g(d_{L_{s_1}, L_{s'}}). \quad (\text{S.437})$$

Because of $i_s \in X^{(\xi_0)}$, $i_{s'} \in Y^{(\xi_0)}$, we have $L_s \cap X \neq \emptyset$ and $L_{s'} \cap Y \neq \emptyset$. Then, from $R = d_{X,Y}$, the distances $d_{L_s, L_{s_1}}$ and $d_{L_{s_1}, L_{s'}}$ should satisfy

$$R \leq d_{L_s, L_{s_1}} + d_{L_{s_1}, L_{s'}} + \text{diam}(L_s) - 1 + \text{diam}(L_{s'}) - 1 + \text{diam}(L_{s_1}) - 1 \leq d_{L_s, L_{s_1}} + d_{L_{s_1}, L_{s'}} + 6\xi_0, \quad (\text{S.438})$$

where we use the inequality (S.424) for $\text{diam}(L_s)$. The above inequality yields

$$\max(d_{L_s, L_{s_1}}, d_{L_{s_1}, L_{s'}}) \geq \frac{R - 6\xi_0}{2} \geq \frac{R}{4} \geq \kappa\xi_0, \quad (\text{S.439})$$

where the second and third inequalities are derived from the condition (S.428) with $m = 2$. From the above inequality, the definition of $g(R)$ in (S.418) implies

$$\min[g(d_{L_s, L_{s_1}}), g(d_{L_{s_1}, L_{s'}})] \leq g(R/4) = \frac{4^\alpha}{\tilde{g}_R/\tilde{g}_{R/4}} g(R) \leq 4^\alpha g(R), \quad (\text{S.440})$$

where the last inequality is given by $\tilde{g}_R \geq \tilde{g}_{R/4}$. Hence, we obtain

$$\sum_{s_1=1}^{\tilde{n}} g(d_{L_s, L_{s_1}}) g(d_{L_{s_1}, L_{s'}}) \leq \sum_{s_1=1}^{\tilde{n}} \min[g(d_{L_s, L_{s_1}}), g(d_{L_{s_1}, L_{s'}})] \cdot [g(d_{L_s, L_{s_1}}) + g(d_{L_{s_1}, L_{s'}})] \leq 2\tilde{\lambda} \cdot 4^\alpha g(R), \quad (\text{S.441})$$

where in the last inequality we use

$$\max_{i_s \in \Lambda^{(\xi_0)}} \sum_{i_{s'} \in \Lambda^{(\xi_0)}} g(d_{L_{s'}, L_s}) \leq \tilde{\lambda} \quad (\text{S.442})$$

with $\tilde{\lambda}$ defined in Eq. (S.420). The proof of the inequality (S.442) is given in Sec. XIV D 3. We thus upper-bound Eq. (S.431) by

$$\mathcal{L}_2 \leq \frac{2(2\gamma^2 9^D |t|)^2}{2!} |X^{(\xi_0)}| \cdot |Y^{(\xi_0)}| 2\tilde{\lambda} \cdot 4^\alpha g(R). \quad (\text{S.443})$$

In the same way, we consider the third term \mathcal{L}_3 :

$$\mathcal{L}_3 = \frac{2(2|t|)^3}{3!} \sum_{Z_1 \cap X \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset} \|\tilde{h}_{Z_2}\| \sum_{Z_3 \cap Z_2 \neq \emptyset, Z_3 \cap Y \neq \emptyset} \|\tilde{h}_{Z_3}\|. \quad (\text{S.444})$$

In order to estimate the summation, we first consider

$$\begin{aligned} & \sum_{Z_1 \cap X \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset} \|\tilde{h}_{Z_2}\| \sum_{Z_3 \cap Z_2 \neq \emptyset, Z_3 \cap Y \neq \emptyset} \|\tilde{h}_{Z_3}\| \\ & \leq \sum_{i_s \in X^{(\xi_0)}} \sum_{i_{s'} \in Y^{(\xi_0)}} \sum_{Z_1 \cap L_s \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset} \|\tilde{h}_{Z_2}\| \sum_{Z_3 \cap Z_2 \neq \emptyset, Z_3 \cap L_{s'} \neq \emptyset} \|\tilde{h}_{Z_3}\| \\ & \leq \sum_{i_s \in X^{(\xi_0)}} \sum_{i_{s'} \in Y^{(\xi_0)}} \sum_{s_1=1}^{\tilde{n}} \sum_{Z_1 \cap L_s \neq \emptyset, Z_1 \cap L_{s_1} \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap L_{s_1} \neq \emptyset} \|\tilde{h}_{Z_2}\| \sum_{Z_3 \cap Z_2 \neq \emptyset, Z_3 \cap L_{s'} \neq \emptyset} \|\tilde{h}_{Z_3}\| \\ & \leq \sum_{i_s \in X^{(\xi_0)}} \sum_{i_{s'} \in Y^{(\xi_0)}} \sum_{s_1=1}^{\tilde{n}} \sum_{Z_1 \cap L_s \neq \emptyset, Z_1 \cap L_{s_1} \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{s_2=1}^{\tilde{n}} \sum_{Z_2 \cap L_{s_1} \neq \emptyset, Z_2 \cap L_{s_2} \neq \emptyset} \|\tilde{h}_{Z_2}\| \sum_{Z_3 \cap L_{s_2} \neq \emptyset, Z_3 \cap L_{s'} \neq \emptyset} \|\tilde{h}_{Z_3}\| \\ & \leq (\gamma^2 9^D)^3 \sum_{i_s \in X^{(\xi_0)}} \sum_{i_{s'} \in Y^{(\xi_0)}} \sum_{s_1=1}^{\tilde{n}} \sum_{s_2=1}^{\tilde{n}} g(d_{L_s, L_{s_1}}) g(d_{L_{s_1}, L_{s_2}}) g(d_{L_{s_2}, L_{s'}}). \end{aligned} \quad (\text{S.445})$$

In the similar manner to the inequality (S.438), we obtain

$$\begin{aligned} R & \leq d_{L_s, L_{s_1}} + d_{L_{s_1}, L_{s_2}} + d_{L_{s_2}, L_{s'}} + \text{diam}(L_s) - 1 + \text{diam}(L_{s'}) - 1 + \text{diam}(L_{s_1}) - 1 + \text{diam}(L_{s_2}) - 1 \\ & \leq d_{L_s, L_{s_1}} + d_{L_{s_1}, L_{s_2}} + d_{L_{s_2}, L_{s'}} + 8\xi_0, \end{aligned} \quad (\text{S.446})$$

which yields

$$\max(d_{L_s, L_{s_1}}, d_{L_{s_1}, L_{s_2}}, d_{L_{s_2}, L_{s'}}) \geq \frac{R - 8\xi_0}{3} \geq \frac{R}{6} \geq \kappa\xi_0 \quad (\text{S.447})$$

from the condition (S.428) with $m = 3$. By combining the above inequality and the definition of $g(R)$ in (S.418), we also obtain

$$\min[g(d_{L_s, L_{s_1}}), g(d_{L_{s_1}, L_{s_2}}), g(d_{L_{s_2}, L_{s'}})] \leq g(R/6) = \frac{6^\alpha}{\tilde{g}_R/\tilde{g}_{R/6}} g(R) \leq 6^\alpha g(R), \quad (\text{S.448})$$

which yields

$$\sum_{s_1=1}^{\tilde{n}} \sum_{s_2=1}^{\tilde{n}} g(d_{L_s, L_{s_1}}) g(d_{L_{s_1}, L_{s_2}}) g(d_{L_{s_2}, L_{s'}}) \leq 3\tilde{\lambda}^2 \cdot 6^\alpha g(R), \quad (\text{S.449})$$

where we use the inequality (S.442). We thus obtain

$$\mathcal{L}_3 \leq \frac{2(2\gamma^2 9^D |t|)^3}{3!} |X^{(\xi_0)}| \cdot |Y^{(\xi_0)}| 3\tilde{\lambda}^2 \cdot 6^\alpha g(R). \quad (\text{S.450})$$

By repeating this process, we obtain

$$\mathcal{L}_m \leq \frac{2(2\gamma^2 9^D |t|)^m}{m!} |X^{(\xi_0)}| \cdot |Y^{(\xi_0)}| m \tilde{\lambda}^{m-1} \cdot (2m)^\alpha g(R) \quad (\text{S.451})$$

for $m \leq m^* - 1$. From the above upper bounds for $\{\mathcal{L}_m\}_{m=1}^{m^*-1}$, we have

$$\begin{aligned} \sum_{m=1}^{m^*-1} \mathcal{L}_m &\leq \frac{2^{\alpha+1}}{\tilde{\lambda}} |X^{(\xi_0)}| \cdot |Y^{(\xi_0)}| g(R) \sum_{m=1}^{\infty} \frac{(2\tilde{\lambda}\gamma^2 9^D |t|)^m}{m!} m^{\alpha+1} \\ &\leq \frac{2^{\alpha+1} e \Gamma(\alpha+2)}{\tilde{\lambda}} |X^{(\xi_0)}| \cdot |Y^{(\xi_0)}| \left(e^{2e\tilde{\lambda}\gamma^2 9^D |t|} - 1 \right) g(R), \end{aligned} \quad (\text{S.452})$$

where the upper bound of the summation is derived as follows:

$$\begin{aligned} \sum_{m=1}^{\infty} \frac{(2\tilde{\lambda}\gamma^2 9^D |t|)^m}{m!} m^{\alpha+1} &\leq \sum_{m=1}^{\infty} \frac{(2\tilde{\lambda}\gamma^2 9^D |t|)^m}{m!} e^m \sum_{\tilde{m}=1}^{\infty} e^{-\tilde{m}} \tilde{m}^{\alpha+1} \\ &\leq \left(e^{2e\tilde{\lambda}\gamma^2 9^D |t|} - 1 \right) \int_0^{\infty} (x+1)^{\alpha+1} e^{-x} \leq e \Gamma(\alpha+2) \left(e^{2e\tilde{\lambda}\gamma^2 9^D |t|} - 1 \right). \end{aligned} \quad (\text{S.453})$$

Note that $\Gamma(x)$ is the gamma function.

Finally, for $m = m^*$, by regarding as $Y = \Lambda$, we can apply the same analyses as in the case of $m \leq m^* - 1$. We obtain

$$\begin{aligned} &\sum_{Z_1 \cap X \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap Z_1 \neq \emptyset} \|\tilde{h}_{Z_2}\| \cdots \sum_{Z_{m^*} \cap Z_{m^*-1} \neq \emptyset} \|\tilde{h}_{Z_{m^*}}\| \\ &\leq \sum_{i_s \in X^{(\xi_0)}} \sum_{i_{s'} \in \Lambda^{(\xi_0)}} \sum_{s_1, s_2, \dots, s_{m^*-1}=1}^{\tilde{n}} \sum_{Z_1 \cap L_{s_1} \neq \emptyset, Z_1 \cap L_{s_1} \neq \emptyset} \|\tilde{h}_{Z_1}\| \sum_{Z_2 \cap L_{s_1} \neq \emptyset, Z_2 \cap L_{s_2} \neq \emptyset} \|\tilde{h}_{Z_2}\| \sum_{Z_3 \cap L_{s_2} \neq \emptyset, Z_3 \cap L_{s'} \neq \emptyset} \|\tilde{h}_{Z_3}\| \\ &\quad \cdots \sum_{Z_{m^*-1} \cap L_{s_{m^*-2}} \neq \emptyset, Z_{m^*-1} \cap L_{m^*-1} \neq \emptyset} \|\tilde{h}_{Z_{m^*-1}}\| \sum_{Z_{m^*} \cap L_{s_{m^*-1}} \neq \emptyset, Z_{m^*} \cap L_{s'} \neq \emptyset} \|\tilde{h}_{Z_{m^*}}\| \\ &\leq (\gamma^2 9^D)^{m^*} \sum_{i_s \in X^{(\xi_0)}} \sum_{s', s_1, s_2, \dots, s_{m^*-1}=1}^{\tilde{n}} g(d_{L_s, L_{s_1}}) g(d_{L_{s_1}, L_{s_2}}) g(d_{L_{s_2}, L_{s_3}}) \cdots g(d_{L_{s_{m^*-2}}, L_{s_{m^*-1}}}) g(d_{L_{s_{m^*-1}}, L_{s'}}) \\ &\leq (\gamma^2 9^D)^{m^*} |X^{(\xi_0)}| \tilde{\lambda}^{m^*}. \end{aligned} \quad (\text{S.454})$$

From the definition of \mathcal{L}_{m^*} as in (S.427), the above inequality yields

$$\mathcal{L}_{m^*} = 2 |X^{(\xi_0)}| \frac{(2\tilde{\lambda}\gamma^2 9^D |t|)^{m^*}}{m^*!} \leq 2 |X^{(\xi_0)}| \left(\frac{2e\tilde{\lambda}\gamma^2 9^D |t|}{m^*} \right)^{m^*}, \quad (\text{S.455})$$

where we use $m^*! \geq (m^*/e)^{m^*}$.

By applying the inequalities (S.452) and (S.455) to (S.425) with Eq. (S.429), we obtain the main inequality (S.419). This completes the proof. \square

3. Derivation of the inequality (S.442)

We here derive the upper bound (S.442) for

$$\max_{i_s \in \Lambda^{(\xi_0)}} \sum_{i_{s'} \in \Lambda^{(\xi_0)}} g(d_{L_{s'}, L_s}). \quad (\text{S.456})$$

Note that we have defined $L_s = i_s[\xi_0]$ and $L_{s'} = i_{s'}[\xi_0]$. First, by using $d_{L_{s'}, L_s} \geq d_{i_{s'}, i_s} - 2\xi_0$ and the definition (S.418), we obtain

$$\sum_{i_{s'} \in \Lambda^{(\xi_0)}} g(d_{L_{s'}, L_s}) \leq \sum_{i_{s'} \in \Lambda^{(\xi_0)}} g(d_{i_{s'}, i_s} - 2\xi_0) \leq \sum_{\substack{i_{s'} \in \Lambda^{(\xi_0)} \\ d_{i_{s'}, i_s} < (\kappa+2)\xi_0}} g(d_{i_{s'}, i_s} - 2\xi_0) + \sum_{\substack{i_{s'} \in \Lambda^{(\xi_0)} \\ d_{i_{s'}, i_s} \geq (\kappa+2)\xi_0}} g(d_{i_{s'}, i_s} - 2\xi_0), \quad (\text{S.457})$$

where in the first inequality we use the condition that $g(R)$ monotonically decreases with R . The first term is bounded from above by

$$\sum_{\substack{i_{s'} \in \Lambda^{(\xi_0)} \\ d_{i_{s'}, i_s} < (\kappa+2)\xi_0}} g(d_{i_{s'}, i_s} - 2\xi_0) \leq \sum_{i \in (i_s[(\kappa+2)\xi_0])^{(\xi_0)}} \tilde{g} \leq \tilde{g} |(i_s[(\kappa+2)\xi_0])^{(\xi_0)}| \leq \tilde{g} \gamma [2(\kappa+3)]^D, \quad (\text{S.458})$$

where the last inequality is derived by using (S.16) as follows:

$$|(i_s[(\kappa+2)\xi_0])^{(\xi_0)}| \leq \gamma \left[\frac{\text{diam}(i_s[(\kappa+2)\xi_0])}{\xi_0} \right]^D \leq \gamma \left[\frac{2(\kappa+2)\xi_0 + 1}{\xi_0} \right]^D \leq \gamma [2(\kappa+3)]^D. \quad (\text{S.459})$$

The second term in (S.457) is bounded by using Lemma 2. By choosing

$$f(z) = g(z - 2\xi_0), \quad x_0 = (\kappa+2)\xi_0, \quad (\text{S.460})$$

we obtain from Ineq. (S.109)

$$\sum_{\substack{i_{s'} \in \Lambda^{(\xi_0)} \\ d_{i_{s'}, i_s} \geq (\kappa+2)\xi_0}} g(d_{i_{s'}, i_s} - 2\xi_0) \leq 2^{D+1} C_{g, (\kappa+2)\xi_0} \gamma D \xi_0^{-D} [(\kappa+2)\xi_0]^D g(\kappa\xi_0), \quad (\text{S.461})$$

where $C_{g, (\kappa+2)\xi_0}$ is defined by Eq. (S.108):

$$C_{g, (\kappa+2)\xi_0} = \sup_{z \in \mathbb{R} | z \geq (\kappa+2)\xi_0} \left(\frac{z^{D+1} g(z - 2\xi_0)}{[(\kappa+2)\xi_0]^{D+1} g(\kappa\xi_0)} \right). \quad (\text{S.462})$$

From the definition (S.418), for $z \geq (\kappa+2)\xi_0$ we have $g(z - 2\xi_0) = \tilde{g}_0 \log^p(z - 2\xi_0 + 1)(z - 2\xi_0)^{-\alpha}$ and hence

$$\begin{aligned} C_{g, (\kappa+2)\xi_0} &= \frac{\kappa^\alpha}{(\kappa+2)^{D+1}} \sup_{z \in \mathbb{R} | z \geq (\kappa+2)\xi_0} \left[(z/\xi_0)^{D+1} [(z - 2\xi_0)/\xi_0]^{-\alpha} \frac{\log^p(z - 2\xi_0 + 1)}{\log^p(\kappa\xi_0 + 1)} \right] \\ &\leq \frac{\kappa^\alpha}{(\kappa+2)^{D+1}} \sup_{z \in \mathbb{R} | z \geq \kappa+2} \left[\frac{(\log(z) + 1)^p}{(z - 2)^{\alpha-D-1}} \right] =: \frac{\kappa^\alpha}{(\kappa+2)^{D+1}} c_{p, \kappa}, \end{aligned} \quad (\text{S.463})$$

where we use the upper bound of

$$\frac{\log(z - 2\xi_0 + 1)}{\log(\kappa\xi_0 + 1)} \leq \frac{\log(z/\xi_0) + \log(\xi_0)}{\log(2\xi_0 + 1)} \leq \frac{\log(z/\xi_0)}{\log(3)} + 1 \leq \log(z/\xi_0) + 1. \quad (\text{S.464})$$

Note that $\kappa \geq 2$ and $\xi_0 \geq 1$. By combining the inequalities (S.461) and (S.463), we have

$$\begin{aligned} \sum_{\substack{i_{s'} \in \Lambda^{(\xi_0)} \\ d_{i_{s'}, i_s} \geq (\kappa+2)\xi_0}} g(d_{i_{s'}, i_s} - 2\xi_0) &\leq 2^{D+1} \gamma D \xi_0^{-D} [(\kappa+2)\xi_0]^D g(\kappa\xi_0) \cdot \frac{\kappa^\alpha}{(\kappa+2)^{D+1}} c_{p, \kappa} \\ &\leq \frac{c_{p, \kappa} \tilde{g}_0 \gamma D 2^{D+1} \log^p(\kappa\xi_0 + 1)}{\kappa + 2} \frac{\xi_0^\alpha}{\xi_0^\alpha}. \end{aligned} \quad (\text{S.465})$$

By applying the inequalities (S.458) and (S.465) to (S.457), we have

$$\sum_{i_{s'} \in \Lambda^{(\xi_0)}} g(d_{L_{s'}, L_s}) \leq \tilde{g} \gamma [2(\kappa+3)]^D + \frac{c_{p, \kappa} \tilde{g}_0 \gamma D 2^{D+1} \log^p(\kappa\xi_0 + 1)}{\kappa + 2} \frac{\xi_0^\alpha}{\xi_0^\alpha} = \tilde{\lambda}, \quad (\text{S.466})$$

where we have defined $\tilde{\lambda}$ in Eq. (S.420). We thus obtain the inequality (S.442). This completes the proof. \square