Construction of Const Dimension Codes from Serval Parallel Lift MRD Code

Xianmang He, Yindong Chen

January 20, 2022

Abstract

In this paper, we generalize the method of using two parallel versions of the lifted MRD code from the existing work [1]. The Delsarte theorem of the rank distribution of MRD codes is an important part to count codewords in our construction. We give a new generalize construction to the following bounds: if $n \geq k \geq d$, then

$$A_q(n+k,k,d) \ge q^{n(k-\frac{d}{2}+1)} + \sum_{r=\frac{d}{2}}^{k-\frac{d}{2}} A_r(Q_q(n,k,\frac{d}{2})).$$

On this basis, we also give a construction of constant-dimension subspace codes from several parallel versions of lifted MRD codes. This construction contributes to a new lower bounds for $A_q((s+1)k+n, d, k)$.

keyword: constant dimension code, subspace code, maximum rank distance code, lifted maximum rank distance code

1 Introduction

Let F_q be the finite field with q>1 elements and let $V\cong F_q^n$ be the set of all vectors of length n over F_q . F_q^n is a vector space with dimension n over F_q . The projective space $\mathcal{P}_q(n)$, is the set of all subspaces of F_q^n , including $\{\mathbf{0}\}$ and F_q^n . For a given integer k, $1\leq k\leq n$, let $\mathcal{G}_q(n,k)$ denote the set of all k-dimensional subspaces of F_q^n . $\mathcal{G}_q(n,k)$ is often referred to as Grassmannian. It is well known that

$$|\mathcal{G}_q(n,k)| = \begin{bmatrix} n \\ k \end{bmatrix}_q \stackrel{\text{def}}{=} \frac{(q^n - 1)(q^{n-1} - 1)\cdots(q^{n-k+1} - 1)}{(q^k - 1)(q^{k-1} - 1)\cdots(q - 1)}$$

where $\begin{bmatrix} n \\ k \end{bmatrix}_q$ is the q-ary Gaussian coefficient.

The set of all subspaces of V forms a metric space associated with the socalled subspace distance $d_S(U,W) = \dim(U+W) - \dim(U\cap W) = 2\dim(U+W) - 2k$. A $(n,M,d,k)_q$ constant dimension code (CDC) C is a subset of V of cardinality M in which for each pair of elements, the subspace distance is lower bounded by d, that is, for all $U \neq W \in C$, we have $d_S(U,W) \geq d$. The main question of subspace coding in the constant dimension case asks for the maximum cardinality M for fixed parameters q, n, d, and k of a $(n,M,d,k)_q$ code. Let $A_q(n,d,k)$ denote the maximum size of an $(n,M,d,k)_q$ code.

One way to construct good subspace codes utilizes rank-metric codes. A linear rank metric code $[k \times n, \varrho, d]_q$ is a subspace C of the vector space of $k \times n$ matrices over F_q , i.e., $F_q^{k \times n}$, for which the distance of two elements is lower bounded via the rank metric $d_r(A, B) = rk(A - B)$, i.e., $d \leq d_r(A, B)$ for all $A \neq B \in C$. For all parameters, $0 \leq k, n, d$ and q prime power, there is a linear rank metric code that attains the maximum cardinality of $\lceil q^{\max\{k,n\}(\min\{k,n\}-d+1)} \rceil$ (see $\lceil 2 \rceil$).

In this paper, we extend the method of using two parallel versions of lifted MRD codes to a general framework [1]. This method gives us a new lower bound, and leads to several new lower bounds from several parallel versions of lifted MRD codes.

2 Preliminaries

2.1 Lifted MRD code

Without loss of generality, we assume that $n \geq k$. For any given MRD code $Q_q(n,k,d)$ with the rank distance d, we have a $(n+k,q^{n(k-d+1)},2d,n)_q$ CDC consisting of $q^{n(k-d+1)}$ subspaces of dimension k in \mathbf{F}_q^{n+k} is defined as $lift(U_A) = \{rowspace[I_k,A]\}$, spanned by rows of (I_k,A) . Here I_k is the $k \times k$ identity matrix. For any A and B, the subspaces U_A and U_B spanned by rows of (I_k,A) and (I_k,B) are the same if and only if A=B. The intersection $U_A \cap U_B$ is the set $\{\alpha A : \alpha A = \alpha B, \alpha \in \mathbf{F}_q^k\}$. Thus $dim(U_A \cap U_B) \leq k - rank(A - B) \leq k - d$. The distance of this CDC is 2d. A CDC constructed as above is called a lifted MRD code C^{MRD} .

2.2 Delsarte Theorem

For $r \in \mathbf{Z}^+$, the rank distribution of a code Q in $Q_q(m, n, d)$ is defined by $A_r(Q) = |\{Q \in Q_q(m, n, d), rk(Q) = r\}|$ (see [3, 4]). We refer the following

result to Theorem 5.6 in [3] or Corollary 26 in [4].

Theorem 1. (Delsarte 1978) Assume that $Q \subseteq Q_q(m, n, d)$ $(m \ge n)$ is a MRD code with rank distance d, then its rank distribution is given by

$$A_r(Q) = \binom{n}{r}_q \sum_{i=0}^{r-d} (-1)^i q^{\binom{i}{2}} \binom{r}{i}_q (\frac{q^{m(n-d+1)}}{q^{m(n+i-r)}} - 1).$$

Here, the value of r varies from d to $min\{m,n\}(A_0(Q)=1)$, and the rank distribution of a MRD code is determined by its parameters m, n, d, q.

Example 1. Assume that m = n = 4, q = 2, d = 2, we have $|Q_2(4,4,2)| = 2^{12}$, and $A_2(Q_2(4,4,2)) = 525, A_3(Q(4,4,2)) = 2250, A_4(Q_2(4,4,2)) = 1320$. Similarly, m = 5, n = 5, d = 2, q = 2, then $A_2(Q_2(5,5,2)) = 4805, A_3(Q_2(5,5,2)) = 124930$. Moreover, $A_2(Q_2(6,5,2)) = 9765, A_3(Q_2(6,5,2)) = 566370, A_2(Q_2(7,5,2)) = 19685, A_3(Q_2(7,5,2)) = 2401570$.

2.3 Previous constructions

There is a variety of lower bounds and upper bounds for constant dimension codes. For numerical values of the known lower and upper bounds on the sizes of subspace codes we refer the reader to the online tables at http://subspacecodes.uni-bayreuth.de associated with [5].

Koetter and Kschischang [6], Etzion and Vardy [7] developed several upper bounds on $A_q(n,d,k)$. Surveys of upper bounds for constant dimension codes can be in the paper [8]. One of the Johnson type bound in [9] (see Theorem 4 in [9]) of CDC is

$$A_{q}(n, 2\delta, k) \leq \frac{\binom{n}{k - \delta + 1}_{q}}{\binom{k}{k - \delta + 1}_{q}}.$$
(1)

The iterative application of the Johnson type bound II ([10, Theorem 3], [9, Theorem 4,5]), which is a q-generalization of [11], gives the upper bound

$$A_q(n,d,k) \le \left\lfloor \frac{q^n - 1}{q^k - 1} \left\lfloor \frac{q^{n-1} - 1}{q^{k-1} - 1} \left\lfloor \dots \left\lfloor \frac{q^{n'+1} - 1}{q^{\frac{d}{2}+1} - 1} A_q(n',d;\frac{d}{2}) \right\rfloor \dots \right\rfloor \right\rfloor \right\rfloor$$
(2)

where $n' = n - k + \frac{d}{2}$. It is attained with equality at n = ak and d = 2k, i.e., and also at n = 13, k = 3, d = 4 with $A_2(13, 4; 3) = 1597245$, see [12].

While a lot of upper bounds for the maximum sizes of CDCs have been proposed in the literature, inequalities dominates most of them. [13] determined the maximum size $A_2(8,6,4)$ of a binary subspace code to be 257. Another known improvement is $A_2(6,4,3) = 77$. Some prior results about general lower bounds for $A_q(n,d,k)$ can see the references [14, 15, 16, 17]. Many CDC's from the multilevel construction based on echelon-Ferrers diagram have been given[18]. Authors of [19] improved the echelon-Ferrers construction by a technology termed as the pending dot. The pending dot was extended to pending block in the paper [14].

The improved linkage construction is one of the most powerful methods in network coding, and it produce currently the best known lower bound about 69.1% of the constant dimension code[16]. The work that is closest to ours is [1], which gives a construction of constant-dimension subspace codes from two parallel versions of lifted MRD codes. If $2t \geq n$, then $A_q(2n, 2(n-t), n) \geq q^{n(t+1)} + \sum_{r=n-t}^n A_r(Q_q(n, n, t))$.

In this paper, we extend the above construction to a general case $A_q(n+k,d,k)$, $n \ge k \ge d$.

3 Our construction

The basic idea is as follows: Define the set $W_1 = \{rowspace[A, I_k] | A \in Q_q(n, k, \frac{d}{2})\}$ be a $(n+k, q^{n(k-\frac{d}{2}+1)}, d, k)_q$ CDC in F_q^{n+k} spanned by the rows of $k \times (n+k)$ matrices (A, I_k) with $q^{n(k-\frac{d}{2}+1)}$ elements. Similar to code W_1 , we have another k dimension subspaces in $\mathbf{F}_q^{n(k-\frac{d}{2}+1)}$ spanned by the rows of $W_2 = (B, I_k)$, where B is in the MRD code $Q_q(n, k, \frac{d}{2})$, can be used to increase the size of the constructed constant-dimension subspace codes. Now the problem is how many different subspaces we can take from these two parallel versions of lifted MRD code such that the subspace distance d is preserved. The key point here is to keep the subspace distances larger than or equal to d by suitable conditions. By using this idea a new lower bound for $A_q(n+k,d,k), k \geq d$ are given with the help from the Delsarte Theorem about the rank distributions of the MRD code $Q_q(n,k,d)$.

3.1 A New Lower Bound

Our construction is

Theorem 2. If $n \ge k \ge d$, then

$$A_q(n+k,k,d) \ge q^{n(k-\frac{d}{2}+1)} + \sum_{r=\frac{d}{2}}^{k-\frac{d}{2}} A_r(Q_q(n,k,\frac{d}{2}))$$

Proof. Let $Q_1 \subset Q_q(n, k, \frac{d}{2})$ be a MRD code with rank distance $\frac{d}{2}$, and let $Q_2 \subset Q_q(n, k, \frac{d}{2})$ be a MRD code with the rank of each element in Q_2 is at most $k - \frac{d}{2}$. The code is defined by

$$\mathbf{C} = \{ (I_k | Q_{12}) : Q_{12} \in Q_1 \} \cup \{ (Q_{21} | I_k) : Q_{21} \in \mathbf{Q_2} \}.$$

From the definition of lift MRD code, the subspace distances of the two codes

$$\mathbf{W}_1 = \{ (I_k | Q_{12}) : Q_{12} \in Q_1 \}$$

and

$$\mathbf{W}_2 = \{ (Q_{21}|I_k) : Q_{22} \in Q_2 \}$$

are at least d. We only need to prove that the subspace distance of $W_1 \in \mathbf{W}_1$ and $W_2 \in \mathbf{W}_2$ is at least d. It is clear that these two codes are disjoint.

It is sufficient to prove that

$$dim(W_1 + W_2) = rank \begin{pmatrix} I_k & Q_{12} \\ Q_{21} & I_k \end{pmatrix} \ge k + \frac{d}{2}$$

We split the Q_{12} into two parts: Q_{121} and Q_{122} , where Q_{121} is a matrix with $(n-k)\times k$ and Q_{122} is a matrix with $k\times k$.

$$dim(W_1 + W_2) = rank \begin{pmatrix} I_k & Q_{121} & Q_{122} \\ Q_{21} & I_k \end{pmatrix} \ge k + \frac{d}{2}$$

The above formula can be transformed into the following by subtracting second row multiplied by Q_{122} :

$$dim(W_1 + W_2) = rank \begin{pmatrix} I_k | Q_{121} - Q_{21} \times Q_{122} & 0 \\ Q_{21} & I_k \end{pmatrix}$$

It is clear that $dim(W_1+W_2) \geq k+rank(I_k|Q_{121}-Q_{21}\times Q_{122})$. We note that the rank of Q_{21} is at most $k-\frac{d}{2}$, then $rank(I_k|Q_{121}-Q_{21}\times Q_{122}) \geq k-(k-\frac{d}{2}) \geq \frac{d}{2}$. In the end, we have $dim(W_1+W_2) \geq k+\frac{d}{2}$. Here $(I_k|Q_{121})$ is a $k\times n$ matrix concatenated from I_k and Q_{121} .

4 Parallel Construction from MRD Codes

4.1 General Construction

Our construction is as follow. For simplify, the subset of a MRD code $Q_q(n, k, \frac{d}{2})$ with the rank at most $k - \frac{d}{2}$ is denoted by $SQ_q(n, k, \frac{d}{2})$.

Theorem 3. If $n \geq k \geq d, s \geq 0$, then $A_q((s+1) \times k + n, d, k) \geq$

$$\sum_{j=0}^{s} q^{((s-j)k+n)(k-\frac{d}{2}+1)} (\sum_{r=\frac{d}{2}}^{k-\frac{d}{2}} A_r(Q_q(k,k,\frac{d}{2})))^j + (\sum_{r=\frac{d}{2}}^{k-\frac{d}{2}} A_r(Q_q(n,k,\frac{d}{2}))) \times (\sum_{r=\frac{d}{2}}^{k-\frac{d}{2}} A_r(Q_q(k,k,\frac{d}{2})))^{s-1}.$$

Proof. For the first block B_1 , we take k dimension subspaces in $F_q^{(s+1)k+n}$ spanned by rows of $(I_k, A_1^1, \ldots, A_s^1)$ where A_1^1, \ldots, A_{s-1}^1 are from the MRD code $Q_q(k, k, \frac{d}{2})$, A_s^1 takes from the MRD code $Q_q(n, k, \frac{d}{2})$. There are $q^{(sk+n)(k-\frac{d}{2}+1)}$ such subspaces.

For the second block B_2 , we take k dimension subspaces in $F_q^{(s+1)k+n}$ spanned by rows of $(A_1^2, I_k, \ldots, A_s^2)$ where A_1^2 takes from $SQ_q(k, k, \frac{d}{2})$. A_2^2, \ldots, A_{s-1}^2 are from the MRD code $Q_q(k, k, \frac{d}{2})$ and $A_s^2 \in Q_q(n, k, \frac{d}{2})$. There are $q^{((s-1)k+n)} \times |SQ_q(n, k, \frac{d}{2})| = q^{((s-1)k+n)} \times (\sum_{r=\frac{d}{2}}^{k-\frac{d}{2}} A_r(Q_q(k, k, \frac{d}{2})))$ such subspaces.

For the third block, we take k dimension subspaces in $F_q^{(s+1)k+n}$ spanned by rows of $(A_1^3, A_2^3, I_k, \ldots, A_s^3)$ where A_1^3, A_2^3 takes from $SQ_q(k, k, \frac{d}{2})$. A_2^3, \ldots, A_{s-1}^3 are from the MRD code $Q_q(k, k, \frac{d}{2})$, and A_s^3 takes from the MRD code $Q_q(n, k, \frac{d}{2})$. There are $q^{((s-2)k+n)(k-\frac{d}{2})} \times (\sum_{i=\frac{d}{2}}^{k-\frac{d}{2}+1} A_i(Q_q(k, k, \frac{d}{2})))^2$ such subspaces. We can continue this process. It is obvious that all these subspaces in $\mathbf{F}_q^{(s+1)k+n}$ are different.

These blocks can be demonstrated as follows:

$$\begin{pmatrix} B_1 = (I_k & Q_q(k,k,\frac{d}{2}) & Q_q(k,k,\frac{d}{2}) & \cdots & Q_q(n,k,\frac{d}{2})) \\ B_2 = (SQ_q(k,k,\frac{d}{2}) & I_k & Q_q(k,k,\frac{d}{2}) & \cdots & Q_q(n,k,\frac{d}{2})) \\ B_3 = (SQ_q(k,k,\frac{d}{2}) & SQ_q(k,k,\frac{d}{2}) & I_k & \cdots & Q_q(n,k,\frac{d}{2})) \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ B_{s+2} = (SQ_q(k,k,\frac{d}{2}) & SQ_q(k,k,\frac{d}{2}) & \cdots & SQ_q(n,k,\frac{d}{2}) & I_k) \end{pmatrix}$$

where all the lower triangles are taken from $SQ_q(k, k, \frac{d}{2})$ or $SQ_q(n, k, \frac{d}{2})$. For any one fixed block position j of I_k , the dimension of the intersection of two different subspaces is at most t since A_1^j, \ldots, A_s^j are in the MRD code $Q_q(k, k, \frac{d}{2})$ or $Q_q(n, k, \frac{d}{2})$. For different block positions $U_j, U_i, j > i$ in the set $\{1, \ldots, s+2\}$ of I_k , the dimension of the sum of two different subspaces is at least $k + \frac{d}{2}$ from theorem 2. We get the conclusion.

4.2 Examples

When s = 0, this is the case, which happens to the theorem 2. When s = 1, in this case we refer to table 1 and the following result can be proved.

Corollary 1. If
$$n \geq k \geq d$$
, we have $A_q(2k+n,d,k) \geq q^{(k+n)(k-\frac{d}{2}+1)} + q^{n(k-\frac{d}{2}+1)} \times (\sum_{r=\frac{d}{2}}^{k-\frac{d}{2}} A_r(Q_q(k,k,\frac{d}{2})) + (\sum_{i=\frac{d}{2}}^{k-\frac{d}{2}} A_r(Q_q(k,k,\frac{d}{2}))) \times (\sum_{i=\frac{d}{2}}^{k-\frac{d}{2}} A_r(Q_q(n,k,\frac{d}{2})).$

For example, when n=5, k=5, d=4, then $A_q(15,4,5) \geq q^{40} + (A_2(Q_q(5,5,2)) + A_3(Q_q(5,5,2)) \times q^{20} + (A_2(Q_q(5,5,2)) + A_3(Q_q(5,5,2)))^2$. Assume that q=2, then $A_2(15,4,5) \geq 2^{40} + (4805 + 124930) \times 2^{20} + (4805 + 124930)^2 = 1252379805361$, exceeds the current known code $A_2(15,4,5) \geq 1235787711790$ which was constructed by the linkage.

When n=6, k=5, d=4, then $A_q(16,4,5) \ge q^{44} + (A_2(Q_q(5,5,2)) + A_3(Q_q(5,5,2)) \times q^{24} + (A_2(Q_q(5,5,2)) + A_3(Q_q(5,5,2))) \times (A_2(Q_q(6,5,2)) + A_3(Q_q(6,5,2)))$. Assume that q=2, then $A_2(16,4,5) \ge 19843523036401$, while the current known code is 19772603404689.

When n = 7, k = 5, d = 4, then $A_q(17,4,5) \ge q^{48} + (A_2(Q_q(5,5,2)) + A_3(Q_q(5,5,2)) \times q^{28} + (A_2(Q_q(5,5,2)) + A_3(Q_q(5,5,2))) \times (A_2(Q_q(7,5,2)) + A_3(Q_q(7,5,2)))$. Assume that q = 2, then $A_2(17,4,5) \ge 316614572112241$, while the current known code is 316361655057323.

When n=8, k=5, d=4, then $A_q(18,4,5) \geq q^{52} + (A_2(Q_q(5,5,2)) + A_3(Q_q(5,5,2)) \times q^{32} + (A_2(Q_q(5,5,2)) + A_3(Q_q(5,5,2))) \times (A_2(Q_q(8,5,2)) + A_3(Q_q(8,5,2)))$. Assume that q=2, then $A_2(18,4,5) \geq 5062094281261681$, while the current known code is 5061786480788587.

When n = 6, k = 6, d = 4, then $A_q(18, 4, 6) \ge q^{60} + (\sum_{r=2}^4 A_r(Q_q(6, 6, 2))) \times q^{30} + (\sum_{r=2}^4 A_r(Q_q(6, 6, 2)))^2$. Assume that q = 2, then $A_2(18, 4, 6) \ge 1321055665352277121$, while the current known code is 1301902384896972957.

When n = 7, k = 6, d = 4, then $A_q(19, 4, 6) \ge q^{65} + (\sum_{r=2}^4 A_r(Q_q(6, 6, 2))) \times q^{35} + (\sum_{r=2}^4 A_r(Q_q(6, 6, 2))) \times (\sum_{r=2}^4 A_r(Q_q(7, 6, 2)))$. Assume that q = 2, then $A_2(19, 4, 6) \ge 41829335877977260673$, while the current known code is 41660876316712223851.

When n=6, k=6, d=6, then $A_q(18,6,6) \ge q^{48} + (A_3(Q_q(6,6,3))) \times q^{24} + (A_3(Q_q(6,6,3)))^2$. Assume that q=2, then $A_2(18,6,6) \ge 1321055665352277121$, while the current known code is 1301902384896972957.

When n=7, k=6, d=6, then $A_q(19,6,6) \geq q^{52} + (A_3(Q_q(6,6,3))) \times q^{28} + (A_3(Q_q(6,6,3))) \times (A_3(Q_q(7,6,3)))$. Assume that q=2, then $A_2(19,6,6) \geq 41829335877977260673$, while the current known code is 41660876316712223851.

Table 1: New subspace codes from parallel linkage

$A_q(n,k,d)$	New	Old
$A_2(15,4,5)$	1252379805361	1235787711790
$A_3(15,4,5)$	12399152568347096641	12394544365887696067
$A_4(15,4,5)$	1215514411238392851780481	1215478900794081741379237
$A_5(15,4,5)$	9113715532351043940956916001	9113676963739967346201192181
$A_7(15,4,5)$	6369953433032789460601458266	6369951878418978850938882154
	169601	998943
$A_8(15,4,5)$	1329603936275508669606118276	1329603830010446369320349184
$A_8(10, 4, 0)$	013276161	800629897
$A_9(15,4,5)$	1478344516592412787455586580	1478344472192502033634129606
A9(10, 4, 0)	29146634561	95716746417
$A_2(16,4,5)$		19772603404689
	1004000504591772043921	1003958093636913086356
$A_4(16,4,5)$	311163172623815098594853761	311162598603284926601722789
$A_5(16,4,5)$	569604907196437900170969666	569604810233747959140019927
715(10, 4, 0)	6001	4056
$A_7(16,4,5)$	152942545364298885085407927	152942544600839682211042601
77(10, 4, 5)	47953212001	25199891012
$A_8(16,4,5)$	544605729453422094767666867	544605728772278832873615029
118(10, 1, 0)	6810298670081	1737261978761
$A_9(16,4,5)$	969941808565538232088309356	969941808205500584267352435
	407880730794721	307639908204424
$A_2(17,4,5)$		316361655057323
$A_3(17,4,5)$		81320605584592333256896
$A_4(17,4,5)$	796576337880209517283482182	796576252424409420394019074
	41	93
$A_5(17,4,5)$	356003008694948362952505652	356003006396092474470158189
	0410001	5765556
$A_7(17,4,5)$	367215049622878118441926070	367215049586616076988713969
	05053113534401	88494253570488
$A_8(17,4,5)$	223070506509377829926955265	223070506505125409945032726
	58406437716147201	04052070322817161

continued table

$\begin{array}{c cccccccc} A_q(n,k,d) & \text{New} & \text{Old} \\ A_9(17,4,5) & 636378820366486886124356752 & 636378820363628933337809933 \\ A_2(18,4,5) & 5062094281261681 & 5061786480788587 \\ A_3(18,4,5) & 5062094281261681 & 5061786480788587 \\ A_4(18,4,5) & 203923520267181222664011458 & 203923520620648811617657149 \\ 32321 & 36261 \\ A_5(18,4,5) & 32321 & 36261 \\ A_5(18,4,5) & 3764166001 & 9564044756 \\ A_7(18,4,5) & 81683334056450410727244356 & 881683334057465200849902241 \\ 50598758561216801 & 56399442519707072 \\ A_8(18,4,5) & 13696794644823698671485722 & 913696794644993679134854045 \\ A_9(18,4,5) & 147528144040561519972171750 & 417528144040576943162937097 \\ A_2(18,4,6) & 1321055665352277121 & 1301902384896972957 \\ A_3(18,4,6) & 1321055665352277121 & 1301902384896972957 \\ A_3(18,4,6) & 13364977346615645679038498701 & 43225562953761729683056546744 \\ A_4(18,4,6) & 1368321 & 15666769 \\ A_7(18,4,6) & 50827312139771315191417379894 & 50827299725042503817812207998 \\ A_7(18,4,6) & 50827312139771315191417379894 & 50827299725042503817812207998 \\ A_7(18,4,6) & 50827312139771315191417379894 & 50827299725042503817812207998 \\ A_7(18,4,6) & 50827312139771359445456241 & 68049806785911717931336256 \\ A_9(18,4,6) & 15329290735337201203431548481 & 15329289509595962385976540015 \\ A_7(18,4,6) & 5082731213977137260673 & 41660876316712223851 \\ A_8(18,4,6) & 14829335877977260673 & 41660876316712223851 \\ A_3(19,4,6) & 41829335877977260673 & 41660876316712223851 \\ A_4(19,4,6) & 41829335877977260673 & 41660876316712223851 \\ A_5(19,4,6) & 41829335877977260673 & 41660876316712223851 \\ A_7(19,4,6) & 41829335879977260673 & 41660876316712223851 \\ A_7(19,4,6) & 41829335879977260673 & 41660876316712223851 \\ A_5(19,4,6) & 4182933587997260673 & 41660876316712223851 \\ A_7(19,4,6) $			continued table
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_q(n,k,d)$		Old
$\begin{array}{c} 307227160504934301\\ A_2(18,4,5) & 5062094281261681 & 5061786480788587\\ A_3(18,4,5) & 6586968939449073017068081 & 6586969052351977742082856\\ A_4(18,4,5) & 203923520267181222664011458 & 203923520620648811617657149\\ 32321 & 36261\\ A_5(18,4,5) & 222501878974454212088292624 & 222501878997557796543846638\\ 3764166001 & 9564044756\\ 881683334056450410727244356 & 881683334057465200849902241\\ 60598758561216801 & 56399442519707072\\ 17565157707505515521 & 86035285169051407881\\ A_9(18,4,5) & 417528144040561519972171750 & 417528144040576943162937097\\ 13157922612937898347041 & 62272976069664226251756\\ A_2(18,4,6) & 1321055665352277121 & 1301902384896972957\\ A_3(18,4,6) & 43241984454039791949376848001 & 43225562953761729683056546744\\ A_4(18,4,6) & 13364977346615645679038498701 & 13364850324721907495003191\\ 19608321 & 15666769\\ A_7(18,4,6) & 86915431345555286301049529274 & 86915062398553386450111346455\\ A_7(18,4,6) & 7508628845999547723521 & 80325862935765420133852250\\ A_8(18,4,6) & 15329290735337201203431548481 & 1532928959595962385976540015\\ A_8(18,4,6) & 17973218529883895304078740000 & 17973217989922197607083648785\\ A_2(19,4,6) & 41829335877977260673 & 41660876316712223851\\ A_3(19,4,6) & 10504270152299418377046931486963 & 10503811797764100313173626438410\\ A_4(19,4,6) & 13685359611640410968142696785811 & 1368533406753251523327488538756\\ A_5(19,4,6) & 7311080812501 & 27160956999473272557435557305\\ A_7(19,4,6) & 854254430749233338554217116630539 & 8542544264787894328644239651424\\ A_5(19,4,6) & 502310159279228668519841248394562 & 5023101586504404954636792632338\\ A_{2}(19,4,6) & 7311080812501 & 7813806156458444082343 & 668612867543533298404046\\ A_{2}(19,4,6) & 502310159279228668519841248394562 & 5023101586504404954636792632338\\ A_{2}(19,4,6) & 502310159279228668519841248394$	$A_9(17,4,5)$		636378820363628933337809933
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		3072271636040343681	8862136580032063628
$\begin{array}{c} A_4(18,4,5) \\ 32321 \\ A_5(18,4,5) \\ 32321 \\ \\ A_5(18,4,5) \\ 32321 \\ \\ 222501878974454212088292624 \\ 3764166001 \\ \\ A_7(18,4,5) \\ 60598758561216801 \\ \\ A_8(18,4,5) \\ 913696794644823698671485722 \\ 17565157707505515521 \\ 417528144040561519972171750 \\ 417528144040576943162937898347041 \\ A_2(18,4,5) \\ 13157922612937898347041 \\ A_2(18,4,6) \\ 132105566535277121 \\ 43(18,4,6) \\ 13608321 \\ 43(18,4,6) \\ 43641844045528838953049829274 \\ 67261010500001 \\ 47(18,4,6) \\ 50827312139771315191417379894 \\ 7508628845999547723521 \\ 48_8(18,4,6) \\ 48_9(18,4$			5061786480788587
$\begin{array}{c} A_4(18,4,5) \\ A_5(18,4,5) \\ \hline \\ A_5(18,4,5) \\ \hline \\ A_7(18,4,5) \\ \hline \\ A_7(18,4,5) \\ \hline \\ A_7(18,4,5) \\ \hline \\ A_7(18,4,5) \\ \hline \\ A_8(18,4,5) \\ \hline \\ A_8(18,4,5) \\ \hline \\ A_8(18,4,5) \\ \hline \\ A_9(18,4,5) \\ \hline \\ A_9(18,4,5) \\ \hline \\ A_9(18,4,5) \\ \hline \\ A_1528144040561519972171750 \\ \hline \\ A_15281440405765515707505515521 \\ \hline \\ A_152814404057652038498071 \\ \hline \\ A_2(18,4,6) \\ \hline \\ \\ A_2(18,4,6) \\ \hline \\ \\ A_2(18,4,6) \\ \hline \\ \\ \\ A_2(18,4,6) \\ \hline \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ $	$A_3(18,4,5)$	6586968939449073017068081	6586969052351977742082856
$\begin{array}{c} 3521\\ A_5(18,4,5) \\ 222501878974454212088292624 \\ 3764166001 \\ 3764160001 \\ 3764160001 \\ 3764160001 \\ 3764160001 \\ 3764160001 \\ 3764160001 \\ 376416001 \\ 376616001 \\ 376616001 \\ 3766001 \\$			203923520620648811617657149
$\begin{array}{c} A_5(18,4,5) \\ A_7(18,4,5) \\ 81683334056450410727244356 \\ 60598758561216801 \\ A_8(18,4,5) \\ 17565157707505515521 \\ 417528144040561519972171750 \\ 13157922612937898347041 \\ 42(18,4,6) \\ 1321055665352277121 \\ 43(18,4,6) \\ 43241984454039791949376848001 \\ 43245667994821937898347041 \\ 43(18,4,6) \\ 43241984454039791949376848001 \\ 432656623953761729683056546744 \\ 43(18,4,6) \\ 43615431345555286301049529274 \\ 46726010500001 \\ 47(18,4,6) \\ 47(18,4,$	$A_4(10, 4, 9)$	32321	36261
$\begin{array}{c} 370410001\\ A_7(18,4,5) \\ 81683334056450410727244356 \\ 60598758561216801 \\ A_8(18,4,5) \\ 17565157707505515521 \\ A_9(18,4,5) \\ 13157922612937898347041 \\ A_2(18,4,6) \\ 1321055665352277121 \\ A_3(18,4,6) \\ 1368334057465200849902241 \\ 62272976069664226251756 \\ A_2(18,4,6) \\ 136908321 \\ A_3(18,4,6) \\ 136908321 \\ A_5(18,4,6) \\ 6726010500001 \\ A_7(18,4,6) \\ 6736184,4,6) \\ 15329290735337201203431548481 \\ A_9(18,4,6) \\ 15329290735337201203431548481 \\ A_9(18,4,6) \\ 15329290735337201203431548481 \\ A_9(18,4,6) \\ 136908321 \\ A_8(18,4,6) \\ 15329290735337201203431548481 \\ A_9(18,4,6) \\ 15329290735337201203431548481 \\ A_9(18,4,6) \\ 13880315113074804045244546241 \\ A_9(19,4,6) \\ 13685359611640410968142696785811 \\ A_{1}(19,4,6) \\ 136853596116404068142696785811 \\ A_{2}(19,4,6) \\ 8814061592799228668519841248394562 \\ 4_{2}(19,4,6) \\ 854254430749233338554217116630539 \\ A_{2}(19,4,6) \\ 854254430749233338554217116630539 \\ 85425442647878943286442366792632338 \\ 4_{2}(19,4,6) \\ 854254430749233338554217116630539 \\ 85425442647878943286442365792632338 \\ 4_{2}(19,4,6) \\ 854254430749233338554217116630539 \\ 854254426478789432864423651424 \\ 668612867543534298440406 \\$	1_(19 1 5)	222501878974454212088292624	222501878997557796543846638
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_5(18,4,5)$	3764166001	9564044756
$\begin{array}{c} 03987350121001\\ A_8(18,4,5) \\ 17565157707505515521 \\ A_9(18,4,5) \\ 13157922612937898347041 \\ A_2(18,4,6) \\ 1321055665352277121 \\ 1301902384896972957\\ A_3(18,4,6) \\ 43241984454039791949376848001 \\ A_4(18,4,6) \\ 19608321 \\ 13666769\\ A_5(18,4,6) \\ 47(18,4,6) \\ 47(18,4,6) \\ 48(18,4,6) \\ $	1 (10 1 5)	881683334056450410727244356	881683334057465200849902241
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_7(18,4,5)$	60598758561216801	56399442519707072
$\begin{array}{c} A_9(18,4,5) \\ A_9(18,4,5) \\ A_2(18,4,6) \\ A_2(18,4,6) \\ A_3(18,4,6) \\ A_3(18,4,6) \\ A_2(18,4,6) \\ A_3(18,4,6) \\ A_3(18,4,6) \\ A_4(18,4,6) \\ A_{13364977346615645679038498701 \\ A_{19608321} \\ A_{1$	1.(19.4.5)	913696794644823698671485722	913696794644993679134854045
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_8(18,4,5)$	17565157707505515521	86035285169051407881
$\begin{array}{c} A_2(18,4,6) \\ A_2(18,4,6) \\ A_3(18,4,6) \\ A_3(18,4,6) \\ A_4(18,4,6) \\ A_4(18,4,6) \\ A_5(18,4,6) \\ A_7(18,4,6) \\ A_8(18,4,6) \\ A_{10}(18,4,6) \\ A_{10}(18$	1-(19 1 5)	417528144040561519972171750	417528144040576943162937097
$\begin{array}{c} A_3(18,4,6) \\ A_3(18,4,6) \\ A_4(18,4,6) \\ \\ A_4(18,4,6) \\ \\ A_5(18,4,6) \\ \\ A_5(18,4,6) \\ \\ A_7(18,4,6) \\ \\ \\ A_7(18,4,6) \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\$	A9(10, 4, 3)	13157922612937898347041	62272976069664226251756
$\begin{array}{c} A_4(18,4,6) \\ A_5(18,4,6) \\ A_5(18,4,6) \\ A_7(18,4,6) \\ A_7(18,4,6) \\ A_8(18,4,6) \\ A_{10}(18,4,6) \\ $	$A_2(18,4,6)$	1321055665352277121	1301902384896972957
$\begin{array}{c} A_4(18,4,6) \\ A_5(18,4,6) \\ \\ A_5(18,4,6) \\ \\ \\ A_7(18,4,6) \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\$	$A_3(18,4,6)$	43241984454039791949376848001	43225562953761729683056546744
$\begin{array}{c} A_5(18,4,6) \\ A_5(18,4,6) \\ \end{array} \\ \begin{array}{c} 86915431345555286301049529274 \\ 6726010500001 \\ \end{array} \\ A_7(18,4,6) \\ \end{array} \\ \begin{array}{c} 50827312139771315191417379894 \\ 7508628845999547723521 \\ \end{array} \\ \begin{array}{c} 50827399725042503817812207998 \\ 9337055565420133852250 \\ \end{array} \\ \begin{array}{c} A_8(18,4,6) \\ \end{array} \\ \begin{array}{c} 15329290735337201203431548481 \\ 57539946320174365857546241 \\ \end{array} \\ \begin{array}{c} 68049806785911717931336256 \\ \end{array} \\ \begin{array}{c} 17973218529883895304078740000 \\ 31880315113074804045244546241 \\ \end{array} \\ \begin{array}{c} 68049806785911717931336256 \\ \end{array} \\ \begin{array}{c} 42(19,4,6) \\ \end{array} \\ \begin{array}{c} 13685359611640410968142696785811 \\ 89910529 \\ \end{array} \\ \begin{array}{c} 43(19,4,6) \\ \end{array} \\ \begin{array}{c} 13685359611640410968142696785811 \\ 89910529 \\ \end{array} \\ \begin{array}{c} 271609616470342511721762551559160 \\ 7311080812501 \\ \end{array} \\ \begin{array}{c} 271609616470342511721762551559160 \\ 7381306156458444082343 \\ \end{array} \\ \begin{array}{c} 854254430749233338554217116630539 \\ 7881306156458444082343 \\ \end{array} \\ \begin{array}{c} 854254426787894328644239651424 \\ 668612867543534298440466 \\ \end{array} \\ \begin{array}{c} 5023101589279228668519841248394562 \\ \end{array} \\ \begin{array}{c} 5023101586504404954636792632338 \\ \end{array} \\ \begin{array}{c} 5023101586504404954636792632338 \\ \end{array} $	1.(19.4.6)	13364977346615645679038498701	13364584050324721907495003191
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_4(10, 4, 0)$	19608321	15666769
$\begin{array}{c} A_7(18,4,6) \\ A_7(18,4,6) \\ \hline \\ A_8(18,4,6) \\ \hline \\ A_8(18,4,6) \\ \hline \\ A_8(18,4,6) \\ \hline \\ A_9(18,4,6) \\ \hline \\ A_1(19,4,6) \\ \hline \\ A_1(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_3(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_3(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_3(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_3(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_3(19,4,6) \\ \hline \\ A_3(19,4,6) \\ \hline \\ A_2(19,4,6) \\ \hline \\ A_3(19,4,6) \\ \hline \\ A_3(19,4,$	1 (19 1 6)	86915431345555286301049529274	86915062398553386450111346455
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_5(10, 4, 0)$	6726010500001	8715816063570
$\begin{array}{c} A_8(18,4,6) \\ A_8(18,4,6) \\ \hline \\ A_9(18,4,6) \\ \hline \\ A_{10}(18,4,6) \\ \hline \\ A_{10}$	1. (19.4.6)	50827312139771315191417379894	50827299725042503817812207998
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_7(10, 4, 0)$	7508628845999547723521	9337055565420133852250
$\begin{array}{c} 57539946320174305857546241 \\ A_9(18,4,6) \end{array} \begin{array}{c} 17973218529883895304078740000 \\ 31880315113074804045244546241 \end{array} \begin{array}{c} 17973217989922197607083648785 \\ 65965820546280222286535998208 \end{array} \\ A_2(19,4,6) \end{array} \begin{array}{c} 41829335877977260673 \\ A_3(19,4,6) \end{array} \begin{array}{c} 41660876316712223851 \\ 10504270152299418377046931486963 \\ 89910529 \end{array} \begin{array}{c} 13685359611640410968142696785811 \\ 89910529 \end{array} \begin{array}{c} 1368533406753251523327488538756 \\ 265820613 \end{array} \\ A_5(19,4,6) \end{array} \begin{array}{c} 271609616470342511721762551559160 \\ 7311080812501 \end{array} \begin{array}{c} 2716095699954793272557435557305 \\ 913288978107256 \end{array} \\ A_7(19,4,6) \end{array} \begin{array}{c} 854254430749233338554217116630539 \\ 7881306156458444082343 \end{array} \begin{array}{c} 8542544264787894328644239651424 \\ 668612867543534298440406 \end{array} \\ \begin{array}{c} 401946 \end{array} \begin{array}{c} 502310159279228668519841248394562 \end{array} \begin{array}{c} 5023101586504404954636792632338 \end{array} \end{array}$	1-(18 1 6)	15329290735337201203431548481	15329289509595962385976540015
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	A8(10, 4, 0)	57539946320174365857546241	68049806785911717931336256
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1. (19.4.6)	17973218529883895304078740000	17973217989922197607083648785
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_9(18,4,0)$	31880315113074804045244546241	65965820546280222286535998208
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_2(19,4,6)$	41829335877977260673	41660876316712223851
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_3(19,4,6)$	10504270152299418377046931486963	10503811797764100313173626438410
$A_5(19,4,6) \begin{bmatrix} 271609616470342511721762551559160 \\ 7311080812501 \end{bmatrix} \begin{bmatrix} 2716095699954793272557435557305 \\ 913288978107256 \end{bmatrix}$ $A_7(19,4,6) \begin{bmatrix} 854254430749233338554217116630539 \\ 7881306156458444082343 \end{bmatrix} \begin{bmatrix} 8542544264787894328644239651424 \\ 668612867543534298440406 \end{bmatrix}$ $A_2(19,4,6) \begin{bmatrix} 502310159279228668519841248394562 \end{bmatrix} \begin{bmatrix} 5023101586504404954636792632338 \end{bmatrix}$		13685359611640410968142696785811	1368533406753251523327488538756
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		89910529	265820613
$A_7(19,4,6) \begin{vmatrix} 854254430749233338554217116630539 \\ 7881306156458444082343 \end{vmatrix} \begin{array}{l} 8542544264787894328644239651424 \\ 668612867543534298440406 \\ \\ A_2(19,4,6) \end{vmatrix} \begin{array}{l} 502310159279228668519841248394562 \\ \hline \\ A_3(19,4,6) \\ \hline \\ A_3($	$A_5(19,4,6)$	271609616470342511721762551559160	2716095699954793272557435557305
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		7311080812501	913288978107256
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	$A_7(19,4,6)$	854254430749233338554217116630539	8542544264787894328644239651424
$A_{2}(10, 4, 6)$		7881306156458444082343	668612867543534298440406
A8(19, 4, 0) 60102624467996257072709633 1856072893849422409772176905	$A_8(19,4,6)$	502310159279228668519841248394562	5023101586504404954636792632338
		60102624467996257072709633	1856072893849422409772176905

$A_q(n,k,d)$	New	Old
$A_9(19,4,6)$	106130054948108187145939004193505	1061300549086915846347321188422
	490502763166947652518585333609	58778340200008478852505231237828
$A_2(18,6,6)$	282957166112041	282206169223861
$A_3(18,6,6)$	79773409708059646924801	7977052899429695519499
$A_4(18,6,6)$	79228596837171602219181433561	79228465213535437618551984193
$A_5(18,6,6)$	355271606149055831666451347994	355271549860537797212597654883
	5761	4375
$A_7(18,6,6)$	367033693031672327723398953389	367033691269048237620480816438
	21195414401	30813838569
$A_8(18,6,6)$	223007453917574044765606725592	223007453646901902254328287720
	19358376203601	81255730905601
$A_9(18,6,6)$	636268545986545104493692756885	636268545755947049961803989258
	8327487086310721	2186036406787787
$A_2(19,6,6)$	4527206647567081	4515298730748862
$A_3(19,6,6)$	6461646138903634303206481	6461417369472937542117973
1 (10 6 6)	202825207897159451521448163	20282487415579548140041494
$A_4(19,6,6)$	21241	597697
$A_5(19,6,6)$	222044753843060819393754856	22204471885174522176980972
	8627695761	29003922001
$A_7(19,6,6)$	881247896969044489775331238	88124789274625593704300569
	06250729756056801	118173789808207533
$A_8(19,6,6)$	913438531246383218917381666	91343853013939625003475366
	27038002732908483921	792854319199699599873
$A_9(19,6,6)$	417455793021772239445991928	41745579287064298367037383
	87121943049164300832481	503578317354686374220297
-		

5 Conclusion

In the paper a parallel construction from MRD codes are given, which is adopted from the existing work [1]. A new lower bounds on $A_q(n,d,k)$ can be proved. In essence, our result is a generalization result of the paper [1] when n is equal to k. In addition, this method is generalized to several parallel versions of maximum rank distance codes, and this method outperform the linkage construction in some cases. These new codes are listed in Table 1.

References

- [1] L. Xu and H. Chen, "New constant-dimension subspace codes from maximum rank-distance codes," *IEEE Transactions on Information Theory*, vol. 64, no. 9, pp. 6315–6319, 2018.
- [2] E. M. Gabidulin, "Theory of codes with maximum rank distance," *Probl.inform.transm*, vol. 21, no. 1, p. 3C16, 1985.
- [3] P. Delsarte, "Bilinear forms over a finite field, with applications to coding theory," *Journal of Combinatorial Theory*, vol. 25, no. 3, pp. 226–241, 1978.
- [4] J. D. L. Cruz, E. Gorla, H. H. Lopez, and A. Ravagnani, "Rank distribution of delsarte codes," *Mathematics*, 2015.
- [5] D. Heinlein, M. Kiermaier, S. Kurz, and A. Wassermann, "Tables of subspace codes," 2016.
- [6] R. Koetter and F. R. Kschischang, "Coding for errors and erasures in random network coding," *IEEE Transactions on Information Theory*, vol. 54, no. 8, pp. 3579–3591, 2008.
- [7] A. V. Tuvi Etzion, "On q-analogs of steiner systems and covering designs," pp. 161–176, 2011.
- [8] A. Khaleghi, D. Silva, and F. R. Kschischang, "Subspace codes," in *Ima International Conference on Cryptography and Coding*, 2009, pp. 1–21.
- [9] Etzion and Vardy, "Error-correcting codes in projective space," *IEEE Transactions on Information Theory*, vol. 57, no. 2, pp. 1165–1173, 2011.
- [10] S. T. Xia and F. W. Fu, "Johnson type bounds on constant dimension codes," *Designs Codes and Cryptography*, vol. 50, no. 2, pp. 163–172, 2009.
- [11] S. Johnson, "A new upper bound for error-correcting codes," *IEEE Trans.inform.theory*, vol. 8, no. 3, pp. 203–207, 1962.
- [12] M. Braun, T. Etzion, P. Ostergård, A. Vardy, and A. Wassermann, "Existence of q-analogs of steiner systems," Forum Math. Pi, 2016.

- [13] D. Heinlein, T. Honold, M. Kiermaier, S. Kurz, and A. Wassermann, "Classifying optimal binary subspace codes of length 8, constant dimension 4 and minimum distance 6," 2017.
- [14] N. Silberstein and A. L. Trautmann, "Subspace codes based on graph matchings, ferrers diagrams, and pending blocks," *IEEE Transactions on Information Theory*, vol. 61, no. 7, pp. 3937–3953, 2015.
- [15] T. Etzion and N. Silberstein, "Codes and designs related to lifted mrd codes," *IEEE Transactions on Information Theory*, vol. 59, no. 2, pp. 1004–1017, 2013.
- [16] H. Gluesingluerssen and C. Troha, "Construction of subspace codes through linkage," *Advances in Mathematics of Communications*, vol. 10, no. 3, pp. 525–540, 2017.
- [17] D. Heinlein and S. Kurz, "symptotic bounds for the sizes of constant dimension codes and an improved lower bound," in *Coding Theory and Applications: 5th International Castle Meeting, ICMCTA 2017, Vihula, Estonia, August 28-31, 2017, Proceedings*, vol. 62, no. 4, 2017, pp. 163–191.
- [18] T. Etzion and N. Silberstein, "Error-correcting codes in projective spaces via rank-metric codes and ferrers diagrams," *IEEE Transactions* on *Information Theory*, vol. 55, no. 7, pp. 2909–2919, 2009.
- [19] A.-L. Trautmann and J. Rosenthal, "New improvements on the echelon-ferrers construction," 2010, pp. 405–408.