THE CYCLIC GROUP AND THE TRANSPOSE OF AN *R*-CYCLIC MATRIX

OCTAVIO ARIZMENDI^(†) AND JAMES A. MINGO^(*)

Dedicated to Dan-Virgil Voiculescu on his 70th birthday.

ABSTRACT. We show that using the cyclic group the transpose of an R-cyclic matrix can be decomposed along diagonal parts into a sum of parts which are freely independent over diagonal scalar matrices. Moreover, if the R-cyclic matrix is self-adjoint then the off-diagonal parts are R-diagonal.

1. INTRODUCTION

After self-adjoint operators there are only a few classes of operators with a good spectral theory; one of them is the class of R-diagonal operators introduced by Nica and Speicher. Indeed, Haagerup and Schultz [9] showed that for R-diagonal operators in a von Neumann algebra which is a factor of type II, one has an abundance of invariant subspaces. The standard examples of R-diagonal operators include Haar unitaries and circular operators. More generally one can consider the product ua where u is a Haar unitary in a *-probability space, *-free from a, see [13, Cor. 15.9]. R-cyclic operators were introduced by Nica, Shlyakhtenko, and Speicher in [15] as a generalization of Rdiagonal operators. A simple way to see this is to consider the following example from Nica-Speicher [13, Thm. 14.18].

Suppose (\mathcal{A}, φ) is a non-commutative *-probability space containing a set $\{E_{ij}\}_{i,j=1}^d$ of matrix units in \mathcal{A} . Letting $\mathcal{B} = E_{11}\mathcal{A}E_{11}$, we may write $\mathcal{A} = M_d(\mathcal{B})$ and any $X \in \mathcal{A}$ as $X = (x_{ij})_{i,j=1}^d$ with $x_{ij} = E_{1i}XE_{j1}$. If X is self-adjoint and free from $\{E_{ij}\}_{i,j=1}^d$, then the off-diagonal entries x_{ij} $(i \neq j)$ are R-diagonal. Let us recall the definition of R-diagonal and R-cyclic operators from [13, Lects. 15 and 20].

^(†)Research supported by Conacyt Grant A1-S-9764 and Simons CRM Scholarin-Residence, March 2019.

^(*)Research supported by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada and Simons CRM Scholar-in-Residence, March 2019.

Definition 1. If (\mathcal{A}, φ) is a *-probability space and $a \in \mathcal{A}$, we shall write $a^{(1)}$ for a and $a^{(-1)}$ for a^* . Then a is R-diagonal if the free cumulant $\kappa_n(a^{(\epsilon_1)}, a^{(\epsilon_2)}, \ldots, a^{(\epsilon_n)}) = 0$ unless: n is even and $\epsilon_i = -\epsilon_{i+1}$ for $1 \leq i \leq n-1$.

Definition 2. Let (\mathcal{A}, φ) be a non-commutative probability space. A matrix $A = (a_{ij})_{ij} \in M_d(\mathcal{A})$ is *R*-cyclic if for all n and all $i_1, \ldots, i_n, j_1, \ldots, j_n \in [d]$ the free cumulant $\kappa_n(a_{i_1j_1}, \ldots, a_{i_nj_n}) = 0$ unless $j_1 = i_2, \ldots, j_n = i_1$.

Example 3. Suppose (\mathcal{A}, τ) is a *-probability space and $x \in \mathcal{A}$. Let $X = \begin{pmatrix} 0 & x \\ x^* & 0 \end{pmatrix}$. Then x is R-diagonal if and only if X is R-cyclic in $M_2(\mathcal{A})$ relative to $\varphi = \operatorname{tr} \otimes \tau$ on $M_2(\mathbb{C}) \otimes \mathcal{A} = M_2(\mathcal{A})$. In [14, Thm. 1.2] it was shown that the R-diagonality of x was equivalent to the freeness of X from $M_2(\mathbb{C})$ over the subalgebra of 2×2 diagonal scalar matrices.

Example 4. In [13, Ex. 20.4] it is shown that if $X \in \mathcal{A}$ is free from a set of matrix units in \mathcal{A} , then X is R-cyclic relative to this set of matrix units. But this is not the most general situation. In [15, Thm. 8.2] it is shown that if $(M_d(\mathcal{A}), \varphi)$ is a non-commutative probability space and $A \in M_d(\mathcal{A})$ is R-cyclic then A is free from $M_d(\mathbb{C})$ over \mathcal{D}_d , where $\mathcal{D}_d \subseteq M_d(\mathbb{C})$ is the subalgebra of diagonal matrices. We shall work with this formulation.

Remark 5. The significance of *R*-diagonal and *R*-cyclic matrices is that orthogonally invariant (and hence unitarily) random matrix models produce R-cyclicity and R-diagonality. In [6, Thm. 6.2] it was shown that orthogonally invariant and constant matrices are asymptotically free provided the constant matrices converge in distribution (another proof was given in [10, Thm. 36]). Suppose X_N is an orthogonally invariant ensemble with a limit distribution. Write $N = d \times p$ and $M_N(\mathbb{C}) = M_d(\mathbb{C}) \otimes M_p(\mathbb{C})$. Let $E_{ij} = e_{ij} \otimes 1_p \in M_d(\mathbb{C}) \otimes M_p(\mathbb{C})$ be the standard matrix units. Then $\{E_{ij}\}_{ij}$ converges to a set of matrix units as $p \to \infty$. Indeed the mixed moments are stationary. Then X_N and $\{E_{ij}\}_{ij}$ are asymptotically free and thus we may realize the limit distribution of X_N as an element x in a non-commutative probability space (\mathcal{A}, φ) where there is a system of matrix units $\{E_{ij}\}_{ij}$ which are free from x. By our earlier discussion such an x is R-cyclic relative to $\{E_{ij}\}_{ij}$. The orthogonal invariance assumption is not a necessary condition, for example constant matrices and Wigner matrices are asymptotically free. A proof for real Wigner matrices is given in [12, Thm. 4.20]. Thus *R*-cyclic operators occur very naturally.

Remark 6. Our main tools will be the two representations of the cyclic group of order d in $d \times d$ matrices. The generators for these actions will

 $\mathbf{2}$

3

be the diagonal matrix D with d^{th} roots of unity on the diagonal and S the matrix that cyclically permutes the standard basis elements. See Notation 13 for the notation. These two representations are intertwined by the Fourier transform matrix. This marks yet another case where the action of a symmetry group produces freeness but the first time the group is the cyclic group. The crucial point for us is that we can write the transpose in terms of these dual actions, see Lemma 17, and S and D are free from our matrix X over the diagonal scalar matrices.

Another place where a group invariance plays a role is in traffic freeness. In this case the group is the symmetric group. A recent paper of Au, Cébron, Dahlqvist, Gabriel, and Male [2] shows the connection to freeness over the diagonal.

Remark 7. The role of the transpose in free probability arose in the work of Aubrun [3] when he showed that in a certain regime the partial transpose of Wishart matrix converged to a semi-circle law. To review this, let G_1, \ldots, G_{d_1} be independent $d_2 \times p$ complex Gaussian random matrices. By this we mean $G_i = (g_{j,k}^{(i)})_{j,k}$ with $\{g_{j,k}^{(i)}\}_{i,j,k}$ independent complex Gaussian $\mathcal{N}(0, 1)$ random variables. We let

(1)
$$W = \frac{1}{d_1 d_2} \left(\frac{G_1}{\vdots} \right) (G_1^* | \cdots | G_{d_1}^*) = \frac{1}{d_1 d_2} (G_i G_j^*)_{ij}$$

and $W^{\mathrm{T}} = \frac{1}{d_1 d_2} (G_j G_i^*)_{ij}$ be the *partial transpose* of W. Aubrun showed that when d_1, d_2 and $p \longrightarrow \infty$ such that $\frac{p}{d_1 d_2} \rightarrow c$, W^{T} converges to a semi-circular operator with free cumulants $\kappa_1 = \kappa_2 = c$. If we fix d_1 and have $d_2, p \rightarrow \infty$ we are in the regime of Banica and Nechita [5]. They showed that $d_1 W^{\mathrm{T}}$ converges in distribution to the free difference of two Marchenko-Pastur laws.

Recall that for each $0 < c < \infty$ there is a probability distribution called the Marchenko-Pastur law with parameter c, denoted MP_c. We let $a = (1 - \sqrt{c})^2$ and $b = (1 + \sqrt{c})^2$; MP_c has density $\frac{\sqrt{(b-t)(t-a)}}{2\pi t}$ on the interval [a, b] for $c \ge 1$ and for 0 < c < 1 has in addition an atom of mass 1 - c at 0, see [12, Def. 2.11]. Note that in Eq. (1) we have used a different normalization than Aubrun and Banica-Nechita, in order to simplify the notation.

So let us suppose that d_1 is fixed. Then W converges to w an operator with distribution MP_c. In [10, Thm. 3.7] the mixed moments of w and w^t were given. If we adopt the convention here that $w^{(1)} = w$ and $w^{(-1)} = w^t$ then

$$\varphi(w^{(\epsilon_1)}\cdots w^{(\epsilon_n)}) = \sum_{\pi \in NC(n)} c^{\#(\pi)} d_1^{f_{\epsilon}(\pi)}$$

where $f_{\epsilon}(\pi) = \#(\epsilon\gamma\delta\gamma^{-1}\epsilon\vee\pi\delta\pi^{-1}) + \#(\pi) - (n+1) \leq 0$ is an integer. See [10, §3] for an explanation of the notation. The point we need here is that W and W^{T} have a joint limit distribution and $\varphi(ww^t) = cd_1^{-1} + c^2 \neq c^2 = \varphi(w)\varphi(w^t)$. In particular w and w^t are not free.

Since W is asymptotically free from the matrix units $\{E_{ij}\}_{ij=1}^{d_1}$ we have that we may realize w as a matrix $(w_{ij})_{ij=1}^{d_1}$ with entries in a noncommutative probability space and w is free from the matrix units. Thus w is R-cyclic.

Banica and Nechita showed that when d_1 is fixed, d_1W^{T} converged in distribution to an operator $x_1 - x_2$ in a non-commutative probability space where x_1 and x_2 are free and which have distribution MP_{c_1} and MP_{c_2} respectively with $c_1 = cd_1\frac{d_1+1}{2}$ and $c_2 = cd_1\frac{d_1-1}{2}$.

Note that if $x = x_1 - x_2$ with $x_i \in MP_{c_i}$ then the free cumulants $\{\kappa_n\}_n$ of x_i are $\kappa_n = c_i$ for all n. Thus the free cumulants of x are given by

$$\kappa_n = c_1 + (-1)^n c_2 = cd_1 \left(\frac{d_1 + 1}{2} + (-1)^n \frac{d_1 - 1}{2} \right) = \begin{cases} cd_1d_1 & n \text{ even} \\ cd_1 & n \text{ odd} \end{cases}$$

One of the motivations for the present work is to present an intrinsic description of this distribution in terms of the matrix W^{T} itself. Indeed we shall show that the diagonal decomposition of d_1W^{T} converges to a free family of $d_1/2 + 1$ self-adjoint operators (assuming d_1 is even) such that all even cumulants are cd_1 and all odd cumulants are 0 except for the first operator which has all cumulants equal to cd_1 .

Let us illustrate this when $d_1 = 2$. We write the limit distribution of W as $w = \frac{1}{d_1} \begin{pmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \end{pmatrix}$. We have that w is MP_c and free from $M_2(\mathbb{C})$. Then the limit distribution of W^{T} is $w^t = \frac{1}{d_1} \begin{pmatrix} w_{11} & w_{21} \\ w_{12} & w_{22} \end{pmatrix}$. We write $d_1 w^t = X_0 + X_1$ with

$$X_0 = \begin{pmatrix} w_{11} & 0 \\ 0 & w_{22} \end{pmatrix}$$
 and $X_1 = \begin{pmatrix} 0 & w_{21} \\ w_{12} & 0 \end{pmatrix}$

In [10, Thm. 6.14] it was shown that X_0 and X_1 are free. X_0 is MP_{d_1c} and X_1 is an even operator with even cumulants d_1c . In this paper we extend this to the general case $d_1 \ge 1$. A precise statement is given in Section 4, Theorem 31. Note that we reach the stronger conclusion of freeness over the scalars because the entries of our matrix have the same

5

distribution-depending on whether they are diagonal or off diagonal. In the absence of this property we only get freeness over diagonal scalar matrices.

2. Preliminaries and Notations

2.1. Operator valued probability spaces. For a non commutative probability space (\mathcal{A}, τ) , let us consider the non commutative probability space $(M_d(\mathcal{A}), \varphi)$ where $\varphi = \text{tr} \otimes \tau$. That is $A \in M_d(\mathcal{A})$, $\varphi(A) = \frac{1}{d}(\tau(A_{11}) + \cdots + \tau(A_{dd}))$.

Let us denote by $\mathcal{D} = \mathcal{D}(\mathbb{C}) \subseteq \mathcal{D}(\mathcal{A}) \subseteq M_d(\mathcal{A})$, the subalgebras of scalar diagonal matrices and diagonal matrices with entries in \mathcal{A} . We shall denote by $\tilde{\varphi} : M_d(\mathcal{A}) \to \mathcal{D}$ and $E : M_d(\mathcal{A}) \to \mathcal{D}(\mathcal{A})$, the unique conditional expectations to \mathcal{D} and, respectively, to $\mathcal{D}(\mathcal{A})$ which are consistent with φ . For $X = (x_{ij})_{i,j} \in M_d(\mathcal{A})$ these are given explicitly by

$$\mathbf{E}(X) = \begin{pmatrix} x_{11} & 0 & \cdots & 0\\ 0 & x_{22} & \cdots & 0\\ \vdots & \vdots & & \vdots\\ 0 & 0 & \cdots & x_{dd} \end{pmatrix}, \quad \tilde{\varphi}(X) = \begin{pmatrix} \tau(x_{11}) & 0 & \cdots & 0\\ 0 & \tau(x_{22}) & \cdots & 0\\ \vdots & \vdots & & \vdots\\ 0 & 0 & \cdots & \tau(x_{dd}). \end{pmatrix}.$$

Let us recall the notion of operator-valued probability space and freenees with amalgamation (for a detailed exposition see [17]).

Definition 8. A \mathcal{B} -valued probability space is a triplet $(\mathcal{A}, \mathcal{B}, \mathbb{E})$, consisting of a unital algebra \mathcal{A} , a unital subalgebra $\mathcal{B} \subset \mathcal{A}$ and a conditional expectation $\mathbb{E} : \mathcal{A} \to \mathcal{B}$, i.e. a unit-preserving linear map such that $\mathbb{E}(b_1 a b_2) = b_1 \mathbb{E}(a) b_2$ for any $a \in \mathcal{A}$ and $b_1, b_2 \in \mathcal{B}$.

Definition 9. Given a \mathcal{B} valued probability space $(\mathcal{A}, \mathcal{B}, \mathbb{E})$), a family of subalgebras $(\mathcal{A}_i)_i$ with $\mathcal{B} \subset \mathcal{A}_i \subset \mathcal{A}$ for each *i* is said to be free with amalgamation over \mathcal{B} , if $\mathbb{E}(x_1 \cdots x_p) = 0$, whenever $x_j \in \mathcal{A}_{i(j)}$, $\mathbb{E}(x_j) =$ 0, for all *j*, and $i(j) \neq i(j+1)$, $j = 1, \ldots, p-1$. A family $\{s_1, \ldots, s_r\}$ of \mathcal{B} -valued random variables in \mathcal{A} are free with amalgamation over \mathcal{B} , if the family of subalgebras $\operatorname{alg}\langle s_i, \mathcal{B} \rangle$, $i = 1, \ldots, r$, are free with amalgamation over \mathcal{B} .

2.2. **Operator-valued free cumulants.** Let us denote by NC(n) the set of non-crossing partitions of [n] ([13, chapter 9], and by $\mathcal{NC} = \bigcup_{n=1}^{\infty} NC(n)$.

For $n \in \mathbb{N}$, a \mathbb{C} -multi-linear map $f : \mathcal{A}^n \to \mathcal{B}$ is called \mathcal{B} -balanced if it satisfies the \mathcal{B} -bilinearity conditions, that for all $b, b' \in \mathcal{B}, a_1, \ldots, a_n \in \mathcal{A}$, and for all $r = 1, \ldots, n-1$,

$$f(ba_1,\ldots,a_nb') = bf(a_1,\ldots,a_n)b'$$

ARIZMENDI AND MINGO

 $f(a_1, \ldots, a_r b, a_{r+1}, \ldots, a_n) = f(a_1, \ldots, a_r, ba_{r+1}, \ldots, a_n)$

A collection of \mathcal{B} -balanced maps $(f_{\pi})_{\pi \in \mathcal{NC}}$ is said to be multiplicative with respect to the lattice of non-crossing partitions if, for every $\pi \in \mathcal{NC}$, f_{π} is computed using the block structure of π in the following way:

1. If $\pi = 1_n \in NC(n)$, we just write $f_n := f_{\pi}$.

2. If $1_n \neq \pi = \{V_1, \ldots, V_k\} \in NC(n)$, then by a known characterization of \mathcal{NC} , there exists a block $V_r = \{s + 1, \ldots, s + l\}$ containing consecutive elements. For any such a block we must have

$$f_{\pi}(a_1, \ldots, a_n) = f_{\pi \setminus V_r}(a_1, \ldots, a_s f_l(a_{s+1}, \ldots, a_{s+l}), a_{s+l+1}, \ldots, a_n),$$

where $\pi \setminus V_r \in NC(n-l)$ is the partition obtained from removing the block V_r .

The operator-valued of \mathcal{B} free cumulants $(\kappa_{\pi}^{\mathcal{B}})_{\pi \in \mathcal{NC}}$ are defined as the unique multiplicative family of \mathcal{B} -balanced maps satisfying the (operator-valued) moment-cumulant formulas

$$\mathbb{E}(a_1 \dots a_n) = \sum_{\pi \in NC(n)} \kappa_{\pi}^{\mathcal{B}}(a_1, \dots, a_n)$$

By the free cumulants of a tuple $(a_1, \ldots, a_k) \in \mathcal{A}^k$, we mean the collection of all cumulant maps

$$\begin{array}{ccccc}
\kappa_{i_1,\ldots,i_n}^{\mathcal{B};a_1,\ldots,a_k} : & \mathcal{B}^{n-1} & \to & \mathcal{B}, \\
& & (b_1,\ldots,b_{n-1}) & \mapsto & \kappa_n^{\mathcal{B}}\left(a_{i_1},b_1a_{i_2},\ldots,b_{i_{n-1}}a_{i_n}\right)
\end{array}$$

for $n \in \mathbb{N}$, $1 \leq i_1, \ldots, i_n \leq k$.

Given subalgebras $(\mathcal{A}_i)_i$ such that $\mathcal{B} \subseteq \mathcal{A}_i \subseteq \mathcal{A}$ for each *i* and elements a_1, \ldots, a_n such that $a_j \in \mathcal{A}_{i_j}$, a free cumulant map $\kappa_{i_1,\ldots,i_n}^{\mathcal{B};a_1,\ldots,a_k}$ is mixed if there exists r < s such that $i_r \neq i_s$. The main feature of the operator-valued cumulants is that they characterize freeness with amalgamation.

Proposition 10 ([17]). The random variables a_1, \ldots, a_n are \mathcal{B} -free if and only if all their mixed cumulants vanish.

Let us finally state the formula for products as arguments, which will be used in the proof of our main results.

Proposition 11. [18] Suppose n_1, \ldots, n_r are positive integers and $n = n_1 + \cdots + n_r$. Given a \mathcal{B} -valued probability space $(\mathcal{A}, \mathcal{B}, \mathbb{E})$ and $a_1, \ldots, a_{n_1}, a_{n_1+1}, \ldots, a_{n_1+n_2}, \ldots, a_{n_1+\dots+n_r} \in \mathcal{A}$, let $A_1 = a_1 \cdots a_{n_1}, A_2 = a_{n_1+1} \cdots a_{n_1+n_2}, \ldots, A_r = a_{n_1+\dots+n_{r-1}+1} \cdots a_{n_1+\dots+n_r}$. Then

(2)
$$\kappa_r(A_1, A_2, \dots, A_{r-1}, A_r) = \sum_{\substack{\pi \in NC(n) \\ \pi \lor \sigma = 1_n}} \kappa_\pi(a_1, \dots, a_n),$$

where $\sigma = \{\{1, 2, \dots, n_1\} \cdots \{n_1 + n_2 + \dots + n_{r-1} + 1, \dots, n_1 + n_2 + \dots + n_r\}\}.$

3. Diagonal decompositions of R-cyclic matrices

3.1. Diagonal Decompositions.

Notation 12. Let $A \in M_d(\mathcal{A})$ be a matrix. We shall write $A = A_0 + A_1 + \cdots + A_{d-1}$ where the A_i 's are shown in Figure 1.

$$A_{0} = \begin{pmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & a_{22} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & a_{dd} \end{pmatrix} A_{1} = \begin{pmatrix} 0 & a_{12} & 0 & \cdots & 0 \\ 0 & 0 & a_{23} & \cdots & 0 \\ \vdots & \vdots & & & \vdots \\ 0 & 0 & 0 & a_{d-1,d} \\ a_{d1} & 0 & 0 & 0 \end{pmatrix}$$
$$A_{k} = \begin{pmatrix} 0 & \cdots & a_{1,k+1} & 0 & \cdots & 0 \\ 0 & \cdots & 0 & a_{2,k+2} & \cdots & 0 \\ \vdots & & \vdots & \vdots & \ddots & \vdots \\ 0 & & & & a_{d-k,d} \\ a_{d-k+1,1} & & & \cdots & 0 \\ & & \ddots & & & & \vdots \\ 0 & 0 & a_{d,k} & \cdots & & 0 \end{pmatrix}$$

FIGURE 1. The first two terms in the diagonal decomposition of A (top row), and the k^{th} term (second row).

We call this the *diagonal decomposition* of A. We interpret all subscripts modulo d, i.e. $A_k = A_l$ if $k \equiv l \pmod{d}$. The same assumption applies to the indices of our matrices, namely $a_{ij} = a_{kl}$ whenever $i \equiv k \pmod{d}$ and $j \equiv l \pmod{d}$. Since d will remain fixed throughout this paper we shall write $i \equiv j$ to mean $i \equiv j \pmod{d}$.

Notation 13. Let S be the matrix that cyclically permutes (backwards) the standard basis of \mathbb{C}^d ,

$$S = \begin{pmatrix} 0 & 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 1 & & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & & \vdots \\ 0 & 0 & \cdots & & 0 & 1 \\ 1 & 0 & \cdots & & 0 & 0 \end{pmatrix},$$

and for $\omega = \exp(2\pi i/d)$, let

$$\boldsymbol{D} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & \omega & 0 & \cdots & 0 \\ 0 & 0 & \omega^2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \omega^{d-1} \end{pmatrix}$$

Notice that the conditional E the expectation onto the diagonal matrices with entries from \mathcal{A} may be written as

(3)
$$E(X) = d^{-1}(X + DXD^{-1} + \dots + D^{d-1}XD^{-(d-1)}),$$

and we have the commutation relations

$$S^k \mathbf{D}^l = \omega^{kl} \mathbf{D}^l S^k$$
 and $\mathbf{D}^l S^k = \omega^{-kl} S^k \mathbf{D}^l$.

Lemma 14. Let $X = (x_{ij})$ be a $d \times d$ matrix. Then $(XS)_{ij} = x_{i,j-1}$, $(SX)_{ij} = x_{i+1,j}$, $(XS^{-1})_{ij} = x_{i,j+1}$, and $(S^{-1}X)_{ij} = x_{i-1,j}$. $E(XS^k) = E(S^{-k}X^t)$.

Proof. The first four equalities follow from the fact that $S_{ij} = 1$ when $j \equiv i + 1$ and 0 otherwise. The last equality follows from the fact that for any matrix X we have $E(X^t) = E(X)$ and $S^{-1} = S^t$.

Let $X = X_0 + X_1 + \cdots + X_{d-1}$ be the diagonal decomposition of X. In matrix notation $(X_k)_{ij} = x_{ij}$ if $j \equiv i + k$ and 0 otherwise. Recall that here $i \equiv j$ means equivalent modulo d, the size of the matrices.

Lemma 15. $X_k = E(XS^{-k})S^k = S^kE(S^{-k}X).$ Proof.

$$(\mathcal{E}(XS^{-k})S^{k})_{ij} = \begin{cases} \mathcal{E}(XS^{-k}))_{ii} & j \equiv i+k\\ 0 & j \not\equiv i+k \end{cases}$$
$$= \begin{cases} x_{i,i+k} & j \equiv i+k\\ 0 & j \not\equiv i+k \end{cases}$$
$$= (X_{k})_{ij},$$

and

$$(S^{k} \mathcal{E}(S^{-k}X))_{ij} = \begin{cases} \mathcal{E}(S^{-k}X))_{jj} & j \equiv i+k \\ 0 & j \not\equiv i+k \end{cases}$$
$$= \begin{cases} x_{j-k,j} & i \equiv j-k \\ 0 & j \not\equiv i+k \end{cases}$$
$$= (X_{k})_{ij}.$$

Notation 16. Let $X \in M_d(\mathcal{A})$ and Y_0, \ldots, Y_{d-1} be the diagonal decomposition of X^t . Then by Lemmas 14 and 15, $Y_k = E(X^t S^{-k})S^k = E(S^k X)S^k$.

Lemma 17. Let $X \in M_d(\mathcal{A})$.

i) If X_0, \ldots, X_{d-1} be the diagonal decomposition of X, then

$$X_k = \frac{1}{d} \sum_{i=1}^d \omega^{ik} \mathbf{D}^i X \mathbf{D}^{-i}.$$

ii) If Y_0, \ldots, Y_{d-1} be the diagonal decomposition of X^t , then

$$Y_k = S^k \left[\frac{1}{d} \sum_{i=1}^d \omega^{-ik} \mathbf{D}^i X \mathbf{D}^{-i} \right] S^k.$$

Proof. For (i) we use Eq. (3) and Lemma 15

$$X_{k} = S^{k} E(S^{-k}X) = S^{k} \frac{1}{d} \sum_{i=1}^{d} D^{i}(S^{-k}X) D^{-i} = \frac{1}{d} \sum_{i=1}^{d} \omega^{ik} D^{i}X D^{-i}$$

Similarly for (ii),

$$Y_{k} = \mathcal{E}(S^{k}X)S^{k} = \frac{1}{d}\sum_{i=1}^{d} \mathcal{D}^{i}(S^{k}X)\mathcal{D}^{-i}S^{k}$$
$$= \frac{1}{d}\sum_{i=1}^{d}\omega^{-ik}S^{k}\mathcal{D}^{i}X\mathcal{D}^{-i}S^{k} = S^{k}\left[\frac{1}{d}\sum_{i=1}^{d}\omega^{-ik}\mathcal{D}^{i}X\mathcal{D}^{-i}\right]S^{k}.$$

3.2. Freeness over $M_d(\mathbb{C})$. From now on we will assume that $X \in M_d(\mathcal{A})$, is free from $M_d(\mathbb{C})$ over \mathcal{D} . The free cumulants $\tilde{\kappa}_n : M_d(\mathcal{A}) \to \mathcal{D}$ refer to operator valued free cumulants over the algebra \mathcal{D} , i.e. free cumulants with respect to $\tilde{\varphi}$.

The following simple observation will be the beginning of our analysis.

Lemma 18. If $X \in M_d(\mathcal{A})$, is free from $M_d(\mathbb{C})$ over \mathcal{D} . Then S is free from the family $\{X_i\}_{i=1}^d$ over \mathcal{D} .

Proof. By Lemma 17, for all i, X_i is in the algebra generated by X and \mathcal{D} and thus since and X is free from $M_d(\mathbb{C})$ over \mathcal{D} , then the $\{X_i\}_{i=1}^d$ is also free from $S \in M_d(\mathbb{C})$ over \mathcal{D} .

Lemma 19. Let $M_0, M_1, M_2, \ldots, M_{d-1}$ be the diagonal decomposition of $M \in M_d(\mathcal{A})$.

Then for $i_1, \ldots, i_r \in \{0, 1, \ldots, d-1\}$ and diagonal matrices D_1, \ldots, D_r , then

$$\tilde{\kappa}_r(M_{i_1}D_1,\ldots,M_{i_r}D_r)=0,$$

whenever $i_1 + \cdots + i_r \not\equiv 0$.

Proof. This follows from cumulant moment formula. Indeed, let μ be the Möbius function for NC(r) (see [13, Lect. 9]) then

$$\tilde{\kappa}_r(M_{i_1}D_1,\ldots,M_{i_r}D_r) = \sum_{\pi \in NC(r)} \mu(0_n,\pi)(\tilde{\varphi})_\pi(M_{i_1}D_1,M_{i_2}D_2,\cdots,M_{i_r}D_r).$$

Suppose $i_1 + i_2 + \cdots + i_r \not\equiv 0$. Then for each partition π there is at least one block $V = \{b_1, \ldots, b_s\}$ such that $i_{b_1} + i_{b_2} + \cdots + i_{b_s} \not\equiv 0$ and thus $\tilde{\varphi}(M_{i_{b_1}}D_{b_1}\cdots M_{i_{b_s}}D_{b_s}) = 0$.

Corollary 20. Let $X \in M_d(\mathcal{A})$ and let

$$\tilde{X}_i = X_{d-i} = \frac{1}{d} \sum_{l=1}^d \omega^{-il} \boldsymbol{D}^l X \boldsymbol{D}^{-l}.$$

Then for $i_1, \ldots, i_r \in [n]$ we have for $D_1, \ldots, D_r \in \mathcal{D}$

$$\tilde{\kappa}_r(\tilde{X}_{i_1}D_1,\ldots,\tilde{X}_{i_r}D_r)=0$$

whenever $i_1 + \cdots + i_r \not\equiv 0 \pmod{n}$.

Proof. Indeed if $i_1 + \dots + i_r \not\equiv 0$ then $d - i_1 + \dots + d - i_r \not\equiv 0$ and hence $\tilde{\kappa}_r(\tilde{X}_{i_1}D_1, \dots, \tilde{X}_{i_r}D_r) = 0 = \tilde{\kappa}_r(X_{d-i_1}D_1, \dots, X_{d-i_r}D_r) = 0$ by Lemma 19.

Lemma 21. Let S be as above. Then for all $D_1, \ldots, D_{2r} \in \mathcal{D}$. i) $\tilde{\kappa}_r(S^{i_1}D_1, S^{i_2}D_2, \cdots, S^{i_r}D_r) = 0$ unless $i_1 + i_2 + \cdots + i_r \equiv 0$ ii) $\tilde{\kappa}_{2r}(S^{i_1}D_1, S^{-i_1}D_2, \cdots, S^{i_r}D_{2r-1}, S^{-i_r}D_{2r}) = 0$, and $\tilde{\kappa}_{2r}(S^{-i_r}D_{2r}, S^{i_1}D_1, S^{-i_1}D_2, \cdots, S^{i_r}D_{2r-1}) = 0$ unless $i_1 \equiv i_2 \equiv \cdots \equiv i_r$.

Proof. (i) Follows from Lemma 19, since $S^i = \mathbb{J}_i$ in the diagonal decomposition of the matrix $\mathbb{J} = (1)_{ij}^d$, i.e. all entries equal to 1.

(*ii*) We use induction on r. First we prove that $\tilde{\kappa}_4(S^kD_1, S^{-k}D_2, S^lD_3, S^{-l}D_4) = 0$ and $\tilde{\kappa}_4(S^{-l}D_4, S^kD_1, S^{-k}D_2, S^lD_3,) = 0$ unless $l \equiv k \pmod{d}$. This will be the base of our induction.

Indeed suppose $k \not\equiv l$ and $k \not\equiv 0 \pmod{d}$, by hypothesis there exists $D \in \mathcal{D}$ such $D = S^k D_1 S^{-k} D_2$, by the formula for products as arguments (2) we have

$$0 = \tilde{\kappa}_3(D, S^l D_3, S^{-l} D_4) = \sum_{\substack{\pi \in NC(4) \\ \pi \lor \sigma = 1_4}} \tilde{\kappa}_\pi(S^k D_1, S^{-k} D_2, S^l D_3, S^{-l} D_4),$$

where $\sigma = \{\{1, 2\}, \{3\}, \{4\}\}\}$. Thus,

$$0 = \tilde{\kappa}_4(S^k D_1, S^{-k} D_2, S^l D_3, S^{-l} D_4) + \tilde{\kappa}_2(S^k D_1 \tilde{\kappa}_2(S^{-k} D_1, S^l D_3), S^{-l} D_4) + \tilde{\kappa}_3(S^k D_1, \tilde{\kappa}_1(S^{-k} D_2) S^l D_3, S^{-l} D_4) + \tilde{\kappa}_1(S^k D_1) \tilde{\kappa}_3(S^{-k} D_2, S^l D_3, S^{-l} D_4).$$

By (i), all of the terms in the second and third line equal 0, and thus $\tilde{\kappa}_4(S^k D_1, S^{-k} D_2, S^l D_3, S^{-l} D_4) = 0.$ Similarly,

$$0 = \tilde{\kappa}_3(S^{-l}D_4, D, S^lD_3) = \sum_{\substack{\pi \in NC(4) \\ \pi \lor \rho = 1_4}} \tilde{\kappa}_\pi(S^{-l}D_4, S^kD_1, S^{-k}D_2, S^lD_3)$$

where $\rho = \{\{1\}, \{2, 3\}, \{4\}\}$. Thus,

$$0 = \tilde{\kappa}_4(S^{-l}D_4, S^kD_1, S^{-k}D_2, S^lD_3) + \tilde{\kappa}_2(S^{-l}D_4, S^kD_1\tilde{\kappa}_2(S^{-k}D_1, S^lD_3)) + \tilde{\kappa}_3(S^kD_1, \tilde{\kappa}_1(S^{-l}D_4, S^{-k}D_2)S^lD_3) + \tilde{\kappa}_3(S^{-l}D_4\kappa_1(S^kD_1), S^{-k}D_2, S^lD_3).$$

Again, by (i), all of the terms in the second and third line equal 0, and thus $\tilde{\kappa}_4(S^{-l}D_4, S^kD_1, S^{-k}D_2, S^lD_3) = 0$, which finishes the base of induction.

Now, we assume that (ii) is true for all $1 \le t \le r$ and prove that it also holds for r + 1. That is, we will prove that

$$\tilde{\kappa}_{2r+2}(S^k D_1, S^{-k} D_2, S^{i_1} D_3, S^{-i_1} D_4, \cdots, S^{i_r} D_{2r}, S^{-i_r} D_{2r+2}) = 0,$$

unless $k \equiv i_1 \equiv i_2 \equiv \cdots \equiv i_r$.

Again we write $D = S^k D_1 S^{-k} D_2$, and use the formula for products as arguments (2), yielding

$$0 = \tilde{\kappa}_{2r+1}(D, S^{i_1}D_3, S^{-i_1}D_4, \cdots, S^{i_r}D_{2r+1}, S^{-i_r}D_{2r+2})$$

$$= \sum_{\substack{\pi \in NC(2r+2)\\ \pi \forall \sigma = 1_{2r+2}}} \tilde{\kappa}_{\pi} (S^k D_1, S^{-k} D_2, S^{i_1} D_3, S^{-i_1} D_4, \cdots, S^{i_r} D_{2r+1}, S^{-i_r} D_{2r+2})$$

where $\sigma = \{\{1, 2\}, \{3\}, \{4\}, \ldots, \{2r+2\}\}$. If we consider the partitions $\pi \in NC(2r+2)$ such that $\pi \vee \sigma = 1_{2r+2}$ we have, apart from 1_{2r+2} , the collection $\pi_2, \ldots, \pi_{2r+2}$ where $\pi_i = \{\{1, i+1, \ldots, 2r+2\}, \{2, 3, \ldots, i\}\}$. This is because $\pi \vee \sigma = 1_{2r+2}$ implies that π has at most two blocks: one containing the element 1 and one containing the element 2.

If i = 2j, then

$$\tilde{\kappa}_{\pi_{2j}}(S^k D_1, S^{-k} D_2, \dots, S^{i_r} D_{2r-1}, S^{-i_r} D_{2r}) = \tilde{\kappa}_{2r-2j+1}(S^k D_1 \Delta_j, S^{-i_j} D_{2j+1}, \cdots, S^{i_r} D_{2r+1}, S^{-i_r} D_{2r+2}),$$

where $\Delta_{2j-1} = \tilde{\kappa}_{2j-1}(S^{-k}D_2, S^{i_1}D_3, S^{-i_1}D_4, \cdots, S^{i_j}D_{2j}) \in \mathcal{D}$. Since $k \neq 0$, then by (i), $\Delta_{2j-1} = 0$.

Second, if j is odd, say j = 2j + 1 then

$$\tilde{\kappa}_{\pi_{2j+1}}(S^k D_1, S^{-k} D_2, \dots, S^{i_r} D_{2r-1}, S^{-i_r} D_{2r}) = \tilde{\kappa}_{2r-2j}(S^k D_1 \Delta_j, S^{-i_j} D_{2j+2}, \cdots, S^{i_r} D_{2r+1}, S^{-i_r} D_{2r+2}),$$

where $\Delta_{2j} = \tilde{\kappa}_{2j}(S^{-k}D_2, S^{i_1}D_3, S^{-i_1}D_4, \cdots, S^{i_j}D_{2j+1}) \in \mathcal{D}.$

If it not the case that $k \equiv i_1 \equiv i_2 \equiv \cdots \equiv i_r$, then the same condition will hold for at least one block of π_j . Hence $\tilde{\kappa}_{\pi_i}(S^k D_1, S^{-k} D_2, \ldots, S^{i_r} D_{2r-1}, S^{-i_r} D_{2r}) = 0$. This proves that the cumulant $\tilde{\kappa}_{2r+2}(S^k D_1, S^{-k} D_2, S^{i_1} D_3, S^{-i_1} D_4, \cdots, S^{i_r} D_{2r}, S^{-i_r} D_{2r+2}) = 0$, as desired. \Box

Definition 22. Let $y_1, y_{-1} \in (\mathcal{A}, \varphi)$ a non-commutative probability space. If for all n and all $\epsilon_1, \epsilon_2, \ldots, \epsilon_n \in \{-1, 1\}$ we have $\tilde{\kappa}_n(y_{\epsilon_1}, y_{\epsilon_2}, \ldots, y_{\epsilon_n}) = 0$ unless n is even and $\epsilon_i = -\epsilon_{i+1}$ for $i = 1, \ldots, n-1$, we say that $\{y_1, y_{-1}\}$ are a *R*-diagonal pair, following [16, Thm. 3.1] (see also [4]).

Remark 23. If (\mathcal{A}, φ) is a *-probability space, then y is R-diagonal if and only if $\{y, y^*\}$ is an R-diagonal pair.

Theorem 24. Suppose X is free from $M_d(\mathbb{C})$ over \mathcal{D} . Let $Y_0, Y_1, \ldots, Y_{d-1}$ be the diagonal decomposition of X^t .

For d even, $Y_0, \{Y_1, Y_{d-1}\}, \{Y_2, Y_{d-2}\}, \ldots, \{Y_{d/2-1}, Y_{d/2+1}\}, Y_{d/2}$ is a free family over \mathcal{D} and $\{Y_i, Y_{d-i}\}$ is a *R*-diagonal pair for $i = 1, \ldots, d/2 - 1$.

For d odd, $Y_0, \{Y_1, Y_{d-1}\}, \{Y_2, Y_{d-2}\}, \ldots, \{Y_{(d-1)/2}, Y_{(d+1)/2}\}$, is a free family over \mathcal{D} and $\{Y_i, Y_{d-i}\}$ is a R-diagonal pair for $i = 1, \ldots, (d-1)/2$.

Proof. We write $Y_i = S^i \tilde{X}_i S^i$ for i = 0, ..., d-1 with \tilde{X}_i as in Lemma 17. Let r > 1 and $i_1, ..., i_r \in \{0, 1, ..., d-1\}$ be given. We must show that

$$\tilde{\kappa}_r(Y_{i_1}D_1,\ldots,Y_{i_r}D_r)=0$$

unless either $i_1 = \cdots = i_r = 0$, or r is even and $i_l + i_{l+1} = d$ for $l = 1, \ldots, r - 1$.

Suppose that for some l_1, \ldots, l_k we have $i_{l_1} = \cdots = i_{l_k} = 0$ and for all other j's we have $i_j \neq 0$. Let $s_j = |\{m \mid 1 \leq m < j \text{ and } i_m = 0\}|$ and $s = |\{m \mid 1 \leq m \leq r \text{ and } i_m = 0\}|$. Let $\sigma = \{V_1, \ldots, V_r\}$ be the interval partition with r blocks constructed as follows. $V_j = \{3j-2s_j-2\}$ if $i_j = 0$ and $V_j = \{3j-2s_j-2, 3j-2s_j-1, 3j-2s_j\}$ if $i_j \geq 1$. In other words, for each k such that $i_{l_k} = 0$, σ has a block of size 1 and for each k such that $i_{l_k} \neq 0$, σ has a block of size 3. For example, if $(i_1, i_2, i_3, i_4, i_5) = (0, 2, 0, 1, 2)$ we have $\sigma = \{(1), (2, 3, 4), (5), (6, 7, 8), (9, 10, 11)\}$.

For $1 \leq m \leq 3r - 2s$, let us define Z_m as follows. If $m \in V_j$ and $i_j = 0$ then $Z_m = X_0D_j$. If $m \in V_j$ and $i_j \geq 1$ then $Z_m = \tilde{X}_{i_j}$ if $m = 3j - 2s_j - 1$, $Z_m = S^{i_j}$ if $m = 3j - 2s_j - 2$ and $Z_m = S^{i_j}D_j$ if $m = 3j - 2s_j$. In the example above Z_1, \ldots, Z_{11} are $X_0D_1, S^2, X_2, S^2D_2, X_0D_3, S^1, X_1, S^1D_4, S^2, X_2, S^2D_5$ respectively. Then, by the formula for products as arguments (2), we have

$$\tilde{\kappa}_r(Y_{i_1}D_1,\ldots,Y_{i_r}D_r) = \sum_{\substack{\pi \in NC(3r-2s)\\ \pi \lor \sigma = 1_{3r-2s}}} \tilde{\kappa}_\pi(Z_1,\ldots,Z_{3r-2s})$$

The proof will consist in analyzing the non-vanishing terms in the formula above. Observe that by Lemma 18, S and the X_i 's are free. Thus for $\tilde{\kappa}_{\pi}(Z_1, \ldots, Z_{3r-2s}) \neq 0$ we must have the blocks of either π to consist of X-blocks (only connecting X's and XD's) and S-blocks (only connecting S's and SD's).

We shall break the proof into two cases. Case 1 is when for some $l \in [r]$ we have $i_l = 0$, i.e. s > 0. Case 2 is when $i_l > 0$ for all $l \in [r]$, i.e. s = 0.

Case 1. Mixed cumulant with at least one Y_0 as an entry, i.e. s > 0.

Let W be an X-block and let $b \in W$ be such that $Z_b = X_0 D$. If W is a singleton then the condition $\pi \vee \sigma = 1_{3r-2s}$ will not be satisfied. So W must have another element.

Without loss of generality, there is at least one element to the right of b (the following argument also applies if there is an element to the left). Now, let $b' \in W$ be the next element to the right of W; it must be a X_i , by freeness. Thus we may label the elements between b and b' as below. By Lemma 21 (i) we have $2i_{t_1} + \cdots + 2i_{t_{k-1}} + i_{t_k} \equiv 0 \pmod{d}$.

$$\cdots |X_0 D_{t_1-1}| \underbrace{S^{i_{t_1}}, \tilde{X}_{i_{t_1}}, S^{i_{t_1}} D_{t_1} | \cdots |S^{i_{t_k}}, \tilde{X}_{i_{t_k}}, S^{i_{t_k}} D_{t_k} | \cdots }_{b} \\ \underline{b} \\ b'$$

By Corollary 20 we must have $i_{t_1} + \cdots + i_{t_{k-1}} \equiv 0 \pmod{d}$. Hence $i_{t_k} \equiv 0 \pmod{d}$. Moreover, the blocks of σ between b and b' will not be joined by π to the other blocks of σ and required by the condition $\pi \vee \sigma = 1_{3r-2s}$. Thus there cannot be any blocks of σ between b and b', and so b' = b + 1.

Finally, if b and b' are the only elements of W, then again the condition $\pi \vee \sigma = 1_{3r-2s}$ will not be satisfied. Proceeding in the same way we may deduce that there is only one X-block, with entries of the form X_0D . Hence $i_1 = i_2 = \cdots = i_r = 0$.

Case 2. Mixed cumulant with no Y_0 as an entry.

In this case, where $i_l \ge 1$ for $1 \le l \le r$. Let $\sigma \in NC(3r)$ be the partition $\{(1, 2, 3), (4, 5, 6), \dots, (3r - 2, 3r - 1, 3r)\}$. Then

$$\tilde{\kappa}_{r}(Y_{i_{1}}D_{1},\ldots,Y_{i_{r}}D_{r}) = \sum_{\substack{\pi \in NC(3r)\\ \pi \lor \sigma = 1_{3r}}} \tilde{\kappa}_{\pi}(S^{i_{1}},\tilde{X}_{i_{1}},S^{i_{1}}D_{1},\ldots,S^{i_{r}},\tilde{X}_{i_{r}},S^{i_{r}}D_{r})$$

Suppose $\pi \in NC(3r)$ and $\tilde{\kappa}_{\pi}(S^{i_1}, \tilde{X}_{i_1}, S^{i_1}D_1, \ldots, S^{i_r}, \tilde{X}_{i_r}, S^{i_r}D_r) \neq 0$. Again, since S if free over \mathcal{D} from the X_i 's, the blocks of π must either be S-blocks, or X-blocks.

We now consider the case of S-blocks. Let $b = 3s_1$ and consider W, the S-block of π and such $b \in W$. Since $i_{s_1} \not\equiv 0$, b must be connected to some other element. Let $b' \in W$ be the next point to the right (if b is the last element in W, then b' is the smallest element of W and a similar argument applies). Then there are $s_1 < \cdots < s_k$ such that either $b' = 3s_k - 2$ or $b' = 3s_k$.

First, suppose that $b' = 3s_k$.

$$\cdots, \tilde{X}_{i_{s_{1}}}, S^{i_{s_{1}}} D_{s_{1}} | \underbrace{S^{i_{s_{2}}}, \cdots, S^{i_{s_{k-1}}} D_{s_{k-1}} | S^{i_{s_{k}}} D_{s_{k}}, \tilde{X}_{i_{s_{k}}}}_{b'}, S^{i_{s_{k}}} D_{s_{k}} | \cdots }_{b'}$$

Then by Lemma 21 (i), $2i_{s_2} + \cdots + 2i_{s_{k-1}} + i_{s_k} \equiv 0 \pmod{d}$. By Corollary 20 we must have $i_{s_1} + \cdots + i_{s_k} \equiv 0 \pmod{d}$. Hence $i_{s_k} \equiv 0 \pmod{d}$, but this case has been ruled out already.

Now, suppose $b' = 3s_k - 2$. If there are any blocks of σ between b and b' they will not be able to joined to the others by π ; contradicting our assumption that $\pi \vee \sigma = 1_{3r}$.

$$\cdots \mid S^{i_{s_1}}, \tilde{X}_{i_{s_1}}, S^{i_{s_1}} D_{s_1} \mid S^{i_{s_2}}, \tilde{X}_{i_{s_2}}, \cdots, S^{i_{s_{k-1}}} D_{s_{k-1}} \mid S^{i_{s_k}}, \tilde{X}_{i_{s_k}}, S^{i_{s_k}} D_{s_k} \mid \cdots \mid \\ \underbrace{ \mid \\ b \mid \\ b' \mid } b'$$

Thus we must have that k = 2 and $s_2 = s_1 + 1$.

Thus, any $b = 3s_1$ must be connected to $3s_1 + 1$. A similar argument shows that any $b = 3s_1 + 1$ must be connected to $3s_1$.

We thus have arrived at the following form for any S-block: $\{3b_1, 3b_1+1, 3b_2, 3b_2+1, \ldots, 3b_l+1\} \pmod{3r}$. Notice that in particular this means that r is even.

By looking at the elements between $3b_i + 1$ and $3b_{i+1}$, using Lemma 21 (*i*), and Corollary 20 we may conclude that $i_{b_r+1} \equiv -i_{b_{r+1}}$, for all $i = 1, \ldots, l$. This implies by Lemma 21 (*ii*) that $i_{b_1} \equiv i_{b_2} \equiv \cdots \equiv i_{b_n}$.

We say a block is of type k if $i_{b_1} = k$ and $0 \le k \le d$. Notice that the block next to W (a block W such that $3b_i - 2$ or $3b_i + 3$ belongs to W', for some $1 \le i \le l$) must have type -k. Thus implies $i_1 \equiv -i_2 \equiv \cdots \equiv i_r$ which proves both claims.

3.3. Scalar case. In this subsection we consider the case where $X \in M_d(\mathcal{A})$ is φ -free from $M_d(\mathbb{C})$. This corresponds to uniform *R*-cyclic matrices. These are matrices such that the non-vanishing cumulants of the entries depend only on the length. The relation with free compression by matrix units may be seen in [13, Theorem 14.20]. The importance of them in random matrix theory is that they appear as limits of unitarily invariant random matrices.

The main difference from the \mathcal{D} -valued case is the analog of Lemma 18, whose proof needs a little more work in the scalar case.

Lemma 25. Suppose $X \in M_d(\mathcal{A})$ is φ -free from $M_d(\mathbb{C})$. Then S is free from the family $\{X_i\}_{i=1}^d$.

In order to prove the lemma above we need a series of results regarding freeness and conjugation with powers of the matrix S.

Lemma 26. Let C be a centred polynomial in S. Then for all k and l we have $\varphi(C\mathbf{D}^l) = \varphi(\mathbf{D}^{-k}C\mathbf{D}^l) = 0$; and for $l \neq 0$ we have $\varphi(\mathbf{D}^l) = 0$ Proof. $\varphi(\mathbf{D}^l) = d^{-1}(1 + \omega^l + \omega^{2l} + \dots + \omega^{(d-1)l}) = 0$ for $l \neq 0$. Also $S^m \mathbf{D}_l$ and $\mathbf{D}^{-k}S^m \mathbf{D}^l$ are 0 on the diagonal for $m \neq 0$. Thus $\varphi(S^m \mathbf{D}_l) = \varphi(\mathbf{D}^{-k}S^m \mathbf{D}^l) = 0$.

Let us recall from [8, Lemma 3.8] the following lemma.

Lemma 27. Suppose (\mathcal{A}, τ) is a non-commutative probability space. Let $C \in \mathcal{A}$ be τ -free from the algebra \mathcal{B} and let $U \in \mathcal{B}$ be a unitary operator with $U^d = 1$ and such that $\varphi(U^k) = 0$ when $k \not\equiv 0 \pmod{d}$. Let $C_i = U^i C U^{-i}$ for $i = 1, \ldots, d$. Then C_1, C_2, \ldots, C_d are τ -free.

Corollary 28. Suppose $X \in M_d(\mathcal{A})$ is φ -free from $M_d(\mathbb{C})$. Then the set $\{\mathbf{D}X\mathbf{D}^{-1}, \mathbf{D}^2X\mathbf{D}^{-2}, \dots, \mathbf{D}^dX\mathbf{D}^{-d}\}$ is a free family.

The following lemma is a generalization of Lemma 18 where we now show that S is free from $\{DXD^{-1}, D^2XD^{-2}, \dots, D^dXD^{-d}\}$.

Lemma 29. Suppose $X \in M_d(\mathcal{A})$ is φ -free from $M_d(\mathbb{C})$. Then the set $\{S, \mathbf{D}X\mathbf{D}^{-1}, \mathbf{D}^2X\mathbf{D}^{-2}, \dots, \mathbf{D}^dX\mathbf{D}^{-d}\}$ is a free family.

Proof. Let $\mathcal{B}_0 = M_d(\mathbb{C})$ and $\mathcal{B}_1 = \operatorname{alg}(1, X)$ be the algebra generated by 1 and X. By assumption \mathcal{B}_0 and \mathcal{B}_1 are free, so any word which is an alternating product of centred elements is centred. For $i = 1, \ldots, d$, let $\mathcal{A}_i = \mathbf{D}^i \operatorname{alg}(1, X) \mathbf{D}^{-i}$ and $\mathcal{A}_0 = \operatorname{alg}(1, S)$. We must show that $\mathcal{A}_0, \mathcal{A}_1, \ldots, \mathcal{A}_d$ are free. Let W be an alternating product of centred elements in the algebras $\mathcal{A}_0, \mathcal{A}_1, \ldots, \mathcal{A}_d$. We will show that W is also an alternating product of centred elements in the algebras \mathcal{B}_0 and \mathcal{B}_1 and thus is centred. This will show that $\mathcal{A}_0, \mathcal{A}_1, \ldots, \mathcal{A}_d$ are free.

So, to this end let W be an alternating product of centred elements in the algebras $\mathcal{A}_0, \mathcal{A}_1, \ldots, \mathcal{A}_d$. We may write $W = V_1 \cdots V_k$ with $V_i \in \mathcal{A}_{j_i}$ such that $j_1, \ldots, j_k \in \{0, 1, \ldots, d\}, j_1 \neq j_2 \neq \cdots \neq j_k$ and $\varphi(V_i) = 0$. For each i for which $j_i > 0$ there is $A_i \in alg(1, X)$ with $\varphi(A_i) = 0$ and $V_i = \mathbf{D}^{j_i} A_i \mathbf{D}^{-j_i}$.

If *i* is such that $j_i \neq 0$ then we can write $V_i V_{i+1}$ as either $\mathbf{D}^{j_i} A_i$ $\mathbf{D}^{-j_i+j_{i+1}} A_{i+1} \cdots$ if $j_{i+1} \neq 0$ or $\mathbf{D}^{j_i} A_i [\mathbf{D}^{-j_i} V_{i+1} \mathbf{D}^{j_{i+2}}] A_{i+2} \cdots$ if $j_{i+1} = 0$ (with $\mathbf{D}^{-j_i} V_{i+1} \mathbf{D}^{j_{i+2}} \in \mathcal{B}_0$). In either case, by Lemma 26, A_i is followed by a centred element of \mathcal{B}_0 . If $j_i = 0$ then $j_{i+1} \neq 0$ so $V_{i-1} V_i V_{i+1} = \cdots A_{i-1} [\mathbf{D}^{-j_{i-1}} V_i \mathbf{D}^{j_{i+1}}] A_{i+1} \cdots$. So again we have a alternating product of centred elements of \mathcal{B}_0 and \mathcal{B}_1 . This proves the claim.

Now we are ready to prove Lemma 25.

Proof of Lemma 25. We have by Lemma 29, $\{S, \mathbf{D}X\mathbf{D}^{-1}, \mathbf{D}^2X\mathbf{D}^{-2}, \dots, \mathbf{D}^d X\mathbf{D}^{-d}\}$ is a free family. In particular, we have that S is free from any linear combination of $\{\mathbf{D}X\mathbf{D}^{-1}, \mathbf{D}^2X\mathbf{D}^{-2}, \dots, \mathbf{D}^dX\mathbf{D}^{-d}\}$ and thus from $\{X_i\}_i$, by Lemma 17.

Next we come to the analogs of Lemmas 19, 20, and 21. The reader will easily convince himself that the proofs of these lemmas are also valid in the case $X \in M_d(\mathcal{A})$ is φ -free from $M_d(\mathbb{C})$, by replacing the diagonal matrices D_i 's by scalars (or by 1), and then use Lemma 25 instead of Lemma 18.

Similarly the proof of the following theorem follows exactly the same steps as the proof of Theorem 24, with the obvious changes.

Theorem 30. Suppose X is free from $M_d(\mathbb{C})$ over \mathbb{C} . Let $Y_0, Y_1, \ldots, Y_{d-1}$ be the diagonal decomposition of X^t .

For d even, $Y_0, \{Y_1, Y_{d-1}\}, \{Y_2, Y_{d-2}\}, \ldots, \{Y_{d/2-1}, Y_{d/2+1}\}, Y_{d/2}$ is a free family over \mathbb{C} and $\{Y_i, Y_{d-i}\}$ is a *R*-diagonal pair for $i = 1, \ldots, d/2 - 1$.

For d odd, Y_0 , $\{Y_1, Y_{d-1}\}$, $\{Y_2, Y_{d-2}\}$, ..., $\{Y_{(d-1)/2}, Y_{(d+1)/2}\}$, is a free family over \mathbb{C} and $\{Y_i, Y_{d-i}\}$ is a R-diagonal pair for i = 1, ..., (d - 1)/2.

4. Return to the Wishart case

Let us conclude by returning to the case of a Wishart matrix W (c.f. Eq. (1)) where d_1 is fixed and $p/d_2 \rightarrow cd_1$, but in the case of arbitrary d_1 . Let w be the limit distribution of the Wishart matrix in Eq. (1). We write d_1w as a $d_1 \times d_1$ matrix: $d_1w = (w_{ij})_{ij}$. Then d_1W^{T} has the limit distribution $d_1w^t = (w_{ji})_{ij}$. When d_1 is even let

(4)
$$d_1 w^t = Y_0 + Y_1 + Y_{d_1-1} + \dots + Y_{d_1/2-1} + Y_{d_1/2+1} + Y_{d_1/2},$$

and when d_1 is odd let

(5)
$$d_1 w^t = Y_0 + Y_1 + Y_{d_1-1} + \dots + Y_{(d_1+1)/2-1} + Y_{(d_1+1)/2+1}.$$

Since all the off-diagonal entries of w are R-diagonal with the same distribution: $\tilde{\kappa}_{2n}(w_{12}, w_{21}, \ldots, w_{12}, w_{21}) = c$ and the diagonal entries are all MP_c, we have that $\tilde{\kappa}_r(Y_{i_1}, \ldots, Y_{i_r}) = \kappa_r(Y_{i_1}, \ldots, Y_{i_r})$.

Thus, we are in the scalar case, as in Section 3.3. Thus, we arrive to the next theorem that states that the diagonal decomposition of d_1w^t is a free decomposition.

Theorem 31. Let w be the limit distribution of the Wishart matrix W in Eq. (1). Suppose d_1 is fixed and $p/d_2 \rightarrow cd_1$. Then $Y_0, Y_1, \ldots, Y_{d_1/2}$ are *-free when d_1 is even and when d_1 is odd we have $Y_0, Y_1, \ldots, Y_{(d_1+1)/2}$ are *-free. Moreover Y_0 has the distribution MP_{cd1}, $Y_{d_1-i} = Y_i^*$, and for $i \geq 1$, Y_i is an R-diagonal operator with $\kappa_{2n}(Y_i, Y_i^*, \ldots, Y_i, Y_i^*) = cd_1$. When d_1 is even $Y_{d_1/2}$ is even with $\kappa_{2n}(Y_{d_1/2}, Y_{d_1/2}^*, \ldots, Y_{d_1/2}, Y_{d_1/2}^*) = cd_1$.

ARIZMENDI AND MINGO

Acknowledgements

This work was carried out during the program 'New Developments in Free Probability and Applications' at the Centre de Recherches Mathématiques (Montreal) March 1 - 31, 2019, as part of the 50th anniversary year program. The authors are very grateful to the director and the staff of the Centre for the funding and hospitality.

References

- O. Arizmendi, I. Nechita, and C. Vargas, On the asymptotic distribution of block-modified random matrices, J. Math. Phys., 57 (2016), no. 1, 015216, 25 pp.
- [2] B. Au, G. Cébron, A. Dahlqvist, F. Gabriel, and C. Male, Large permutation invariant random matrices are asymptotically free over the diagonal, arXiv:1805.07045.
- G. Aubrun, Partial Transposition Of Random States And Non-Centered Semicircular Distributions, *Random Matrix Theory and Applications*, 1 (2012), no. 2, 1250001, 29 pp.
- [4] M. Boedihardjo and K. Dykema, On algebra-valued *R*-diagonal elements, *Houston Math. J.*, 44 (2018), 209-252.
- [5] T. Banica and I. Nechita, Asymptotic Eigenvalue Distributions of Block-Transposed Wishart Matrices, J. Theor. Probab., 26 (2013), 855–869.
- [6] M. Capitaine and M. Casalis, Cumulants for random matrices as convolutions on the symmetric group, II, J. Theor. Probab., 20, 505-533 (2007).
- [7] M. Fukuda and P. Śniady, Partial Transpose Of Random Quantum States: Exact Formulas And Meanders, J. Math. Phys. 54 (2013), no. 4, 042202, 23 pp.
- [8] Haagerup, U., Larsen, F. Brown's spectral distribution measure for *R*-diagonal elements in finite von Neumann algebras. J. Funct. Analy., 176(2), 331–367 (2000).
- [9] U. Haagerup and H. Schultz, Invariant subspaces for operators in a general II₁-factor. Publ. Math. Inst. Hautes Études Sci., 109 (2009), 19–111.
- [10] J. A. Mingo, M. Popa, Real second order freeness and Haar orthogonal matrices, J. Math. Phys. 54 (2013), no. 5, 051701, 35 pp.
- [11] J. A. Mingo and M. Popa, Freeness and the partial transposes of Wishart random matrices, *Canad. J. Math.* **71** (2019), 659-681.
- [12] J. A. Mingo and R. Speicher, Free Probability and Random Matrices, Fields Institute Monographs 35, Springer, 2017
- [13] A. Nica and R. Speicher, *Lectures on the Combinatorics of Free Probability*, Cambridge Univ. Press, 2006.
- [14] A. Nica, D. Shlyakhtenko and R. Speicher, *R*-Diagonal Elements and Freeness with Amalgamation, *Canad. J. Math.* 53 (2001), 355-381.
- [15] A. Nica, D. Shlyakhtenko and R. Speicher, *R*-Cyclic Families of Matrices in Free Probability, *J. Funct. Anal.* 188 (2002), 227–271.
- [16] R. Speicher and P. Śniady, Continuous family of invariant subspaces for *R*diagonal operators. *Invent. Math.* **104** (2001), 329–363.

- [17] R. Speicher, Combinatorial theory of the free product with amalgamation and operator-valued free probability theory, Memoirs of the American Math. Society, vol. 132, 1998.
- [18] R. Speicher, A conceptual proof of a basic result in the combinatorial approach to freeness, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* 3 (2000), 213– 222.
- [19] C. E. I. Redelmeier, Real second-order freeness and the asymptotic real secondorder freeness of several real matrix ensembles, *Int. Math. Res. Not.* 2014, no. 12, pp. 3353-3395.

CENTRO DE INVESTIGACIÓN EN MATEMÁTICAS, GUANAJUATO, MEXICO *E-mail address:* octavius@cimat.mx

DEPARTMENT OF MATHEMATICS AND STATISTICS, QUEEN'S UNIVERSITY, JEFFRY HALL, KINGSTON, ONTARIO, K7L 3N6, CANADA

E-mail address: mingo@mast.queensu.ca