# Obstructions for bounded shrub-depth and rank-depth

O-joung Kwon\* $^{\dagger 1,2}$ , Rose McCarty<sup>3</sup>, Sang-il Oum\* $^{2,4}$ , and Paul Wollan<sup>5</sup>

<sup>1</sup>Department of Mathematics, Incheon National University, Incheon, South Korea.

<sup>2</sup>Discrete Mathematics Group, Institute for Basic Science (IBS),
Daejeon, South Korea.

<sup>3</sup>Department of Combinatorics and Optimization, University of Waterloo, Waterloo, Canada.

<sup>4</sup>Department of Mathematical Sciences, KAIST, Daejeon, South Korea.

January 19, 2022

#### Abstract

Shrub-depth and rank-depth are dense analogues of the tree-depth of a graph. It is well known that a graph has large tree-depth if and only if it has a long path as a subgraph. We prove an analogous statement for shrub-depth and rank-depth, which was conjectured by Hliněný, Kwon, Obdržálek, and Ordyniak [Tree-depth and vertex-minors, European J. Combin. 2016]. Namely, we prove that a graph has large rank-depth if and only if it has a vertex-minor isomorphic to a long path. This implies that for every integer t, the class of graphs with no vertex-minor isomorphic to the path on t vertices has bounded shrub-depth.

<sup>&</sup>lt;sup>5</sup>Department of Computer Science, University of Rome, "La Sapienza", Rome, Italy.

<sup>\*</sup>Supported by IBS-R029-C1.

<sup>&</sup>lt;sup>†</sup>Supported by the National Research Foundation of Korea (NRF) grant funded by the Ministry of Education (No. NRF-2018R1D1A1B07050294).

<sup>&</sup>lt;sup>1</sup>E-mail addresses: ojoungkwon@gmail.com (Kwon), rose.mccarty@uwaterloo.ca (McCarty), sangil@ibs.re.kr (Oum), wollan@di.uniroma1.it (Wollan).

### 1 Introduction

Nešetřil and Ossona de Mendez [17] introduced the tree-depth of a graph G, which is defined as the minimum height of a rooted forest whose closure contains the graph G as a subgraph. This concept has been proved to be very useful, in particular in the study of graph classes of bounded expansion [18]. Similar to the grid theorem for tree-width of Robertson and Seymour [24], it is known that a graph has large tree-depth if and only if it has a long path as a subgraph, see [17, Proposition 6.1]. For more information on tree-depth, the readers are referred to the surveys [20, 17] by Nešetřil and Ossona de Mendez.

There have been attempts to define an analogous concept suitable for dense graphs. For tree-width, this line of research has resulted in width parameters such as clique-width [3] and rank-width [22]. In a conference paper published in 2012, Ganian, Hliněný, Nešetřil, Obdržálek, Ossona de Mendez, and Ramadurai [8] introduced the *shrub-depth* of a graph class, as an extension of tree-depth for dense graphs. Recently, DeVos, Kwon, and Oum [6] introduced the *rank-depth* of a graph as an alternative to shrub-depth and showed that shrub-depth and rank-depth are equivalent in the following sense.

**Theorem 1.1** (DeVos, Kwon, and Oum [6]). A class of graphs has bounded rank-depth if and only if it has bounded shrub-depth.

Theorem 1.1 allows us to work exclusively with rank-depth going forward, and we omit the definition of shrub-depth. The definition of rank-depth is presented in Section 2.

One useful feature of rank-depth is that it does not increase under taking vertex-minors. In other words, if H is a vertex-minor of G, then the rank-depth of H is at most that of G. This allows us to consider obstructions for having small rank-depth in terms of vertex-minors. DeVos, Kwon, and Oum [6] showed that the rank-depth of the n-vertex path is larger than  $\log n/\log(1+4\log n)$  for  $n \geq 2$  and thus graphs having a long path as a vertex-minor have large rank-depth. Hliněný, Kwon, Obdržálek, and Ordyniak [11] conjectured that the converse is also true. Their original conjecture was stated in terms of shrub-depth but is equivalent by Theorem 1.1. We prove their conjecture as follows.

**Theorem 1.2.** For every positive integer t, there exists an integer N(t) such that every graph of rank-depth at least N(t) contains a vertex-minor isomorphic to the path on t vertices.

Courcelle and Oum [4] showed that there is a CMSO<sub>1</sub> transduction that maps a graph to its vertex-minors. Therefore, Theorem 1.2 implies that a class  $\mathcal{G}$  of graphs has bounded rank-depth if and only if for every CMSO<sub>1</sub> transduction  $\tau$ , there exists an integer t such that  $P_t \notin \tau(\mathcal{G})$ , which was conjectured by Ganian, Hliněný, Nešetřil, Obdržálek, and Ossona de Mendez [7].

If we apply the same proof for bipartite graphs, then we prove the following theorem on pivot-minors of graphs. Pivot-minors are more restricted in a sense that every pivot-minor of a graph is a vertex-minor but not every vertex-minor is a pivot-minor. This theorem allows us to deduce a corollary for binary matroids of large branch-depth.

**Theorem 1.3.** For every positive integer t, there exists an integer N(t) such that every bipartite graph of rank-depth at least N(t) contains a pivot-minor isomorphic to  $P_t$ .

The paper is organized as follows. In Section 2, we review vertex-minors and rank-depth and prove a few useful properties related to rank-depth. In Section 3, we present the proof of Theorem 1.2. In Section 4, we obtain Theorem 1.3 and discuss its consequence to binary matroids of large branch-depth. Finally, in Section 5 we conclude the paper by giving some remarks on linear  $\chi$ -boundedness of graphs with no  $P_t$  vertex-minors.

# 2 Preliminaries and basic lemmas

All graphs in this paper are simple, meaning that neither loops nor parallel edges are allowed. For two sets X and Y, we write  $X\Delta Y$  for  $(X\backslash Y)\cup (Y\backslash X)$ .

Let G be a graph. We write V(G) and E(G) for the vertex set and the edge set of G, respectively. For a vertex v of G, we write  $N_G(v)$  to denote the set of all neighbors of v in G. For a vertex v of G, let G-v denote the graph obtained from G by removing v and all edges incident with v. For an edge e of G, let G-e denote the graph obtained from G by removing e. For a vertex subset G of G, we write G[S] for the subgraph of G induced by G. We write G for the complement of G; that is, G and G are adjacent in G if and only if they are not adjacent in G.

We write A(G) for the adjacency matrix of G over the binary field, that is, the  $V(G) \times V(G)$  matrix over the binary field such that the (x,y)-entry is one if  $x \neq y$  and x is adjacent to y in G, and zero otherwise. For an  $X \times Y$  matrix M and  $X' \subseteq X$ ,  $Y' \subseteq Y$ , we write M[X', Y'] for the  $X' \times Y'$  submatrix of M.




Figure 1: The graph  $K_4 \square K_4$ .

Let  $P_n$  denote the path on n vertices, and let  $K_n$  denote the complete graph on n vertices. The *radius* of a tree is the minimum r such that there is a node having distance at most r from every node.

For two *n*-vertex graphs G and H with fixed orderings  $\{v_1, v_2, \ldots, v_n\}$  and  $\{w_1, w_2, \ldots, w_n\}$  on their respective vertex sets, let  $G \boxtimes H$  be the graph with vertex set  $V(G) \cup V(H)$  such that  $(G \boxtimes H)[V(G)] = G$ ,  $(G \boxtimes H)[V(H)] = H$ , and for all  $i, j \in \{1, 2, \ldots, n\}$ ,  $v_i w_j \in E(G \boxtimes H)$  if and only if  $i \geqslant j$ . See Figure 1 for an example. An induced subgraph isomorphic to  $G \boxtimes H$  for some G and H is called a *semi-induced half-graph* in [19].

#### 2.1 Vertex-minors

For a vertex v of a graph G, local complementation at v is an operation which results in a new graph G \* v on V(G) such that

$$E(G * v) = E(G)\Delta\{xy : x, y \in N_G(v), x \neq y\}.$$

For an edge uv of a graph G, the operation of pivoting uv, denoted  $G \wedge uv$ , is defined as  $G \wedge uv := G * u * v * u$ . See Oum [21] for further background and properties of local complementation and pivoting. In particular, note that if G is bipartite, then so is  $G \wedge uv$ .

A graph H is locally equivalent to G if H can be obtained from G by a sequence of local complementations. A graph H is pivot equivalent to G if H can be obtained from G by a sequence of pivots. A graph H is a vertexminor of G if H is an induced subgraph of a graph that is locally equivalent to G. Finally, a graph H is a pivot-minor of G if H is an induced subgraph of a graph that is pivot equivalent to G.

For a subset S of V(G), let  $\rho_G(S)$  be the rank of the  $S \times (V(G)\backslash S)$  submatrix of A(G). This function is called the *cut-rank* function of G. It is easy to show that the cut-rank function is invariant under taking local complementations, again see Oum [21]. Thus we have the following fact.

**Lemma 2.1.** If H is a vertex-minor of G and  $X \subseteq V(G)$ , then

$$\rho_H(X \cap V(H)) \leq \rho_G(X)$$
.

The following lemmas will be used to find a long path.

**Lemma 2.2** (Kim, Kwon, Oum, and Sivaraman [12, Lemma 5.6]). The graph  $K_n \boxtimes \overline{K_n}$  has a pivot-minor isomorphic to  $P_{n+1}$ .

**Lemma 2.3** (Kwon and Oum [14, Lemma 2.8]). The graph  $\overline{K_n} \boxtimes \overline{K_n}$  has a pivot-minor isomorphic to  $P_{2n}$ .

# 2.2 Rank-depth

We now review the notion of rank-depth, which was introduced by DeVos, Kwon, and Oum [6]. A decomposition of a graph G is a pair  $(T, \sigma)$  of a tree T and a bijection  $\sigma$  from V(G) to the set of leaves of T. The radius of a decomposition  $(T, \sigma)$  is the radius of the tree T. For a non-leaf node  $v \in V(T)$ , the components of the graph T - v give rise to a partition  $\mathcal{P}_v$  of V(G) by  $\sigma$ . The width of v is defined to be

$$\max_{\mathcal{P}' \subseteq \mathcal{P}_v} \rho_G \left( \bigcup_{X \in \mathcal{P}'} X \right).$$

The width of the decomposition  $(T, \sigma)$  is the maximum width of a non-leaf node of T. We say that a decomposition  $(T, \sigma)$  is a (k, r)-decomposition of G if the width is at most k and the radius is at most r. The rank-depth of a graph G is the minimum integer k such that G admits a (k, k)-decomposition. If |V(G)| < 2, then there is no decomposition and the rank-depth is zero. Note that every tree in a decomposition has radius at least one and therefore the rank-depth of a graph is at least one if  $|V(G)| \ge 2$ .

By Lemma 2.1, it is easy to see the following.

**Lemma 2.4** (DeVos, Kwon, and Oum [6]). If H is a vertex-minor of G, then the rank-depth of H is at most the rank-depth of G.

The next two lemmas will serve as a base case for induction in the proof of Theorem 1.2.

**Lemma 2.5.** Let G be a graph of rank-depth m. Then G has a connected component of rank-depth at least m-1.

*Proof.* If m < 2, then it is trivial, as the one-vertex graph has rank-depth zero. Thus, we may assume that  $m \ge 2$ .

Suppose for contradiction that every connected component of G has rank-depth at most m-2. Let  $C_1, C_2, \ldots, C_t$  be the connected components of G. For each  $i \in \{1, 2, \ldots, t\}$ ,

- if  $C_i$  contains at least two vertices, then we take an (m-2, m-2)decomposition  $(T_i, \sigma_i)$  where  $r_i$  is a node of  $T_i$  having distance at most m-2 to every node of  $T_i$ , and
- if  $C_i$  consists of one vertex, then let  $T_i$  be the one-node graph on  $\{r_i\}$  and let  $\sigma_i: V(C_i) \to \{r_i\}$  be the uniquely possible function.

We obtain a new decomposition  $(T, \sigma)$  of G by taking the disjoint union of  $T_i$ 's and adding a new node r and adding edges  $rr_i$  for all  $i \in \{1, 2, ..., t\}$ . For every vertex v of G, define  $\sigma(v) = \sigma_i(v)$  if v is a vertex of  $C_i$ . Then  $(T, \sigma)$  has depth at most m - 1 and width at most m - 2. This contradicts the assumption that G has rank-depth m.

We conclude that G has a connected component of rank-depth at least m-1.

The following lemma can be proven similarly to Lemma 2.5. For a graph G of rank-depth m and a non-empty vertex set A, it is easy to check that G-A has rank-depth at least m-|A|, and by Lemma 2.5, G-A has a connected component of rank-depth at least m-|A|-1. But, by a direct argument, we can guarantee that there is a connected component of G-A of rank-depth at least m-|A|. We include the full proof for completeness.

**Lemma 2.6.** Let G be a graph of rank-depth m and A be a non-empty proper subset of V(G). Then G - A has a connected component of rank-depth at least m - |A|.

*Proof.* If  $|A| \ge m$ , then any connected component has rank-depth at least zero. Thus, we may assume that |A| < m. This implies that  $m \ge 2$  as A is non-empty.

Suppose for contradiction that every connected component of G-A has rank-depth at most m-|A|-1. Let  $C_1, C_2, \ldots, C_t$  be the connected components of G-A. For each  $i \in \{1, 2, \ldots, t\}$ ,

• if  $C_i$  contains at least two vertices, then we take an (m-|A|-1, m-|A|-1)-decomposition  $(T_i, \sigma_i)$  where  $r_i$  is a node of  $T_i$  having distance at most m-|A|-1 to every node of  $T_i$ , and

• if  $C_i$  consists of one vertex, we set  $T_i$  to be the one-node graph on  $\{r_i\}$  and let  $\sigma_i: V(C_i) \to \{r_i\}$  be the uniquely possible function.

We obtain a new decomposition  $(T, \sigma)$  of G by taking the disjoint union of  $T_i$ 's and adding a new node r and adding edges  $rr_i$  for all  $i \in \{1, 2, \ldots, t\}$ , and additionally appending |A| leaves to r and assigning each vertex of A to a distinct leaf with the map  $\sigma$ . For every vertex v of G - A, define  $\sigma(v) = \sigma_i(v)$  if v is a vertex of  $C_i$ . Then  $(T, \sigma)$  has depth at most m - |A| and width at most m - 1. Because  $|A| \ge 1$ , this contradicts the assumption that G has rank-depth m.

We conclude that G-A has a connected component of rank-depth at least m-|A|.

**Lemma 2.7.** Let m and d be positive integers. Let G be a graph with a vertex partition (A, B) such that connected components of G[A] and G[B] have rank-depth at most m and  $\rho_G(A) \leq d$ . Then G has rank-depth at most m+d+1.

*Proof.* Let  $C_1, \ldots, C_p$  be the connected components of G[A], and  $D_1, \ldots, D_q$  be the connected components of G[B]. For each  $i \in \{1, 2, \ldots, p\}$ ,

- if  $C_i$  contains at least two vertices, then we take an (m, m)-decomposition  $(T_i, \sigma_i)$  where  $r_i$  is a node of  $T_i$  having distance at most m to every node of  $T_i$ , and
- if  $C_i$  consists of one vertex, then set  $T_i$  as the one-node graph on  $\{r_i\}$  and  $\sigma_i: V(C_i) \to \{r_i\}$  as the uniquely possible function.

Similarly, we define  $(F_j, \mu_j)$  for each  $D_j$  where  $f_j$  is a node of  $F_j$  having distance at most m to every node of  $F_j$ .

Now, we obtain a new decomposition  $(T, \sigma)$  of G as follows. Let T be the tree obtained by taking the disjoint union of all of  $T_i$ 's and  $F_j$ 's, adding new vertices x and y, an edge xy, edges  $xr_i$  for all  $i \in \{1, \ldots, p\}$ , and edges  $yf_j$  for all  $j \in \{1, \ldots, q\}$ . Define  $\sigma(v) = \sigma_i(v)$  if v is a vertex of  $C_i$ , and  $\sigma(v) = \mu_j(v)$  if v is a vertex of  $D_j$ . Then  $(T, \sigma)$  has depth at most m + 2 and width at most m + d. Because  $d \ge 1$ , G has rank-depth at most  $\max\{m + 2, m + d\} \le m + d + 1$ .

# 2.3 Rank-width

We now review the definition of rank-width. A rank-decomposition of a graph G is a pair (T, L) of a tree T whose vertices each have degree either

one or three, and a bijection L from V(G) to the set of leaves of T. The width of an edge e of T is the cut-rank in G of the set of all leaves assigned to one of the components of T-e. The width of the rank-decomposition (T,L) is the maximum width of an edge of T. Finally, the rank-width of G is the minimum width over all rank-decompositions of G. Graphs with at most one vertex do not admit rank-decompositions and we define their rank-width to be zero.

# 3 The proof

We write R(n;k) to denote the minimum number N such that every coloring of the edges of  $K_N$  with k colors induces a monochromatic complete subgraph on n vertices. The classical theorem of Ramsey [23] implies that R(n;k) exists.

The following lemma is well known. We include its proof for the sake of completeness.

**Lemma 3.1.** Let G be a graph of rank-width at most q and let  $M \subseteq V(G)$ . If  $|M| \ge 3k + 1$  for a positive integer k, then there is a vertex partition (X,Y) of G such that  $\rho_H(X) \le q$  and  $\min(|M \cap X|, |M \cap Y|) > k$ .

Proof. Suppose that there is no such vertex partition. Let (T,L) be a rank-decomposition of width at most q. For each edge uv of T, let us orient e towards v if the component of T-e containing u has at most k vertices in L(M). By the assumption, every edge is oriented. Since T is acyclic, there is a node w of T such that all edges of T incident with w are oriented towards w. But this implies that  $|M| \leq 3k$ , a contradiction.  $\square$ 

For a path P with an endpoint x and a graph H and a non-empty subset of vertices  $S \subseteq V(H)$ , we denote by (P, x) + (H, S) the graph obtained from the disjoint union of P and H by adding all edges between x and S. We now prove our main proposition; Theorem 1.2 will follow quickly after.

**Proposition 3.2.** For all positive integers a, b, t, q, there exists an integer f(a, b, t, q) such that every graph of rank-width at most q and rank-depth at least f(a, b, t, q) has a vertex-minor isomorphic to either  $P_t$  or  $(P_a, x) + (H, S)$  where x is an endpoint of  $P_a$ , H is a connected graph of rank-depth at least b, and S is a non-empty subset of V(H).

*Proof.* For all positive integers b, t, q, we set

$$f(1, b, t, q) := b + 2,$$




Figure 2: The graph  $G_1[A_1 \cup \cdots \cup A_r \cup V(H_r)]$ , when r=3 and a=5.

and for  $a \ge 2$ , we set

$$u := \max(3 \cdot (2^{q} - 1) + 1, t - 1),$$

$$r := R(u + 1; 2^{a - 1}),$$

$$g_{i} := \begin{cases} b + q + 2 & \text{if } i = r, \\ f(a - 1, g_{i + 1}, t, q) & \text{if } i \in \{0, 1, 2, \dots, r - 1\}, \end{cases}$$

$$f(a, b, t, q) := g_{0}.$$

We prove the proposition by induction on a. Let G be a graph whose rank-depth is at least f(a, b, t, q) and rank-width is at most q. If a = 1, then it has a component G' of rank-depth at least b + 1 by Lemma 2.5. Let  $v \in V(G')$ . By Lemma 2.6, G' - v has a connected component H of rank-depth at least b. So,  $(G'[\{v\}], v) + (H, N_G(v) \cap V(H))$  is the second outcome.

Thus, we may assume that  $a \ge 2$ . Suppose that G has no vertex-minor isomorphic to  $P_t$ . We claim that G contains the second outcome.

Let  $H_0 := G$ . Observe that  $H_0$  has rank-depth at least  $f(a, b, t, q) = g_0$ . For  $i \in \{1, 2, ..., r\}$ , we recursively find tuples  $(A_i, x_i, H_i, S_i)$  from  $H_{i-1}$  such that

- $A_i$  is isomorphic to  $P_{a-1}$  and  $x_i$  is an endpoint of  $A_i$ ,
- $H_i$  is a connected graph of rank-depth at least  $g_i$ ,
- $S_i$  is a non-empty subset of  $V(H_i)$ , and
- $(A_i, x_i) + (H_i, S_i)$  is a vertex-minor of  $H_{i-1}$ .

Let  $i \in \{1, 2, ..., r\}$  and assume that  $H_{i-1}$  is a given graph of rank-depth at least  $g_{i-1}$ . Then by the induction hypothesis,  $H_{i-1}$  has a vertex-minor  $(A_i, x_i) + (H_i, S_i)$  where  $A_i$  is isomorphic to  $P_{a-1}$ ,  $x_i$  is an endpoint of  $A_i$ ,

 $H_i$  is a connected graph of rank-depth at least  $g_i$ , and  $S_i$  is a non-empty subset of  $V(H_i)$ . By the choice of functions  $g_0, g_1, \ldots, g_r$ , we can obtain the tuples for all  $i \in \{1, 2, \ldots, r\}$ .

Observe that for i < j, no vertex in  $V(A_i) \setminus \{x_i\}$  has a neighbor in  $H_i$ , and therefore, the sequence of local complementations to obtain  $(A_j, x_j) + (H_j, S_j)$  from  $H_{j-1}$  does not change previous paths  $A_1, \ldots, A_{j-1}$ , but may change the edges between  $x_1, x_2, \ldots, x_{j-1}$ .

By definition,  $H_r$  is connected and has rank-depth at least  $g_r$ . Let  $G_1$  be the graph obtained from G by following the sequence of local complementations to obtain  $(A_1, x_1) + (H_1, S_1), \ldots, (A_r, x_r) + (H_r, S_r)$ . See Figure 2 for a depiction. Note that  $G_1[V(H_i) \cup \{x_i\}]$  is connected for each i, as  $H_i$  is connected,  $S_i$  is non-empty, and we apply local complementations only inside  $H_i$  to obtain later  $H_j$ 's.

If for some  $i \in \{1, 2, ..., r\}$ ,  $N_{G_1}(x_i) \cap V(H_r) = \emptyset$ , then by taking a shortest path from  $x_i$  to  $V(H_r)$  in the graph  $G_1[V(H_i) \cup \{x_i\}]$ , we can directly obtain the second outcome. So, we may assume that each of  $\{x_1, x_2, ..., x_r\}$  has a neighbor in  $V(H_r)$ .

Note that for i < j, only the endpoint  $x_i$  in  $A_i$  can have a neighbor in  $A_j$  in  $G_1$ , and therefore, there are  $2^{a-1}$  possible ways of having edges between  $A_i$  and  $A_j$  in  $G_1$ . Since  $r = R(u+1; 2^{a-1})$ , by applying the theorem of Ramsey, we deduce that there exists a subset  $W \subseteq \{1, 2, \ldots, r\}$  of size u+1 such that for all i < j with  $i, j \in W$ ,  $\{\ell : \text{the } \ell\text{-th vertex of } A_j \text{ is adjacent to } x_i \text{ in } G_1\}$  are identical

If  $x_i$  has a neighbor in  $V(A_j - x_j)$  in  $G_1$  for some i < j with  $i, j \in W$ , then  $G_1$  has  $\overline{K_u} \boxtimes \overline{K_u}$  or  $\overline{K_u} \boxtimes K_u$  as an induced subgraph. Since  $u \ge t-1$ , by Lemmas 2.2 and 2.3,  $G_1$  contains a pivot-minor isomorphic to  $P_t$ , contradicting the assumption. So, for all i < j with  $i, j \in W$ ,  $x_i$  has no neighbors in  $V(A_j - x_j)$ .

Note that  $\{x_i : i \in W\}$  is an independent set or a clique in  $G_1$ . If it is an independent set, then for some  $i' \in W$ , we set

•  $G_2 := G_1$  and  $W' := W \setminus \{i'\}.$ 

If  $\{x_i : i \in W\}$  is a clique, then we choose a vertex  $x_{i'}$  for some  $i' \in W$  and locally complement at  $x_{i'}$ . Then  $\{x_i : i \in W \setminus \{i'\}\}$  becomes an independent set. We set

•  $G_2 := G_1 * x_{i'}$  and  $W' := W \setminus \{i'\}.$ 

Let  $M := \{x_i : i \in W'\}$  and  $H := G_2[V(H_r) \cup M \cup \{x_{i'}\}].$ 

By definition, H is locally equivalent to the graph  $G_1[V(H_r) \cup M \cup \{x_{i'}\}]$ . Thus, as the latter is connected, H is also connected. Similarly, since  $H_r =$   $G_1[V(H_r)]$  has rank-depth at least  $g_r$ , H has rank-depth at least  $g_r$ . Also, note that H has rank-width at most q and M is an independent set of size  $u \ge 3 \cdot (2^q - 1) + 1$  in H. Thus, by Lemma 3.1, H admits a vertex partition (X,Y) such that  $|M \cap X| > 2^q - 1$ ,  $|M \cap Y| > 2^q - 1$ , and  $\rho_H(X) \le q$ .

Since H has rank-depth at least  $g_r = b + q + 2$  and  $\rho_H(X) \leq q$ , by Lemma 2.7, H[X] or H[Y] has a connected component of rank-depth at least b+1. Without loss of generality, we assume that H[X] has a connected component Q of rank-depth at least b+1.

Now, if  $M \cap Y$  has a vertex  $x_i$  that has no neighbor in Q, then by taking a shortest path from  $x_i$  to Q in H, along with  $A_i$ , we can find the second outcome.

Thus, we may assume that in H, all vertices in  $M \cap Y$  have a neighbor in Q. Since  $\rho_H(X) \leq q$ , there are at most  $2^q - 1$  distinct non-zero rows in the matrix  $A(H)[M \cap Y, V(Q)]$ . As  $|M \cap Y| \geq 2^q$ , by the pigeon-hole principle, H has two vertices  $x_{i_1}$  and  $x_{i_2}$  in  $M \cap Y$  for some  $i_1, i_2 \in W'$  that have the same neighborhood in Q.

First assume that  $x_{i_1}$  has exactly one neighbor in Q, say w. As Q has rank-depth at least b+1, Q-w has a connected component Q' having rank-depth at least b by Lemma 2.6. Then

$$(G_2[V(A_{i_1}) \cup \{w\}], w) + (Q', N_{G_2}(w) \cap V(Q'))$$

is the required second outcome. So, we may assume that  $x_{i_1}$  has at least two neighbors in Q. Let w be a neighbor of  $x_{i_1}$  in Q.

Since  $x_{i_1}$  and  $x_{i_2}$  have the same neighborhood in Q and they are not adjacent, if we pivot  $x_{i_2}w$ , then the edges between  $x_{i_1}$  and  $N_H(x_{i_1}) \cap V(Q)$  are removed and  $x_{i_2}$  becomes the unique neighbor of  $x_{i_1}$  in  $V(Q) \cup \{x_{i_2}\}$ . Note that  $G_2[V(Q) \cup \{x_{i_2}\}]$  is connected, and thus  $(G_2 \wedge x_{i_2}w)[V(Q) \cup \{x_{i_2}\}]$  is also connected. As Q has rank-depth at least b+1,  $(G_2 \wedge x_{i_2}w)[V(Q) \cup \{x_{i_2}\}] - x_{i_2}$  has a connected component Q' that has rank-depth at least b. Then

$$((G_2 \wedge x_{i_2}w)[V(A_{i_1}) \cup \{x_{i_2}\}], x_{i_2}) + (Q', N_{G_2 \wedge x_{i_2}w}(x_{i_2}) \cap V(Q'))$$

is the second outcome. This proves the proposition.

Proposition 3.2 implies the following result.

**Theorem 3.3.** For all positive integers t and q, there exists an integer F(t,q) such that every graph of rank-width at most q and rank-depth at least F(t,q) contains a vertex-minor isomorphic to  $P_t$ .

*Proof.* We take F(t,q) := f(t-1,1,t,q) where f is the function in Proposition 3.2.

A circle graph is the intersection graph of chords on a circle. It is easy to see that  $P_t$  is a circle graph. We can derive Theorem 1.2 by taking  $q = \beta(P_t)$  from the following recent theorem.

**Theorem 3.4** (Geelen, Kwon, McCarty, and Wollan [10]). For every circle graph H, there exists an integer  $\beta(H)$  such that every graph of rank-width more than  $\beta(H)$  contains a vertex-minor isomorphic to H.

**Theorem 1.2.** For every positive integer t, there exists an integer N(t) such that every graph of rank-depth at least N(t) contains a vertex-minor isomorphic to  $P_t$ .

Proof. We take  $N(t) := F(t, \beta(P_t))$  where  $\beta$  is the function given in Theorem 3.4 and F is the function from Theorem 3.3. If a graph has rank-width more than  $\beta(P_t)$ , then by Theorem 3.4, it contains a vertex-minor isomorphic to  $P_t$ . So, we may assume that a graph has rank-width at most  $\beta(P_t)$ . Then by Theorem 3.3, it contains a vertex-minor isomorphic to  $P_t$ .

# 4 Pivot-minors

We can prove a stronger result on bipartite graphs, by slightly modifying the proof of Proposition 3.2. Suppose that a given graph G is bipartite in the proof of Proposition 3.2. The only place that we have to apply local complementation instead of pivoting is when the set  $\{x_i : i \in W\}$  is a clique, and we want to change it into an independent set. But if G is bipartite, then the obtained set  $\{x_i : i \in W\}$  has no triangle, and so it is an independent set since  $|W| \ge 3$ . Therefore, we can proceed only with pivoting. For bipartite graphs, we can use the following theorem due to Oum [21], obtained as a consequence of the grid theorem for binary matroids [9].

**Theorem 4.1** (Oum [21]). For every bipartite circle graph H, there exists an integer  $\gamma(H)$  such that every bipartite graph of rank-width more than  $\gamma(H)$  contains a pivot-minor isomorphic to H.

Thus we deduce the following theorem for bipartite graphs.

**Theorem 1.3.** For every positive integer t, there exists an integer N(t) such that every bipartite graph of rank-depth at least N(t) contains a pivot-minor isomorphic to  $P_t$ .



Figure 3: The fan graph  $F_5$ .

Theorem 1.3 allows us to obtain the following corollary for binary matroids, solving a special case of a conjecture of DeVos, Kwon, and Oum [6] on general matroids. We need a few terms to state the corollary. The branch-depth of a matroid is defined analogously to the definition of the rank-depth obtained by replacing the cut-rank function with the matroid connectivity function [6]. Let  $F_t$  be the fan graph, that is the union of  $P_t$  with one vertex adjacent to all vertices of  $P_t$ , see Figure 3. As usual,  $M(F_t)$  denotes the cycle matroid of  $F_t$ .

Corollary 4.2. For every positive integer t, there exists an integer N(t) such that every binary matroid of branch-depth at least N(t) contains a minor isomorphic to  $M(F_t)$ .

Proof. It is known [2, 21] that the connectivity function of a binary matroid is equal to the cut-rank function of a corresponding bipartite graph, called a fundamental graph. Furthermore for two binary matroids M and N, if N is connected, and a fundamental graph of N is a pivot-minor of a fundamental graph of M, then either N or  $N^*$  is a minor of M, see Oum [21, Corollary 3.6]. Since  $(M(F_t))^*$  has a minor isomorphic to  $M(F_{t-1})$ , we deduce the corollary from Theorem 1.3, because the path graph  $P_{2t-1}$  is a fundamental graph of  $M(F_t)$ .

We show that the class  $\{K_n \boxtimes K_n : n \ge 1\}$  has unbounded rank-depth, while for every positive integer n,  $K_n \boxtimes K_n$  has no pivot-minor isomorphic to  $P_5$ . It implies that contrary to Theorem 1.3, the class of graphs having no  $P_n$  pivot-minor has unbounded rank-depth for each  $n \ge 5$ .

Kwon and Oum [16, Lemma 6.5] showed that for every integer  $n \ge 2$ ,  $K_n \boxtimes K_n$  contains a vertex-minor isomorphic to  $P_{2n-2}$ . Thus,  $\{K_n \boxtimes K_n : n \ge 1\}$  has unbounded rank-depth.

Now, we show that for  $n \ge 1$ ,  $K_n \square K_n$  has no pivot-minor isomorphic to  $P_5$ . We prove a stronger statement that  $K_n \square K_n$  has no pivot-minor isomorphic to  $K_{1,3}$ . Dabrowski et al. [5] characterized the class of graphs having no pivot-minor isomorphic to  $K_{1,3}$  in terms of forbidden induced subgraphs. See Figure 4 for bull,  $W_4$ , and  $\overline{BW_3}$ .




Figure 4: The graphs bull,  $W_4$ , and  $\overline{BW_3}$ .

**Theorem 4.3** (Dabrowski et al. [5]). A graph has a pivot-minor isomorphic to  $K_{1,3}$  if and only if it has an induced subgraph isomorphic to one of  $K_{1,3}$ ,  $P_5$ , bull,  $W_4$ , and  $\overline{BW_3}$ .

**Lemma 4.4.** For  $n \ge 1$ ,  $K_n \boxtimes K_n$  has no induced subgraph isomorphic to one of  $K_{1,3}$ ,  $P_5$ , bull,  $W_4$ , and  $\overline{BW_3}$ .

*Proof.* As the maximum size of an independent set in  $K_n \square K_n$  is 2,  $K_n \square K_n$  has no induced subgraph isomorphic to one of  $K_{1,3}$ ,  $P_5$ , and bull.

Also  $K_n \square K_n$  has no induced cycle of length 4 because such a cycle should contain two vertices in each  $K_n$  but the edges between two  $K_n$ 's have no induced matching of size 2. Therefore, it has no induced subgraph isomorphic to  $W_4$  or  $\overline{BW_3}$ .

By Theorem 4.3 and Lemma 4.4,  $K_n \square K_n$  has no pivot-minor isomorphic to  $K_{1,3}$ , and to  $P_5$ . Thus, for all  $n \ge 5$ , the class of graphs having no  $P_n$  pivot-minor includes  $\{K_n \square K_n : n \ge 1\}$ , which has unbounded rank-depth. It may be interesting to see whether every graph with sufficiently large rank-depth contains either  $P_n$  or  $K_n \square K_n$  as a pivot-minor. We leave it as an open question.

**Question 1.** Does there exist a function f such that for every n, every graph with rank-depth at least f(n) contains a pivot-minor isomorphic to  $P_n$  or  $K_n \boxtimes K_n$ ?

# 5 Concluding remarks

#### 5.1 Linear $\chi$ -boundedness

We define linear rank-width. For an ordering  $(v_1, v_2, \ldots, v_n)$  of the vertex set of a graph G, its width is defined as the maximum of  $\rho_G(\{v_1, \ldots, v_i\})$  for all  $i \in \{1, 2, \ldots, n-1\}$ , and the linear rank-width of G is defined as the minimum width of all orderings of G. If |V(G)| < 2, then the linear rank-width of G is defined as 0.

Graphs of bounded rank-depth have bounded linear rank-width, which was already known through the notions of shrub-depth and linear clique-width [7]. Kwon and Oum [15] proved it directly as follows.

**Proposition 5.1** (Kwon and Oum [15]). Every graph of rank-depth k has linear rank-width at most  $k^2$ .

We write  $\chi(G)$  to denote the chromatic number of G and  $\omega(G)$  to denote the maximum size of a clique of G. A class  $\mathcal{C}$  of graphs is  $\chi$ -bounded if there is a function f such that  $\chi(H) \leq f(\omega(H))$  for all induced subgraphs H of a graph in  $\mathcal{C}$ . In addition, if f can be taken as a polynomial function, then  $\mathcal{C}$  is polynomially  $\chi$ -bounded. If f can be taken as a linear function, then  $\mathcal{C}$  is linearly  $\chi$ -bounded.

**Proposition 5.2** (Nešetřil, Ossona de Mendez, Rabinovich, and Siebertz [19]). For every positive integer r, there exists an integer c(r) such that for every graph G of linear rank-width at most r,

$$\chi(G) \leqslant c(r) \,\omega(G).$$

By combining Proposition 5.1 and Proposition 5.2, we can prove the following, which answers a previous question by Kim, Kwon, Oum, and Sivaraman [12].

**Theorem 5.3.** For every positive integer t, the class of graphs with no vertex-minor isomorphic to  $P_t$  is linearly  $\chi$ -bounded.

We remark that there is an alternative way to prove Theorem 5.3 without using linear rank-width. First, DeVos, Kwon, and Oum [6, Lemma 4.10] showed that if a graph has rank-depth k, then it has an (a, k)-shrubbery where

$$a = (1 + o(1))2^{(2^{2k+1}(2^{2k+2}-1)+1)k/2}.$$

(Please see [6] for the definition of an (a,k)-shrubbery.) Lemma 2.16 of Nešetřil, Ossona de Mendez, Rabinovich, and Siebertz [19] states that every class of bounded shrub-depth can be partitioned into bounded number of vertex-disjoint induced subgraphs, each of which is a cograph. Its (short and easy) proof shows that a graph with an (a,k)-shrubbery can be partitioned into at most a vertex-disjoint induced subgraphs, each of which is a cograph. Since cographs are perfect, we deduce that if G has rank-depth at most k, then  $\chi(G) \leq \omega(G)(1 + o(1))2^{(2^{2k+1}(2^{2k+2}-1)+1)k/2}$ .




Figure 5: Obstructions for being a vertex-minor of a path.

# 5.2 When does the class of $\mathcal{H}$ -vertex-minor-free graphs have bounded rank-depth?

For a set  $\mathcal{H}$  of graphs, we say that G is  $\mathcal{H}$ -minor-free if G has no minor isomorphic to a graph in  $\mathcal{H}$ , and G is  $\mathcal{H}$ -vertex-minor-free if G has no vertex-minor isomorphic to a graph in  $\mathcal{H}$ . Robertson and Seymour [24] showed that  $\mathcal{H}$ -minor-free graphs have bounded tree-width if and only if  $\mathcal{H}$  contains a planar graph. As an analogue, Geelen, Kwon, McCarty, and Wollan [10] showed that  $\mathcal{H}$ -vertex-minor-free graphs have bounded rank-width if and only if  $\mathcal{H}$  contains a circle graph. Interestingly, Theorem 1.2 allows us to characterize the classes  $\mathcal{H}$  such that  $\mathcal{H}$ -vertex-minor-free graphs have bounded rank-depth. This is due to the following theorem; the equivalence between (a) and (b) was shown by Kwon and Oum [13] and the equivalence between (a) and (c) was shown by Adler, Farley, and Proskurowski [1].

**Theorem 5.4** (Kwon and Oum [13]; Adler, Farley, and Proskurowski [1]). Let H be a graph. The following are equivalent.

- (a) H has linear rank-width at most one.
- (b) H is a vertex-minor of a path.
- (c) H has no vertex-minor isomorphic to  $C_5$ , N, or Q in Figure 5.

We define linear rank-width in the next subsection. Here, we only need the fact that linear rank-width does not increase when we take vertex-minors and that paths have linear rank-width 1 and arbitrary large rank-depth to deduce the following corollary from Theorems 5.4 and 1.2.

Corollary 5.5. Let  $\mathcal{H}$  be a set of graphs. Then the following are equivalent.

- (a) The class of H-vertex-minor-free graphs has bounded rank-depth.
- (b) H contains a graph of linear rank-width at most one.
- (c)  $\mathcal{H}$  contains a graph with no vertex-minor isomorphic to  $C_5$ , N, or Q.

**Acknowledgement.** The authors would like to thank anonymous reviewers for their helpful suggestions.

# References

- [1] I. Adler, A. M. Farley, and A. Proskurowski. Obstructions for linear rank-width at most 1. *Discrete Appl. Math.*, 168:3–13, 2014.
- [2] A. Bouchet. Connectivity of isotropic systems. In Combinatorial Mathematics: Proceedings of the Third International Conference (New York, 1985), volume 555 of Ann. New York Acad. Sci., pages 81–93. New York Acad. Sci., New York, 1989.
- [3] B. Courcelle and S. Olariu. Upper bounds to the clique width of graphs. Discrete Appl. Math., 101(1-3):77–114, 2000.
- [4] B. Courcelle and S. Oum. Vertex-minors, monadic second-order logic, and a conjecture by Seese. *J. Combin. Theory Ser. B*, 97(1):91–126, 2007.
- [5] K. K. Dabrowski, F. Dross, J. Jeong, M. M. Kanté, O. Kwon, S. Oum, and D. Paulusma. Computing small pivot-minors. In *Graph-theoretic concepts in computer science*, volume 11159 of *Lecture Notes in Comput. Sci.*, pages 125–138. Springer, Cham, 2018.
- [6] M. DeVos, O. Kwon, and S. Oum. Branch-depth: Generalizing tree-depth of graphs. *European J. Combin.*, 90:103186, 23, 2020.
- [7] R. Ganian, P. Hliněný, J. Nešetřil, J. Obdržálek, and P. Ossona de Mendez. Shrub-depth: Capturing height of dense graphs. *Log. Methods Comput. Sci.*, 15(1):7:1–7:25, 2019.
- [8] R. Ganian, P. Hliněný, J. Nešetřil, J. Obdržálek, P. Ossona de Mendez, and R. Ramadurai. When trees grow low: shrubs and fast MSO<sub>1</sub>. In Mathematical foundations of computer science 2012, volume 7464 of Lecture Notes in Comput. Sci., pages 419–430. Springer, Heidelberg, 2012.
- [9] J. Geelen, B. Gerards, and G. Whittle. Excluding a planar graph from GF(q)-representable matroids. *J. Combin. Theory Ser. B*, 97(6):971–998, 2007.

- [10] J. Geelen, O. Kwon, R. McCarty, and P. Wollan. The grid theorem for vertex-minors. *J. Combin. Theory Ser. B*, 2020. https://doi.org/10.1016/j.jctb.2020.08.004.
- [11] P. Hliněný, O. Kwon, J. Obdržálek, and S. Ordyniak. Tree-depth and vertex-minors. *European J. Combin.*, 56:46–56, 2016.
- [12] R. Kim, O. Kwon, S. Oum, and V. Sivaraman. Classes of graphs with no long cycle as a vertex-minor are polynomially χ-bounded. J. Combin. Theory, Ser. B, 140:372–386, 2020.
- [13] O. Kwon and S. Oum. Graphs of small rank-width are pivot-minors of graphs of small tree-width. *Discrete Appl. Math.*, 168:108–118, 2014.
- [14] O. Kwon and S. Oum. Unavoidable vertex-minors in large prime graphs. European J. Combin., 41:100–127, 2014.
- [15] O. Kwon and S. Oum. Graphs of bounded depth-2 rank-brittleness. *J. Graph Theory*, 2020. https://doi.org/10.1002/jgt.22619.
- [16] O. Kwon and S. Oum. Scattered classes of graphs. SIAM J. Discrete Math., 34(1):972–999, 2020.
- [17] J. Nešetřil and P. Ossona de Mendez. Tree-depth, subgraph coloring and homomorphism bounds. *European J. Combin.*, 27(6):1022–1041, 2006.
- [18] J. Nešetřil and P. Ossona de Mendez. *Sparsity*, volume 28 of *Algorithms and Combinatorics*. Springer, Heidelberg, 2012.
- [19] J. Nešetřil, P. Ossona de Mendez, R. Rabinovich, and S. Siebertz. Classes of graphs with low complexity: the case of classes with bounded linear rankwidth. *European J. Combin.*, 91:103223, 29, 2021.
- [20] J. Nešetřil and P. Ossona de Mendez. On low tree-depth decompositions. *Graphs Combin.*, 31(6):1941–1963, 2015.
- [21] S. Oum. Rank-width and vertex-minors. J. Combin. Theory Ser. B, 95(1):79–100, 2005.
- [22] S. Oum and P. Seymour. Approximating clique-width and branch-width. J. Combin. Theory Ser. B, 96(4):514–528, 2006.
- [23] F. P. Ramsey. On a problem of formal logic. *Proc. London Math. Soc.*, 30(s2):264-286, 1930.

[24] N. Robertson and P. Seymour. Graph minors. X. Obstructions to tree-decomposition. *J. Combin. Theory Ser. B*, 52(2):153–190, 1991.