

**REMARK ON A LOWER BOUND OF PERTURBATIONS IN 2D
INVISCID SHEAR FLOW IN A PERIODIC BOX**

TSUYOSHI YONEDA

ABSTRACT. Bedrossian and Masmoudi (2015) showed nonlinear stability of 2D inviscid parallel shear flows in the infinite stripe $(\mathbb{R}/2\mathbb{Z}) \times \mathbb{R}$. Inspired by their result, combining a diffeomorphism group result by Misiólek (1993), in this paper we give a lower bound of perturbations in 2D inviscid shear flow in a periodic box.

1. INTRODUCTION AND THE MAIN THEOREM

For $m \in \mathbb{N}$, let us define a stripe type of periodic box as $\mathbb{T}^2 := (\mathbb{R}/2\mathbb{Z}) \times (\mathbb{R}/(2m\mathbb{Z}))$. The incompressible Euler equations in \mathbb{T}^2 are described as follows:

$$(1.1) \quad \begin{aligned} \partial_t u + (u \cdot \nabla)u &= -\nabla p, \quad x = (x_1, x_2) \in \mathbb{T}^2, \quad t > 0, \\ \nabla \cdot u &= 0, \quad u|_{t=0} = u_0, \end{aligned}$$

where $u = u(t, x)$ is the velocity and $p = p(t, x)$ is the pressure. Throughout this paper we always handle smooth solutions, that is, we take initial velocity $u_0 \in C^\infty(\mathbb{T}^2)$. It is well known that the corresponding unique smooth solution u exists globally in time (see [3] for example). Next we define the Lagrangian flow η . For $u \in C^\infty([0, \infty) \times \mathbb{T}^2)$, let $\eta(t, x)$ be a solution to the following ODE:

$$\frac{d}{dt}\eta(t, x) = u(t, \eta(t, x)) =: u \circ \eta, \quad \eta(0, x) =: e(x) = x.$$

We can express it in Riemannian geometric language using the exponential map of the right-invariant metric (defined by the L^2 kinetic energy of the fluid) on the group $\mathcal{D}_\mu^s(\mathbb{T}^2)$ of volume-preserving diffeomorphisms of \mathbb{T}^2 of Sobolev class H^s with $s > 2$, namely $\eta(t) = \exp_e t u_0$ (see for example [2] or [4]). Clearly, η uniquely exists globally in time and is smooth. Now let us define the inviscid parallel shear flow (stationary Euler flow):

$$u_\infty(x) = (f_m(x_2), 0), \quad f_m \in C^\infty(\mathbb{R}/(2m\mathbb{Z})).$$

Here we choose f_m in the following way: First let us take $f \in C^\infty(\mathbb{R})$ with $\|f'\|_{L^\infty} < \infty$, and then we take f_m satisfying

$$(1.2) \quad \begin{cases} \lim_{m \rightarrow \infty} \sup_{-\frac{m}{2} < x_2 \leq \frac{m}{2}} (|f_m(x_2) - f(x_2)| + |f'_m(x_2) - f'(x_2)|) = 0, \\ \sup_m \|f'_m\|_{L^\infty} < \infty. \end{cases}$$

Date: November 12, 2019.

2010 Mathematics Subject Classification. Primary 76E05; Secondary 35Q31.

Key words and phrases. inviscid fluid flow, parallel shear flow.

For any $U_{in} \in C^\infty$ with $\nabla \cdot U_{in} = 0$, from now on, we consider the following family of L^2 geodesics in \mathcal{D}_μ^s , or equivalently, fluid flows in \mathbb{T}^2 , starting at the identity configuration e in the direction $u_\infty + \sigma U_{in}$:

$$\tilde{\eta}(\sigma, t) := \exp_e(t(u_\infty + \sigma U_{in})), \quad \sigma \in (-1, 1), \quad t \geq 0.$$

Note that $\partial_\sigma \tilde{\eta}(\sigma, t)|_{t=0} = 0$. Let $\tilde{u} = \tilde{u}(\sigma, t)$ be the corresponding solution to the Euler equations (1.1) for the initial data $u_\infty + \sigma U_{in}$, and let us set

$$\text{perturbation:} \quad U = (U_1, U_2) = U(\sigma, t) := \tilde{u}(\sigma, t) - u_\infty.$$

Note that $U(0, t) = 0$ ($t > 0$). In the infinite stripe case $(\mathbb{R}/2\mathbb{Z}) \times \mathbb{R}$ (formally $m = \infty$), Bedrossian and Masmoudi [1] showed the following nonlinear stability result (we rewrite their statement into our setting): For any $U_{in} \in \mathcal{G}^{\lambda_0}$ with $\|U_{in}\|_{\mathcal{G}^{\lambda_0}} \sim 1$ (for the precise definition of the Gevrey-norm $\|\cdot\|_{\mathcal{G}^{\lambda_0}}$, see [1]). In this paper, we just use the embeddings: $\|\cdot\|_{L^2} \leq \|\cdot\|_{\mathcal{G}^\lambda}$ and $\|\nabla \cdot\|_{L^\infty} \leq \|\cdot\|_{H^3} \leq \|\cdot\|_{\mathcal{G}^\lambda}$ and $\sigma > 0$, then there is \tilde{f} sufficiently smooth and appropriately decaying (depending on σ , but controlled by $\|\tilde{f}'\|_{L^\infty} \leq \|\tilde{f}\|_{\mathcal{G}^{\lambda'}} < \sigma \ll 1$), such that if

$$u_\infty(x) = \begin{pmatrix} f(x_2) \\ 0 \end{pmatrix} \quad \text{with} \quad f(x_2) = x_2 + \tilde{f}(x_2),$$

then we have the following stability estimate:

$$(1.3) \quad \|U_1(\sigma, t)\|_{L^2} \lesssim \frac{\sigma}{\langle t \rangle} \quad \text{and} \quad \|U_2(\sigma, t)\|_{L^2} \lesssim \frac{\sigma}{\langle t \rangle^2},$$

where $\langle t \rangle := (1 + |t|^2)^{1/2}$.

On the other hand, inspired by their result, we constructed the following lower bound.

Theorem 1.1. *Let us consider the perturbation U of the shear flow $(f_m, 0)$ in $(\mathbb{R}/2\mathbb{Z}) \times (\mathbb{R}/(2m\mathbb{Z}))$ satisfying (1.2). For any $\epsilon > 0$ and $T > 0$, there is $\sigma_0 > 0$ such that if $\sigma < \sigma_0$, we have*

$$\|U_1(\sigma, t)\|_{L^2(\mathbb{T}^2)} \gtrsim \sigma \quad \text{or} \quad \|U_2(\sigma, t)\|_{L^2(\mathbb{T}^2)} \gtrsim \frac{\sigma}{\langle t \rangle^{1+\epsilon}}$$

for $m \in \mathbb{N}$ and $t \in [0, T]$. In particular, this lower bound is independent of m and T .

Remark 1. If the energy of the initial velocity $u_0 = u_\infty + \sigma U_{in}$ is not equal to the energy of the shear flow, namely, $\|u_0\|_{L^2} \neq \|u_\infty\|_{L^2}$, then we can easily see that the energy of perturbation $\|U(\sigma, t)\|_{L^2}$ is not zero for any $t > 0$. This is just due to the triangle inequality (the following is the case when $\|u_0\|_{L^2} > \|u_\infty\|_{L^2}$):

$$\|u_0\|_{L^2} = \|\tilde{u}(\sigma, t)\|_{L^2} \leq \|u_\infty\|_{L^2} + \|U(\sigma, t)\|_{L^2} \Rightarrow \|u_0\|_{L^2} - \|u_\infty\|_{L^2} \leq \|U(\sigma, t)\|_{L^2}.$$

However, this estimate cannot exclude the case $\|u_0\|_{L^2} = \|u_\infty\|_{L^2}$ (with $u_0 \neq u_\infty$).

2. PROOF OF THE MAIN THEOREM

First we show $\partial_\sigma^k U(\sigma, \cdot) \in C^\infty([0, \infty) \times \mathbb{T}^2)$ ($\sigma \in (-1, 1)$, $k = 1, 2, \dots$). In order to do so, just multiply ∂_σ^k to (1.1) on both sides (replace u with \tilde{u}), and just apply inductively the existence theory of the usual Euler equations (for the existence theory itself, see [3] example). Since $\tilde{u}(\sigma, t) = u_\infty + U(\sigma, t)$, we see

$$\partial_t \tilde{\eta} = u_\infty \circ \tilde{\eta} + U \circ \tilde{\eta}.$$

Thus by differentiation in σ on both sides, we have

$$\partial_t \partial_\sigma \tilde{\eta} = ((\nabla u_\infty) \circ \tilde{\eta}) \partial_\sigma \tilde{\eta} + (\partial_\sigma U) \circ \tilde{\eta} + (\nabla U \circ \tilde{\eta}) \partial_\sigma \tilde{\eta}$$

and then

$$(2.1) \quad \partial_t \partial_\sigma \tilde{\eta}|_{\sigma=0} = ((\nabla u_\infty) \circ \tilde{\eta}) \partial_\sigma \tilde{\eta}|_{\sigma=0} + (\partial_\sigma U) \circ \tilde{\eta}|_{\sigma=0}.$$

We see that

$$((\nabla u_\infty) \circ \tilde{\eta}) \partial_\sigma \tilde{\eta} = \begin{pmatrix} \partial_\sigma \tilde{\eta}_2 (\partial_2 f) \circ \tilde{\eta}_2 \\ 0 \end{pmatrix},$$

thus multiplying $\partial_\sigma \tilde{\eta}_2$ to the second component of (2.1), and take integration in space, we have

$$(1/2) \partial_t \|\partial_\sigma \tilde{\eta}_2(0, t)\|_{L^2}^2 \leq \|\partial_\sigma U_2(0, t)\|_{L^2} \|\partial_\sigma \tilde{\eta}_2(0, t)\|_{L^2},$$

and this means

$$\|\partial_\sigma \tilde{\eta}_2(0, t)\|_{L^2} \lesssim \int_0^t \|\partial_\sigma U_2(0, t')\|_{L^2} dt'.$$

We plug the above estimate into the first component of (2.1), we obtain

$$(1/2) \partial_t \|\partial_\sigma \tilde{\eta}_1(0, t)\|_{L^2}^2 \leq \int_0^t \|\partial_\sigma U_2(0, t')\|_{L^2} dt' \|f'_m\|_{L^\infty} \|\partial_\sigma \tilde{\eta}_1(0, t)\|_{L^2} \\ + \|\partial_\sigma U_1(0, t)\|_{L^2} \|\partial_\sigma \tilde{\eta}_1(0, t)\|_{L^2},$$

and this means

$$\|\partial_\sigma \tilde{\eta}_1(0, t)\|_{L^2} \lesssim \int_0^t \int_0^{t'} \|\partial_\sigma U_2(0, t'')\|_{L^2} dt'' dt' \|f'_m\|_{L^\infty} + t \sup_{0 < t' < t} \|\partial_\sigma U_1(0, t')\|_{L^2}.$$

On the other hand, observe that by construction $\partial_\sigma \tilde{\eta}(\sigma, t)|_{\sigma=0}$ is a Jacobi field along the geodesic $\tilde{\eta}(0, t)$ and therefore we have (see [4], and, for the recent development on the inviscid parallel shear flow on compact manifolds, see [5])

$$\|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2} \geq Ct \quad \text{for any } t > 0,$$

where $C = \partial_t|_{t=0} \|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2}$. In our case this positive constant C is nothing more than $\|U_{in}\|_{L^2}$, which is (essentially) independent of m . In fact, since

$$\frac{\partial_\sigma \tilde{\eta}(0, t)}{t} \sim \frac{1}{t} \int_0^t (\partial_\sigma U \circ \tilde{\eta})(0, t') dt' \sim U_{in},$$

from (2.1), we see that, at least $\partial_t|_{t=0} \|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2}$ is nonzero (for each m). Second we show that $C = \|U_{in}\|_{L^2}$ rigorously. By

$$\partial_t \|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2} = \frac{\langle \partial_t \partial_\sigma \tilde{\eta}(0, t), \partial_\sigma \tilde{\eta}(0, t) \rangle_{L^2}}{\|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2}}$$

and since the Jacobi field vanishes at $t = 0$, i.e., $\partial_\sigma \tilde{\eta}(0, 0) = 0$, then, by L'Hôpital's rule, we have

$$\partial_t|_{t=0} \|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2} = \frac{\langle \partial_t|_{t=0} \partial_\sigma \tilde{\eta}(0, t), \partial_t|_{t=0} \partial_\sigma \tilde{\eta}(0, t) \rangle_{L^2}}{\partial_t|_{t=0} \|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2}} = \frac{\|U_{in}\|_{L^2}^2}{\partial_t|_{t=0} \|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2}}$$

Thus we have $\partial_t|_{t=0} \|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2} = \|U_{in}\|_{L^2}$.

Therefore we have

$$\begin{aligned} Ct &\leq \|\partial_\sigma \tilde{\eta}(0, t)\|_{L^2} \sim \|\partial_\sigma \tilde{\eta}_1(0, t)\|_{L^2} + \|\partial_\sigma \tilde{\eta}_2(0, t)\|_{L^2} \\ &\lesssim \int_0^t \|\partial_\sigma U_2(0, t')\|_{L^2} dt' + \int_0^t \int_0^{t'} \|\partial_\sigma U_2(0, t'')\|_{L^2} dt'' dt' \|f'_m\|_{L^\infty} \\ &\quad + t \sup_{0 < t' < t} \|\partial_\sigma U_1(0, t')\|_{L^2}. \end{aligned}$$

For any sufficiently small $\epsilon > 0$, let us assume that both $\partial_\sigma U_1$ and $\partial_\sigma U_2$ are controlled by

$$\|\partial_\sigma U_1(0, t)\|_{L^2} \lesssim 1 \quad \text{and} \quad \|\partial_\sigma U_2(\sigma, t)\|_{L^2} \lesssim \langle t \rangle^{-1-\epsilon}$$

for $t > 0$. Then this assumption breaks the lower bound estimate, and it is in contradiction. Finally, for any sufficiently large $T > 0$, there is $\sigma_0 > 0$ such that for any $\sigma < \sigma_0$ and $t \in [0, T]$,

$$\|U_1(\sigma, t)\|_{L^2} \sim \sigma \|U_1(0, t)\|_{L^2} \gtrsim \sigma \quad \text{or} \quad \|U_2(\sigma, t)\|_{L^2} \sim \sigma \|U_2(0, t)\|_{L^2} \gtrsim \sigma \langle t \rangle^{-1-\epsilon}$$

by the Taylor expansion along σ -direction.

Acknowledgments. The author would like to thank Professors Herbert Koch and Gerard Misiolek for inspiring communications. Research of TY was partially supported by Grant-in-Aid for Young Scientists A (17H04825), Grant-in-Aid for Scientific Research B (15H03621, 17H02860, 18H01136 and 18H01135).

REFERENCES

1. J. Bedrossian and N. Masmoudi, *Inviscid damping and the asymptotic stability of planar shear flows in the 2D Euler equations*, Publ. Math. IHÉS, **122** (2015), 195–300.
2. D. Ebin and J. Marsden, *Groups of diffeomorphisms and the motion of an incompressible fluid*, Ann. Math. **92** (1970), 102–163.
3. A. Majda and A. Bertozzi, *Vorticity and Incompressible Flow*, Cambridge University Press, Cambridge 2002.
4. G. Misiolek, *Stability of flows of ideal fluids and the geometry of the group of diffeomorphisms*, Indiana Univ. Math. J., **42** (1993), 215–235.
5. T. Tauchi and T. Yoneda, *Existence of a conjugate point in the incompressible Euler flow on an ellipsoid*, arXiv:1907.08365

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA 3-8-1 ME-GURO, TOKYO 153-8914, JAPAN

E-mail address: yoneda@ms.u-tokyo.ac.jp