

# LINEAR INVISCID DAMPING IN SOBOLEV AND GEVREY SPACES

CHRISTIAN ZILLINGER

ABSTRACT. In a recent article [Jia19] Jia established linear inviscid damping in Gevrey regularity for compactly supported Gevrey regular shear flows in a finite channel, which is of great interest in view of existing nonlinear results [DM18], [BM14], [LJ18]. In this article we provide an alternative very short proof of stability in Gevrey regularity as a consequence of stability in high Sobolev regularity [Zil17a], [Zil16]. Here, we consider both the setting of a finite channel with compactly supported perturbations and of an infinite channel without this restriction. Furthermore, we consider the setting where perturbations vanish only of finite order.

In recent years the asymptotic stability of the Euler equations

$$\partial_t v + v \cdot \nabla v + \nabla p = 0,$$

near shear flow solutions  $v = (U(y), 0)$  has been an area of very active research. Following the works of Mouhot and Villani [MV10] on Landau damping in plasma physics, in a seminal work Bedrossian and Masmoudi [BM14] for the first time established nonlinear asymptotic stability and damping for the prototypical case  $U(y) = y$ , known as Couette flow. Here Gevrey regularity plays a crucial role in controlling nonlinear resonances, so called *echoes* [DM18], [DZ19], [BMM16]. In contrast, the linear problem is known to stable in (arbitrary) Sobolev regularity [Zil17a], [GNRS19], [WZZ18] for the setting without boundary, but only stable in (optimal) low Sobolev regularity for the setting with boundary [Zil16], [WZZ18] unless shear perturbation vanishes on the boundary. In a recent work [Jia19] Jia thus studied the problem of linear asymptotic stability of compactly supported perturbations to Couette flow in Gevrey regularity.

As the main results of this article we show that:

- Stability in Gevrey regularity corresponds to a quantitative control of the stability in Sobolev spaces. In particular, we show that the control established in [Zil17a] yields a very short proof of stability in Gevrey classes for the setting of an infinite channel. Furthermore, only a quantitative stability result in  $L^2$  is needed, which then implies all higher stability results.
- In [Jia19] Jia considers the question of stability in Gevrey regularity for the setting of a finite channel, where the shear flow and the vorticity perturbation are compactly supported away from the boundary. In view of the boundary instabilities established in [Zil16] such a restriction might be necessary. In this work we show that under such a support condition stability in Gevrey regularity for the setting of a finite channel essentially reduces to the setting without boundary with minor correction terms similar to the  $H^1$  stability problem considered in [Zil16].

- As a further result, we establish stability in Sobolev spaces  $H^j$ ,  $j \leq N$  in the setting of a finite channel if  $U''$  and  $\omega_0$  vanish up to order  $N$  on the boundary.
- In this article we restrict ourselves to considering small, smooth bilipschitz shear flows and circular flows close to (Taylor-)Couette flow. We expect a further extension to more general and degenerate shear flows and circular flows in weighted spaces in an analogous way to [CZZ19] to be possible with some technical effort.

In this sense the core problem of linear inviscid damping lies in establishing  $L^2$  stability and  $H^1$  stability (which has to account for some boundary effects). The setting of higher regularity then follows by an iteration scheme.

We remark that our theorems impose a smallness condition which is sufficient but not necessary. As shown in [WZZ18] a more precise condition is given by requiring that there are no embedding eigenvalues of the associated Rayleigh problem. Our stronger condition allows us to construct a Lyapunov functional using perturbative methods.

The linearized Euler equations around a shear flow  $U(y)$  are given by

$$\partial_t \omega + U(y) \partial_x \omega + U''(y) \partial_x \Delta^{-1} \omega = 0.$$

Here,  $\omega_0$  and hence  $\omega$  is understood to without loss of generality have zero mean in  $x$  and in the setting of a finite channel  $\mathbb{T}_L \times [0, 1]$ ,  $\Delta^{-1}$  imposes zero Dirichlet boundary conditions in  $y$ . Assuming that  $U(y)$  is Bilipschitz, we change variables by  $y = U^{-1}(z)$  and denote

$$\begin{aligned} f(z) &= U''(U^{-1}(z)), \\ g(z) &= U'(U^{-1}(z)), \end{aligned}$$

and further pass to Lagrangian coordinates  $(t, x + tz, z)$ . With respect to these coordinates our problem is given by

$$(1) \quad \begin{aligned} \partial_t \omega + f \partial_x L_t \omega &= 0, \\ L_t &= (\partial_x^2 + (g(\partial_z - t \partial_x))^2)^{-1}, \end{aligned}$$

where again in the setting of a finite channel  $L_t$  satisfies zero Dirichlet boundary conditions.

We recall that Gevrey classes measure the growth of  $C^j$  or  $H^j$  norms as  $j \rightarrow \infty$ . See [IJ18], [Gev] and [Hör15, page 281].

**Definition 1.** *Let  $f \in C^\infty$  and  $s \in [1, \infty)$ . We then introduce the following three related but distinct definitions of the Gevrey class  $\mathcal{G}_s$ :*

- (1) *We say  $f$  is in the  $L^\infty$  based Gevrey class  $\mathcal{G}_s^\infty$  if there exists a constant  $C > 0$  such that*

$$\|f\|_{C^j} \leq C^{1+j} (1+j)^{sj}$$

*for all  $j \in \mathbb{N}$ .*

- (2) *We say that  $f$  is in the (Sobolev based) Gevrey class  $\mathcal{G}_s$  if there exists a constant  $C > 0$  such that*

$$\|f\|_{H^j} \leq C^{1+j} (1+j)^{sj}$$

*for any  $j \in \mathbb{N}$*

- (3) We say that  $f$  is in the second Sobolev based Gevrey class if there exists a constant  $\lambda > 0$  such that

$$\int \exp(\lambda \langle \xi \rangle^{\frac{1}{s}}) |\tilde{f}(\xi)|^2 d\xi < \infty.$$

**Remark 2.** • We remark that in the literature also a parametrization in terms of  $\frac{1}{s} \in (0, 1]$  is common.

- A more general version of the first definition considers the restriction of  $f$  to compact sets  $K$  and constants  $C_K$ . For example,  $f(x) = x$  does not satisfy our definition, since we impose that  $f(x)$  is bounded uniformly. However, since our theorems impose these constraints on derivatives of  $U(y)$ ,  $U(y)$  itself may be close to affine.
- We may use a Sobolev embedding to estimate  $\|f\|_{C^j} \leq \|f\|_{H^{j+N}}$ , where  $N > 0$  depends on the dimension. Increasing  $s$  slightly and increasing  $C$ , we thus see that every  $f$  in the first Sobolev based Gevrey class is also contained in the  $L^\infty$  based Gevrey class.
- Expressing  $\exp(\lambda \langle \xi \rangle^{\frac{1}{s}})$  as a series, the last definition implies that

$$\lambda^j \frac{1}{j!} \|f\|_{H^{\frac{j}{2s}}}^2 \leq C$$

for all  $j \in \mathbb{N}$ . Expressing the factorial using the Stirling approximation and considering  $\sigma = \frac{j}{2s}$ , we thus see that any such function also satisfies the second definition.

- Conversely, we may round up  $\frac{j}{2s}$  in the series expansion to show that the second definition also implies the third with an arbitrarily small loss in  $s$ .
- In our analysis we consider the regularity of coefficient functions according to the first definition and the regularity of the vorticity according to the second definition.

The following three theorems summarize our main results. We first consider the case of an infinite channel  $\mathbb{T}_L \times \mathbb{R}$ , for which we had previously established non-quantitative stability results in [Zil17a]. The following theorem improves this to quantitative estimates for each  $H^j$  and thus to Gevrey regularity. Here, as we will see in Section 1, the core of the proof is given by establishing quantitative stability in  $L^2$ , from which higher regularity follows by a short inductive argument.

**Theorem 3** (Summary infinite channel). *Consider the linearized Euler equations (1) on  $\mathbb{T}_L \times \mathbb{R}$  around a bilipschitz shear flow  $(U(y), 0)$ . There exists  $c > 0$  such that if*

$$(2) \quad \|f\|_{W^{1,\infty}L} < c,$$

then for any  $s \in [1, \infty)$  if  $f, g \in \mathcal{G}_s^\infty$  and  $\omega_0 \in \mathcal{G}_s$ , the problem (1) is stable in Gevrey regularity. That is, if for all  $j \in \mathbb{N}$  it holds that

$$\begin{aligned} \|f\|_{C^j} + \|g\|_{C^j} &\leq D_1^{1+j} (1+j)^{js}, \\ \|\omega_0\|_{H^j} &\leq D_2^{1+j} (1+j)^{js}, \end{aligned}$$

then there exists  $C = C(D_1, D_2, c)$  such that for all times  $t \geq 0$

$$(3) \quad \|\omega(t)\|_{H^j} \leq C^{1+j} (1+j)^{js}.$$

The smallness condition (2) here is imposed in order to allow a perturbative construction in our stability proof for  $L^2$  and sufficient but not necessary (see Section 1 for further discussion).

While the setting of a finite channel  $\mathbb{T}_L \times [0, 1]$  is in general unstable in higher Sobolev regularity [Zil16], it turns out that in the setting of compactly supported perturbations studied in [Jia19] all boundary effects can be easily controlled and an analogous stability result hold.

**Theorem 4** (Summary finite channel). *Consider the linearized Euler equations (1) on  $\mathbb{T}_L \times [0, 1]$  around a bilipschitz shear flow  $(U(y), 0)$ , with  $U' \geq 1$ . There exists  $c > 0$  such that if*

$$(4) \quad \|f\|_{W^{1,\infty}L} < c,$$

then for any  $s \in [1, \infty)$  if  $f, g \in \mathcal{G}_s^\infty$  and  $\omega_0 \in \mathcal{G}_s$  are compactly supported away from the boundary, then problem (1) on  $\mathbb{T}_L \times [0, 1]$  is stable in Gevrey regularity in the sense of Theorem 3.

Here, the core of the problem lies in establishing stability in  $H^1$  as in [Zil16], from which the desired quantitative higher Sobolev and Gevrey regularity results then follow by induction.

Finally, we note that it is not necessary to impose the condition of compact support, but that a high order of vanishing on the boundary is sufficient to establish stability in Sobolev regularity (or Gevrey regularity).

**Theorem 5** (Finite regularity for a finite channel). *Let  $g$  satisfy the same assumptions as in the previous theorem and suppose that there exists  $N \in \mathbb{N}_0$  such that  $f$  and the initial vorticity perturbation  $\omega_0$  vanish to order  $N$  on the boundary. Suppose further that*

$$\|f\|_{C^j} + \|g\|_{C^j} \leq D_1^{1+j}(1+j)^{j^s}$$

for all  $j \leq N$ . Then there exists  $C = C(D_1, L)$  such that for all  $j \leq N$  and all  $t \geq 0$  it holds that

$$\|\omega(t)\|_{H^j} \leq C^{1+j}\|\omega_0\|_{H^j}.$$

In particular, the Gevrey stability result of Theorem 4 is still valid if we only assume that  $f$  and  $\omega_0$  vanish of infinite order.

## 1. THE INFINITE CHANNEL CASE

As a starting point we consider the problem (1) in the infinite channel  $\mathbb{T}_L \times \mathbb{R}$  and establish the following quantitative improvement of Theorem 4.5 in [Zil17a]:

**Theorem 6.** *Suppose that  $U \in C^2(\mathbb{R})$  is bilipschitz and let  $g(z) = U'(U^{-1}(z))$ ,  $f(z) = U''(U^{-1}(z))$ . Consider the linearized Euler equations around  $U$  on  $\mathbb{T}_L \times \mathbb{R}$  and suppose that*

$$\|f\|_{W^{1,\infty}L} \ll 1.$$

Suppose further that for some  $j \in \mathbb{N}$ ,  $f, g \in W^{1+j,\infty}$ . Then the solution  $\omega$  (in Lagrangian coordinates) satisfies

$$\|\omega(t)\|_{\dot{H}^j}^2 \leq C\|\omega_0\|_{\dot{H}^j}^2 + C2^j \sum_{j_1+j_2 \leq j} \|\omega_0\|_{\dot{H}^{j_1}}^2 \|(f, g)\|_{j_2},$$

for all times  $t \geq 0$ . Here, we used the short notations:

$$\begin{aligned} \|f\|_j &:= \sup_{j_1+j_2+\dots+j_N=j} \prod \|\partial_z^{j_i} f\|_{L^\infty}, \\ \|(f, g)\|_j &:= \sum_{j_1+j_2=j} (1 + \|f\|_{j_1})(1 + \|g\|_{j_2}). \end{aligned}$$

We note that by definition of  $\|\cdot\|_j$  it holds that

$$\|f\|_{j_1} \|f\|_{j_2} \leq \|f\|_{j_1+j_2},$$

which helps to simplify recursive (commutator) estimates of the form

$$\begin{aligned} a_0 &= 1, \\ a_{j+1} &\leq \sum_{j_1+j_2=j} \|f\|_{j_1} a_{j_2}. \end{aligned}$$

We further remark that if  $f \in \mathcal{G}_s^\infty$ , then  $\|f\|_j$  satisfies analogous estimates to  $\|f\|_{C^j}$ :

$$\begin{aligned} \|f\|_j &= 2^j \sup_{j_1+j_2+\dots+j_N=j} \prod \|\partial_z^{j_i} f\|_{L^\infty} \\ &\leq 2^j \sup \prod C^{1+j_i} (1 + j_i)^{j_i} \\ &\leq 2^j C^{2j} (1 + j)^j. \end{aligned}$$

In [Zil17a] we subsumed the precise bound into a control by  $C_j \|\omega_0\|_{H^j}^2$  for a non-explicit constant  $C_j$  and we imposed the stronger constraint that  $\|f\|_{W^{j+1, \infty} L} \ll 1$ . However, as already noted and proven in [Zil16], [Zil17b] only smallness in  $W^{1, \infty}$  is actually used in the proof.

This quantitative control of constants then immediately allows us to establish the stability in Gevrey classes expressed in Theorem 3.

*Proof of Theorem 3.* Let  $s \in [1, \infty)$  be given and  $U'' \in \mathcal{G}_s^\infty, \omega_0 \in \mathcal{G}_s$ . There thus exists constants  $D_1, D_2$  such that

$$\begin{aligned} \|(f, g)\|_j &\leq D_1^{1+j} (1 + j)^{js}, \\ \|\omega_0\|_{H^j} &\leq D_2^{1+j} (1 + j)^{js}, \end{aligned}$$

for any  $j \in \mathbb{N}$ . Applying Theorem 6 we hence obtain that

$$\begin{aligned} \|\omega(t)\|_{H^j}^2 &\leq C(\|\omega_0\|_{H^j}^2 + 2^j \sum_{j_1+j_2=m} C_{j_1} \|\omega_0\|_{H^{j_2}}^2) \\ &\leq CD_1^{2(1+j)} + C2^j \sum_{j_1+j_2=j} C(1 + C^{1+j_1} D_2^{2(1+j_1)} (1 + j_1)^{2j_1 s}) D_2^{2(1+j_2)} (1 + j_2)^{2j_2 s}. \end{aligned}$$

We now note that

$$(1 + j_1)^{2j_1 s} (1 + j_2)^{2j_2 s} \leq (1 + j)^{2(j_1+j_2)s} = (1 + j)^{2js}$$

and (very) roughly estimate all other powers involved in terms of

$$D = 100 \max(C, D_1, D_2)^2.$$

□

*Proof of Theorem 6.* In the following we retrace and improve the proof in [Zil16] and [Zil17a, Section 4] in order to obtain a quantitative control of the constants in the stability estimate.

We iteratively construct a family of Lyapunov functionals. That is, we claim that for all  $j \in \mathbb{N}$  there exist non-increasing energies  $E_j(t)$  such that

$$(5) \quad C \|\omega(t)\|_{H^j}^2 \leq E_j(t) \leq 2 \|\omega(t)\|_{H^j}^2 + C^j \sum_{j_1+j_2=j, j_2 \neq j} \|(f, g)\|_j \|\omega(t)\|_{H^{j_2}}^2.$$

The statement of the theorem then immediately follows by estimating

$$C \|\omega(t)\|_{H^j}^2 \leq E_j(t) \leq E_j(0) \leq 2 \|\omega_0\|_{H^j}^2 + C^j \sum_{j_1+j_2 < j} \|(f, g)\|_{W^{j_1, \infty}} \|\omega_0\|_{H^{j_2}}^2.$$

Here, it turns out that the main challenge lies in constructing the first energy functional  $E_0(t)$  and establishing sufficiently good control of  $\partial_t E_0(t)$ . Energies  $E_j(t)$  with larger  $j$  can then be constructed inductively.

The case  $j = 0$ :

In order to introduce ideas, let us recall the damping mechanism, known as the Orr mechanism, in case of Couette flow  $U(y) = y$ . In this case  $\omega(t, x, y) = \omega_0(x - ty, y)$  and as a result

$$\partial_x \Delta^{-1} \omega \rightsquigarrow \frac{ik}{k^2 + \eta^2} \tilde{\omega}_0(k, \eta + kt),$$

where  $\tilde{\omega}$  denotes the Fourier transform. Changing to coordinates moving with the flow  $(x + ty, y)$  and thus  $(k, \eta - kt)$  we thus obtain the multiplier

$$\frac{k}{k^2 + (\eta - kt)^2} \tilde{\omega}_0(k, \eta).$$

This multiplier illustrates the main properties of the damping mechanism:

- As  $t \rightarrow \infty$  the multiplier  $\frac{k}{k^2 + (\eta - kt)^2}$  tends to zero (at an algebraic rate) and the velocity hence asymptotically converges in  $L^2$ .
- In contrast  $\omega(t, x, y) = \omega_0(x - ty, y)$  does not converge strongly in  $L^2$  but only weakly.
- While  $\frac{k}{k^2 + (\eta - kt)^2}$  decays after the time  $t_c = \frac{\eta}{k}$  before that time the multiplier is actually increasing. Furthermore the operator norm on  $L^2$ ,

$$\sup_{k, \eta} \frac{k}{k^2 + (\eta - kt)^2} = \sup_{k, \eta} \frac{k}{k^2 + \eta^2}$$

does not improve in time.

- However, if we can fix  $k$  and  $\eta$ , then  $\frac{k}{k^2 + (\eta - kt)^2}$  is integrable in time.

Building in particular on the last property we thus aim to construct an energy  $E_0(t)$  such that  $-\partial_t E_0(t) \geq 0$  controls

$$|\langle \omega, U''(y) \partial_x \Delta^{-1} \omega \rangle|.$$

As the coefficient functions in problem (1) do not depend on  $x$  the problem decouples with respect to Fourier modes  $k$  in  $x$  and we may without loss of generality

restrict to considering  $\omega$  being restricted to a single arbitrary but fixed mode  $k$ . If  $g \geq C > 0$  we define the  $H_t^1$  energy by

$$\|u\|_{H_t^1}^2 = \int k^2 |u|^2 + C^2 |(\partial_z - ikt)u|^2$$

and we define the dual  $H_t^{-1}$  energy of a function  $u \in L^2$  in terms of a Fourier weight

$$\|u\|_{H_t^{-1}}^2 := \sum_k \int |\tilde{u}|^2 \frac{1}{k^2 + C^2(\eta - kt)^2} d\eta.$$

In particular, we note that this multiplier is integrable in time and hence define the Fourier multiplier  $A(t)$  by

$$\mathcal{F}(Au) = \exp(c \arctan(C(\eta - kt))) \tilde{u}(k, \eta),$$

where  $0 < c < 1$  is a constant. This multiplier is non-increasing and it holds that for any  $u$  not depending on time

$$\|u\|_{H_t^{-1}}^2 \exp(-c\pi) \leq -\partial_t \langle u, Au \rangle \leq \exp(c\pi) \|u\|_{H_t^{-1}}^2.$$

We then make the ansatz

$$E_0(t) := \langle \omega(t), A(t)\omega(t) \rangle.$$

As  $\exp(c \arctan(C(\eta - kt)))$  is bounded above and below it holds that

$$\exp(-c\pi) \|\omega(t)\|_{L^2}^2 \leq E_0(t) \leq \exp(c\pi) \|\omega(t)\|_{L^2}^2,$$

so (5) holds. It remains to verify that  $E_0(t)$  is non-increasing. We estimate

$$\begin{aligned} \frac{d}{dt} E_0(t) &= \langle \omega(t), (\partial_t A)\omega(t) \rangle + 2 \langle f \partial_x L_t \omega(t), A(t)\omega(t) \rangle \\ &\leq -\exp(-c\pi) \|\omega(t)\|_{H_t^{-1}}^2 + 2 \langle f \partial_x L_t \omega(t), A(t)\omega(t) \rangle. \end{aligned}$$

Using duality we then estimate

$$\begin{aligned} |\langle f \partial_x L_t \omega(t), A(t)\omega(t) \rangle| &\leq \|A(t)\omega(t)\|_{H_t^{-1}} \|f \partial_x L_t \omega(t)\|_{H_t^1} \\ &\leq \exp(c\pi) \|\omega(t)\|_{H_t^{-1}} \|f\|_{W^{1,\infty}} \|\partial_x L_t \omega(t)\|_{H_t^1}. \end{aligned}$$

Lastly, recall that  $L_t \omega$  solves

$$(-k^2 + (g(\partial_y - ikt))^2) L_t \omega = \omega.$$

Testing this equation with  $-\frac{1}{g} L_t \omega$  and using that  $g$  is bounded below we thus obtain that

$$\|L_t \omega\|_{H_t^1}^2 \leq \left\langle -\frac{1}{g L_t \omega}, \omega \right\rangle \leq \|\omega\|_{H_t^{-1}} \left\| \frac{1}{g} \right\|_{C^1} \|L_t \omega\|_{H_t^1}.$$

and thus

$$\|\partial_x L_t \omega(t)\|_{H_t^1} \leq \left\| \frac{1}{g} \right\|_{C^1} \|\omega(t)\|_{H_t^{-1}}.$$

Thus, if  $\|f\|_{W^{1,\infty}}$  is sufficiently small, it holds that

$$\frac{d}{dt} E_0(t) + C \|\omega(t)\|_{H_t^{-1}}^2 \leq 0$$

and in particular it follows that  $E_0(t)$  non-increasing.

The induction step:

Based on the above estimate we claim that in addition to (5) it holds that

$$(6) \quad \begin{aligned} \frac{d}{dt} E_j(t) &\leq -C \|\partial_y^j \omega(t)\|_{H_t^{-1}}^2 - C \|(f, g)\|_1^2 \|\partial_y^{j-1} \omega(t)\|_{H_t^{-1}}^2 \\ &\quad - \cdots - C \|(f, g)\|_j^2 \|\omega(t)\|_{H_t^{-1}}^2, \end{aligned}$$

which we have just established for  $j = 0$ . We then make the ansatz

$$(7) \quad E_0(t) = \langle \omega, A\omega \rangle, \\ E_{j+1}(t) := 2\langle \partial_y^{j+1} \omega, A\partial_y^{j+1} \omega \rangle + 4C \sum_{j_1+j_2=j} \|(f, g)\|_{j_1}^2 E_{j_2}(t).$$

In particular, by construction this satisfies (5) for every  $j$ . It remains to be shown that  $E_{j+1}(t)$  satisfies (6) and hence is non-increasing.

Thus, consider the  $\partial_y^{j+1}$  derivative of the linearized Euler equations:

$$\partial_t \partial_y^{j+1} \omega + f \partial_x L_t \partial_y^{j+1} \omega = [f \partial_x L_t, \partial_y^{j+1}] \omega.$$

By the construction of  $A(t)$  we obtain that

$$\begin{aligned} \frac{d}{dt} 2\langle \partial_y^{j+1} \omega, A\partial_y^{j+1} \omega \rangle &= 2\langle \partial_y^{j+1} \omega, \dot{A} \partial_y^{j+1} \omega \rangle \\ &\quad + 4\langle -f \partial_x L_t \partial_y^{j+1} \omega, A\partial_y^{j+1} \omega \rangle + 4\langle [f \partial_x L_t, \partial_y^{j+1}] \omega, A\partial_y^{j+1} \omega \rangle \\ &\leq -C \|\partial_y^j \omega(t)\|_{H_t^{-1}}^2 + 4\langle [f \partial_x L_t, \partial_y^{j+1}] \omega, A\partial_y^{j+1} \omega \rangle. \end{aligned}$$

Using (6) up to  $j$  and our ansatz (7) it thus suffices to show that

$$\langle [f \partial_x L_t, \partial_y^{j+1}] \omega, A\partial_y^{j+1} \omega \rangle \leq C \|\partial_y^{j+1} \omega\|_{H_t^{-1}} \sum_{j_1+j_2=j} \|(f, g)\|_{j_1} \|\partial_y^{j_2} \omega\|_{H_t^{-1}},$$

at which point we can then conclude our estimate by using Young's inequality.

Indeed, by duality we may control

$$\langle [f \partial_x L_t, \partial_y^{j+1}] \omega, A\partial_y^{j+1} \omega \rangle \leq \|A\partial_y^{j+1} \omega\|_{H_t^{-1}} \|[f \partial_x L_t, \partial_y^{j+1}] \omega\|_{H_t^1},$$

and by construction of  $A$

$$\|A\partial_y^{j+1} \omega\|_{H_t^{-1}} \leq c \|\partial_y^{j+1} \omega\|_{H_t^{-1}}.$$

We may thus focus on computing and estimating the commutator. Here, the  $j+1$  derivatives may fall either on  $f$  or on  $\partial_x L_t$  and we can estimate

$$\|(\partial_y^{j_1} f) \partial_y^{j_2} \partial_x L_t \omega\|_{H_t^1} \leq \|\partial_y^{j_1} f\|_{L^\infty} \|\partial_y^{j_2} \partial_x L_t \omega\|_{H_t^1} + \|\partial_y^{j_1+1} f\|_{L^\infty} \|\partial_y^{j_2} \partial_x L_t \omega\|_{L^2}.$$

Therefore using the structure of  $\|(f, g)\|_j$  we may further reduce to studying  $\|\partial_y^{j_2} \partial_x L_t \omega\|_{H_t^1}$ .

Using the definition of  $L_t$  and the fact that in this setting of an infinite channel we need not worry about boundary conditions, we observe that  $\partial_y^{j_2} L_t \omega$  is the unique solution of

$$(-k^2 + (g(\partial_z - ikt))^2) \partial_y^{j_2} L_t \omega = \partial_y^{j_2} \omega + [(-k^2 + (g(\partial_z - ikt))^2), \partial_y^{j_2}] L_t \omega.$$

Again using the ellipticity of this problem (see also [Zil17a]) it thus follows that

$$\|\partial_y^{j_2} L_t \omega\|_{H_t^1}^2 \leq \|\partial_y^{j_2} \omega\|_{H_t^{-1}}^2 + \| [(-k^2 + (g(\partial_z - ikt))^2), \partial_y^{j_2}] L_t \omega \|_{H_t^{-1}}^2.$$

Inductively repeating this argument for the commutator on the right-hand-side, we may estimate

$$\|\partial_y^{j_2} L_t \omega\|_{H_t^1}^2 \leq \|\partial_y^{j_2} \omega\|_{H_t^{-1}}^2 + \sum_{j_3+j_4=j_2} \|g\|_{j_3}^2 \|\partial_y^{j_4} \omega\|_{H_t^{-1}}^2,$$



which concludes the proof.  $\square$

We thus observe that the main challenge of the proof in this infinite channel setting is given by establishing the result at the level of  $L^2$ . Higher Sobolev space estimates may then be obtained inductively by using commutator estimates. Furthermore Gevrey stability estimates then correspond to good quantitative control of the constants in these estimates.

As one of the main results of [Zil16] we showed that in the setting of a finite channel additional corrections due to boundary effects have to be taken into account and are generally not negligible, leading to asymptotic instability. However, in the special case where  $\omega$  and  $f$  vanish to sufficiently high order on the boundary this instability does not manifest (up to this order). In particular, as we show in the following section, if the perturbation and  $f$  are compactly supported as in the setting considered by Jia [Jia19] the above proof essentially extends to the setting with boundary with some minor modifications.

## 2. THE FINITE CHANNEL CASE

In this section we consider the setting of a finite channel  $\mathbb{T}_L \times [0, 1]$

$$(8) \quad \begin{aligned} \partial_t \omega + f \partial_x L_t \omega &= 0, \\ (\partial_x^2 + (g(\partial_y - t \partial_x))^2) L_t \omega &= 0, \\ L_t \omega|_{y=0,1} &= 0. \end{aligned}$$

Here, in addition to technical challenges such finding suitable basis representations to replace the Fourier transform a major obstacle is given by the boundary conditions imposed on the stream function. Indeed, as one of the main results of [Zil16] we showed that these corrections are generically not integrable in time and results in blow-up in  $H^s$ ,  $s > \frac{3}{2}$  if  $f \omega_0$  does not vanish on the boundary.

That is, while  $L_t \omega(t)$  is prescribed to satisfy impermeable wall conditions (which equals zero Dirichlet conditions after removing the  $x$  average),  $\partial_y^j L_t \omega$  is not given by the unique solution to

$$\begin{aligned} (-k^2 + (g(\partial_y - ikt))^2) \psi &= \partial_y^j \omega + [(-k^2 + (g(\partial_y - ikt))^2), \partial_y^j] L_t \omega, \\ \psi|_{y=0,1} &= 0, \end{aligned}$$

since generically  $\partial_y^j L_t \omega|_{y=0,1} \neq 0$ .

Hence, in order to compute  $\partial_y^j L_t \omega$  we need to include additional boundary corrections:

$$(9) \quad \begin{aligned} \partial_y^j L_t \omega &= L_t \partial_y^j \omega + L_t [(g(\partial_y - ikt))^2, \partial_y^j] L_t \omega \\ &\quad + (\partial_y^j L_t \omega)(0) e^{ikty} u_1 + (\partial_y^j L_t \omega)(1) e^{ikt(y-1)} u_2, \end{aligned}$$

where  $e^{ikty} u_1, e^{ikt(y-1)} u_2$  are homogeneous solutions of the stream function problem (see Proposition 8).

However, if  $\omega_0$  and  $f$  happen to be supported in  $I \subset (0, 1)$

$$(10) \quad \text{supp}(\omega_0) \subset \mathbb{T} \times I, \text{supp}(f) \subset I,$$

this instability can be avoided as shown in [Jia19] and [IJ18] for the linear and nonlinear Euler equations, respectively.

In the following we show that under this support assumption linear stability in arbitrary Sobolev spaces and Gevrey regularity follow as an extension of the  $H^1$

stability results in [Zil16] and thus provide a new short proof of the former result (for a different class of shear flows). Furthermore, we also consider the setting where  $f$  and  $\omega_0$  vanish of (at least) a finite order  $N$ :

$$(V_N) \quad \forall j \leq N, \partial_y^j \omega_0(0) = \partial_y^j \omega_0(1) = \partial_y^j f(0) = \partial_y^j f(1) = 0.$$

As a first observation we note that (10) and  $(V_N)$  are preserved under the evolution.

**Lemma 7.** *Let  $I \subset (0, 1)$  be a closed subinterval and suppose that  $f \in C^1$ ,  $\omega_0 \in L^2$  satisfy (10). Then for any  $t \geq 0$  the solution  $\omega$  of (8) satisfies*

$$\text{supp}(\omega(t) - \omega_0) \subset \mathbb{T} \times I.$$

*Similarly, if  $f \in C^{N+1}$ ,  $\omega_0 \in H^{N+1}$  satisfy  $(V_N)$ , then  $\omega(t) - \omega_0$  vanishes to order at least  $N$  on the boundary.*

*Proof of Lemma 7.* Suppose that  $f$  is supported in the interval  $I$ . Then

$$\partial_t \omega = -f \partial_x L_t \omega$$

vanishes if  $f$  vanishes and hence

$$\omega(t) - \omega_0 = \int_0^t \partial_t \omega$$

is supported in  $\mathbb{T}_L \times I$ .

Concerning the finite order of vanishing we note that  $\partial_t \omega = -f \partial_x L_t \omega$  vanishes of order at least  $N$ , since  $L_t \omega$  vanishes on the boundary (though it might vanish arbitrarily slowly) and  $f$  by assumption vanishes to order at least  $N$ . Integrating over the compact time interval  $[0, t]$  it thus follows that  $\omega(t) - \omega_0$  vanishes of order at least  $N$ .  $\square$

This allows us to establish improved estimates on  $\partial_y^j L_t \omega|_{y=0,1}$ .

**Proposition 8.** *Suppose that for some  $N \in \mathbb{N}$ ,  $f$  and  $\omega_0$  satisfy  $(V_N)$ . For any  $j \leq N$  it holds that*

$$(11) \quad |\partial_y^j L_t \omega|_{y=0,1}| \leq (1 + \|g\|_j) \langle t \rangle^{j-1} \partial_y L_t \omega|_{y=0,1}.$$

*Furthermore, for any  $\delta > 0$  we may estimate*

$$(12) \quad \partial_y L_t \omega|_{y=0,1} \leq C_\delta \langle t \rangle^{-j} \sum_{j_1+j_2=j} \|g\|_{j_1} \sqrt{\sum_{\eta} \frac{1}{\langle \eta - kt \rangle^{1-\delta}} |\partial_y^{j_2} \omega(\eta)|^2}.$$

*Proof of Proposition 8.* In the case  $j = 1$  the estimate (11) is an equality and we recall that  $L_t \omega|_{y=0,1} = 0$  by definition of  $L_t$ . For  $j \geq 2$  we may reduce to the above estimates by noting that

$$(g(\partial_y - ikt))^j L_t \omega = k^2 (g(\partial_y - ikt))^{j-2} L_t \omega + (g(\partial_y - ikt))^{j-2} \omega.$$

If  $j - 2 \leq N$  the last term vanishes and we hence obtain a recursion formula

$$(g(\partial_y - ikt))^j L_t \omega = k^2 (g(\partial_y - ikt))^{j-2} L_t \omega,$$

which we can solve for  $\partial_y^j L_t \omega$  since  $g$  is bounded above and below. The estimate (11) thus immediately follows by induction.

It thus remains to study  $\partial_y L_t \omega|_{y=0,1}$ . As shown in [Zil17a, Lemma 3] we may compute this Neumann data as integrals against  $\omega$ . That is, if  $u_0, u_1$  are solutions of the adjoint problem

$$(13) \quad (-k^2 + ((\partial_y - ikt)g)^2)u = 0$$

with boundary conditions

$$\begin{pmatrix} u_0(0) & u_1(0) \\ u_0(1) & u_1(1) \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix},$$

then it follows that

$$(14) \quad \int u\omega = \int (-k^2 + (g(\partial_y - ikt))^2)L_t\omega = ug^2(\partial_y - ikt)L_t\omega|_{y=0,1} \\ = \begin{cases} g^2(0)\partial_y L_t\omega(0) & \text{if } u = u_0, \\ g^2(1)\partial_y L_t\omega(1) & \text{if } u = u_1, \end{cases}$$

where we again used that  $L_t\omega|_{y=0,1}$ . We note that by the structure of (13) it holds that  $u_0(t, y) = e^{ikty}u_0(0, y)$  and  $u_1(t, y) = e^{ikt(y-1)}u_1(0, y)$  and that

$$u_0(0, y) = \frac{1 \sinh(k(U^{-1}(y) - U^{-1}(1)))}{g \sinh(k(U^{-1}(0) - U^{-1}(1)))}, \\ u_1(0, y) = \frac{1 \sinh(k(U^{-1}(y) - U^{-1}(0)))}{g \sinh(k(U^{-1}(0) - U^{-1}(1)))},$$

can be explicitly computed and are smooth functions.

As  $\omega$  vanishes on the boundary we may integrate by parts  $j$  times in (14) and thus obtain that

$$\partial_y L_t\omega(0) = \frac{k}{g^2(0)} \frac{1}{(-ikt)^j} \int e^{ikty} \partial_y^j (u_0(0, y)\omega).$$

Expanding  $\partial_y^j (u_1(y)\omega)$  by the product rule and  $\partial_y^j \omega$  in terms of its Fourier series, the estimate then follows by noting that

$$\frac{1}{k^j} \left| \int e^{i(kt-\eta)y} \partial_y^j u_1 \right| \leq \frac{1}{\langle \eta - kt \rangle}$$

and that by the Cauchy-Schwarz inequality for any sequence  $R \in l^2$  it holds that

$$\sum_{\eta} \frac{1}{\langle \eta - kt \rangle} |R_{\eta}| \leq \sqrt{\sum_{\eta} \frac{1}{\langle \eta - kt \rangle^{1+\delta}}} \sqrt{\sum_{\eta} \frac{1}{\langle \eta - kt \rangle^{1-\delta}} |R_{\eta}|^2}.$$

The result hence follows with  $C_{\delta} = \sqrt{\sum_{\eta} \frac{1}{\langle \eta - kt \rangle^{1+\delta}}} < \infty$ .  $\square$

With these preparations we are now ready to prove Theorem 4.

*Proof of Theorem 4.* As in Section 1 we iteratively construct a family of Lyapunov functionals. Our basic building block is given by the weight  $A(t)$  from [Zil17a, Lemma 5.3]

$$A(t) : e^{iny} \exp(\arctan(\frac{\eta}{k} - t) - \int_0^t \frac{1}{\langle \tau \rangle^{2\beta}} \frac{1}{(1 + (\eta/k - \tau)^2)^{2\gamma}} d\tau) e^{iny},$$

which clearly satisfies

$$(15) \quad \langle u, \dot{A}(t)u \rangle \leq -C \sum_{\eta} |u(\eta)|^2 \left( \frac{1}{1 + (\eta/k - t)^2} + \frac{1}{\langle t \rangle^{2\beta}} \frac{1}{(1 + (\eta/k - t)^2)^{2\gamma}} \right).$$

Here the second term will be used to control contributions due to  $\partial_y^j L_t \omega|_{y=0,1}$ .

Following the same strategy as in the proof of Theorem 6 we make the ansatz

$$(16) \quad \begin{aligned} E_0(t) &= \langle \omega, A\omega \rangle, \\ E_{j+1}(t) &:= 2\langle \partial_y^{j+1}\omega, A\partial_y^{j+1}\omega \rangle + 4C \sum_{j_1+j_2=j} \|(f, g)\|_{j_1}^2 E_{j_2}(t). \end{aligned}$$

and claim that for all  $j$

$$(17) \quad \begin{aligned} \frac{d}{dt} E_j(t) &\leq -C \sum_{\eta} \left( \frac{1}{1 + (\eta/k - t)^2} + \frac{1}{\langle t \rangle^{2\beta} (1 + (\eta/k - t)^2)^{\gamma}} \right) |\partial_y^j \omega(\eta)|^2 \\ &\quad - C \sum_{j_1+j_2=j-1} \|\partial_y(f, g)\|_{j_1} \\ &\quad \times \sum_{\eta} \left( \frac{1}{1 + (\eta/k - t)^2} + \frac{1}{\langle t \rangle^{2\beta} (1 + (\eta/k - t)^2)^{\gamma}} \right) |\partial_y^{j_2} \omega(\eta)|^2 \leq 0. \end{aligned}$$

By construction it then again holds that

$$C\|\omega(t)\|_{H^j}^2 \leq E_j(t) \leq E_j(0) \leq 2\|\omega_0\|_{H^j}^2 + C^j \sum_{j_1+j_2 < j} \|(f, g)\|_{W^{j_1, \infty}} \|\omega_0\|_{H^{j_2}}^2,$$

which implies the result.

The case  $j = 0$ : In the following we recall the construction of  $E_0(t)$  and  $E_1(t)$  from [Zil17a] and subsequently extend our proof to the case of general  $j$ . We claim that there exists  $c > 0$  such

$$(18) \quad \langle A(t)\omega, ikfL_t\omega \rangle \leq ck\|f\| \sum_{\eta} |\tilde{\omega}(\eta)|^2 \frac{1}{1 + (\eta/k - t)^2}.$$

Then if  $\|f\|kc < C/2$  with  $C$  as in (15) it immediately follows that  $E_0(t)$  is non-increasing and furthermore

$$\frac{d}{dt} E_0(t) \leq -C/2 \sum_{\eta} |\tilde{\omega}(\eta)|^2 \frac{1}{1 + (\eta/k - t)^2}.$$

Indeed, suppose that  $g > c > 0$  and for any  $u \in L^2$  and any  $t > 0$  define  $\Lambda_t[u]$  to be the unique solution of

$$\begin{aligned} (-k^2 + c^2(\partial_y - ikt)^2)\Lambda_t[u] &= u, \\ \Lambda_t[u]|_{y=0,1} &= 0. \end{aligned}$$

That is, we replaced  $g$  by a constant. We then define

$$\|u\|_{H_t^1}^2 := k^2\|u\|_{L^2}^2 + c^2\|(\partial_y - ikt)u\|_{L^2}^2$$

and

$$\|u\|_{H_t^{-1}}^2 := -\langle \Lambda_t[u], u \rangle = \|\Lambda_t[u]\|_{H_t^1}^2.$$

These are by construction dual norms, so we may estimate

$$\langle A(t)\omega, ikfL_t\omega \rangle \leq \|A(t)\omega\|_{H_t^{-1}} k\|f\|_{C^1} \|L_t\omega\|_{H_t^1}.$$

Since  $L_t\omega$  is defined in terms of an elliptic operator we may further estimate

$$\|L_t\omega\|_{H_t^1}^2 \leq -\langle \omega, L_t\omega \rangle \leq \|\omega\|_{H_t^{-1}} \|L_t\omega\|_{H_t^1}^2.$$

Our estimate thus follows if we can show that  $\|u\|_{H_t^{-1}}^2$  is controlled by a Fourier multiplier as in (18). In the whole-space setting of Section 1 such a result is trivial since  $(-k^2 + c^2(\partial_y - ikt)^2)$  is given by a Fourier multiplier and thus  $H_t^{-1}$  is as well. In the present setting with boundary this poses some technical challenges and one can show by explicit computation of  $\Lambda_t[e^{i\eta y}]$  (see [Zil17a] Lemma 5.2) that indeed

$$\|u\|_{H_t^{-1}}^2 \leq c \sum_{\eta} |u(\eta)|^2 \frac{1}{1 + (\eta/k - t)^2}.$$

This concludes the proof for the case  $j = 0$ . As shown in [Zil16] a similar result is also valid in fractional Sobolev spaces.

The induction step: Having established the base case of (17) we now consider the induction step.

We may write the  $\partial_y^{j+1}$  derivative of (8) as

$$(19) \quad \partial_t \partial_y^j \omega + \sum_{j_1+j_2=j} \binom{j}{j_1} (\partial_y^{j_1} f) \partial_y^{j_2} L_t \omega = 0.$$

Furthermore, we may split

$$\begin{aligned} \partial_y^{j_2} L_t \omega &= (\partial_y^{j_2} L_t \omega - \partial_y^{j_2} L_t \omega(0)u_0 - \partial_y^{j_2} L_t \omega(1)u_1) + \partial_y^{j_2} L_t \omega(0)u_0 + \partial_y^{j_2} L_t \omega(1)u_1 \\ &=: \psi_j + \partial_y^{j_2} L_t \omega(0)u_0 + \partial_y^{j_2} L_t \omega(1)u_1. \end{aligned}$$

As the  $H_t^1$  norms of  $u_0(t, y) = e^{ikt y} u_0(0, y)$  and  $u_1(t, y) = e^{ikt(y-1)} u_1(0, y)$  are independent of  $t$  it follows that

$$(20) \quad \|\partial_y^j L_t \omega\|_{H_t^1} \leq \|\psi_j\|_{H_t^1} + C|\partial_y^j L_t \omega|_{(0)} + C|\partial_y^j L_t \omega|_{(1)}.$$

As  $\psi_j$  satisfies zero Dirichlet boundary conditions, we may test

$$(-k^2 + (g(\partial_y - ikt))^2) \psi_j = \partial_y^j \omega(t) + [(-k^2 + (g(\partial_y - ikt))^2), \partial_y^j] L_t \omega$$

with  $-\psi_j$  and integrate by parts to obtain that

$$(21) \quad \|\psi_j\|_{H_t^1} \leq \|\partial_y^j \omega(t)\|_{H_t^{-1}} + \sum_{j_1+j_2=j, j_2 < j} \|g\|_{j_1} \|\partial_y^{j_2} L_t \omega\|_{H_t^1}.$$

Combining (20) and (21) it follows that

$$(22) \quad \|\psi_j\|_{H_t^{-1}} \leq \|\partial_y^j \omega\|_{H_t^{-1}} + \sum_{j_1+j_2=j, j_2 < j} \|g\|_{j_1} (\|\partial_y^{j_2} \omega\|_{H_t^{-1}} + |\partial_y^{j_2} L_t \omega|_{y=0,1}).$$

We may now further invoke Proposition 8 to estimate

$$(23) \quad |\partial_y^{j_2} L_t \omega|_{y=0,1} \leq C_\delta \langle t \rangle^{-1} \sum_{j_3+j_4=j_2} \|g\|_{j_3} \sqrt{\sum_{\eta} \frac{1}{\langle \eta - kt \rangle^{1-\delta}} |\partial_y^{j_4} \omega(\eta)|^2}.$$

As a final tool, we note that for  $u \in \{u_0, u_1\}$  due to the oscillatory structure it holds that

$$(24) \quad |\langle A \partial_y^{j+1} \omega, u \rangle| \leq C_\delta \sqrt{\sum_{\eta} \frac{1}{\langle \eta/k - t \rangle^{1-\delta}} |\partial_y^{j+1} \omega(\eta)|^2}$$

With all these estimates at hand, we may integrate (19) against  $A\partial_y^{j+1}\omega$  and control

$$(25) \quad |\langle \partial_t \partial_y^{j+1} \omega, A \partial_y^{j+1} \omega \rangle| \leq \|f\|_{W^{1,\infty}} \|\partial_y^{j+1} \omega\|_{H_t^{-1}}^2$$

$$(26) \quad + \|f\|_{W^{1,\infty}} C_\delta \langle t \rangle^{-1} \sum_\eta \frac{1}{\langle \eta/k - t \rangle^{1-\delta}} |\partial_y^{j+1} \omega(\eta)|^2$$

$$(27) \quad + \sum_{j_1+j_2=j+1, j_2 < j+1} \|g\|_{j_1} (\|\partial_y^{j_2} \omega\|_{H_t^{-1}} + |\partial_y^{j_2} \omega(\eta)|^2).$$

The first two terms are exactly such that (15) shows that they can be absorbed into

$$\langle \partial_y^{j+1} \omega, \dot{A} \partial_y^{j+1} \omega \rangle,$$

provided  $f$  satisfies the smallness assumption. The remaining terms are all of lower order and using (17) can be absorbed into

$$\frac{d}{dt} 4C \sum_{j_1+j_2=j} \|(f, g)\|_{j_1}^2 E_{j_2}(t)$$

by the induction assumption. Thus, indeed  $E_{j+1}(t)$  satisfies (17), which concludes the proof.  $\square$

### 3. DISCUSSION

In this article we show that stability in Gevrey regularity corresponds to a quantitative control of stability in Sobolev spaces. Furthermore, this quantitative essentially reduces to establishing good estimates in  $L^2$  and  $H^1$  as in [Zil16], [Zil17a], which then extend to arbitrary Sobolev regularity. We thus provide a new perspective on and a very short alternative proof of the results of Jia [Jia19].

Furthermore, we consider the settings of both infinite and finite channels and the milder constraint of a high finite order of vanishing instead of requiring compact support. In particular, vanishing of infinite order is shown to be sufficient to establish stability in Gevrey regularity. On the other hand, our perturbative construction of the energy functional  $E_0$  imposes a smallness condition instead of a sharper non-resonance condition.

A natural question in view of the existing instability results in  $H^{3/2+}$  ([Zil16]) for perturbations not vanishing on the boundary and the stability results of Theorem 5 is to which extent the condition  $(V_N)$  is necessary for (asymptotic) stability to hold, both in the linear and nonlinear setting. Here, the analysis in [Zil16] suggests to consider boundary corrections and search for  $j$  such that

$$\partial_t \partial_y^j \omega|_{y=0,1} = -\partial_y^j (f \partial_x L_t \omega)$$

is not integrable in time. This would then imply an instability in  $W^{j,\infty}$  and by the Sobolev embedding also an instability in higher Sobolev regularity. In this view a sharper formulation of  $(V_N)$  hence might be to impose an order of vanishing  $N$  on the product  $f\omega_0$  instead. However, condition  $(V_N)$  allows for a simple formulation of Proposition 8.

**Acknowledgments.** Christian Zillinger's research is supported by the ERCEA under the grant 014 669689-HADE and also by the Basque Government through the BERC 2014-2017 program and by Spanish Ministry of Economy and Competitiveness MINECO: BCAM Severo Ochoa excellence accreditation SEV-2013-0323.

## REFERENCES

- [BM14] Jacob Bedrossian and Nader Masmoudi. Asymptotic stability for the Couette flow in the 2D Euler equations. *Applied Mathematics Research eXpress*, 2014(1):157–175, 2014.
- [BMM16] Jacob Bedrossian, Nader Masmoudi, and Clément Mouhot. Landau damping: para-products and Gevrey regularity. *Annals of PDE*, 2(1):4, 2016.
- [CZZ19] Michele Coti Zelati and Christian Zillinger. On degenerate circular and shear flows: the point vortex and power law circular flows. *Communications in Partial Differential Equations*, 44(2):110–155, 2019.
- [DM18] Yu Deng and Nader Masmoudi. Long time instability of the Couette flow in low Gevrey spaces. *arXiv preprint arXiv:1803.01246*, 2018.
- [DZ19] Yu Deng and Christian Zillinger. Echo chains as a linear mechanism: Norm inflation, modified exponents and asymptotics. *arXiv preprint arXiv:1910.12914*, 2019.
- [Gev] Gevrey class, Encyclopedia of Mathematics. [http://www.encyclopediaofmath.org/index.php?title=Gevrey\\_class&oldid=14111](http://www.encyclopediaofmath.org/index.php?title=Gevrey_class&oldid=14111)
- [GNRS19] Emmanuel Grenier, Toan T Nguyen, Frédéric Rousset, and Avy Soffer. Linear inviscid damping and enhanced viscous dissipation of shear flows by using the conjugate operator method. *Journal of Functional Analysis*, page 108339, 2019.
- [Hör15] Lars Hörmander. *The analysis of linear partial differential operators I: Distribution theory and Fourier analysis*. Springer, 2015.
- [IJ18] Alexandru Ionescu and Hao Jia. Inviscid damping near shear flows in a channel. *arXiv preprint arXiv:1808.04026*, 2018.
- [Jia19] Hao Jia. Linear inviscid damping in Gevrey spaces. *arXiv preprint arXiv:1904.01188*, 2019.
- [MV10] Clément Mouhot and Cédric Villani. Landau damping. *Journal of Mathematical Physics*, 51(1):015204, 2010.
- [WZZ18] Dongyi Wei, Zhifei Zhang, and Weiren Zhao. Linear inviscid damping for a class of monotone shear flow in sobolev spaces. *Communications on Pure and Applied Mathematics*, 71(4):617–687, 2018.
- [Zil16] Christian Zillinger. Linear inviscid damping for monotone shear flows in a finite periodic channel, boundary effects, blow-up and critical Sobolev regularity. *Arch. Ration. Mech. Anal.*, 221(3):1449–1509, 2016.
- [Zil17a] Christian Zillinger. Linear inviscid damping for monotone shear flows. *Trans. Amer. Math. Soc.*, 369(12):8799–8855, 2017.
- [Zil17b] Christian Zillinger. On circular flows: linear stability and damping. *J. Differential Equations*, 263(11):7856–7899, 2017.