

Parameter uncertainties in weighted unbinned maximum likelihood fits

Christoph Langenbruch^a,

^a I. Physikalisches Institut B, RWTH Aachen, Sommerfeldstr. 14, 52074 Aachen, Germany

E-Mail: christoph.langenbruch@cern.ch.

Abstract

Parameter estimation via unbinned maximum likelihood fits is central for many analyses performed in high energy physics. Unbinned maximum likelihood fits using event weights, for example to statistically subtract background contributions via the *sPlot* formalism, or to correct for acceptance effects, have recently seen increasing use in the community. However, it is well known that the naive approach to the estimation of parameter uncertainties via the second derivative of the logarithmic likelihood does not yield confidence intervals with the correct coverage in the presence of event weights. This paper derives the asymptotically correct expression and compares it with several commonly used approaches for the determination of parameter uncertainties, some of which are shown to not generally be asymptotically correct. In addition, the effect of uncertainties on event weights is discussed, including uncertainties that can arise from the presence of nuisance parameters in the determination of *sWeights*.

Contents

1	Introduction	1
2	Unbinned maximum likelihood fits and event weights	1
2.1	Asymptotically correct uncertainties in the presence of per-event weights .	2
2.2	Commonly used approaches to uncertainties in weighted fits	5
3	Event weights and inclusion of weight uncertainties	7
3.1	Acceptance corrections	7
3.1.1	Weight uncertainties	8
3.2	The <i>sPlot</i> formalism	10
3.2.1	<i>sWeight</i> uncertainties from nuisance parameters	13
4	Examples	14
4.1	Correcting for an acceptance effect with event weights	14
4.2	Background subtraction using <i>sWeights</i>	16
5	Conclusions	28
6	Acknowledgements	29
A	Expectation value Eq. 26 for examples correcting for acceptance effects	30
B	Expectation value Eq. 26 for examples using <i>sWeights</i>	32
C	Binned <i>sWeighted</i> fits	33
D	Results from pseudoexperiments correcting acceptance effects	36
E	Results from pseudoexperiments using <i>sWeights</i>	38
F	<i>sWeights</i> with negligible nuisance parameter correlations	40

1 Introduction

Unbinned maximum likelihood fits are an essential tool for parameter estimation in high energy physics, due to the desirable features of the maximum likelihood estimator. In the asymptotic limit the maximum likelihood estimator is normally distributed around the true parameter value and its variance is equal to the minimum variance bound [1, 2]. Furthermore, in the unbinned approach no information is lost due to binning.

The inclusion of weights into the maximum likelihood formalism is desirable in many applications. Examples are the statistical subtraction of background events in the *sPlot* formalism [3] through the use of per-event weights, and the correction of acceptance effects via weighting by the inverse efficiency. However, with the inclusion of per-event weights the confidence intervals determined by the inverse second derivative of the negative logarithmic likelihood (in the multidimensional case the inverse of the Hessian matrix of the negative logarithmic likelihood) are no longer asymptotically correct.¹ There are several approaches that are commonly used to determine confidence intervals in the presence of event weights. However, as will be shown below, not all of these techniques are guaranteed to give asymptotically correct coverage. In this paper, the asymptotically correct expression for the determination of parameter uncertainties will be derived and then compared with these approaches.

This paper is structured as follows: In Sec. 2 the unbinned maximum likelihood formalism is briefly summarised and the inclusion of event weights is discussed. The asymptotically correct expression is derived in Sec. 2.1 and compared with several commonly used approaches to determine uncertainties in weighted maximum likelihood fits in Sec. 2.2. Section 3 details the correction of acceptance effects and the *sPlot* method as two sources of event weights. The inclusion of uncertainties on event weights, in particular uncertainties originating from nuisance parameters present in the determination of the *sWeights*, is also discussed. The different approaches are compared and contrasted using two specific examples in Sec. 4, an angular fit correcting for an acceptance effect (Sec. 4.1) and the determination of a lifetime when statistically subtracting background events using *sWeights* (Sec. 4.2). Finally, conclusions are given in Sec. 5.

2 Unbinned maximum likelihood fits and event weights

The maximum likelihood estimator for a set of N_P parameters $\vec{\lambda} = (\lambda_1, \dots, \lambda_{N_P})$, given N measurements $\vec{x} = \{x_1, \dots, x_N\}$, is determined by solving (typically numerically using a software package like MINUIT [5]) the maximum likelihood condition

$$\begin{aligned} \frac{\partial}{\partial \lambda_j} \ln \mathcal{L}(x_1, \dots, x_N | \vec{\lambda}) \Big|_{\hat{\vec{\lambda}}} &= 0 \\ \sum_{e=1}^N \frac{\partial}{\partial \lambda_j} \ln \mathcal{P}(x_e | \vec{\lambda}) \Big|_{\hat{\vec{\lambda}}} &= 0, \end{aligned} \tag{1}$$

¹It should further be noted, that the inclusion of event weights involves some loss of information [4].

where $\mathcal{P}(x_e|\vec{\lambda})$ denotes the probability density function evaluated for the event x_e and parameters $\vec{\lambda}$. Maximising the logarithmic likelihood $\ln \mathcal{L}$ finds the parameters $\hat{\vec{\lambda}}$ for which the measured data \vec{x} becomes the most likely. The covariance matrix V_{ij} for the parameters in the absence of event weights can be calculated from the inverse matrix of second derivatives (the Hessian matrix) of the negative logarithmic likelihood

$$\begin{aligned} V_{ij} &= - \left(\frac{\partial^2}{\partial \lambda_i \partial \lambda_j} \ln \mathcal{L}(x_1, \dots, x_N | \vec{\lambda}) \Big|_{\hat{\vec{\lambda}}} \right)^{-1} \\ &= - \left(\sum_{e=1}^N \frac{\partial^2}{\partial \lambda_i \partial \lambda_j} \ln \mathcal{P}(x_e | \vec{\lambda}) \Big|_{\hat{\vec{\lambda}}} \right)^{-1}. \end{aligned} \quad (2)$$

evaluated at $\vec{\lambda} = \hat{\vec{\lambda}}$. When including even weights $w_{e=1, \dots, N}$ to give each measurement a specific weight the maximum likelihood condition becomes²

$$\sum_{e=1}^N w_e \frac{\partial}{\partial \lambda_j} \ln \mathcal{P}(x_e | \vec{\lambda}) \Big|_{\hat{\vec{\lambda}}} = 0. \quad (3)$$

It should be noted, that the weighted inverse Hessian matrix

$$V_{ij} = - \left(\sum_{e=1}^N w_e \frac{\partial^2}{\partial \lambda_i \partial \lambda_j} \ln \mathcal{P}(x_e | \vec{\lambda}) \Big|_{\hat{\vec{\lambda}}} \right)^{-1}. \quad (4)$$

will generally not give asymptotically correct confidence intervals. This can be most easily seen when assuming constant weights $w_e = w$ which will result in an over-estimation ($w > 1$) or under-estimation ($w < 1$) of the statistical power of the sample and confidence intervals that thus under- or overcover.

2.1 Asymptotically correct uncertainties in the presence of per-event weights

To derive the parameter variance in the presence of event weights we first discuss the simple case of a single parameter λ . In this case, the estimator $\hat{\lambda}$ is defined implicitly by the condition

$$\sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda} \Big|_{\hat{\lambda}} = 0 \quad (5)$$

which constitutes an unbiased estimating equation (see *e.g.* Ref. [6]), as

$$\left\langle \sum_{e=1}^N w_e \frac{\partial \mathcal{P}(x_e | \lambda)}{\partial \lambda} \Big|_{\lambda_0} \right\rangle = 0, \quad (6)$$

²It should be noted that the left-hand side of Eq. 3 strictly speaking is no longer a standard logarithmic likelihood, however that does not preclude its use in parameter estimation as an estimating function.

which is shown for event weights to correct an acceptance effect in Sec. 3.1 (Eq. 29), and for *sWeights* in Sec. 3.2 (Eq. 43), respectively. Using the fact that $\langle w \partial^2 \mathcal{P} / \partial \lambda^2 |_{\lambda} \rangle < 0$ (see Eqs. 30 and 44) it can be shown that the estimator $\hat{\lambda}$ defined by Eq. 5 is consistent [6]. We can then Taylor-expand Eq. 5 to first order around the (unknown) true value λ_0 , to which $\hat{\lambda}$ converges in the asymptotic limit of large N :

$$\sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda} \Big|_{\lambda_0} + (\hat{\lambda} - \lambda_0) \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda^2} \Big|_{\lambda_0} = 0. \quad (7)$$

This equation can be rewritten as

$$\hat{\lambda} - \lambda_0 = - \frac{\sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda} \Big|_{\lambda_0}}{\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda^2} \Big|_{\lambda_0}}, \quad (8)$$

giving the deviation of the estimator $\hat{\lambda}$ from the true value λ_0 . Due to the central limit theorem, the numerator converges to a Gaussian distribution with mean zero (according to Eq. 6) and variance

$$NV \left(w \frac{\partial \ln \mathcal{P}(x | \lambda)}{\partial \lambda} \Big|_{\lambda_0} \right) = N \left\langle w^2 \frac{\partial \ln \mathcal{P}(x | \lambda)}{\partial \lambda} \Big|_{\lambda_0} \frac{\partial \ln \mathcal{P}(x | \lambda)}{\partial \lambda} \Big|_{\lambda_0} \right\rangle. \quad (9)$$

The sum in the denominator goes to

$$\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda^2} \Big|_{\lambda_0} \rightarrow N \left\langle w \frac{\partial^2 \ln \mathcal{P}(x | \lambda)}{\partial \lambda^2} \Big|_{\lambda_0} \right\rangle \quad (10)$$

in the asymptotic limit due to the law of large numbers. Using Eqs. 9 and 10, the variance in the asymptotic limit is then given by

$$\begin{aligned} V(\hat{\lambda} - \lambda_0) &= \langle (\hat{\lambda} - \lambda_0)^2 \rangle \\ &= \frac{\left\langle w^2 \frac{\partial \ln \mathcal{P}(x | \lambda)}{\partial \lambda} \Big|_{\lambda_0} \frac{\partial \ln \mathcal{P}(x | \lambda)}{\partial \lambda} \Big|_{\lambda_0} \right\rangle}{N \left\langle w \frac{\partial^2 \ln \mathcal{P}(x | \lambda)}{\partial \lambda^2} \Big|_{\lambda_0} \right\rangle^2} \end{aligned} \quad (11)$$

The right-hand side of Eq. 11 is the inverse Godambe information [7, 8], which is central to the theory of estimating equations. As the estimator is consistent, we replace λ_0 with $\hat{\lambda}$ in the asymptotic limit, and further estimate the expectation values through the sample, resulting in

$$V(\hat{\lambda} - \lambda_0) = \frac{\sum_{e=1}^N w_e^2 \frac{\partial \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda} \Big|_{\hat{\lambda}} \frac{\partial \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda} \Big|_{\hat{\lambda}}}{\left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda^2} \Big|_{\hat{\lambda}} \right) \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e | \lambda)}{\partial \lambda^2} \Big|_{\hat{\lambda}} \right)}. \quad (12)$$

This expression is also known as the *sandwich estimator*. In the case where event weights are absent ($w_e = 1$), the numerator in Eq. 11 cancels with one of the inverse Hessian matrices as in this case

$$\left\langle \frac{\partial \ln \mathcal{P}(x|\lambda)}{\partial \lambda} \Big|_{\lambda_0} \frac{\partial \ln \mathcal{P}(x|\lambda)}{\partial \lambda} \Big|_{\lambda_0} \right\rangle = - \left\langle \frac{\partial^2 \ln \mathcal{P}(x|\lambda)}{\partial \lambda^2} \Big|_{\lambda_0} \right\rangle. \quad (13)$$

For $w_e = 1$ the Godambe information thus simplifies to the well known Fisher information.

For the multidimensional case we analogously Taylor-expand Eq. 3 to first order, resulting in

$$\sum_{e=1}^N w_e \frac{\partial}{\partial \lambda_i} \ln \mathcal{P}(x_e|\vec{\lambda}) \Big|_{\vec{\lambda}_0} + \sum_{e=1}^N w_e \sum_{j=1}^{N_P} (\hat{\lambda}_j - \lambda_{0j}) \frac{\partial^2}{\partial \lambda_j \partial \lambda_i} \ln \mathcal{P}(x_e|\vec{\lambda}) \Big|_{\vec{\lambda}_0} = 0, \quad (14)$$

which can be written as a matrix equation

$$\begin{pmatrix} \sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1} \Big|_{\vec{\lambda}_0} \\ \vdots \\ \sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P}} \Big|_{\vec{\lambda}_0} \end{pmatrix} = - \begin{pmatrix} \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1^2} \Big|_{\vec{\lambda}_0} & \cdots & \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1 \partial \lambda_{N_P}} \Big|_{\vec{\lambda}_0} \\ \vdots & \ddots & \vdots \\ \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P} \partial \lambda_1} \Big|_{\vec{\lambda}_0} & \cdots & \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P}^2} \Big|_{\vec{\lambda}_0} \end{pmatrix} \begin{pmatrix} \hat{\lambda}_1 - \lambda_{01} \\ \vdots \\ \hat{\lambda}_{N_P} - \lambda_{0N_P} \end{pmatrix}. \quad (15)$$

Matrix inversion yields an expression for the deviation of the estimator $\hat{\lambda}_i$ from the true value λ_{0i}

$$\begin{pmatrix} \hat{\lambda}_1 - \lambda_{01} \\ \vdots \\ \hat{\lambda}_{N_P} - \lambda_{0N_P} \end{pmatrix} = - \begin{pmatrix} \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1^2} \Big|_{\vec{\lambda}_0} & \cdots & \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1 \partial \lambda_{N_P}} \Big|_{\vec{\lambda}_0} \\ \vdots & \ddots & \vdots \\ \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P} \partial \lambda_1} \Big|_{\vec{\lambda}_0} & \cdots & \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P}^2} \Big|_{\vec{\lambda}_0} \end{pmatrix}^{-1} \begin{pmatrix} \sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1} \Big|_{\vec{\lambda}_0} \\ \vdots \\ \sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P}} \Big|_{\vec{\lambda}_0} \end{pmatrix}. \quad (16)$$

The covariance matrix V_{ij} is then given by

$$\begin{aligned} V_{ij} &= \langle (\hat{\lambda}_i - \lambda_{0i})(\hat{\lambda}_j - \lambda_{0j}) \rangle \\ &= \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_i \partial \lambda_k} \Big|_{\hat{\lambda}} \right)^{-1} \times \left\langle \left(\sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \Big|_{\hat{\lambda}} \right) \left(\sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \Big|_{\hat{\lambda}} \right) \right\rangle \\ &\quad \times \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l \partial \lambda_j} \Big|_{\hat{\lambda}} \right)^{-1} \\ &= \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_i \partial \lambda_k} \Big|_{\hat{\lambda}} \right)^{-1} \times \left(\sum_{e=1}^N w_e^2 \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \Big|_{\hat{\lambda}} \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \Big|_{\hat{\lambda}} \right) \\ &\quad \times \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l \partial \lambda_j} \Big|_{\hat{\lambda}} \right)^{-1}, \end{aligned} \quad (17)$$

or explicitly written in matrix notation as

$$\begin{aligned}
V_{ij} = & \left(\begin{array}{ccc} \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1^2} \Big|_{\hat{\lambda}} & \cdots & \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1 \partial \lambda_{N_P}} \Big|_{\hat{\lambda}} \\ \vdots & \ddots & \vdots \\ \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P} \partial \lambda_1} \Big|_{\hat{\lambda}} & \cdots & \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P}^2} \Big|_{\hat{\lambda}} \end{array} \right)^{-1} \\
& \times \left(\begin{array}{ccc} \sum_{e=1}^N w_e^2 \left[\frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1} \Big|_{\hat{\lambda}} \right]^2 & \cdots & \sum_{e=1}^N w_e^2 \left[\frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1} \Big|_{\hat{\lambda}} \right] \left[\frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P}} \Big|_{\hat{\lambda}} \right] \\ \vdots & \ddots & \vdots \\ \sum_{e=1}^N w_e^2 \left[\frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P}} \Big|_{\hat{\lambda}} \right] \left[\frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1} \Big|_{\hat{\lambda}} \right] & \cdots & \sum_{e=1}^N w_e^2 \left[\frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P}} \Big|_{\hat{\lambda}} \right]^2 \end{array} \right) \\
& \times \left(\begin{array}{ccc} \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1^2} \Big|_{\hat{\lambda}} & \cdots & \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_1 \partial \lambda_{N_P}} \Big|_{\hat{\lambda}} \\ \vdots & \ddots & \vdots \\ \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P} \partial \lambda_1} \Big|_{\hat{\lambda}} & \cdots & \sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_{N_P}^2} \Big|_{\hat{\lambda}} \end{array} \right)^{-1}. \tag{18}
\end{aligned}$$

The above expressions are familiar from the derivation of Eq. 2 (in the absence of event weights) in standard textbooks (*e.g.* Ref. [9]). Equation 18 has been previously discussed in Ref. [4] in the context of event weights for efficiency correction. However, it does not seem to be commonly used and often one of the approaches detailed below in Sec. 2.2 is employed instead.

2.2 Commonly used approaches to uncertainties in weighted fits

Instead of using the asymptotically correct approach given by Eq. 18, often other techniques are used to determine parameter uncertainties in weighted unbinned maximum likelihood fits which are presented below.

- (a) A simple approach used sometimes is to rescale the weights w_e according to

$$w'_e = w_e \frac{\sum_{e=1}^N w_e}{\sum_{e=1}^N w_e^2} \tag{19}$$

and to use Eq. 4 with the weights w'_e . This will rescale the weights such that the total statistical power of the sample corresponds to N events, however, it will not generally reproduce the result in Eq. 18.

- (b) A method proposed in Refs. [1, 2] is to correct the covariance matrix according to

$$V'_{ij} = V_{ik} C_{kl}^{-1} V_{lj}, \tag{20}$$

where $V_{ik(lj)}$ is given by Eq. 4 and C_{kl} is the inverse Hessian matrix determined using squared weights w_e^2 according to

$$C_{kl} = - \left(\sum_{e=1}^N w_e^2 \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k \partial \lambda_l} \Big|_{\hat{\lambda}} \right)^{-1}. \tag{21}$$

This method is the nominal method used in the ROOFIT software package when using weighted events [10] and is thus widely used in particle physics. It corresponds to the result in Eq. 18 only if

$$\left\langle \sum_{e=1}^N w_e^2 \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \bigg|_{\hat{\lambda}} \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \bigg|_{\hat{\lambda}} \right\rangle = - \left\langle \sum_{e=1}^N w_e^2 \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k \partial \lambda_l} \bigg|_{\hat{\lambda}} \right\rangle \quad (22)$$

This is however not generally the case. This becomes more clear when rewriting the left- and right-hand side of Eq. 22 according to

$$\sum_{e=1}^N w_e^2 \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \bigg|_{\hat{\lambda}} \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \bigg|_{\hat{\lambda}} = \sum_{e=1}^N \frac{w_e^2}{\mathcal{P}^2(x_e|\vec{\lambda})} \frac{\partial \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \bigg|_{\hat{\lambda}} \frac{\partial \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \bigg|_{\hat{\lambda}} \quad \text{and} \quad (23)$$

$$\begin{aligned} - \sum_{e=1}^N w_e^2 \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k \partial \lambda_l} \bigg|_{\hat{\lambda}} &= \sum_{e=1}^N \frac{w_e^2}{\mathcal{P}^2(x_e|\vec{\lambda})} \frac{\partial \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \bigg|_{\hat{\lambda}} \frac{\partial \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \bigg|_{\hat{\lambda}} \\ &\quad - \sum_{e=1}^N \frac{w_e^2}{\mathcal{P}(x_e|\vec{\lambda})} \frac{\partial^2 \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k \partial \lambda_l} \bigg|_{\hat{\lambda}} \end{aligned} \quad (24)$$

The expectation value of the second part on the right-hand side of Eq. 24 is not generally zero. While Refs. [1, 2] correctly derive that the expectation value

$$\left\langle \frac{w}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_k \partial \lambda_l} \right\rangle = 0 \quad (25)$$

for an efficiency correction $\epsilon_e = 1/w_e$, this is not generally the case for the expression with squared weights

$$\left\langle \frac{w^2}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_k \partial \lambda_l} \right\rangle \quad (26)$$

resulting in confidence intervals that are not generally asymptotically correct when using this approach. This will be detailed for efficiency corrections in Sec. 3.1 and for *sWeights* in Sec. 3.2. For the specific examples discussed in Secs. 4.1 and 4.2 in this paper, the corresponding expectation values are calculated explicitly in Apps. A and B.

- (c) A general approach for the determination of parameter uncertainties is to bootstrap the data [11]. Repeatedly resampling the data set with replacement allows new samples to be generated that can in turn be used to estimate the parameters $\vec{\lambda}$ using the maximum likelihood method. The width of the distribution of estimated parameter values can then be used as estimator for the parameter uncertainty. This approach is generally valid, however repeatedly (typically $\mathcal{O}(10^3)$ times) solving Eq. 3 numerically can be computationally expensive and thus this approach is often unfeasible.

3 Event weights and inclusion of weight uncertainties

3.1 Acceptance corrections

Acceptance of events with a certain probability ϵ , depending on the measurements x_e and y_e , can be accounted for in unbinned maximum likelihood fits by using event weights $w_e = 1/\epsilon(x_e, y_e)$ in Eq. 3. This can be advantageous when it is difficult or computationally expensive to determine the norm of the probability density function when including the efficiency as an explicit additional multiplicative factor $\epsilon(x, y)$. The covariance in this case can be estimated using Eq. 18 as previously suggested in Ref. [4].

To determine expectation values it is necessary to include the acceptance effect in the probability density function. Following the notation in Refs. [1,2], the probability density function $\mathcal{P}(x, y|\vec{\lambda})$ gives the probability to find the measurements x and y depending on the parameters $\vec{\lambda}$ with

$$\mathcal{P}(x, y|\vec{\lambda}) = \mathcal{P}(x|\vec{\lambda})\mathcal{Q}(y|x) \quad (27)$$

and the proper normalisation $\int \mathcal{P}(x|\vec{\lambda})dx = 1$ and $\int \mathcal{Q}(y|x)dy = 1$. The efficiency $\epsilon(x, y)$ can depend on both variables x and y . The resulting total probability density function including this acceptance effect is then given by

$$\mathcal{G}(x, y|\vec{\lambda}) = \frac{\mathcal{P}(x|\vec{\lambda})\mathcal{Q}(y|x)\epsilon(x, y)}{\int \mathcal{P}(x|\vec{\lambda})\mathcal{Q}(y|x)\epsilon(x, y)dxdy} = \mathcal{P}(x|\vec{\lambda})\mathcal{Q}(y|x)\epsilon(x, y)/\mathcal{N} \quad (28)$$

with normalisation \mathcal{N} . This is the probability density function that needs to be used when determining expectation values. For the likelihood condition we find

$$\begin{aligned} \left\langle w(x, y) \frac{\partial \ln \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle &= \left\langle w(x, y) \frac{1}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle \\ &= \int \frac{1}{\epsilon(x, y)} \frac{1}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \mathcal{P}(x|\vec{\lambda})\mathcal{Q}(y|x)\epsilon(x, y)dxdy/\mathcal{N} \\ &= \frac{\partial}{\partial \lambda_j} \int \mathcal{P}(x|\vec{\lambda})\mathcal{Q}(y|x)dxdy/\mathcal{N} = \frac{\partial}{\partial \lambda_j} 1/\mathcal{N} = 0, \end{aligned} \quad (29)$$

as expected. Further, we obtain

$$\begin{aligned} \left\langle w(x, y) \frac{\partial^2 \ln \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i \partial \lambda_j} \right\rangle &= \left\langle w(x, y) \frac{1}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i \partial \lambda_j} \right\rangle - \left\langle w(x, y) \frac{1}{\mathcal{P}^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle \\ &= \int \frac{1}{\epsilon(x, y)} \frac{1}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i \partial \lambda_j} \mathcal{P}(x|\vec{\lambda})\mathcal{Q}(y|x)\epsilon(x, y)dxdy/\mathcal{N} \\ &\quad - \int \frac{1}{\epsilon(x, y)} \frac{1}{\mathcal{P}^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \mathcal{P}(x|\vec{\lambda})\mathcal{Q}(y|x)\epsilon(x, y)dxdy/\mathcal{N} \\ &= - \int \frac{1}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} dx/\mathcal{N}, \end{aligned} \quad (30)$$

confirming the expectation value in Eq. 25, and

$$\begin{aligned}
\left\langle w(x, y) \frac{\partial \ln \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \ln \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle &= \left\langle w(x, y) \frac{1}{\mathcal{P}^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle \\
&= \int \frac{1}{\epsilon(x, y)} \frac{1}{\mathcal{P}^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \mathcal{P}(x|\vec{\lambda}) \mathcal{Q}(y|x) \epsilon(x, y) dx dy / \mathcal{N} \\
&= - \left\langle w(x, y) \frac{\partial^2 \ln \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i \partial \lambda_j} \right\rangle \tag{31}
\end{aligned}$$

However, the equality derived above is not generally fulfilled for squared weights. In this case, we find

$$\begin{aligned}
\left\langle w^2(x, y) \frac{\partial^2 \ln \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i \partial \lambda_j} \right\rangle &= \left\langle w^2(x, y) \frac{1}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i \partial \lambda_j} \right\rangle - \left\langle w^2(x, y) \frac{1}{\mathcal{P}^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle \\
&= \frac{\partial^2}{\partial \lambda_i \partial \lambda_j} \int \frac{1}{\epsilon(x, y)} \mathcal{P}(x|\vec{\lambda}) \mathcal{Q}(y|x) dx dy / \mathcal{N} \\
&\quad - \int \frac{1}{\epsilon(x, y)} \frac{1}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \mathcal{Q}(y|x) dx dy / \mathcal{N} \tag{32}
\end{aligned}$$

and

$$\begin{aligned}
\left\langle w^2(x, y) \frac{\partial \ln \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \ln \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle &= \left\langle w^2(x, y) \frac{1}{\mathcal{P}^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle \\
&= \int \frac{1}{\epsilon(x, y)} \frac{1}{\mathcal{P}(x|\vec{\lambda})} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i} \frac{\partial \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_j} \mathcal{Q}(y|x) dx dy / \mathcal{N} \\
&= - \left\langle w^2(x, y) \frac{\partial^2 \ln \mathcal{P}(x|\vec{\lambda})}{\partial \lambda_i \partial \lambda_j} \right\rangle \\
&\quad + \frac{\partial^2}{\partial \lambda_i \partial \lambda_j} \int \frac{1}{\epsilon(x, y)} \mathcal{P}(x|\vec{\lambda}) \mathcal{Q}(y|x) dx dy / \mathcal{N}, \tag{33}
\end{aligned}$$

where the term in the last line of Eq. 33, which corresponds to the expectation value in Eq. 26, is not generally zero, as the integral in the numerator can retain a dependence on $\vec{\lambda}$. For the example discussed in Sec. 4.1 this is explicitly calculated in App. A. This shows that parameter uncertainties determined using Eq. 20 are not generally asymptotically correct when performing weighted fits to account for acceptance corrections.

3.1.1 Weight uncertainties

If the weights to correct for an acceptance effect are only known to a certain precision, this induces an additional variance that is not included in Eq. 18 and that needs to be accounted for. If an efficiency histogram is used, *i.e.* the efficiency is given in N_B bins

with uncertainty $\sigma(w_e)_{m=1,\dots,N_B}$, weights inside a bin are fully correlated, but uncorrelated with other bins. In this case, the additional term that needs to be added to account for the weight uncertainties is given by

$$\begin{aligned}
V''_{ij} &= \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_i \partial \lambda_k} \Big|_{\hat{\lambda}} \right)^{-1} \\
&\times \sum_{m=1}^{N_B} \left(\left[\sum_{e \in \text{bin } m} \sigma(w_e)_m \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \Big|_{\hat{\lambda}} \right] \left[\sum_{e \in \text{bin } m} \sigma(w_e)_m \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \Big|_{\hat{\lambda}} \right] \right) \\
&\times \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l \partial \lambda_j} \Big|_{\hat{\lambda}} \right)^{-1}
\end{aligned} \tag{34}$$

If the efficiency is modelled analytically, for example by a parameterisation that is fit to simulated samples, the impact of the uncertainty of the parameters \vec{p} describing the efficiency needs to be accounted for. The weights $w_e(\vec{p})$ can be expanded to first order according to

$$w_e(\vec{p}) = w_e(\vec{p})|_{\hat{p}} + \sum_{m=1}^{N_T} (p_m - \hat{p}_m) \frac{\partial w_e(\vec{p})}{\partial p_m} \Big|_{\hat{p}}, \tag{35}$$

where \vec{p} are the N_T parameters that model the efficiency and $w_e(\vec{p}) = 1/\epsilon(\vec{p})$ and \hat{p} denotes their best estimate. For brevity, the explicit dependence on \vec{p} will be omitted in the following. Accounting for the additional variance due to the uncertainties of the event weights in Eq. 18 requires the calculation of the expectation value

$$\begin{aligned}
&\left\langle \left(\sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \Big|_{\hat{\lambda}} \right) \left(\sum_{e=1}^N w_e \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \Big|_{\hat{\lambda}} \right) \right\rangle \\
&= \sum_{e=1}^N w_e|_{\hat{p}}^2 \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \Big|_{\hat{\lambda}} \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \Big|_{\hat{\lambda}} \\
&+ \sum_{m,n=1}^{N_T} \left(\sum_{e=1}^N \frac{\partial w_e}{\partial p_m} \Big|_{\hat{p}} \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_k} \Big|_{\hat{\lambda}} \right) \Sigma_{mn} \left(\sum_{e=1}^N \frac{\partial w_e}{\partial p_n} \Big|_{\hat{p}} \frac{\partial \ln \mathcal{P}(x_e|\vec{\lambda})}{\partial \lambda_l} \Big|_{\hat{\lambda}} \right)
\end{aligned} \tag{36}$$

Here, Σ_{mn} denotes the covariance matrix for the parameters \vec{p} that analytically describe the efficiency. This results in an additional term that needs to be added to the covariance

matrix Eq. 18 given by

$$\begin{aligned}
V_{ij}'' &= \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_i \partial \lambda_k} \Big|_{\hat{\lambda}} \right)^{-1} \\
&\times \sum_{m,n=1}^{N_T} \left(\sum_{e=1}^N \frac{\partial w_e}{\partial p_m} \Big|_{\hat{p}} \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_k} \Big|_{\hat{\lambda}} \right) \Sigma_{mn} \left(\sum_{e=1}^N \frac{\partial w_e}{\partial p_n} \Big|_{\hat{p}} \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_l} \Big|_{\hat{\lambda}} \right) \\
&\times \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_l \partial \lambda_j} \Big|_{\hat{\lambda}} \right)^{-1}, \tag{37}
\end{aligned}$$

3.2 The *sPlot* formalism

The *sPlot* formalism was introduced in Ref. [3] to statistically separate different event species in a data sample using per-event weights, the so-called *sWeights*, that are determined using a *discriminating variable*. The *sWeights* allow to reconstruct the distribution of the different species in a *control variable*. In this section, only a brief recap of the *sPlot* formalism is given, it is described in more detail in Ref. [3].

The *sWeights* are determined using an extended unbinned maximum likelihood fit of the *discriminating variable* y where the N_S different event species are well separated. An example of a discriminating variable (which will be discussed in more detail in Sec. 4.2) would be the reconstructed mass of a particle which is flat for the background components and peaks clearly for the signal component. The *sWeight* assigned to event e for component i is then given by

$$w_i(y_e) = \frac{\sum_{j=1}^{N_S} \Sigma_{ij}^{\text{yields}} \mathcal{P}_j(y_e)}{\sum_{j=1}^{N_S} N_j \mathcal{P}_j(y_e)} \tag{38}$$

with the probability density functions $\mathcal{P}_j(y_e)$. Here, $\Sigma_{ij}^{\text{yields}}$ denotes the covariance matrix from the extended maximum likelihood fit of the yields which can be calculated according to

$$\left(\Sigma_{ij}^{\text{yields}} \right)^{-1} = \frac{\partial^2 - \ln \mathcal{L}}{\partial N_i \partial N_j} = \sum_{e=1}^N \frac{\mathcal{P}_i(y_e) \mathcal{P}_j(y_e)}{\left(\sum_{j=1}^{N_S} N_j \mathcal{P}_j(y_e) \right)^2}. \tag{39}$$

Using the *sWeights* in a weighted unbinned maximum likelihood fit allows to statistically subtract events originating from species not of interest (such as backgrounds) [12]. However, for the determination of the uncertainties one cannot rely on Eq. 4 and this paper advocates the use of the method presented in Sec. 2.1 (namely Eq. 18) instead.

For *sWeights* the full probability density function, including its dependence on the

control variable y is given by

$$\mathcal{P}(x, y|\vec{\lambda}) = \frac{\sum_{j=1}^{N_s} N_j \mathcal{P}_j(x, y|\vec{\lambda})}{\sum_{j=1}^{N_s} N_j} = \frac{\sum_{j=1}^{N_s} N_j \mathcal{P}_j(x|\vec{\lambda}) \mathcal{P}_j(y)}{\sum_{j=1}^{N_s} N_j}$$

$$\stackrel{e.g.}{=} f_{\text{sig}} \mathcal{P}_{\text{sig}}(x|\vec{\lambda}) \mathcal{P}_{\text{sig}}(y) + (1 - f_{\text{sig}}) \mathcal{P}_{\text{bkg}}(x) \mathcal{P}_{\text{bkg}}(y) \quad (40)$$

with all probability density functions properly normalised. The $sWeights$ have the property

$$\int w_i(y) \mathcal{P}_j(y) dy = \int \frac{\sum_k \Sigma_{ik}^{\text{yields}} \mathcal{P}_k(y)}{\sum_k N_k \mathcal{P}_k(y)} \mathcal{P}_j(y) dy = \sum_k \Sigma_{ik}^{\text{yields}} \Sigma_{kj}^{\text{yields}-1} = \delta_{ij}, \quad e.g. \quad (41)$$

$$\int w_{\text{sig}}(y) \mathcal{P}_{\text{sig}}(y) dy = 1 \quad \text{and} \quad \int w_{\text{sig}}(y) \mathcal{P}_{\text{bkg}}(y) dy = 0. \quad (42)$$

The expectation values need to be calculated using the full probability density function, keeping in mind that only a part of the total probability density function, $\mathcal{P}_i(x|\vec{\lambda})$ (typically the signal part), is used in the $sWeighted$ fit. We find for the likelihood condition

$$\begin{aligned} \left\langle w_i(y) \frac{\partial \ln \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle &= \left\langle w_i(y) \frac{1}{\mathcal{P}_i(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \right\rangle \\ &= \int w_i(y) \frac{1}{\mathcal{P}_i(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\sum_{j=1}^{N_s} N_j \mathcal{P}_j(x|\vec{\lambda}) \mathcal{P}_j(y)}{\sum_{j=1}^{N_s} N_j} dx dy \\ &= \int \frac{1}{\mathcal{P}_i(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} f_i \mathcal{P}_i(x|\vec{\lambda}) dx \\ &= \frac{\partial}{\partial \lambda_j} \int f_i \mathcal{P}_i(x|\vec{\lambda}) dx = \frac{\partial}{\partial \lambda_j} f_i = 0, \end{aligned} \quad (43)$$

with fraction $f_i = N_i / \sum_{j=1}^{N_s} N_j$. Similar to Sec. 3.1, we further find

$$\begin{aligned} \left\langle w_i(y) \frac{\partial^2 \ln \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \right\rangle &= \left\langle w_i(y) \frac{1}{\mathcal{P}_i(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \right\rangle - \left\langle w_i(y) \frac{1}{\mathcal{P}_i^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} \right\rangle \\ &= \int w_i(y) \frac{1}{\mathcal{P}_i(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \mathcal{P}(x, y|\vec{\lambda}) dx dy \\ &\quad - \int w_i(y) \frac{1}{\mathcal{P}_i^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} \mathcal{P}(x, y|\vec{\lambda}) dx dy \\ &= - \int \frac{f_i}{\mathcal{P}_i(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} dx, \end{aligned} \quad (44)$$

confirming the expectation value in Eq. 25, and

$$\begin{aligned}
\left\langle w_i(y) \frac{\partial \ln \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \ln \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} \right\rangle &= \left\langle w_i(y) \frac{1}{\mathcal{P}_i^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} \right\rangle \\
&= \int w_i(y) \frac{1}{\mathcal{P}_i^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} \mathcal{P}(x, y|\vec{\lambda}) dx dy \\
&= - \left\langle w_i(y) \frac{\partial^2 \ln \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \right\rangle. \tag{45}
\end{aligned}$$

For squared weights, the corresponding expectation values become

$$\begin{aligned}
\left\langle w_i^2(y) \frac{\partial^2 \ln \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \right\rangle &= \left\langle w_i^2(y) \frac{1}{\mathcal{P}_i(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \right\rangle - \left\langle w_i^2(y) \frac{1}{\mathcal{P}_i^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} \right\rangle \\
&= \int w_i^2(y) \frac{1}{\mathcal{P}_i(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \mathcal{P}(x, y|\vec{\lambda}) dx dy \\
&\quad - \int w_i^2(y) \frac{1}{\mathcal{P}_i^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} \mathcal{P}(x, y|\vec{\lambda}) dx dy \tag{46}
\end{aligned}$$

and

$$\begin{aligned}
\left\langle w_i^2(y) \frac{\partial \ln \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \ln \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} \right\rangle &= \left\langle w_i^2(y) \frac{1}{\mathcal{P}_i^2(x|\vec{\lambda})} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j} \frac{\partial \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_k} \right\rangle \\
&= - \left\langle w_i^2(y) \frac{\partial^2 \ln \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \right\rangle \\
&\quad + \int w_i^2(y) \frac{1}{\mathcal{P}_i(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}_i(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \mathcal{P}(x, y|\vec{\lambda}) dx dy. \tag{47}
\end{aligned}$$

The integral in the last line of Eq. 47, which corresponds to the expectation value in Eq. 26, is not generally zero, for the example of a single signal and background component we find

$$\begin{aligned}
\int w_{\text{sig}}^2(y) \frac{1}{\mathcal{P}_{\text{sig}}(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}_{\text{sig}}(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \mathcal{P}(x, y|\vec{\lambda}) dx dy &= \int w_{\text{sig}}^2(y) \frac{1}{\mathcal{P}_{\text{sig}}(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}_{\text{sig}}(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} \\
&\quad \times \left[f_{\text{sig}} \mathcal{P}_{\text{sig}}(x|\vec{\lambda}) \mathcal{P}_{\text{sig}}(y) + (1 - f_{\text{sig}}) \mathcal{P}_{\text{bkg}}(x) \mathcal{P}_{\text{bkg}}(y) \right] dx dy \\
&= \underbrace{\int \frac{1}{\mathcal{P}_{\text{sig}}(x|\vec{\lambda})} \frac{\partial^2 \mathcal{P}_{\text{sig}}(x|\vec{\lambda})}{\partial \lambda_j \partial \lambda_k} (1 - f_{\text{sig}}) \mathcal{P}_{\text{bkg}}(x) dx}_{\text{not generally 0}} \\
&\quad \times \underbrace{\int w_{\text{sig}}^2(y) \mathcal{P}_{\text{bkg}}(y) dy}_{>0}, \tag{48}
\end{aligned}$$

where the integral over x is explicitly shown to be not generally zero for the *sWeighted* examples in this paper in App. B. This shows that the method to determine parameter uncertainties according to Eq. 20, is not generally asymptotically correct for *sWeighted* fits.

3.2.1 *sWeight* uncertainties from nuisance parameters

A potential complication arises from nuisance parameters used in the fit of the discriminating variable. The covariance-weighting in Eq. 38 does not account for the impact of these nuisance parameters on the *sWeights*, including their potential correlations with the event yields. Indeed, Ref. [3] stresses to fix all nuisance parameters when deriving the covariance matrix in Eq. 39 via an unbinned maximum likelihood fit. If the nuisance parameters however have for example significant correlations with the event yields, this has a systematic effect on the event weights resulting in an underestimation of the covariance when using Eq. 18, that needs to be accounted for.

To study the impact on the event weights w_e it is useful to expand the *sWeights* around their nominal value $w_e|_{\hat{\vec{p}}}$ in full analogy to Eq. 35, according to

$$w_e(\vec{p}) = w_e(\vec{p})|_{\hat{\vec{p}}} + \sum_{m=1}^{N_T} (p_m - \hat{p}_m) \left. \frac{\partial w_e(\vec{p})}{\partial p_m} \right|_{\hat{\vec{p}}}. \quad (49)$$

The full set of parameters needed for the fit of the discriminating variable is denoted by \vec{p} and their best estimate from the fit by $\hat{\vec{p}}$. The parameter set consists of the N_S yields and N_N additional nuisance parameters, resulting in $N_T = N_S + N_N$ parameters in total.

Analogously to Eq. 36, to account for the effect of the uncertainties of the nuisance parameters on the *sWeights* (which is fully correlated amongst all events) in Eq. 18 necessitates the calculation of the expectation value

$$\begin{aligned} & \left\langle \left(\sum_{e=1}^N w_e \left. \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_k} \right|_{\hat{\vec{\lambda}}} \right) \left(\sum_{e=1}^N w_e \left. \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_l} \right|_{\hat{\vec{\lambda}}} \right) \right\rangle \\ &= \sum_{e=1}^N w_e|_{\hat{\vec{p}}}^2 \left. \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_k} \right|_{\hat{\vec{\lambda}}} \left. \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_l} \right|_{\hat{\vec{\lambda}}} \\ &+ \sum_{m,n=1}^{N_T} \left(\sum_{e=1}^N \left. \frac{\partial w_e}{\partial p_m} \right|_{\hat{\vec{p}}} \left. \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_k} \right|_{\hat{\vec{\lambda}}} \right) \Sigma_{mn} \left(\sum_{e=1}^N \left. \frac{\partial w_e}{\partial p_n} \right|_{\hat{\vec{p}}} \left. \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_l} \right|_{\hat{\vec{\lambda}}} \right) \end{aligned} \quad (50)$$

Here, the covariance matrix Σ_{mn} is given by the difference of the covariance matrices $\Sigma_{mn}^{\text{full}}$, which is the covariance matrix of the fit of the discriminating variable including all parameters, and $\Sigma_{mn}^{\text{yields}}$ which only includes the yields and which corresponds to Eq. 39. Assuming the yields are ordered the same way in the first rows and columns of the

covariance matrices, Σ_{mn} is given by

$$\begin{aligned} \Sigma_{mn} &= \Sigma_{mn}^{\text{full}} - \Sigma_{mn}^{\text{yields}} \\ &= \begin{pmatrix} \Sigma_{00}^{\text{full}} & \cdots & \cdots & \cdots & \cdots & \Sigma_{N_T 0}^{\text{full}} \\ \vdots & \ddots & & & & \vdots \\ \vdots & & \ddots & & & \vdots \\ \vdots & & & \ddots & & \vdots \\ \vdots & & & & \ddots & \vdots \\ \Sigma_{0N_T}^{\text{full}} & \cdots & \cdots & \cdots & \cdots & \Sigma_{N_T N_T}^{\text{full}} \end{pmatrix} - \begin{pmatrix} \Sigma_{00}^{\text{yields}} & \cdots & \Sigma_{N_S 0}^{\text{yields}} & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & & \vdots \\ \Sigma_{0N_S}^{\text{yields}} & \cdots & \Sigma_{N_S N_S}^{\text{yields}} & 0 & \cdots & 0 \\ 0 & \cdots & 0 & 0 & & 0 \\ \vdots & & \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & \cdots & 0 \end{pmatrix} \end{aligned} \quad (51)$$

The fixing of the nuisances in the derivation of the *sWeights* thus results in the additional term

$$\begin{aligned} V_{ij}'' &= \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_i \partial \lambda_k} \Big|_{\hat{\lambda}} \right)^{-1} \\ &\times \sum_{m,n=1}^{N_T} \left(\sum_{e=1}^N \frac{\partial w_e}{\partial p_m} \Big|_{\hat{p}} \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_k} \Big|_{\hat{\lambda}} \right) \Sigma_{mn} \left(\sum_{e=1}^N \frac{\partial w_e}{\partial p_n} \Big|_{\hat{p}} \frac{\partial \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_l} \Big|_{\hat{\lambda}} \right) \\ &\times \left(\sum_{e=1}^N w_e \frac{\partial^2 \ln \mathcal{P}(x_e | \vec{\lambda})}{\partial \lambda_l \partial \lambda_j} \Big|_{\hat{\lambda}} \right)^{-1}, \end{aligned} \quad (52)$$

with Σ_{mn} given by Eq. 51, that needs to be added to the covariance in Eq. 18 when nuisance parameters with significant correlations with the event yields are present in the fit of the control variable to determine the *sWeights*. While binned fits are not the main topic of this paper, it should be noted that a similar expression also can be necessary when performing a binned fit (for example a binned χ^2 fit) of *sWeighted* histograms. This is discussed in detail in App. C.

4 Examples

4.1 Correcting for an acceptance effect with event weights

The first example discussed in this paper is the fit of an angular distribution to determine angular coefficients, using event weights to correct for an acceptance effect. The probability density function used to generate and fit the pseudoexperiments is a simple second order polynomial in the angle $\cos \theta$:

$$\begin{aligned} \mathcal{P}(\cos \theta | c_0, c_1) &= (1 + c_0 \cos \theta + c_1 \cos^2 \theta) / \mathcal{N} \quad \text{with} \\ \mathcal{N} &= \int_{-1}^{+1} (1 + c_0 \cos \theta + c_1 \cos^2 \theta) d\cos \theta = 2 + \frac{2}{3}c_1 \end{aligned} \quad (53)$$




Figure 1: Angular $\cos \theta$ distribution of (black) data and (red) efficiency corrected events for 10 000 pseudoexperiments consisting of 1 000 events each.

In the generation, the values $c_0^{\text{gen}} = 0$ and $c_1^{\text{gen}} = 0$ are used. Events are generated using a $\cos \theta$ -dependent efficiency $\epsilon(\cos \theta)$. Two efficiencies shapes are studied, given by

- (a) $\epsilon(\cos \theta) = 1.0 - 0.7 \cos^2 \theta$ and
- (b) $\epsilon(\cos \theta) = 0.3 + 0.7 \cos^2 \theta$.

For simplicity, no uncertainty is assumed on the description of the acceptance effect by $\epsilon(\cos \theta)$, otherwise the effect of uncertainties on event weights would need to be included as described in Sec. 3.1. Figure 1 shows the generated data (including the acceptance effect) in black and the efficiency corrected distributions, weighted by $w_e = 1/\epsilon(\cos \theta_e)$ in red.

The parameters c_0 and c_1 are determined using a weighted unbinned maximum likelihood fit, solving Eq. 3. The uncertainties on the parameters c_0 and c_1 are determined using the approaches to determine parameter uncertainties that are discussed in Sec. 2. The following methods are studied:

- (a) The method of using the uncertainties determined according to Eq. 4 without any correction, denoted as *wFit* in this section.
- (b) Scaling the weights according to Eq. 19. This approach is referred to as *scaled weights*.
- (c) Determining the covariance matrix using Eq. 20. This method is referred to as *squared correction* in the following.
- (d) Bootstrapping the data (using 1000 bootstraps) with replacement, denoted as *bootstrapping*.

- (e) The method to determine the covariance according to Eq. 18 as discussed in Sec. 2.1, referred to as *asymptotic* method.
- (f) A conventional fit (*cFit*) modelling the efficiency correction effect in the probability density function (and its normalisation) instead of using event weights.


The performance of the methods is compared using pseudoexperiments, with each study consisting of 10 000 toy data samples. The same data samples are used for every method. The distribution of the pull, defined as $p_i(c_0) = (c_{0,i} - c_0^{\text{gen}})/\sigma_i(c_0)$ (and analogously for parameter c_1), is used to test the different methods for uncertainty estimation. Here, the fitted value for parameter c_0 in pseudoexperiment i is denoted as $c_{0,i}$, the corresponding uncertainty is denoted as $\sigma_i(c_0)$ and the generated value as c_0^{gen} . If the fit is unbiased and the uncertainties are determined correctly, the pull distribution is expected to be a Gaussian distribution with a mean compatible with zero and a width compatible with one. Different event yields per data sample ($N = 1\,000, 2\,000, 5\,000, 10\,000, 20\,000, 50\,000$) are studied to investigate the influence of statistics.

The pull distributions for the parameters c_0 and c_1 for 2 000 events are shown in Figs. 2 and 3. The pull means and widths depending on statistics are given in Figs. 4 and 5. Numerical values are given in Tabs. 3 and 4 in App. D. A few remarks are in order.


The *wFit* method is unbiased but shows significant undercoverage for c_0 and c_1 for both acceptance corrections tested. The *scaled weights* approach shows significant undercoverage for both c_0 and c_1 for the acceptance (a) and overcoverage for acceptance (b). In both cases, the coverage remains incorrect even for high statistics. Both the use of the *wFit* as well as the *scaled weights* methods are therefore strongly disfavoured to determine the parameter uncertainties in this example for a simple efficiency correction. The *squared method* shows good behaviour for parameter c_0 but incorrect coverage for parameter c_1 . For parameter c_1 the method shows overcoverage for acceptance (a) and very significant undercoverage (more severe than even the *wFit*) for acceptance (b). The reason for this behaviour is the different expectation value of Eq. 26 with respect to the second derivatives to c_0 and c_1 , as detailed in App. A. This illustrates that the *squared correction* method, which is widely used in particle physics, in general does not provide asymptotically correct confidence intervals when using event weights to correct for acceptance effects. *Bootstrapping* the data sample or using the *asymptotic* approach results in unbiased pull distributions with correct coverage already at low statistics for both c_0 and c_1 and both acceptance effects. This paper therefore advocates for the use of the *asymptotic* method (or alternatively *bootstrapping*) when using event weights to account for acceptance corrections. The pull distributions for the *cFit* also show, as expected, good behaviour. As there is no loss of information for the *cFit*, it can result in better sensitivity and its use should be strongly considered, where feasible.

4.2 Background subtraction using *sWeights*

As second specific example for the determination of confidence intervals in the presence of event weights, the determination of the lifetime τ of an exponential decay in the presence




(a) c_0 pull distributions




(b) c_1 pull distributions

Figure 2: Pull distributions from 10 000 pseudoexperiments for the different approaches to the uncertainty estimation for the efficiency correction $\epsilon(\cos \theta) = 1.0 - 0.7 \cos^2 \theta$ at a total yield of 2 000 events for each pseudoexperiment.




(a) c_0 pull distributions




(b) c_1 pull distributions

Figure 3: Pull distributions from 10 000 pseudoexperiments for the different approaches to the uncertainty estimation for the efficiency correction $\epsilon(\cos \theta) = 0.3 + 0.7 \cos^2 \theta$ at a total yield of 2000 events for each pseudoexperiment.




(a) c_0 pull mean and width




(b) c_1 pull mean and width

Figure 4: (Left) pull means and (right) pull widths for the efficiency correction $\epsilon(\cos \theta) = 1.0 - 0.7 \cos^2 \theta$, depending on total event yield N_{tot} .



(a) c_0 pull mean and width



(b) c_1 pull mean and width

Figure 5: (Left) pull means and (right) pull widths for the efficiency correction $\epsilon(\cos \theta) = 0.3 + 0.7 \cos^2 \theta$, depending on total event yield N_{tot} .




Figure 6: Discriminating mass distribution for (black) the full data, (blue) signal and (red) background.

parameter	value
$f_{\text{sig}} = N_{\text{sig}}/(N_{\text{sig}} + N_{\text{bkg}})$	0.5
α_{bkg}	0.0
$m(B)$	$5.367 \text{ GeV}/c^2$
$\sigma(m)$	$23 \text{ MeV}/c^2$
mass range	$[5\ 337, 5\ 567] \text{ MeV}/c^2$
$\tau_{\text{sig}}^{\text{gen}}$	1.5 ps
t range	$[0, 10] \text{ ps}$

(a) Parameters used in the generation

	N_{sig}	N_{bkg}	α_{bkg}
N_{sig}	1.00	-0.63	-0.67
N_{bkg}	-0.63	1.00	0.67
α_{bkg}	-0.67	0.67	1.00

(b) Mean correlation matrix

Table 1: (Left) the parameters used in the generation of the pseudoexperiments. Only N_{sig} , N_{bkg} , and the background slope α_{bkg} are varied in the mass fit. The background slope α_{bkg} is then fixed for the determination of the *sWeights*. (Right) the mean correlation matrix from the mass fit when both the yields and α_{bkg} are allowed to float.

of background is discussed. The *sPlot* method [3] is used to statistically subtract the background component. As *discriminating variable*, the reconstructed mass is used. In this example, the signal is distributed according to a Gaussian in the reconstructed mass, and the background is described by a single Exponential function with slope α_{bkg} . Figure 6 shows the mass distribution for signal and background components. The parameters used in the generation of the pseudoexperiments are listed in Tab. 1a. The configuration is purposefully chosen such that there is a significant correlation between the yields and the slope of the background exponential, to illustrate the effect of fixing nuisance parameters in the *sPlot* formalism, as discussed in Sec. 3.2. The resulting mean correlation matrix for the mass fit is shown in Tab. 1b. The simpler case, where no significant correlation between α_{bkg} and the event yields is present, due to a different choice of mass range, is discussed in App. F.

The decay time distribution (the *control variable*) that is used to determine the




Figure 7: Decay time distributions for the four different background options for (black) the full data, (blue) signal and (red) background. The signal and background components are obtained using *sWeights*.

lifetime is a single Exponential for the signal. For the background component, several different shapes were tested: (a) A single Exponential with long lifetime, (b) a Gaussian distribution, (c) a triangular distribution, and (d) a flat distribution in the decay time. Figure 7 shows the decay time distribution for the different options. The decay time distributions for signal and background components shown are obtained using the *sPlot* formalism [3] described in Sec. 3.2.

The parameter τ is determined using a weighted unbinned maximum likelihood fit solving the maximum likelihood condition Eq. 3 numerically. Its uncertainty $\sigma(\tau)$ is determined using the different methods for weighted unbinned maximum likelihood fits discussed in Sec. 2. The following approaches are studied:

- (a) A weighted fit determining the uncertainties according to Eq. 4 without any correction. This method is denoted as *sFit* in the following.


- (b) Scaling the weights according to Eq. 19. The approach is denoted as *scaled weights*.
- (c) Determining the covariance matrix using Eq. 20. This method is referred to as *squared correction*.
- (d) Bootstrapping the data (using 1000 bootstraps) with replacement, without rederiving the *sWeights* (*i.e.* keeping the original *sWeights* for each event). Denoted as *bootstrapping* in the following.
- (e) Bootstrapping the data (again using 1000 bootstraps) and rederiving the *sWeights* for every bootstrapped sample, in the following denoted as *full bootstrapping*.
- (f) The asymptotic method to determine the covariance according to Eq. 18 as discussed in Sec. 2.1, but not accounting for the uncertainty of the *sWeights*. This approach is referred to as *asymptotic* method.
- (g) The asymptotic method to determine the covariance, including the additional covariance due to the uncertainty of the *sWeights* as given by Eq. 52. This method is denoted as *asymptotic corrected*.
- (h) A conventional fit (*cFit*) modelling both signal and background components in two dimensions (mass and decay time) for comparison. As the main point of using *sWeights* is to remove the need to model the background contribution in the fit, this method is given purely for comparison.

The performance of the different methods is evaluated using pseudoexperiments. Every study consists of 10 000 data samples generated and then fit for an initial determination of the *sWeights*. For every method, the same data samples are used.


The performance of the different methods is compared using the distribution of the pull, defined as $p_i(\tau) = (\tau_i - \tau_{\text{sig}}^{\text{gen}})/\sigma_i(\tau)$. Here, τ_i is the central value determined by the weighted maximum likelihood fit and $\sigma_i(\tau)$ the uncertainty determined by the above methods. The lifetime used in the generation is denoted as $\tau_{\text{sig}}^{\text{gen}}$. To study the influence of statistics, pseudoexperiments are performed for different numbers of events. The total yields $N_{\text{tot}} = N_{\text{sig}} + N_{\text{bkg}}$ generated correspond to 1 000, 2 000, 4 000, 10 000 and 20 000 events. The signal fraction used in the generation is $f_{\text{sig}} = N_{\text{sig}}/(N_{\text{sig}} + N_{\text{bkg}}) = 0.5$.

The pull distributions from 10 000 pseudoexperiments, each with a total yield of 2000 events, are shown in Fig. 8 and 9. The pull means and widths are shown in Figs. 10 and 11. Numerical values for the different configurations are given in Tabs. 5 and 6 in App. E.

As expected, both the *sFit* as well as the approach using *scaled weights* perform quite poorly, as they show large undercoverage, both for low statistics as well as for high statistics. Furthermore, they exhibit significant bias at low statistics (which reduces at large statistics) due to a strong correlation of the uncertainty with the parameter τ . This strongly disfavours the use of these methods for these *sWeighted* examples. The *squared correction* method shows better performance, nevertheless also exhibits significant bias (which reduces for higher statistics) and undercoverage. It should be stressed




(a) Exponential background model




(b) Gaussian background model

Figure 8: Pull distributions from 10 000 pseudoexperiments for the different approaches to the uncertainty estimation for a total yield of 2 000 events in each pseudoexperiment. The different figures shown correspond to the different background models as specified in Sec. 4.




(a) Triangular background model




(b) Flat background model

Figure 9: Pull distributions from 10 000 pseudoexperiments for the different approaches to the uncertainty estimation for a total yield of 2 000 events in each pseudoexperiment. The different figures shown correspond to the different background models as specified in Sec. 4.




(a) Exponential background model




(b) Gaussian background model

Figure 10: (Left) pull means and (right) pull widths depending on total event yield N_{tot} .



(a) Triangular background model



(b) Flat background model

Figure 11: (Left) pull means and (right) pull widths depending on total event yield N_{tot} .

that significant undercoverage is still present at large statistics. This shows that the *squared correction* method in general does not provide asymptotically correct confidence intervals. Both *bootstrapping* as well as the *asymptotic* methods perform better for the examples studied here. However, both methods show some remaining undercoverage even at high statistics. It is instructive that *bootstrapping* the data without redetermining the *sWeights* performs identically to the *asymptotic method* without accounting for the uncertainty due to nuisance parameters. However, when performing a *full bootstrapping* including rederiving the *sWeights* for the bootstrapped samples or when using the *corrected asymptotic* method the confidence intervals cover correctly already at low statistics and no significant biases are observed. This paper therefore advocates the use of the *corrected asymptotic* method, or alternatively, if computationally possible, the *full bootstrapping* approach for the determination of uncertainties in unbinned maximum likelihood fits using *sWeights*. If nuisance parameters have no large impact on the the *sWeights*, the *asymptotic* method is also appropriate, as shown in App. F.

The conventional fit describing the background component in the decay time explicitly instead of using *sWeights* also shows good behaviour, as expected. When the background distribution is known, a conventional fit is generally advantageous as it has improved sensitivity due to the additional available information. For this example, where the background pollution and parameter correlations are large, the parameter sensitivity is significantly improved when using the conventional (unweighted) fit.

5 Conclusions

This paper derives the asymptotically correct method to determine parameter uncertainties in the presence of event weights, which was previously discussed in Ref. [4] for acceptance corrections but does not currently see widespread use in high energy particle physics. The performance of this approach is compared with several commonly used methods, using two examples that are typical use cases: The correction of an acceptance effect using event weights and statistically subtracting background events using the *sPlot* formalism [3]. For both examples, the asymptotically correct method performs well while several of the commonly used methods are shown to not generally result in correct coverage, even for large statistics. If statistics are sufficiently large this paper therefore advocates the use of the asymptotically correct method in weighted unbinned maximum likelihood fits, in particular over the current nominal method used in the ROOFIT [10] fitting framework, which was proposed in Refs. [1, 2], and is shown to not generally result in asymptotically correct uncertainties. If computationally feasible, the bootstrapping approach [11] can be a useful alternative. In addition, the inclusion of event weight uncertainties in weighted unbinned maximum likelihood fits for both the *sPlot* formalism and when using weights to account for an acceptance effect is discussed. In particular, an expression is given that corrects the covariance for the presence of nuisance parameters that can affect the *sWeights*. A similar expression is also given for the case of a χ^2 fit of binned *sWeighted* data. A patch for ROOFIT to allow the determination of the covariance matrix according to Eq. 18 is available from the author on request.

6 Acknowledgements

C. L. gratefully acknowledges support by the Emmy Noether programme of the Deutsche Forschungsgemeinschaft (DFG), grant identifier LA 3937/1-1. Furthermore, C. L. would like to thank Roger Barlow for helpful comments and questions on an early version of this paper.

References

- [1] W. T. Eadie, D. Drijard, F. E. James, M. Roos and B. Sadoulet, *Statistical methods in experimental physics*. North-Holland, Amsterdam, 1971.
- [2] F. James, *Statistical methods in experimental physics*. Hackensack, USA: World Scientific (2006) 345 p, 2006.
- [3] M. Pivk and F. R. Le Diberder, *SPlot: A Statistical tool to unfold data distributions*, *Nucl. Instrum. Meth.* **A555** (2005) 356 [physics/0402083].
- [4] F. T. Solmitz, *Analysis of experiments in particle physics*, *Annual Review of Nuclear Science* **14** (1964) 375.
- [5] F. James and M. Roos, *Minuit: A System for Function Minimization and Analysis of the Parameter Errors and Correlations*, *Comput. Phys. Commun.* **10** (1975) 343.
- [6] A. C. Davison, *Statistical Models*, Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2003, 10.1017/CBO9780511815850.
- [7] V. P. Godambe, *An optimum property of regular maximum likelihood estimation*, *Ann. Math. Statist.* **31** (1960) 1208.
- [8] V. Godambe and M. Thompson, *Some aspects of the theory of estimating equations*, *Journal of Statistical Planning and Inference* **2** (1978) 95 .
- [9] R. J. Barlow, *Statistics: a guide to the use of statistical methods in the physical sciences*, Manchester physics series. Wiley, Chichester, 1989.
- [10] W. Verkerke and D. P. Kirkby, *The RooFit toolkit for data modeling*, *eConf* **C0303241** (2003) MOLT007 [physics/0306116].
- [11] B. Efron, *Bootstrap methods: Another look at the jackknife*, *The Annals of Statistics* **7** (1979) 1.
- [12] Y. Xie, *sFit: a method for background subtraction in maximum likelihood fit*, *arXiv e-prints* (2009) arXiv:0905.0724 [0905.0724].

A Expectation value Eq. 26 for examples correcting for acceptance effects

As mentioned in Sec. 2.2, Eq. 22 is not generally asymptotically valid. To demonstrate this with an example, the expectation value in Eq. 26 is explicitly calculated below for the angular fit in Sec. 4.1. Using the probability density function

$$\mathcal{P}(\cos \theta | c_0, c_1) = \frac{1 + c_0 \cos \theta + c_1 \cos^2 \theta}{2 + \frac{2}{3}c_1} \quad (54)$$

and the efficiency correction (b)

$$\epsilon(\cos \theta) = \frac{3}{10} + \frac{7}{10} \cos^2 \theta \quad (55)$$

we derive the expectation value in Eq. 26 according to (showing the asymptotic behaviour for the double partial derivative to c_1 is sufficient):

$$\begin{aligned} & \left\langle \frac{1}{\epsilon^2(\cos \theta_e) \mathcal{P}(\cos \theta_e | c_0, c_1)} \frac{\partial^2 \mathcal{P}(\cos \theta_e | c_0, c_1)}{\partial c_1^2} \right\rangle \\ &= \frac{\int_{-1}^{+1} \frac{1}{\epsilon^2(\cos \theta) \mathcal{P}(\cos \theta | c_0, c_1)} \frac{\partial^2 \mathcal{P}(\cos \theta | c_0, c_1)}{\partial c_1^2} \mathcal{P}(\cos \theta | c_0, c_1) \epsilon(\cos \theta) d\cos \theta}{\int_{-1}^{+1} \epsilon(\cos \theta) \mathcal{P}(\cos \theta | c_0, c_1) d\cos \theta} \\ &= \frac{\frac{\partial^2}{\partial c_1^2} \int_{-1}^{+1} \frac{1}{\frac{3}{10} + \frac{7}{10} \cos^2 \theta} \frac{1 + c_0 \cos \theta + c_1 \cos^2 \theta}{2 + \frac{2}{3}c_1} d\cos \theta}{\int_{-1}^{+1} \left(\frac{3}{10} + \frac{7}{10} \cos^2 \theta \right) \frac{1 + c_0 \cos \theta + c_1 \cos^2 \theta}{2 + \frac{2}{3}c_1} d\cos \theta}. \end{aligned} \quad (56)$$

Using computer algebra, these integrals can be easily evaluated analytically. For the denominator we obtain

$$\int_{-1}^{+1} \left(\frac{3}{10} + \frac{7}{10} \cos^2 \theta \right) \frac{1 + c_0 \cos \theta + c_1 \cos^2 \theta}{2 + \frac{2}{3}c_1} d\cos \theta = \frac{36}{50} \frac{c_1}{c_1 + 3} + \frac{48}{30} \frac{1}{c_1 + 3} \stackrel{c_1=0}{=} \frac{8}{15}. \quad (57)$$

The expression for the numerator is slightly more complicated, it results in

$$\begin{aligned} & \frac{\partial^2}{\partial c_1^2} \int_{-1}^{+1} \frac{1}{\frac{3}{10} + \frac{7}{10} \cos^2 \theta} \frac{1 + c_0 \cos \theta + c_1 \cos^2 \theta}{2 + \frac{2}{3}c_1} d\cos \theta \\ &= \frac{\partial^2}{\partial c_1^2} \left(\frac{1}{\frac{2}{3}c_1 + 2} \left[-\frac{20\sqrt{21}}{49} \tan^{-1} \left(\frac{\sqrt{21}}{3} \right) c_1 + \frac{20}{7} c_1 + \frac{20}{\sqrt{21}} \tan^{-1} \left(\frac{\sqrt{21}}{3} \right) \right] \right) \\ &\stackrel{c_1=0}{=} \frac{20}{7\sqrt{21}} \tan^{-1} \left(\frac{\sqrt{21}}{3} \right) + \frac{140}{3 \cdot 21^{3/2}} \tan^{-1} \left(\frac{\sqrt{21}}{3} \right) - \frac{20}{21} \\ &\approx 0.146. \end{aligned} \quad (58)$$

We thus find

$$\left\langle \frac{1}{\epsilon^2(\cos \theta_e)\mathcal{P}(\cos \theta_e|c_0, c_1)} \frac{\partial^2 \mathcal{P}(\cos \theta_e|c_0, c_1)}{\partial c_1^2} \right\rangle \approx 0.274 \neq 0. \quad (59)$$

This shows clearly, that Eq. 22 is not generally asymptotically correct. The result in Eq. 59 has been crosschecked using the pseudoexperiments described in Sec. 4.1 and indeed the additional term fluctuates around $N \times 0.2742$ as derived above.

For acceptance (a) we find similarly

$$\left\langle \frac{1}{\epsilon^2(\cos \theta_e)\mathcal{P}(\cos \theta_e|c_0, c_1)} \frac{\partial^2 \mathcal{P}(\cos \theta_e|c_0, c_1)}{\partial c_1^2} \right\rangle \approx -0.135 \neq 0, \quad (60)$$

which is also confirmed using the pseudoexperiments. The fact that the expectation value is negative for acceptance (a) and positive for acceptance (b) indicates that, as observed using the pseudoexperiments, the *squared correction* method overcovers for acceptance (a) and undercovers for acceptance (b).

It is instructive to note that for the double partial derivative to c_0 , $\langle \epsilon^{-2} \mathcal{P}^{-1} \partial^2 / \partial c_0^2 \mathcal{P} \rangle$, we find an expectation value of zero, as the integration of the numerator removes the c_0 dependence in that case. This is the reason why the *squared correction* method results in compatible results with the *asymptotic* method for c_0 , but shows incorrect coverage for c_1 .

B Expectation value Eq. 26 for examples using *sWeights*

For the examples in this paper using *sWeights* discussed in Sec. 4.2 we explicitly calculate the expectation value in Eq. 26 as detailed below. With the total probability density function

$$\mathcal{P}(t, m|\tau) = f_{\text{sig}}\mathcal{P}_{\text{sig}}(t|\tau)\mathcal{P}_{\text{sig}}(m) + (1 - f_{\text{sig}})\mathcal{P}_{\text{bkg}}(t)\mathcal{P}_{\text{bkg}}(m) \quad (61)$$

we find

$$\begin{aligned} \left\langle w_{\text{sig}}^2(m) \frac{1}{\mathcal{P}_{\text{sig}}(t|\tau)} \frac{\partial^2 \mathcal{P}_{\text{sig}}(t|\tau)}{\partial \tau^2} \right\rangle &= \int w_{\text{sig}}^2(m) \frac{1}{\mathcal{P}_{\text{sig}}(t|\tau)} \frac{\partial^2 \mathcal{P}_{\text{sig}}(t|\tau)}{\partial \tau^2} \\ &\quad \times [f_{\text{sig}}\mathcal{P}_{\text{sig}}(t|\tau)\mathcal{P}_{\text{sig}}(m) + (1 - f_{\text{sig}})\mathcal{P}_{\text{bkg}}(t)\mathcal{P}_{\text{bkg}}(m)] dt dm \\ &= \int \frac{1}{\mathcal{P}_{\text{sig}}(t|\tau)} \frac{\partial^2 \mathcal{P}_{\text{sig}}(t|\tau)}{\partial \tau^2} (1 - f_{\text{sig}})\mathcal{P}_{\text{bkg}}(t) dt \\ &\quad \times \int w_{\text{sig}}^2(m)\mathcal{P}_{\text{bkg}}(m) dm. \end{aligned} \quad (62)$$

The integral over the decay time depends on the background model $\mathcal{P}_{\text{bkg}}(t)$, and the (common) signal PDF is given by

$$\mathcal{P}_{\text{sig}}(t|\tau_{\text{sig}}) = \frac{e^{-t/\tau_{\text{sig}}}}{\tau_{\text{sig}} (e^{-t_{\text{min}}/\tau_{\text{sig}}} - e^{-t_{\text{max}}/\tau_{\text{sig}}})}. \quad (63)$$

The evaluation of the integral over the decay time results in numerical values of

$$\begin{aligned} \int \frac{1}{\mathcal{P}_{\text{sig}}(t|\tau)} \frac{\partial^2 \mathcal{P}_{\text{sig}}(t|\tau)}{\partial \tau^2} (1 - f_{\text{sig}})\mathcal{P}_{\text{bkg}}^{\text{exponential}}(t) dt &= 0.328, \\ \int \frac{1}{\mathcal{P}_{\text{sig}}(t|\tau)} \frac{\partial^2 \mathcal{P}_{\text{sig}}(t|\tau)}{\partial \tau^2} (1 - f_{\text{sig}})\mathcal{P}_{\text{bkg}}^{\text{Gaussian}}(t) dt &= 0.559, \\ \int \frac{1}{\mathcal{P}_{\text{sig}}(t|\tau)} \frac{\partial^2 \mathcal{P}_{\text{sig}}(t|\tau)}{\partial \tau^2} (1 - f_{\text{sig}})\mathcal{P}_{\text{bkg}}^{\text{triangular}}(t) dt &= 0.608, \\ \int \frac{1}{\mathcal{P}_{\text{sig}}(t|\tau)} \frac{\partial^2 \mathcal{P}_{\text{sig}}(t|\tau)}{\partial \tau^2} (1 - f_{\text{sig}})\mathcal{P}_{\text{bkg}}^{\text{flat}}(t) dt &= 0.775. \end{aligned} \quad (64)$$

The integral over the mass is identical for all background models. It results in a numerical value of 0.47 for the case of significant correlations discussed in Sec. 4.2 and 0.77 for the case of negligible correlations discussed in App. F. This shows that, as expected from the pseudoexperiments, the *squared correction* undercovers for all background models studied in this paper. The amount of undercoverage can be particularly easily compared for the case of negligible correlations, and is indeed found to be correctly reproduced using the values above.

C Binned *sWeighted* fits

Instead of performing an *sWeighted* unbinned maximum likelihood fit, an alternative is to perform a binned χ^2 fit of an *sWeighted* distribution. The χ^2 to minimise in the fit is given by

$$\begin{aligned}\chi^2 &= \sum_{k=1}^{N_B} \frac{[N_k^{\text{data}} - N_k^{\text{pred.}}(\vec{\lambda})]^2}{\sigma_i^2} \\ &= \sum_{k=1}^{N_B} \frac{[\sum_{e \in \text{bin } k} w_e - N_k^{\text{pred.}}(\vec{\lambda})]^2}{\sum_{e \in \text{bin } k} w_e^2},\end{aligned}\quad (65)$$

where N_B denotes the number of bins. The χ^2 minimisation corresponds to the condition

$$\left. \frac{\partial}{\partial \lambda_i} \chi^2 \right|_{\hat{\lambda}} = 0. \quad (66)$$

Expanding this equation around the true value $\vec{\lambda}_0$ yields

$$\begin{aligned}\frac{\partial}{\partial \lambda_i} \left[\chi^2(\vec{\lambda}_0) + \sum_{j=1}^{N_P} (\hat{\lambda}_j - \lambda_{0j}) \frac{\partial}{\partial \lambda_j} \chi^2(\vec{\lambda}_0) \right] &= 0 \\ \sum_{j=1}^{N_P} (\hat{\lambda}_j - \lambda_{0j}) \frac{\partial^2}{\partial \lambda_i \partial \lambda_j} \chi^2(\vec{\lambda}_0) &= -\frac{\partial}{\partial \lambda_i} \chi^2(\vec{\lambda}_0).\end{aligned}\quad (67)$$

The deviation of the estimator from the true value is thus given by

$$\begin{pmatrix} \hat{\lambda}_1 - \lambda_{01} \\ \vdots \\ \hat{\lambda}_{N_P} - \lambda_{0N_P} \end{pmatrix} = \begin{pmatrix} \frac{\partial^2}{\partial \lambda_1^2} \chi^2 & \cdots & \frac{\partial^2}{\partial \lambda_1 \partial \lambda_{N_P}} \chi^2 \\ \vdots & \ddots & \vdots \\ \frac{\partial^2}{\partial \lambda_{N_P} \partial \lambda_1} \chi^2 & \cdots & \frac{\partial^2}{\partial \lambda_{N_P}^2} \chi^2 \end{pmatrix}^{-1} \begin{pmatrix} \frac{\partial}{\partial \lambda_1} \chi^2 \\ \vdots \\ \frac{\partial}{\partial \lambda_{N_P}} \chi^2 \end{pmatrix}, \quad (68)$$

and the covariance matrix by

$$\begin{aligned}V_{ij} &= \langle (\hat{\lambda}_i - \lambda_{0i})(\hat{\lambda}_j - \lambda_{0j}) \rangle \\ &= \left(\frac{\partial^2}{\partial \lambda_i \partial \lambda_k} \chi^2 \right)^{-1} \left\langle \left(\frac{\partial}{\partial \lambda_k} \chi^2 \right) \left(\frac{\partial}{\partial \lambda_l} \chi^2 \right) \right\rangle \left(\frac{\partial^2}{\partial \lambda_l \partial \lambda_j} \chi^2 \right)^{-1}.\end{aligned}\quad (69)$$

The expectation value can be calculated according to

$$\begin{aligned}
\left\langle \frac{\partial}{\partial \lambda_i} \chi^2 \frac{\partial}{\partial \lambda_j} \chi^2 \right\rangle &= \left\langle \left(\frac{\partial}{\partial \lambda_i} \sum_{k=1}^{N_B} \frac{(N_k^{\text{data}} - N_k^{\text{pred.}})^2}{\sigma_k^2} \right) \left(\frac{\partial}{\partial \lambda_j} \sum_{l=1}^{N_B} \frac{(N_l^{\text{data}} - N_l^{\text{pred.}})^2}{\sigma_l^2} \right) \right\rangle \\
&= \left\langle 4 \left(\sum_{k=1}^{N_B} \frac{N_k^{\text{data}} - N_k^{\text{pred.}}}{\sigma_k^2} \frac{\partial N_k^{\text{pred.}}}{\partial \lambda_i} \right) \left(\sum_{l=1}^{N_B} \frac{N_l^{\text{data}} - N_l^{\text{pred.}}}{\sigma_l^2} \frac{\partial N_l^{\text{pred.}}}{\partial \lambda_j} \right) \right\rangle \\
&= \left\langle 4 \sum_{k=1}^{N_B} \left(\frac{N_k^{\text{data}} - f_k^{\text{pred.}} \sum_{r=1}^{N_B} N_r^{\text{data}}}{\sigma_k^2} \frac{\partial f_k^{\text{pred.}}}{\partial \lambda_i} \sum_{r=1}^{N_B} N_r^{\text{data}} \right) \right. \\
&\quad \left. \times \sum_{l=1}^{N_B} \left(\frac{N_l^{\text{data}} - f_l^{\text{pred.}} \sum_{r=1}^{N_B} N_r^{\text{data}}}{\sigma_l^2} \frac{\partial f_l^{\text{pred.}}}{\partial \lambda_j} \sum_{r=1}^{N_B} N_r^{\text{data}} \right) \right\rangle. \tag{70}
\end{aligned}$$

In the last line, the predicted fraction for each bin, $f_{k(l)}(\vec{\lambda})$ is introduced, as the total normalisation for the χ^2 fit is given by the sum of the *sWeights* in all bins, $\sum_{r=1}^{N_B} N_r^{\text{data}}$, and therefore $N_{k(l)}^{\text{data}} = f_{k(l)}(\vec{\lambda}) \sum_{r=1}^{N_B} N_r^{\text{data}} = f_{k(l)}(\vec{\lambda}) \sum_{e=1}^N w_e$.

In full analogy to the unbinned case one can expand the *sWeights* using Eq. 49 to account for the uncertainties of nuisance parameters. This results in the corrected covariance matrix

$$\begin{aligned}
V_{ij} &= \left(\frac{\partial^2}{\partial \lambda_i \partial \lambda_k} \chi^2 \right)^{-1} \left\langle \left(\frac{\partial}{\partial \lambda_k} \chi^2 \Big|_{\hat{p}} \right) \left(\frac{\partial}{\partial \lambda_l} \chi^2 \Big|_{\hat{p}} \right) \right\rangle \left(\frac{\partial^2}{\partial \lambda_l \partial \lambda_j} \chi^2 \right)^{-1} \\
&\quad + \left(\frac{\partial^2}{\partial \lambda_i \partial \lambda_k} \chi^2 \right)^{-1} 4 \sum_{r=1}^{N_B} \left(\frac{\sum_{e \in \text{bin } r} \frac{\partial w_e}{\partial p_m} - f_r \sum_{e=1}^N \frac{\partial w_e}{\partial p_m}}{\sigma_r^2} \frac{\partial f_r^{\text{pred.}}}{\partial \lambda_k} \sum_{e=1}^N w_e \right) \times \Sigma_{mn} \\
&\quad \times \sum_{s=1}^{N_B} \left(\frac{\sum_{e \in \text{bin } s} \frac{\partial w_e}{\partial p_n} - f_s \sum_{e=1}^N \frac{\partial w_e}{\partial p_n}}{\sigma_s^2} \frac{\partial f_s^{\text{pred.}}}{\partial \lambda_l} \sum_{e=1}^N w_e \right) \left(\frac{\partial^2}{\partial \lambda_l \partial \lambda_j} \chi^2 \right)^{-1} \tag{71}
\end{aligned}$$

Here, Σ_{mn} is the covariance matrix difference given in Eq. 51, and the sum over the indices m and n is implicit.

This expression is studied using the same pseudoexperiments as described in Sec. 4.2. The lifetime is determined by performing a binned χ^2 fit using ten bins of the decay time. When performing this χ^2 fit a constant offset, c , is allowed in the fit as otherwise a bias will be present due to the fact that an Exponential function will always be positive, whereas the *sWeighted* bins can become negative due to statistical fluctuations. The fraction f_k in bin k (spanning the range $[t_k, t_{k+1})$) is thus given by

$$f_k(\tau, c) = \frac{\int_{t_k}^{t_{k+1}} \frac{e^{-t/\tau}}{\tau} + c}{\int_{t_{\min}}^{t_{\max}} \frac{e^{-t/\tau}}{\tau} + c} \tag{72}$$

and both the parameters τ and c are determined in the fit. The performance of the χ^2 fit without the correction due to the nuisance parameters is compared with the corrected

method	pull	1 k	2 k	4 k	10 k	20 k
χ^2	mean	-0.12 ± 0.01	-0.08 ± 0.01	-0.07 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01
	width	1.04 ± 0.01	1.04 ± 0.01	1.02 ± 0.01	1.02 ± 0.01	1.02 ± 0.01
χ^2 corrected	mean	-0.11 ± 0.01	-0.08 ± 0.01	-0.06 ± 0.01	-0.04 ± 0.01	-0.02 ± 0.01
	width	1.02 ± 0.01	1.02 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01

(a) Exponential background model

method	pull	1 k	2 k	4 k	10 k	20 k
χ^2	mean	-0.20 ± 0.01	-0.12 ± 0.01	-0.09 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01
	width	1.11 ± 0.01	1.14 ± 0.01	1.14 ± 0.01	1.15 ± 0.01	1.14 ± 0.01
χ^2 corrected	mean	-0.20 ± 0.01	-0.12 ± 0.01	-0.09 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01
	width	1.02 ± 0.01	1.02 ± 0.01	1.01 ± 0.01	1.01 ± 0.01	1.01 ± 0.01

(b) Gaussian background model

method	pull	1 k	2 k	4 k	10 k	20 k
χ^2	mean	-0.19 ± 0.03	-0.17 ± 0.01	-0.12 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01
	width	3.33 ± 0.02	1.14 ± 0.01	1.19 ± 0.01	1.22 ± 0.01	1.22 ± 0.01
χ^2 corrected	mean	-0.21 ± 0.01	-0.16 ± 0.01	-0.12 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01
	width	1.06 ± 0.01	1.01 ± 0.01	1.01 ± 0.01	1.02 ± 0.01	1.02 ± 0.01

(c) Triangular background model

method	pull	1 k	2 k	4 k	10 k	20 k
χ^2	mean	-0.12 ± 0.01	-0.07 ± 0.01	-0.06 ± 0.01	-0.03 ± 0.01	-0.01 ± 0.01
	width	1.02 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01
χ^2 corrected	mean	-0.12 ± 0.01	-0.07 ± 0.01	-0.06 ± 0.01	-0.03 ± 0.01	-0.01 ± 0.01
	width	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01

(d) Flat background model

Table 2: Results from a binned χ^2 fit of the pseudoexperiments discussed in Sec. 4.2.

covariance given by Eq. 71. Table 2 gives the results for the four different background models. As is apparent, the corrected covariance yields asymptotically correct coverage in this example, whereas the uncorrected approach shows undercoverage for the Gaussian and triangular background model.

As an aside, it should be noted that a binned χ^2 fit can be advantageous if the background events that are statistically subtracted using the *sPlot* formalism would have high impact on the signal parameters. In the example of the lifetime determination this is the case when the background events exhibit large decay times. The maximum likelihood estimator in this case can show larger variance than a binned χ^2 approach.

D Results from pseudoexperiments correcting acceptance effects

method	pull	1 k	2 k	5 k	10 k	20 k	50 k
wFit	mean	0.01 ± 0.01	0.01 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01	0.02 ± 0.01	0.01 ± 0.01
	width	1.40 ± 0.01	1.40 ± 0.01	1.37 ± 0.01	1.38 ± 0.01	1.38 ± 0.01	1.37 ± 0.01
scaled weights	mean	0.01 ± 0.01	0.01 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01
	width	1.16 ± 0.01	1.17 ± 0.01	1.14 ± 0.01	1.15 ± 0.01	1.15 ± 0.01	1.14 ± 0.01
squared correction	mean	0.01 ± 0.01	0.01 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.00 ± 0.01
	width	1.01 ± 0.01	1.01 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	0.99 ± 0.01
bootstrapping	mean	0.01 ± 0.01	0.01 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.00 ± 0.01
	width	0.99 ± 0.01	1.01 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	0.99 ± 0.01
asymptotic	mean	0.01 ± 0.01	0.01 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.00 ± 0.01
	width	1.01 ± 0.01	1.01 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	0.99 ± 0.01
cFit	mean	0.01 ± 0.01	0.01 ± 0.01	-0.00 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.00 ± 0.01
	width	1.01 ± 0.01	1.01 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	0.99 ± 0.01

(a) Parameter c_0

method	pull	1 k	2 k	5 k	10 k	20 k	50 k
wFit	mean	-0.11 ± 0.01	-0.07 ± 0.01	-0.06 ± 0.01	-0.05 ± 0.01	-0.02 ± 0.01	-0.03 ± 0.01
	width	1.40 ± 0.01	1.39 ± 0.01	1.38 ± 0.01	1.38 ± 0.01	1.38 ± 0.01	1.39 ± 0.01
scaled weights	mean	-0.10 ± 0.01	-0.06 ± 0.01	-0.05 ± 0.01	-0.04 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01
	width	1.17 ± 0.01	1.16 ± 0.01	1.15 ± 0.01	1.15 ± 0.01	1.15 ± 0.01	1.16 ± 0.01
squared correction	mean	-0.07 ± 0.01	-0.04 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01	-0.01 ± 0.01	-0.02 ± 0.01
	width	0.80 ± 0.01	0.79 ± 0.01	0.79 ± 0.01	0.79 ± 0.01	0.79 ± 0.01	0.79 ± 0.01
bootstrapping	mean	-0.03 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01	-0.03 ± 0.01	-0.01 ± 0.01	-0.02 ± 0.01
	width	0.96 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.01 ± 0.01
asymptotic	mean	-0.07 ± 0.01	-0.05 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01	-0.01 ± 0.01	-0.02 ± 0.01
	width	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01
cFit	mean	-0.05 ± 0.01	-0.03 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01	-0.02 ± 0.01
	width	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01

(b) Parameter c_1

Table 3: Means and widths of the pull distribution for the different approaches to the uncertainty estimation for the efficiency correction $\epsilon(\cos\theta) = 1.0 - 0.7 \cos^2\theta$, depending on the number of events.

method	pull	1 k	2 k	5 k	10 k	20 k	50 k
wFit	mean	0.00 ± 0.01	-0.00 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01	-0.01 ± 0.01	-0.00 ± 0.01
	width	1.22 ± 0.01	1.23 ± 0.01	1.24 ± 0.01	1.24 ± 0.01	1.22 ± 0.01	1.21 ± 0.01
scaled weights	mean	0.00 ± 0.01	-0.00 ± 0.01	-0.00 ± 0.01	0.00 ± 0.01	-0.01 ± 0.01	-0.00 ± 0.01
	width	0.83 ± 0.01	0.83 ± 0.01	0.84 ± 0.01	0.84 ± 0.01	0.83 ± 0.01	0.82 ± 0.01
squared correction	mean	0.00 ± 0.01	-0.00 ± 0.01	-0.00 ± 0.01	0.00 ± 0.01	-0.01 ± 0.01	-0.00 ± 0.01
	width	1.00 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	0.99 ± 0.01
bootstrapping	mean	0.00 ± 0.01	-0.00 ± 0.01	-0.00 ± 0.01	0.00 ± 0.01	-0.01 ± 0.01	-0.00 ± 0.01
	width	0.99 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	0.99 ± 0.01
asymptotic	mean	0.00 ± 0.01	-0.00 ± 0.01	-0.00 ± 0.01	0.00 ± 0.01	-0.01 ± 0.01	-0.00 ± 0.01
	width	0.99 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	0.99 ± 0.01
cFit	mean	0.00 ± 0.01	-0.00 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01	-0.01 ± 0.01	-0.01 ± 0.01
	width	0.99 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	0.99 ± 0.01

(a) Parameter c_0

method	pull	1 k	2 k	5 k	10 k	20 k	50 k
wFit	mean	-0.03 ± 0.01	-0.01 ± 0.01	-0.01 ± 0.01	0.00 ± 0.01	-0.02 ± 0.01	-0.00 ± 0.01
	width	1.36 ± 0.01	1.37 ± 0.01	1.40 ± 0.01	1.38 ± 0.01	1.37 ± 0.01	1.37 ± 0.01
scaled weights	mean	-0.02 ± 0.01	-0.00 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01	-0.01 ± 0.01	-0.00 ± 0.01
	width	0.93 ± 0.01	0.93 ± 0.01	0.95 ± 0.01	0.94 ± 0.01	0.93 ± 0.01	0.93 ± 0.01
squared correction	mean	0.29 ± 0.03	0.21 ± 0.03	0.12 ± 0.03	0.11 ± 0.03	0.04 ± 0.03	0.04 ± 0.03
	width	3.05 ± 0.02	2.92 ± 0.02	2.91 ± 0.02	2.86 ± 0.02	2.84 ± 0.02	2.82 ± 0.02
bootstrapping	mean	0.03 ± 0.01	0.03 ± 0.01	0.01 ± 0.01	0.02 ± 0.01	-0.00 ± 0.01	0.00 ± 0.01
	width	0.98 ± 0.01	0.99 ± 0.01	1.02 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01
asymptotic	mean	-0.03 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01	0.00 ± 0.01	-0.01 ± 0.01	-0.00 ± 0.01
	width	0.99 ± 0.01	1.00 ± 0.01	1.02 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01
cFit	mean	-0.04 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01	-0.00 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01
	width	1.00 ± 0.01	1.00 ± 0.01	1.02 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01

(b) Parameter c_1

Table 4: Means and widths of the pull distribution for the different approaches to the uncertainty estimation for the efficiency correction $\epsilon(\cos\theta) = 0.3 + 0.7 \cos^2\theta$, depending on the number of events.

E Results from pseudoexperiments using $sWeights$

method	pull	1 k	2 k	4 k	10 k	20 k
sFit	mean	-0.50 ± 0.03	-0.35 ± 0.03	-0.25 ± 0.03	-0.16 ± 0.02	-0.13 ± 0.02
	width	2.77 ± 0.02	2.61 ± 0.02	2.53 ± 0.02	2.48 ± 0.02	2.47 ± 0.02
scaled weights	mean	-0.34 ± 0.02	-0.24 ± 0.02	-0.17 ± 0.02	-0.11 ± 0.02	-0.09 ± 0.02
	width	2.16 ± 0.02	2.08 ± 0.01	2.03 ± 0.01	2.00 ± 0.01	1.99 ± 0.01
squared correction	mean	-0.13 ± 0.01	-0.09 ± 0.01	-0.07 ± 0.01	-0.04 ± 0.01	-0.04 ± 0.01
	width	1.44 ± 0.01	1.44 ± 0.01	1.43 ± 0.01	1.42 ± 0.01	1.42 ± 0.01
bootstrapping	mean	-0.06 ± 0.01	-0.04 ± 0.01	-0.04 ± 0.01	-0.02 ± 0.01	-0.03 ± 0.01
	width	1.23 ± 0.01	1.26 ± 0.01	1.27 ± 0.01	1.27 ± 0.01	1.27 ± 0.01
full bootstrapping	mean	-0.05 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01
	width	0.95 ± 0.01	0.98 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	1.00 ± 0.01
asymptotic	mean	-0.06 ± 0.01	-0.04 ± 0.01	-0.04 ± 0.01	-0.02 ± 0.01	-0.03 ± 0.01
	width	1.23 ± 0.01	1.26 ± 0.01	1.27 ± 0.01	1.27 ± 0.01	1.27 ± 0.01
asymptotic corrected	mean	0.00 ± 0.01	0.00 ± 0.01	-0.00 ± 0.01	-0.00 ± 0.01	-0.01 ± 0.01
	width	0.96 ± 0.01	0.98 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	0.99 ± 0.01
cFit	mean	-0.05 ± 0.01	-0.03 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01
	width	1.00 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01

(a) Exponential background model

method	pull	1 k	2 k	4 k	10 k	20 k
sFit	mean	-1.41 ± 0.08	-0.83 ± 0.05	-0.57 ± 0.04	-0.30 ± 0.04	-0.24 ± 0.04
	width	7.86 ± 0.06	4.64 ± 0.03	4.26 ± 0.03	4.05 ± 0.03	4.01 ± 0.03
scaled weights	mean	-0.96 ± 0.06	-0.58 ± 0.04	-0.40 ± 0.03	-0.21 ± 0.03	-0.17 ± 0.03
	width	5.52 ± 0.04	3.64 ± 0.03	3.40 ± 0.02	3.26 ± 0.02	3.24 ± 0.02
squared correction	mean	-0.24 ± 0.02	-0.16 ± 0.02	-0.12 ± 0.02	-0.05 ± 0.02	-0.05 ± 0.02
	width	2.03 ± 0.01	1.93 ± 0.01	1.90 ± 0.01	1.87 ± 0.01	1.87 ± 0.01
bootstrapping	mean	-0.02 ± 0.02	-0.02 ± 0.02	-0.02 ± 0.02	0.00 ± 0.02	-0.01 ± 0.02
	width	1.57 ± 0.01	1.61 ± 0.01	1.63 ± 0.01	1.64 ± 0.01	1.65 ± 0.01
full bootstrapping	mean	-0.04 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01	-0.00 ± 0.01	-0.01 ± 0.01
	width	0.98 ± 0.01	0.98 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	1.00 ± 0.01
asymptotic	mean	-0.02 ± 0.02	-0.02 ± 0.02	-0.02 ± 0.02	0.00 ± 0.02	-0.01 ± 0.02
	width	1.59 ± 0.01	1.63 ± 0.01	1.64 ± 0.01	1.64 ± 0.01	1.64 ± 0.01
asymptotic corrected	mean	0.04 ± 0.01	0.03 ± 0.01	0.01 ± 0.01	0.02 ± 0.01	0.01 ± 0.01
	width	0.97 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	0.99 ± 0.01
cFit	mean	-0.07 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01	-0.01 ± 0.01	-0.01 ± 0.01
	width	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01

(b) Gaussian background model

Table 5: Means and widths of the pull distribution for the different approaches to the uncertainty estimation, depending on the total yield N_{tot} . The different tables shown correspond to the different background models as specified in Sec. 4.

method	pull	1 k	2 k	4 k	10 k	20 k
sFit	mean	-1.36 ± 0.07	-0.83 ± 0.05	-0.56 ± 0.04	-0.28 ± 0.04	-0.18 ± 0.04
	width	6.83 ± 0.05	4.63 ± 0.03	4.25 ± 0.03	4.08 ± 0.03	4.05 ± 0.03
scaled weights	mean	-0.92 ± 0.05	-0.57 ± 0.04	-0.39 ± 0.03	-0.19 ± 0.03	-0.12 ± 0.03
	width	4.97 ± 0.04	3.62 ± 0.03	3.39 ± 0.02	3.29 ± 0.02	3.27 ± 0.02
squared correction	mean	-0.24 ± 0.02	-0.16 ± 0.02	-0.12 ± 0.02	-0.04 ± 0.02	-0.02 ± 0.02
	width	2.01 ± 0.01	1.91 ± 0.01	1.88 ± 0.01	1.88 ± 0.01	1.89 ± 0.01
bootstrapping	mean	-0.02 ± 0.02	-0.02 ± 0.02	-0.02 ± 0.02	0.01 ± 0.02	0.02 ± 0.02
	width	1.56 ± 0.01	1.58 ± 0.01	1.60 ± 0.01	1.64 ± 0.01	1.65 ± 0.01
full bootstrapping	mean	-0.04 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01	0.00 ± 0.01	0.01 ± 0.01
	width	0.97 ± 0.01	0.97 ± 0.01	0.98 ± 0.01	1.00 ± 0.01	1.00 ± 0.01
asymptotic	mean	-0.02 ± 0.02	-0.02 ± 0.02	-0.02 ± 0.02	0.01 ± 0.02	0.02 ± 0.02
	width	1.57 ± 0.01	1.59 ± 0.01	1.61 ± 0.01	1.63 ± 0.01	1.65 ± 0.01
asymptotic corrected	mean	0.04 ± 0.01	0.02 ± 0.01	0.01 ± 0.01	0.03 ± 0.01	0.02 ± 0.01
	width	0.97 ± 0.01	0.97 ± 0.01	0.98 ± 0.01	0.99 ± 0.01	1.00 ± 0.01
cFit	mean	-0.08 ± 0.01	-0.05 ± 0.01	-0.04 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01
	width	1.00 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01

(a) Triangular background model

method	pull	1 k	2 k	4 k	10 k	20 k
sFit	mean	-0.99 ± 0.05	-0.59 ± 0.04	-0.41 ± 0.04	-0.31 ± 0.03	-0.19 ± 0.03
	width	4.67 ± 0.03	3.87 ± 0.03	3.59 ± 0.03	3.48 ± 0.02	3.42 ± 0.02
scaled weights	mean	-0.67 ± 0.04	-0.40 ± 0.03	-0.29 ± 0.03	-0.22 ± 0.03	-0.14 ± 0.03
	width	3.52 ± 0.02	3.05 ± 0.02	2.87 ± 0.02	2.80 ± 0.02	2.77 ± 0.02
squared correction	mean	-0.20 ± 0.02	-0.12 ± 0.02	-0.09 ± 0.02	-0.08 ± 0.02	-0.05 ± 0.02
	width	1.84 ± 0.01	1.79 ± 0.01	1.76 ± 0.01	1.74 ± 0.01	1.73 ± 0.01
bootstrapping	mean	-0.04 ± 0.01	-0.01 ± 0.01	-0.03 ± 0.01	-0.04 ± 0.01	-0.02 ± 0.01
	width	1.38 ± 0.01	1.42 ± 0.01	1.43 ± 0.01	1.44 ± 0.01	1.44 ± 0.01
full bootstrapping	mean	-0.04 ± 0.01	-0.01 ± 0.01	-0.02 ± 0.01	-0.03 ± 0.01	-0.01 ± 0.01
	width	0.96 ± 0.01	0.98 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	1.00 ± 0.01
asymptotic	mean	-0.04 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01	-0.04 ± 0.01	-0.02 ± 0.01
	width	1.40 ± 0.01	1.43 ± 0.01	1.43 ± 0.01	1.43 ± 0.01	1.43 ± 0.01
asymptotic corrected	mean	0.03 ± 0.01	0.03 ± 0.01	0.01 ± 0.01	-0.01 ± 0.01	0.00 ± 0.01
	width	0.97 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	0.99 ± 0.01
cFit	mean	-0.07 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01
	width	1.00 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	0.99 ± 0.01

(b) Flat background model

Table 6: Means and widths of the pull distribution for the different approaches to the uncertainty estimation, depending on the total yield N_{tot} . The different tables shown correspond to the different background models as specified in Sec. 4.

F *sWeights* with negligible nuisance parameter correlations

In Sec. 4.2 the mass range which is used to determine the *sWeights* is chosen such that the background slope α_{bkg} is significantly correlated with the event yields. This results in an additional uncertainty on the *sWeights* that needs to be accounted for using Eq. 52. For completeness, in this section the different methods are studied in the absence of any significant impact of nuisance parameters on the *sWeights*. To this end, the mass range $[5\,267, 5\,467] \text{ MeV}/c^2$ is chosen, which is a symmetrical mass window around the peak at $5\,367 \text{ MeV}/c^2$, as shown in Fig. 12. This choice results in negligible correlation of α_{bkg} with N_{sig} and N_{bkg} . All other settings are kept as in Sec. 4.2.




Figure 12: Discriminating mass distribution for (black) the full data, (blue) signal and (red) background using the mass range $[5\,267, 5\,467] \text{ MeV}/c^2$.

The means and widths of the resulting pull distributions are given in Tabs. 7 and 8. As for the case discussed in Sec. 4.2, the *sFit*, the *scaled weights* method and the *squared weights* method show significant undercoverage. All other methods, namely the (*full bootstrapping*), the *asymptotic (corrected)* method and the *cFit* show correct coverage. In particular, the correction according to Eq. 52 does not negatively affect the coverage properties when no significant correlations with nuisance parameters are present in the fit to determine the *sWeights*.

method	pull	1 k	2 k	4 k	10 k	20 k
sFit	mean	-0.29 ± 0.02	-0.18 ± 0.02	-0.14 ± 0.02	-0.10 ± 0.02	-0.04 ± 0.02
	width	2.47 ± 0.02	2.41 ± 0.02	2.37 ± 0.02	2.36 ± 0.02	2.37 ± 0.02
scaled weights	mean	-0.19 ± 0.02	-0.12 ± 0.02	-0.09 ± 0.02	-0.07 ± 0.02	-0.02 ± 0.02
	width	1.76 ± 0.01	1.72 ± 0.01	1.70 ± 0.01	1.69 ± 0.01	1.70 ± 0.01
squared correction	mean	-0.06 ± 0.01	-0.03 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01	-0.00 ± 0.01
	width	1.14 ± 0.01	1.14 ± 0.01	1.13 ± 0.01	1.13 ± 0.01	1.14 ± 0.01
bootstrapping	mean	-0.01 ± 0.01	0.00 ± 0.01	-0.00 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01
	width	0.98 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.01 ± 0.01
full bootstrapping	mean	0.00 ± 0.01	0.01 ± 0.01	0.00 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01
	width	0.99 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	1.02 ± 0.01
asymptotic	mean	-0.00 ± 0.01	0.00 ± 0.01	-0.00 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01
	width	0.99 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	1.01 ± 0.01
asymptotic corrected	mean	-0.00 ± 0.01	0.00 ± 0.01	-0.00 ± 0.01	-0.01 ± 0.01	0.01 ± 0.01
	width	0.99 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	1.01 ± 0.01
cFit	mean	-0.05 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01	0.00 ± 0.01
	width	1.02 ± 0.01	1.01 ± 0.01	1.01 ± 0.01	1.01 ± 0.01	1.01 ± 0.01

(a) Exponential background model

method	pull	1 k	2 k	4 k	10 k	20 k
sFit	mean	-0.42 ± 0.03	-0.25 ± 0.03	-0.16 ± 0.03	-0.11 ± 0.03	-0.08 ± 0.03
	width	3.29 ± 0.02	3.08 ± 0.02	3.01 ± 0.02	2.94 ± 0.02	2.93 ± 0.02
scaled weights	mean	-0.27 ± 0.02	-0.15 ± 0.02	-0.10 ± 0.02	-0.07 ± 0.02	-0.05 ± 0.02
	width	2.32 ± 0.02	2.19 ± 0.02	2.15 ± 0.02	2.10 ± 0.01	2.10 ± 0.01
squared correction	mean	-0.06 ± 0.01	-0.03 ± 0.01	-0.01 ± 0.01	-0.01 ± 0.01	-0.01 ± 0.01
	width	1.15 ± 0.01	1.13 ± 0.01	1.13 ± 0.01	1.11 ± 0.01	1.11 ± 0.01
bootstrapping	mean	0.02 ± 0.01	0.02 ± 0.01	0.02 ± 0.01	0.01 ± 0.01	0.00 ± 0.01
	width	0.96 ± 0.01	0.97 ± 0.01	0.98 ± 0.01	0.97 ± 0.01	0.97 ± 0.01
full bootstrapping	mean	0.04 ± 0.01	0.04 ± 0.01	0.03 ± 0.01	0.01 ± 0.01	0.01 ± 0.01
	width	0.99 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01
asymptotic	mean	0.03 ± 0.01	0.03 ± 0.01	0.03 ± 0.01	0.01 ± 0.01	0.01 ± 0.01
	width	0.97 ± 0.01	0.98 ± 0.01	0.98 ± 0.01	0.97 ± 0.01	0.97 ± 0.01
asymptotic corrected	mean	0.03 ± 0.01	0.03 ± 0.01	0.03 ± 0.01	0.01 ± 0.01	0.01 ± 0.01
	width	0.97 ± 0.01	0.98 ± 0.01	0.98 ± 0.01	0.97 ± 0.01	0.97 ± 0.01
cFit	mean	-0.06 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01
	width	1.01 ± 0.01	1.02 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	1.00 ± 0.01

(b) Gaussian background model

Table 7: Means and widths of the pull distribution for the different approaches to the uncertainty estimation, depending on the total yield N_{tot} . The different tables shown correspond to the different background models as specified in Sec. 4. In this case, the mass range $[5\,267, 5\,467]$ MeV/ c^2 is chosen, such that the nuisance parameter α_{bkg} has negligible correlation with the event yields.

method	pull	1 k	2 k	4 k	10 k	20 k
sFit	mean	-0.42 ± 0.03	-0.29 ± 0.03	-0.20 ± 0.03	-0.13 ± 0.03	-0.10 ± 0.03
	width	3.25 ± 0.02	3.08 ± 0.02	3.01 ± 0.02	3.02 ± 0.02	2.96 ± 0.02
scaled weights	mean	-0.27 ± 0.02	-0.18 ± 0.02	-0.13 ± 0.02	-0.08 ± 0.02	-0.06 ± 0.02
	width	2.30 ± 0.02	2.19 ± 0.02	2.15 ± 0.02	2.16 ± 0.02	2.12 ± 0.01
squared correction	mean	-0.06 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01
	width	1.15 ± 0.01	1.13 ± 0.01	1.13 ± 0.01	1.14 ± 0.01	1.13 ± 0.01
bootstrapping	mean	0.01 ± 0.01	0.01 ± 0.01	0.00 ± 0.01	0.00 ± 0.01	-0.00 ± 0.01
	width	0.96 ± 0.01	0.97 ± 0.01	0.97 ± 0.01	0.99 ± 0.01	0.98 ± 0.01
full bootstrapping	mean	0.03 ± 0.01	0.02 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.00 ± 0.01
	width	0.98 ± 0.01	0.99 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	1.00 ± 0.01
asymptotic	mean	0.03 ± 0.01	0.02 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.00 ± 0.01
	width	0.96 ± 0.01	0.97 ± 0.01	0.97 ± 0.01	0.99 ± 0.01	0.98 ± 0.01
asymptotic corrected	mean	0.03 ± 0.01	0.02 ± 0.01	0.01 ± 0.01	0.01 ± 0.01	0.00 ± 0.01
	width	0.96 ± 0.01	0.97 ± 0.01	0.97 ± 0.01	0.99 ± 0.01	0.98 ± 0.01
cFit	mean	-0.07 ± 0.01	-0.05 ± 0.01	-0.04 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01
	width	1.02 ± 0.01	1.01 ± 0.01	1.00 ± 0.01	1.01 ± 0.01	1.00 ± 0.01

(a) Triangular background model

method	pull	1 k	2 k	4 k	10 k	20 k
sFit	mean	-0.46 ± 0.03	-0.36 ± 0.03	-0.28 ± 0.03	-0.17 ± 0.03	-0.09 ± 0.03
	width	3.15 ± 0.02	3.04 ± 0.02	2.97 ± 0.02	2.92 ± 0.02	2.90 ± 0.02
scaled weights	mean	-0.30 ± 0.02	-0.24 ± 0.02	-0.19 ± 0.02	-0.12 ± 0.02	-0.06 ± 0.02
	width	2.23 ± 0.02	2.17 ± 0.02	2.13 ± 0.02	2.09 ± 0.01	2.07 ± 0.01
squared correction	mean	-0.09 ± 0.01	-0.08 ± 0.01	-0.07 ± 0.01	-0.04 ± 0.01	-0.02 ± 0.01
	width	1.22 ± 0.01	1.23 ± 0.01	1.22 ± 0.01	1.21 ± 0.01	1.20 ± 0.01
bootstrapping	mean	-0.01 ± 0.01	-0.02 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01	-0.00 ± 0.01
	width	0.96 ± 0.01	0.98 ± 0.01	0.98 ± 0.01	0.98 ± 0.01	0.98 ± 0.01
full bootstrapping	mean	0.00 ± 0.01	-0.01 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01	0.00 ± 0.01
	width	0.98 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01
asymptotic	mean	-0.01 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01	-0.00 ± 0.01
	width	0.97 ± 0.01	0.98 ± 0.01	0.98 ± 0.01	0.98 ± 0.01	0.98 ± 0.01
asymptotic corrected	mean	-0.01 ± 0.01	-0.02 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01	-0.00 ± 0.01
	width	0.97 ± 0.01	0.98 ± 0.01	0.98 ± 0.01	0.98 ± 0.01	0.98 ± 0.01
cFit	mean	-0.04 ± 0.01	-0.03 ± 0.01	-0.03 ± 0.01	-0.02 ± 0.01	-0.01 ± 0.01
	width	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01	1.00 ± 0.01

(b) Flat background model

Table 8: Means and widths of the pull distribution for the different approaches to the uncertainty estimation, depending on the total yield N_{tot} . The different tables shown correspond to the different background models as specified in Sec. 4. In this case, the mass range $[5\,267, 5\,467]$ MeV/ c^2 is chosen, such that the nuisance parameter α_{bkg} has negligible correlation with the event yields.