

Discrete logarithm and Diffie-Hellman problems in identity black-box groups

Gábor Ivanyos^{*} Antoine Joux[†] Miklos Santha[‡]

November 6, 2019

Abstract

We investigate the computational complexity of the discrete logarithm, the computational Diffie-Hellman and the decisional Diffie-Hellman problems in some identity black-box groups $G_{p,t}$, where p is a prime number and t is a positive integer. These are defined as quotient groups of vector space \mathbb{Z}_p^{t+1} by a hyperplane H given through an identity oracle. While in general black-box groups with unique encoding these computational problems are classically all hard and quantumly all easy, we find that in the groups $G_{p,t}$ the situation is more contrasted. We prove that while there is a polynomial time probabilistic algorithm to solve the decisional Diffie-Hellman problem in $G_{p,1}$, the probabilistic query complexity of all the other problems is $\Omega(p)$, and their quantum query complexity is $\Omega(\sqrt{p})$. Our results therefore provide a new example of a group where the computational and the decisional Diffie-Hellman problems have widely different complexity.

1 Introduction

Black-box groups we introduced by Babai and Szemerédi [5] for studying the structure of finite matrix groups. In a black-box group, the group elements are encoded by binary strings of certain length, the multiplication and inverse group operations are given by oracles. Similarly, identity testing, that is checking whether an element is equal to the identity element, is also done with a special identity oracle. These oracles are also called black-boxes, giving their names to the groups. Identity testing is required when several strings may encode the same group element. In this case we speak about non-unique encoding, in opposition to unique encoding when every group element is encoded by a unique string. Black-box groups with non-unique encoding are motivated by their ability to capture factor groups of subgroups of matrix groups by certain normal subgroups which admit efficient algorithms for membership testing. An important example for such a subgroup is the solvable radical, that is the largest solvable normal subgroup.

A black-box group problem might only have the oracle inputs if the problem concerns some global property of the group. In some other cases it might also have standard inputs (or shortly

^{*}Institute for Computer Science and Control, Budapest, Hungary (Gabor.Ivanyos@sztaki.hu).

[†]CISPA Helmholtz Center for Information Security, Saarbrücken, Germany and Sorbonne Université, Institut de Mathématiques de Jussieu–Paris Rive Gauche, CNRS, INRIA, Univ Paris Diderot, Campus Pierre et Marie Curie, F-75005 Paris, France (antoine.joux@m4x.org).

[‡]CNRS, IRIF, Université de Paris, F-75013 Paris, France; and Centre for Quantum Technologies and MajuLab, National University of Singapore, Singapore 117543 (miklos.santha@gmail.com).

inputs), a finite set of group elements represented by their encodings. A black-box group algorithm is allowed to call the oracles for the group operations and for the identity test and it might also perform arbitrary bit operations. The query complexity of an algorithm is the number of oracle calls, while the running time or computational complexity is the number of oracle calls together with the number of other bit operations, when we are maximizing over both oracle and standard inputs. In the quantum setting, the oracles are given by unitary operators.

Many classical algorithms have been developed for computations with black-box groups [6, 4, 14], for example the identification of the composition factors, even the non-commutative ones. When the oracle operations can be simulated by efficient procedures, efficient black-box algorithms automatically produce efficient algorithms. Permutation groups, finite matrix groups over finite fields and over algebraic number fields fit in this context. There has been also considerable effort to design quantum algorithms in black-box groups. In the case of unique encoding efficient algorithms have been conceived for the decomposition of Abelian groups into a direct sum of cyclic groups of prime power order [9], order computing and membership testing in solvable groups [22] and solving the hidden subgroup problem in Abelian groups [17].

The discrete logarithm problem DLOG, and various Diffie-Hellman type problems are fundamental tasks in computational number theory. They are equally important in cryptography, since the security of many cryptographic systems is based on their computational difficulty. Let G be a cyclic group (denoted additively). Given two group elements g and h , where g is a generator, DLOG asks to compute an integer d such that $h = dg$. Given three group elements g, ag and bg , where g is a generator, the computational Diffie-Hellman problem CDH is to compute $(ab)g$. Given three group elements g, ag, bg and cg , where g is a generator, the decisional Diffie-Hellman problem DDH is to decide whether $c = ab$ modulo the order of g .

The problems are enumerated in decreasing order of difficulty: DDH can not be harder than CDH and CDH is not harder than DLOG. While there are groups where even DLOG is easy (for example \mathbb{Z}_m , the additive group of integers modulo m), in general all three problems are thought to be computationally hard. We are not aware of any group where CDH is easy while DLOG is hard. In fact, Maurer and Wolf have proven in [16] that under a seemingly reasonable number-theoretic assumption, the two problems are equivalent in the case of unique-encoding groups. Based on this, Joux and Nguyen [13] have constructed a cryptographic group where DDH is easy to compute while CDH is as hard as DLOG. In generic black-box groups we have provable query lower bounds for these problems, even in the case of unique encoding. Shoup has proven [21] that in \mathbb{Z}_p , given as a black-box group with unique encoding, to solve DLOG and CDH require $\Omega(p^{1/2})$ group operations. Subsequently, Damgård, Hazay and Zottare [10] have established a lower bound of the same order for DDH. We remark that the Pohlig-Hellman [18] algorithm reduces DLOG in arbitrary cyclic groups to DLOG in its prime order subgroups. Furthermore, in prime order groups (with unique encodings), Shanks's baby-step giant-step algorithm solve the problem using $O(p^{1/2})$ group operation, thus matching the lower bound for black-box groups.

Though, as we said, all three problems are considered computationally intractable on a classical computer, there is a polynomial time quantum algorithm for DLOG due to Shor [20]. Since DLOG is an instance of the Abelian hidden subgroup problem, Mosca's result [17] implies that by a quantum computer it can also be solved efficiently in black-box groups with unique encoding.

We are concerned here with identity black-box group, a special class of black-box groups where only the identity test is given by an oracle. These groups are quotient groups of some explicitly given ambient group. An identity black box group G is specified by an ambient group G' and an

identity oracle Id which tests membership in some (unknown) normal subgroup H of G' . In G the group operations are explicitly defined by the group operations in G' , and therefore it is the quotient group G'/H .

Let p be a prime number. More specifically we will study the problems DLOG, CDH and DDH in identity black-box groups whose ambient group is \mathbb{Z}_p^{t+1} , for some positive integer t , and where the normal subgroup H , specified by the identity oracle, is isomorphic to \mathbb{Z}_p^t . We denote such an identity black box group by $G_{p,t}$. We fully characterize the complexity of the three problems in these groups. Our results are mainly query lower bounds: the probabilistic query complexity of all these problems, except DDH in level 1 groups, is $\Omega(p)$, and their quantum query complexity is $\Omega(\sqrt{p})$. These lower bounds are obviously tight since DLOG($G_{p,t}$) can be solved, for all $t \geq 1$, by exhaustive search and by Grover's algorithm in respective query complexity p and $O(\sqrt{p})$. We have also one, maybe surprising, algorithmic result: the computational complexity of DDH($G_{p,1}$) is polynomial. Our results can be summarized in the following theorems.

Theorem (Lower bounds)

1. For all $t \geq 1$, the randomized query complexity of DLOG($G_{p,t}$) and CDH($G_{p,1}$) is $\Omega(p)$.
2. For all $t \geq 1$, the quantum query complexity of DLOG($G_{p,t}$) and CDH($G_{p,1}$) is $\Omega(\sqrt{p})$.
3. For all $t \geq 2$, the randomized query complexity of DDH($G_{p,t}$) is $\Omega(p)$.
4. For all $t \geq 2$, the quantum query complexity of DDH($G_{p,t}$) is $\Omega(\sqrt{p})$.

Theorem (Upper bound) DDH($G_{p,1}$) can be solved in probabilistic polynomial time in $\log p$.

2 Preliminaries

Formally, a *black-box group* G is a 4-tuple $G = (C, \text{Mult}, \text{Inv}, \text{Id})$ where C is the set of admissible codewords, $\text{Mult} : C \times C \mapsto C$ is a binary operation, $\text{Inv} : C \rightarrow C$ is a unary operation and $\text{Id} : C \rightarrow \{0, 1\}$ is a unary Boolean function. The operations Mult, Inv and the function Id are given by *oracles*. We require that there exists a finite group \tilde{G} and a surjective map $\phi : C \rightarrow \tilde{G}$ for which, for every $x, y \in C$, we have $\phi(\text{Mult}(x, y)) = \phi(x)\phi(y)$, $\phi(\text{Inv}(x)) = \phi(x)^{-1}$, and $\phi(x) = 1_{\tilde{G}}$ if and only if $\text{Id}(x) = 1$. We say that x is the identity element in G or that $x = 1$ if $\text{Id}(x) = 1$. With the identity oracle we can test equality since $x = y$ in G exactly when $\text{Id}(\text{Mult}(x, \text{Inv}(y))) = 1$. We say that a black-box group has *unique encoding* if ϕ is a bijection. For abelian groups we also use the additive notation in which case the binary operation of G is denoted by Add and its identity element is denoted by 0 .

We are concerned here with a special class of black-box groups which are quotient groups of some explicitly given group. An *identity black-box group* is a couple $G = (G', \text{Id})$ where G' is group and the *identity oracle* $\text{Id} : G' \rightarrow \{0, 1\}$ is the characteristic function of some (unknown) normal subgroup H of G' . We call G' the *ambient group* of G . In G the group operations Mult and Inv are defined by the group operations in the ambient group G' modulo H . As a consequence, G is the quotient group G'/H .

Let p be a prime number and let $t \geq 1$ be a positive integer. We denote by \mathbb{Z}_p the additive group of integers modulo p , by \mathbb{F}_p the finite field of size p , and by \mathbb{Z}_p^t the t -dimensional vector space over \mathbb{F}_p . For $h, k \in \mathbb{Z}_p^t$, we denote their scalar product modulo p by $h \cdot k$.

We will work with identity black-box groups whose ambient group is \mathbb{Z}_p^{t+1} , and the subgroup H is isomorphic to \mathbb{Z}_p^t , that is a hyperplane of \mathbb{Z}_p^{t+1} . We will specify the identity oracle by a non-zero normal vector $n \in \mathbb{Z}_p^{t+1}$ of H . By permuting coordinates and multiplying by some non-zero constant, we can suppose without loss of generality that it is of the form $n = (1, n_1, \dots, n_t)$. We call such a vector *t-suitable*. We define the function $\text{Id}_n : \mathbb{Z}_p^{t+1} \rightarrow \{0, 1\}$ by $\text{Id}_n(h) = 1$ if $h \cdot n = 0$. Clearly Id_n is the characteristic function of the hyperplane $H_n = \{h \in \mathbb{Z}_p^{t+1} : h \cdot n = 0\}$. We define the identity black-box group $G_{p,t} = (\mathbb{Z}_p^{t+1}, \text{Id})$, where the identity oracle Id satisfies $\text{Id} = \text{Id}_n$, for some (unknown) *t-suitable* vector n . We call t the *level* of the group $G_{p,t}$. We emphasize again that the group operations of $G_{p,t}$ are performed as group operations in \mathbb{Z}_p^{t+1} . Therefore, for $h, k \in \mathbb{Z}_p^{t+1}$, the equality $h = k$ in $G_{p,t}$ means equality in \mathbb{Z}_p^{t+1} modulo H_n , where H_n is identified by Id_n . To be short, we will refer to $G_{p,t}$ as the *hidden cyclic group* of level t .

Proposition 2.1. *The groups $G_{p,t}$, and \mathbb{Z}_p are isomorphic and the map $\phi : G_{p,t} \rightarrow \mathbb{Z}_p$ defined by $\phi(h) = h \cdot n \in \mathbb{Z}_p$ is a group isomorphism.*

Proof. The maps from \mathbb{Z}_p^{t+1} to $G_{p,t}$ (respectively to \mathbb{Z}_p) mapping $h \in \mathbb{Z}_p^{t+1}$ to its class in the quotient $G_{p,t}$ (respectively to $h \cdot n$) are group homomorphisms with the same kernel H_n . \square

We recall now the basic notions of query complexity for the specific case of boolean oracle functions. Let m be a positive integer. An *oracle function* is a function $A : S \rightarrow \{0, 1\}^M$, where $S \subseteq \{0, 1\}^m$ and $M \geq 1$ is a positive integer. If $M = 1$, then we call the oracle function *boolean*. The input $f \in S$ is given by an oracle, that is $f(x)$ can be accessed by the query x . The output on f is $A(f)$. Each query adds one to the complexity of an algorithm, but all other computations are free. The state of the computation is represented by three registers, the query register $1 \leq x \leq m$, the answer register $a \in \{0, 1\}$, and the work register z . The computation takes place in the vector space spanned by all basis states $|x\rangle|a\rangle|z\rangle$. In the *quantum query* model introduced by Beals et al. [7] the state of the computation is a complex combination of all basis states which has unit length in the norm l_2 . In the *randomized query* model it is a non-negative real combination of unit length in the norm l_1 , and in the deterministic model it is always one of the basis states.

The query operation O_f maps the basis state $|x\rangle|a\rangle|z\rangle$ into the state $|x\rangle|(a + f(x)) \bmod 2\rangle|z\rangle$. Non-query operations do not depend on f . A *k-query algorithm* is a sequence of $(k + 1)$ operations (U_0, U_1, \dots, U_k) where U_i is unitary in the quantum, and stochastic in the randomized model. Initially the state of the computation is set to some fixed value $|0\rangle|0\rangle|0\rangle$, and then the sequence of operations $U_0, O_f, U_1, O_f, \dots, U_{k-1}, O_f, U_k$ is applied. A quantum or randomized algorithm *computes* A on input f if the observation of the last M bits of the work register yields $A(f)$ with probability at least $2/3$. Then $Q(A)$ (respectively $R(A)$) is the smallest k for which there exists a k -query quantum (respectively randomized) algorithm which computes A on every input f . We have $R(A) \leq Q(A) \leq m$.

We define now our problems we are concerned with, the *discrete logarithm* problem DLOG, the *computational Diffie-Hellman* problem CDH and the *decisional Diffie-Hellman* problem DDH in hidden cyclic groups $G_{p,t}$. We say that a quadruple $(g, h, k, \ell) \in G_{p,t}^4$ where g is a generator of $G_{p,t}$ is a *DH-quadruple* if $h = ag, k = bg$ and $\ell = cg$ for some integers a, b, c such that $c = ab$ modulo p .

DLOG($G_{p,t}$)

Oracle input: Id_n for some t -suitable vector n .

Input: A couple $(g, h) \in G_{p,t}^2$ such that g is a generator of $G_{p,t}$.

Output: A non-negative integer d such that $dg = h$.

CDH($G_{p,t}$)

Oracle input: Id_n for some t -suitable vector n .

Input: A triple $(g, h, k) \in G_{p,t}^3$ such that g is a generator of $G_{p,t}$.

Output: $\ell \in G_{p,t}$ such that (g, h, k, ℓ) is a DH-quadruple.

DDH($G_{p,t}$)

Oracle input: Id_n for some t -suitable vector n .

Input: A quadruple $(g, h, k, \ell) \in G_{p,t}^4$ such that g is a generator of $G_{p,t}$.

Question: Is (g, h, k, ℓ) a DH-quadruple?

An algorithm for these problems has access to the input and oracle access to the oracle input, and every query is counted as one computational step. We say that it solves the problem *efficiently* if it works in time polynomial in $\log p$ and t . For any fixed input, the problems become oracle functions, where we consider only those identity oracles for which the input is legitimate. By their *query complexity* we mean, both in the quantum and in the randomized model, the maximum, over all inputs, of the respective query complexity of these oracle function.

The problems are enumerated in decreasing order of difficulty. The existence of an efficient algorithm for DLOG($G_{p,t}$) implies the existence of an efficient algorithm for CDH($G_{p,t}$), which in turn gives rise to an efficient algorithm for DDH($G_{p,t}$). For query complexity we have $Q(\text{DDH}(G_{p,t})) \leq Q(\text{CDH}(G_{p,t})) \leq Q(\text{DLOG}(G_{p,t}))$, and the same inequalities hold for the randomized model. The problems are getting harder as the level of the hidden cyclic group increases, as the almost trivial reductions in the next Proposition show. To ease notation, for $h = (h_0, \dots, h_t)$ in \mathbb{Z}_p^{t+1} , we denote by h' the element $(h_0, \dots, h_t, 0) \in \mathbb{Z}_p^{t+2}$.

Proposition 2.2. *For every $t \geq 1$, DLOG($G_{p,t}$) and DDH($G_{p,t}$) are polynomial time many-one reducible to respectively DLOG($G_{p,t+1}$) and DDH($G_{p,t+1}$); and CDH($G_{p,t}$) is computable in polynomial time with a single query to CDH($G_{p,t+1}$).*

Proof. First observe that the identity oracle $\text{Id}_{n'}$ in $G_{p,t+1}$ can be simulated by the identity oracle Id_n of $G_{p,t}$. Indeed, for an arbitrary element h^* in $G_{p,t+1}$, where $h^* = (h_0, h_1, \dots, h_t, h_{t+1})$, set $h = (h_0, h_1, \dots, h_t)$. Then $h^* \cdot n' = h \cdot n$. Let g be a generator of $G_{p,t}$ with identity oracle Id_n , that is $g \cdot n \neq 0$. Then $g' = (g, 0)$ is a generator of $G_{p,t+1}$ with identity oracle $\text{Id}_{n'}$, since $g' \cdot n' = g \cdot n$, and therefore $g' \cdot n' \neq 0$.

For arbitrary $h, k, \ell \in G_{p,t}$, and for every integer d , we have $dg = h$ if and only if $dg' = h'$. Similarly, (g, h, k, ℓ) is a DH-quadruple if and only if (g', h', k', ℓ') is a DH-quadruple. This gives the many-one reductions for DLOG and DDH. In the CDH reduction, on an instance (g, h, k) , we call CDH($G_{p,t+1}$) on instance (g', h', k') . Suppose that it gives the answer $\ell^* = (\ell_0, \ell_1, \dots, \ell_t, \ell_{t+1})$. We set $\ell = (\ell_0, \ell_1, \dots, \ell_t)$, and observe that (g, h, k, ℓ) is a DH-quadruple because (g', h', k', ℓ^*) is a DH-quadruple and $\ell^* \cdot n' = \ell \cdot n$. \square

3 The complexity in groups of level 1

In most parts of this section we restrict ourselves to the case $t = 1$. To simplify notation, we set $n = (1, s)$ and we denote the identity oracle Id_n by Id_s and the line H_n by H_s . Also, we refer to s as the *secret*. As it turns out, solving $\text{DLOG}(G_{p,1})$ or $\text{CDH}(G_{p,1})$ is essentially as hard as finding the secret, therefore we define formally this problem as

$\text{SECRET}(G_{p,1})$
Oracle input: Id_s for some $s \in \mathbb{Z}_p$.
Output: s .

What is the complexity of finding s , that is how many calls to the identity oracle are needed for that task? To answer this question, we consider US, the well studied unstructured search problem. For this, let C be an arbitrary set, and let $s \in C$ be an arbitrary distinguished element. Then the *Grover oracle* $\Delta_s : C \rightarrow \{0, 1\}$ is the boolean function such that $\Delta_s(x) = 1$ if and only if $x = s$. The *unstructured search* problem over C is defined as

$\text{US}(C)$
Oracle input: Δ_s for some $s \in C$.
Output: s .

Suppose that the size of C is N . It is easily seen that probabilistic query complexity of $\text{US}(C)$ is linear in N . The quantum query complexity of the problem is also well studied. Grover [11] has determined that it can be solved with $O(\sqrt{N})$ queries, while Bennett et al. [8] have shown that $\Omega(\sqrt{N})$ queries are also necessary.

Fact 3.1. *For $|C| = N$, the randomized query complexity of $\text{US}(C)$ is $\Theta(N)$ and its quantum query complexity is $\Theta(\sqrt{N})$.*

The relationship between US and the problem SECRET is given by the fact that the identity oracle Id_s and the Grover oracle Δ_s can simulate each other with a single query.

Proposition 3.2. *The identity oracle Id_s of $G_{p,s}$ and the Grover oracle Δ_s , defined over \mathbb{Z}_p , can simulate each other with at most one query.*

Proof. The simulation of the Grover oracle by the identity oracle is simple: for $x \in \mathbb{Z}_p$ just query Id_s on $(x, -1)$.

For the reverse direction, let $h = (h_0, h_1)$ be an input to the identity oracle. Then h is the identity element, that is $\text{Id}_s(h) = 1$ if and only if $-h_0 = sh_1$. When h_1 is invertible in \mathbb{Z}_p we can check by the Grover oracle if $-h_0h_1^{-1} = s$. For $h_1 = 0$ the only possible value to put h into H_s is 0. Therefore we have

$$\text{Id}(h) = \begin{cases} 1 & \text{if } h = (0, 0) \\ 0 & \text{if } h_1 = 0 \text{ and } h_0 \neq 0 \\ \Delta_s(-h_0h_1^{-1}) & \text{otherwise.} \end{cases}$$

□

Corollary 3.3. *The randomized query complexity of $\text{SECRET}(G_{p,1})$ is $\Theta(p)$ and its quantum query complexity is $\Theta(\sqrt{p})$.*

We will now consider the reductions of $\text{SECRET}(G_{p,1})$ to $\text{DLOG}(G_{p,1})$ and $\text{CDH}(G_{p,1})$. The case of $\text{DLOG}(G_{p,1})$ in fact follows from the case of $\text{CDH}(G_{p,1})$, but it is so simple that it is worth to describe it explicitly.

Lemma 3.4. *The secret s in $G_{p,1}$ can be found with a single oracle call to $\text{DLOG}(G_{p,1})$*

Proof. First observe that $(1, 0)$ is a generator of $G_{p,1}$, for every s . The algorithm calls $\text{DLOG}(G_{p,1})$ on input $(g, h) = ((1, 0), (0, 1))$. Since $\phi(g) = 1$ and $\phi(h) = s$, the oracle's answer is the secret s itself. \square

We remark that with overwhelming probability we could have given also a random couple $(g, h) \in G_{p,1}^2$ to the oracle, where g is a generator. Indeed, let's suppose that d is the discrete logarithm. Then $h - dg \in H_s$, and therefore $s = -(h_0 - dg_0)(h_1 - dg_1)^{-1}$, where the operations are done in \mathbb{Z}_p , under the condition that $h_1 - dg_1 \neq 0$, which happens with probability $(p-1)/p$.

The reduction of $\text{SECRET}(G_{p,1})$ to $\text{CDH}(G_{p,1})$ requires more work. The main idea of the reduction is to extend \mathbb{Z}_p to a field and use the multiplication for the characterization of DH-quadruples. Indeed, since \mathbb{Z}_p is the additive group of the field \mathbb{F}_p , we can use the isomorphism ϕ of Proposition 2.1 between $G_{p,t}$ and \mathbb{Z}_p to define appropriate multiplication and multiplicative inverse operations. This extends $G_{p,t}$ to a field isomorphic to \mathbb{F}_p which we denote by $F_{p,t}$. This process is completely standard but we describe it for completeness. The definitions of these two operations are:

$$\begin{aligned} hk &= \phi^{-1}(\phi(h)\phi(k)), \\ h^{-1} &= \phi^{-1}(\phi(h)^{-1}). \end{aligned}$$

With these operations the map ϕ becomes a field isomorphism between $F_{p,t}$ and \mathbb{F}_p .

Proposition 3.5. *The map ϕ of Proposition 2.1 is an isomorphism between $F_{p,t}$ and \mathbb{F}_p .*

Proof. By definition $\phi(hk) = \phi(h)\phi(k)$ and $\phi(h^{-1}) = \phi(h)^{-1}$. \square

The field structure of $F_{p,t}$ yields a very useful characterization of DH-quadruples.

Proposition 3.6. *Let g be a generator of $G_{p,t}$. In $F_{p,t}$ the quadruple (g, h, k, ℓ) is a DH-quadruple if and only if*

$$g\ell - hk = 0.$$

Proof. Let $h = ag, k = bg$ and $\ell = cg$ for some integers a, b, c . Using the field structure of $F_{p,t}$, it is true that $g\ell - hk = 0$ if and only if $(c - ab)g^2 = 0$. Since $F_{p,t}$ is isomorphic to \mathbb{F}_p , an element g is a generator of the additive group \mathbb{Z}_p exactly when $g \neq 0$, and therefore when g^2 is a generator. Therefore $(c - ab)g^2 = 0$ if and only if $c = ab$. \square

We define the application $\chi : \mathbb{Z}_p^{t+1} \rightarrow \mathbb{F}_p[x_1, \dots, x_t]$, from \mathbb{Z}_p^{t+1} to the ring of t -variate polynomials over \mathbb{F}_p , where the image $\chi(h)$ of $h = (h_0, h_1, \dots, h_t) \in \mathbb{Z}_p^{t+1}$ is the polynomial $p_h(x_1, \dots, x_t) = h_0 + \sum_{i=1}^t h_i x_i$. Observe that $p_h(n_1, \dots, n_t) = h \cdot n$, therefore the isomorphism ϕ between $G_{p,t}$ with identity oracle Id_n and \mathbb{Z}_p can be also expressed as $\phi(h) = p_h(n_1, \dots, n_t)$.

Proposition 3.7. *Let g be a generator of $G_{p,1}$ and let h, k, ℓ be arbitrary elements. Then (g, h, k, ℓ) is a DH-quadruple if and only if s is a root of the polynomial $p_g(x)p_\ell(x) - p_h(x)p_k(x)$.*

Proof. By Proposition 3.6 we know that (g, h, k, ℓ) is a DH-quadruple if and only if $g\ell - hk = 0$, that is when $p_{g\ell-hk}(s) = 0$. Now Proposition 3.5 implies that this happens exactly when $p_g(s)p_\ell(s) - p_h(s)p_k(s) = 0$. □

Lemma 3.8. *There is a probabilistic polynomial time algorithm which, given oracle access to $\text{CDH}(G_{p,1})$, solves $\text{SECRET}(G_{p,1})$. The algorithm asks a single query to $\text{CDH}(G_{p,1})$. If we are also given a quadratic non-residue in \mathbb{Z}_p , the algorithm can be made deterministic.*

Proof. The algorithm sets $g = (1, 0), h = (0, 1), k = (1, 1)$ and presents it to the oracle. Let the oracle's answer be $\ell = (\ell_0, \ell_1)$. Since (g, h, k, ℓ) is a DH-quadruple, by Proposition 3.7 we have that s is the root of the second degree equation

$$x^2 + (1 - \ell_1)x + \ell_0 = 0.$$

Assuming that a quadratic non-residue in \mathbb{Z}_p is available then the (not necessarily distinct) roots x_1, x_2 can be computed in deterministic polynomial time using the Shanks-Tonelli algorithm [19]. Without this assumption, a quadratic non-residue can always be computed in probabilistic polynomial time. Finally, we make at most two calls to Id_s on $(x_1, -1)$ and on $(x_2, -1)$. The positive answer tells us which one of the roots is the secret s . □

Similarly to the DLOG case, we could have presented with overwhelming probability also a random triple $(g, h, k) \in G_{p,s}^3$ to $\text{CDH}(G_{p,1})$, where g is a generator. Indeed, if the oracle answer is $\ell = (\ell_0, \ell_1)$ then s is a root of the at most second degree equation

$$(g_0 + g_1x)(\ell_0 + \ell_1x) = (h_0 + h_1x)(k_0 + k_1x).$$

If the equation is of degree 2 then we can proceed as in the proof of Lemma 3.8. This happens exactly when $h_1k_1 \neq g_1\ell_1$. But for every possible fixed value a for $g_1\ell_1$, the probability, over random h_1 and k_1 , that $h_1k_1 = a$ is at most $2/p$, the worst case being $a = 0$. Therefore a random triple (g, h, k) would be suitable for the proof with probability at least $(p - 2)/p$.

Theorem 3.9. *The following lower bounds hold for the query complexity of DLOG and CDH:*

- (1) *The classical query complexity of both $\text{DLOG}(G_{p,s})$ and $\text{CDH}(G_{p,s})$ is $\Omega(p)$.*
- (2) *The quantum query complexity of both $\text{DLOG}(G_{p,s})$ and $\text{CDH}(G_{p,s})$ is $\Omega(\sqrt{p})$.*

Proof. Let us suppose that with m queries to the identity oracle Id_s , one can solve $\text{DLOG}(G_{p,1})$ or $\text{CDH}(G_{p,1})$. Respectively Lemma 3.4 and Lemma 3.8 imply that $\text{SECRET}(G_{p,1})$ can be solved with m queries. The result then follows from the lower bounds of Corollary 3.3. □

Theorem 3.10. *The $\text{DDH}(G_{p,1})$ problem can be solved in probabilistic polynomial time. If we are given a quadratic non-residue in \mathbb{Z}_p the algorithm can be made deterministic.*

Proof. Let (g, h, k, ℓ) be an input to $\text{DDH}(G_{p,1})$ where g is a generator of $G_{p,1}$. By Proposition 3.7 it is a DH-quadruple if and only if s is a root of the polynomial $p_g(x)p_\ell(x) - p_h(x)p_k(x)$, and that is what the algorithm checks. When the polynomial is constant, then the answer is yes if the constant is zero, and otherwise it is no. When the polynomial is non constant, the algorithm essentially proceeds as the one in Lemma 3.8. It solves the at most second degree equation and then checks with the identity oracle if one root is equal to s . □

4 The complexity of DDH in groups of level 2

There are several powerful means to prove quantum query lower bounds, most notably the adversary and the polynomial method [7]. The quantum adversary method initiated by Ambainis [2] has been extended in several ways. The most powerful of those, the method using negative weights [12], turned out to be an exact characterization of the quantum query complexity [15]. We use here a special case of the positive weighted adversary method that also gives probabilistic lower bounds.

Fact 4.1. [1, 3, 23] *Let $A : S \rightarrow \{0, 1\}$ be a boolean oracle function, where $S \subseteq \{0, 1\}^m$. For any $S \times S$ matrix M , set*

$$\sigma(M, f) = \sum_{g \in S} M[f, g].$$

Let Γ be an arbitrary $S \times S$ nonnegative symmetric matrix that satisfies $\Gamma[f, g] = 0$ whenever $A(f) \neq A(g)$. For $1 \leq x \leq m$, let Γ_x be the matrix

$$\Gamma_x[f, g] = \begin{cases} 0 & \text{if } f(x) \neq g(x), \\ \Gamma[f, g] & \text{otherwise.} \end{cases}$$

Then

$$\begin{aligned} \mathbf{Q}(A) &= \Omega \left(\min_{\Gamma[f, g] \neq 0, f(x) \neq g(x)} \sqrt{\frac{\sigma(\Gamma, f)\sigma(\Gamma, g)}{\sigma(\Gamma_x, f)\sigma(\Gamma_x, g)}} \right), \\ \mathbf{R}(A) &= \Omega \left(\min_{\Gamma[f, g] \neq 0, f(x) \neq g(x)} \max \left\{ \frac{\sigma(\Gamma, f)}{\sigma(\Gamma_x, f)}, \frac{\sigma(\Gamma, g)}{\sigma(\Gamma_x, g)} \right\} \right). \end{aligned}$$

Theorem 4.2. *The following lower bounds hold for the query complexity of DDH in level 2 hidden cyclic groups:*

$$\mathbf{Q}(\text{DDH}(G_{p,2})) = \Omega(\sqrt{p}) \quad \text{and} \quad \mathbf{R}(\text{DDH}(G_{p,2})) = \Omega(p).$$

Proof. Let $i = ((1, 0, 0), (0, 1, 0), (0, 0, 1), (0, 1, 1))$. Observe that the element $(1, 0, 0)$ is a generator of $G_{p,2}$, for any 2-suitable vector $n = (1, n_1, n_2)$. By Proposition 3.6, we know that i is a DH-quadruple if and only if $n_1 + n_2 = n_1 n_2$. We say that n is *positive* if this equality holds, otherwise we say that it is *negative*. Let $m = p^3$ and let $S = \{\text{Id}_n : n \in \mathbb{Z}_p^2\}$. We will apply Fact 4.1 to the boolean oracle function DDH defined in $G_{p,2}$ on input i with the oracle input being the identity oracle $\text{Id}_n : \mathbb{Z}_p^3 \rightarrow \{0, 1\}$. For simplicity we will refer to this boolean oracle function just by DDH(n). We define the symmetric $p^2 \times p^2$ boolean adversary matrix Γ as follows:

$$\Gamma[n, n'] = \begin{cases} 1 & \text{if } \text{DDH}(n) \neq \text{DDH}(n'), \\ 0 & \text{otherwise,} \end{cases}$$

where again $\Gamma[n, n']$ is a shorthand notation for $\Gamma[\text{Id}_n, \text{Id}_{n'}]$.

We first determine $\sigma(\Gamma, n)$. If $n_1 = 1$ then there is no n_2 such that $n_1 + n_2 = n_1 n_2$. Otherwise, for every fixed $n_1 \neq 1$, there is a unique n_2 that makes this equality hold, in particular $n_2 = n_1(n_1 - 1)^{-1}$. Therefore the number of positive n is $p - 1$ and the number of negative n is $p^2 - p + 1$. Thus we have the following values for $\sigma(\Gamma, n)$:

$$\sigma(\Gamma, n) = \begin{cases} p^2 - p + 1 & \text{if } n \text{ is positive,} \\ p - 1 & \text{otherwise.} \end{cases}$$

Let us recall, that by definition, for every $h \in G_{p,2}$,

$$\Gamma_h[n, n'] = \begin{cases} 1 & \text{if } \text{DDH}(n) \neq \text{DDH}(n') \text{ and } \text{Id}_n(h) \neq \text{Id}_{n'}(h), \\ 0 & \text{otherwise.} \end{cases} \quad (1)$$

We fix now n and n' such that $\text{DDH}(n) \neq \text{DDH}(n')$, we will suppose without loss of generality that n is positive and n' is negative. We also fix $h = (h_0, h_1, h_2)$ in \mathbb{Z}_p^3 such that $\text{Id}_n(h) \neq \text{Id}_{n'}(h)$. This implies that $(h_1, h_2) \neq (0, 0)$. We want to lower bound $\sigma(\Gamma, n)/\sigma(\Gamma_h, n)$ and $\sigma(\Gamma, n')/\sigma(\Gamma_h, n')$. Obviously both fractions are at least 1. We distinguish two cases according to whether $\text{Id}_n(h) = 0$ or $\text{Id}_{n'}(h) = 0$.

Case 1: $\text{Id}_{n'}(h) = 0$. Then

$$\sigma(\Gamma_h, n') = \left| \left\{ (m_1, m_2) \in \mathbb{Z}_p^2 : \begin{array}{l} m_1 + m_2 = m_1 m_2 \text{ and} \\ h_0 + h_1 m_1 + h_2 m_2 = 0 \end{array} \right\} \right|.$$

We claim that the cardinality at the right hand side is at most 2. We know already that $m_1 \neq 1$ and $m_2 = m_1(m_1 - 1)^{-1}$. Therefore m_1 satisfies the second degree equation

$$h_1 x^2 + (h_0 - h_1 + h_2 - 1)x + (1 - h_0) = 0.$$

The number of roots of this equation is at most 2, unless the polynomial is 0. But this can not be the case, because then $h_1 = h_2 = 0$, a contradiction. Therefore, taking into account (1), we have

$$\frac{\sigma(\Gamma, n')}{\sigma(\Gamma_h, n')} = \Omega\left(\frac{p}{1}\right) = \Omega(p). \quad (2)$$

Case 2: $\text{Id}_n(h) = 0$. Then

$$\sigma(\Gamma_h, n) = |\{(m_1, m_2) \in \mathbb{Z}_p^2 : h_0 + h_1 m_1 + h_2 m_2 = 0\}|.$$

Since $(h_1, h_2) \neq (0, 0)$, the number of roots of this linear equation with two variables is p . Therefore, again taking into account (1), we have

$$\frac{\sigma(\Gamma, n)}{\sigma(\Gamma_h, n)} = \Omega\left(\frac{p^2}{p}\right) = \Omega(p). \quad (3)$$

The statements of the theorem immediately follow from equations (2) and (3). \square

Similarly to the remarks after Lemmas 3.4 and 3.8, we could have used in the proof instead of i a random input (g, h, k, ℓ) , with high probability. Indeed, let us suppose that we fix any $u = (u_0, u_1, u_2)$ in \mathbb{Z}_p^3 , with $(u_1, u_2) \neq (0, 0)$, to evaluate the properties of the matrix Γ_u . If we can show that the number of solutions of the system of equations

$$\begin{cases} (g_0 + g_1 x + g_2 y)(\ell_0 + \ell_1 x + \ell_2 y) \\ \quad - (h_0 + h_1 x + h_2 y)(k_0 + k_1 x + k_2 y) = 0 \\ u_0 + u_1 x + u_2 y = 0 \end{cases}$$

is at most 2, then the same proof works. Without loss of generality we can suppose that $u_2 \neq 0$. We express y from the second equation as function of x and substitute it in the first equation. We claim that the second degree polynomial we get, in particular its constant member, is non-zero with probability at least $(p-2)/p$. The value of the constant member besides u depends on $g_0, g_2, \ell_0, \ell_2, h_0, h_2, k_0, k_2$. We fix arbitrarily all these random variables except g_0 and ℓ_0 . Then, for some constants $a, b, c \in \mathbb{Z}_p$, we need to evaluate the probability of the event $(g_0 + a)(\ell_0 + b) \neq c$, which is at least $(p-2)/p$.

Acknowledgments

Parts of this research was accomplished while the first two authors were visiting the Centre for Quantum Technologies at the National University of Singapore and also while the last two authors were Fellows in November 2018 at the Stellenbosch Institute for Advanced Study. They would like to thank STIAS for support and hospitality. The research at CQT was supported by the Singapore National Research Foundation, the Prime Minister's Office, Singapore and the Ministry of Education, Singapore under the Research Centres of Excellence programme under research grant R 710-000-012-135. This research was also partially funded by the Hungarian National Research, Development and Innovation Office – NKFIH. In addition, this work has been supported in part by the European Union as H2020 Programme under grant agreement number ERC-669891 and by the QuantERA ERA-NET Cofund project QuantAlgo.

References

- [1] S. Aaronson, Lower bounds for local search by quantum arguments, in: Proceedings of the 36th Annual ACM Symposium on the Theory of Computing (STOC), pp. 465-474, 2004.
- [2] A. Ambainis, Quantum lower bounds by quantum arguments, *Journal of Computer and System Sciences* 64 (2010), 750-767.
- [3] A. Ambainis, Polynomial degree vs. quantum query complexity. *Journal of Computer and System Sciences* 72:2 (2006), 220-238, 2006.
- [4] L. Babai and R. Beals, A polynomial-time theory of black-box groups I, in: *Groups St. Andrews 1997 in Bath, I*, London Math. Soc. Lecture Notes Ser., vol. 260, Cambridge Univ. Press, pp. 30-64, 1999.
- [5] L. Babai and E. Szemerédi, On the complexity of matrix group problems I, in: Proceedings of the 25th IEEE Symposium on Foundations of Computer Science (FOCS), pp. 229-240, 1984.
- [6] R. Beals and L. Babai, Las Vegas algorithms for matrix groups, in: Proceedings of the 34th IEEE Foundations of Computer Science (FOCS), pp. 427-436, 1993.
- [7] R. Beals, H. Buhrman, R. Cleve, M. Mosca, and R. de Wolf, Quantum lower bounds by polynomials, *Journal of the ACM* 48:4 (2001), 778-797.
- [8] C. Bennett, E. Bernstein, G. Brassard and U. Vazirani, The strengths and weaknesses of quantum computation, *SIAM Journal on Computing* 26:5 (1997), 1510-1523.
- [9] K. Cheung and M. Mosca, Decomposing finite abelian groups, *Quantum Information and Computation* 1:3 (2001), 26-32.
- [10] I. Damgård, C. Hazay and A. Zottarel, Short paper on the generic hardness of DDH-II, Manuscript, <http://cs.au.dk/~angela/Hardness.pdf>, 2014.
- [11] L. Grover, A fast quantum mechanical algorithm for database search, in: Proceedings of the 28th Annual ACM Symposium on the Theory of Computing (STOC), pp. 212-219, 1996.

- [12] P. Høyer, T. Lee and R. Špalek, Negative weights make adversaries stronger, in: Proceedings of the 39th ACM Symposium on Theory of Computing (STOC), pp. 526-535, 2007.
- [13] A. Joux and K. Nguyen, Separating Decision Diffie-Hellman from Computational Diffie-Hellman in Cryptographic Groups, *Journal of Cryptology* 1 (2003), 239-247.
- [14] W. Kantor and Á. Seress, Black box classical groups, American Mathematical Society, Issue 708, 2001.
- [15] T. Lee, R. Mittal, B. Reichardt, R. Špalek and M. Szegedy, Quantum query complexity of state conversion, in: Proceedings of the 34th IEEE Foundations of Computer Science (FOCS), pp. 344-353, 2011.
- [16] U. Maurer and S. Wolf, The relationship between breaking the Diffie–Hellman protocol and computing discrete logarithms, *SIAM J. Comput* 28:5 (1999), 1689-1721.
- [17] M. Mosca, Quantum Computer Algorithms, PhD thesis, University of Oxford, 1999.
- [18] S. Pohlig and M. Hellman, An improved algorithm for computing logarithms over $GF(p)$ and its cryptographic significance, *IEEE Transactions on Information Theory* 24 (1978), 106-110.
- [19] D. Shanks, Five number-theoretic algorithms, In: Proceedings of the 2nd Manitoba Conference on Numerical Mathematics, pp. 51-70, 1972.
- [20] P. Shor, Algorithms for quantum computation: Discrete logarithm and factoring, *SIAM Journal on Computing*, 26:5 (1997), pp. 1484-1509.
- [21] V. Shoup, Lower bounds for discrete logarithms and related problems, in: Proceedings of the 16th Annual International Conference on Theory and Application of Cryptographic Techniques (EUROCRYPT), pp. 256-266, 1997.
- [22] J. Watrous, Quantum algorithms for solvable groups, In: Proceedings of the 33rd ACM Symposium on Theory of Computing (STOC), pp. 60-67, 2001.
- [23] S. Zhang, On the power of Ambainis’s lower bounds, *Theoretical Computer Science* 339 (2005), 241-256.