

The second law of thermodynamics from concavity of energy eigenvalues

C. Itoi and M. Amano

*Department of Physics, GS & CST, Nihon University,
Kandasurugadai, Chiyoda, Tokyo 101-8308, Japan*

(Dated: December 6, 2019)

Quantum dynamics controlled by a time-dependent coupling constant are studied. It is proven that an energy eigenstate expectation value of work done by the system in a quench process cannot exceed the work in the corresponding quasi-static process, if the energy eigenvalue is a concave function of the coupling constant. We propose this concavity of energy eigenvalues as a new universal criterion for quantum dynamical systems to satisfy the second law of thermodynamics. This criterion can be checked for a concerned quantum system within the framework of time-independent quantum mechanics. We give a simple universal reason for the concavity, and prove that every energy eigenvalue is indeed concave in some specific quantum systems. These results agree with the maximal work principle in the adiabatic environment as an expression of the second law of thermodynamics. Our result gives a simple example of an integrable system satisfying an analogue to the strong eigenstate thermalization hypothesis (ETH) with respect to the principle of maximum work.

PACS numbers:

a. Introduction Prove the second law of thermodynamics within the framework of mechanics. This fundamental problem has attracted many physicists for more than hundred years. Many attempts to clarify the second law of thermodynamics in classical or quantum mechanics have been made. Roughly speaking, there are two categories of these studies. One is to clarify the thermalization in isolated mechanical systems, and another one is to verify the impossibility to make any perpetual motion of second kind. In the first category, a remarkable philosophy of typical pure quantum state for thermally equilibrium states has been proposed recently. This states that a typical pure state in all possible mechanical states can be a thermally equilibrium state, and any state approaches to a typical state belonging to a set of all states most likely¹⁻⁵. The eigenstate thermalization hypothesis (ETH) assumes that almost all energy eigenstates are thermally equilibrium states⁶⁻¹³. The strong version of ETH claims that all energy eigenstates are thermally equilibrium states^{10,14-23}. There are several studies of ETH for integrable and non-integrable systems, and some of these have shown that the strong ETH fails for integrable systems^{10,14,15,23-28}. Interactions among many particles are essential to understand thermalization.

In another category, there have been fewer studies than those in the first category. Planck's principle is well-known as a specific expression of the impossibility of perpetual motion of second kind. This principle claims that a positive amount of work cannot be extracted from thermodynamic systems in an arbitrary cyclic operation in the adiabatic environment. Any realization of a perpetual motion of second kind implies the violation of this principle. Passivity is known as a sufficient condition on a quantum state for the impossibility of work extraction from quantum systems. We say that a quantum state is passive, if a positive amount of work cannot be extracted from this state as an initial state in any cyclic unitary evolution. Some mixed states, such as the Gibbs

states are passive^{29,30}, but general pure states except the ground state are not passive. The passivity on quantum states is stronger than Planck's principle, since a set of all unitary transformations contains unphysical time evolution. Therefore, it is worth studying possibility or impossibility of work extraction for an arbitrary cyclic process given by a physical evolution from an initial pure state. There have been several studies on the work extraction by a physical evolution from an initial pure state in an adiabatic environment³¹⁻³⁹. From the view point of the second law of thermodynamics, however, a few studies of the work extraction have been reported. Kaneko, Ioda and Sagawa study the validity of Planck's principle analogue to the strong ETH in quantum spin systems³⁹. They obtain numerical evaluations of amounts of works from energy eigenstates of quantum spin systems in a composite cyclic process which consists of instantaneous changes of the interaction and leaving the system a while. They find the strong ETH like behavior only in non-integrable systems. In integrable systems, however, they find only weak ETH behavior.

In the present paper, we propose a new universal criterion "concavity of energy eigenvalues" for quantum mechanical systems to satisfy the principle of maximum work in the adiabatic environment as an expression of the second law of thermodynamics. This criterion is expected to be useful, since it can be checked in each concerned system within the framework of time-independent quantum mechanics. The principle of maximum work in the adiabatic environment states that work done by the system in an arbitrary process cannot exceed the work in the corresponding quasi-static process. This is equivalent to Planck's principle, since the quasi-static process is reversible. We study quantum dynamical processes controlled by a time-dependent coupling constant. We evaluate the changes of the energy under quench and quasi-static processes solving the time-dependent Schrödinger equation and using the adiabatic theorem. We derive

several formulae among expectation values of important physical quantities in an arbitrary energy eigenstate using the Hellmann-Feynman theorem^{40,41}. These formulae enable us to prove that an energy eigenstate expectation value of work done by the system in a quench process cannot exceed the work in the corresponding quasi-static process, if its energy eigenvalue is a concave function of the coupling constant. We give several simple examples in a specific many-particle systems confined in a bounded region by a confining potential with a time-dependent coupling constant where every energy eigenvalue is a concave function of the coupling constant. Particularly in free harmonic oscillators, it is shown that the amount of work in the process with an arbitrary speed cannot exceed that in the quasi-static process. A system of free particles is an example of integrable system which satisfies an analogue to the strong ETH for the principle of maximum work in the adiabatic environment. Finally, we point out that a possible reason for the fail of the strong ETH like behavior of integrable systems reported in preceding studies is violation of the adiabatic theorem by the level crossing of the systems.

b. Quantum dynamics of a time-dependent Hamiltonian Here we consider a general quantum many-body system under quench and quasi-static changes of the coupling constant. The time-dependent state $\Psi(t)$ satisfies the following Schrödinger equation

$$i\hbar \frac{\partial}{\partial t} \Psi(t) = \mathcal{H}(\lambda(t)) \Psi(t). \quad (1)$$

The time-dependent Hamiltonian $\mathcal{H}(\lambda(t))$ consists of a time-dependent term $\lambda(t)H_I$ with a time-dependent coupling constant $\lambda(t)$ and another time-independent term H

$$\mathcal{H}(\lambda(t)) := H + \lambda(t)H_I, \quad (2)$$

where H and H_I are self-adjoint operators. Assume that H has no upper bound, and H_I is positive semi-definite $H_I \geq 0$ with a positive coupling constant $\lambda > 0$, where the bound on an arbitrary self-adjoint operator O is defined by the expectation value $\langle O \rangle_\chi$ optimized by a suitable normalized state χ .

Let Φ_k be a normalized energy eigenstate which belongs to the eigenvalue $\mathcal{E}_k(\lambda)$ as a function of λ

$$\mathcal{H}(\lambda)\Phi_k = \mathcal{E}_k(\lambda)\Phi_k. \quad (3)$$

For $t < 0$, let $\Psi(t) = \exp[-it\mathcal{E}_k(\lambda_0)/\hbar]\Phi_k$ be an initial state, and consider a process by a quench $\lambda_0 \rightarrow \lambda_1$ at $t = 0$. The continuity of the state at $t = 0$ implies that the state after the quench is given by

$$\Psi(t) = \exp\left[-it\mathcal{H}(\lambda_1)/\hbar\right]\Phi_k, \quad (4)$$

The work done by the system in this quench process is

$$\begin{aligned} & \mathcal{E}_k(\lambda_0) - \langle \mathcal{H}(\lambda_1) \rangle_\Psi \\ &= \langle (H + \lambda_0 H_I) \rangle_{\Phi_k} - \langle (H + \lambda_1 H_I) \rangle_{\Phi_k} \\ &= (\lambda_0 - \lambda_1) \langle H_I \rangle_{\Phi_k}. \end{aligned} \quad (5)$$

The derivative of the eigenvalue equation with respect to the coupling constant and internal products between Φ_k and both sides give

$$\langle H_I \rangle_{\Phi_k} = \mathcal{E}'_k(\lambda). \quad (6)$$

This identity is known as the Hellmann-Feynman theorem^{40,41}. The assumption of the positive semi-definiteness of $\langle H_I \rangle_{\Phi_k}$ implies that the energy eigenvalue is monotonically increasing in the coupling constant. Increasing λ gives a positive amount of work to the system. We have the following formula for quantum interacting systems

$$\mathcal{E}'_k(\lambda) = \frac{\mathcal{E}_k(\lambda) - \langle H \rangle_{\Phi_k}}{\lambda}. \quad (7)$$

and another one

$$\mathcal{E}''_k(\lambda) = -\frac{1}{\lambda} \frac{d\langle H \rangle_{\Phi_k}}{d\lambda}. \quad (8)$$

The work in the quench process becomes

$$(\lambda_0 - \lambda_1) \langle H_I \rangle_{\Phi_k} = (\lambda_0 - \lambda_1) \mathcal{E}'_k(\lambda_0). \quad (9)$$

On the other hand, the work done by the particle in the quasi-static process is

$$\mathcal{E}_k(\lambda_0) - \mathcal{E}_k(\lambda_1)$$

since a quasi-static process preserves a quantum number of the energy eigenstate by the quantum adiabatic theorem. Therefore, we have the following theorem. If $\mathcal{E}_k(\lambda)$ is a concave function of λ , the work in the quench process from the initial energy eigenstate Φ_k cannot exceed that in the corresponding quasi-static process

$$(\lambda_0 - \lambda_1) \mathcal{E}'_k(\lambda_0) \leq \mathcal{E}_k(\lambda_0) - \mathcal{E}_k(\lambda_1). \quad (10)$$

The identity (8) implies that the concavity of $\mathcal{E}_k(\lambda)$ is guaranteed by the monotonically increasing $\langle H \rangle_{\Phi_k}$ in λ . In the quasi-static change $\lambda_0 \rightarrow \lambda_1$ with $0 < \lambda_0 < \lambda_1$, the change of energy is positive semi-definite

$$\Delta \mathcal{E}_k = \mathcal{E}_k(\lambda_1) - \mathcal{E}_k(\lambda_0) \geq 0,$$

since $\mathcal{E}'_k(\lambda) = \langle H_I \rangle_{\Phi_k} \geq 0$. The increasing energy given by this operation is shared by the following two terms

$$\Delta \mathcal{E}_k = \Delta \langle H \rangle_{\Phi_k} + \Delta(\lambda \langle H_I \rangle_{\Phi_k}).$$

It is intuitively obvious that both terms are positive semi-definite for any k

$$\Delta \langle H \rangle_{\Phi_k} \geq 0, \quad \Delta(\lambda \langle H_I \rangle_{\Phi_k}) \geq 0.$$

under the assumption that the time-independent Hamiltonian H has no upper bound. This implies that $\langle H \rangle_{\Phi_k}$ is a monotonically increasing function of λ for any k . We assume the monotonicity of $\langle H \rangle_{\Phi_k}$ which implies the concavity of the eigenvalue $\mathcal{E}_k(\lambda)$ for any k . Therefore, the principle of maximum work in the adiabatic environment is valid in the strong sense in general many-body systems.

c. Many-particle systems Here we consider a quantum dynamical system of N particles confined in a bounded region in one dimension. Let $x = (x_1, \dots, x_N)$ be a set of positions of N particles, and let $\Psi(x, t)$ be a time-dependent state of particles satisfying the Schrödinger equation. The time-dependent Hamiltonian $\mathcal{H}(\lambda(t))$ has a positive semi-definite confining potential $u(x_n)$ with a time-dependent coupling constant $\lambda(t)$ and another time-independent term H , which consists of kinetic energy and interaction $V(x)$ among N particles

$$H := -\frac{\hbar^2}{2} \sum_{n=1}^N \frac{\partial^2}{\partial x_n^2} + V(x), \quad (11)$$

$$H_I := \sum_{n=1}^N u(x_n). \quad (12)$$

First, consider the case $V(x) = 0$. Let $\phi_{k_n}(x_n)$ be a single particle energy eigenstate of n -th particle with an eigenvalue $E_{k_n}(\lambda)$. If this function is concave function of λ for any n , the theorem is valid also for any eigenstate of N free particles for a set $k = (k_1, \dots, k_N)$ of quantum numbers

$$\Phi_k(x) = \prod_{n=1}^N \phi_{k_n}(x_n),$$

with the eigenvalue

$$\mathcal{E}_k(\lambda) = \sum_{n=1}^N E_{k_n}(\lambda). \quad (13)$$

Note that the energy of each particle is conserved for a fixed λ . This system is integrable, since it has N coordinates and N conserved quantities. Let us give a simple example where all energy eigenvalues are concave functions of λ . Consider N free particles in a potential

$$u(x) = \frac{1}{2j} x^{2j}, \quad (14)$$

with a positive integer j . This potential $u(x)$ is positive semi-definite. The virial theorem⁴² and the identity (6) imply

$$\langle H \rangle_{\Phi_k} = \sum_n \frac{\lambda}{2} \langle x_n u'(x_n) \rangle_{\Phi_k} = \lambda j \sum_{n=1}^N \langle u(x_n) \rangle_{\Phi_k} = \lambda j \mathcal{E}'_k(\lambda). \quad (15)$$

This and the identity (7) give the following differential equation for $\mathcal{E}_k(\lambda)$

$$(j+1)\lambda \mathcal{E}'_k(\lambda) = \mathcal{E}_k(\lambda). \quad (16)$$

The solution of this differential equation is given by

$$\mathcal{E}_k(\lambda) = C_k \lambda^{\frac{1}{j+1}}, \quad (17)$$

with a positive constant C_k independent of λ . This shows that the function $\mathcal{E}_k(\lambda)$ is concave. Therefore, the the

work in the quench process cannot exceed that in the quasi-static process for any initial eigenstate Φ_k . This result suggests an analogue to the strong ETH, despite the integrability of free particles. The same argument based on the virial theorem is possible for a model with an interaction

$$V(x) := \sum_{1 \leq m < n \leq N} \frac{g}{(x_m - x_n)^2}. \quad (18)$$

In this system, we have the solution (17) and every energy eigenvalue is a concave function of λ . The generalization of our argument based on the virial theorem to models in an arbitrary dimension is also possible.

d. An arbitrary process in harmonic oscillators In the specific case of $j = 1$ and $V = 0$, the system becomes a set of N independent harmonic oscillators. In this case, we can prove that the quasi-static work gives the upper bound on the work done by any operation with an arbitrary speed including the quench operation. Define a time-dependent parameter $\omega_t := \sqrt{\lambda(t)}$, and define the following time-dependent annihilation and creation operators for each particle

$$a_t := \sqrt{\frac{\omega_t}{2\hbar}} x + \sqrt{\frac{\hbar}{2\omega_t}} \frac{\partial}{\partial x}, \quad (19)$$

$$a_t^\dagger := \sqrt{\frac{\omega_t}{2\hbar}} x - \sqrt{\frac{\hbar}{2\omega_t}} \frac{\partial}{\partial x}. \quad (20)$$

These have a commutation relation $[a_t, a_t^\dagger] = 1$. These time-dependent operators can be represented in terms of initial operators. The Hamiltonian for one particle is given by

$$\mathcal{H}(\lambda(t)) = -\frac{\hbar^2}{2} \frac{\partial^2}{\partial x^2} + \frac{\lambda(t)}{2} x^2 = \hbar\omega_t \left(a_t^\dagger a_t + \frac{1}{2} \right) \quad (21)$$

Let $U(t)$ be a unitary operator which represents the time-dependent solution $\psi(x, t) = U(t)\phi_k(x)$ of the Schrödinger equation with an initial state $\phi_k(x)$ with an energy eigenvalue $E_k = \hbar\omega_0(k + \frac{1}{2})$, ($k = 0, 1, 2, \dots$) Define unitary transformed oscillators for the time evolution by represent them in the following Bogoliubov transformation of the initial operators

$$U(t)^\dagger a_t U(t) = X(t)a_0 + Y(t)a_0^\dagger, \quad (22)$$

$$U(t)^\dagger a_t^\dagger U(t) = X(t)a_0^\dagger + Y(t)^* a_0. \quad (23)$$

Note that $|X(t)|^2 - |Y(t)|^2 = 1$. These satisfy the following differential equations

$$i\dot{X}(t) = \omega_t X(t) - i\frac{\dot{\omega}_t}{2\omega_t} Y(t)^*, \quad (24)$$

$$i\dot{Y}(t) = \omega_t Y(t) - i\frac{\dot{\omega}_t}{2\omega_t} X(t)^*, \quad (25)$$

with an initial condition $X(0) = 1, Y(0) = 0$. The energy of the particle at t is given by

$$\begin{aligned} \langle \mathcal{H}(\lambda(t)) \rangle_\psi &= \langle \phi_k, U(t)^\dagger \mathcal{H}(\lambda(t)) U(t) \phi_k \rangle \\ &= \hbar\omega_t \left(k + \frac{1}{2} \right) + 2\hbar\omega_t |Y(t)|^2 \left(k + \frac{1}{2} \right) \end{aligned} \quad (26)$$

Since $Y(t) = 0$ in a quasi-static process, the work done by the particle is $-2\hbar\omega_t\left(k + \frac{1}{2}\right)$. For N independent particles, the difference between the work done by N independent particles in an arbitrary process and that in a quasi-static process is given by the summation over the contributions of N particles

$$-2\hbar\omega_t \sum_{n=1}^N |Y_n(t)|^2 \left(k_n + \frac{1}{2}\right) \leq 0.$$

This shows that the work done by N particles in the quasi-static process gives the maximum work in the adiabatic environment.

e. Summary We prove that the concavity of the energy eigenvalue in the coupling constant is necessary for the principle of maximum work. This criterion can be checked in each concerned system within the frame work

of time-independent quantum mechanics. We give an example with a many-particle system in a specific potential where every energy eigenvalue becomes a concave function of the coupling constant. This is an example of an integrable system which satisfies the strong ETH analogue with respect to Planck's principle.

Finally, we comment on quantum systems with level crossing of energy eigenvalues by the change of coupling constant. In such systems, the adiabatic theorem fails, and concavity of the energy eigenvalues does not yields Planck's principle in strong sense as pointed out by Kaneko, Ioda and Sagawa³⁹. We consider that level crossing likely occurs in integrable quantum systems.

Acknowledgments

We are grateful to R. M. Woloshyn for a careful reading of the manuscript. It is pleasure to thank T. Sako for helpful discussions.

-
- ¹ H. Tasaki, Phys. Rev. Lett. **80**, 1373 (1998).
² S. Goldstein, T. Hara and H. Tasaki, Phys. Rev. Lett. **111**, 140401(2013).
³ H. Tasaki, J. Stat. Phys. **163**, 937 (2016).
⁴ S. Goldstein, J. L. Lebowitz, C. Mastrodonato, R. Tumulka, and N. Zanghì, Proc. R. Soc. A **466**, 3203 (2010).
⁵ S. Sugiura and A. Shimizu, Phys. Rev. Lett. **111**, 010401 (2013)
⁶ M. V. Berry, J. Phys. A **10**, 2083 (1977).
⁷ J. M. Deutsch, Phys. Rev. A **43**, 2046 (1991).
⁸ B. Li, and M. Robnik, J. Phys. A **27**, 5509 (1994).
⁹ M. Srednicki, Phys. Rev. E **50**, 888 (1994).
¹⁰ M. Rigol, V. Dunjko and M. Olshanii, Nature **452**, 854 (2008).
¹¹ A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, Rev. Mod. Phys. **83**, 863 (2011).
¹² L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, Adv. Phys. **65**, 239 (2016).
¹³ H. Wilming, M. Goihl, I. Roth, and J. Eisert, Phys. Rev. Lett. **123**, 200604 (2019)
¹⁴ R. V. Jensen and R. Shankar, Phys. Rev. Lett. **54**, 1879 (1985).
¹⁵ R. Steinigeweg, J. Herbrych, and P. Prelovšek, Phys. Rev. E **87**, 012118 (2013).
¹⁶ R. Steinigeweg, A. Khodja, H. Niemeyer, C. Gogolin, and J. Gemmer, Phys. Rev. Lett. **112**, 130403 (2014).
¹⁷ W. Beugeling, R. Moessner, and M. Haque, Phys. Rev. E **89**, 042112 (2014).
¹⁸ H. Kim, T. N. Ikeda, and D. A. Huse, Phys. Rev. E **90**, 052105 (2014).
¹⁹ A. Khodja, R. Steinigeweg, and J. Gemmer, Phys. Rev. E **91**, 012120 (2015).
²⁰ R. Mondaini, K. R. Fratus, M. Srednicki, and M. Rigol, Phys. Rev. E **93**, 032104 (2016).
²¹ J. R. Garrison and T. Grover, Phys. Rev. X **8**, 021026 (2018).
²² A. Dymarsky, N. Lashkari, and H. Liu, Phys. Rev. E **97**, 012140 (2018).
²³ T. Yoshizawa, E. Iyoda and T. Sagawa, Phys. Rev. Lett. **120**, 200604 (2018).
²⁴ M. Rigol, Phys. Rev. Lett. **103**, 100403 (2009).
²⁵ G. Biroli, C. Kollath, and A. M. Läuchli, Phys. Rev. Lett. **105**, 250401 (2010).
²⁶ V. Alba, Phys. Rev. B **91**, 155123 (2015).
²⁷ F. H. L. Essler and M. Fagotti, J. Stat. Mech. **064002** (2016).
²⁸ L. Vidmar and M. Rigol, J. Stat. Mech. **064007** (2016).
²⁹ W. Pusz and S. L. Woronowicz, Comm. Math. Phys. **58**, 273 (1978).
³⁰ A. Lenard, J. Stat. Phys. **19**, 575 (1978).
³¹ R. Dorner, S. R. Clark, L. Heaney, R. Fazio, J. Goold, and V. Vedral, Phys. Rev. Lett. **110**, 230601 (2013).
³² R. Gallego, A. Riera, and J. Eisert, New J. Phys. **16**, 125009 (2014).
³³ M. Perarnau-Llobet, A. Riera, R. Gallego, H. Wilming, and J. Eisert, New J. Phys. **18**, 123035 (2016).
³⁴ R. Modak and M. Rigol, Phys. Rev. E **95**, 062145 (2017).
³⁵ T. P. Le, J. Levinsen, K. Modi, M. M. Parish, and F. A. Pollock, Phys. Rev. A **97**, 022106 (2018).
³⁶ F. Jin, R. Steinigeweg, H. De Raedt, K. Michielsen, M. Campisi, and J. Gemmer, Phys. Rev. E **94**, 012125 (2016).
³⁷ D. Schmidtke, L. Knipschild, M. Campisi, R. Steinigeweg, and J. Gemmer, Phys. Rev. E **98**, 012123 (2018).
³⁸ S. Deffner, J. P. Paz, and W. H. Zurek, Phys. Rev. E **94**, 010103(R) (2016).
³⁹ K. Kaneko, E. Iyoda and T. Sagawa, Phys. Rev. E **99**, 032128 (2019).
⁴⁰ H. Hellmann, Einführung in die Quantenchemie (Deuticke, Leipzig, Germany, 1937)
⁴¹ R. P. Feynman, Phys. Rev. **56**, 340 (1939).
⁴² V. Fock, Z. Phys. A. **63**, 855 (1930).