

HILBERT SERIES AND INVARIANTS IN EXTERIOR ALGEBRAS

ELITZA HRISTOVA

ABSTRACT. In this paper, we consider the exterior algebra $\Lambda(W)$ of a polynomial $\mathrm{GL}(n)$ -module W and use previously developed methods to determine the Hilbert series of the algebra of invariants $\Lambda(W)^G$, where G is one of the classical complex subgroups of $\mathrm{GL}(n)$, namely $\mathrm{SL}(n)$, $\mathrm{O}(n)$, $\mathrm{SO}(n)$, or $\mathrm{Sp}(2d)$ (for $n = 2d$). Since $\Lambda(W)^G$ is finite dimensional, we apply the described method to compute a lot of explicit examples. For $\Lambda(S^3\mathbb{C}^3)^{\mathrm{SL}(3)}$, using the computed Hilbert series, we obtain an explicit set of generators.

1. INTRODUCTION

Let $\mathrm{GL}(n)$ denote the complex general linear group and let W be a finite dimensional polynomial $\mathrm{GL}(n)$ -module. Then W can be written as a direct sum of its irreducible components

$$W = \bigoplus_{\lambda} k(\lambda)V_{\lambda}^n,$$

where $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{N}_0^n$, $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$, is a non-negative integer partition (here $\mathbb{N}_0 = \{0, 1, 2, \dots\}$) and V_{λ}^n denotes the irreducible $\mathrm{GL}(n)$ -module with highest weight λ . Let $A = \bigoplus_{i \geq 0} A^i$ be a finitely generated graded algebra over \mathbb{C} for which each homogeneous component A^i is a polynomial $\mathrm{GL}(n)$ -module. In the papers [5, 6], Drensky and the author develop a method for determining the Hilbert series of the algebra of invariants A^G , where G is one of the classical complex groups $\mathrm{O}(n)$, $\mathrm{SO}(n)$, and $\mathrm{Sp}(2d)$ (the last in the case when $n = 2d$). These results can be easily extended to the case $G = \mathrm{SL}(n)$ using an earlier work of Drensky and other authors [1]. A general class of examples of an algebra A with the above properties is given by the quotient $T(W)/I$, where $T(W)$ denotes the tensor algebra of a polynomial $\mathrm{GL}(n)$ -module W and I is a $\mathrm{GL}(n)$ -invariant ideal in $T(W)$. This class includes the symmetric and exterior algebras of W , $S(W)$ and $\Lambda(W)$, as well as the class of relatively free algebras in varieties of associative algebras. In the papers [5, 6], Drensky and the author use the developed method to study the Hilbert series $H(A^G, t)$ for A being the symmetric algebra $S(W)$ and for certain relatively free algebras with the property that their varieties are the only minimal varieties of exponent 2. In the present paper, we consider the case $A = \Lambda(W)$ and study the algebra of invariants $\Lambda(W)^G$, where G is again one of $\mathrm{SL}(n)$, $\mathrm{O}(n)$, $\mathrm{SO}(n)$, and $\mathrm{Sp}(2d)$. The case $A = \Lambda(W)$ is remarkable with the property that the considered Hilbert series are polynomials, which allows for explicit computation of a lot of examples. Invariants in exterior algebras for different actions of classical groups and related problems are the topic of many recent papers, see, e.g., [3, 4, 8].

2010 *Mathematics Subject Classification.* 13A50; 15A72; 15A75; 20G05.

Key words and phrases. invariant theory, Hilbert series, multiplicity series, exterior algebras.

The present paper is organized as follows. In Section 2 we give a brief introduction to the general method for determining Hilbert series which is developed in [6]. In Section 3 we describe how the general method is applied to $A = \Lambda(W)$ and in Section 4 we take concrete examples of W and compute explicitly the respective Hilbert series. The results are given in Tables 1-9 and in Example 4.2. In some cases, we use the computed Hilbert series to describe also a set of generators of the respective algebra of invariants (see Example 4.3). Since for most of the examples we use computer programs written in Mathematica, in the Appendix we provide a part of the source code.

2. PRELIMINARIES

The ground field is \mathbb{C} . We start by recalling the notion of Hilbert series, which applies to finitely generated graded algebras (resp., vector spaces) over \mathbb{C} .

Definition 2.1. Let $A = \bigoplus_{i \geq 0} A^i$ be a finitely generated graded algebra (or a vector space) over \mathbb{C} such that $A^0 = \mathbb{C}$ or $A^0 = 0$. The **Hilbert series** of A is the formal power series

$$H(A, t) = \sum_{i \geq 0} (\dim A^i) t^i.$$

The next definition is a generalization of the notion of Hilbert series for multigraded algebras (resp., for multigraded vector spaces).

Definition 2.2. Let

$$A = \bigoplus_{\mu \in \mathbb{N}_0^n} A(\mu)$$

be a finitely generated algebra or vector space with an \mathbb{N}_0^n -grading. The Hilbert series of A with respect to this grading is the formal power series $H(A, x_1, \dots, x_n) \in \mathbb{Z}[[x_1, \dots, x_n]]$ defined by

$$H(A, x_1, \dots, x_n) = \sum_{\mu = (\mu_1, \dots, \mu_n) \in \mathbb{N}_0^n} \dim A(\mu) x_1^{\mu_1} \cdots x_n^{\mu_n}.$$

One example of a vector space with an \mathbb{N}_0^n -grading is the $\mathrm{GL}(n)$ -module V_λ^n together with its weight space decomposition. The Hilbert series of V_λ^n with respect to this grading coincides with the character $\chi_{V_\lambda^n}(x_1, \dots, x_n)$ of V_λ^n and consequently has the form

$$H(V_\lambda^n, x_1, \dots, x_n) = \chi_{V_\lambda^n}(x_1, \dots, x_n) = S_\lambda(x_1, \dots, x_n),$$

where $S_\lambda(x_1, \dots, x_n)$ is the Schur polynomial corresponding to the partition λ . Similarly, each polynomial $\mathrm{GL}(n)$ -module W also has an \mathbb{N}_0^n -grading and a corresponding Hilbert series which is equal to the character of W .

Next, as in the Introduction, we consider the special class of finitely generated graded algebras $A = \bigoplus_{i \geq 0} A^i$, such that each homogeneous component A^i for $i \geq 0$ is a polynomial $\mathrm{GL}(n)$ -module. Equivalently, A can be defined as an arbitrary direct sum of polynomial $\mathrm{GL}(n)$ -modules. Then, we can write the decomposition of A in the following way:

$$A = \bigoplus_{i \geq 0} A^i = \bigoplus_{i \geq 0} \bigoplus_{\lambda} m_i(\lambda) V_\lambda,$$

where $A^i = \bigoplus_{\lambda} m_i(\lambda) V_{\lambda}$ is the decomposition of A^i into irreducible $\mathrm{GL}(n)$ -modules and the sum runs over all partitions λ in \mathbb{N}_0^n . Hence, A has two gradings – the \mathbb{N}_0 -grading coming from the decomposition into homogeneous components and an \mathbb{N}_0^n -grading coming from the \mathbb{N}_0^n -grading of each homogeneous component. Following [1, 5, 6], we introduce a Hilbert series of A which takes into account both gradings:

$$H(A; x_1, \dots, x_n, t) = \sum_{i \geq 0} H(A^i, x_1, \dots, x_n) t^i = \sum_{i \geq 0} \left(\sum_{\lambda} m_i(\lambda) S_{\lambda}(x_1, \dots, x_n) \right) t^i.$$

Let S_n denote the symmetric group in n variables. Then, $H(A; x_1, \dots, x_n, t) \in \mathbb{Z}[[x_1, \dots, x_n]]^{\mathrm{S}_n}[[t]]$, i.e., the coefficient in front of t^i for each $i \geq 0$ is a symmetric function in the variables x_1, \dots, x_n . For any symmetric function $f(x_1, \dots, x_n) \in \mathbb{C}[[x_1, \dots, x_n]]^{\mathrm{S}_n}$, one can define the notion of multiplicity series (see, e.g., [1]). Following [1, 5, 6], we introduce a generalization of the notion of multiplicity series for functions $f(x_1, \dots, x_n, t)$ in $\mathbb{C}[[x_1, \dots, x_n]]^{\mathrm{S}_n}[[t]]$. Let $f(x_1, \dots, x_n, t) \in \mathbb{C}[[x_1, \dots, x_n]]^{\mathrm{S}_n}[[t]]$, i.e., $f(x_1, \dots, x_n, t)$ can be written as

$$f(x_1, \dots, x_n, t) = \sum_{i \geq 0} \left(\sum_{\lambda} c_i(\lambda) S_{\lambda}(x_1, \dots, x_n) \right) t^i,$$

where $c_i(\lambda) \in \mathbb{C}$ and the second sum runs over all partitions $\lambda \in \mathbb{N}_0^n$. The multiplicity series of $f(x_1, \dots, x_n, t)$ is defined by:

$$(1) \quad M(f; x_1, \dots, x_n, t) = \sum_{i \geq 0} \left(\sum_{\lambda} c_i(\lambda) x_1^{\lambda_1} \cdots x_n^{\lambda_n} \right) t^i.$$

By the change of variables $v_1 = x_1, v_2 = x_1 x_2, \dots, v_n = x_1 \cdots x_n$, we rewrite the above multiplicity series in the following form:

$$(2) \quad M'(f; v_1, \dots, v_n, t) = \sum_{i \geq 0} \left(\sum_{\lambda} c_i(\lambda) v_1^{\lambda_1 - \lambda_2} v_2^{\lambda_2 - \lambda_3} \cdots v_{n-1}^{\lambda_{n-1} - \lambda_n} v_n^{\lambda_n} \right) t^i.$$

For any subgroup G of $\mathrm{GL}(n)$, the algebra of invariants A^G inherits the \mathbb{N}_0 -grading of A and therefore one can study the Hilbert series $H(A^G, t)$. The following theorem, given in [6], shows the relation between $H(A^G, t)$ and the multiplicity series of $H(A; x_1, \dots, x_n, t)$ defined by (1) and (2).

Theorem 2.3 ([6]). *Let $A = \bigoplus_{i \geq 0} A^i$ be a finitely generated graded algebra, such*

that each homogeneous component A^i for $i \geq 0$ is a polynomial $\mathrm{GL}(n)$ -module.

(i) *The Hilbert series of the algebra of invariants $A^{\mathrm{SL}(n)}$ is given by*

$$H(A^{\mathrm{SL}(n)}, t) = M'(H(A); 0, \dots, 0, 1, t).$$

(ii) *The Hilbert series of $A^{\mathrm{Sp}(2d)}$ (where $n = 2d$) is given by*

$$H(A^{\mathrm{Sp}(2d)}, t) = M'(H(A); 0, 1, 0, 1, \dots, 0, 1, t).$$

(iii) *The Hilbert series of $A^{\mathrm{O}(n)}$ is*

$$H(A^{\mathrm{O}(n)}, t) = M_n(t),$$

where M_n is defined iteratively in the following way:

$$M_1(x_2, \dots, x_n, t) = \frac{1}{2} (M(H(A); -1, x_2, \dots, x_n, t) + M(H(A); 1, x_2, \dots, x_n, t)),$$

$$M_2(x_3, \dots, x_n, t) = \frac{1}{2}(M_1(-1, x_3, \dots, x_n, t) + M_1(1, x_3, \dots, x_n, t))$$

.....

$$M_n(t) = \frac{1}{2}(M_{n-1}(-1, t) + M_{n-1}(1, t)).$$

(iv) *The Hilbert series of $A^{\text{SO}(n)}$ is*

$$H(A^{\text{SO}(n)}, t) = M'_n(t),$$

where

$$M'_1(v_2, \dots, v_n, t) = \frac{1}{2}(M'(H(A); -1, v_2, \dots, v_n, t) + M'(H(A); 1, v_2, \dots, v_n, t)),$$

$$M'_2(v_3, \dots, v_n, t) = \frac{1}{2}(M'_1(-1, v_3, \dots, v_n, t) + M'_1(1, v_3, \dots, v_n, t))$$

.....

$$M'_{n-1}(v_n, t) = \frac{1}{2}(M'_{n-2}(-1, v_n, t) + M'_{n-2}(1, v_n, t)),$$

$$M'_n(t) = M'_{n-1}(1, t).$$

3. THE ALGEBRA OF INVARIANTS $\Lambda(W)^G$

In this section we show how Theorem 2.3 can be applied to determine the Hilbert series of A^G for $A = \Lambda(W)$.

Let W be a p -dimensional polynomial $\text{GL}(n)$ -module, let ξ_1, \dots, ξ_p be a basis of W consisting of weight vectors, and let $\alpha_1 = (\alpha_{11}, \dots, \alpha_{1n}), \dots, \alpha_p = (\alpha_{p1}, \dots, \alpha_{pn})$ denote the corresponding weights (with possible repetitions). The Hilbert series of W with respect to the \mathbb{N}_0^n -grading given by the weight space decomposition of W is equal to the character χ_W of W and hence

$$H(W, x_1, \dots, x_n) = \sum_{j=1}^p x_1^{\alpha_{j1}} \cdots x_n^{\alpha_{jn}}.$$

Similarly, the Hilbert series $H(\Lambda^k(W), x_1, \dots, x_n)$ of $\Lambda^k(W)$ for $0 \leq k \leq p$ is equal to the character $\chi_{\Lambda^k(W)}$. It is a standard fact (see, e.g., [7]) that if $W = V$, where V denotes the natural representation of $\text{GL}(n)$, the character of $\Lambda^k(V)$ is equal to the k -th elementary symmetric polynomial, i.e.,

$$\chi_{\Lambda^k(V)}(x_1, \dots, x_n) = E_k(x_1, \dots, x_n) = \sum_{1 \leq s_1 < \dots < s_k \leq n} x_{s_1} \cdots x_{s_k}.$$

Using the above formula, one shows that

$$H(\Lambda^k(W), x_1, \dots, x_n) = \sum_{1 \leq s_1 < \dots < s_k \leq n} (x_1^{\alpha_{s_1 1}} \cdots x_n^{\alpha_{s_1 n}}) \cdots (x_1^{\alpha_{s_k 1}} \cdots x_n^{\alpha_{s_k n}}).$$

Therefore,

$$(3) \quad H(\Lambda(W), x_1, \dots, x_n, t) = \prod_{j=1}^p (1 + x_1^{\alpha_{j1}} \cdots x_n^{\alpha_{jn}} t).$$

Lemma 3 from [2] describes a way to determine the multiplicity series of each symmetric function $f(x_1, \dots, x_n) \in \mathbb{C}[[x_1, \dots, x_n]]^{\text{S}_n}$ (see also Lemma 1.1 from [1]). The following lemma is a direct generalization for functions in $\mathbb{C}[[x_1, \dots, x_n]]^{\text{S}_n}[[t]]$.

Lemma 3.1. *Let $X = \{x_1, \dots, x_n\}$ and let $f(X, t) \in \mathbb{C}[[X]]^{\mathbb{S}_n}[[t]]$. Let*

$$g(X, t) = f(X, t) \prod_{i < j} (x_i - x_j) = \sum_{i \geq 0} \left(\sum_{r_{i_j} \geq 0} \alpha_i(r_{i_1}, \dots, r_{i_n}) x_1^{r_{i_1}} \cdots x_n^{r_{i_n}} \right) t^i,$$

for some $\alpha_i(r_{i_1}, \dots, r_{i_n}) \in \mathbb{C}$. Then the multiplicity series of $f(X, t)$ is given by

$$M(f; x_1, \dots, x_n, t) = \frac{1}{x_1^{n-1} x_2^{n-2} \cdots x_{n-2}^2 x_{n-1}} \sum_{i \geq 0} \left(\sum_{r_{i_j} > r_{i_{j+1}}} \alpha_i(r_{i_1}, \dots, r_{i_n}) x_1^{r_{i_1}} \cdots x_n^{r_{i_n}} \right) t^i,$$

where the sum is over all $r_i = (r_{i_1}, \dots, r_{i_n})$ such that $r_{i_1} > r_{i_2} > \cdots > r_{i_n}$.

In the next section, we use this general procedure for concrete choices of W .

4. EXAMPLES

Let $V = \mathbb{C}^n$ denote the natural $\mathrm{GL}(n)$ -module and let again G be one of $\mathrm{SL}(n)$, $\mathrm{O}(n)$, $\mathrm{SO}(n)$, and $\mathrm{Sp}(2d)$ (for $n = 2d$). In the first set of examples, we consider the module $W = S^k V$, for $k \geq 3$, and determine the Hilbert series $H(\Lambda(W)^G, t)$ for fixed values of n and k . The case $k = 2$ is not included in the consideration, since explicit expressions for $H(\Lambda(S^2 V)^G, t)$ and $H(\Lambda(\Lambda^2 V)^G, t)$ for all n are already obtained in [5].

We notice first that when $W = S^k V$, Equation (3) implies that

$$H(\Lambda(S^k V), x_1, \dots, x_n, t) = \prod_{i_1 + \cdots + i_n = k} (1 + x_1^{i_1} \cdots x_n^{i_n} t).$$

In the simplest case, when $n = 2$ and $k = 3$, we show the computation of $H(\Lambda(S^k V)^G, t)$ in the following example.

Example 4.1. Let $n = 2$ and $W = S^3 V$, where $V = \mathbb{C}^2$. Then

$$H(\Lambda(S^3 V), x_1, x_2, t) = (1 + x_1^3 t)(1 + x_2^3 t)(1 + x_1^2 x_2 t)(1 + x_1 x_2^2 t).$$

Using Lemma 3.1 we obtain that

$$M(H(\Lambda(S^3 V)); x_1, x_2, t) = 1 + x_1^3 t + x_1^3 x_2^3 t^2 + x_1^5 x_2 t^2 + x_1^6 x_2^3 t^3 + x_1^6 x_2^6 t^4.$$

By definition, the multiplicity series $M(H(\Lambda(S^3 V)); x_1, x_2, t)$ carries the information about the $\mathrm{GL}(2)$ -structure of $\Lambda(S^3 V)$. Hence, in the notations from the Introduction, we obtain the following decomposition of $\Lambda(S^3 V)$ as a $\mathrm{GL}(2)$ -module

$$\Lambda(S^3 V) \cong \mathbb{C} \oplus S^3 V \oplus (V_{(3,3)}^2 \oplus V_{(5,1)}^2) \oplus V_{(6,3)}^2 \oplus V_{(6,6)}^2.$$

In particular, $\Lambda^2(S^3 V) \cong V_{(3,3)}^2 \oplus V_{(5,1)}^2$, $\Lambda^3(S^3 V) \cong V_{(6,3)}^2$ and $\Lambda^4(S^3 V) \cong V_{(6,6)}^2$.

Next, using Theorem 2.3 we determine the following expressions for $H(\Lambda(S^3 V)^G, t)$:

$$H(\Lambda(S^3 V)^{\mathrm{SL}(2)}, t) = H(\Lambda(S^3 V)^{\mathrm{Sp}(2)}, t) = 1 + t^2 + t^4;$$

$$H(\Lambda(S^3 V)^{\mathrm{O}(2)}, t) = 1 + t^4;$$

$$H(\Lambda(S^3 V)^{\mathrm{SO}(2)}, t) = 1 + 2t^2 + t^4.$$

For computing further examples we have written short programs in Mathematica (respectively for $n = 2, 3, 4$, and 5 variables) and in Tables 1-6 below and in Example 4.2 we give the corresponding expressions for the Hilbert series.

TABLE 1. Hilbert series for $n = 2$

k	$H(\Lambda(S^k V)^{\text{SL}(2)}, t)$
3	$1 + t^2 + t^4$
4	$1 + t^5$
5	$1 + t^2 + t^4 + t^6$
6	$1 + t^3 + t^4 + t^7$
7	$1 + t^2 + t^4 + t^6 + t^8$
8	$1 + t^4 + t^5 + t^9$
9	$1 + t^2 + 2t^4 + 2t^6 + t^8 + t^{10}$
10	$1 + t^3 + t^4 + t^7 + t^8 + t^{11}$
11	$1 + t^2 + 2t^4 + 3t^6 + 2t^8 + t^{10} + t^{12}$
12	$1 + 2t^4 + 2t^5 + 2t^8 + 2t^9 + t^{13}$
13	$1 + t^2 + 2t^4 + 4t^6 + 4t^8 + 2t^{10} + t^{12} + t^{14}$
14	$1 + t^3 + 2t^4 + 4t^7 + 4t^8 + 2t^{11} + t^{12} + t^{15}$
15	$1 + t^2 + 3t^4 + 6t^6 + 7t^8 + 6t^{10} + 3t^{12} + t^{14} + t^{16}$
16	$1 + 2t^4 + 3t^5 + 8t^8 + 8t^9 + 3t^{12} + 2t^{13} + t^{17}$
17	$1 + t^2 + 3t^4 + 8t^6 + 12t^8 + 12t^{10} + 8t^{12} + 3t^{14} + t^{16} + t^{18}$
18	$1 + t^3 + 3t^4 + 10t^7 + 13t^8 + 5t^9 + 5t^{10} + 13t^{11} + 10t^{12} + 3t^{15} + t^{16} + t^{19}$
19	$1 + t^2 + 3t^4 + 10t^6 + 20t^8 + 24t^{10} + 20t^{12} + 10t^{14} + 3t^{16} + t^{18} + t^{20}$
20	$1 + 3t^4 + 4t^5 + t^6 + 4t^7 + 22t^8 + 28t^9 + 13t^{10} + 13t^{11} + 28t^{12} + 22t^{13} + 4t^{14} + t^{15} + 4t^{16} + 3t^{17} + t^{21}$

TABLE 2. Hilbert series for $n = 2$

k	$H(\Lambda(S^k V)^{\text{O}(2)}, t)$	$H(\Lambda(S^k V)^{\text{SO}(2)}, t)$
3	$1 + t^4$	$1 + 2t^2 + t^4$
4	$1 + t + t^4 + t^5$	$1 + t + 2t^2 + 2t^3 + t^4 + t^5$
5	$1 + 3t^4$	$1 + 3t^2 + 3t^4 + t^6$
6	$1 + t + t^3 + 4t^4 + 3t^5$	$1 + t + 3t^2 + 5t^3 + 5t^4 + 3t^5 + t^6 + t^7$
7	$1 + 7t^4 + t^8$	$1 + 4t^2 + 8t^4 + 4t^6 + t^8$
8	$1 + t + 2t^3 + 9t^4 + 9t^5 + 2t^6 + t^8 + t^9$	$1 + t + 4t^2 + 8t^3 + 12t^4 + 12t^5 + 8t^6 + 4t^7 + t^8 + t^9$
9	$1 + 14t^4 + 4t^6 + 5t^8$	$1 + 5t^2 + 18t^4 + 18t^6 + 5t^8 + t^{10}$
10	$1 + t + 4t^3 + 17t^4 + 21t^5 + 11t^6 + 7t^7 + 9t^8 + 5t^9$	$1 + t + 5t^2 + 13t^3 + 24t^4 + 32t^5 + 32t^6 + 24t^7 + 13t^8 + 5t^9 + t^{10} + t^{11}$
11	$1 + 24t^4 + 19t^6 + 24t^8 + t^{12}$	$1 + 6t^2 + 33t^4 + 58t^6 + 33t^8 + 6t^{10} + t^{12}$
12	$1 + t + 6t^3 + 29t^4 + 44t^5 + 37t^6 + 37t^7 + 44t^8 + 29t^9 + 6t^{10} + t^{12} + t^{13}$	$1 + t + 6t^2 + 18t^3 + 43t^4 + 73t^5 + 94t^6 + 94t^7 + 73t^8 + 43t^9 + 18t^{10} + 6t^{11} + t^{12} + t^{13}$
13	$1 + 38t^4 + 58t^6 + 93t^8 + 17t^{10} + 7t^{12}$	$1 + 7t^2 + 55t^4 + 151t^6 + 151t^8 + 55t^{10} + 7t^{12} + t^{14}$
14	$1 + t + 9t^3 + 45t^4 + 81t^5 + 96t^6 + 127t^7 + 162t^8 + 131t^9 + 60t^{10} + 24t^{11} + 16t^{12} + 7t^{13}$	$1 + t + 7t^2 + 25t^3 + 69t^4 + 141t^5 + 227t^6 + 289t^7 + 289t^8 + 227t^9 + 141t^{10} + 69t^{11} + 25t^{12} + 7t^{13} + t^{14} + t^{15}$

TABLE 3. Hilbert series for $n = 3$

k	$H(\Lambda(S^k V)^{\text{SL}(3)}, t)$
3	$1 + t^3 + t^7 + t^{10}$
4	$1 + t^6 + t^9 + t^{15}$
5	$1 + t^3 + t^6 + t^9 + t^{12} + t^{15} + t^{18} + t^{21}$
6	$1 + t^5 + 2t^6 + t^7 + t^8 + 6t^9 + 7t^{10} + 6t^{11} + 8t^{12} + 13t^{13} + 16t^{14} + 13t^{15} + 8t^{16} + 6t^{17} + 7t^{18} + 6t^{19} + t^{20} + t^{21} + 2t^{22} + t^{23} + t^{28}$

TABLE 4. Hilbert series for $n = 3$

k	$H(\Lambda(S^k V)^{\text{O}(3)}, t)$	$H(\Lambda(S^k V)^{\text{SO}(3)}, t)$
3	$1 + 2t^4 + 2t^6 + t^{10}$	$1 + 3t^3 + 2t^4 + 2t^6 + 3t^7 + t^{10}$
4	$1 + t + 3t^4 + 12t^5 + 15t^6 + 8t^7 + 8t^8 + 15t^9 + 12t^{10} + 3t^{11} + t^{14} + t^{15}$	$1 + t + 3t^4 + 12t^5 + 15t^6 + 8t^7 + 8t^8 + 15t^9 + 12t^{10} + 3t^{11} + t^{14} + t^{15}$
5	$1 + 10t^4 + 60t^6 + 158t^8 + 294t^{10} + 210t^{12} + 125t^{14} + 15t^{16} + 7t^{18}$	$1 + 7t^3 + 10t^4 + 15t^5 + 60t^6 + 125t^7 + 158t^8 + 210t^9 + 294t^{10} + 294t^{11} + 210t^{12} + 158t^{13} + 125t^{14} + 60t^{15} + 15t^{16} + 10t^{17} + 7t^{18} + t^{21}$

TABLE 5. Hilbert series for $n = 4$

k	$H(\Lambda(S^k V)^{\text{SL}(4)}, t)$	$H(\Lambda(S^k V)^{\text{Sp}(4)}, t)$
3	$1 + t^4 + t^8 + t^{12} + t^{16} + t^{20}$	$1 + t^2 + 3t^4 + 4t^6 + 7t^8 + 7t^{10} + 7t^{12} + 4t^{14} + 3t^{16} + t^{18} + t^{20}$
4	$1 + t^5 + t^8 + t^9 + t^{10} + 4t^{13} + 3t^{14} + t^{15} + 3t^{16} + 4t^{17} + 4t^{18} + 3t^{19} + t^{20} + 3t^{21} + 4t^{22} + t^{25} + t^{26} + t^{27} + t^{30} + t^{35}$	$1 + t^4 + 5t^5 + 2t^6 + 2t^7 + 22t^8 + 55t^9 + 79t^{10} + 106t^{11} + 207t^{12} + 383t^{13} + 490t^{14} + 528t^{15} + 651t^{16} + 804t^{17} + 804t^{18} + 651t^{19} + 528t^{20} + 490t^{21} + 383t^{22} + 207t^{23} + 106t^{24} + 79t^{25} + 55t^{26} + 22t^{27} + 2t^{28} + 2t^{29} + 5t^{30} + t^{31} + t^{35}$

TABLE 6. Hilbert series for $n = 4$

k	$H(\Lambda(S^k V)^{\text{O}(4)}, t)$	$H(\Lambda(S^k V)^{\text{SO}(4)}, t)$
3	$1 + 2t^4 + 3t^6 + 15t^8 + 8t^{10} + 15t^{12} + t^{14} + 5t^{16}$	$1 + 7t^4 + 4t^6 + 30t^8 + 16t^{10} + 30t^{12} + 4t^{14} + 7t^{16} + t^{20}$

Example 4.2. Hilbert series of $\Lambda(S^3 V)^G$ for $n = 5$:

$$H(\Lambda(S^3 V)^{\text{SL}(5)}, t) = 1 + t^5 + t^{30} + t^{35};$$

$$H(\Lambda(S^3 V)^{\text{O}(5)}, t) = 1 + 2t^4 + 3t^6 + 24t^8 + 79t^{10} + 198t^{12} + 461t^{14} + 592t^{16} + 741t^{18} + 488t^{20} + 359t^{22} + 106t^{24} + 56t^{26} + 3t^{28} + 7t^{30};$$

$$\begin{aligned}
H(\Lambda(S^3V)^{\text{SO}(5)}, t) &= 1 + 2t^4 + 7t^5 + 3t^6 + 3t^7 + 24t^8 + 56t^9 + 79t^{10} + 106t^{11} + \\
&198t^{12} + 359t^{13} + 461t^{14} + 488t^{15} + 592t^{16} + 741t^{17} + 741t^{18} + 592t^{19} + \\
&488t^{20} + 461t^{21} + 359t^{22} + 198t^{23} + 106t^{24} + 79t^{25} + 56t^{26} + 24t^{27} + 3t^{28} + \\
&3t^{29} + 7t^{30} + 2t^{31} + t^{35}.
\end{aligned}$$

Example 4.3. The results in Table 3 show that $\Lambda(S^3V)^{\text{SL}(3)}$ is generated by a pair $\{v, *v\}$, where v is a certain element in $\Lambda^3(S^3V)$ and $*v$ is the Hodge dual of v , i.e. the unique element in $\Lambda^7(S^3V)$ such that

$$v \wedge *v = \langle v, v \rangle \text{vol},$$

where $\langle \cdot, \cdot \rangle$ is the inner product on $\Lambda^3(S^3V)$, which is induced by the standard inner product on $V = \mathbb{C}^3$, and vol is a chosen volume form on S^3V . In this example, we give explicit expressions for v and $*v$.

Let $\{e_1, e_2, e_3\}$ denote the standard basis for $V = \mathbb{C}^3$. Then a basis for $V^{\otimes 3}$ is given by the elements $\{e_i \otimes e_j \otimes e_k\}_{\{1 \leq i, j, k \leq 3\}}$. We consider the standard symmetrization mapping on $V^{\otimes 3}$, defined by

$$\begin{aligned}
S : V^{\otimes 3} &\rightarrow V^{\otimes 3} \\
S(e_i \otimes e_j \otimes e_k) &= \frac{1}{3!} \sum_{\sigma \in S_3} e_{\sigma(i)} \otimes e_{\sigma(j)} \otimes e_{\sigma(k)},
\end{aligned}$$

where S_3 denotes again the symmetric group in 3 variables. Using the standard notation $e_i e_j e_k = S(e_i \otimes e_j \otimes e_k)$, a basis for $S^3V = \text{im}S$ is given by the following ten elements:

$$\begin{aligned}
a_1 &= e_1^3, & a_2 &= e_2^3, & a_3 &= e_3^3, & a_4 &= e_1^2 e_2, & a_5 &= e_1^2 e_3, \\
a_6 &= e_2^2 e_3, & a_7 &= e_1 e_2^2, & a_8 &= e_1 e_3^2, & a_9 &= e_2 e_3^2, & a_{10} &= e_1 e_2 e_3.
\end{aligned}$$

Next, the following map

$$f : V^{\otimes 3p} \rightarrow \Lambda^p(S^3V)$$

given by

$$e_{i_1} \otimes e_{j_1} \otimes e_{k_1} \otimes \cdots \otimes e_{i_p} \otimes e_{j_p} \otimes e_{k_p} \mapsto e_{i_1} e_{j_1} e_{k_1} \wedge \cdots \wedge e_{i_p} e_{j_p} e_{k_p}$$

is a surjective homomorphism of $\text{GL}(3)$ -modules for any $p \geq 1$. Thus, any $\text{SL}(3)$ -invariant in $\Lambda^p(S^3V)$ comes from an $\text{SL}(3)$ -invariant in $V^{\otimes 3p}$. In particular, for $p = 3$, we take v to be the image of the element $\text{Std}_3(e_1, e_2, e_3) \otimes \text{Std}_3(e_1, e_2, e_3) \otimes \text{Std}_3(e_1, e_2, e_3)$, where

$$\text{Std}_3(e_1, e_2, e_3) = \sum_{\sigma \in S_3} \text{sign}(\sigma) e_{\sigma(1)} \otimes e_{\sigma(2)} \otimes e_{\sigma(3)}$$

is the standard polynomial of degree 3. Explicitly, we obtain

$$\begin{aligned}
v &= a_1 \wedge a_2 \wedge a_3 - 3a_3 \wedge a_4 \wedge a_7 - 3a_1 \wedge a_6 \wedge a_9 + 3a_2 \wedge a_5 \wedge a_8 + \\
&6a_7 \wedge a_8 \wedge a_{10} - 6a_4 \wedge a_9 \wedge a_{10} + 6a_5 \wedge a_6 \wedge a_{10} + \\
&3a_5 \wedge a_7 \wedge a_9 + 3a_4 \wedge a_6 \wedge a_8.
\end{aligned}$$

Then, if we choose $\text{vol} = a_1 \wedge a_2 \wedge \cdots \wedge a_{10}$, for $*v$ we obtain

$$\begin{aligned} *v = & a_4 \wedge a_5 \wedge a_6 \wedge a_7 \wedge a_8 \wedge a_9 \wedge a_{10} - \frac{1}{3}a_1 \wedge a_2 \wedge a_5 \wedge a_6 \wedge a_8 \wedge a_9 \wedge a_{10} - \\ & - \frac{1}{3}a_2 \wedge a_3 \wedge a_4 \wedge a_5 \wedge a_7 \wedge a_8 \wedge a_{10} - \frac{1}{3}a_1 \wedge a_3 \wedge a_4 \wedge a_6 \wedge a_7 \wedge a_9 \wedge a_{10} - \\ & - \frac{1}{9}a_1 \wedge a_2 \wedge a_3 \wedge a_4 \wedge a_5 \wedge a_6 \wedge a_9 + \frac{1}{9}a_1 \wedge a_2 \wedge a_3 \wedge a_5 \wedge a_6 \wedge a_7 \wedge a_8 - \\ & - \frac{1}{9}a_1 \wedge a_2 \wedge a_3 \wedge a_4 \wedge a_7 \wedge a_8 \wedge a_9 - \frac{1}{9}a_1 \wedge a_2 \wedge a_3 \wedge a_4 \wedge a_6 \wedge a_8 \wedge a_{10} + \\ & + \frac{1}{9}a_1 \wedge a_2 \wedge a_3 \wedge a_5 \wedge a_7 \wedge a_9 \wedge a_{10}. \end{aligned}$$

Remark 4.4. Similarly, the results in Tables 1, 3, and 5 show that $\Lambda(S^5V)^{\text{SL}(2)}$, $\Lambda(S^6V)^{\text{SL}(2)}$, $\Lambda(S^8V)^{\text{SL}(2)}$, $\Lambda(S^4V)^{\text{SL}(3)}$, and $\Lambda(S^3V)^{\text{SL}(5)}$ are generated each by a pair $\{v, *v\}$ and these pairs are obtained in an analogous way to Example 4.3.

In the next set of examples, we consider $W = \Lambda^3V$. Then, Equation (3) implies that

$$H(\Lambda(\Lambda^3V), x_1, \dots, x_n, t) = \prod_{1 \leq i < j < k \leq n} (1 + x_i x_j x_k t).$$

In Tables 7-9 below, we give the expressions for $H(\Lambda(\Lambda^3V)^G, t)$, for $n = 5, 6$.

TABLE 7. Hilbert series for $n = 5$

$H(\Lambda(\Lambda^3V)^{\text{SL}(5)}, t)$	$H(\Lambda(\Lambda^3V)^{\text{O}(5)}, t)$	$H(\Lambda(\Lambda^3V)^{\text{SO}(5)}, t)$
$1 + t^{10}$	$1 + t^{10}$	$1 + t^3 + t^7 + t^{10}$

TABLE 8. Hilbert series for $n = 6$

$H(\Lambda(\Lambda^3V)^{\text{SL}(6)}, t)$	$H(\Lambda(\Lambda^3V)^{\text{Sp}(6)}, t)$
$1 + t^2 + t^4 + t^6 + t^8 + t^{10} + t^{12} + t^{14} + t^{16} + t^{18} + t^{20}$	$1 + 2t^2 + 4t^4 + 5t^6 + 5t^8 + 5t^{10} + 5t^{12} + 5t^{14} + 4t^{16} + 2t^{18} + t^{20}$

TABLE 9. Hilbert series for $n = 6$

$H(\Lambda(\Lambda^3V)^{\text{O}(6)}, t)$	$H(\Lambda(\Lambda^3V)^{\text{SO}(6)}, t)$
$1 + t^4 + 2t^8 + 3t^{10} + 2t^{12} + t^{16} + t^{20}$	$1 + t^2 + t^4 + 2t^6 + 2t^8 + 8t^{10} + 2t^{12} + 2t^{14} + t^{16} + t^{18} + t^{20}$

Remark 4.5. In a similar way, further results for $H(\Lambda(S^kV)^G, t)$ and $H(\Lambda(\Lambda^kV)^G, t)$ for larger k and for larger n can be obtained. These results are not included in the paper only in order to avoid too complicated expressions for the respective Hilbert series.

5. APPENDIX

Below we provide the source code of the program in Mathematica for determining the Hilbert series $H(\Lambda(S^k V)^G, t)$ for $V = \mathbb{C}^2$ and $G = \text{SL}(2)$, $\text{O}(2)$, and $\text{SO}(2)$. The value of k should be chosen in the beginning of the program.

```

k = 3; H = x - y;
For[i = 0, i <= k, i++, H = Expand[H*(1 + x^(k - i)*y^i*t)]];
M = CoefficientRules[H, {x, y}]; L = {};
For[i = 1, i <= Length[M], i++,
  If[M[[i]][[1]][[1]] > M[[i]][[1]][[2]], L = Append[L, M[[i]]]];
MultSeries = FromCoefficientRules[L, {x, y}];
MultSeries = Expand[MultSeries*1/x];
Print["Multiplicity Series = " MultSeries];
M1 = CoefficientRules[MultSeries, {x, y}]; L1 = {};
For[i = 1, i <= Length[M1], i++,
  If[M1[[i]][[1]][[1]] == M1[[i]][[1]][[2]], L1 = Append[L1, M1[[i]]]];
InvariantsSL = FromCoefficientRules[L1, {x, y}];
InvariantsSL = InvariantsSL /. x -> 1;
InvariantsSL = InvariantsSL /. y -> 1;
Print["Hilbert series of SL-invariants =" InvariantsSL];
L2 = {}; For[i = 1, i <= Length[M1], i++,
  If[EvenQ[M1[[i]][[1]][[1]]] && EvenQ[M1[[i]][[1]][[2]]],
    L2 = Append[L2, M1[[i]]]];
Invariants0 = FromCoefficientRules[L2, {x, y}];
Invariants0 = Invariants0 /. x -> 1;
Invariants0 = Invariants0 /. y -> 1;
Print["Hilbert series of 0-invariants =" Invariants0];
For[i = 1, i <= Length[M1], i++,
  If[OddQ[M1[[i]][[1]][[1]]] && OddQ[M1[[i]][[1]][[2]]],
    L2 = Append[L2, M1[[i]]]];
InvariantsS0 = FromCoefficientRules[L2, {x, y}];
InvariantsS0 = InvariantsS0 /. x -> 1;
InvariantsS0 = InvariantsS0 /. y -> 1;
Print["Hilbert series of S0-invariants =" InvariantsS0]

```

ACKNOWLEDGEMENTS

I am very grateful to Vesselin Drensky for introducing me to the topic of Hilbert series and for his advices and stimulating discussions.

REFERENCES

- [1] F. Benanti, S. Boumova, V. Drensky, G. K. Genov, P. Koev, *Computing with rational symmetric functions and applications to invariant theory and PI-algebras*, Serdica Math. J. **38** (2012), Nos 1-3, 137-188.
- [2] A. Berele, *Applications of Below's theorem to the cocharacter sequence of p.i. algebras*, J. Algebra **298** (2006), 208-214.
- [3] S. Dolce, *On certain modules of covariants in exterior algebras*, Algebr Represent Theor **18** (2015), 1299-1319.
- [4] S. Dolce, *On covariants in exterior algebras for the even special orthogonal group*, Journal of Algebra **462** (2016), 163-180.

- [5] V. Drensky, E. Hristova, *Invariants of symplectic and orthogonal groups acting on $GL(n, \mathbb{C})$ -modules*, arXiv:1707.05893 [math.AC].
- [6] V. Drensky, E. Hristova, *Noncommutative invariant theory of symplectic and orthogonal groups*, *Linear Algebra and its Applications* **581** (2019), 198-213.
- [7] W. Fulton, J. Harris, *Representation Theory. A First Course*. Graduate Texts in Mathematics, **129**, Readings in Mathematics, Springer-Verlag, New York, 1991.
- [8] M. Itoh, *Invariant theory in exterior algebras and Amitsur-Levitzki type theorems*, *Adv. Math.* **288** (2016), 679-701.

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD.
G. BONCHEV STR., BLOCK 8, 1113 SOFIA, BULGARIA

E-mail address: e.hristova@math.bas.bg