

# On Connectivity and Robustness in Random Intersection Graphs

Jun Zhao, *Student Member, IEEE*, Osman Yağan, *Member, IEEE*, and Virgil Gligor, *Senior Member, IEEE*

**Abstract**—Random intersection graphs have received much attention recently and been used in a wide range of applications ranging from key predistribution in wireless sensor networks to modeling social networks. For these graphs, each node is equipped with a set of objects in a random manner, and two nodes have an undirected edge in between if they have at least one object in common. In this paper, we investigate connectivity and robustness in a *general random intersection graph* model. Specifically, we establish sharp asymptotic zero-one laws for  $k$ -connectivity and  $k$ -robustness, as well as the asymptotically exact probability of  $k$ -connectivity, for any positive integer  $k$ . The  $k$ -connectivity property quantifies how resilient is the connectivity of a graph against node or edge failures, while  $k$ -robustness measures the effectiveness of *local-information-based* consensus algorithms (that do not use global graph topology information) in the presence of adversarial nodes. In addition to presenting the results under the general random intersection graph model, we consider two special cases of the general model, a *binomial* random intersection graph and a *uniform* random intersection graph, which both have numerous applications as well. For these two specialized graphs, our results on asymptotically exact probabilities of  $k$ -connectivity and asymptotic zero-one laws for  $k$ -robustness are also novel in the literature.

**Index Terms**—Complex networks, connectivity, consensus, random intersection graphs, robustness.

## I. INTRODUCTION

### A. Background

Since random intersection graphs were introduced by Singer-Cohen [20], different classes of these graphs have received considerable attention [3]–[5], [13], [17]–[19], [24] recently. In these graphs, each node is assigned a set of objects selected by some random mechanism. An undirected edge exists between any two nodes that have at least one object in common. Random intersection graphs have been used in modeling and analyzing real-world networks in a wide variety of applications. Examples include secure wireless sensor networks [5], [13], [24], social networks [17]–[19], classification analysis [14], and cryptanalysis [35]. Several properties such as clustering [4], component evolution [5] and degree distribution [14] have been analyzed for different classes of random intersection graphs.

The graph model in this paper, hereafter referred to as a *general random intersection graph*, represents a generalization of the

random intersection graphs studied by Bloznelis *et al.* [4], [5], and is defined on a node set  $\mathcal{V}_n = \{v_1, v_2, \dots, v_n\}$  as follows. Each node  $v_i$  ( $i = 1, 2, \dots, n$ ) is assigned an object set  $S_i$  from an object pool  $\mathcal{P}_n$  consisting of  $P_n$  distinct objects, where  $P_n$  is a function of  $n$ . Each object set  $S_i$  is constructed from the following two steps: First, the size of  $S_i$ ,  $|S_i|$ , is determined according to some probability distribution  $\mathcal{D}_n : \{1, 2, \dots, P_n\} \rightarrow [0, 1]$ . Of course, we have  $\sum_{x=1}^{P_n} \mathbb{P}[|S_i| = x] = 1$ , with  $\mathbb{P}[A]$  denoting the probability that event  $A$  occurs. Next,  $S_i$  is formed by selecting  $|S_i|$  distinct objects uniformly at random from the object pool  $\mathcal{P}_n$ . In other words, conditioning on  $|S_i| = s_i$ , set  $S_i$  is chosen uniformly among all  $s_i$ -size subsets of  $\mathcal{P}_n$ . This process is repeated independently for all object sets  $S_1, \dots, S_n$ . Finally, an undirected edge is assigned between two nodes if and only if their corresponding object sets have at least one object in common; namely, distinct nodes  $v_i$  and  $v_j$  have an undirected edge in between if and only if  $S_i \cap S_j \neq \emptyset$ . The graph defined through this adjacency notion is denoted by  $G(n, P_n, \mathcal{D}_n)$ .

A specific case of the general model  $G(n, P_n, \mathcal{D}_n)$ , known as the *binomial* random intersection graph, has been widely explored to date [9]–[14]. Under this model, each object set  $S_i$  is constructed by a Bernoulli-like mechanism; i.e., by adding each object to  $S_i$  independently with probability  $p_n$ . Like integer  $P_n$ , probability  $p_n$  is also a function of  $n$ . The term “binomial” accounts for the fact that  $|S_i|$  now follows a binomial distribution with  $P_n$  as the number of trials and  $p_n$  as the success probability in each trial. We denote the binomial random intersection graph by  $G_b(n, P_n, p_n)$ , where the subscript “b” stands for “binomial”.

Another well-known special case of the general model  $G(n, P_n, \mathcal{D}_n)$  is the *uniform* random intersection graph [3], [13], [17], [24]. Under the uniform model, the probability distribution  $\mathcal{D}_n$  concentrates on a single integer  $K_n$ , where  $1 \leq K_n \leq P_n$ ; i.e., for each node  $v_i$ , the object set size  $|S_i|$  equals  $K_n$  with probability 1. Note that  $P_n$  and  $K_n$  are both integer functions of  $n$ . We denote by  $G_u(n, P_n, K_n)$  the uniform random intersection graph, with the subscript “u” meaning “uniform”.

### B. Applications of Random Intersection Graphs

A concrete example for the application of random intersection graphs can be given in the context of secure wireless sensor networks. As explained in detail in numerous other places [5], [17], [22]–[24], [31], the uniform random intersection graph model  $G_u(n, P_n, K_n)$  is induced naturally by the Eschenauer–Gligor random key predistribution scheme [13], which is a typical solution to ensure secure communications in wireless sensor networks. In particular, let the set of  $n$  nodes in graph  $G_u(n, P_n, K_n)$  stand for the  $n$  sensors in the wireless network. Also, let the object pool  $\mathcal{P}_n$  (with size  $P_n$ ) represent the set of cryptographic keys available to the network and let  $K_n$  be the number of keys assigned to each sensor (selected uniformly at random from the

The material in this paper was presented in part at the 2014 IEEE Conference on Decision and Control (CDC), Los Angeles, CA, USA [30].

J. Zhao was, and O. Yağan and V. Gligor are with the Cybersecurity Lab (CyLab) and the Department of Electrical and Computer Engineering, Carnegie Mellon University, Pittsburgh, PA 15213, USA. (Emails: junzhao@alumni.cmu.edu, oyagan@ece.cmu.edu, gligor@cmu.edu). J. Zhao is now with Arizona State University.

This research was supported in part by CyLab, and Department of Electrical & Computer Engineering at Carnegie Mellon University, and also by Grants CCF-0424422 and CNS-0831440 from the National Science Foundation to the Berkeley TRUST STC.

key pool  $\mathcal{P}_n$ ). Then, the edges in  $G_u(n, P_n, K_n)$  represent pairs of sensors that share at least one cryptographic key and thus that can securely communicate over existing wireless links in the Eschenauer–Gligor scheme. In the above application, objects that nodes have are cryptographic keys, so uniform random intersection graphs are also referred to as random key graphs [22]–[24], [31].

In the secure sensor network area, the general random intersection graph model captures the differences that may exist among the number of keys assigned to each sensor. These differences appear for a variety of reasons including (a) the number may vary from sensor to sensor in a heterogeneous sensor network due to differences in the sizes of sensor memories [5]; (b) the number may decrease due to the revocation of compromised nodes and keys [8]; and (c) the number may increase due to the establishment of path keys, where new keys are generated and distributed to participating sensors after deployment [13].

Random intersection graphs can also be used to model social networks, where a node represents an individual, and an object could be an hobby of individuals, a book being read, or a movie being watched, etc. [4], [7], [19], [31]. Then a link between two individuals characterizes a common-interest relation; e.g., two individuals have a connection if they have a common hobby, read the same book, or watch the same movie. In this setting, binomial/uniform/general random intersection graphs represent common-interest networks where the sets of interests that individuals have are constructed in different ways. Specifically, in binomial random intersection graphs, each interest is attached to each person independently with the same probability; in uniform random intersection graphs, all individuals have the same number of interests; and general random intersection graphs provide general possibilities for assigning individuals' interest sets; e.g., without probability or number-of-interest restrictions.

### C. Problem Formulation

We now introduce the graph properties that we are interested in. First,  $k$ -connectivity is formally defined as follows.

**Definition 1 ( $k$ -Connectivity [6])** *A graph is said to be  $k$ -connected if each pair of nodes has at least  $k$  internally node-disjoint path(s) in between, where two paths are internally node-disjoint if except the source and destination, the intermediate nodes are different. Equivalently, by Menger's theorem, a graph is  $k$ -connected if it cannot be disconnected by deleting at most  $(k - 1)$  nodes or edges, where a graph is connected if there exists at least a path of edges between any two nodes.*

Clearly,  $k$ -connectivity quantifies well-established measures of strength. For instance, it captures the resiliency of graphs against node or edge failures. It also captures the resiliency of consensus protocols in the presence of  $h$  adversarial nodes in a graph with node size greater than  $3h$ ; i.e., a necessary and sufficient condition is that the graph is  $(2h + 1)$ -connected [10].

Many graph algorithms rely on sufficient connectivity; e.g., algorithms to achieve consensus [2], [26], [27]. However, these algorithms typically assume that nodes have full knowledge of the graph topology, which is often impractical [26]. To account for the lack of full topology knowledge in the general case, Zhang and Sundaram introduce the notion of *graph robustness* [26], which

has received much attention recently [15], [16], [25], [27], [28]. Formally,  $k$ -robustness is defined as follows.

**Definition 2 ( $k$ -Robustness)** *A graph with a node set  $\mathcal{V}$  is  $k$ -robust if at least one of (a) and (b) below hold for every pair of non-empty, disjoint subsets  $A$  and  $B$  of  $\mathcal{V}$ : (a) there exists at least a node  $v_a \in A$  such that  $v_a$  has no less than  $k$  neighbors inside  $\mathcal{V} \setminus A$ ; and (b) there exists at least a node  $v_b \in B$  such that  $v_b$  has no less than  $k$  neighbors inside  $\mathcal{V} \setminus B$ .*

Zhang and Sundaram [26] show that when nodes have local topology knowledge, consensus can still be reached in a sufficiently robust graph in the presence of adversarial nodes, but not in a sufficiently connected and insufficiently robust graph.

Graph robustness provides a different notion of strength than  $k$ -connectivity. That is, it quantifies the effectiveness and resiliency of local-information-based consensus algorithms in the presence of adversarial nodes. We detail the application of robustness to consensus in the next subsection. Robustness also has broad relevance in graph processes beyond consensus; e.g., robustness plays a key role in information cascades [26].

### D. Application of Robustness to Consensus

To study consensus in a graph, we consider that all nodes are synchronous and the time is divided into different time slots. From one time slot to the next time slot, each node updates its value. Let  $x_i[t]$  denote the value of node  $v_i$  at time slot  $t$  for  $t = 0, 1, \dots$ . We first suppose all nodes are benign. Then consensus means  $\lim_{t \rightarrow \infty} |x_i[t] - x_j[t]| = 0$  for each pair of nodes  $v_i$  and  $v_j$ . The updating process of each node's value is as follows. With  $V_i$  denoting the neighborhood set of each node  $v_i$ , from time slot  $t$  to  $t + 1$ ,  $v_i$  updates its value  $x_i[t]$  to  $x_i[t + 1]$  by incorporating node  $v_j$ 's value  $x_j[t]$  that  $v_j$  sends to  $v_i$ , for  $v_j \in V_i$ ; i.e., there is a function  $f_i(\cdot)$  such that

$$x_i[t + 1] = f_i(\{x_j[t] \mid v_j \in V_i \cup \{v_i\}\}).$$

Now we consider the case where there might exist adversarial nodes, i.e., nodes that are not benign. A node  $v_i$  is said to be benign if it sends  $x_i[t]$  to all of its neighbors and applies  $f_i(\cdot)$  at every time slot  $t$ , and is called adversarial otherwise. In the presence of adversarial nodes, consensus means  $\lim_{t \rightarrow \infty} |x_i[t] - x_j[t]| = 0$  for each pair of *benign* nodes  $v_i$  and  $v_j$ .

Under the adversary model that the total number of adversarial node(s) in the graph is upper bounded by  $h$ , then consensus can be achieved if and only if the graph is  $(2h + 1)$ -connected, given the graph has more than  $3h$  nodes [10]. However, the algorithms often assume that all nodes know the entire network topology [26]. Suppose each node does not know the entire network topology and only knows the number of adversarial nodes in its neighborhood, Zhang and Sundaram [26] show the usefulness of robustness in studying consensus. Specifically, under the adversary model that each benign node has at most  $h$  adversarial node(s) as neighbors, then consensus can be achieved if the graph is  $(2h + 1)$ -robust [26]. With the above, we can use consensus dynamics to motivate the study of both connectivity and robustness, where connectivity (resp., robustness) is applicable to the case where each node knows the global (resp., local) network topology.

### E. Related Work

For connectivity (i.e.,  $k$ -connectivity with  $k = 1$ ) in binomial random intersection graph  $G_b(n, P_n, p_n)$ , Rybarczyk establishes the exact probability [19] and a zero-one law [18], [19]. She further shows a zero-one law for  $k$ -connectivity [18], [19]. Our Theorem 1 provides not only a zero-one law, but also the exact probability to understand  $k$ -connectivity precisely.

For connectivity in uniform random intersection graph  $G_u(n, P_n, K_n)$ , Rybarczyk [17] derives the exact probability and a zero-one law, while Blackburn and Gerke [3], and Yağan and Makowski [24] also obtain zero-one laws. Rybarczyk [18] implicitly shows a zero-one law for  $k$ -connectivity in  $G_u(n, P_n, K_n)$ . Our Theorem 2 also gives a zero-one law. In addition, it gives the exact probability to provide an accurate understanding of  $k$ -connectivity.

For general random intersection graph  $G(n, P_n, \mathcal{D}_n)$ , Godehardt and Jaworski [14] investigate its degree distribution and Bloznelis *et al.* [5] explore its component evolution. Recently, Yağan [23] obtains a zero-one law for connectivity.

Since asymptotic probability results of  $k$ -connectivity in random graphs are often established by first showing the corresponding results for the property of having minimum degree at least  $k$ , and then proving the probability of having minimum degree at least  $k$  yet not being  $k$ -connected converges to zero asymptotically, all the above references on  $k$ -connectivity (resp., connectivity) also establish the corresponding results for the property of having minimum degree at least  $k$  (resp., 1).

To date, there have not been results on  $(k)$ -robustness of random intersection graphs reported by others. As noted in Lemma 3, Zhang and Sundaram [26] present a zero-one law for  $k$ -robustness in an Erdős-Rényi graph.

For random intersection graphs in this paper, two nodes have an edge in between if their object sets share at least one object. A natural variant is to define graphs with edges only between nodes which have at least  $s$  objects in common (instead of just 1) for some positive integer  $s$ . Recent researches [7], [32] investigate  $k$ -connectivity in graphs under this definition.

### F. Contributions and Organization

With the above notions of  $k$ -connectivity and  $k$ -robustness in mind, a natural questions to ask is when will random intersection graphs become  $k$ -connected or  $k$ -robust? We answer this question and summarize the key contributions as follows:

- i) We derive sharp zero-one laws and asymptotically exact probabilities for  $k$ -connectivity in general random intersection graphs.
- ii) We establish sharp zero-one laws for  $k$ -robustness in general random intersection graphs.
- iii) For the two specific instances of the general graph model, a binomial random intersection graph and a uniform random intersection graph, we provide the first results on asymptotically exact probabilities of  $k$ -connectivity and zero-one laws for  $k$ -robustness.

This paper extends the conference version [30] in several ways:

- i) We strengthen the known results on binomial/uniform/general random intersection graphs. Specifically, Theorems 1–6 in

this paper eliminate the condition  $|\alpha_n| = o(\ln n)$  in [30, Theorems 1–6].

- ii) For  $k$ -connectivity of a uniform random intersection graph, we provide a complete proof in Section VI. Note that this result serves as the building block for all other results.
- iii) We enhance numerical experiments to better confirm the theoretical results; see Section III.
- iv) We discuss the parameter conditions of the theorems in detail; see Section II-C.
- v) We compare our results of binomial/uniform/general random intersection graphs with those of Erdős-Rényi graphs; see the last paragraph of Section II.

The rest of the paper is organized as follows. Section II presents the main results as Theorems 1–6. To improve the readability of the paper, we defer the proofs of the theorems to the end of the paper. We provide numerical experiments in Section III. Afterwards, we introduce some auxiliary lemmas in Section IV, before establishing the main results in Sections V, VI and VII. Section VIII details the proofs of the lemmas. Finally, we conclude the paper Section IX.

## II. THE RESULTS

Our main results are presented in Theorems 1–6 below. We defer the proofs of all theorems to Sections V–VII. Throughout the paper,  $k$  is a positive integer and does not scale with  $n$ ; and  $e$  is the base of the natural logarithm function,  $\ln$ . All limits are understood with  $n \rightarrow \infty$ . We use the standard Landau asymptotic notation  $o(\cdot)$ ,  $O(\cdot)$ ,  $\omega(\cdot)$ ,  $\Omega(\cdot)$ ,  $\Theta(\cdot)$  and  $\sim$ ; see [31, Page 2-Footer 1]. In particular, for two positive sequences  $f_n$  and  $g_n$ , the relation  $f_n \sim g_n$  signifies  $\lim_{n \rightarrow \infty} (f_n/g_n) = 1$ . For a random variable  $X$ , the terms  $\mathbb{E}[X]$  and  $\text{Var}[X]$  stand for its expected value and variance, respectively.

As noted in Section I-A, we denote a binomial (resp., uniform) random intersection graph by  $G_b(n, P_n, p_n)$  (resp.,  $G_u(n, P_n, K_n)$ ). Clearly,  $G_b(n, P_n, 0)$  (resp.,  $G_u(n, P_n, 0)$ ) is an empty graph, while  $G_b(n, P_n, 1)$  (resp.,  $G_u(n, P_n, P_n)$ ) being a complete graph is  $k$ -connected for  $n \geq k + 1$  and is  $k$ -robust for  $n \geq 2k$ . Then for each  $n \geq 2k$ , with  $P_n$  fixed and  $p_n$  increasing from 0 to 1 (resp.,  $K_n$  increasing from 0 to  $P_n$ ), the probabilities of  $k$ -connectivity and  $k$ -robustness of  $G_b(n, P_n, p_n)$  (resp.,  $G_u(n, P_n, K_n)$ ) increase from 0 to 1. In addition, for random graphs, results are often obtained in the asymptotic sense since the analysis becomes intractable in the finite regime [11], [12], [20]–[22].

Given the above, it is natural to anticipate that our results are presented in the form of zero-one laws, where a zero-one law means that the probability of a graph having a certain property asymptotically converges to 0 under some conditions and to 1 under some other conditions. Moreover, it is useful to have a complete picture by obtaining the asymptotically exact probability result [21]. For binomial/uniform/general random intersection graphs, we derive asymptotically exact probabilities for  $k$ -connectivity in Theorems 1–3, and zero-one laws for  $k$ -robustness in Theorems 4–6. A future work is to establish asymptotically exact probabilities for  $k$ -robustness.

Noting that for any graph/network,  $k$ -connectivity implies that the minimum node degree is at least  $k$  [11], we often present

results for the property of minimum node degree being at least  $k$  together with  $k$ -connectivity results.

#### A. Asymptotically Exact Probabilities for $k$ -Connectivity and the Property of Minimum Node Degree Being at Least $k$

##### 1) $k$ -Connectivity and Minimum Node Degree in Binomial Random Intersection Graphs:

For a binomial random intersection graph, Theorem 1 below shows asymptotically exact probabilities for  $k$ -connectivity and the property of minimum node degree being at least  $k$ .

**Theorem 1** For a binomial random intersection graph  $G_b(n, P_n, p_n)$ , with a sequence  $\alpha_n$  for all  $n$  defined through

$$p_n^2 P_n = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}, \quad (1)$$

if  $P_n = \omega(n(\ln n)^5)$ ,

$$\lim_{n \rightarrow \infty} \mathbb{P}[\text{Graph } G_b(n, P_n, p_n) \text{ is } k\text{-connected.}]$$

$$= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \end{cases} \quad (2a)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty), \end{cases} \quad (2b)$$

and

$$\lim_{n \rightarrow \infty} \mathbb{P}[\text{Graph } G_b(n, P_n, p_n) \text{ has a minimum node degree at least } k.]$$

$$= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \end{cases} \quad (3a)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty). \end{cases} \quad (3b)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty). \end{cases} \quad (3c)$$

**Remark 1** As we will explain in Section V-A within the proof of Theorem 1, for (2a) (2b) (3a) (3b) (i.e., the zero-one laws), the condition  $P_n = \omega(n(\ln n)^5)$  can be weakened as  $P_n = \Omega(n(\ln n)^5)$ , while we enforce  $P_n = \omega(n(\ln n)^5)$  for (2c) (3c).

##### 2) $k$ -Connectivity and Minimum Node Degree in Uniform Random Intersection Graphs:

For a uniform random intersection graph, Theorem 2 below gives asymptotically exact probabilities for  $k$ -connectivity and the property of minimum node degree being at least  $k$ .

**Theorem 2** For a uniform random intersection graph  $G_u(n, P_n, K_n)$ , with a sequence  $\alpha_n$  for all  $n$  defined through

$$\frac{K_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}, \quad (4)$$

if  $K_n = \Omega(\sqrt{\ln n})$ , then

$$\lim_{n \rightarrow \infty} \mathbb{P}[\text{Graph } G_u(n, P_n, K_n) \text{ is } k\text{-connected.}]$$

$$= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \end{cases} \quad (5a)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty), \end{cases} \quad (5b)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty), \end{cases} \quad (5c)$$

and

$$\lim_{n \rightarrow \infty} \mathbb{P}[\text{Graph } G_u(n, P_n, K_n) \text{ has a minimum node degree at least } k.]$$

$$= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \end{cases} \quad (6a)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty). \end{cases} \quad (6b)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty). \end{cases} \quad (6c)$$

##### 3) $k$ -Connectivity and Minimum Node Degree in General Random Intersection Graphs:

For a general random intersection graph, Theorem 3 below provides asymptotically exact probabilities for  $k$ -connectivity and the property of minimum node degree being at least  $k$ .

**Theorem 3** Consider a general random intersection graph  $G(n, P_n, \mathcal{D}_n)$ . Let  $X_n$  be a random variable following probability distribution  $\mathcal{D}_n$ . With a sequence  $\alpha_n$  for all  $n$  defined through

$$\frac{\{\mathbb{E}[X_n]\}^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}, \quad (7)$$

if  $\mathbb{E}[X_n] = \Omega(\sqrt{\ln n})$  and  $\text{Var}[X_n] = o\left(\frac{\{\mathbb{E}[X_n]\}^2}{n(\ln n)^2}\right)$ , then

$$\lim_{n \rightarrow \infty} \mathbb{P}[\text{Graph } G(n, P_n, \mathcal{D}_n) \text{ is } k\text{-connected.}]$$

$$= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \end{cases} \quad (8a)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty), \end{cases} \quad (8b)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty), \end{cases} \quad (8c)$$

and

$$\lim_{n \rightarrow \infty} \mathbb{P}[\text{Graph } G(n, P_n, \mathcal{D}_n) \text{ has a minimum node degree at least } k.]$$

$$= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \end{cases} \quad (9a)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty). \end{cases} \quad (9b)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty). \end{cases} \quad (9c)$$

#### B. Asymptotic Zero-One Laws for $k$ -Robustness

##### 1) $k$ -Robustness in Binomial Random Intersection Graphs:

Theorem 4 below gives an asymptotic zero-one law for  $k$ -robustness in a binomial random intersection graph.

**Theorem 4** For a binomial random intersection graph  $G_b(n, P_n, p_n)$ , with a sequence  $\alpha_n$  for all  $n$  defined through

$$p_n^2 P_n = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}, \quad (10)$$

if  $P_n = \Omega(n(\ln n)^5)$ , then

$$\lim_{n \rightarrow \infty} \mathbb{P}[\text{Graph } G_b(n, P_n, p_n) \text{ is } k\text{-robust.}]$$

$$= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty. \end{cases} \quad (11a)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty. \end{cases} \quad (11b)$$

##### 2) $k$ -Robustness in Uniform Random Intersection Graphs:

Theorem 5 below presents an asymptotic zero-one law for  $k$ -robustness in a uniform random intersection graph.

**Theorem 5** For a uniform random intersection graph  $G_u(n, P_n, K_n)$ , with a sequence  $\alpha_n$  for all  $n$  defined through

$$\frac{K_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}, \quad (12)$$

if  $K_n = \Omega((\ln n)^3)$ , then

$$\lim_{n \rightarrow \infty} \mathbb{P}[\text{Graph } G_u(n, P_n, K_n) \text{ is } k\text{-robust.}]$$

$$= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty. \end{cases} \quad (13a)$$

$$= \begin{cases} 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty. \end{cases} \quad (13b)$$

##### 3) $k$ -Robustness in General Random Intersection Graphs:

Theorem 6 as follows provides an asymptotic zero-one law for  $k$ -robustness in a general random intersection graph.

**Theorem 6** Consider a general random intersection graph  $G(n, P_n, \mathcal{D}_n)$ . Let  $X_n$  be a random variable following probability distribution  $\mathcal{D}_n$ . With a sequence  $\alpha_n$  for all  $n$  defined through

$$\frac{\{\mathbb{E}[X_n]\}^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}, \quad (14)$$

if  $\mathbb{E}[X_n] = \Omega((\ln n)^3)$  and  $\text{Var}[X_n] = o\left(\frac{\{\mathbb{E}[X_n]\}^2}{n(\ln n)^2}\right)$ , then

$$\lim_{n \rightarrow \infty} \mathbb{P}[\text{Graph } G(n, P_n, \mathcal{D}_n) \text{ is } k\text{-robust.}] = \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty. \end{cases}$$

In view of Theorems 1–6, for each binomial/uniform/general random intersection graph, its  $k$ -connectivity,  $k$ -robustness and the property of minimum node degree being at least  $k$  asymptotically obey the same zero–one laws. Moreover, these zero–one laws are all *sharp* since  $|\alpha_n|$  can be much smaller compared to  $\ln n$ ; e.g., even  $\alpha_n = \pm \cdot \ln \ln \dots \ln n$  satisfies  $\lim_{n \rightarrow \infty} \alpha_n = \pm \infty$ .

We compare our results of random intersection graphs with those of Erdős–Rényi graphs below. From [9, Section 1.1],  $p_n^2 P_n$  in the scaling conditions (1) and (10) of Theorems 1 and 4 is an asymptotics of the edge probability in a binomial random intersection graph  $G_b(n, P_n, p_n)$ . Also, by [4, Lemma 1],  $\frac{K_n^2}{P_n}$  in the scaling conditions (4) and (12) of Theorems 2 and 5 (resp.,  $\frac{\{\mathbb{E}[X_n]\}^2}{P_n}$  in the scaling conditions (7) and (14) of Theorems 3 and 6) is an asymptotics of the edge probability in a uniform random intersection graph  $G_u(n, P_n, K_n)$  (resp., a general random intersection graph  $G(n, P_n, \mathcal{D}_n)$ ). Then comparing Theorems 1–3 with Lemma 2, and comparing Theorems 4–6 with Lemma 3, we conclude binomial/uniform/general random intersection graphs under certain parameter conditions<sup>1</sup> exhibit the same behavior with Erdős–Rényi graphs in the sense that for each of (i)  $k$ -connectivity, (ii) the property of minimum node degree being at least  $k$ , and (iii)  $k$ -robustness, a common point for the transition from a zero-law to a one-law occurs when the edge probability equals  $\frac{\ln n + (k-1) \ln \ln n}{n}$ . The term  $\alpha_n$  in Equations (1) (resp., (4) and (7)), or Equations (10) (resp., (12) and (14)) measures how much  $p_n^2 P_n$  (resp.,  $\frac{K_n^2}{P_n}$  and  $\frac{\{\mathbb{E}[X_n]\}^2}{P_n}$ ) is away from the critical value  $\frac{\ln n + (k-1) \ln \ln n}{n}$ .

### C. A Discussion of Parameter Conditions

Note that we impose conditions on the parameters in the theorems; e.g.,  $P_n = \omega(n(\ln n)^5)$  in Theorem 1, and  $K_n = \Omega(\sqrt{\ln n})$  in Theorem 2. These conditions are enforced to have the proofs get through and are not that conservative as explained below. We take a binomial random intersection graph as an example and note that Theorem 1 for  $k$ -connectivity in a binomial random intersection graph does not hold if the condition  $P_n = \omega(n(\ln n)^5)$  in Theorem 1 is replaced by  $P_n = n^\tau$  for a positive constant  $\tau < 1$ . Specifically, we use [18, Theorem 4] and [18, Conjecture 1] confirmed by later work [19] to have the following claim:

**Claim 1** Under  $P_n = n^\tau$  for a positive constant  $\tau < 1$ , with a sequence  $\gamma_n$  for all  $n$  defined through

$$p_n P_n = \ln n + \gamma_n, \quad (15)$$

<sup>1</sup>Under other parameter conditions, the conclusion may not hold as in the case of binomial random intersection graphs shown by Rybarczyk [18], [19].

then

$$\lim_{n \rightarrow \infty} \mathbb{P}[G_b(n, P_n, p_n) \text{ is } k\text{-connected.}] = \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \gamma_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \gamma_n = \infty. \end{cases}$$

Note that different from (1), the scaling condition (15) above does not depend on  $k$ .

Claim 1 has  $P_n = n^\tau$  for a positive constant  $\tau < 1$ , while Theorem 1 has  $P_n = \omega(n(\ln n)^5)$ . We let  $\delta$  denote an arbitrary constant with  $\tau < \delta < \frac{\tau+1}{2}$  below. Claim 1 shows that the probability of  $G_b(n, n^\tau, n^{-\delta})$  (i.e.,  $G_b(n, P_n, p_n)$  with  $P_n = n^\tau$  and  $p_n = n^{-\delta}$ ) being  $k$ -connected asymptotically converges to 0 since  $\gamma_n$  specified by (15) satisfies

$$\gamma_n = p_n P_n - \ln n = n^{\tau-\delta} - \ln n \rightarrow -\infty, \text{ as } n \rightarrow \infty.$$

In contrast, Theorem 1 with  $P_n = \omega(n(\ln n)^5)$  replaced by  $P_n = n^\tau$  for a positive constant  $\tau < 1$  presents that the probability of  $G_b(n, n^\tau, n^{-\delta})$  (i.e.,  $G_b(n, P_n, p_n)$  with  $P_n = n^\tau$  and  $p_n = n^{-\delta}$ ) being  $k$ -connected asymptotically approaches to 1 because  $\alpha_n$  defined by (1) satisfies

$$\begin{aligned} \alpha_n &= n p_n^2 P_n - [\ln n + (k-1) \ln \ln n] \\ &= n^{1+\tau-2\delta} - [\ln n + (k-1) \ln \ln n] \rightarrow \infty, \text{ as } n \rightarrow \infty. \end{aligned}$$

Hence, Claim 1 shows that if the condition  $P_n = \omega(n(\ln n)^5)$  of Theorem 1 is replaced by  $P_n = n^\tau$  for a positive constant  $\tau < 1$ , we will not obtain the  $k$ -connectivity result of Theorem 1. A future work is to investigate the intermediate range  $\omega(n^\tau) = P_n = O(n(\ln n)^5)$ .

We have discussed the parameter conditions for binomial random intersection graphs. It is unclear whether  $K_n = \Omega(\sqrt{\ln n})$  in Theorem 2 and  $K_n = \Omega((\ln n)^3)$  in Theorem 5 for uniform random intersection graphs can be weakened since these conditions are also often enforced in related work [5], [24], [31]. Moreover, these conditions are applicable to secure sensor networks since it has been shown that  $K_n$  is at least on the order of  $\ln n$  to have reasonable connectivity and resiliency [21], [33], [34]. For a general random intersection graph, Yağan [23] recently obtains a zero–one law for connectivity and shows in [23, Section 3.3] that Theorem 3 with  $\text{Var}[X_n] = o\left(\frac{\{\mathbb{E}[X_n]\}^2}{n(\ln n)^2}\right)$  replaced by a broader condition does not hold.

To conclude, the parameter conditions in our theorems are not that conservative.

## III. NUMERICAL EXPERIMENTS

We present numerical experiments in the non-asymptotic regime to confirm our theoretical results.

Figure 1 depicts the probability that a binomial random intersection graph  $G_b(n, P, p)$  has  $k$ -connectivity or  $k$ -robustness, for  $k = 1, 2$ . Similarly, Figure 2 illustrates the probability of  $k$ -connectivity or  $k$ -robustness for  $k = 2, 3$  in a uniform random intersection graph  $G_u(n, P, K)$ . In all set of experiments, we fix the number of nodes at  $n = 2000$  and the object pool size  $P = 20000$ . For each pair  $(n, P, p)$  (resp.,  $(n, P, K)$ ), we generate 1000 independent samples of  $G_b(n, P, p)$  (resp.,  $G_u(n, P, K)$ ) and count the number of times that the obtained graphs are  $k$ -connected or  $k$ -robust. Then the counts divided by 1000 become the corresponding empirical probabilities. As illustrated in Figures 1 and 2, there is an evident transition in the




Fig. 1: A plot of the empirical probabilities that a binomial random intersection graph  $G_b(n, P, p)$  has  $k$ -connectivity or  $k$ -robustness as a function of  $p$ , with  $n = 2000$ ,  $P = 20000$  and  $k = 2, 6$ .




Fig. 2: A plot of the empirical probabilities that a uniform random intersection graph  $G_u(n, P, K)$  has  $k$ -connectivity or  $k$ -robustness as a function of  $K$ , with  $n = 2000$ ,  $P = 20000$  and  $k = 3, 4$ .

probabilities of  $k$ -connectivity and  $k$ -robustness. Also, for each  $k$ , the curves of  $k$ -connectivity and  $k$ -robustness are close to each other. Furthermore, the vertical lines in Figure 1 specify  $p$  such that  $p^2 P$  equals  $\frac{\ln n + (k-1) \ln \ln n}{n}$ , while the vertical lines in Figure 2 specify  $K$  such that  $\frac{K^2}{P}$  is closest to  $\frac{\ln n + (k-1) \ln \ln n}{n}$  (since  $K$  and  $P$  are both integers, there might not exist  $K$  satisfying  $\frac{K^2}{P} = \frac{\ln n + (k-1) \ln \ln n}{n}$ ).

The vertical lines in Figure 1 are at  $4.4 \times 10^{-4}$  and  $4.9 \times 10^{-4}$  because under  $n = 2000$  and  $P = 20000$ ,  $\sqrt{\frac{\ln n + (k-1) \ln \ln n}{nP}}$  is  $\sqrt{\frac{\ln 2000}{2000 \times 20000}} \approx 4.4 \times 10^{-4}$  for  $k = 1$  and is  $\sqrt{\frac{\ln 2000 + \ln \ln 2000}{2000 \times 20000}} \approx 4.9 \times 10^{-4}$  for  $k = 2$ . The vertical lines in Figure 2 are at 10 and 11 because under  $n = 2000$  and  $P = 20000$ ,  $\arg \min_K \left| \frac{K^2}{P} - \frac{\ln n + (k-1) \ln \ln n}{n} \right|$  equals 10 for  $k = 2$  from  $\frac{\ln 2000 + \ln \ln 2000}{2000} \approx 0.00481$ ,  $\frac{9^2}{20000} \approx 0.00405$  and  $\frac{10^2}{20000} \approx 0.005$ , and equals 11 for  $k = 3$  from  $\frac{\ln 2000 + 2 \ln \ln 2000}{2000} \approx 0.00583$ ,  $\frac{10^2}{20000} \approx 0.005$  and  $\frac{11^2}{20000} \approx 0.00605$ .

#### IV. AUXILIARY LEMMAS

We present lemmas that are used in proving the theorems.

##### A. Relationships between $k$ -Robustness, $k$ -Connectivity, and Minimum Node Degree

Lemma 1 below, taken from [26, Lemma 1], provides relationships between  $k$ -robustness,  $k$ -connectivity, and minimum node degree.

**Lemma 1 ([26, Lemma 1])** *For any graph/network,  $k$ -robustness implies  $k$ -connectivity, which further implies that the minimum node degree is at least  $k$ .*

To prove that  $k$ -robustness implies  $k$ -connectivity, [26, Lemma 1] shows that a graph  $G$  that is not  $k$ -connected is also not  $k$ -robust. The idea is that for  $G$  being not  $k$ -connected, there exists a set of  $k - 1$  nodes, whose deletion gives two disjoint subgraphs with node sets  $V_a$  and  $V_b$ , respectively. Then in graph  $G$ , each node in  $V_a$  has less than  $k$  neighbors outside of  $V_a$ , and each node in  $V_b$  has less than  $k$  neighbors outside of  $V_b$ , so  $G$  is not  $k$ -robust. Then it follows that  $k$ -robustness implies  $k$ -connectivity. In addition, it is clear that  $k$ -connectivity implies that the minimum node degree is at least  $k$ .

Lemma 1 is used in the proofs of Theorems 4 and 5.

##### B. Results of Erdős-Rényi Graphs

Lemma 2 below by Erdős and Rényi [12] investigates  $k$ -connectivity and minimum node degree in Erdős-Rényi graphs. An Erdős-Rényi graph  $G(n, \hat{p}_n)$  [11] is defined on a set of  $n$  nodes such that any two nodes have an edge in between independently with probability  $\hat{p}_n$ .

**Lemma 2 (Erdős and Rényi [12])** *For an Erdős-Rényi graph  $G(n, \hat{p}_n)$ , with a sequence  $\alpha_n$  for all  $n$  through*

$$\hat{p}_n = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n},$$

*then it holds that*

$$\begin{aligned} & \lim_{n \rightarrow \infty} \mathbb{P}[G(n, \hat{p}_n) \text{ is } k\text{-connected.}] \\ &= \lim_{n \rightarrow \infty} \mathbb{P}[G(n, \hat{p}_n) \text{ has a minimum node degree at least } k.] \\ &= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty. \end{cases} \end{aligned}$$

Lemma 2 is used in the comparison of random intersection graphs and Erdős-Rényi graphs.

Lemma 3 below presents the result on  $k$ -robustness in Erdős-Rényi graphs.

**Lemma 3** *For an Erdős-Rényi graph  $G(n, \hat{p}_n)$ , with a sequence  $\alpha_n$  for all  $n$  through*

$$\hat{p}_n = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n},$$

*then it holds that*

$$\lim_{n \rightarrow \infty} \mathbb{P}[G(n, \hat{p}_n) \text{ is } k\text{-robust.}] = \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty. \end{cases} \quad (16)$$

Lemma 3 is applied to Section VII-A for proving Theorem 4. Lemma 3 is also used in the comparison of random intersection graphs and Erdős-Rényi graphs.

To prove Lemma 3, we note the following three facts. (a) The desired result (16) with  $|\alpha_n| = o(\ln \ln n)$  is demonstrated in [26, Theorem 3]. (b) By [18, Facts 3 and 7], for any monotone

increasing graph property  $\mathcal{I}$ , the probability that graph  $G(n, \hat{p}_n)$  has property  $\mathcal{I}$  is non-decreasing as  $\hat{p}_n$  increases, where a graph property is called monotone increasing if it holds under the addition of edges. (c)  $k$ -Robustness is a monotone increasing graph property according to [15, Lemma 3]. In view of (a) (b) and (c) above, we obtain Lemma 3.

### C. Lemmas for Graph Coupling

We present several lemmas for graph coupling below. Formally, a coupling [18], [19], [30] of two random graphs  $G_1$  and  $G_2$  means a probability space on which random graphs  $G'_1$  and  $G'_2$  are defined such that  $G'_1$  and  $G'_2$  have the same distributions as  $G_1$  and  $G_2$ , respectively. If  $G'_1$  is a spanning subgraph (resp., spanning supergraph) of  $G'_2$ , we say that under the graph coupling,  $G_1$  is a spanning subgraph (resp., spanning supergraph) of  $G_2$ , where a spanning subgraph (resp., spanning supergraph) is a subgraph (resp., supergraph) that has the same node set with the original graph.

Following Rybarczyk's notation [18], we write

$$G_1 \preceq G_2 \quad (\text{resp., } G_1 \preceq_{1-o(1)} G_2) \quad (17)$$

if there exists a coupling under which  $G_1$  is a spanning subgraph of  $G_2$  with probability 1 (resp.,  $1 - o(1)$ ). We write

$$G_2 \succeq G_1 \quad (\text{resp., } G_2 \succeq_{1-o(1)} G_1) \quad (18)$$

if there exists a coupling under which  $G_2$  is a spanning supergraph of  $G_1$  with probability 1 (resp.,  $1 - o(1)$ ). According to the definitions above,  $G_1 \preceq G_2$  and  $G_2 \succeq G_1$  are equivalent, while  $G_1 \preceq_{1-o(1)} G_2$  and  $G_2 \succeq_{1-o(1)} G_1$  are equivalent.

In view that  $k$ -connectivity and  $k$ -robustness are monotone increasing graph properties [12], [15], it is natural to obtain that under  $G_1 \preceq G_2$  or  $G_1 \preceq_{1-o(1)} G_2$ , if  $G_1$  is  $k$ -connected (resp.  $k$ -robust) with a probability at least  $1 - o(1)$ , then  $G_2$  is also  $k$ -connected (resp.  $k$ -robust) with a probability at least  $1 - o(1)$ . This result is formally presented in Lemma 4 below given by Rybarczyk [18]. Lemma 4 considers any monotone increasing graph property for generality.

**Lemma 4 (Rybarczyk [18])** *For two random graphs  $G_1$  and  $G_2$ , the following results hold for any monotone increasing graph property  $\mathcal{I}$ .*

(a) *If  $G_1 \preceq G_2$ , then*

$$\mathbb{P}[G_2 \text{ has } \mathcal{I}] \geq \mathbb{P}[G_1 \text{ has } \mathcal{I}].$$

(b) *If  $G_1 \preceq_{1-o(1)} G_2$ , then*

$$\mathbb{P}[G_2 \text{ has } \mathcal{I}] \geq \mathbb{P}[G_1 \text{ has } \mathcal{I}] - o(1).$$

Lemma 4 is used in many places of this paper. We then present Lemmas 5–8. Except Lemma 7 which is from [5, Lemma 4], the proofs of other lemmas are deferred to Section VIII.

1) *Coupling between general random intersection graphs and uniform random intersection graphs:*

**Lemma 5** *Let  $X_n$  be a random variable with probability distribution  $\mathcal{D}_n$ . If  $\text{Var}[X_n] = o\left(\frac{\{\mathbb{E}[X_n]\}^2}{n(\ln n)^2}\right)$ , then there exists  $\epsilon_n = o\left(\frac{1}{\ln n}\right)$  such that*

$$\begin{aligned} G_u(n, P_n, (1-\epsilon_n)\mathbb{E}[X_n]) &\preceq_{1-o(1)} G(n, P_n, \mathcal{D}_n) \\ &\preceq_{1-o(1)} G_u(n, P_n, (1+\epsilon_n)\mathbb{E}[X_n]). \end{aligned} \quad (19)$$

Lemma 5 is shown in Section VIII-A, and is used to prove Theorems 3 and 6.

2) *Coupling between binomial random intersection graphs and Erdős–Rényi graphs:*

**Lemma 6** *If  $p_n = O\left(\frac{1}{n \ln n}\right)$  and  $p_n^2 P_n = O\left(\frac{1}{\ln n}\right)$ , then there exists  $\hat{p}_n = p_n^2 P_n \cdot [1 - O\left(\frac{1}{\ln n}\right)]$  such that*

$$G(n, \hat{p}_n) \preceq_{1-o(1)} G_b(n, P_n, p_n). \quad (20)$$

Lemma 6 is shown in Section VIII-B, and is used to prove Theorem 4.

3) *Coupling between binomial random intersection graphs and uniform random intersection graphs:*

**Lemma 7 ([5, Lemma 4])** *If  $p_n P_n = \omega(\ln n)$ , and for all  $n$  sufficiently large,*

$$\begin{aligned} K_{n,-} &\leq p_n P_n - \sqrt{3(p_n P_n + \ln n) \ln n}, \\ K_{n,+} &\geq p_n P_n + \sqrt{3(p_n P_n + \ln n) \ln n}, \end{aligned}$$

then

$$\begin{aligned} G_u(n, P_n, K_{n,-}) &\preceq_{1-o(1)} G_b(n, P_n, p_n) \\ &\preceq_{1-o(1)} G_u(n, P_n, K_{n,+}). \end{aligned}$$

Lemma 7 is used in the proof of Theorem 1.

**Lemma 8** *If  $K_n = \omega(\ln n)$  and  $p_n = \frac{K_n}{P_n} \left(1 - \sqrt{\frac{3 \ln n}{K_n}}\right)$ , then*

$$G_u(n, P_n, K_n) \succeq_{1-o(1)} G_b(n, P_n, p_n).$$

Lemma 8 is established in Section VIII-C, and is used to prove Theorem 5.

We will use each of Lemmas 5–8 along with Lemma 4. For simplicity, we just use Lemma 5 as an example to explain its implication with Lemma 4. From property (b) of Lemma 4 and result (19) of Lemma 5, we obtain for any monotone increasing graph property  $\mathcal{I}$  that

$$\begin{aligned} &\mathbb{P}[\text{Graph } G_u(n, P_n, (1-\epsilon_n)\mathbb{E}[X_n]) \text{ has } \mathcal{I}] - o(1) \\ &\leq \mathbb{P}[\text{Graph } G(n, P_n, \mathcal{D}_n) \text{ has } \mathcal{I}] \\ &\leq \mathbb{P}[\text{Graph } G_u(n, P_n, (1+\epsilon_n)\mathbb{E}[X_n]) \text{ has } \mathcal{I}] + o(1). \end{aligned}$$

## V. ESTABLISHING THEOREMS 1 AND 3

Theorems 1–3 describe results on  $k$ -connectivity for binomial/uniform/general random intersection graphs. We prove Theorems 1 and 3 in this section, and present the proof of Theorem 2 separately as Section VI next due to the length of the proof.

We briefly explain the idea of proving Theorems 1 and 3 from Theorem 2 below. First, we demonstrate Theorem 1 from Theorem 2 using the coupling between binomial random intersection graphs and uniform random intersection graphs given by Lemma 7 of Section IV-C3. Second, we establish Theorem 3 from Theorem 2 using the coupling between general random intersection graphs and uniform random intersection graphs given by Lemma 5 of Section IV-C1.

### A. The Proof of Theorem 1

As explained in Appendix A, we can introduce an extra condition  $|\alpha_n| = o(\ln n)$  in proving Theorem 1. Then from Theorem 2, Lemmas 4 and 7, and the fact that both  $k$ -connectivity and the property of minimum node degree being at least  $k$  are monotone increasing graph properties, the proof of Theorem 1 is completed once we show that with  $K_{n,\pm}$  given by

$$K_{n,\pm} = p_n P_n \pm \sqrt{3(p_n P_n + \ln n) \ln n}, \quad (21)$$

under conditions of Theorem 1 and  $|\alpha_n| = o(\ln n)$ , we have  $K_{n,\pm} = \Omega(\sqrt{\ln n})$  and with  $\alpha_{n,\pm}$  defined by

$$\frac{K_{n,\pm}^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_{n,\pm}}{n}, \quad (22)$$

then

$$\alpha_{n,\pm} = \alpha_n \pm o(1). \quad (23)$$

From conditions (1) and  $|\alpha_n| = o(\ln n)$ , it is clear that

$$p_n^2 P_n \sim \frac{\ln n}{n}. \quad (24)$$

Substituting (24) and condition  $P_n = \omega(n(\ln n)^5)$  into (21), we obtain

$$p_n P_n = \sqrt{p_n^2 P_n \cdot P_n} = \omega\left(\frac{\ln n}{n} \cdot n(\ln n)^5\right) = \omega((\ln n)^3),$$

$$K_{n,\pm} = \omega((\ln n)^3) = \Omega(\sqrt{\ln n}), \quad (25)$$

and

$$\begin{aligned} \frac{K_{n,\pm}^2}{P_n} &= p_n^2 P_n \cdot \left[1 \pm \sqrt{3 \left(1 + \frac{\ln n}{p_n P_n}\right) \frac{\ln n}{p_n P_n}}\right] \\ &= p_n^2 P_n \cdot \left[1 \pm o\left(\frac{1}{\ln n}\right)\right]. \end{aligned} \quad (26)$$

Then from (1) (22) and (26), we obtain (23). As explained before, with (22) (23) and (25), Theorem 1 is proved from Theorem 2 and Lemmas 4 and 7.

Finally, as noted in Remark 1 after Theorem 1, to prove zero-one laws (2a) (2b) (3a) (3b) but not (2c) (3c) in Theorem 1, condition  $P_n = \omega(n(\ln n)^5)$  can be weakened as  $P_n = \Omega(n(\ln n)^5)$ . This is seen by the argument that under  $P_n = \Omega(n(\ln n)^5)$ ,  $K_{n,\pm} = \Omega(\sqrt{\ln n})$  still holds and (23) is weakened as  $\alpha_{n,\pm} = \alpha_n \pm O(1)$ , so we still have zero-one laws (2a) (2b) (3a) (3b).

### B. The Proof of Theorem 3

Given Lemmas 4 and 5 and the fact that both  $k$ -connectivity and the property of minimum node degree being at least  $k$  are monotone increasing graph properties, we will show Theorem 3 once proving for any  $\epsilon_n = o(\frac{1}{\ln n})$  that

$$\begin{aligned} &\lim_{n \rightarrow \infty} \mathbb{P}[G_u(n, P_n, (1 \pm \epsilon_n)\mathbb{E}[X_n]) \text{ is } k\text{-connected.}] \\ &= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty), \end{cases} \end{aligned} \quad (27)$$

and

$$\begin{aligned} &\lim_{n \rightarrow \infty} \mathbb{P}\left[\text{Graph } G_u(n, P_n, (1 \pm \epsilon_n)\mathbb{E}[X_n]) \text{ has a} \right. \\ &\quad \left. \text{minimum node degree at least } k. \right] \\ &= \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ e^{-\frac{e^{-\alpha^*}}{(k-1)!}}, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty). \end{cases} \end{aligned} \quad (28)$$

Under  $\mathbb{E}[X_n] = \Omega(\sqrt{\ln n})$  and  $\epsilon_n = o(\frac{1}{\ln n})$ , it follows that  $(1 \pm \epsilon_n)\mathbb{E}[X_n] = \Omega(\sqrt{\ln n})$ . From Theorem 2, we will have (27) and (28) once we prove that sequences  $\gamma_n^+$  and  $\gamma_n^-$  defined through

$$\frac{\{(1 \pm \epsilon_n)\mathbb{E}[X_n]\}^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \gamma_n^\pm}{n} \quad (29)$$

satisfy

$$\lim_{n \rightarrow \infty} \gamma_n^\pm = \begin{cases} -\infty, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ \infty, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty, \\ \alpha^*, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty). \end{cases} \quad (30)$$

Now we establish (30). From (7) (29) and  $\epsilon_n = o(\frac{1}{\ln n})$ , it follows that

$$\begin{aligned} \gamma_n^\pm &= n \cdot \frac{\{(1 \pm \epsilon_n)\mathbb{E}[X_n]\}^2}{P_n} - [\ln n + (k-1) \ln \ln n] \\ &= (1 \pm \epsilon_n)^2 [\ln n + (k-1) \ln \ln n + \alpha_n] \\ &\quad - [\ln n + (k-1) \ln \ln n] \\ &= \alpha_n + \epsilon_n(\epsilon_n \pm 2) [\ln n + (k-1) \ln \ln n + \alpha_n] \\ &= \alpha_n \pm \left[ o\left(\frac{\alpha_n}{\ln n}\right) + o(1) \right], \end{aligned} \quad (31)$$

where the last step uses  $\epsilon_n = o(\frac{1}{\ln n})$ . Then (31) clearly implies (30). Therefore, as mentioned above, we establish (27) (28) and finally Theorem 3.

## VI. THE PROOF OF THEOREM 2

As explained in Appendix B, we can introduce an extra condition  $|\alpha_n| = o(\ln n)$  in proving Theorem 2. Then since a necessary condition for a graph to be  $k$ -connected is that the minimum degree is at least  $k$ , (6a) implies (5a), and we have

$$\begin{aligned} &\mathbb{P}[\text{Graph } G_u(n, P_n, K_n) \text{ is } k\text{-connected.}] \\ &= \mathbb{P}\left[\text{Graph } G_u(n, P_n, K_n) \text{ has a} \right. \\ &\quad \left. \text{minimum node degree at least } k. \right] \\ &\quad - \mathbb{P}\left[G_u(n, P_n, K_n) \text{ has a minimum degree} \right. \\ &\quad \left. \text{at least } k, \text{ but is not } k\text{-connected.} \right]. \end{aligned} \quad (32)$$

From (32), we know that (5b) (resp., (5c)) will follow from the combination of Lemma 9 below and (6b) (resp., (6c)), where Lemma 9 uses the extra condition  $|\alpha_n| = o(\ln n)$  explained above. Also as mentioned before, (6a) implies (5a). Therefore, the proof of Theorem 2 will be completed once we demonstrate (6a) (6b) (6c) and Lemma 9, where we also use the extra condition  $|\alpha_n| = o(\ln n)$  in proving (6a) (6b) (6c). We let  $e^{-\infty} = 0$  and  $e^\infty = \infty$ , so  $e^{-\frac{e^{-\lim_{n \rightarrow \infty} \alpha_n}}{(k-1)!}}$  equals 0 if  $\lim_{n \rightarrow \infty} \alpha_n = -\infty$ , 1 if  $\lim_{n \rightarrow \infty} \alpha_n = \infty$  and  $e^{-\frac{e^{-\alpha^*}}{(k-1)!}}$  if  $\lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty)$ . Then (6a) (6b) (6c) under  $|\alpha_n| = o(\ln n)$  can be compactly presented by Lemma 10 below. Hence, the proof of Theorem 2 finally reduces to proving Lemmas 9 and 10.

**Lemma 9** For a uniform random intersection graph  $G_u(n, P_n, K_n)$  under  $K_n = \Omega(\sqrt{\ln n})$  and  $\frac{K_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}$ , where  $\lim_{n \rightarrow \infty} \alpha_n$  exists and  $|\alpha_n| = o(\ln n)$ , it follows that

$$\lim_{n \rightarrow \infty} \mathbb{P}\left[G_u(n, P_n, K_n) \text{ has a minimum degree} \right. \\ \left. \text{at least } k, \text{ but is not } k\text{-connected.} \right] = 0. \quad (33)$$

**Lemma 10** For a uniform random intersection graph  $G_u(n, P_n, K_n)$  under  $K_n = \Omega(\sqrt{\ln n})$  and  $\frac{K_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}$ , where  $\lim_{n \rightarrow \infty} \alpha_n$  exists and  $|\alpha_n| = o(\ln n)$ , it follows that

$$\begin{aligned} &\lim_{n \rightarrow \infty} \mathbb{P}[G_u(n, P_n, K_n) \text{ has a minimum degree at least } k.] \\ &= e^{-\frac{e^{-\lim_{n \rightarrow \infty} \alpha_n}}{(k-1)!}}. \end{aligned}$$



To prove Lemma 9, we use the following Lemma 11 on  $G_u(n, P_n, K_n) \cap G(n, \hat{p}_n)$ , where  $G(n, \hat{p}_n)$  is an Erdős-Rényi graph with  $n$  nodes and edge probability  $\hat{p}_n$ , and the intersection of two graphs  $G_A$  and  $G_B$  defined on the same node set is constructed on the node set with the edge set being the intersection of the edge sets of  $G_A$  and  $G_B$ .

**Lemma 11 (Our work [31, Propositions 3 and 4])** *For a graph  $G_u(n, P_n, K_n) \cap G(n, \hat{p}_n)$  under  $P_n = \Omega(n)$ ,  $\frac{K_n}{P_n} = o(1)$  and  $\frac{K_n^2}{P_n} \cdot \hat{p}_n = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}$ , where  $\lim_{n \rightarrow \infty} \alpha_n$  exists and  $|\alpha_n| = o(\ln n)$ , it follows that*

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[ G_u(n, P_n, K_n) \cap G(n, \hat{p}_n) \text{ has a minimum degree at least } k, \text{ but is not } k\text{-connected.} \right] = 0. \quad (34)$$

Lemma 11 is from our work [31, Propositions 3 and 4]. Setting  $\hat{p}_n = 1$ , we have  $G_u(n, P_n, K_n) \cap G(n, \hat{p}_n) = G_u(n, P_n, K_n)$  and obtain results on  $G_u(n, P_n, K_n)$  from Lemma 11:

*For  $G_u(n, P_n, K_n)$  under  $P_n = \Omega(n)$ ,  $\frac{K_n}{P_n} = o(1)$  and  $\frac{K_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n}$ , where  $\lim_{n \rightarrow \infty} \alpha_n$  exists and  $|\alpha_n| = o(\ln n)$ , result (33) holds.*

Then clearly, Lemma 9 will be proved once we show conditions in Lemma 9 imply  $P_n = \Omega(n)$  and  $\frac{K_n}{P_n} = o(1)$ . From conditions in Lemma 9, we have  $K_n = \Omega(\sqrt{\ln n})$  and  $\frac{K_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n} \sim \frac{\ln n}{n}$  given  $|\alpha_n| = o(\ln n)$ . Then we further get  $P_n = K_n^2 / \frac{K_n^2}{P_n} = \Omega(\ln n / \frac{\ln n}{n}) = \Omega(n)$  and  $\frac{K_n}{P_n} = \frac{K_n^2}{P_n} / K_n = O(\frac{\ln n}{n} / \sqrt{\ln n}) = o(1)$ . Hence, as mentioned above, Lemma 9 is established.

Now we prove Lemma 10. We let  $q_n$  be the edge probability in a uniform random intersection graph  $G_u(n, P_n, K_n)$ ; i.e., two nodes in  $G_u(n, P_n, K_n)$  have an edge in between with probability  $q_n$ . Under conditions of Lemma 10, given  $|\alpha_n| = o(\ln n)$ , we have

$$\frac{K_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \alpha_n}{n} \sim \frac{\ln n}{n}. \quad (35)$$

Hence, from [31, Lemma 8-Property (a)], it follows that

$$q_n = \frac{K_n^2}{P_n} \left[ 1 \pm O\left(\frac{K_n^2}{P_n}\right) \right] \sim \frac{\ln n}{n}. \quad (36)$$

Then, by [29, Section 3], Lemma 10 will follow once we show Lemma 12 below, where  $\mathcal{V}_n = \{v_1, v_2, \dots, v_n\}$  is the set of nodes in graph  $G_u(n, P_n, K_n)$ .

**Lemma 12** *For a uniform random intersection graph  $G_u(n, P_n, K_n)$  under  $K_n = \Omega(\sqrt{\ln n})$  and  $q_n \sim \frac{\ln n}{n}$ , it follows for integers  $m \geq 1$  and  $h \geq 0$  that*

$$\begin{aligned} & \mathbb{P}[\text{Nodes } v_1, v_2, \dots, v_m \text{ have degree } h] \\ & \sim (h!)^{-m} (nq_n)^{hm} e^{-mnq_n}. \end{aligned} \quad (37)$$

The rest of this section is devoted to proving Lemma 12.

In a uniform random intersection graph  $G_u(n, P_n, K_n)$ , recalling that  $\mathcal{V}_n = \{v_1, v_2, \dots, v_n\}$  is the set of nodes, we let  $S_i$  be the set of  $K_n$  distinct objects assigned to node  $v_i \in \mathcal{V}_n$ . We further define  $\mathcal{V}_m$  as  $\{v_1, v_2, \dots, v_m\}$  and  $\overline{\mathcal{V}_m}$  as  $\mathcal{V}_n \setminus \mathcal{V}_m$ . Among nodes in  $\overline{\mathcal{V}_m}$ , we denote by  $N_i$  the set of nodes neighboring to  $v_i$  for  $i = 1, 2, \dots, m$ . We denote  $N_i \cap N_j$  by  $N_{ij}$ , and  $S_i \cap S_j$  by  $S_{ij}$ .

We have the following two observations:

- i) If node  $v_i$  has degree  $h$ , then  $|N_i| \leq h$ , where the equal sign holds if and only if  $v_i$  is directly connected to none of nodes in  $\mathcal{V}_m \setminus \{v_i\}$ ; i.e., if and only if  $\bigcap_{j \in \{1, 2, \dots, m\} \setminus \{i\}} (S_{ij} = \emptyset)$

happens.

- ii) If  $|N_i| \leq h$  for any  $i = 1, 2, \dots, m$ , then

$$\left| \bigcup_{1 \leq i \leq m} N_i \right| \leq \sum_{1 \leq i \leq m} |N_i| \leq hm, \quad (38)$$

where the two equal signs in (38) both hold if and only if

$$\left( \bigcap_{1 \leq i < j \leq m} (N_{ij} = \emptyset) \right) \cap \left( \bigcap_{1 \leq i \leq m} (|N_i| = h) \right). \quad (39)$$

From i) and ii) above, if nodes  $v_1, v_2, \dots, v_m$  have degree  $h$ , we have either of the following two cases:

- (a) Any two of  $v_1, v_2, \dots, v_m$  have no edge in between (namely,  $\bigcap_{1 \leq i < j \leq m} (S_{ij} = \emptyset)$ ); and event (39) happens.
- (b)  $|\bigcup_{1 \leq i \leq m} N_i| \leq hm - 1$ .

In addition, if case (a) happens, then nodes  $v_1, v_2, \dots, v_m$  have degree  $h$ . However, if case (b) occurs, there is no such conclusion. With  $P_a$  (resp.,  $P_b$ ) denoting the probability of case (a) (resp., case (b)), we obtain

$$P_a \leq \mathbb{P}[\text{Nodes } v_1, v_2, \dots, v_m \text{ have degree } h] \leq P_a + P_b,$$

where

$$P_a = \mathbb{P} \left[ \left( \bigcap_{1 \leq i < j \leq m} (S_{ij} = \emptyset) \right) \cap \left( \bigcap_{1 \leq i < j \leq m} (N_{ij} = \emptyset) \right) \cap \left( \bigcap_{1 \leq i \leq m} (|N_i| = h) \right) \right],$$

and

$$P_b = \mathbb{P} \left[ \left| \bigcup_{1 \leq i \leq m} N_i \right| \leq hm - 1 \right].$$

Hence, (37) holds after we prove the following (40) and (41):

$$P_b = o((nq_n)^{hm} e^{-mnq_n}). \quad (40)$$

and

$$P_a \sim (h!)^{-m} (nq_n)^{hm} e^{-mnq_n} \cdot [1 + o(1)], \quad (41)$$

We will prove (40) and (41) below. We let  $\mathbb{S}_m$  denote the tuple  $(S_1, S_2, \dots, S_m)$ . The expression “ $|\mathbb{S}_m = \mathbb{S}_m^*$ ” means “given  $S_1 = S_1^*, S_2 = S_2^*, \dots, S_m = S_m^*$ ”, where  $\mathbb{S}_m^* = (S_1^*, S_2^*, \dots, S_m^*)$  with  $S_1^*, S_2^*, \dots, S_m^*$  being arbitrary  $K_n$ -size subsets of the object pool  $\mathcal{P}_n$  (see Page 1 in the graph definition for the meaning of  $\mathcal{P}_n$ ). Note that  $S_{ij}^* := S_i^* \cap S_j^*$ . For two different nodes  $v$  and  $w$  in  $G_u(n, P_n, K_n)$ , we use  $v \leftrightarrow w$  to denote the event that there is an edge between  $v$  and  $w$ ; i.e., the symbol “ $\leftrightarrow$ ” means “is directly connected with”.

#### A. The Proof of (40)

Let  $w$  be an arbitrary node in  $\overline{\mathcal{V}_m}$ . The event  $w \in \bigcup_{1 \leq i \leq m} N_i$  means  $w \leftrightarrow$  at least one of nodes in  $\mathcal{V}_m$ , which for different  $w$  would be independent given  $\mathbb{S}_m = \mathbb{S}_m^*$ . Then we have

$$\mathbb{P} \left[ \left| \bigcup_{1 \leq i \leq m} N_i \right| = t \mid \mathbb{S}_m = \mathbb{S}_m^* \right] \quad (42)$$

$$\begin{aligned} &= \frac{(n-m)!}{t!(n-m-t)!} \\ &\times \{ \mathbb{P}[w \leftrightarrow \text{at least one of nodes in } \mathcal{V}_m \mid \mathbb{S}_m = \mathbb{S}_m^*] \}^t \\ &\times \{ \mathbb{P}[w \leftrightarrow \text{none of nodes in } \mathcal{V}_m \mid \mathbb{S}_m = \mathbb{S}_m^*] \}^{n-m-t}. \end{aligned} \quad (43)$$

By the union bound, it holds that

$$\begin{aligned} & \mathbb{P}[w \leftrightarrow \text{at least one of nodes in } \mathcal{V}_m | \mathbb{S}_m = \mathbb{S}_m^*] \\ & \leq \sum_{1 \leq i \leq m} \mathbb{P}[w \leftrightarrow v_i | \mathbb{S}_m = \mathbb{S}_m^*] = mq_n, \end{aligned} \quad (44)$$

which yields

$$\mathbb{P}[w \leftrightarrow \text{none of nodes in } \mathcal{V}_m | \mathbb{S}_m = \mathbb{S}_m^*] \geq 1 - mq_n. \quad (45)$$

In addition, we find

$$\begin{aligned} & \mathbb{P}[w \leftrightarrow \text{none of nodes in } \mathcal{V}_m | \mathbb{S}_m = \mathbb{S}_m^*] \\ & = \frac{\binom{P_n - |\bigcup_{1 \leq i \leq m} S_i^*|}{K_n}}{\binom{P_n}{K_n}} \\ & \leq (1 - q_n)^{K_n - |\bigcup_{1 \leq i \leq m} S_i^*|} \quad (\text{by [22, Lemma 5.1]}) \\ & \leq e^{-K_n^{-1} q_n |\bigcup_{1 \leq i \leq m} S_i^*|} \quad (\text{by } 1 + x \leq e^x \text{ for any real } x). \end{aligned} \quad (46)$$

We will prove

$$\begin{aligned} & \sum_{\mathbb{S}_m^*} \left\{ \mathbb{P}[\mathbb{S}_m = \mathbb{S}_m^*] \times \left\{ \mathbb{P} \left[ w \leftrightarrow \text{none of nodes in } \mathcal{V}_m \mid \mathbb{S}_m = \mathbb{S}_m^* \right] \right\}^{n-m-hm} \right\} \\ & \leq e^{-mnq_n} \cdot [1 + o(1)]. \end{aligned} \quad (47)$$

$$(48)$$

From (43) (44) and (48), we derive

$$\begin{aligned} P_b &= \mathbb{P} \left[ \left| \bigcup_{1 \leq i \leq m} N_i \right| \leq hm - 1 \right] \\ &= \sum_{t=0}^{hm-1} \sum_{\mathbb{S}_m^*} \left\{ \mathbb{P}[\mathbb{S}_m = \mathbb{S}_m^*] \cdot (42) \right\} \\ &\leq \sum_{t=0}^{hm-1} \left[ n^t \cdot (mq_n)^t \cdot (47) \right] \\ &\leq (nq_n)^{hm} e^{-mnq_n} \cdot [1 + o(1)] \cdot m^{hm} \sum_{t=0}^{hm-1} (mnq_n)^{t-hm}. \end{aligned} \quad (49)$$

Applying (36) to (49), we obtain (40). Hence, we complete the proof of (40) once showing (48), whose proof is detailed below.

From (36) (45) and (46), we have

$$\begin{aligned} (47) &\leq (1 - mq_n)^{-m-hm} \\ &\quad \times \sum_{\mathbb{S}_m^*} \left\{ \mathbb{P}[\mathbb{S}_m = \mathbb{S}_m^*] \cdot e^{-K_n^{-1} nq_n |\bigcup_{1 \leq i \leq m} S_i^*|} \right\} \\ &\leq [1 + o(1)] \cdot \sum_{\mathbb{S}_m^*} \left\{ \mathbb{P}[\mathbb{S}_m = \mathbb{S}_m^*] \cdot e^{-K_n^{-1} nq_n |\bigcup_{1 \leq i \leq m} S_i^*|} \right\}, \end{aligned} \quad (50)$$

so (48) holds once we demonstrate

$$\begin{aligned} & \sum_{\mathbb{S}_m^*} \left\{ \mathbb{P}[\mathbb{S}_m = \mathbb{S}_m^*] \cdot e^{-K_n^{-1} nq_n |\bigcup_{1 \leq i \leq m} S_i^*|} \right\} \\ & \leq e^{-mnq_n} \cdot [1 + o(1)]. \end{aligned} \quad (51)$$

We denote the left hand side of (51) by  $Z_{m,n}$ . Dividing  $\mathbb{S}_m^*$  into

two parts  $\mathbb{S}_{m-1}^*$  and  $S_m^*$ , we derive

$$\begin{aligned} Z_{m,n} &= \sum_{\substack{\mathbb{S}_{m-1}^* \\ S_m^*}} \left\{ \mathbb{P}[(\mathbb{S}_{m-1} = \mathbb{S}_{m-1}^*) \cap (S_m = S_m^*)] \right. \\ &\quad \times \left. e^{-K_n^{-1} nq_n |\bigcup_{1 \leq i \leq m} S_i^*|} \right\} \\ &= \sum_{\mathbb{S}_{m-1}^*} \mathbb{P}[\mathbb{S}_{m-1} = \mathbb{S}_{m-1}^*] \left\{ e^{-K_n^{-1} nq_n |\bigcup_{1 \leq i \leq m-1} S_i^*|} \right. \\ &\quad \times \left. \sum_{S_m^*} \mathbb{P}[S_m = S_m^*] e^{-K_n^{-1} nq_n |S_m^* \setminus \bigcup_{1 \leq i \leq m-1} S_i^*|} \right\}, \end{aligned} \quad (52)$$

where

$$\begin{aligned} & \sum_{S_m^*} \mathbb{P}[S_m = S_m^*] e^{-K_n^{-1} nq_n |S_m^* \setminus \bigcup_{1 \leq i \leq m-1} S_i^*|} \\ & \leq e^{-nq_n} \sum_{S_m^*} \mathbb{P}[S_m = S_m^*] e^{K_n^{-1} nq_n |S_m^* \cap (\bigcup_{i=1}^{m-1} S_i^*)|} \\ & = e^{-nq_n} \sum_{r=0}^{K_n} \mathbb{P} \left[ \left| S_m \cap \left( \bigcup_{i=1}^{m-1} S_i^* \right) \right| = r \right] e^{K_n^{-1} nq_n r}. \end{aligned} \quad (53)$$

Denoting  $|\bigcup_{i=1}^{m-1} S_i^*|$  by  $v$ , then for  $r$  satisfying the conditions  $0 \leq r \leq |S_m^*| = K_n$  and  $S_m^* \cup (\bigcup_{i=1}^{m-1} S_i^*) = K_n + v - r \leq P_n$  (i.e., for  $r \in [\max\{0, K_n + v - P_n\}, K_n]$ ), we obtain

$$\mathbb{P} \left[ \left| S_m \cap \left( \bigcup_{i=1}^{m-1} S_i^* \right) \right| = r \right] = \binom{v}{r} \binom{P_n - v}{K_n - r} / \binom{P_n}{K_n}, \quad (54)$$

which together with  $K_n \leq v \leq mK_n$  yields

$$\begin{aligned} & \text{L.H.S. of (54)} \\ & \leq \frac{(mK_n)^r}{r!} \cdot \frac{(P_n - K_n)^{K_n - r}}{(K_n - r)!} \cdot \frac{K_n!}{(P_n - K_n)^{K_n}} \\ & \leq \frac{1}{r!} \left( \frac{mK_n^2}{P_n - K_n} \right)^r \quad \text{for } r \in [\max\{0, K_n + v - P_n\}, K_n]. \end{aligned} \quad (55)$$

Also, it is clear that

$$\text{L.H.S. of (54)} = 0 \quad \text{for } r \notin [\max\{0, K_n + v - P_n\}, K_n]. \quad (56)$$

Applying (55) and (56) to (53), we establish

$$\begin{aligned} & \sum_{\mathbb{S}_m^*} \mathbb{P}[S_m = S_m^*] e^{-K_n^{-1} nq_n |S_m^* \setminus \bigcup_{1 \leq i \leq m-1} S_i^*|} \\ & \leq e^{-nq_n} \sum_{r=0}^{K_n} \frac{1}{r!} \left( \frac{mK_n^2}{P_n - K_n} \right)^r e^{K_n^{-1} nq_n r} \\ & \leq e^{-nq_n} \cdot e^{\frac{mK_n^2}{P_n - K_n} \cdot e^{K_n^{-1} nq_n}}. \end{aligned} \quad (57)$$

From  $K_n = \Omega(\sqrt{\ln n})$  and (35) (i.e.,  $\frac{K_n^2}{P_n} \sim \frac{\ln n}{n}$ ), we have  $P_n = \omega(K_n)$  and further

$$\frac{mK_n^2}{P_n - K_n} \sim \frac{mK_n^2}{P_n} \sim \frac{m \ln n}{n}. \quad (58)$$

For an arbitrary  $\epsilon > 0$ , from (36), we obtain  $q_n \leq (1 + \epsilon) \frac{\ln n}{n}$  for all  $n$  sufficiently large, which with  $K_n = \Omega(\sqrt{\ln n}) \geq 2$  yields that for all  $n$  sufficiently large,

$$e^{K_n^{-1} nq_n} \leq e^{\frac{1}{2}(1+\epsilon) \ln n} = n^{\frac{1}{2}(1+\epsilon)}. \quad (59)$$

From (58) and (59), we get

$$\frac{mK_n^2}{P_n - K_n} \cdot e^{K_n^{-1} nq_n} \leq m \ln n \cdot n^{\frac{1}{2}(\epsilon-1)} \cdot [1 + o(1)]. \quad (60)$$

Since  $\epsilon > 0$  is arbitrary, it follows from (60) that for arbitrary  $0 < c < \frac{1}{2}$ , then for all  $n$  sufficiently large, it is clear that

$$\frac{mK_n^2}{P_n - K_n} \cdot e^{K_n^{-1}nq_n} \leq n^{-c}. \quad (61)$$

Using (61) in (57), for all  $n$  sufficiently large, it follows that

$$\sum_{S_m^*} \mathbb{P}[S_m = S_m^*] e^{-K_n^{-1}nq_n | S_m^* \cup \bigcup_{1 \leq i \leq m-1} S_i^*|} \leq e^{-nq_n} \cdot e^{n^{-c}}. \quad (62)$$

Substituting (62) into (52), for all  $n$  sufficiently large, we obtain

$$\begin{aligned} Z_{m,n} &\leq e^{-nq_n} \cdot e^{n^{-c}} \cdot \sum_{S_{m-1}^*} \mathbb{P}[S_{m-1} = S_{m-1}^*] e^{-K_n^{-1}nq_n | \bigcup_{1 \leq i \leq m-1} S_i^*|} \\ &\leq e^{-nq_n} \cdot e^{n^{-c}} \cdot Z_{m-1,n}. \end{aligned} \quad (63)$$

We then evaluate  $Z_{2,n}$ . By (51), it holds that

$$\begin{aligned} Z_{2,n} &= \sum_{S_1^*} \sum_{S_2^*} \left\{ \mathbb{P}[(S_1 = S_1^*) \cap (S_2 = S_2^*)] \cdot e^{-K_n^{-1}nq_n | S_1^* \cup S_2^*|} \right\} \\ &= \sum_{S_1^*} \mathbb{P}[S_1 = S_1^*] \sum_{S_2^*} \mathbb{P}[S_2 = S_2^*] e^{-K_n^{-1}nq_n | S_1^* \cup S_2^*|.} \end{aligned} \quad (64)$$

Setting  $m = 2$  in (62), for all  $n$  sufficiently large, we derive

$$\sum_{S_2^*} \mathbb{P}[S_2 = S_2^*] e^{-K_n^{-1}nq_n | S_2^* \setminus S_1^*|} \leq e^{-nq_n} \cdot e^{n^{-c}}.$$

Then for all  $n$  sufficiently large, it follows that

$$\begin{aligned} \sum_{S_2^*} \mathbb{P}[S_2 = S_2^*] e^{-K_n^{-1}nq_n | S_1^* \cup S_2^*|} &= e^{-nq_n} \sum_{S_2^*} \mathbb{P}[S_2 = S_2^*] e^{-K_n^{-1}nq_n | S_2^* \setminus S_1^*|} \\ &\leq e^{-2nq_n} \cdot e^{n^{-c}}. \end{aligned} \quad (65)$$

From (64) and (65), for all  $n$  sufficiently large, we obtain

$$\begin{aligned} Z_{m,n} &\leq (e^{-nq_n} \cdot e^{n^{-c}})^{m-2} \cdot Z_{2,n} \\ &\leq (e^{-nq_n} \cdot e^{n^{-c}})^{m-2} \cdot e^{-2nq_n} \cdot e^{n^{-c}} \\ &\leq e^{-mnq_n} \cdot e^{(m-1)n^{-c}}. \end{aligned} \quad (66)$$

Letting  $n \rightarrow \infty$ , we finally establish

$$Z_{m,n} \leq e^{-mnq_n} \cdot [1 + o(1)];$$

i.e., (51) is proved. As explained, (48) and then (40) follow.

### B. The Proof of (41)

Again let  $w$  be an arbitrary node in  $\overline{\mathcal{V}_m}$ . We have

$$\mathbb{P}\left[\left(\bigcap_{1 \leq i < j \leq m} (N_{ij} = \emptyset)\right) \cap \left(\bigcap_{1 \leq i \leq m} (|N_i| = h)\right) \middle| \mathbb{S}_m = \mathbb{S}_m^*\right] \quad (67)$$

$$\begin{aligned} &= \frac{(n-m)!}{(h!)^m (n-m-hm)!} \\ &\quad \times \prod_{1 \leq i \leq m} \left( \left\{ \mathbb{P}\left[ \begin{array}{l} w \leftrightarrow v_i, \\ \text{but } w \leftrightarrow \text{none of} \\ \text{nodes in } \mathcal{V}_m \setminus \{v_i\} \end{array} \middle| \mathbb{S}_m = \mathbb{S}_m^* \right] \right\}^h \right) \\ &\quad \times \left\{ \mathbb{P}[w \leftrightarrow \text{none of nodes in } \mathcal{V}_m \mid \mathbb{S}_m = \mathbb{S}_m^*] \right\}^{n-m-hm} \end{aligned} \quad (68)$$

and

$$P_a = \sum_{\mathbb{S}_m^* : \bigcap_{1 \leq i < j \leq m} (S_{ij}^* = \emptyset)} \left\{ \mathbb{P}[\mathbb{S}_m = \mathbb{S}_m^*] \cdot (67) \right\}, \quad (69)$$

where  $S_{ij}^* := S_i^* \cap S_j^*$ .

For  $i = 1, 2, \dots, m$ , under  $\mathbb{S}_m^* : \bigcap_{1 \leq i < j \leq m} (S_{ij}^* = \emptyset)$ , it follows that

$$\begin{aligned} &\mathbb{P}[w \leftrightarrow v_i, \text{ but none of nodes in } \mathcal{V}_m \setminus \{v_i\} \mid \mathbb{S}_m = \mathbb{S}_m^*] \\ &\geq \mathbb{P}[w \leftrightarrow v_i \mid \mathbb{S}_m = \mathbb{S}_m^*] \\ &\quad - \sum_{\substack{1 \leq j \leq m \\ j \neq i}} \mathbb{P}[w \leftrightarrow \text{both } v_i \text{ and } v_j \mid \mathbb{S}_m = \mathbb{S}_m^*], \end{aligned} \quad (70)$$

where we note

$$\mathbb{P}[w \leftrightarrow v_i \mid \mathbb{S}_m = \mathbb{S}_m^*] = q_n, \quad (71)$$

and

$$\begin{aligned} &\mathbb{P}[w \leftrightarrow \text{both } v_i \text{ and } v_j \mid \mathbb{S}_m = \mathbb{S}_m^*] \\ &= \mathbb{P}[w \leftrightarrow v_i \mid \mathbb{S}_m = \mathbb{S}_m^*] + \mathbb{P}[w \leftrightarrow v_j \mid \mathbb{S}_m = \mathbb{S}_m^*] \\ &\quad - \mathbb{P}[(w \leftrightarrow v_i) \cup (w \leftrightarrow v_j) \mid \mathbb{S}_m = \mathbb{S}_m^*] \\ &= q_n + q_n - \binom{P_n - 2K_n}{K_n} / \binom{P_n}{K_n} \end{aligned} \quad (72)$$

given  $\mathbb{S}_m^* : \bigcap_{1 \leq i < j \leq m} (S_{ij}^* = \emptyset)$ . From [22, Lemma 5.1], we get  $\binom{P_n - 2K_n}{K_n} / \binom{P_n}{K_n} \leq (1 - q_n)^2$ , which with (71) and (72) are used in (70) to derive

$$\begin{aligned} &\mathbb{P}[w \leftrightarrow v_i, \text{ but none of nodes in } \mathcal{V}_m \setminus \{v_i\} \mid \mathbb{S}_m = \mathbb{S}_m^*] \\ &\geq q_n - (m-1) \cdot 2q_n^2. \end{aligned} \quad (73)$$

Substituting (45) and (73) to (68), and then from (69), we obtain

$$\begin{aligned} P_a &\geq \frac{(n-m-hm)^{hm}}{(h!)^m} \cdot [q_n - 2(m-1)q_n^2]^{hm} \\ &\quad \times (1 - mq_n)^{n-m-hm} \sum_{\mathbb{S}_m^* : \bigcap_{1 \leq i < j \leq m} (S_{ij}^* = \emptyset)} \mathbb{P}[\mathbb{S}_m = \mathbb{S}_m^*]. \end{aligned}$$

Then from (36), it further hold that

$$\begin{aligned} P_a &\geq \frac{n^{hm}}{(h!)^m} \cdot (q_n)^{hm} \cdot e^{-mnq_n} \\ &\quad \times [1 - o(1)] \cdot \mathbb{P}\left[\bigcap_{1 \leq i < j \leq m} (S_{ij} = \emptyset)\right]. \end{aligned} \quad (74)$$

From (46), under  $\mathbb{S}_m^* : \bigcap_{1 \leq i < j \leq m} (S_{ij}^* = \emptyset)$ , it holds that

$$\mathbb{P}[w \leftrightarrow \text{none of nodes in } \mathcal{V}_m \mid \mathbb{S}_m = \mathbb{S}_m^*] \leq e^{-mq_n}. \quad (75)$$

For each  $i = 1, 2, \dots, m$ , we have

$$\begin{aligned} &\mathbb{P}[w \leftrightarrow v_i, \text{ but } w \leftrightarrow \text{none of nodes in } \mathcal{V}_m \setminus \{v_i\} \mid \mathbb{S}_m = \mathbb{S}_m^*] \\ &\leq \mathbb{P}[w \leftrightarrow v_i \mid \mathbb{S}_m = \mathbb{S}_m^*] = q_n. \end{aligned} \quad (76)$$

Substituting (76) and (75) to (68), and then from (69), we obtain

$$\begin{aligned} P_a &\leq \frac{n^{hm}}{(h!)^m} \cdot (q_n)^{hm} \cdot e^{-mnq_n} \cdot \sum_{\mathbb{S}_m^* : \bigcap_{1 \leq i < j \leq m} (S_{ij}^* = \emptyset)} \mathbb{P}[\mathbb{S}_m = \mathbb{S}_m^*] \\ &= \frac{n^{hm}}{(h!)^m} \cdot (q_n)^{hm} \cdot e^{-mnq_n} \cdot \mathbb{P}\left[\bigcap_{1 \leq i < j \leq m} (S_{ij} = \emptyset)\right]. \end{aligned} \quad (77)$$

From (74) and (77), we obtain

$$P_a \sim \frac{n^{hm}}{(h!)^m} \cdot (q_n)^{hm} \cdot e^{-mnq_n} \cdot \mathbb{P}\left[\bigcap_{1 \leq i < j \leq m} (S_{ij} = \emptyset)\right]. \quad (78)$$

By the union bound, it is clear that

$$\begin{aligned} & \mathbb{P}\left[\bigcap_{1 \leq i < j \leq m} (S_{ij} = \emptyset)\right] \\ & \geq 1 - \sum_{1 \leq i < j \leq m} \mathbb{P}[S_{ij} \neq \emptyset] = 1 - \binom{m}{2} q_n. \end{aligned} \quad (79)$$

From (36) and (79), since a probability is at most 1, we get

$$\lim_{n \rightarrow \infty} \mathbb{P}\left[\bigcap_{1 \leq i < j \leq m} (S_{ij} = \emptyset)\right] = 1. \quad (80)$$

Using (80) in (78), we establish (41).

## VII. ESTABLISHING THEOREMS 4–6

Theorems 4–6 present results on  $k$ -robustness for binomial/uniform/general random intersection graphs. We prove Theorems 4–6 in this section and start with explaining the idea below. First, the zero-law of Theorem 4 is established from the zero-law of Theorem 1 since  $k$ -robustness implies the property of minimum node degree being at least  $k$  from Lemma 1 above, while the one-law of Theorem 4 is proven from the coupling between binomial random intersection graphs and Erdős–Rényi graphs given by Lemma 6 of Section IV-C2. Second, the zero-law of Theorem 5 is demonstrated from the zero-law of Theorem 2 because  $k$ -robustness implies the property of minimum node degree being at least  $k$  from Lemma 1 above, while the one-law of Theorem 5 is established from the coupling between binomial random intersection graphs and uniform random intersection graphs given by Lemma 7 of Section IV-C3. Finally, both the zero-law and one-law of Theorem 6 are proved from the coupling between general random intersection graphs and uniform random intersection graphs given by Lemma 5 of Section IV-C1.

### A. The Proof of Theorem 4

Since  $k$ -robustness implies the property of minimum node degree being at least  $k$  from Lemma 1, the zero-law of Theorem 4 is clear from (6a) of Theorem 1 in view that under conditions of Theorem 4, if  $\lim_{n \rightarrow \infty} \alpha_n = -\infty$ ,

$$\begin{aligned} & \mathbb{P}[G_b(n, P_n, p_n) \text{ is } k\text{-robust.}] \\ & \leq \mathbb{P}\left[G_b(n, P_n, p_n) \text{ has a} \right. \\ & \quad \left. \text{minimum node degree at least } k. \right] \rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned} \quad (81)$$

Note that Theorem 1 uses  $P_n = \omega(n(\ln n)^5)$  while Theorem 4 uses  $P_n = \Omega(n(\ln n)^5)$ . Above we can use (6a) since (6a) still holds under  $P_n = \Omega(n(\ln n)^5)$  as given in Remark 1 after Theorem 1.

Below we prove the one-law of Theorem 4. As explained in Appendix A, we can introduce an extra condition  $|\alpha_n| = o(\ln n)$  in proving Theorem 4. Given (10) and  $|\alpha_n| = o(\ln n)$ , we have

$$p_n^2 P_n \sim \frac{\ln n}{n},$$

which together with condition  $P_n = \Omega(n(\ln n)^5)$  leads to

$$p_n \sim \sqrt{\frac{\ln n}{n P_n}} = O\left(\sqrt{\frac{\ln n}{n^2 (\ln n)^5}}\right) = O\left(\frac{1}{n(\ln n)^2}\right). \quad (82)$$

Noting that (82) implies condition  $p_n = O\left(\frac{1}{n \ln n}\right)$  in Lemma 6, we apply Lemmas 3, 4 and 6, and condition (10) to derive the following: there exists  $\hat{p}_n = \frac{\ln n + (k-1) \ln \ln n + \alpha_n - O(1)}{n}$  such that

if  $\lim_{n \rightarrow \infty} \alpha_n = \infty$ ,

$$\begin{aligned} & \mathbb{P}[\text{Graph } G_b(n, P_n, p_n) \text{ is } k\text{-robust.}] \\ & \geq \mathbb{P}[\text{Graph } G(n, \hat{p}_n) \text{ is } k\text{-robust.}] - o(1) \rightarrow 1, \text{ as } n \rightarrow \infty. \end{aligned} \quad (83)$$

The proof of Theorem 4 is completed via (81) and (83).

### B. The Proof of Theorem 5

As explained in Appendix B, we can introduce an extra condition  $|\alpha_n| = o(\ln n)$  in proving Theorem 5. Since  $k$ -robustness implies that the minimum node degree is at least  $k$  from Lemma 1, the zero-law of Theorem 5 is clear from Lemma 10 in view that under conditions of Theorem 5 with the extra condition  $|\alpha_n| = o(\ln n)$ , if  $\lim_{n \rightarrow \infty} \alpha_n = -\infty$ ,

$$\begin{aligned} & \mathbb{P}[G_u(n, P_n, K_n) \text{ is } k\text{-robust.}] \\ & \leq \mathbb{P}\left[G_u(n, P_n, K_n) \text{ has a} \right. \\ & \quad \left. \text{minimum node degree at least } k. \right] \rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned} \quad (84)$$

Below we establish the one-law of Theorem 5 with the help of Theorem 4. Given  $K_n = \Omega((\ln n)^3) = \omega(\ln n)$ , we use Lemma 8 to obtain that with  $p_n$  set by

$$p_n = \frac{K_n}{P_n} \left(1 - \sqrt{\frac{3 \ln n}{K_n}}\right), \quad (85)$$

it holds that

$$\begin{aligned} & \mathbb{P}[\text{Graph } G_u(n, P_n, K_n) \text{ is } k\text{-robust.}] \\ & \geq \mathbb{P}[\text{Graph } G_b(n, P_n, p_n) \text{ is } k\text{-robust.}] - o(1). \end{aligned} \quad (86)$$

From (12) and  $|\alpha_n| = o(\ln n)$ , we obtain  $\frac{K_n^2}{P_n} \sim \frac{\ln n}{n}$ , which together with  $K_n = \Omega((\ln n)^3)$  results in

$$P_n \sim \frac{n K_n^2}{\ln n} = \Omega(n(\ln n)^5), \quad (87)$$

From  $K_n = \Omega((\ln n)^3)$  and (85), it follows that

$$\begin{aligned} p_n^2 P_n &= \left[\frac{K_n}{P_n} \left(1 - \sqrt{\frac{3 \ln n}{K_n}}\right)\right]^2 \cdot P_n \\ &= \frac{K_n^2}{P_n} \cdot \left[1 - O\left(\frac{1}{\ln n}\right)\right]. \end{aligned} \quad (88)$$

By (12) and (88), it is clear that

$$p_n^2 P_n = \frac{\ln n + (k-1) \ln \ln n + \alpha_n - O(1)}{n}. \quad (89)$$

Given (87) (89) and  $|\alpha_n| = o(\ln n)$ , we use Theorem 4 and (86) to get that if  $\lim_{n \rightarrow \infty} \alpha_n = \infty$ , then

$$\mathbb{P}[G_u(n, P_n, K_n) \text{ is } k\text{-robust.}] \rightarrow 1, \text{ as } n \rightarrow \infty. \quad (90)$$

The proof of Theorem 5 is completed via (84) and (90).

### C. The Proof of Theorem 6

Similar to the process of proving Theorem 3 with the help of Theorem 2, we demonstrate Theorem 6 using Theorem 5, which has been proved above.

Given Lemmas 4 and 5 and the fact that  $k$ -robustness is a monotone increasing graph property, we will show Theorem 6 once proving for any  $\epsilon_n = o\left(\frac{1}{\ln n}\right)$  that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \mathbb{P}[G_u(n, P_n, (1 \pm \epsilon_n) \mathbb{E}[X_n]) \text{ is } k\text{-robust.}] \\ & = \begin{cases} 0, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ 1, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty. \end{cases} \end{aligned} \quad (91)$$

Under  $\mathbb{E}[X_n] = \Omega(\sqrt{\ln n})$  and  $\epsilon_n = o(\frac{1}{\ln n})$ , it follows that  $(1 \pm \epsilon_n)\mathbb{E}[X_n] = \Omega(\sqrt{\ln n})$ . From Theorem 5, we will have (27) once we prove that sequences  $\gamma_n^+$  and  $\gamma_n^-$  defined through

$$\frac{\{(1 \pm \epsilon_n)\mathbb{E}[X_n]\}^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \gamma_n^\pm}{n} \quad (92)$$

satisfy

$$\lim_{n \rightarrow \infty} \gamma_n^\pm = \begin{cases} -\infty, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = -\infty, \\ \infty, & \text{if } \lim_{n \rightarrow \infty} \alpha_n = \infty. \end{cases} \quad (93)$$

Note that (14) and (92) are exactly the same as (7) and (29), while (93) is a subset of (30). Since (31) follows from (7) (29) and  $\epsilon_n = o(\frac{1}{\ln n})$ , we use (14) (92) and  $\epsilon_n = o(\frac{1}{\ln n})$  to obtain (31), which further yields (93). Therefore, as mentioned above, we establish (91) and finally Theorem 6.

### VIII. ESTABLISHING LEMMAS IN SECTION IV

Lemmas 3 and 7 are clear in Section IV. Below we prove Lemmas 5, 6 and 8.

#### A. The Proof of Lemma 5

According to [5, Lemma 3], for any monotone increasing graph property  $\mathcal{I}$  and any  $|\epsilon_n| < 1$ ,

$$\begin{aligned} & \mathbb{P}[G(n, P_n, \mathcal{D}_n) \text{ has } \mathcal{I}] - \mathbb{P}[G_u(n, P_n, (1 - \epsilon_n)\mathbb{E}[X_n]) \text{ has } \mathcal{I}] \\ & \geq \{1 - \mathbb{P}[X_n < (1 - \epsilon_n)\mathbb{E}[X_n]]\}^n - 1, \end{aligned} \quad (94)$$

and

$$\begin{aligned} & \mathbb{P}[G(n, P_n, \mathcal{D}_n) \text{ has } \mathcal{I}] - \mathbb{P}[G_u(n, P_n, (1 + \epsilon_n)\mathbb{E}[X_n]) \text{ has } \mathcal{I}] \\ & \leq 1 - \{1 - \mathbb{P}[X_n > (1 + \epsilon_n)\mathbb{E}[X_n]]\}^n. \end{aligned} \quad (95)$$

By (94) (95) and the fact that  $\lim_{n \rightarrow \infty} (1 - m_n)^n = 1$  for  $m_n = o(\frac{1}{\ln n})$  (this can be proved by a simple Taylor series expansion as in [31, Fact 2]), the proof of Lemma 5 is completed once we demonstrate that with  $\text{Var}[X_n] = o(\frac{\{\mathbb{E}[X_n]\}^2}{n(\ln n)^2})$ , there exists  $\epsilon_n = o(\frac{1}{\ln n})$  such that

$$\mathbb{P}[X_n < (1 - \epsilon_n)\mathbb{E}[X_n]] = o\left(\frac{1}{n}\right), \quad (96)$$

and

$$\mathbb{P}[X_n > (1 + \epsilon_n)\mathbb{E}[X_n]] = o\left(\frac{1}{n}\right). \quad (97)$$

To prove (96) and (97), Chebyshev's inequality yields

$$\mathbb{P}[|X_n - \mathbb{E}[X_n]| > \epsilon_n \mathbb{E}[X_n]] \leq \frac{\text{Var}[X_n]}{\{\epsilon_n \mathbb{E}[X_n]\}^2}. \quad (98)$$

We set  $\epsilon_n$  by  $\epsilon_n = \sqrt[4]{\frac{n \text{Var}[X_n]}{\{\mathbb{E}[X_n]\}^2}} \cdot \frac{1}{\sqrt{\ln n}}$ . Then given condition

$\text{Var}[X_n] = o(\frac{\{\mathbb{E}[X_n]\}^2}{n(\ln n)^2})$ , we obtain

$$\epsilon_n = o\left(\sqrt[4]{\frac{1}{(\ln n)^2}}\right) \cdot \frac{1}{\sqrt{\ln n}} = o\left(\frac{1}{\ln n}\right), \quad (99)$$

and

$$\frac{\text{Var}[X_n]}{\{\epsilon_n \mathbb{E}[X_n]\}^2} = \sqrt{\frac{\text{Var}[X_n]}{n \{\mathbb{E}[X_n]\}^2}} \cdot \ln n = o\left(\frac{1}{n}\right). \quad (100)$$

By (98) (99) and (100), it is straightforward to see that (96) and (97) hold with  $\epsilon_n = o(\frac{1}{\ln n})$ . Therefore, we have completed the proof of Lemma 5.

#### B. The Proof of Lemma 6

In view of [18, Theorem 1], if  $p_n^2 P_n < 1$  and  $p_n = o(\frac{1}{n})$ , with  $\hat{p}_n := p_n^2 P_n \cdot \left(1 - np_n + 2p_n - \frac{p_n^2 P_n}{2}\right)$ , then (20) follows. Given conditions  $p_n = O(\frac{1}{n \ln n})$  and  $p_n^2 P_n = O(\frac{1}{\ln n})$  in Lemma 6,  $p_n^2 P_n < 1$  and  $p_n = o(\frac{1}{n})$  clearly hold. Then Lemma 6 is proved once we show  $\hat{p}_n = p_n^2 P_n \cdot [1 - O(\frac{1}{\ln n})]$ , which is easy to see via

$$\begin{aligned} & -np_n + 2p_n - \frac{p_n^2 P_n}{2} \\ & = (-n+2) \cdot O\left(\frac{1}{n \ln n}\right) - \frac{1}{2} \cdot O\left(\frac{1}{\ln n}\right) = -O\left(\frac{1}{\ln n}\right). \end{aligned}$$

Hence, the proof of Lemma 6 is completed.

#### C. The Proof of Lemma 8

We use Lemma 7 to prove Lemma 8. From  $K_n = \omega(\ln n)$  and  $p_n = \frac{K_n}{P_n} \left(1 - \sqrt{\frac{3 \ln n}{K_n}}\right)$ , we first obtain  $p_n P_n = \omega(\ln n)$  and then for all  $n$  sufficiently large,

$$\begin{aligned} & K_n - \left[p_n P_n + \sqrt{3(p_n P_n + \ln n) \ln n}\right] \\ & = K_n \sqrt{\frac{3 \ln n}{K_n}} - \sqrt{3 \left[K_n \left(1 - \sqrt{\frac{3 \ln n}{K_n}}\right) + \ln n\right] \ln n} \\ & = \sqrt{3 K_n \ln n} - \sqrt{3 \left[K_n + \sqrt{\ln n} \left(\sqrt{\ln n} - \sqrt{3 K_n}\right)\right] \ln n} \\ & \geq 0. \end{aligned}$$

Then by Lemma 7, Lemma 8 is now established.

### IX. CONCLUSION AND FUTURE WORK

Under a general random intersection graph model, we derive sharp zero-one laws for  $k$ -connectivity and  $k$ -robustness, as well as the asymptotically exact probability of  $k$ -connectivity, where  $k$  is an arbitrary positive integer. A future direction is to obtain the asymptotically exact probability of  $k$ -robustness for a precise characterization on the robustness strength.

### REFERENCES

- [1] J. Zhao, O. Yağan, and V. Gligor, "On Connectivity and Robustness in Random Intersection Graphs," 2016. Full version of this paper, available online at <https://sites.google.com/site/publicworkofzhao/TAC2016.pdf>
- [2] N. Abaid and M. Porfiri, "Consensus over numerosity-constrained random networks," *IEEE Transactions on Automatic Control*, 56(3):649–654, 2011.
- [3] S. Blackburn and S. Gerke, "Connectivity of the uniform random intersection graph," *Discrete Mathematics*, 309(16), 2009.
- [4] M. Bloznelis, "Degree and clustering coefficient in sparse random intersection graphs," *The Annals of Applied Probability*, 23(3):1254–1289, 2013.
- [5] M. Bloznelis, J. Jaworski, and K. Rybarczyk, "Component evolution in a secure wireless sensor network," *Networks*, 53:19–26, January 2009.
- [6] B. Bollobás, *Random graphs*. Cambridge Studies in Advanced Mathematics (No. 73). Cambridge University Press, 2001.
- [7] M. Bradonjić, A. Hagberg, N. Hengartner, and A. Percus, "Component evolution in general random intersection graphs," in *Workshop on Algorithms and Models for the Web Graph (WAW)*, pp. 36–49, 2010.
- [8] H. Chan, V. Gligor, A. Perrig, and G. Muralidharan, "On the distribution and revocation of cryptographic keys in sensor networks," *IEEE Transactions on Dependable and Secure Computing (TDSC)*, 2(3):233–247, 2005.
- [9] M. Deijfen and W. Kets, "Random intersection graphs with tunable degree distribution and clustering," *Probability in the Engineering and Information Sciences*, 23:661–674, 2009.

- [10] D. Dolev. The byzantine generals strike again. *Journal of Algorithms*, 3(1):14–30, 1982.
- [11] P. Erdős and A. Rényi. On random graphs, I. *Publicationes Mathematicae (Debrecen)*, 6:290–297, 1959.
- [12] P. Erdős and A. Rényi. On the strength of connectedness of random graphs. *Acta Math. Acad. Sci. Hungar.*, pages 261–267, 1961.
- [13] L. Eschenauer and V. Gligor. A key-management scheme for distributed sensor networks. In *ACM Conference on Computer and Communications Security (CCS)*, 2002.
- [14] E. Godehardt and J. Jaworski. Two models of random intersection graphs for classification. *Exploratory Data Analysis in Empirical Research*, pages 67–81, 2003.
- [15] H. LeBlanc, H. Zhang, X. Koutsoukos, and S. Sundaram. Resilient asymptotic consensus in robust networks. *IEEE Journal on Selected Areas in Communications (JSAC)*, 31(4):766–781, April 2013.
- [16] H. LeBlanc, H. Zhang, S. Sundaram, and X. Koutsoukos. Resilient continuous-time consensus in fractional robust networks. In *American Control Conference (ACC)*, 2013.
- [17] K. Rybarczyk. Diameter, connectivity and phase transition of the uniform random intersection graph. *Discrete Mathematics*, 311, 2011.
- [18] K. Rybarczyk. Sharp threshold functions for the random intersection graph via a coupling method. *The Electronic Journal of Combinatorics*, 18:36–47, 2011.
- [19] K. Rybarczyk. The coupling method for inhomogeneous random intersection graphs. *arXiv*, 1301.0466 [math.CO], January 2013.
- [20] K. Singer-Cohen. *Random intersection graphs*. PhD thesis, Department of Mathematical Sciences, The Johns Hopkins University, 1995.
- [21] O. Yağan. *Random Graph Modeling of Key Distribution Schemes in Wireless Sensor Networks*. PhD thesis, Dept. of ECE, University of Maryland, June 2011.
- [22] O. Yağan. Performance of the Eschenauer-Gligor key distribution scheme under an on/off channel. *IEEE Transactions on Information Theory*, 58(6):3821–3835, June 2012.
- [23] O. Yağan. Zero-one laws for connectivity in inhomogeneous random key graphs. *ArXiv e-prints*, August 2015. Available online at <http://arxiv.org/pdf/1508.02407v1.pdf>.
- [24] O. Yağan and A. M. Makowski. Zero-one laws for connectivity in random key graphs. *IEEE Transactions on Information Theory*, 58(5):2983–2999, May 2012.
- [25] H. Zhang, E. Fata, and S. Sundaram. A notion of robustness in complex networks. *IEEE Transactions on Control of Network Systems*, 2015.
- [26] H. Zhang and S. Sundaram. Robustness of complex networks with implications for consensus and contagion. In *IEEE Conference on Decision and Control (CDC)*, pages 3426–3432, December 2012.
- [27] H. Zhang and S. Sundaram. Robustness of information diffusion algorithms to locally bounded adversaries. In *IEEE American Control Conference (ACC)*, pages 5855–5861, 2012.
- [28] H. Zhang and S. Sundaram. A simple median-based resilient consensus algorithm. In *Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 1734–1741, 2012.
- [29] J. Zhao, O. Yağan, and V. Gligor. On asymptotically exact probability of  $k$ -connectivity in random key graphs intersecting Erdős–Rényi graphs. 2014. Available online at <http://arxiv.org/pdf/1409.6022v1.pdf>
- [30] J. Zhao, O. Yağan, and V. Gligor. On the strengths of connectivity and robustness in general random intersection graphs. In *IEEE Conference on Decision and Control (CDC)*, pages 3661–3668, Dec 2014.
- [31] J. Zhao, O. Yağan, and V. Gligor.  $k$ -Connectivity in random key graphs with unreliable links. *IEEE Transactions on Information Theory*, 61(7):3810–3836, July 2015.
- [32] J. Zhao, O. Yağan, and V. Gligor. On  $k$ -connectivity and minimum vertex degree in random  $s$ -intersection graphs. In *ACM-SIAM Meeting on Analytic Algorithmics and Combinatorics (ANALCO)*, pages 1–15, January 2015.
- [33] R. Di Pietro, L. Mancini, A. Mei, A. Panconesi, and J. Radhakrishnan. Sensor networks that are provably resilient. In *International Conference on Security and Privacy for Emerging Areas in Communication Networks (Securecomm)*, pages 1–10, Aug 2006.
- [34] R. Di Pietro, L. Mancini, A. Mei, A. Panconesi, and J. Radhakrishnan. Redoubtable sensor networks. *ACM Transactions on Information and System Security*, 11(3):13:1–13:22, 2008.
- [35] S. Blackburn, D. Stinson, and J. Upadhyay. On the complexity of the herding attack and some related attacks on hash functions. *Designs, Codes and Cryptography*, 64(1-2):171–193, 2012.

## APPENDIX

A. A lemma to confine  $|\alpha_n|$  in Theorems 1 and 4 as  $o(\ln n)$

We present Lemma 13 to confine  $|\alpha_n|$  in Theorems 1 and 4 as  $o(\ln n)$ ; i.e., if Theorems 1 and 4 hold under an extra condition  $|\alpha_n| = o(\ln n)$ , then they also hold regardless of this condition.

**Lemma 13 (a)** For graph  $G_b(n, P_n, p_n)$  under

$$p_n^2 P_n = \frac{\ln n + (k-1) \ln \ln n + \beta_n}{n} \quad (101)$$

with  $\lim_{n \rightarrow \infty} \beta_n = -\infty$ , there exists graph  $G_b(n, P_n, \widehat{p}_n)$  under

$$\widehat{p}_n^2 P_n = \frac{\ln n + (k-1) \ln \ln n + \widetilde{\beta}_n}{n} \quad (102)$$

with  $\lim_{n \rightarrow \infty} \widetilde{\beta}_n = -\infty$  and  $\widehat{\beta}_n = -o(\ln n)$  such that  $G_b(n, P_n, p_n) \preceq G_b(n, P_n, \widehat{p}_n)$ .

**(b)** For graph  $G_b(n, P_n, p_n)$  under

$$p_n^2 P_n = \frac{\ln n + (k-1) \ln \ln n + \beta_n}{n} \quad (103)$$

with  $\lim_{n \rightarrow \infty} \beta_n = \infty$ , there exists graph  $G_b(n, P_n, \widehat{p}_n)$  under

$$\widehat{p}_n^2 P_n = \frac{\ln n + (k-1) \ln \ln n + \widehat{\beta}_n}{n} \quad (104)$$

with  $\lim_{n \rightarrow \infty} \widehat{\beta}_n = \infty$  and  $\widehat{\beta}_n = o(\ln n)$  such that  $G_b(n, P_n, \widehat{p}_n) \preceq G_b(n, P_n, p_n)$ .

The proof of Lemma 13 is given in Section C.

We now explain that given Lemma 13, if Theorems 1 and 4 hold under the extra condition  $|\alpha_n| = o(\ln n)$ , then they also hold regardless of the extra condition. Note that results (2c) and (3c) both have a condition  $\lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty)$ , which clearly implies  $|\alpha_n| = o(\ln n)$ . Hence, we only need to look at results (2a) (2b) (3a) (3b) (11a) and (11b). In particular, we will show that

if (2a) (3a) and (11a) hold under condition  $\alpha_n = -o(\ln n)$ , then they also hold regardless of the condition. (105)

and

if (2b) (3b) and (11b) hold under condition  $\alpha_n = o(\ln n)$ , then they also hold regardless of the condition. (106)

To see (105), we use Lemma 13-Property (a) and Lemma 4, and note that  $k$ -connectivity, the property of minimum node degree being at least  $k$ , and  $k$ -robustness are all monotone increasing graph properties. Then with  $\mathcal{J}$  denoting any one of the above three properties, for graph  $G_b(n, P_n, p_n)$  under (101) with  $\lim_{n \rightarrow \infty} \beta_n = -\infty$ , there exists graph  $G_b(n, P_n, \widehat{p}_n)$  under (102) with  $\lim_{n \rightarrow \infty} \widetilde{\beta}_n = -\infty$  and  $\widehat{\beta}_n = -o(\ln n)$  such that

$$\begin{aligned} \mathbb{P}[G_b(n, P_n, p_n) \text{ has } \mathcal{J}] \\ \leq \mathbb{P}[G_b(n, P_n, \widehat{p}_n) \text{ has } \mathcal{J}]. \end{aligned} \quad (107)$$

If (2a) and (11a) hold under condition  $\alpha_n = -o(\ln n)$ , then we use them on graph  $G_b(n, P_n, \widehat{p}_n)$  to obtain

$$\lim_{n \rightarrow \infty} \mathbb{P}[G_b(n, P_n, \widehat{p}_n) \text{ has } \mathcal{J}] = 0, \quad (108)$$

Therefore, (107) and (108) yield

$$\lim_{n \rightarrow \infty} \mathbb{P}[G_b(n, P_n, p_n) \text{ has } \mathcal{J}] = 0.$$

In other words, for graph  $G_b(n, P_n, p_n)$ , with  $\alpha_n$  in Theorems 1 and 4 replaced by  $\beta_n$ , (2a) (3a) and (11a) hold. Note that we do not have any constraint on whether  $\beta_n$  can be expressed as  $-o(\ln n)$ . Hence, the arguments above establish (105). The proof

of (106) using Lemma 13-Property (b) is similar to that of (105) using Lemma 13-Property (a). We omit the details for simplicity.

*B. A lemma to confine  $|\alpha_n|$  in Theorems 2 and 5 as  $o(\ln n)$*

We present Lemma 14 to confine  $|\alpha_n|$  in Theorems 2 and 5 as  $o(\ln n)$ ; i.e., if Theorems 2 and 5 hold under an extra condition  $|\alpha_n| = o(\ln n)$ , then they also hold regardless of this condition.

**Lemma 14 (a)** *For graph  $G_u(n, P_n, K_n)$  under  $P_n = \Omega(n)$  and*

$$\frac{K_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \beta_n}{n} \quad (109)$$

*with  $\lim_{n \rightarrow \infty} \beta_n = -\infty$ , there exists graph  $G_u(n, P_n, \widetilde{K}_n)$  under  $P_n = \Omega(n)$  and*

$$\frac{\widetilde{K}_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \widetilde{\beta}_n}{n} \quad (110)$$

*with  $\lim_{n \rightarrow \infty} \widetilde{\beta}_n = -\infty$  and  $\widetilde{\beta}_n = -o(\ln n)$ , such that  $G_u(n, P_n, K_n) \preceq G_u(n, P_n, \widetilde{K}_n)$ .*

**(b)** *For graph  $G_u(n, P_n, K_n)$  under  $P_n = \Omega(n)$  and*

$$\frac{K_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \beta_n}{n} \quad (111)$$

*with  $\lim_{n \rightarrow \infty} \beta_n = \infty$ , there exists graph  $G_u(n, P_n, \widehat{K}_n)$  under  $P_n = \Omega(n)$  and*

$$\frac{\widehat{K}_n^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \widehat{\beta}_n}{n} \quad (112)$$

*with  $\lim_{n \rightarrow \infty} \widehat{\beta}_n = \infty$  and  $\widehat{\beta}_n = o(\ln n)$ , such that  $G_u(n, P_n, \widehat{K}_n) \preceq G_u(n, P_n, K_n)$ .*

The proof of Lemma 14 is given in Section D.

We now explain that given Lemma 14, if Theorems 2 and 5 hold under the extra condition  $|\alpha_n| = o(\ln n)$ , then they also hold regardless of the extra condition. Note that results (5c) and (6c) both have a condition  $\lim_{n \rightarrow \infty} \alpha_n = \alpha^* \in (-\infty, \infty)$ , which clearly implies  $|\alpha_n| = o(\ln n)$ . Hence, we only need to look at results (5a) (5b) (6a) (6b) (13a) and (13b). In particular, we need to show that

if (5a) (6a) and (13a) hold under condition  $\alpha_n = -o(\ln n)$ , then they also hold regardless of the condition.

(113)

and

if (5b) (6b) and (13b) hold under condition  $\alpha_n = o(\ln n)$ , then they also hold regardless of the condition.

(114)

The process of proving (113) and (114) using Lemma 14 is the same as the above process of proving (105) and (106) using Lemma 13. For brevity, we do not repeat the details here.

### C. Proof of Lemma 13

**Proving property (a):**

We set

$$\widetilde{\beta}_n = \max\{\beta_n, -\ln \ln n\}. \quad (115)$$

Given (115) and  $\lim_{n \rightarrow \infty} \beta_n = -\infty$ , we obtain  $\lim_{n \rightarrow \infty} \widetilde{\beta}_n = -\infty$  and  $\widetilde{\beta}_n = -o(\ln n)$ . We use  $\beta_n = -o(\ln n)$  and (102) to have  $\widetilde{p}_n^2 P_n \sim \frac{\ln n}{n}$ , so it is clear for all  $n$  sufficiently large that  $\widetilde{p}_n$  is less than 1 and can be used as a probability. Under  $p_n \leq \widetilde{p}_n$ , by [18, Section 3], there exists a graph coupling under

which  $G_b(n, P_n, p_n)$  is a spanning subgraph of  $G_b(n, P_n, \widetilde{p}_n)$ ; i.e.,  $G_b(n, P_n, p_n) \preceq G_b(n, P_n, \widetilde{p}_n)$ .

**Proving property (b):**

We set

$$\widehat{\beta}_n = \min\{\beta_n, \ln \ln n\}. \quad (116)$$

Given (116) and  $\lim_{n \rightarrow \infty} \beta_n = \infty$ , we clearly obtain  $\lim_{n \rightarrow \infty} \widehat{\beta}_n = \infty$  and  $\widehat{\beta}_n = o(\ln n)$ .

It holds from (116) that  $\widehat{\beta}_n \leq \beta_n$ , which along with (103) and (104) yields  $p_n \geq \widehat{p}_n$ . Under  $p_n \geq \widehat{p}_n$ , by [18, Section 3], there exists a graph coupling under which  $G_b(n, P_n, p_n)$  is a spanning supergraph of  $G_b(n, P_n, \widehat{p}_n)$ ; i.e.,  $G_b(n, P_n, \widehat{p}_n) \preceq G_b(n, P_n, p_n)$ .

### D. The Proof of Lemma 14

**Proving property (a):**

We define  $\widetilde{\beta}_n^*$  by

$$\widetilde{\beta}_n^* = \max\{\beta_n, -\ln \ln n\}, \quad (117)$$

and define  $\widetilde{K}_n^*$  such that

$$\frac{(\widetilde{K}_n^*)^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \widetilde{\beta}_n^*}{n}. \quad (118)$$

Note that  $\widetilde{K}_n^*$  might or might not be an integer. We set

$$\widetilde{K}_n := \lfloor \widetilde{K}_n^* \rfloor, \quad (119)$$

where the floor function  $\lfloor x \rfloor$  means the largest integer not greater than  $x$ .

From (111) (117) and (118), it holds that

$$K_n \leq \widetilde{K}_n^*. \quad (120)$$

Then by (119) (120) and the fact that  $K_n$  and  $\widetilde{K}_n$  are both integers, it follows that

$$K_n \leq \widetilde{K}_n. \quad (121)$$

From (121), by [5, Lemma 3], there exists a graph coupling under which  $G_u(n, P_n, K_n)$  is a spanning subgraph of  $G_u(n, P_n, \widetilde{K}_n)$ ; i.e.,  $G_u(n, P_n, K_n) \preceq G_u(n, P_n, \widetilde{K}_n)$ . Therefore, the proof of property (a) is completed once we show  $\widetilde{\beta}_n$  defined in (110) satisfies

$$\lim_{n \rightarrow \infty} \widetilde{\beta}_n = -\infty, \quad (122)$$

$$\text{and } \widetilde{\beta}_n = -o(\ln n). \quad (123)$$

We first prove (122). From (110) (118) and (119), it holds that

$$\widetilde{\beta}_n \leq \widetilde{\beta}_n^*, \quad (124)$$

which together with (117) and  $\lim_{n \rightarrow \infty} \beta_n = -\infty$  yields (122).

Now we establish (123). From (119), we have  $\widetilde{K}_n > \widetilde{K}_n^* - 1$ . Then from (110), it holds that

$$\begin{aligned} \widetilde{\beta}_n &= n \cdot \frac{\widetilde{K}_n^2}{P_n} - [\ln n + (k-1) \ln \ln n] \\ &> n \cdot \frac{(\widetilde{K}_n^*)^2 - 2\widetilde{K}_n^*}{P_n} - [\ln n + (k-1) \ln \ln n]. \end{aligned} \quad (125)$$

By  $\lim_{n \rightarrow \infty} \beta_n = -\infty$ , it holds that  $\beta_n \leq 0$  for all  $n$  sufficiently large. Then from (117), it follows that

$$\widetilde{\beta}_n^* = -O(\ln \ln n), \quad (126)$$

which along with (118) yields

$$\frac{\widetilde{K}_n^*}{P_n} \sim \sqrt{\frac{\ln n}{nP_n}} = O\left(\frac{\sqrt{\ln n}}{n}\right). \quad (127)$$

Applying (118) (127) and  $P_n = \Omega(n)$  to (125), we obtain

$$\begin{aligned}\widetilde{\beta}_n &> \left\{ n \cdot \frac{(\widetilde{K}_n^*)^2}{P_n} - [\ln n + (k-1) \ln \ln n] \right\} - 2n \cdot \frac{\widetilde{K}_n^*}{P_n} \\ &= \widetilde{\beta}_n^* - O(\sqrt{\ln n}).\end{aligned}\quad (128)$$

Thus, from (124) (126) and (128), clearly  $\widetilde{\beta}_n$  can be written as  $-O(\sqrt{\ln n})$  and further  $-o(\ln n)$ ; i.e., (123) is proved. Then as explained above, since we have shown (122) and (123), property (a) of Lemma 14 is established.

**Proving property (b):**

We define  $\widehat{\beta}_n^*$  by

$$\widehat{\beta}_n^* = \min\{\beta_n, \ln \ln n\}, \quad (129)$$

and define  $\widehat{K}_n^*$  such that

$$\frac{(\widehat{K}_n^*)^2}{P_n} = \frac{\ln n + (k-1) \ln \ln n + \widehat{\beta}_n^*}{n}. \quad (130)$$

We set

$$\widehat{K}_n := \lceil \widehat{K}_n^* \rceil. \quad (131)$$

From (111) (129) and (130), it holds that

$$K_n \geq \widehat{K}_n^*. \quad (132)$$

Then by (131) (132) and the fact that  $K_n$  and  $\widehat{K}_n$  are both integers, it follows that

$$K_n \geq \widehat{K}_n. \quad (133)$$

From (133), by [5, Lemma 3], there exists a graph coupling under which  $G_u(n, P_n, K_n)$  is a spanning supergraph of  $G_u(n, P_n, \widehat{K}_n)$ ; i.e.,  $G_u(n, P_n, \widehat{K}_n) \preceq G_u(n, P_n, K_n)$ . Therefore, the proof of property (b) is completed once we show  $\widehat{\beta}_n$  defined in (112) satisfies

$$\lim_{n \rightarrow \infty} \widehat{\beta}_n = \infty, \quad (134)$$

$$\text{and } \widehat{\beta}_n = o(\ln n). \quad (135)$$

We first prove (134). From (112) (130) and (131), it holds that

$$\widehat{\beta}_n \geq \widehat{\beta}_n^*, \quad (136)$$

which together with (129) and  $\lim_{n \rightarrow \infty} \beta_n = \infty$  yields (134).

Now we establish (135). From (131), we have  $\widehat{K}_n < \widehat{K}_n^* + 1$ . Then from (112), it holds that

$$\begin{aligned}\widehat{\beta}_n &= n \cdot \frac{\widehat{K}_n^2}{P_n} - [\ln n + (k-1) \ln \ln n] \\ &\leq n \cdot \frac{(\widehat{K}_n^*)^2 + 3\widehat{K}_n^*}{P_n} - [\ln n + (k-1) \ln \ln n].\end{aligned}\quad (137)$$

By  $\lim_{n \rightarrow \infty} \beta_n = \infty$ , it holds that  $\beta_n \geq 0$  for all  $n$  sufficiently large. Then from (129), it follows that

$$\widehat{\beta}_n^* = O(\ln \ln n), \quad (138)$$

which along with (130) and condition  $P_n = \Omega(n)$  induces

$$\frac{\widehat{K}_n^*}{P_n} \sim \sqrt{\frac{\ln n}{nP_n}} = O\left(\frac{\sqrt{\ln n}}{n}\right). \quad (139)$$

Hence, we have  $\lim_{n \rightarrow \infty} \widehat{K}_n^* = \infty$  and it further holds for all  $n$  sufficient large that

$$(\widehat{K}_n^* + 1)^2 < (\widehat{K}_n^*)^2 + 3\widehat{K}_n^*. \quad (140)$$

Applying (130) (139) and  $P_n = \Omega(n)$  to (137), we obtain

$$\begin{aligned}\widehat{\beta}_n &< \left\{ n \cdot \frac{(\widehat{K}_n^*)^2}{P_n} - [\ln n + (k-1) \ln \ln n] \right\} + 3n \cdot \frac{\widehat{K}_n^*}{P_n} \\ &= \widehat{\beta}_n^* + O(\sqrt{\ln n}).\end{aligned}\quad (141)$$

Thus, from (136) (138) and (141), clearly  $\widehat{\beta}_n$  can be written as  $O(\sqrt{\ln n})$  and further  $o(\ln n)$ ; i.e., (135) is proved. Then as explained above, since we have shown (134) and (135), property (b) of Lemma 14 is established.