

A HYBRIDIZED DISCONTINUOUS GALERKIN METHOD FOR POISSON-TYPE PROBLEMS WITH SIGN-CHANGING COEFFICIENTS *

JEONGHUN J. LEE[†] AND SANDER RHEBERGEN[‡]

Abstract. In this paper, we present a hybridized discontinuous Galerkin (HDG) method for Poisson-type problems with sign-changing coefficients. We introduce a sign-changing stabilization parameter that results in a stable HDG method independent of domain geometry and the ratio of the negative and positive coefficients. Since the Poisson-type problem with sign-changing coefficients is not elliptic, standard techniques with a duality argument to analyze the HDG method cannot be applied. Hence, we present a novel error analysis exploiting the stabilized saddle-point problem structure of the HDG method. Numerical experiments in two dimensions and for varying polynomial degree verify our theoretical results.

Key words. HDG methods, sign-changing coefficients, error analysis

AMS subject classifications. 65N12, 65N15, 65N30

1. Introduction. Partial differential equations (PDE) with sign-changing coefficients are used to model physical phenomena of meta-materials, for example, wave transmission problems between classical materials and meta-materials. Therefore, there is an emerging interest in the development of numerical methods for such PDEs. However, the bilinear forms for problems with sign-changing coefficients are not coercive, so standard techniques relying on coercivity cannot be applied. The lack of coercivity indeed poses fundamental challenges to study the well-posedness of such PDEs, as well as the development of numerical methods.

A popular approach in the study of numerical methods for PDEs with sign-changing coefficients is the “ T -coercivity” approach. This approach utilizes a linear operator T that recovers the coercivity of the bilinear form. The existence of such an operator T is a natural assumption because the existence of T is equivalent to the well-posedness of the PDE in the sense of Banach–Nečas–Babuška in the functional analysis framework [7]. The application of the T operator to develop numerical methods was first proposed in [2]. Since then, the T -coercivity approach has been used in the development of numerical methods for Poisson-type problems [7], Helmholtz-type problems [8], and Maxwell-type problems [2, 5, 4].

In numerical methods that use T -coercivity, the T operator cannot be used directly, because the domain and the range of T are not the discrete spaces in general. For this reason, T_h is introduced, a discrete approximation of T , which has the coercivity recovery property in the discrete setting. The existence of such T_h is non-trivial and difficult to guarantee because the form of T is unknown in general. For general geometry and meshes sufficient conditions for the existence of T_h are proposed in [7], but these include non-quantitative constants such as a norm of the discrete interpolation of T and the ratio of the positive and negative coefficients (the contrast). It is furthermore shown in [7] that locally symmetric meshes across the transmission inter-

*

Funding: SR gratefully acknowledges support from the Natural Sciences and Engineering Research Council of Canada through the Discovery Grant program (RGPIN-05606-2015).

[†]Department of Mathematics, Baylor University, Waco, TX, USA (jeonghun.lee@baylor.edu), <http://orcid.org/0000-0001-5201-8526>

[‡]Department of Applied Mathematics, University of Waterloo, Canada (srheberg@uwaterloo.ca), <http://orcid.org/0000-0001-6036-0356>

face, where the coefficient changes sign, improve the quality of numerical solutions. However, the generation of such meshes is a non-trivial restriction in mesh generation for complex interfaces.

Hybridization (or static condensation) was originally introduced for mixed finite element methods to reduce computational cost [16], and it was further analyzed in [1]. More recently, hybridization was combined with the discontinuous Galerkin (DG) method. This resulted in the hybridizable discontinuous Galerkin (HDG) method which was shown to be more efficient than standard DG methods. The HDG method was systematically presented in [10] for elliptic partial differential equations. Since its introduction, it has been extended to many different problems. These include HDG methods for the Helmholtz problem, e.g., [13, 18] and the Maxwell problem, e.g., [6, 15].

In this paper, we develop a numerical method for Poisson-type problems with a sign-changing coefficient that *avoids* the T -coercivity approach. The numerical method we present in this paper is always well-posed *without* any conditions on the domain geometry or the ratio between the negative and positive coefficients. This is achieved by employing two key ideas for the problem: a mixed formulation of the problem and using a hybridized discontinuous Galerkin (HDG) method with *sign-changing* stabilization parameters. We will see that the sign-changing stabilization parameter and the discontinuous test function space in discontinuous Galerkin methods allow us to obtain stable numerical methods without any non-quantitative assumptions on mesh size or coefficients.

We also carry out an error analysis of the HDG method for this problem. The error analysis of HDG methods typically uses the Aubin–Nitsche duality argument with the full elliptic regularity assumption. However, this assumption is not feasible in our setting, because the PDE is not elliptic. By revealing a stabilized saddle point problem structure of HDG methods, we circumvent this obstacle and we can show an error estimate without using the duality argument. We note that such analysis has been applied also to HDG methods for standard Poisson-type problems to avoid the full elliptic regularity assumption in the a priori error analysis (see [12]).

The outline of this paper is as follows. In [section 2](#) we introduce the Poisson problem with a sign-changing coefficient. We introduce the HDG method and discuss its well-posedness in [section 3](#). An a priori error analysis is presented in [section 4](#). In [section 5](#) we show that a superconvergence result can be obtained when a suitable regularity assumption is available. Our analysis is verified by numerical examples in [section 6](#), while conclusions are drawn in [section 7](#).

2. The Poisson problem with a sign-changing coefficient. Let $\Omega \subset \mathbb{R}^d$, $d \geq 2$, be a polygonal domain that is divided into two subdomains Ω_+ and Ω_- such that $\overline{\Omega} = \overline{\Omega_+} \cup \overline{\Omega_-}$ and $\Omega_+ \cap \Omega_- = \emptyset$. The boundaries of Ω , Ω_+ and Ω_- are denoted by, respectively, $\partial\Omega$, $\partial\Omega_+$ and $\partial\Omega_-$. The interface separating the domains is denoted by $\Gamma_I := \overline{\Omega_+} \cap \overline{\Omega_-}$ and we define furthermore $\Gamma_+ := \partial\Omega_+ \setminus \Gamma_I$ and $\Gamma_- := \partial\Omega_- \setminus \Gamma_I$. We assume that Γ_D and Γ_N are disjoint subsets of $\partial\Omega$ such that $\partial\Omega = \overline{\Gamma_D} \cup \overline{\Gamma_N}$. The outward unit normal vector field on $\partial\Omega$ is denoted by \mathbf{n} .

Let σ be a scalar function defined as

$$(2.1) \quad \sigma := \begin{cases} \sigma_+ & \text{on } \Omega_+, \\ \sigma_- & \text{on } \Omega_-, \end{cases}$$

where $\sigma_+ > 0$ and $\sigma_- < 0$ are constants. The *contrast* is defined as $\kappa_\sigma := \sigma_-/\sigma_+$.

Throughout this paper we assume that

$$(2.2) \quad 0 < \sigma_{\min} \leq \sigma_+, \quad -\sigma_- \leq \sigma_{\max} < +\infty.$$

We consider the following partial differential equation (PDE) for the scalar $u : \Omega \rightarrow \mathbb{R}$:

$$\begin{aligned} (2.3a) \quad & \nabla \cdot (\sigma \nabla u) = f && \text{in } \Omega, \\ (2.3b) \quad & u = u_D && \text{on } \Gamma_D, \\ (2.3c) \quad & -\sigma \nabla u \cdot \mathbf{n} = u_N && \text{on } \Gamma_N, \end{aligned}$$

where $u_D : \Gamma_D \rightarrow \mathbb{R}$ and $u_N : \Gamma_N \rightarrow \mathbb{R}$ are given boundary data and $f : \Omega \rightarrow \mathbb{R}$ is a given source term. This is not a standard second order PDE, because the sign of σ is indefinite. Let $H_0^1(\Omega) := \{v \in H^1(\Omega) : v = 0 \text{ on } \partial\Omega\}$ where $H^1(\Omega)$ is the standard Sobolev space of L^2 functions such that the L^2 -norm of their gradient is bounded. The variational formulation for the problem (when $u_D = u_N = 0$) is given by: Find $u \in H_0^1(\Omega)$ such that

$$(2.4) \quad \int_{\Omega} \sigma \nabla u \cdot \nabla v \, dx = \int_{\Omega} f v \, dx \quad \forall v \in H_0^1(\Omega).$$

It is known that this problem is not always well-posed, for example, when $\kappa_{\sigma} = -1$, the problem is ill-posed. The conditions of well-posedness of the problem depends on the values of σ and the geometry of Ω_+ and Ω_- . For more information on the well-posedness of this problem we refer to [7, 3].

3. The HDG method. Introducing the auxiliary variable $\mathbf{q} = -\sigma \nabla u$, (2.3) can be written as a system of first-order equations:

$$\begin{aligned} (3.1a) \quad & \sigma^{-1} \mathbf{q} + \nabla u = 0 && \text{in } \Omega, \\ (3.1b) \quad & \nabla \cdot \mathbf{q} = -f && \text{in } \Omega, \\ (3.1c) \quad & u = u_D && \text{on } \Gamma_D, \\ (3.1d) \quad & -\sigma \nabla u \cdot \mathbf{n} = u_N && \text{on } \Gamma_N. \end{aligned}$$

In this section we introduce the HDG method for (3.1).

3.1. Preliminaries. Let \mathcal{T}_h be a triangulation of Ω and \mathcal{F}_h be the set of faces in the triangulation \mathcal{T}_h with co-dimension 1. By \mathcal{T}_h^+ we denote the subset of \mathcal{T}_h such that its elements are in Ω_+ . \mathcal{T}_h^- is defined similarly. We assume that \mathcal{T}_h is a conforming triangulation with respect to Γ_I , i.e., there is a subset \mathcal{F}_h^i of \mathcal{F}_h such that $\Gamma_I = \cup_{F \in \mathcal{F}_h^i} \bar{F}$. For later reference we define \mathcal{F}_h^+ and \mathcal{F}_h^- as the subsets of interior facets of \mathcal{F}_h such that the facets are in Ω_+ and Ω_- , respectively. We also define \mathcal{F}_h^{∂} as the facets on $\partial\Omega$, i.e., $\partial\Omega = \cup_{F \in \mathcal{F}_h^{\partial}} \bar{F}$. In summary, \mathcal{F}_h is a disjoint union of \mathcal{F}_h^{∂} , \mathcal{F}_h^+ , \mathcal{F}_h^- , \mathcal{F}_h^i . We also define \mathcal{F}_h^D as the subset of \mathcal{F}_h such that $\bar{\Gamma}_D = \cup_{F \in \mathcal{F}_h^D} \bar{F}$.

Let $K \in \mathcal{T}_h$, denote the diameter of K by h_K , and let $h = \max_{K \in \mathcal{T}_h} h_K$. We assume that there is no $K \in \mathcal{T}_h$ such that $\partial K \subset \Gamma_I$. For scalar functions $u, v \in L^2(K)$ and vector functions $\mathbf{q}, \mathbf{r} \in L^2(K; \mathbb{R}^d)$ we denote, respectively, $(u, v)_K := \int_K uv \, dx$ and $(\mathbf{q}, \mathbf{r})_K := \int_K \mathbf{q} \cdot \mathbf{r} \, dx$. Furthermore, for two functions \mathbf{r} and v with well-defined traces on ∂K , we define $\langle \mathbf{r} \cdot \mathbf{n}, v \rangle_{\partial K} := \int_{\partial K} \mathbf{r} \cdot \mathbf{n}_K v \, ds$, where \mathbf{n}_K is the unit outward normal vector field on ∂K . Additionally, we define $(u, v)_D := \sum_{K \subset D} (u, v)_K$, $(\mathbf{q}, \mathbf{r})_D := \sum_{K \subset D} (\mathbf{q}, \mathbf{r})_K$, and $\langle \mathbf{r} \cdot \mathbf{n}, v \rangle_{\partial \mathcal{T}_h} := \sum_{K \in \mathcal{T}_h} \langle \mathbf{r} \cdot \mathbf{n}_K, v \rangle_{\partial K}$, with similar definitions on \mathcal{T}_h^+ and \mathcal{T}_h^- .

We use the standard notation for Sobolev spaces based on the L^2 -norm, i.e., $H^s(D)$, $s \geq 0$, denotes the Sobolev space based on the L^2 -norm with s -differentiability on the domain D . We refer to [14] for a rigorous definition of this space. The norm on $H^s(D)$ is denoted by $\|\cdot\|_{s,D}$. When $s = 0$ we drop the subscript s .

To define the HDG method let

$$\mathbf{V}_h(K) = \mathcal{P}_k(K; \mathbb{R}^d), \quad W_h(K) = \mathcal{P}_k(K), \quad M_h(F) = \mathcal{P}_k(F),$$

where $\mathcal{P}_k(D)$ and $\mathcal{P}_k(D; \mathbb{R}^d)$ are the spaces of scalar and d -dimensional vector valued polynomials of degree at most k on a domain D . We will use the following finite element spaces:

$$(3.2) \quad \mathbf{V}_h = \{\mathbf{r} \in L^2(\mathcal{T}_h; \mathbb{R}^d) : \mathbf{r}|_K \in \mathbf{V}_h(K), \quad \forall K \in \mathcal{T}_h\},$$

$$(3.3) \quad W_h = \{v \in L^2(\mathcal{T}_h) : v|_K \in W_h(K), \quad \forall K \in \mathcal{T}_h\},$$

$$(3.4) \quad M_h = \{\mu \in L^2(\mathcal{F}_h) : \mu|_F \in M_h(F), \quad \forall F \in \mathcal{F}_h, \quad \mu|_{\mathcal{F}_h^D} = 0\}.$$

3.2. The discrete formulation. For a sufficiently regular solution (u, \mathbf{q}) of the mixed problem (3.1), and for \bar{u} the restriction of u on \mathcal{F}_h , we can derive a system of variational equations from (3.1) with test functions in $\mathbf{V}_h \times W_h \times M_h$ as

$$(3.5a) \quad (\sigma^{-1} \mathbf{q}, \mathbf{r})_\Omega + (\nabla u, \mathbf{r})_\Omega - \langle u - \bar{u}, \mathbf{r} \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h} = 0 \quad \mathbf{r} \in \mathbf{V}_h,$$

$$(3.5b) \quad (\mathbf{q}, \nabla v)_\Omega - \langle \mathbf{q} \cdot \mathbf{n} + \tau(u - \bar{u}), v \rangle_{\partial \mathcal{T}_h} = (f, v)_\Omega \quad v \in W_h,$$

$$(3.5c) \quad \langle \mathbf{q} \cdot \mathbf{n} + \tau(u - \bar{u}), \bar{v} \rangle_{\partial \mathcal{T}_h \setminus \mathcal{F}_h^D} = \langle u_N, \bar{v} \rangle_{\mathcal{F}_h^N} \quad \bar{v} \in M_h.$$

Here we use τ to denote a piecewise constant function adapted to \mathcal{F}_h . Throughout this paper we assume that

$$(3.6) \quad \tau|_F = \begin{cases} 0 & \text{if } F \subset \Gamma_I, \\ > 0 & \text{if } F \subset \partial K, K \subset \Omega_+, F \not\subset \Gamma_I, \\ < 0 & \text{if } F \subset \partial K, K \subset \Omega_-, F \not\subset \Gamma_I. \end{cases}$$

Our HDG method for (3.1) is the discrete counterpart of this system of variational equations, i.e., we find $(\mathbf{q}_h, u_h, \bar{u}_h) \in \mathbf{V}_h \times W_h \times M_h$ such that

$$(3.7a) \quad (\sigma^{-1} \mathbf{q}_h, \mathbf{r})_\Omega + (\nabla u_h, \mathbf{r})_\Omega - \langle u_h - \bar{u}_h, \mathbf{r} \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h} = - \langle u_D, \mathbf{r} \cdot \mathbf{n} \rangle_{\mathcal{F}_h^D} \quad \mathbf{r} \in \mathbf{V}_h,$$

$$(3.7b) \quad (\mathbf{q}_h, \nabla v)_\Omega - \langle \mathbf{q}_h \cdot \mathbf{n} + \tau(u_h - \bar{u}_h), v \rangle_{\partial \mathcal{T}_h} = - \langle \tau u_D, v \rangle_{\mathcal{F}_h^D} + (f, v)_\Omega \quad v \in W_h,$$

$$(3.7c) \quad \langle \mathbf{q}_h \cdot \mathbf{n} + \tau(u_h - \bar{u}_h), \bar{v} \rangle_{\partial \mathcal{T}_h} = \langle u_N, \bar{v} \rangle_{\mathcal{F}_h^N} \quad \bar{v} \in M_h.$$

Here equations (3.5a), (3.5b) and the discrete equations (3.7a), (3.7b) look inconsistent. This is because we impose the restriction $\mu|_{\mathcal{F}_h^D} = 0$ on M_h so that we can take the same trial and test function spaces in our numerical method. This choice will be useful to reveal a stabilized saddle point structure of our numerical method, and also allow us to obtain optimal error estimates without the Aubin–Nitsche duality argument. To clarify the consistency between (3.5a), (3.5b) and (3.7a), (3.7b), we point out that (3.7a), (3.7b) with $\bar{u}_h \in \tilde{M}_h$ are equivalent to the discretized forms of (3.5a), (3.5b) with \tilde{M}_h defined by

$$\tilde{M}_h =$$

$$\{\mu \in L^2(\mathcal{F}_h) : \mu|_F \in M_h(F) \quad \forall F \in \mathcal{F}_h, \quad \langle \mu, \lambda \rangle_{\mathcal{F}_h^D} = \langle u_D, \lambda \rangle_{\mathcal{F}_h^D} \quad \forall \lambda \in M_h(\mathcal{F}_h^D)\}.$$

For later use we define

$$(3.8) \quad \begin{aligned} a_h(\mathbf{q}, \mathbf{r}) &:= (\sigma^{-1} \mathbf{q}, \mathbf{r})_{\Omega}, \\ b_h(v, \bar{v}; \mathbf{r}) &:= (\nabla v, \mathbf{r})_{\Omega} - \langle v - \bar{v}, \mathbf{r} \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h}, \\ c_h(u, \bar{u}; v, \bar{v}) &:= \langle \tau(u - \bar{u}), v - \bar{v} \rangle_{\partial \mathcal{T}_h}. \end{aligned}$$

Then the sum of the left-hand sides of (3.7) can be rewritten as

$$(3.9) \quad B_h(\mathbf{q}_h, u_h, \bar{u}_h; \mathbf{r}, v, \bar{v}) := a_h(\mathbf{q}_h, \mathbf{r}) + b_h(u_h, \bar{u}_h; \mathbf{r}) + b_h(v, \bar{v}; \mathbf{q}_h) - c_h(u_h, \bar{u}_h; v, \bar{v}).$$

For brevity, but without loss of generality, we assume $u_D = u_N = 0$ and $\Gamma_D \neq \emptyset$ in the remainder of this paper.

In the following lemma we prove well-posedness of the HDG method (3.7).

LEMMA 3.1 (Well-posedness). *Let $k \geq 0$. If τ satisfies (3.6), then there exists a unique solution $(\mathbf{q}_h, u_h, \bar{u}_h) \in \mathbf{V}_h \times W_h \times M_h$ to (3.7).*

Proof. To show well-posedness of (3.7) assume that $f = 0$. We will show that $\mathbf{q}_h, u_h, \bar{u}_h$ vanish.

Let χ_D be the characteristic function that has the value 1 in domain D and 0 elsewhere. Then let $\mathbf{r} = \chi_{\Omega_+} \mathbf{q}_h - \chi_{\Omega_-} \mathbf{q}_h$, $v = -\chi_{\Omega_+} u_h + \chi_{\Omega_-} u_h$, $\bar{v} = \chi_{\Omega_+} \bar{u}_h - \chi_{\Omega_-} \bar{u}_h$ in (3.7), and add all the equations. (Note that $\bar{v} = 0$ on Γ_I .) The evaluation of (3.7) with the first components of these test functions gives

$$(\sigma^{-1} \mathbf{q}_h, \mathbf{q}_h)_{\Omega_+} + \langle \tau(u_h - \bar{u}_h), u_h - \bar{u}_h \rangle_{\partial \mathcal{T}_h^+} = 0.$$

Similarly, the evaluation of (3.7) with the second components of the above test functions gives

$$-(\sigma^{-1} \mathbf{q}_h, \mathbf{q}_h)_{\Omega_-} - \langle \tau(u_h - \bar{u}_h), u_h - \bar{u}_h \rangle_{\partial \mathcal{T}_h^-} = 0.$$

Since σ^{-1} and τ are positive and negative on Ω_+ and Ω_- , respectively, we can conclude that $\mathbf{q}_h = 0$ on Ω and $u_h = \bar{u}_h$ on $\mathcal{F}_h^+ \cup \mathcal{F}_h^- \cup \mathcal{F}_h^\partial$. This also implies that u_h is continuous on Ω_+ and on Ω_- , respectively. It then follows from (3.7a) that $\nabla u_h = 0$, i.e., u_h is constant on Ω_+ and on Ω_- , respectively. Since $\mathbf{q}_h = 0$, $\nabla u_h = 0$, and $u_D = 0$, (3.7a) gives $\langle u_h - \bar{u}_h, \mathbf{r} \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h} = 0$ for all $\mathbf{r} \in \mathbf{V}_h$, implying $u_h = \bar{u}_h$ also on Γ_I . Given now that u_h is constant on Ω , we obtain that $u_h = \bar{u}_h = 0$ because $u_h = \bar{u}_h = 0$ on Γ_D . \square

We consider the following norm and seminorm on \mathbf{V}_h and $W_h \times M_h$:

$$\begin{aligned} \|\mathbf{r}\|_{\mathbf{V}_h}^2 &= (\sigma^{-1} \mathbf{r}, \mathbf{r})_{\Omega_+} - (\sigma^{-1} \mathbf{r}, \mathbf{r})_{\Omega_-}, \\ |(v, \bar{v})|_{W_h \times M_h}^2 &= (v, v)_{\Omega} + \langle |\tau| (v - \bar{v}), v - \bar{v} \rangle_{\partial \mathcal{T}_h}, \end{aligned}$$

where $|\tau|$ is the absolute value of τ . Additionally, we define $\mathbf{X}_h := \mathbf{V}_h \times W_h \times M_h$ with seminorm

$$|(\mathbf{r}, v, \bar{v})|_{\mathbf{X}_h} := (\|\mathbf{r}\|_{\mathbf{V}_h}^2 + |(v, \bar{v})|_{W_h \times M_h}^2)^{\frac{1}{2}}.$$

Furthermore, in the remainder of this paper we let $C > 0$ be a constant independent of the mesh size h .

LEMMA 3.2. *There exists $C_{\Omega} > 0$ which depends only on Ω and Γ_D such that for $v \in W_h$ there exists $\mathbf{w} \in H^1(\Omega; \mathbb{R}^d)$ satisfying $\nabla \cdot \mathbf{w} = v$ and $\|\mathbf{w}\|_{1, \Omega} \leq C_{\Omega} \|v\|_{\Omega}$ with $\mathbf{w} \cdot \mathbf{n} = 0$ on Γ_N .*

Proof. We will use a result in [17, p.176] which claims that for any given $f \in L^2(\Omega)$ and $\mathbf{a} \in H^{\frac{1}{2}}(\partial\Omega)$ satisfying $\int_{\partial\Omega} \mathbf{a} \cdot \mathbf{n} \, ds = \int_{\Omega} f \, dx$, there exists $\mathbf{w} \in H^1(\Omega; \mathbb{R}^d)$ such that $\nabla \cdot \mathbf{w} = f$, $\mathbf{w} = \mathbf{a}$ on $\partial\Omega$, and $\|\mathbf{w}\|_{H^1(\Omega)} \leq C\|f\|_{L^2(\Omega)} + \|\mathbf{a}\|_{H^{\frac{1}{2}}(\partial\Omega)}$.

We first note that if $z \in L^2(\Omega)$ has mean value zero, then there exists $\mathbf{w} \in H_0^1(\Omega; \mathbb{R}^d)$ such that $\nabla \cdot \mathbf{w} = z$ and $\|\mathbf{w}\|_{H^1(\Omega)} \leq C\|z\|_{L^2(\Omega)}$ by taking $\mathbf{a} = \mathbf{0}$ in the above result.

Consider now a decomposition $v = v_0 + v_m$ with $v_m = \chi_{\Omega} \frac{1}{|\Omega|} \int_{\Omega} v \, dx$ where $|\Omega|$ is the Lebesgue measure of Ω , and where v_0 is the component of v with mean-value zero. It is clear that $\|v_0\|_{L^2(\Omega)} + \|v_m\|_{L^2(\Omega)} \leq C\|v\|_{L^2(\Omega)}$. As just noted, there exists $\mathbf{w}_0 \in H_0^1(\Omega; \mathbb{R}^d)$ such that $\nabla \cdot \mathbf{w}_0 = v_0$ and $\|\mathbf{w}_0\|_{H^1(\Omega)} \leq C\|v_0\|_{L^2(\Omega)}$. Since Ω is a polygonal/polyhedral domain, one can find a function $\phi \in H^{1/2}(\partial\Omega)$ whose support resides on Γ_D , $\int_{\Gamma_D} \phi \cdot \mathbf{n} \, ds = c_0 > 0$, and $\|\phi\|_{H^{\frac{1}{2}}(\partial\Omega)} = 1$. We can find ϕ_0 , a renormalization of ϕ , satisfying $\int_{\Gamma_D} \phi_0 \cdot \mathbf{n} \, ds = \int_{\Omega} v_m \, dx$, and it holds that

$$\|\phi_0\|_{H^{\frac{1}{2}}(\partial\Omega)} \leq \frac{\|v_m\|_{L^1(\Omega)}}{c_0} \leq C\|v_m\|_{L^2(\Omega)},$$

with a constant that depends on $|\Omega|$ and c_0 . Applying the result in [17, p.176] there exists $\mathbf{w}_1 \in H^1(\Omega; \mathbb{R}^d)$ such that $\nabla \cdot \mathbf{w}_1 = v_m$, $\mathbf{w}_1 = \phi_0$ on $\partial\Omega$, and $\|\mathbf{w}_1\|_{H^1(\Omega)} \leq C\|\phi_0\|_{H^{1/2}(\partial\Omega)} + \|v_m\|_{L^2(\Omega)} \leq C\|v_m\|_{L^2(\Omega)}$. Then $\mathbf{w} = \mathbf{w}_0 + \mathbf{w}_1$ is a function in $H^1(\Omega; \mathbb{R}^d)$ satisfying the desired conditions. \square

Before we prove an inf-sup condition, that we use in the error analysis in section 4, let us recall a result from [9, Proposition 2.1]: For a simplex K and a fixed facet F_K of K , there is an interpolation operator $\Pi'_h : H^1(K; \mathbb{R}^d) \rightarrow \mathbf{V}_h(K)$ such that

$$\begin{aligned} (\Pi'_h \mathbf{r}, \mathbf{r}')_K &= (\mathbf{r}, \mathbf{r}')_K & \forall \mathbf{r}' \in \mathcal{P}_{k-1}(K; \mathbb{R}^d) \quad \text{if } k > 0, \\ \langle \Pi'_h \mathbf{r} \cdot \mathbf{n}, w \rangle_F &= \langle \mathbf{r} \cdot \mathbf{n}, w \rangle_F & \forall w \in \mathcal{P}_k(F) \quad \text{if } F \neq F_K. \end{aligned}$$

This interpolation operator Π'_h also satisfies $\|(\mathbf{r} - \Pi'_h \mathbf{r}) \cdot \mathbf{n}\|_{\partial K} \leq Ch_K^{1/2} \|\mathbf{r}\|_{H^1(K)}$.

THEOREM 3.3 (inf-sup condition). *Suppose that $k \geq 0$. If τ satisfies (3.6) and $\gamma_0 \leq |\tau| \leq \gamma_1$ for $\tau \neq 0$ on \mathcal{F}_h with positive constants γ_0 and γ_1 independent of h , then there exists a positive constant β independent of h such that*

$$(3.10) \quad \inf_{(\mathbf{p}, z, \bar{z}) \in \mathbf{X}_h} \sup_{(\mathbf{r}, v, \bar{v}) \in \mathbf{X}_h} \frac{B_h(\mathbf{p}, z, \bar{z}; \mathbf{r}, v, \bar{v})}{|(\mathbf{p}, z, \bar{z})|_{\mathbf{X}_h} |(\mathbf{r}, v, \bar{v})|_{\mathbf{X}_h}} \geq \beta > 0.$$

Here we assume that $|(\mathbf{p}, z, \bar{z})|_{\mathbf{X}_h}, |(\mathbf{r}, v, \bar{v})|_{\mathbf{X}_h} > 0$.

Proof. We first prove the following weak inf-sup condition: There exist $C_0, C_1 > 0$ independent of h such that

$$(3.11) \quad \inf_{(z, \bar{z}) \in W_h \times M_h} \sup_{\mathbf{r} \in \mathbf{V}_h} \frac{b_h(z, \bar{z}; \mathbf{r})}{\|\mathbf{r}\|_{\mathbf{V}_h}} \geq C_0 \|z\|_{\Omega} - C_1 \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial\mathcal{T}_h}^{\frac{1}{2}}.$$

We show this by proving an equivalent condition, i.e., there exist $C'_0, C'_1, C'_2 > 0$ independent of h such that for any $0 \neq (z, \bar{z}) \in W_h \times M_h$ there exists $\mathbf{r} \in \mathbf{V}_h$ such that $\|\mathbf{r}\|_{\mathbf{V}_h} \leq C'_2 \|z\|_{\Omega}$ and

$$(3.12) \quad b_h(z, \bar{z}; \mathbf{r}) \geq C'_0 \|z\|_{\Omega}^2 - C'_1 \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial\mathcal{T}_h}^{\frac{1}{2}} \|z\|_{\Omega}.$$

By Lemma 3.2 there exists $\mathbf{w} \in H^1(\Omega; \mathbb{R}^d)$ with $\mathbf{w} \cdot \mathbf{n}|_{\Gamma_N} = 0$ such that $\nabla \cdot \mathbf{w} = -z$ and $\|\mathbf{w}\|_{1,\Omega} \leq C_\Omega \|z\|_\Omega$ with C_Ω depending only on Ω and Γ_D . We define $\Pi_h \mathbf{w}$ on K as follows: If $\partial K \cap \Gamma_I = \emptyset$, then $\Pi_h \mathbf{w}$ is the L^2 orthogonal projection of \mathbf{w} into \mathbf{V}_h . If $\partial K \cap \Gamma_I \neq \emptyset$, then $\Pi_h \mathbf{w} = \Pi'_h \mathbf{w}$ with the interpolation Π'_h associated to a facet $F \subset \partial K$, $F \not\subset \Gamma_I$. It then holds that

$$(3.13) \quad \begin{aligned} (\mathbf{w}, \mathbf{r})_K &= (\Pi_h \mathbf{w}, \mathbf{r})_K & \forall \mathbf{r} \in \mathcal{P}_{k-1}(K; \mathbb{R}^d) \text{ for all } K \in \mathcal{T}_h, \\ \langle \mathbf{w} \cdot \mathbf{n}, w \rangle_F &= \langle \Pi_h \mathbf{w} \cdot \mathbf{n}, w \rangle_F & \forall w \in \mathcal{P}_k(F) \text{ for all } F \in \mathcal{F}_h^i. \end{aligned}$$

Then

$$(3.14) \quad \begin{aligned} -\|z\|_\Omega^2 &= (z, \nabla \cdot \mathbf{w})_\Omega \\ &= -(\nabla z, \mathbf{w})_\Omega + \langle z, \mathbf{w} \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h} \\ &= -(\nabla z, \Pi_h \mathbf{w})_\Omega + \langle z - \bar{z}, \mathbf{w} \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h} \\ &= -b_h(z, \bar{z}; \Pi_h \mathbf{w}) + \langle z - \bar{z}, (\mathbf{w} - \Pi_h \mathbf{w}) \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h \setminus \Gamma_I}, \end{aligned}$$

where the third equality holds because $\mathbf{w} \cdot \mathbf{n}$ and \bar{z} are single-valued on facets, $\bar{z} = 0$ on Γ_D by the definition of M_h , and $\mathbf{w} \cdot \mathbf{n} = 0$ on Γ_N , and the fourth equality holds because of (3.13).

The element-wise trace inequality and the Cauchy-Schwarz inequality give

$$(3.15) \quad \begin{aligned} \langle z - \bar{z}, (\mathbf{w} - \Pi_h \mathbf{w}) \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h \setminus \Gamma_I} &\geq -C \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial \mathcal{T}_h}^{\frac{1}{2}} \|\mathbf{w}\|_{1,\Omega} \\ &\geq -C' \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial \mathcal{T}_h}^{\frac{1}{2}} \|z\|_\Omega. \end{aligned}$$

Combining (3.14) and (3.15) we obtain

$$(3.16) \quad b_h(z, \bar{z}; \Pi_h \mathbf{w}) \geq \|z\|_\Omega^2 - C' \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial \mathcal{T}_h}^{\frac{1}{2}} \|z\|_\Omega.$$

Moreover,

$$(3.17) \quad \|\Pi_h \mathbf{w}\|_\Omega \leq C \|\mathbf{w}\|_{1,\Omega} \leq C_\Omega \|z\|_\Omega$$

holds, so the weak inf-sup condition (3.12) follows.

To prove (3.10) suppose that $\mathbf{r}_0 \in \mathbf{V}_h$ is an element satisfying (3.12) for given z . For given $(\mathbf{p}, z, \bar{z}) \in \mathbf{X}_h$ we take $\mathbf{r} = \chi_{\Omega_+} \mathbf{p} - \chi_{\Omega_-} \mathbf{p} + \varepsilon \mathbf{r}_0$, $v = -\chi_{\Omega_+} z + \chi_{\Omega_-} z$, $\bar{v} = -\chi_{\Omega_+} \bar{z} + \chi_{\Omega_-} \bar{z}$ in (3.9), with $\varepsilon > 0$ to be determined later. Then by (3.12) and Young's inequality,

$$\begin{aligned} B_h(\mathbf{p}, z, \bar{z}; \mathbf{r}, v, \bar{v}) &= \|\mathbf{p}\|_{\mathbf{V}_h}^2 + \varepsilon (\sigma^{-1} \mathbf{p}, \mathbf{r}_0)_\Omega + \varepsilon b_h(z, \bar{z}; \mathbf{r}_0) \\ &\quad + \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial \mathcal{T}_h} \\ &\geq \|\mathbf{p}\|_{\mathbf{V}_h}^2 - \frac{1}{2} \|\mathbf{p}\|_{\mathbf{V}_h}^2 - \frac{1}{2} \varepsilon^2 (C'_2)^2 \|z\|_\Omega^2 \\ &\quad + \varepsilon C'_0 \|z\|_\Omega^2 - \varepsilon C'_1 \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial \mathcal{T}_h}^{\frac{1}{2}} \|z\|_\Omega \\ &\quad + \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial \mathcal{T}_h} \\ &\geq \|\mathbf{p}\|_{\mathbf{V}_h}^2 - \frac{1}{2} \|\mathbf{p}\|_{\mathbf{V}_h}^2 - \frac{1}{2} \varepsilon^2 (C'_2)^2 \|z\|_\Omega^2 \\ &\quad + \varepsilon C'_0 \|z\|_\Omega^2 - \frac{1}{2} \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial \mathcal{T}_h} \\ &\quad - \frac{1}{2} \varepsilon^2 (C'_1)^2 \|z\|_\Omega^2 + \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial \mathcal{T}_h}. \end{aligned}$$

Choosing ε sufficiently small, we obtain

$$B_h(\mathbf{p}, z, \bar{z}; \mathbf{r}, v, \bar{v}) \geq C (\|\mathbf{p}\|_{\mathbf{V}_h}^2 + \|z\|_{\Omega}^2 + \langle |\tau|(z - \bar{z}), z - \bar{z} \rangle_{\partial\mathcal{T}_h}).$$

Finally, it is not difficult to show, by the choice of (\mathbf{r}, v, \bar{v}) , that

$$|(\mathbf{r}, v, \bar{v})|_{\mathbf{X}_h} \leq C |(\mathbf{p}, z, \bar{z})|_{\mathbf{X}_h}.$$

This proves (3.10). \square

4. Error analysis. In this section we present the *a priori* error estimates of our HDG method. For this we recall an interpolation result (cf. [11, Theorem 2.1]): For $K \in \mathcal{T}_h$, $\tau|_{\partial K}$ is a piecewise constant function which is nonnegative if $K \subset \Omega_+$ and nonpositive if $K \subset \Omega_-$ such that $\tau|_{\partial K} \neq 0$. Then there exists a bounded interpolation operator $\Pi := (\Pi_{\mathbf{V}}, \Pi_W)$ from $H^1(K; \mathbb{R}^d) \times H^1(K)$ to $\mathbf{V}_h(K) \times W_h(K)$ such that

$$(4.1a) \quad (\Pi_{\mathbf{V}} \mathbf{r}, \tilde{\mathbf{r}})_K = (\mathbf{r}, \tilde{\mathbf{r}})_K \quad \forall \tilde{\mathbf{r}} \in [\mathcal{P}_{k-1}(K)]^d,$$

$$(4.1b) \quad (\Pi_W v, \tilde{v})_K = (v, \tilde{v})_K \quad \forall \tilde{v} \in \mathcal{P}_{k-1}(K),$$

$$(4.1c) \quad \langle \Pi_{\mathbf{V}} \mathbf{r} \cdot \mathbf{n} + \tau \Pi_W v, \lambda \rangle_F = \langle \mathbf{r} \cdot \mathbf{n} + \tau v, \lambda \rangle_F \quad \forall \lambda \in \mathcal{P}_k(F),$$

for $F \subset \partial K$ and $(\mathbf{r}, v) \in H^1(K; \mathbb{R}^d) \times H^1(K)$, and

$$(4.2) \quad \|\mathbf{r} - \Pi_{\mathbf{V}} \mathbf{r}\|_K \leq Ch_K^{k_r+1} |\mathbf{r}|_{k_r+1, K} + h_K^{k_v+1} \tau_K^* |v|_{k_v+1, K},$$

$$(4.3) \quad \|v - \Pi_W v\|_K \leq Ch_K^{k_v+1} |v|_{k_v+1, K} + \frac{h_K^{k_r+1}}{\tau_K^{\max}} |\nabla \cdot \mathbf{r}|_{k_r, K},$$

hold with $0 \leq k_v, k_r \leq k$. Here,

$$\tau_K^{\max} = \begin{cases} \max_{F \subset \partial K} \tau|_F & \text{if } \tau \geq 0, \\ \max_{F \subset \partial K} (-\tau|_F) & \text{if } \tau \leq 0, \end{cases}$$

$$\tau_K^* = \begin{cases} \max_{F \subset \partial K \setminus F^*} \tau|_F & \text{if } \tau \geq 0, \\ \max_{F \subset \partial K \setminus F^*} (-\tau|_F) & \text{if } \tau \leq 0, \end{cases}$$

where F^* is the face where τ_K^{\max} is attained, and the implicit constants are independent of K and τ . We remark that the proof in [11] is only for nonnegative τ . The proof, however, can easily be adapted to nonpositive τ , so we omit its proof here. We will also require P_M , the facet-wise L^2 projection from $L^2(\mathcal{F}_h \setminus \mathcal{F}_h^D)$ to M_h .

For brevity of notation, we will denote the difference between an unknown ϕ and its approximation ϕ_h by $e_\phi := \phi - \phi_h$. It will be convenient to also split the error into interpolation and approximation errors:

$$(4.4) \quad \begin{aligned} e_{\mathbf{q}} &= e_{\mathbf{q}}^I + e_{\mathbf{q}}^h := (\mathbf{q} - \Pi_{\mathbf{V}} \mathbf{q}) + (\Pi_{\mathbf{V}} \mathbf{q} - \mathbf{q}_h), \\ e_u &= e_u^I + e_u^h := (u - \Pi_W u) + (\Pi_W u - u_h), \\ e_{\bar{u}} &= e_{\bar{u}}^I + e_{\bar{u}}^h := (\bar{u} - P_M \bar{u}) + (P_M \bar{u} - \bar{u}_h). \end{aligned}$$

The error equations are obtained by subtracting (3.7) from (3.5) and integration by parts,

$$(4.5a) \quad (\sigma^{-1} e_{\mathbf{q}}, \mathbf{r})_{\Omega} - (e_u, \nabla \cdot \mathbf{r})_{\Omega} + \langle e_{\bar{u}}, \mathbf{r} \cdot \mathbf{n} \rangle_{\partial\mathcal{T}_h} = 0 \quad \mathbf{r} \in \mathbf{V}_h,$$

$$(4.5b) \quad (e_{\mathbf{q}}, \nabla v)_{\Omega} - \langle e_{\mathbf{q}} \cdot \mathbf{n} + \tau(e_u - e_{\bar{u}}), v \rangle_{\partial\mathcal{T}_h} = 0 \quad v \in W_h,$$

$$(4.5c) \quad \langle e_{\mathbf{q}} \cdot \mathbf{n} + \tau(e_u - e_{\bar{u}}), \bar{v} \rangle_{\partial\mathcal{T}_h \setminus \mathcal{F}_h^D} = 0 \quad \bar{v} \in M_h.$$

THEOREM 4.1 (A priori error estimates). *Let $(\mathbf{q}, u) \in H^s(\Omega; \mathbb{R}^d) \times H^s(\Omega)$, $s \geq 1$ be the solution of (3.5) and let \bar{u} be the trace of u on $\mathcal{F}_h \setminus \mathcal{F}_h^D$ and $\bar{u} = 0$ on \mathcal{F}_h^D . We assume that τ satisfies the same conditions as in Theorem 3.3, and let $(\mathbf{q}_h, u_h, \bar{u}_h) \in \mathbf{V}_h \times W_h \times M_h$ solve (3.7). Then,*

$$(4.6a) \quad \|\mathbf{q} - \mathbf{q}_h\|_{\mathbf{V}_h} \leq Ch^s (\|\mathbf{q}\|_{H^s(\Omega)} + \|u\|_{H^s(\Omega)}), \quad 1 \leq s \leq k+1,$$

$$(4.6b) \quad \|u - u_h\|_{\Omega} \leq Ch^s (\|\mathbf{q}\|_{H^s(\Omega)} + \|u\|_{H^s(\Omega)}), \quad 1 \leq s \leq k+1,$$

$$(4.6c) \quad \langle P_M u - \bar{u}_h, P_M u - \bar{u}_h \rangle_{\partial \mathcal{T}_h \setminus \mathcal{F}_h^D}^{\frac{1}{2}} \leq Ch^{s-\frac{1}{2}} (\|\mathbf{q}\|_{H^s(\Omega)} + \|u\|_{H^s(\Omega)}), \quad 1 \leq s \leq k+1.$$

Proof. By definition of the interpolation operator (4.1), the error equations (4.5) can be reduced to

$$(4.7a) \quad (\sigma^{-1} e_{\mathbf{q}}^h, \mathbf{r})_{\Omega} - (e_u^h, \nabla \cdot \mathbf{r})_{\Omega} + \langle e_{\bar{u}}^h, \mathbf{r} \cdot \mathbf{n} \rangle_{\partial \mathcal{T}_h} = -(\sigma^{-1} e_{\mathbf{q}}^I, \mathbf{r})_{\Omega} \quad \mathbf{r} \in \mathbf{V}_h,$$

$$(4.7b) \quad (e_{\mathbf{q}}^h, \nabla v)_{\Omega} - \langle e_{\mathbf{q}}^h \cdot \mathbf{n} + \tau(e_u^h - e_{\bar{u}}^h), v \rangle_{\partial \mathcal{T}_h} = 0 \quad v \in W_h,$$

$$(4.7c) \quad \langle e_{\mathbf{q}}^h \cdot \mathbf{n} + \tau(e_u^h - e_{\bar{u}}^h), \bar{v} \rangle_{\partial \mathcal{T}_h \setminus \mathcal{F}_h^D} = 0 \quad \bar{v} \in M_h.$$

To show (4.6a), take $\mathbf{r} = \chi_{\Omega_+} e_{\mathbf{q}}^h - \chi_{\Omega_-} e_{\mathbf{q}}^h$, $v = -\chi_{\Omega_+} e_u^h + \chi_{\Omega_-} e_u^h$, $\bar{v} = \chi_{\Omega_+} e_{\bar{u}}^h - \chi_{\Omega_-} e_{\bar{u}}^h$ in (4.7), and add all the equations. We find

$$\begin{aligned} (|\sigma|^{-1} e_{\mathbf{q}}^h, e_{\mathbf{q}}^h)_{\Omega} + \langle \tau | (e_u^h - e_{\bar{u}}^h), e_u^h - e_{\bar{u}}^h \rangle_{\partial \mathcal{T}_h} &= -(\sigma^{-1} e_{\mathbf{q}}^I, \mathbf{r})_{\Omega} \\ &\leq (|\sigma|^{-1} e_{\mathbf{q}}^I, e_{\mathbf{q}}^I)_{\Omega}^{\frac{1}{2}} (|\sigma|^{-1} e_{\mathbf{q}}^h, e_{\mathbf{q}}^h)_{\Omega}^{\frac{1}{2}}. \end{aligned}$$

By this inequality, the triangle inequality, and (4.2), one can obtain (4.6a).

To show (4.6b), we note a result from Theorem 3.3: There exists $(\mathbf{r}, v, \bar{v}) \in \mathbf{X}_h$ such that $B_h(e_{\mathbf{q}}^h, e_u^h, e_{\bar{u}}^h; \mathbf{r}, v, \bar{v}) \geq |(e_{\mathbf{q}}^h, e_u^h, e_{\bar{u}}^h)|_{\mathbf{X}_h}$ with $|\mathbf{r}, v, \bar{v}|_{\mathbf{X}_h} \leq C$. Together with (4.7) and (4.2) this results in

$$|(e_{\mathbf{q}}^h, e_u^h, e_{\bar{u}}^h)|_{\mathbf{X}_h} \leq C \|e_{\mathbf{q}}^I\|_{\mathbf{V}_h} \leq Ch^s (\|\mathbf{q}\|_{H^s(\Omega)} + \|u\|_{H^s(\Omega)}), \quad 1 \leq s \leq k+1.$$

Then (4.6b) follows by the triangle inequality.

We next show (4.6c). For each $K \in \mathcal{T}_h$ there exists $\mathbf{r} \in \mathbf{V}_h(K)$ such that $\mathbf{r} \cdot \mathbf{n} = e_{\bar{u}}^h$ on ∂K and $\|\mathbf{r}\|_K \leq Ch_K^{\frac{1}{2}} \|e_{\bar{u}}^h\|_{\partial K}$ with C independent of K . Taking this \mathbf{r} in (4.7a), we obtain

$$\begin{aligned} \langle e_{\bar{u}}^h, e_{\bar{u}}^h \rangle_{\partial \mathcal{T}_h} &= |-(\sigma^{-1} e_{\mathbf{q}}^I, \mathbf{r})_{\Omega} - (e_u^h, \nabla \cdot \mathbf{r})_{\Omega}| \\ &\leq C \|e_{\mathbf{q}}^I\|_{\Omega} \|\mathbf{r}\|_{\Omega} + \|e_u^h\|_{\Omega} \|\nabla \cdot \mathbf{r}\|_{\Omega} \\ &\leq C \|e_{\mathbf{q}}^I\|_{\Omega} h^{\frac{1}{2}} \langle e_{\bar{u}}^h, e_{\bar{u}}^h \rangle_{\partial \mathcal{T}_h}^{\frac{1}{2}} + Ch^{-\frac{1}{2}} \|e_u^h\|_{\Omega} \langle e_{\bar{u}}^h, e_{\bar{u}}^h \rangle_{\partial \mathcal{T}_h}^{\frac{1}{2}} \end{aligned}$$

where we used an inverse inequality and the properties of the chosen \mathbf{r} in the last inequality. Then (4.6c) follows by (4.6a) and (4.6b). \square

REMARK 4.1. *We remark that the proof of (4.6a) is not new. However, to the best of our knowledge, the proofs of the estimates (4.6b) and (4.6c) for sign-changing stabilization parameters without a duality argument have not been reported in the literature.*

5. Superconvergence by a duality argument. In this section we discuss superconvergence of u_h by a duality argument. For $\theta \in L^2(\Omega)$, let us consider a dual problem,

$$\begin{aligned} (5.1a) \quad & \sigma^{-1} \Phi + \nabla \phi = 0 && \text{in } \Omega, \\ (5.1b) \quad & \nabla \cdot \Phi = \theta && \text{in } \Omega, \\ (5.1c) \quad & \phi = 0 && \text{on } \partial\Omega. \end{aligned}$$

We assume that the solution (Φ, ϕ) satisfies the regularity assumption

$$(5.2) \quad \|\Phi\|_{1,\Omega_+} + \|\Phi\|_{1,\Omega_-} + \|\phi\|_{2,\Omega_+} + \|\phi\|_{2,\Omega_-} \leq C_{\text{reg}} \|\theta\|_{\Omega}.$$

This assumption holds, for example, when both Ω_+ and Ω_- are convex (cf. [7, Proposition 2]).

The following lemma is an analogue of Lemma 4.2 in [11]. The proof is the same as in [11], but we include a detailed proof here to show that the proof still holds with sign-changing coefficients.

THEOREM 5.1 (duality argument). *Suppose that $(\mathbf{q}, u) \in H^s(\Omega; \mathbb{R}^d) \times H^s(\Omega)$, $s \geq 1$, and $(\mathbf{q}_h, u_h, \bar{u}_h) \in \mathbf{V}_h \times W_h \times M_h$ are the solutions of (3.5) and (3.7), respectively. Let e_u^h be as defined in (4.4). If (5.2) holds for the solution (Φ, ϕ) of (5.1), then,*

$$(5.3) \quad (e_u^h, \theta)_{\Omega} = (\sigma^{-1}(\mathbf{q} - \mathbf{q}_h), \Pi_{\mathbf{V}} \Phi - \Phi)_{\Omega} + (\Pi_{\mathbf{V}} \mathbf{q} - \mathbf{q}, \nabla(\phi - \phi_h))_{\Omega}$$

for any $\phi_h \in W_h$. Additionally, the following superconvergence result holds:

$$(5.4) \quad \|\Pi_W u - u_h\|_{\Omega} \leq C C_{\text{reg}} h (\|\mathbf{q} - \mathbf{q}_h\|_{\mathbf{V}_h} + \|\mathbf{q} - \Pi_{\mathbf{V}} \mathbf{q}\|_{\Omega})$$

if $k \geq 1$.

Proof. In this proof we abbreviate ‘integration by parts’ by IBP. We will furthermore use (\cdot, \cdot) and $\langle \cdot, \cdot \rangle$ instead of $(\cdot, \cdot)_{\Omega}$ and $\langle \cdot, \cdot \rangle_{\partial\mathcal{T}_h}$ for brevity of notation.

We first note that

$$(5.5) \quad \langle e_u^h, \Phi \cdot \mathbf{n} \rangle = 0.$$

From (5.1) one can obtain

$$\begin{aligned} (5.6) \quad (e_u^h, \theta) &= (e_u^h, \nabla \cdot \Phi) \\ &= -(\nabla e_u^h, \Phi) + \langle e_u^h, \Phi \cdot \mathbf{n} \rangle && \text{by IBP} \\ &= -(\nabla e_u^h, \Pi_{\mathbf{V}} \Phi) + \langle e_u^h, \Phi \cdot \mathbf{n} \rangle && \text{by (4.1a)} \\ &= (e_u^h, \nabla \cdot \Pi_{\mathbf{V}} \Phi) + \langle e_u^h, (\Phi - \Pi_{\mathbf{V}} \Phi) \cdot \mathbf{n} \rangle && \text{by IBP} \\ &= (e_u^h, \nabla \cdot \Pi_{\mathbf{V}} \Phi) - \langle e_u^h, \tau(\phi - \Pi_W \phi) \cdot \mathbf{n} \rangle && \text{by (4.1c)} \\ &= (\sigma^{-1} e_{\mathbf{q}}, \Pi_{\mathbf{V}} \Phi) + \langle e_u^h, \Pi_{\mathbf{V}} \Phi \cdot \mathbf{n} \rangle - \langle e_u^h, \tau(\phi - \Pi_W \phi) \cdot \mathbf{n} \rangle && \text{by (4.7a)} \\ &= (\sigma^{-1} e_{\mathbf{q}}, \Pi_{\mathbf{V}} \Phi) + \langle e_u^h, (\Pi_{\mathbf{V}} \Phi - \Phi) \cdot \mathbf{n} \rangle - \langle e_u^h, \tau(\phi - \Pi_W \phi) \cdot \mathbf{n} \rangle && \text{by (5.5)} \\ &= (\sigma^{-1} e_{\mathbf{q}}, \Pi_{\mathbf{V}} \Phi) - \langle e_u^h - e_{\bar{u}}^h, \tau(\phi - \Pi_W \phi) \rangle && \text{by (4.1c)} \\ &= (\sigma^{-1} e_{\mathbf{q}}, \Pi_{\mathbf{V}} \Phi) - \langle e_u^h - e_{\bar{u}}^h, \tau P_M \phi \rangle + \langle e_u^h - e_{\bar{u}}^h, \tau \Pi_W \phi \rangle \end{aligned}$$

where we used the property of the L^2 projection P_M in the last identity.

We next note that by (4.7c) and the property of P_M ,

$$\langle e_u^h - e_{\bar{u}}^h, \tau P_M \phi \rangle = -\langle e_q^h \cdot \mathbf{n}, P_M \phi \rangle = -\langle e_q^h \cdot \mathbf{n}, \phi \rangle.$$

We also have

$$\begin{aligned} \langle \tau (e_u^h - e_{\bar{u}}^h), \Pi_W \phi \rangle &= (e_q^h, \nabla \Pi_W \phi) - \langle e_q^h \cdot \mathbf{n}, \Pi_W \phi \rangle && \text{by (4.7b)} \\ &= -(\nabla \cdot e_q^h, \Pi_W \phi) && \text{by IBP} \\ &= -(\nabla \cdot e_q^h, \phi) && \text{by (4.1a)} \\ &= (e_q^h, \nabla \phi) - \langle e_q^h \cdot \mathbf{n}, \phi \rangle. && \text{by IBP} \end{aligned}$$

These two identities combined with (5.6) result in

$$(5.7) \quad (e_u^h, \theta) = (\sigma^{-1} e_q, \Pi_V \Phi) + (e_q^h, \nabla \phi).$$

Note that $(\sigma^{-1} e_q, \Phi) + (e_q, \nabla \phi) = 0$ by (5.1a). Using this identity and (4.1a), we may write (5.7) as

$$\begin{aligned} (e_u^h, \theta) &= (\sigma^{-1} e_q, \Pi_V \Phi - \Phi) + (e_q^h - e_q, \nabla \phi) \\ &= (\sigma^{-1} (\mathbf{q} - \mathbf{q}_h), \Pi_V \Phi - \Phi) + (\Pi_V \mathbf{q} - \mathbf{q}, \nabla (\phi - \phi_h)), \end{aligned}$$

for any $\phi_h \in W_h$. This completes the proof of (5.3).

To see (5.4), choose $\theta = e_u^h := \Pi_W u - u_h$. Furthermore, since $k \geq 1$ and ϕ_h is arbitrary, $\|\nabla (\phi - \phi_h)\|_\Omega \leq Ch \|\phi\|_{2,\Omega}$. Now, applying the Cauchy-Schwarz inequality to (5.3) gives

$$\begin{aligned} \|\Pi_W u - u_h\|_\Omega^2 &\leq Ch (\|\mathbf{q} - \mathbf{q}_h\|_{\mathbf{V}_h} \|\Phi\|_{1,\Omega} + \|\mathbf{q} - \Pi_V \mathbf{q}\|_\Omega \|\phi\|_{2,\Omega}) \\ &\leq CC_{\text{reg}} h \|\Pi_W u - u_h\|_\Omega (\|\mathbf{q} - \mathbf{q}_h\|_{\mathbf{V}_h} + \|\mathbf{q} - \Pi_V \mathbf{q}\|_\Omega), \end{aligned}$$

where we used (5.2) in the last inequality. The conclusion follows by dividing both sides by $\|\Pi_W u - u_h\|_\Omega$. \square

Following [20], we define a post-processed solution of u_h, u_h^* , by

$$(5.8a) \quad (\nabla u_h^*, \nabla v)_K = -(\sigma^{-1} \mathbf{q}_h, \nabla v)_K, \quad v \in \mathcal{P}_{k+1}(K),$$

$$(5.8b) \quad (u_h^*, 1)_K = (u_h, 1)_K,$$

for all $K \in \mathcal{T}_h$. The next result shows that the error in the post-processed solution u_h^* superconverges. Although the proof is standard, we include it here for completeness.

THEOREM 5.2 (superconvergence). *Suppose that $(\mathbf{q}, u) \in H^k(\Omega; \mathbb{R}^d) \times H^k(\Omega)$, $k \geq 1$, and $(\mathbf{q}_h, u_h, \bar{u}_h) \in \mathbf{V}_h \times W_h \times M_h$ are the solutions of (3.5) and (3.7), respectively. Let u_h^* be the post-processed solution defined by (5.8). The following result holds:*

$$\|u - u_h^*\|_\Omega \leq Ch (\|\mathbf{q} - \mathbf{q}_h\|_{\mathbf{V}_h} + \|\mathbf{q} - \Pi_V \mathbf{q}\|_\Omega) + Ch^{k+2} \|u\|_{k+2,\Omega}.$$

Proof. Let $\mathcal{P}_m(\mathcal{T}_h) := \{v \in L^2(\Omega) : v|_K \in \mathcal{P}_m(K) \forall K \in \mathcal{T}_h\}$. Let $\Pi : H^1(\Omega) \rightarrow \mathcal{P}_{k+1}(\mathcal{T}_h)$ be an interpolation such that $P_0 \Pi = P_0$ and $\|v - \Pi v\|_\Omega + h \|\nabla v - \nabla \Pi v\|_\Omega \leq Ch^{k+2} \|v\|_{k+2,\Omega}$, where P_0 is the L^2 projection to $\mathcal{P}_0(\mathcal{T}_h)$. By the triangle inequality,

it suffices to estimate $\|\Pi u - u_h^*\|_\Omega$. Again by the triangle inequality, it is sufficient to estimate $\|P_0(\Pi u - u_h^*)\|_\Omega$ and $\|(I - P_0)(\Pi u - u_h^*)\|_\Omega$.

Since $P_0(\Pi u - u_h^*) = P_0 u - P_0 u_h = P_0(\Pi_W u - u_h)$ by (5.8b) and the definition of Π_W , we find $\|P_0(\Pi u - u_h^*)\|_\Omega \leq Ch(\|\mathbf{q} - \mathbf{q}_h\|_{\mathbf{V}_h} + \|\mathbf{q} - \Pi \mathbf{V} \mathbf{q}\|_\Omega)$ by Theorem 5.1.

To estimate $\|(I - P_0)(\Pi u - u_h^*)\|_\Omega$, recall that $\mathbf{q} = -\sigma \nabla u$. By (5.8a) we have

$$(\nabla(u - u_h^*), \nabla v)_\Omega = -(\sigma^{-1}(\mathbf{q} - \mathbf{q}_h), \nabla v)_\Omega, \quad v \in \mathcal{P}_{k+1}(\mathcal{T}_h).$$

Taking $v = (I - P_0)(\Pi u - u_h^*)$, we obtain

$$\begin{aligned} \|\nabla(I - P_0)(\Pi u - u_h^*)\|_\Omega^2 &= -(\nabla(u - \Pi u), \nabla(I - P_0)(\Pi u - u_h^*))_\Omega \\ &\quad - (\sigma^{-1}(\mathbf{q} - \mathbf{q}_h), \nabla(I - P_0)(\Pi u - u_h^*))_\Omega. \end{aligned}$$

The Cauchy–Schwarz inequality and the assumption on Π gives

$$\begin{aligned} \|\nabla(I - P_0)(\Pi u - u_h^*)\|_\Omega &\leq \|\nabla(u - \Pi u)\|_\Omega + \|\sigma^{-1}(\mathbf{q} - \mathbf{q}_h)\|_\Omega \\ &\leq h^{k+1}\|u\|_{k+2,\Omega} + C\|\mathbf{q} - \mathbf{q}_h\|_{\mathbf{V}_h}. \end{aligned}$$

The desired estimate follows by combining this estimate and the estimate

$$\|(I - P_0)(\Pi u - u_h^*)\|_\Omega \leq Ch\|\nabla(I - P_0)(\Pi u - u_h^*)\|_\Omega$$

obtained by an element-wise Poincare inequality. \square

6. Numerical examples. In this section we verify the analysis of section 4 and section 5 via numerical experiments. All the examples have been implemented using the NGSolve finite element library [19].

6.1. The cavity problem. We consider the symmetric cavity problem of [7, Section 6].

Let $\Omega = (-1, 1) \times (0, 1)$, $\Omega_+ := (-1, 0) \times (0, 1)$, and $\Omega_- := (0, 1) \times (0, 1)$. We choose the source term $f \in L^2(\Omega)$ such that the exact solution to (2.4) is given by (6.1)

$$u(x_1, x_2) := \begin{cases} ((x_1 + 1)^2 - (\sigma_+ + \sigma_-)^{-1}(2\sigma_+ + \sigma_-)(x_1 + 1)) \sin(\pi x_2) & \text{on } \Omega_+, \\ (\sigma_+ + \sigma_-)^{-1} \sigma_+ (x_1 - 1) \sin(\pi x_2) & \text{on } \Omega_-. \end{cases}$$

We set $\sigma_+ = 1$ and $\kappa_\sigma = -1.001$.

We consider both a symmetric and a non-symmetric mesh, see Figure 6.1. We will compare the results obtained by the HDG method (3.7) to those obtained by an H^1 -conforming finite element method (CG) as analyzed in [7]. For the HDG method the penalty parameter is set to $\tau = 1$ on Ω_+ and $\tau = -1$ on Ω_- . We take $k = 0, 1, 2, 3$ as polynomial degree.

On the symmetric mesh we observe from Table 6.1 that the primal variable computed using the H^1 -conforming finite element method converges optimally in the L^2 -norm for $k \geq 1$. Optimal rates of convergence in the L^2 -norm are obtained with the HDG method for the primal variable as well as the vector variable for $k \geq 0$. Furthermore, we observe superconvergence in the post-processed primal variable for $k \geq 1$ when the HDG method is used. These HDG results verify our analysis in section 4 and section 5.

On the non-symmetric mesh we observe from Table 6.2 irregular rates of convergence in the L^2 -norm in the primal variable when using the H^1 -conforming finite




Fig. 6.1: A symmetric mesh and non-symmetric mesh used in the numerical examples described in [section 6](#).

element method for $k = 1$ and $k = 2$. Similar behaviour is observed also in [7]. Near optimal rates of convergence in the L^2 -norm are obtained with this method for $k = 3$. Using the HDG method we observe optimal rates of convergence in the L^2 -norm for the errors in u_h and \mathbf{q}_h and superconvergence for the error in u_h^* for $k = 2$ and $k = 3$. The rates of convergence in the L^2 -norm using the HDG method when $k = 0$ or $k = 1$ show the same irregular behaviour as when using the H^1 -conforming finite element method. We remark, though, that the HDG method always shows a smaller error on the same mesh compared to the H^1 -conforming finite element method.

In [Figure 6.2](#) we compare the solution u_h computed using the H^1 -conforming finite element method and HDG method with the exact solution for different values of κ_σ . We use $k = 2$ and compute the solution on a non-symmetric mesh consisting of 8960 simplicial elements. We observe that the closer κ_σ is to -1 (for which the problem is not well-posed), the worse the H^1 -conforming finite element compares to the exact solution. The HDG method compares well with the exact solution for all tested values of κ_σ . We observe the same behaviour in [Figure 6.3](#) in which we plot the solution along the line $x_2 = 0.5$. We indeed saw in [Table 6.2](#) that the HDG method outperforms the H^1 -conforming finite element method on the non-symmetric mesh.

6.2. Meta-material layer. We next consider a problem motivated by that given in [8, Section 6.1]. For this, let $\Omega = (0, 5) \times (0, 2)$. The definition of Ω_+ and Ω_- are shown in [Figure 6.4](#). For this setup it is known that (2.3) is well-posed if $\kappa_\sigma \in (-\infty, 0) \setminus [\kappa_{\min}, \kappa_{\max}]$ where $\kappa_{\min} \approx -1.46$ and $\kappa_{\max} \approx -0.69$, [7, Section 3.2].

We set $\sigma_+ = 1$ and

$$(6.2) \quad f(x_1, x_2) := \begin{cases} \sin(\pi x_2/2) & \text{in } \Omega_+ \text{ if } x_1 < 1.3, \\ 0 & \text{otherwise.} \end{cases}$$

We consider three different values for κ_σ : $\kappa_\sigma = -1.5$, $\kappa_\sigma = -1.6$, and $\kappa_\sigma = -2$.

We compute the solution using both the HDG method (3.7) and an H^1 -conforming finite element method. For this we take $k = 3$ and consider the solution on a mesh consisting of 24576 simplicial elements. We plot and compare contour plots of u_h obtained by both methods in [Figure 6.5](#). We observe that the closer κ_σ is to the interval $[\kappa_{\min}, \kappa_{\max}]$, the larger the difference in solution between the HDG and H^1 -conforming finite element method. This is observed also when plotting the solution on the line $x_2 = 1$. When $\kappa_\sigma = -1.5$ we observe a larger undershoot at $x_1 = 3.3$ with the H^1 -conforming finite element method than when using the HDG method.

Cells	CG e_u	r	HDG e_u	r	HDG e_q	r	HDG e_{u^*}	r
$k = 0$								
256	-	-	9.9e+0	-	1.9e+2	-	9.9e+0	-
1024	-	-	5.0e+0	1.0	9.9e+1	1.0	5.0e+0	1.0
4096	-	-	2.5e+0	1.0	5.0e+1	1.0	2.5e+0	1.0
16384	-	-	1.3e+0	1.0	2.5e+1	1.0	1.3e+0	1.0
65536	-	-	6.4e-1	1.0	1.3e+1	1.0	6.4e-1	1.0
262144	-	-	3.2e-1	1.0	6.3e+0	1.0	3.2e-1	1.0
$k = 1$								
256	1.0e+1	-	6.9e+0	-	1.5e+1	-	2.6e-1	-
1024	2.5e+0	2.0	1.7e+0	2.0	3.9e+0	2.0	3.3e-2	3.0
4096	6.3e-1	2.0	4.1e-1	2.0	9.7e-1	2.0	4.2e-3	3.0
16384	1.6e-1	2.0	1.0e-1	2.0	2.4e-1	2.0	5.3e-4	3.0
65536	3.9e-2	2.0	2.5e-2	2.0	6.1e-2	2.0	6.6e-5	3.0
262144	9.8e-3	2.0	6.4e-3	2.0	1.5e-2	2.0	8.3e-6	3.0
$k = 2$								
256	2.9e-1	-	2.5e-1	-	5.3e-1	-	7.3e-3	-
1024	3.7e-2	3.0	3.1e-2	3.0	6.7e-2	3.0	4.6e-4	4.0
4096	4.6e-3	3.0	3.8e-3	3.0	8.5e-3	3.0	2.9e-5	4.0
16384	5.8e-4	3.0	4.7e-4	3.0	1.1e-3	3.0	1.8e-6	4.0
65536	7.2e-5	3.0	5.9e-5	3.0	1.3e-4	3.0	1.1e-7	4.0
262144	9.0e-6	3.0	7.3e-6	3.0	1.7e-5	3.0	7.1e-9	4.0
$k = 3$								
256	7.3e-3	-	7.1e-3	-	1.5e-2	-	1.6e-4	-
1024	4.6e-4	4.0	4.4e-4	4.0	9.3e-4	4.0	5.1e-6	5.0
4096	2.9e-5	4.0	2.7e-5	4.0	5.8e-5	4.0	1.6e-7	5.0
16384	1.8e-6	4.0	1.7e-6	4.0	3.6e-6	4.0	5.0e-9	5.0
65536	2.2e-7	3.0	1.0e-7	4.0	2.3e-7	4.0	1.9e-10	4.7

Table 6.1: Rates of convergence in the L^2 -norm of CG and HDG on a symmetric mesh for different polynomial degree k , see [subsection 6.1](#).

When $\kappa_\sigma = -1.6$ and $\kappa_\sigma = -2$, i.e., κ_σ is further away from the interval $[\kappa_{\min}, \kappa_{\max}]$, the difference in solution between both discretizations is smaller. These results are in agreement with those observed in [subsection 6.1](#).

7. Conclusions. In this paper we introduced an HDG method for the solution of Poisson-type problems with a sign-changing coefficient. Well-posedness of our method does not rely on conditions on domain geometry and the ratio of the negative and positive coefficients. We presented a novel error analysis of the HDG method that avoids the Aubin–Nitsche duality argument. Under suitable regularity assumptions, a superconvergence result is obtained by a duality argument. Numerical experiments show that the HDG method outperforms an H^1 -conforming finite element method for this problem.

References.

- [1] D. N. Arnold and F. Brezzi. Mixed and nonconforming finite element methods: implementation, postprocessing and error estimates. *RAIRO Modél. Math. Anal. Numér.*, 19(1):7–32, 1985. doi: 10.1051/m2an/1985190100071.
- [2] A.-S. Bonnet-Ben Dhia, P. Ciarlet Jr., and C. M. Zwölf. Time harmonic wave diffraction problems in materials with sign-shifting coefficients. *J. Comput. Appl.*

Cells	CG e_u	r	HDG e_u	r	HDG e_q	r	HDG e_{u^*}	r
$k = 0$								
140	-	-	3.0e+2	-	1.5e+3	-	3.0e+2	-
560	-	-	2.7e+2	0.2	3.3e+3	-1.1	2.7e+2	0.2
2240	-	-	3.0e+2	-0.2	2.4e+3	0.4	3.0e+2	-0.2
8960	-	-	2.8e+2	0.1	2.2e+3	0.1	2.8e+2	0.1
35840	-	-	4.4e+3	-4.0	5.2e+4	-4.6	4.4e+3	-4.0
143360	-	-	1.8e+2	4.6	5.8e+3	3.2	1.8e+2	4.6
573440	-	-	1.6e+1	3.5	5.6e+2	3.4	1.6e+1	3.5
$k = 1$								
140	2.7e+3	-	2.5e+1	-	2.4e+2	-	2.2e+1	-
560	1.3e+2	4.3	7.6e+1	-1.6	1.1e+3	-2.3	7.6e+1	-1.8
2240	2.5e+1	2.4	1.5e+0	5.7	3.6e+1	5.0	1.2e+0	6.0
8960	1.2e+1	1.1	8.7e-1	0.8	3.3e+1	0.1	8.5e-1	0.5
35840	6.1e+0	1.0	1.2e+0	-0.4	7.9e+1	-1.3	1.2e+0	-0.5
143360	2.1e+0	1.5	1.3e-2	6.5	4.2e-1	7.5	2.4e-3	8.9
$k = 2$								
140	2.3e+2	-	1.1e+0	-	2.0e+1	-	9.1e-1	-
560	1.0e+3	-2.1	1.0e-1	3.3	3.1e+0	2.7	5.5e-2	4.0
2240	7.8e-1	10.3	1.1e-2	3.2	2.6e-1	3.6	2.7e-3	4.3
8960	7.9e-2	3.3	1.4e-3	3.0	1.8e-2	3.8	9.5e-5	4.9
35840	5.2e-2	0.6	1.7e-4	3.0	1.6e-3	3.5	4.1e-6	4.5
143360	7.8e-4	6.1	2.2e-5	3.0	2.6e-4	2.6	3.8e-7	3.4
$k = 3$								
140	7.1e-1	-	3.0e-2	-	8.6e-1	-	2.0e-2	-
560	7.0e-2	3.3	2.5e-3	3.6	8.5e-2	3.4	1.5e-3	3.8
2240	2.6e-3	4.7	1.3e-4	4.3	3.9e-3	4.5	3.5e-5	5.4
8960	7.9e-5	5.1	7.7e-6	4.1	1.7e-4	4.5	7.6e-7	5.5
35840	3.3e-6	4.6	4.8e-7	4.0	7.3e-6	4.5	1.6e-8	5.5
143360	2.7e-7	3.6	3.0e-8	4.0	3.2e-7	4.5	7.1e-10	4.5

Table 6.2: Rates of convergence in the L^2 -norm of CG and HDG on a non-symmetric mesh for different polynomial degree k , see [subsection 6.1](#).

- Math.*, 234(6):1912–1919, 2010. doi: 10.1016/j.cam.2009.08.041.
- [3] A.-S. Bonnet-Ben Dhia, L. Chesnel, and P. Ciarlet Jr. T-coercivity for scalar interface problems between dielectric and metamaterials. *ESAIM: M2AN*, 46: 1363–1387, 2012. doi: 10.1051/m2an/2012006.
 - [4] A.-S. Bonnet-Ben Dhia, L. Chesnel, and P. Ciarlet Jr. T-coercivity for the Maxwell problem with sign-changing coefficients. *Comm. Partial Differential Equations*, 39(6):1007–1031, 2014. doi: 10.1080/03605302.2014.892128.
 - [5] A.-S. Bonnet-Ben Dhia, L. Chesnel, and P. Ciarlet Jr. Two-dimensional Maxwell’s equations with sign-changing coefficients. *Appl. Numer. Math.*, 79:29–41, 2014. doi: 10.1016/j.apnum.2013.04.006.
 - [6] H. Chen, W. Qiu, K. Shi, and M. Solano. A superconvergent hdg method for the maxwell equations. *J. Sci. Comput.*, 70(3):1010–1029, 2017. doi: 10.1007/s10915-016-0272-z.
 - [7] L. Chesnel and P. Ciarlet Jr. T-coercivity and continuous Galerkin methods: application to transmission problems with sign changing coefficients. *Numer. Math.*, 124:1–29, 2013. doi: 10.1007/s00211-012-0510-8.




Fig. 6.2: A contour plot of u_h comparing the H^1 -conforming finite element method and HDG method (dashed lines) with the exact solution (solid lines) for different values of κ_σ , see [subsection 6.1](#).

- [8] E. T. Chung and P. Ciarlet Jr. A staggered discontinuous Galerkin method for wave propagation in media with dielectrics and meta-materials. *J. Comput. Appl. Math.*, 239:189–207, 2013. doi: 10.1016/j.cam.2012.09.033.
- [9] B. Cockburn, B. Dong, and J. Guzmán. A superconvergent LDG-hybridizable Galerkin method for second-order elliptic problems. *Math. Comp.*, 77(264):1887–1916, 2008. doi: 10.1090/S0025-5718-08-02123-6.
- [10] B. Cockburn, J. Gopalakrishnan, and R. Lazarov. Unified hybridization of discontinuous Galerkin, mixed, and continuous Galerkin methods for second order elliptic problems. *SIAM J. Numer. Anal.*, 47(2):1319–1365, 2009. doi: 10.1137/070706616.
- [11] B. Cockburn, J. Gopalakrishnan, and F. Sayas. A projection-based error analysis of HDG methods. *Math. Comp.*, 79:1351–1367, 2010. doi: 10.1090/S0025-5718-10-02334-3.
- [12] B. Cockburn, O. Dubois, J. Gopalakrishnan, and S. Tan. Multigrid for an HDG method. *IMA J. Numer. Anal.*, 34(4):1386–1425, 2014. doi: 10.1093/imanum/drt024.




Fig. 6.3: Comparing the solution exact solution u (solid lines) along the line $x_2 = 0.5$ with that computed using the H^1 -conforming finite element method and the HDG method (dashed lines) for different values of κ_σ , see [subsection 6.1](#).




Fig. 6.4: The domain $\Omega = (0, 5) \times (0, 2)$ is subdivided in Ω_+ and Ω_- , see [subsection 6.2](#).

- [13] J. Cui and W. Zhang. An analysis of HDG methods for the Helmholtz equation. *IMA Journal of Numerical Analysis*, 34(1):279–295, 2013. doi: 10.1093/imanum/drt005.
- [14] L. C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, Second edition, 2010. doi: 10.1090/gsm/019.
- [15] X. Feng, P. Lu, and X. Xu. A hybridizable discontinuous galerkin method for the time-harmonic maxwell equations with high wave number. *Computational Methods in Applied Mathematics*, 16(3):429–445, 2016. doi: 10.1515/cmam-2016-0021.
- [16] B. M. Fraeijs de Veubeke. Stress function approach. In *World Congress on the Finite Element Method in Structural Mechanics*. Bournemouth, 1975.
- [17] G. P. Galdi. *An introduction to the mathematical theory of the Navier–Stokes equations. Steady-state problems*. Springer Monographs in Mathematics. Springer, New York, Second edition, 2011. doi: 10.1007/978-0-387-09620-9.
- [18] R. Griesmaier, , and P. Monk. Error analysis for a hybridizable discontinuous Galerkin method for the Helmholtz equation. *J. Sci. Comput.*, 49(3):291–310, 2011. doi: 10.1007/s10915-011-9460-z.
- [19] J. Schöberl. C++11 implementation of finite elements in NGSolve. Technical Report ASC Report 30/2014, Institute for Analysis and Scientific Computing, Vienna University of Technology, 2014. URL <http://www.asc.tuwien.ac.at/~schoeberl/wiki/publications/ngs-cpp11.pdf>.
- [20] R. Stenberg. Postprocessing schemes for some mixed finite elements. *RAIRO Modél. Math. Anal. Numér.*, 25(1):151–167, 1991. doi: 10.1051/m2an/1991250101511.




Fig. 6.5: A contour plot of u_h a slice through $x_2 = 1$ comparing the solution obtained by the HDG method (solid lines) to an H^1 -conforming finite element method (dashed lines) for different values of κ_σ , see [subsection 6.2](#).