## Very slow heating for weakly driven quantum many-body systems

Wojciech De Roeck<sup>1</sup> and Victor Verreet<sup>1</sup>

<sup>1</sup>KU Leuven, 3001 Leuven, Belgium<sup>\*</sup> (Dated: 8th December 2021)

It is well understood that many-body systems driven at high frequency heat up only exponentially slowly and exhibit a long prethermalization regime. We prove rigorously that a certain relevant class of systems heat very slowly under weak periodic driving at intermediate frequency as well. This class of systems are those whose time-dependent, possibly translation-invariant, Hamiltonian is a weak perturbation of a sum of mutually commuting, terms. This condition covers several periodically kicked systems that have been considered in the literature recently, in particular kicked Ising models. In contrast to the high-frequency regime, the prethermalization dynamics of our systems is in general not related to any time-independent effective Floquet Hamiltonian. Our results also have nontrivial implications for closed (time-independent) systems. We use the example of an Ising model with transversal and longitudinal field to show how they imply confinement of excitations. More generally, they show how "glassy" kinetically constrained models emerge naturally from simple many-body Hamiltonians, thus connecting to the topic of 'translation-invariant localization'.

PACS numbers:

Introduction The phenomenon of a long-lived quasistationary state, also known as a prethermal state, has become an important paradigm in the theory of nonequilibrium many-body systems. Such quasi-stationary states often exhibit interesting features that cannot be realized in equilibrium states [17, 30, 34, 55]. One crucial ingredient to have a long-lived prethermal regime is of course that the system resists thermalization for a long time. In this letter, we are interested in cases where the thermalization rate is beyond perturbation theory. In particular, we do not consider the oft-discussed case of prethermalization in weakly perturbed integrable systems, where the prethermalization regime lasts in general for a time  $\propto 1/g^2$  with g the perturbation strength [6, 7, 16, 33, 44], but instead we look for thermalization times superpolynomial in 1/q.

Several instances of such prethermal phases have been identified. The most obvious class consists of systems periodically driven at high frequency, where a Magnus-Floquet expansion yields an effective Hamiltonian [1, 2, 8, 10, 28, 36]. It is natural to inquire whether a similar scenario might also be realized away from the high-frequency regime and in this letter we aim to provide a positive and general answer to this question.

Weakly driven systems Given a Hamiltonian  $H = H^0 + gW(t)$  describing spins on a spatial lattice of arbitrary dimension, with periodic but otherwise generic driving W(t) = W(t + T). When is the heating rate non-perturbatively small in g, i.e. taking g small compared to other local energy scales, regardless of the initial state? This question remains meaningful if we allow  $H^0$  to depend periodically on t as well, as long as its one-cylce propagator  $U_0 = \mathcal{T} e^{-i \int_0^T dt H^0(t)} (\mathcal{T} \dots$  denotes time-ordering of the epxonential), or one of its powers  $U_0^p, p \in \mathbb{N}$ , has a meaningful (i.e. local) conservation law, i.e. there is a Q such that  $[Q, U_0^p] = 0$  and Q is a sum of

local terms. Such a conservation law has observable consequences and the question of its persistence at  $g \neq 0$  is well-posed. In this setting, we argue that heating is nonpertubatively slow whenever we can represent the *p*-cycle propagator as  $U_0^p = \mathcal{T} e^{-i \int_0^{pT} dt \tilde{H}^0(t)}$  with  $\tilde{H}^0(t)$  a sum of local terms  $\tilde{H}_i^0(t)$  that mutually commute at all times  $[\tilde{H}_i^0(t), \tilde{H}_j^0(t')] = 0$ , and a certain *Diophantine condition* is satisfied.

Static systems Now we take a time-independent local many-body Hamiltonian  $H = H^0 + gW$  with W generic and g again small compared to local energy scales. Instead of 'heating', the appropriate question is now to what extent conservation laws of  $H^0$  (in particular  $H^0$ itself) are broken by gW. We claim that a sufficient conditon for superpolynomial persistence of these conservation laws is again that  $H^0$  is a sum of mutually commuting terms. The larger the set of Bohr frequencies (energy differences) defined by these local commuting terms, the smaller we need to take g to see the effect, but the more numerous the number of quasi-conserved charges Q. This multitude of conserved quantities can lead to local frustration and emergent kinetic constraints, as we will demonstrate in an example.

**Ergodicity-breaking phases** The most well-known examples of the above claims are the cases where  $H^0, \tilde{H}^0$ , respectively, are sums of commuting disordered terms (known as 'LIOMs' [22, 24, 45, 47]). Then, those claims are weakened versions of the stability of many-body localization (MBL) w.r.t. small perturbations, be it in static systems [4, 20, 37], periodically driven [3, 29, 39] or time-crystals (case p > 1)[26, 27, 53]. Our claims are weaker because they only state that the thermalization time is very large instead of infinite and also because they posit the existance of a few ( $\mathcal{O}(1)$  regardless of volume) quasi-conserved charges Q. In this letter, we concentrate on translation invariant  $H^0, \tilde{H}^0$  and we exhibit rigorously

 $\mathcal{O}(1)$  number of charges. Depending on the initial state, it may actually happen that there emerge additional  $\mathcal{O}(L)$  quasi-conserved charges (L= number of degrees of freedom) via frustration and effective kinetic constraints. This is related to *fractons* [38, 42] and, more generally, to so-called translation invariant quasi-localization, see e.g. [12, 21, 25, 46, 54].

Intuitive picture Let us start from the static setting  $H = H^0 + gW$ . As announced, we assume that  $H^0 = \sum_i H_i^0$  with  $[H_i^0, H_j^0] = 0$ , therefore every operator of the form  $\sum_i f(H_i^0)$  commutes with  $H^0$  as well. If now all  $H_i^0$  have a finite spatial range R and norm bound  $\gamma$ , and the number of degrees of freedom per site (local Hilbert space dimension) is finite as well, then we can represent

$$H^0 = \mathbf{J} \cdot \mathbf{N} = \sum_{\alpha=1}^q J_\alpha N^{(\alpha)}$$

where  $N^{(\alpha)} = \sum_{i} N_i^{(\alpha)}$  with *i* labelling lattice sites and such that **a**)  $[N_i^{(\alpha)}, N_j^{(\alpha')}] = 0$ , **b**)  $N_i^{(\alpha)}$  has integer spectrum, **c**)  $N_i^{(\alpha)}$  acts only within a distance *R* of site *i* and is uniformly bounded  $||N_i^{\alpha}|| \leq \gamma$ .

If the conservation laws  $N^{(\alpha)}$  are to be broken by local terms of strength g, we have to find *local transitions*  $\mathbf{n} \to \mathbf{n} + \Delta \mathbf{n}$  with  $\Delta \mathbf{n} \in \mathbb{Z}^q$  that are on-resonance up to an energy offset of order g, i.e.

$$|\mathbf{J} \cdot \Delta \mathbf{n}| \le \mathcal{O}(g) \tag{1}$$

Since gW acts locally and the  $N_i^{\alpha}$  are bounded, the components of  $\Delta \mathbf{n}$  have to be smaller than some fixed value. This means that, for generic  $\mathbf{J}$  and small enough g, this constraint allows no nonzero solution  $\mathbf{n} \neq 0$  and hence we are led to believe that all  $N^{(\alpha)}$  are conserved. This is of course merely first-order reasoning but it turns out (see later) that higher orders do not change the picture, upon taking non-dissipative effects into account by a unitary frame rotation. If we add periodic time-dependence (frequency  $2\pi/T$ ) to the problem, then similar reasoning applies with the role of  $H^0$  now played by  $\frac{1}{T} \int_0^{pT} dt H^0(t)$ . To allow for the possibility of absorbing/emitting  $\Delta n_0$ quanta from the drive, we modify the condition (1) to

$$|\mathbf{J} \cdot \Delta \mathbf{n} + (2\pi/T)\Delta n_0| \le g$$

The same considerations as above apply, except that now we need a constraint on the q + 1-tuple  $(2\pi/T, \mathbf{J})$ . Such constraints are well-known from KAM-phenomena [41] and Nekoroshev estimates, where, just as in our case, they express that resonances are absent. Since the frequency  $2\pi/T$  enters the above formulas on the same footing as the couplings  $J^{\alpha}$ , the same formalism applies to quasi-periodically driven systems as well (long prethermalization observed in [15]), see [18].

We stress that the above considerations involve *spatial* locality, as opposed to locality in momentum space. Indeed, consider the relevant case of free fermions:  $H^0 =$ 

 $\sum_k \omega(k) n_k$  with  $n_k = c_k^{\dagger} c_k$  the occupation operator of momentum mode k, and  $\omega(k)$  the dispersion relation. Hence  $H^0$  is a sum of very simple and mutually commuting operators, but they are **not** local and they do not satisfy our requirements, Indeed, for such  $H^0$  one expects kinetic theory based on non-linear Boltzmann equations to describe the heating process, with rate  $\propto g^2$ , see [31, 49].

Result for static systems Our systems live on a large but finite graph  $\Lambda$ , with a finite Hilbert space  $\mathbb{C}^d$  attached to each site  $i \in \Lambda$ . The total Hilbert space  $\mathcal{H}$  is hence  $(\mathbb{C}^d)^{\otimes_{\Lambda}}$  and we say that an operator  $O = O_S$  is supported in a set S if it is of the form  $O_S \otimes \mathbb{1}_{S^c}$ . We consider a Hamiltonian of the form

 $H = H^0 + qW,$ 

with

$$H^{0} = \mathbf{J} \cdot \mathbf{N} = \sum_{\alpha=1}^{q} J_{\alpha} N^{(\alpha)}$$
(2)

where the operators  $N^{(\alpha)}$  satisfy the conditions  $\mathbf{a}, \mathbf{b}, \mathbf{c}$ ) in the previous section, and, as a concrete generalization of condition (1) we assume the following *Diophantine* condition expressing that the **J** are sufficiently incommensurable at all orders in perturbation theory: there exist positive numbers  $\tau, x > 0$ , such that

$$|\mathbf{m} \cdot \mathbf{J}| \ge \frac{x|\mathbf{J}|}{|\mathbf{m}|^{\tau}}, \qquad \forall \mathbf{m} \in \mathbb{Z}^q$$
(3)

Finally, we need the Hamiltonian gW to have local terms of strength  $\mathcal{O}(g)$ , decaying fast in the size of their support. To that end, we define (see SM) the *local norms*  $|| \cdot ||_{\kappa}$  with spatial decay parameter  $\kappa > 0$  and we choose a  $\kappa_0 > 0$  such that  $||gW||_{\kappa_0} \leq g$ .

**Theorem 1.** There is a unitary base change  $\hat{Y}$  such that, in the rotated frame, the Hamiltonian takes the form

$$\hat{Y}H\hat{Y}^{\dagger} = H^0 + \hat{D} + \hat{V}$$

where, with constants c, C depending only on the parameters  $\kappa_0, \gamma, q, p$  and in particular not on the volume  $|\Lambda|$ ,

1. The driving  $\hat{V}$  is very weak:

$$||\hat{V}||_{\kappa} \leq g \exp\left(-c[1/\epsilon]^{1/p}\right), \qquad \epsilon \equiv \frac{g}{|\mathbf{J}|}(1+\frac{1}{x})$$
  
for any  $p > q + \tau$  and  $\kappa = c\kappa_0 |\log \epsilon|^{-1}$ .

2.  $||\hat{D}||_{\kappa} \leq Cg$  and  $[N^{(\alpha)}, \hat{D}] = 0$  for any  $\alpha = 1, \dots, q$ .

We see hence that  $\hat{D}$  inherits the conservation laws of  $H^0$  and  $\hat{V}$  should be considered as a remaining weak driving, enforcing thermalization at quasi-exponential times. Importantly, the unitary conjugation  $\hat{Y} \cdot \hat{Y}^{\dagger}$  is close to identity: for local operators O with support size  $\mathcal{O}(1)$ , we have  $||\hat{Y}O\hat{Y}^{\dagger} - O|| \leq C\epsilon ||O||$  (see SM for a more precise statement and construction). One obvious consequence is that, in the original frame, the dressed operators  $\hat{N}^{(\alpha)} \equiv \hat{Y}N^{(\alpha)}\hat{Y}^{\dagger}$ , which are sums of quasi-ocal terms, are quasi-conserved:

$$\frac{1}{|\Lambda|} ||\mathbf{e}^{\mathbf{i}tH} \hat{N}^{(\alpha)} \mathbf{e}^{-\mathbf{i}tH} - \hat{N}^{(\alpha)}|| \le g \exp\left(-c[1/\epsilon]^{1/p}\right)$$

up to stretched-exponential long time  $\exp(c[1/\epsilon]^{1/p})$ . Hence in particular thermalization is obstructed until that time.

**Comments** If we imagine **J** to be chosen uniformly on the unit hypersphere  $S^q$ , then the probability of the Diophantine condition (3) being violated, decays as Cx when  $x \to 0$ , provided that  $\tau > q - 1$ . In particular, (3) is satisfied with probability 1 for some x. It follows that the power p in the stretched exponential can be chosen  $p = 2q - 1 + \delta$ , with  $\delta$  arbitrarily small. In particular, for the case q = 1, this yields almost an exponential in  $1/\epsilon$ , as was already proven in [1, 17].

If the condition (3) (for a given x) fails for some  $\mathbf{m}$  with  $|\mathbf{m}| \sim R\gamma n$ , then this simply means that our perturbative reduction of the driving  $\hat{V}$  can only be carried out to a power n instead of a power diverging as  $\epsilon \to 0$ , giving  $||V||_{\kappa} \leq g\epsilon^n$ . Therefore, it is really only low-order resonances that can hamper the slow thermalization. However, such low-order resonances can almost always be lifted by redefining  $H^0$ , as we also illustrate in the example. **Example: static Ising** Let

$$H = \sum_{i} J\sigma_i^x \sigma_{i+1}^x + h_x \sigma_i^x + h_z \sigma_i^z$$

and we consider  $h_z$  as the weak-driving parameter g, i.e. we set  $H^0 \equiv JN^{(1)} + h_x N^{(2)}$  with  $N^{(1)} \sum_i \sigma_i^x \sigma_{i+1}^x$  the number of domain walls (up to a constant) and  $N^{(2)} = \sum_i \sigma_i^x$  the magnetization.

If the pair  $(J, h_x)$  is sufficiently incommensurable, i.e. condition (3) holds, then our theory yields that, in the rotated frame, the Hamiltonian is approximately  $\hat{H} =$  $JN^{(1)} + h_x N^{(2)} + \hat{D}$  (we neglected the very weak driving  $\hat{V}$ ), with both  $N^{(1,2)}$  commuting with  $\hat{D}$  and hence conserved locally. For example, let us consider a configuration of the form

The only transition that preserves both magnetization and the number of domains consists of a shift of the entire block  $\uparrow\uparrow \cdot \uparrow\uparrow$ . However, since the operator  $\hat{D}$  is a sum of local terms, the largest term that can cause such a shift, is of size  $g\epsilon^{c\ell_0}$  with c a constant of order 1. In our formalism, this follows by the local bound  $||\hat{D}||_{\kappa} \leq Cg$ , expressing exponential decay of local terms in  $\hat{D}$ . Hence, the domains acquire a very large mass and and are nearly static. An even more drastic example is

$$\cdots$$

with the configuration extending infinitely far in both directions. Here, no single transition is allowed by the conservation laws and there should be therefore no dissipative dynamics up to (quasi-)exponential time. These phenomena were numerically studied and explained in [35], our theorem adds a controlled proof and a novel point of view. If  $(J, h_x)$  are commensurable, say  $J = 2h_x$ , then of course our theorem does not apply in the above way. However, in that case, one can choose  $H^0 = h_x N^{(1')}$ with  $N^{(1')} \equiv 2N^{(1)} + N^{(2)}$ , giving again a meaningful constraint. Let us finally return to the case of incommensurable  $(J, h_x)$ . More generally, if we consider a system with a small density of domain walls, the above considerations show that the local dynamics is highly constrained and appears many-body-localized for a long time. It can hence be a considered as an emergent kinetically constrained model [52]. However, as was argued in [13], the long time scale that emerges here is in general not beyond perturbation theory in the parameter q.

**Result for periodic driving** We are inspired by the setup proposed in the section "weakly driven systems", with  $U_0^p$  generated by  $\tilde{H}^0(t)$ . However, for the sake of notational simplicity, we write now  $H^0$  instead of  $\tilde{H}^0$  and we redefine  $T \mapsto pT$ . This means that we potentially describe the original system of interest only at stroboscopic times  $t \in pT\mathbb{N}$ , but this suffices for the sake of obstructions to thermalization.

Concretely, we consider a  $T\mbox{-}{\rm periodic}$  Hamiltonian of the form

$$H(t) = H^{0}(t) + gW(t), \qquad H(t) = H(t+T)$$

with

$$H^{0}(t) = \mathbf{J}(t) \cdot \mathbf{N} = \sum_{\alpha=1}^{q} J_{\alpha}(t) N^{(\alpha)}$$
(4)

Here, the operators  $N^{(\alpha)}$  are exactly as in the static setup with the sole difference that we allow *T*-periodic coupling  $\mathbf{J}(t) = \mathbf{J}(t+T)$ . Furthermore, we assume that the timedependent (local terms of) W(t) and the couplings  $\mathbf{J}(t)$ are piecewise-continuous. Just as in the static case, we assume the local bound  $||gW||_{\kappa_0} \leq g$ , except that now this bound includes a supremum over t, see SM. Finally, the *Diophantine* condition is slighly modified: there are  $0 < \tau, x < \infty$  such that, with  $\bar{J} = \frac{1}{T} \int_0^T dt \mathbf{J}(t)$ 

$$\inf_{n \in \mathbb{Z}} |T\mathbf{m} \cdot \bar{\mathbf{J}} - 2\pi n| \ge \frac{x}{|\mathbf{m}|^{\tau}}, \qquad \forall \mathbf{m} \in \mathbb{Z}^q \qquad (5)$$

We let U(t) be the solution of the Schrödinger equation

$$i\partial_t U(t) = H(t)U(t), \qquad U(0) = 1$$

**Theorem 2.** There is a *T*-periodic unitary base-change  $\hat{Y}(t)$  such that  $\hat{U}(t) \equiv \hat{Y}(t)U(t)$  solves the Schrödinger equation

$$i\partial_t \hat{U}(t) = (H^0(t) + \hat{D}(t) + \hat{V}(t))\hat{U}(t),$$




Figure 1: (Taken From [43], Figure 13). Kicked Ising chain at J = 1. White regions are slowly thermalising: an appropriately defined thermalization rate is found to be smaller than  $10^{-6}$  per cycle. The colored lines and

letters demarcate our perturbative regimes

and (with c, C depending only on  $\gamma$ ,  $\kappa_0$ , q and p)

- 1. The driving  $\hat{V}(t)$  is very weak
- $||\hat{V}(t)||_{\kappa} \le g \exp\left(-c[1/\epsilon]^{1/p}\right), \qquad \epsilon \equiv gT\left(1+\frac{1}{x}\right)$

with  $p > q + \tau + 1$  and  $\kappa = c\kappa_0 |\log \epsilon|^{-1}$ 

2.  $||\hat{D}(t)||_{\kappa} \leq Cg$  and  $[N^{(\alpha)}, \hat{D}] = 0$  for any  $\alpha = 1, \ldots, q$ .

Just as in the static case, this theorem shows that the dressed operators  $\hat{N}^{(\alpha)}(t) = \hat{Y}(t)N^{(\alpha)}\hat{Y}^*(t)$  are quasi-conserved  $\frac{1}{|\Lambda|} \left( ||\hat{N}^{(\alpha)}(t) - U(t)\hat{N}^{(\alpha)}(0)U^{\dagger}(t)|| \right) \leq g \exp\left(-c[1/\epsilon]^{1/p}\right)$  for  $t \leq (1/g) \exp\left(c[1/\epsilon]^{1/p}\right)$  and they obstruct heating.

Our scheme does not yield any effective Hamiltonian  $H_F$ , satisfying  $U \approx e^{-iTH_F}$ , with  $\approx$  indicating validity up to long times. Instead, we obtain  $U = \mathcal{T}e^{-i\int_0^T dt(H^0(t)+\hat{D}(t))}$ . This would yield a  $H_F = \frac{1}{T}\int_0^T dt(H^0(t) + \hat{D}(t))$  provided that the local terms of  $H^0(t) + \hat{D}(t)$  commute between different times t, but this is not true in general[56] and so existence of some  $H_F$ seems unlikely.

**Example: kicked Ising chain** We consider the one-cycle unitary given by

$$U \equiv e^{-i\sum_i J\sigma_i^x \sigma_{i+1}^x} e^{-i\sum_i \mathbf{h} \cdot \boldsymbol{\sigma}_i}$$

where  $\boldsymbol{\sigma}_i = (\sigma_i^x, \sigma_i^y, \sigma_i^z)$  are the Pauli-matrices acting on site  $i = 1, \ldots, L$ . This model was studied numerically in

detail in [43]. Figure 1 (copied from Figure 13 in [43]) shows the regimes where one observes very slow thermalization numerically. We view this model as originating from a time-dependent  $H(t) = H_1 \mathbb{1}_{[0,1]}(t) + H_2 \mathbb{1}_{[1,2]}(t)$ with period T = 2. Our Theorem 2 applies to this model in several cases (not mutually exclusive), distinghuished by which Hamiltonian plays the role of  $H^0(t)$ .

**Case A** ( $|\mathbf{h}| \ll 1$ ). Here, we let  $H^0(t) = J(t) \sum_i \sigma_i^x \sigma_{i+1}^x$ with  $J(t) = J1_{[0,1]}(t)$ . Theorem 2 posits that heating is non-perturbatively slow in  $|\mathbf{h}| \ll 1$ .

**Case B** ( $|J| \ll 1$ ). Here, we take  $H^0(t) = \sum_i \mathbf{h}(t) \cdot \boldsymbol{\sigma}_i$  with  $\mathbf{h}(t) = \mathbf{h}\mathbf{1}_{[1,2]}$ . Theorem 2 says that heating is no-perturbatively slow in  $|J| \ll 1$ .

**Case C** ( $|h_z| \ll 1$ ). Here  $H^0(t) = \sum_i (J1_{[0,1]}(t)\sigma_i^x \sigma_{i+1}^x + h_x 1_{[1,2]}(t)\sigma_i^x)$  and we take  $|h_z| \ll 1$ , again Theorem 2 shows slow heating in that regime.

**Case D** ( $|h_x| \ll 1$ ). Here  $U_0$  can be mapped to a free fermion expression with non-trivial dispersion relation by the Jordan-Wigner transformation. However, as we argued in section 'intuitive picture', this does not meet our criteria. The figure shows indeed that there is **no** very slow heating for  $|h_x| \ll 1$  (provided other couplings are not in a perturbative regime).

**Case E**  $(|\boldsymbol{h} - (0, 0, \frac{\pi}{2})| \ll 1)$ . Here we need to take the second power of the propagator. Indeed, now  $U_0 = e^{-i\sum_i J\sigma_i^x \sigma_{i+1}^x} e^{-i\frac{\pi}{2}\sum_i \sigma_i^z} = e^{-i\sum_i J\sigma_i^x \sigma_{i+1}^x} F$  with the flip  $F = \prod_i (-i\sigma_i^z)$ . Therefore,  $U_0^2 = e^{-2iJ\sum_i \sigma_i^x \sigma_{i+1}^x}$ , i.e. the setup applies simply setting  $H^0(t) = 2J\sum_i \sigma_i^x \sigma_{i+1}^x$ .

The claims in cases  $\mathbf{A}, \mathbf{B}$  have also been confirmed recently in [51] by numerics and a replica expansion for an effective Hamiltonian  $H_F$  satisfying  $U = e^{-iTH_F}$ . As explained above, the existence of  $H_F$  is however not suggested by our treatment. More precisely,[51] predict slow thermalization with the stretched exponential with p = 2. In our theorem, taking q = 1 and  $\tau > 1$  so that the Diophantine condition (5) is satisfied almost surely for some x (see SM), we prove these claims with  $p = 3 + \delta$ , for arbitrarily small  $\delta$ .

**Conclusion** We have identified a class of conditions under which non-perturbative rigorous lower bounds on the thermalization time can be proven, both in static and periodically driven systems. In the driven case, our theorems allow to understand the phase diagram of the kicked Ising model. In the static case, we provide a rigorous underpinning of the observed very slow dynamics of domain walls in Ising models.

**Note** The manuscript [18] appeared while we were finishing our paper. It is based on the same ideas as the present letter, has very analogous results, and goes beyond our analysis in many respects. Since it however focuses on the case of quasiperiodic driving, its analogue of our Theorem 2 has technical restrictions on the smoothness of the driving protocol, in particular ruling out our main example "kicked Ising models". Acknowledgements W.D.R. acknowledges the support of the Flemish Research Fund FWO undergrants G098919N and G076216N, and the support of KULeuven University under internal grantC14/16/062.

\* Electronic address: wojciech.deroeck@kuleuven.be

- Dmitry Abanin, Wojciech De Roeck, Wen Wei Ho, and François Huveneers. A rigorous theory of manybody prethermalization for periodically driven and closed quantum systems. *Communications in Mathematical Physics*, 354(3):809–827, 2017.
- [2] Dmitry A. Abanin, Wojciech De Roeck, Wen Wei Ho, and F. Huveneers. Effective hamiltonians, prethermalization, and slow energy absorption in periodically driven manybody systems. *Phys. Rev. B*, 95:014112, Jan 2017.
- [3] Dmitry A Abanin, Wojciech De Roeck, and François Huveneers. Theory of many-body localization in periodically driven systems. *Annals of Physics*, 372:1–11, 2016.
- [4] DM Basko, IL Aleiner, and BL Altshuler. Metalinsulator transition in a weakly interacting many-electron system with localized single-particle states. Annals of physics, 321(5):1126–1205, 2006.
- [5] Giancarlo Benettin, Jürg Fröhlich, and Antonio Giorgilli. A nekhoroshev-type theorem for hamiltonian systems with infinitely many degrees of freedom. *Communications in mathematical physics*, 119(1):95–108, 1988.
- [6] Jürgen Berges, Sz Borsányi, and Christof Wetterich. Prethermalization. *Physical review letters*, 93(14):142002, 2004.
- [7] Bruno Bertini, Fabian HL Essler, Stefan Groha, and Neil J Robinson. Prethermalization and thermalization in models with weak integrability breaking. *Physical re*view letters, 115(18):180601, 2015.
- [8] Marin Bukov, Luca D'Alessio, and Anatoli Polkovnikov. Universal high-frequency behavior of periodically driven systems: from dynamical stabilization to floquet engineering. Adv. Phys., 64(2):139–226, 2015.
- [9] Noé Cuneo, Jean-Pierre Eckmann, and C Eugene Wayne. Energy dissipation in hamiltonian chains of rotators. *Nonlinearity*, 30(11):R81, 2017.
- [10] L. D'Alessio and M. Rigol. Long-time behavior of isolated periodically driven interacting lattice systems. *Phys. Rev.* X, 4:041048, Dec 2014.
- [11] Nilanjana Datta, Roberto Fernández, and Jürg Fröhlich. Low-temperature phase diagrams of quantum lattice systems. i. stability for quantum perturbations of classical systems with finitely-many ground states. *Journal of statistical physics*, 84(3-4):455–534, 1996.
- [12] Wojciech De Roeck and François Huveneers. Asymptotic quantum many-body localization from thermal disorder. *Communications in Mathematical Physics*, 332(3):1017– 1082, 2014.
- [13] Wojciech De Roeck and François Huveneers. Scenario for delocalization in translation-invariant systems. *Physical Review B*, 90(16):165137, 2014.
- [14] Wojciech De Roeck and François Huveneers. Asymptotic localization of energy in nondisordered oscillator chains. *Communications on Pure and Applied Mathematics*, 68(9):1532–1568, 2015.

- [15] Philipp T Dumitrescu, Romain Vasseur, and Andrew C Potter. Logarithmically slow relaxation in quasiperiodically driven random spin chains. *Physical review letters*, 120(7):070602, 2018.
- [16] Martin Eckstein, Marcus Kollar, and Philipp Werner. Thermalization after an interaction quench in the hubbard model. *Physical review letters*, 103(5):056403, 2009.
- [17] Dominic V Else, Bela Bauer, and Chetan Nayak. Prethermal phases of matter protected by time-translation symmetry. *Physical Review X*, 7(1):011026, 2017.
- [18] Dominic V Else, Wen Wei Ho, and Philipp T Dumitrescu. Long-lived interacting phases of matter protected by multiple time-translation symmetries in quasiperiodicallydriven ssystems. arXiv preprint arXiv:1910.03584, 2019.
- [19] Antonio Giorgilli, Simone Paleari, and Tiziano Penati. An extensive adiabatic invariant for the klein–gordon model in the thermodynamic limit. In Annales Henri Poincaré, volume 16, pages 897–959. Springer, 2015.
- [20] IV Gornyi, AD Mirlin, and DG Polyakov. Interacting electrons in disordered wires: Anderson localization and low-t transport. *Physical review letters*, 95(20):206603, 2005.
- [21] James M Hickey, Sam Genway, and Juan P Garrahan. Signatures of many-body localisation in a system without disorder and the relation to a glass transition. Journal of Statistical Mechanics: Theory and Experiment, 2016(5):054047, 2016.
- [22] David A Huse, Rahul Nandkishore, and Vadim Oganesyan. Phenomenology of fully many-body-localized systems. *Physical Review B*, 90(17):174202, 2014.
- [23] François Huveneers. Drastic fall-off of the thermal conductivity for disordered lattices in the limit of weak anharmonic interactions. *Nonlinearity*, 26(3):837, 2013.
- [24] John Z Imbrie. On many-body localization for quantum spin chains. Journal of Statistical Physics, 163(5):998– 1048, 2016.
- [25] Yu Kagan and LA Maksimov. Localization in a system of interacting particles diffusing in a regular crystal. *Zhurnal Eksperimentalnoi i Teoreticheskoi Fiziki*, 87:348–365, 1984.
- [26] Vedika Khemani, Achilleas Lazarides, Roderich Moessner, and Shivaji L Sondhi. Phase structure of driven quantum systems. *Physical review letters*, 116(25):250401, 2016.
- [27] Vedika Khemani, CW Von Keyserlingk, and Shivaji Lal Sondhi. Defining time crystals via representation theory. *Physical Review B*, 96(11):115127, 2017.
- [28] Achilleas Lazarides, Arnab Das, and Roderich Moessner. Equilibrium states of generic quantum systems subject to periodic driving. *Phys. Rev. E*, 90:012110, Jul 2014.
- [29] Achilleas Lazarides, Arnab Das, and Roderich Moessner. Fate of many-body localization under periodic driving. *Physical review letters*, 115(3):030402, 2015.
- [30] Netanel H Lindner, Erez Berg, and Mark S Rudner. Universal chiral quasisteady states in periodically driven many-body systems. *Physical Review X*, 7(1):011018, 2017.
- [31] Jani Lukkarinen and Herbert Spohn. Weakly nonlinear schrödinger equation with random initial data. *Inven*tiones mathematicae, 183(1):79–188, 2011.
- [32] RS MacKay and S Aubry. Proof of existence of breathers for time-reversible or hamiltonian networks of weakly coupled oscillators. *Nonlinearity*, 7(6):1623, 1994.
- [33] Krishnanand Mallayya, Marcos Rigol, and Wojciech

De Roeck. Prethermalization and thermalization in generic isolated quantum systems. *arXiv preprint* arXiv:1810.12320, 2018.

- [34] Ivar Martin, Gil Refael, and Bertrand Halperin. Topological frequency conversion in strongly driven quantum systems. *Physical Review X*, 7(4):041008, 2017.
- [35] Paolo Pietro Mazza, Gabriele Perfetto, Alessio Lerose, Mario Collura, and Andrea Gambassi. Suppression of transport in nondisordered quantum spin chains due to confined excitations. *Physical Review B*, 99(18):180302, 2019.
- [36] Takashi Mori, Tomotaka Kuwahara, and Keiji Saito. Rigorous bound on energy absorption and generic relaxation in periodically driven quantum systems. *Phys. Rev. Lett.*, 116:120401, Mar 2016.
- [37] Rahul Nandkishore and David A Huse. Many body localization and thermalization in quantum statistical mechanics. Annual Review of Condensed Matter Physics, 6:15–38, 2015.
- [38] Shriya Pai, Michael Pretko, and Rahul M Nandkishore. Localization in fractonic random circuits. *Physical Review X*, 9(2):021003, 2019.
- [39] Pedro Ponte, Z Papić, François Huveneers, and Dmitry A Abanin. Many-body localization in periodically driven systems. *Physical review letters*, 114(14):140401, 2015.
- [40] Jürgen Pöschel. Nekhoroshev estimates for quasiconvex hamiltonian systems. Mathematische Zeitschrift, 213(1):187–216, 1993.
- [41] Jurgen Poschel. A lecture on the classical kam theorem. In Proc. Symp. Pure Math, volume 69, pages 707–732, 2001.
- [42] Abhinav Prem, Jeongwan Haah, and Rahul Nandkishore. Glassy quantum dynamics in translation invariant fracton models. *Physical Review B*, 95(15):155133, 2017.
- [43] Tomaž Prosen. Chaos and complexity of quantum motion. Journal of Physics A: Mathematical and Theoretical, 40(28):7881, 2007.
- [44] Peter Reimann and Lennart Dabelow. Typicality of prethermalization. *Physical review letters*, 122(8):080603, 2019.
- [45] V Ros, M Müller, and A Scardicchio. Integrals of motion in the many-body localized phase. *Nuclear Physics B*, 891:420–465, 2015.
- [46] Mauro Schiulaz, Alessandro Silva, and Markus Müller. Dynamics in many-body localized quantum systems without disorder. *Physical Review B*, 91(18):184202, 2015.
- [47] Maksym Serbyn, Z Papić, and Dmitry A Abanin. Local conservation laws and the structure of the many-body localized states. *Physical review letters*, 111(12):127201, 2013.
- [48] Barry Simon. The statistical mechanics of lattice gases, volume 1. Princeton University Press, 2014.
- [49] Herbert Spohn. Kinetic equations for quantum manyparticle systems. arXiv preprint arXiv:0706.0807, 2007.
- [50] Daniel Ueltschi. Cluster expansions and correlation functions. Moscow Math. J, 4(2):511–522, 2004.
- [51] Szabolcs Vajna, Katja Klobas, Tomaž Prosen, and Anatoli Polkovnikov. Replica resummation of the baker-campbell-hausdorff series. *Physical review letters*, 120(20):200607, 2018.
- [52] Merlijn van Horssen, Emanuele Levi, and Juan P Garrahan. Dynamics of many-body localization in a translation-invariant quantum glass model. *Physical Re*-

view B, 92(10):100305, 2015.

- [53] Frank Wilczek. Quantum time crystals. *Physical review letters*, 109(16):160401, 2012.
- [54] N. Y. Yao, C. R. Laumann, J. I. Cirac, M. D. Lukin, and J. E. Moore. Quasi-many-body localization in translation-invariant systems. *Phys. Rev. Lett.*, 117:240601, 2016.
- [55] Norman Y Yao, Andrew C Potter, I-D Potirniche, and Ashvin Vishwanath. Discrete time crystals: rigidity, criticality, and realizations. *Physical review letters*, 118(3):030401, 2017.
- [56] The way to realize this is to convince oneself that the local terms in  $\hat{D}(t)$  can be completely arbitrary in general, up to the commutation with  $N^{(\alpha)}$

# Supplemental Material

The aim of this SM is to provide a full proof of our results. For reasons of clarity, we repeat the full setup here. Some parameters are defined in a slightly different way, but we hope that the path to the statements in the main text is yet sufficiently direct. Our proofs are based on rigorous implementations of Schrieffer-Wolff transformations. These have a ppeared a lot in mathematical physics, and we have been in particular influenced by [11, 24]. Cluster expansions are then used to resum the resulting expansions, and we use the neat formalism of [50]. From a more direct point of view, the proof is based on techniques used in [1, 17] that ultimately are descendants of KAM theory and Nekoroshev estimates [40]. The application of such ideas in systems with a few degrees of freedom is standard and we do not review it. There are also a few rigorous works where such ideas are used to describe many-body systems at spectral edges (i.e. low energy, low density) where the situation effectively reduces to few-body theory, see e.g. [5, 9, 32]. Rigorous results where such techniques are applied to constrain the dynamics in a genuine many-body setup have come into focus in the last years, see e.g. [12, 14, 17, 19, 23, 24]

## TECHNICAL PRELIMINARIES AND SETUP

#### Spaces

We consider a large but finite graph  $\Lambda$ , equipped with the graph distance. The vertices i of this graph are our 'sites'. There is a finite Hilbert space  $\mathbb{C}^d$  attached to each site  $i \in \Lambda$ , and we take d to be fixed. The total Hilbert space  $\mathcal{H}$  is hence  $(\mathbb{C}^d)^{\otimes_{\Lambda}}$  and we say that an operator  $O = O_S$  in  $\mathcal{B} \equiv \mathcal{B}(\mathcal{H})$  (space of bounded operators on  $\mathcal{H}$ ) is supported in a set S if it is of the form  $O_S \otimes \mathbb{1}_{S^c}$ , with a slight abuse of notation. This is the setting for quantum spin systems. One can also consider lattice fermions, if one makes some modifications in the definition, see . For operators  $O \in \mathcal{B}$ , we use the standard operator norm  $||O|| = \sup_{\psi \in \mathcal{H}, \psi \neq 0} \frac{||O\psi||}{||\psi||}$  where, on the right hand side,  $||\cdot||$  is the Hilbert space norm on  $\mathcal{H}$ . For an operator A, we will freely use the notation  $\mathrm{ad}_A$  to denote the superoperator acting on  $\mathcal{B}$  as

$$\operatorname{ad}_A(B) = [A, B]$$

The 'number' operators  $N^{(\alpha)}, \alpha = 1, \ldots, q$ 

These operators play a central role in our analysis. They are given as sums of local terms  $N = \sum_{S \subset \Lambda} N_S^{\alpha}$  satisfying the following conditions

- 1. All local terms mutually commute:  $[N_S^{\alpha}, N_{S'}^{\alpha'}]$ .
- 2. All of the  $N_S^{\alpha}$  have integer spectrum.
- 3. There is a fixed range R such that  $N_S^{\alpha} = 0$  whenever diam(S) > R.
- 4. There is a local bound  $\sup_{i \in \Lambda} \sum_{S \ni i} ||N_S^{\alpha}|| \le \gamma$ .

As explained in the main text, item 1,2) actually follow from the assumption of having a single operator  $H^0$  with commuting local terms.

With these definitions in hand, we need to refine the notion of support of operators, following [17]. We say that  $O \in \mathcal{B}$  is 'strongly supported' in S if O is supported in S and, for any  $S' \not\subset S$  we have  $[O, N_{S'}^{\alpha}] = 0$ . Here are the important consequences

- 1. For any function f, if O is strongly supported in S, then  $f(ad_N)O$  is strongly supported in S.
- 2. If A, B are strongly supported in  $S_A, S_B$ , then [A, B] is strongly supported in  $S_A \cup S_B$ .

We wirte  $\mathcal{B}_S \subset \mathcal{B}$  for the algebra of operators strongly supported in S.

## The projectors $\mathcal{P}_{\mathbf{m}}$

First, we introduce the superoperator  $\mathcal{N}^{(\alpha)} = \mathrm{ad}_{\mathcal{N}^{(\alpha)}}$  acting on  $\mathcal{B}$ , and, for any  $\mathbf{m} \in \mathbb{Z}^q$ 

$$\mathcal{P}_{\mathbf{m}}(O) = \prod_{\alpha=1}^{q} \frac{1}{2\pi} \int_{0}^{2\pi} \mathrm{se}^{\mathrm{i}s(\mathcal{N}^{(\alpha)} - m_{\alpha})}(O).$$

One checks that

- 1.  $\mathcal{P}_{\mathbf{m}} = \mathcal{P}_{\mathbf{m}}^2$ , i.e.  $\mathcal{P}_{\mathbf{m}}$  is a projector.
- 2.  $\mathcal{P}_{\mathbf{m}}$  is a contraction in the operator norm:  $||\mathcal{P}_{m}(O)|| \leq ||O||$ .
- 3. If O is strongly supported in S, then  $\mathcal{P}_{\mathbf{m}}(O)$  is strongly supported in S.

#### Norms on Hamiltonians

To handle operators G that are sums of local terms, like many-body Hamiltonians, we introduce *local norms*  $|| \cdot ||_{\kappa}$  that are defined pertaining to a representation of G as a sum of local terms

$$G = \sum_{S \in \mathcal{P}_{\Lambda}} G_S, \qquad G_S \in \mathcal{B}_S$$

where  $\mathcal{P}_{\Lambda}$  stands for the set of connected (by adjacency) subsets of  $\Lambda$ . Since a given operators G can always be represented in different ways as a sum over local terms (e.g.  $\sigma_i^z + \sigma_{i+1}^z$  can be viewed as one term with  $S = \{i, i+1\}$ or as two terms with  $S = \{i\}, \{i+1\}$ ), the above definition does not immediately yield a well-defined norm. One could try to take the supremum over all representations but that is cumbersome in practice. Therefore, the standard solution to this problem in mathematical statistical mechanics is to define the functions  $S \mapsto G_S$  as the central objects ("*Potentials*" or "*Interactions*", see e.g. [48]) and to have norms on them, and we will do this here. However, to keep the notation light, we denote these objects by the same symbol G. A further complication is that we consider timedependent Hamiltonians G(t) and hence potentials, but here one can simply take the supremum over t and mostly drop t from the notation. The norm is then, for any decay rate  $\kappa$ ,

$$||G||_{\kappa} \equiv \max_{i \in \Lambda} \sum_{S \in \mathcal{P}_{\Lambda}, S \ni i} e^{\kappa|S|} \sup_{t} ||G_{S}(t)||$$

## STATEMENT OF RESULTS

## Time-dependent results

We assume that our time-dependent Hamiltonian is of the form

$$H(t) = H^0(t) + gW(t),$$

with

$$H^{0}(t) = \mathbf{J}(t) \cdot \mathbf{N} = \sum_{\alpha=1}^{q} J_{\alpha}(t) N^{(\alpha)}$$
(A1)

where

- 1. The operators  $N^{(\alpha)}$  satisfy the conditions above, with parameters R (range) and  $\gamma$  (local bound).
- 2. The local constituents  $W_S(t)$  and the parameters J(t) are periodic in t and piecewise-continuous.
- 3.  $||W||_{\kappa_0} \leq 1.$

4. There are  $\tau, x > 0$  such that, with  $\bar{J} \equiv \frac{1}{T} \int_0^T dt J(t)$ 

$$\inf_{n \in \mathbb{Z}} |T\mathbf{m} \cdot \bar{\mathbf{J}} - 2\pi n| \ge \frac{x}{|\mathbf{m}|^{\tau}}, \qquad \forall \mathbf{m} \in \mathbb{Z}^q$$
(A2)

If  $\mathbf{J}/|\mathbf{J}|$  is sampled uniformly on the unit hypersphere, then such a constant x can be found with probability 1 provided that  $q < \tau$ , see the standard reasoning in the Appendix at the end

The genuine dimensionless small parameter in our theory is then

 $\epsilon \equiv gT(1+\frac{1}{r})$ 

We let U(t) be the solution of the Schrödinger equation

$$i\partial_t U(t) = H(t)U(t), \qquad U(0) = 1$$

**Theorem 3.** There is a T-periodic unitary  $\hat{Y}$  and T-periodic Hamiltonians  $\hat{D}(t), \hat{V}(t)$  such that  $\hat{U}(t) \equiv \hat{Y}(t)U(t)$  solves the Schrödinger equation

$$i\partial_t \hat{U} = (H^0(t) + \hat{D}(t) + \hat{V}(t))\hat{U}(t),$$

where  $\hat{D}$  conserves **N** and  $\hat{V}$  is very weak:

- 1.  $[N^{(\alpha)}, \hat{D}] = 0$  for any  $\alpha = 1, ..., q$ .
- 2.  $||\hat{D}||_{\kappa} \leq Cg$  and  $||\hat{V}||_{\kappa} \leq ge^{-C(1/\epsilon)^{1/p}}$ , with p any number larger than  $q + \tau + 1$ ,  $\kappa = C\kappa_0 |\log \epsilon|^{-1}$  and constants C, c depending only on the parameters  $\kappa_0, \gamma, q$  and on p. (These constants can have different values from line to line.)
- 3. The unitary transformation  $\hat{Y}$  is quasi-local and close to identity in the sense that  $||\hat{Y}G\hat{Y}^{\dagger} G||_{\kappa} \leq C\epsilon ||G||_{\kappa}$  for any Hamiltonian G.

This theorem shows in particular that the dressed operators  $\hat{N}^{(\alpha)}(t) = \hat{Y}(t)N^{(\alpha)}\hat{Y}^*(t)$  are conserved up to very small errors, i.e.

$$\frac{1}{|\Lambda|} \left( ||\hat{N}^{(\alpha)}(t) - U(t)\hat{N}^{(\alpha)}(0)U^{\dagger}(t)|| \right) \le Cg \mathrm{e}^{-c(1/\epsilon)^{1/p}}, \qquad \text{for times } t \le (c/g) \mathrm{e}^{c(1/\epsilon)^{1/p}}$$

This follows by a simple Duhamel estimate, see [1].

#### Static results

We assume that our time-independent Hamiltonian is of the form

$$H = H^0 + gW,$$

with

$$H^{0}(t) = \mathbf{J} \cdot \mathbf{N} = \sum_{\alpha=1}^{q} J_{\alpha}(t) N^{(\alpha)}$$
(A3)

where

- 1. The operators  $N^{(\alpha)}$  satisfy the conditions above, with parameters R (range) and  $\gamma$  (local bound).
- 2.  $||W||_{\kappa_0} \leq 1.$
- 3. There are tau, x > 0y such that

$$\frac{1}{|\mathbf{J}|}|\mathbf{m} \cdot \mathbf{J}| \ge \frac{x}{|\mathbf{m}|^{\tau}}, \qquad \forall \mathbf{m} \in \mathbb{Z}^q$$

If we sample  $\mathbf{J}/|\mathbf{J}|$  uniformly on the hypersphere, then a constant x can be found with probability 1 provided that  $q - 1 < \tau$ . This is of course essentially the same consideration as that following condition (A2) (it would have been exactly the same, save for the dimension, if in(A2) we were to treat  $2\pi/T$  on the same footing as the  $J_{\alpha}$ ). The genuine dimensionless small parameter in our theory is then

$$\epsilon \equiv \frac{g}{|\mathbf{J}|} \left(1 + \frac{1}{x}\right)$$

**Theorem 4.** There is a unitary  $\hat{Y}$  and Hamiltonians  $\hat{D}, \hat{V}$  such that

$$\hat{Y}H\hat{Y}^{\dagger} = H^0 + \hat{D} + \hat{V}$$

where again  $\hat{D}$  conserves **N** and  $\hat{V}$  is very weak:

- 1.  $[N^{(\alpha)}, \hat{D}] = 0$  for any  $\alpha = 1, ..., q$ .
- 2.  $||\hat{D}||_{\kappa} \leq Cg$  and  $||\hat{V}||_{\kappa} \leq g e^{-c(1/\epsilon)^{1/p}}$  with p any number larger than  $q + \tau + 1$ ,  $\kappa = C\kappa_0 |\log \epsilon|^{-1}$  and C; c depending only on the parameters  $\kappa_0, \gamma, q$  and on p.
- 3. The unitary transformation  $\hat{Y}$  is quasi-local and close to identity in the sense that  $||\hat{Y}G\hat{Y}^{\dagger} G||_{\kappa} \leq C\epsilon ||G||_{\kappa}$  for any Hamiltonian G.

This theorem shows in particular that the dressed operators  $\hat{N}^{(\alpha)}(t) = \hat{Y}(t)N^{(\alpha)}\hat{Y}^{\dagger}(t)$  are conserved up to very small errors, i.e.

$$\frac{1}{|\Lambda|} \left( \left| \left| \hat{N}^{(\alpha)}(t) - U(t) \hat{N}^{(\alpha)}(0) U^{\dagger}(t) \right| \right| \right) \le Cg \mathrm{e}^{-c(1/\epsilon)^{1/p}}, \qquad \text{for times } t \le c/g \mathrm{e}^{c(1/\epsilon)^{1/p}},$$

This follows by a simple Duhamel estimate, see [1].

#### **PROOFS: ALGEBRA**

We primarily treat the time-periodic case, and afterwards we do the static case, focusing on the few differences.

## **Renormalization of Hamiltonians**

We start from

$$H = H^0 + gW = H^0 + V_1 + D_1$$

where we put  $D_1 = \mathcal{P}_0(gW), V_1 = gW - D_1$  Since we anticipate an interative procedure, we put immediately

$$H_n = H^0 + V_n + D_n,$$

where  $\mathcal{P}_0(V_n) = 0$  and  $\mathcal{P}_0(D_n) = D_n$ . We perform a unitary transformation  $e^{A_n}$  with  $A_n$  anti-Hermitian, and we write

$$e^{A_n} \left( i\partial_t + H^0 + D_n + V_n \right) e^{-A_n} = i\partial_t + H^0 + D_n + (i\partial_t A_n + [A_n, H^0] + V_n) + W_{n+1}$$
(A4)

where  $W_n$  has been simply defined so as to the make the last equation correct, and we will later derive a precise expression.

We now determine  $A_n$  by the equation (written here immediately for arbitrary n)

$$i\partial_t A_n + [H^0, A_n] + V_n = 0.$$
 (A5)

choosing the solution that makes it periodic, namely

$$A_n(t) = \mathcal{V}_{0,t}(1 - \mathcal{V}_{0,T})^{-1} \int_0^T s \mathcal{V}_{s,T} V_n(s) + \int_0^t s \mathcal{V}_{s,t} V_n(s)$$
(A6)

where  $\mathcal{V}_{t_0,t}$  is the superoperator that is the solution of the equation

$$\partial_t \mathcal{V}_{t_0,t} = \mathrm{iad}_{H^0} \mathcal{V}_{t_0,t}$$

The inverse of  $1 - \mathcal{V}_{0,T}$  is well-defined on operators O such that  $\mathcal{P}_0(O) = 0$ , by the Diophantine condition. This is the reason we consider  $V_n$  instead of  $W_n$ . To proceed, we introduce some more notation. We define the transformations (O is an arbitrary operator)

$$\gamma_n(O) := \mathrm{e}^{-A_n} O \mathrm{e}^{A_n} = \mathrm{e}^{-\mathrm{ad}_{A_n}} O.$$

and

$$\alpha_n(O) := \int_0^1 \operatorname{s} \operatorname{e}^{-sA_n} O \operatorname{e}^{sA_n} = \int_0^1 \operatorname{s} \operatorname{e}^{-s\operatorname{ad}_{A_n}} O.$$

The latter involves a dummy time s that has nothing to do with the cycle time t; the transformation  $\alpha_k$  is defined pointwisely for any t in the cycle. The use of  $\alpha_n$  is that

$$\mathrm{e}^{-A_n}\partial_t \mathrm{e}^{A_n} = \alpha_n(\partial_t A_n)$$

as one easily checks by an explicit calculation. If  $A_n(t)$  for different t would commute among themselves, then we would simply find back the familiar expression  $e^{-A_n} \partial_t e^{A_n} = \partial_t A_n$ . With the help of the above notation, we get

$$W_{n+1} = \alpha_n (\mathrm{ad}_{A_n} H^0) - \mathrm{ad}_{A_n} H^0 + \gamma_n (D_n) - D_n + \mathrm{i}(\alpha_n (\partial_t A_n) - \partial_t A_n)$$
  
=  $-(\alpha_n (V_n) - V_n) + \gamma_n (V_n + D_n) - (V_n + D_n)$  (A7)

where we used (A5) to get the second line. Finally, we see that

$$D_{n+1} = D_n + \mathcal{P}_0(W_{n+1}), \qquad V_{n+1} = (1 - \mathcal{P}_0)(W_{n+1})$$

## **PROOFS: ANALYSIS**

#### Bound on $A_n$ from $V_n$

The most important ingredient is the bound on  $A_n$  from properties of  $V_n$ . We recall that  $V_n = \sum_{S \in \mathcal{P}_\Lambda} V_{n,S}$ , with  $V_{n,S}$  strongly supported in S. Then we have

**Lemma 1.** Write  $||O(\cdot)|| := \sup_t ||O(t)||$ . Then

$$\sup_{t} ||A_{n,S}|| \le C_q (1 + (1/x)(\gamma|S|)^{\tau+q}) T \sup_{t} ||V_{n,S}||$$

where  $A_{n,S}$  is defined by substituting  $V_{n,S}$  for  $V_n$  in (A6) and  $C_q$  only depends on q.

*Proof.* Call  $O = V_{n,S}$  and decompose

$$O = \sum_{\mathbf{m}} O_{\mathbf{m}} := \sum_{\mathbf{m}} \mathcal{P}_{\mathbf{m}}(O)$$

There are norms on local operators that make  $\mathcal{N}$  into a q-tuple of Hermitian operators, and hence, by spectral calculus, for any function  $f: \mathbb{Z}^z \to \mathbb{C}$ 

$$f(\mathcal{N})O = \sum_{\mathbf{m}} f(\mathbf{m})O_{\mathbf{m}}$$
(A8)

Moreover, since  $\mathcal{P}_{\mathbf{m}}$  is contractive in operator norm, we have

$$||f(\mathcal{N})O|| \leq \sum_{\mathbf{m}: O_{\mathbf{m}} \neq 0} |f(\mathbf{m})|||O||.$$

Obviously  $(\mathcal{V}_{0,T}-1)^{-1} = f(\mathcal{N})$  and, by our Diophantine assumption on **J**:

$$|f(\mathbf{m})| \le \max_{n \in \mathbb{Z}} |T\mathbf{m} \cdot \bar{\mathbf{J}} - 2\pi n|^{-1} \le \frac{|\mathbf{m}|^{\tau}}{x}$$

We get then

$$||f(\mathcal{N})O|| \le C_q (1/x) (\gamma|S|)^{\tau+q} ||O|$$

Apart from this, we simply use the estimate  $||\mathcal{V}_{t_0,t}O|| \leq ||O||$  in (A6). This yields the claim.

Note that (A8) seems wasteful at first sight because one would want to replace the sum over **m** by a supremum over **m**, which would be justfied, for example, if the decomposition  $O = \sum_{\mathbf{m}} O_{\mathbf{m}}$  were orthogonal with respect to a scalar product associated to  $|| \cdot ||$ . It is not clear to us to what extent one could improve on this bound.

## Main lemma's

The following lemma is our prime tool. It was proven by cluster expansions in [1]. Since the only difference here is that we consider strong supports rather than the normal support, we omit the proof which carries over line per line.

**Lemma 2.** Let Z, Q be potentials and assume that  $3||Q||_{\kappa} \leq \delta \kappa := \kappa - \kappa'$ . Then

$$||\mathbf{e}^{Q} Z \mathbf{e}^{-Q} - Z||_{\kappa'} \le \frac{18}{\delta \kappa \kappa'} ||Z||_{\kappa} ||Q||_{\kappa}.$$

Since  $||Z||_{\kappa'} \leq ||Z||_{\kappa}$ , we also get

$$||\mathbf{e}^{Q} Z \mathbf{e}^{-Q}||_{\kappa'} \le \left(1 + \frac{18}{\delta \kappa \kappa'} ||Q||_{\kappa}\right) ||Z||_{\kappa}.$$

Furthermore, we also need the following estimate, of which we omit the obvious proof.

**Lemma 3.** Let  $Z, \widetilde{Z}$  be potentials such that for some  $p \geq 1$ 

$$||\tilde{Z}_S|| \le |S|^p ||Z_S||,$$

then, for  $\tilde{\kappa} < \kappa$ 

$$||\widetilde{Z}||_{\widetilde{\kappa}} \le \frac{(p/e)^p}{(\kappa - \widetilde{\kappa})^p} ||Z||_{\kappa}.$$

## Iterating bounds

We define  $\kappa(n)$  for  $n \ge 1$  by

$$\kappa(n) = \frac{\kappa_0}{1 + \log(n)}$$

and

$$\delta\kappa(n) \equiv \min(\kappa(n+\frac{1}{2}) - \kappa(n+1), \kappa(n) - \kappa(n+\frac{1}{2}))$$

so that  $\frac{1}{\delta\kappa(n)} \leq 2n\log^2(n+1)$ . We abbreviate  $||\cdot||_{\kappa(n)}$  by  $||\cdot||_n$  and  $||\cdot||_{\kappa(n+\frac{1}{2})}$  by  $||\cdot||_{n+\frac{1}{2}}$ . We set

 $w(n) := ||W_n||_n, \qquad d(n) := ||D_n||_n.$ 

Using the expression (A7) for  $W_{n+1}$  and Lemma 2, we then get, provided that  $6||A_n||_{n+\frac{1}{2}} \leq \delta\kappa(n)$ 

$$||W_n||_{n+1} \le \frac{18}{\delta\kappa(n)\kappa(n+1)} ||A_n||_{n+\frac{1}{2}} (||D_n||_n + ||V_n||_n)$$

Using Lemma 1 and 3, we have

$$||A_n||_{n+\frac{1}{2}} \le \frac{C(1+\frac{1}{x})T}{(\delta\kappa(n))^{\tau+q}}||V_n||_n$$

where C depends only on  $\kappa_0, \gamma, q$  (also in the equations below). Combining the two previous bounds and using  $||V_n||_n \leq ||D_n||_n$  and  $||V_n||_n \leq ||W_n||_n$ , we get

$$w(n+1) \le C \log^3(n+1)n^{\tau+q+1} [(1+\frac{1}{r})T]w(n)d(n)$$

Similarly, we have

$$d(n+1) \le d(n) + w(n+1)$$

We can continue the iteration provided that  $d(n) \leq Cd(1)$ . Finally, from the definition of the parameter g we have that w(1) = d(1) = g, and so we obtain

$$\frac{w(n+1)}{w(n)} \le C \log^3(n+1)n^{\tau+q+1} [(1+\frac{1}{x})gT]$$

Therefore, the iteration can be continued up to  $n = n_*$  with  $n_*$  the maximal number satisfying  $C \log^3(n_*+1)n_*^{\tau+q+1}\epsilon < 1$ . We then obtain the bound on  $||\hat{V}||_{\kappa} = ||V_{n_*}||_{n_*}$  in the theorem, choosing  $n_* = \lfloor \epsilon^{-1/p} \rfloor$  with  $p > \tau + q + 1$ . The bound showing the proximity of  $\hat{Y}$  to identity follows just as in [1].

#### PROOFS FOR THE STATIC CASE

Recall that here we have (no time-dependence)

$$H = H^0 + V_1 + D_1, \qquad H_n = H^0 + V_n + D_n$$

with  $V_n, D_n$  satisfying  $\mathcal{P}_0(V_n) = 0$  and  $\mathcal{P}_0(D_n) = D_n$ . We perform a unitary transformation  $e^{A_n}$  and we write

$$e^{A_n} \left( H^0 + D_n + V_n \right) e^{-A_n} = H^0 + D_n + \left( [A_n, H^0] + V_n \right) + W_{n+1}$$
(A9)

with  $W_{n+1}$  to be identified later. We determine  $A_n$  by

$$[A_n, H^0] + V_n = 0. (A10)$$

and from here onwards the algebra is identical. Namely we get

$$W_{n+1} = -(\alpha_n(V_n) - V_n) + \gamma_n(V_n + D_n) - (V_n + D_n)$$

Just as in the time-dependent case, the main point is to get a bound on  $||A_n||_n$  from  $||V_n||$ . From (A10), we see that a solution for  $A_n$  is given by

$$A_{n,S} = -(\mathrm{ad}_{H^0})^{-1} V_{n,S}$$

Proceeding similarly as in the time-dependent case, we obtain here

$$||A_{n,S}|| \le ||V_{n,S}|| \sum_{\mathbf{m} \in \mathbb{Z}^{q}, |\mathbf{m}| \le \gamma|S|} \frac{1}{|\mathbf{J} \cdot \mathbf{m}|} \le C(1 + \frac{1}{x}) \frac{1}{|\mathbf{J}|} (\gamma|S|)^{\tau+q-1} ||V_{n,S}||$$

where the last bound follows from the Diophantine condition  $\frac{|\mathbf{J}|}{|\mathbf{J}\cdot\mathbf{m}|} \leq |\mathbf{m}|^{\tau}/x$  and we have written  $(1+\frac{1}{x})$  instead of  $\frac{1}{x}$  simply to remind ourselves that it is impossible to choose x large. If we now copy the steps done in the section, we obtain

$$\frac{w(n+1)}{w(n)} \le C \log^3(n+1) n^{\tau+q} [(1+\frac{1}{x})\frac{g}{|\mathbf{J}|}]$$

and so we can now take  $p > \tau + q$  instead of  $p > \tau + q + 1$ .

## APPENDIX

## **Diophantine conditions**

We recall the Diophantine condition

$$\inf_{n \in \mathbb{Z}} |T\mathbf{m} \cdot \bar{\mathbf{J}} - 2\pi n| \ge \frac{x}{|\mathbf{m}|^{\tau}}$$

To continue, let us fix  $\mathbf{b} = \frac{T}{2\pi} \bar{\mathbf{J}}$  and  $\hat{\mathbf{b}} = \frac{\mathbf{b}}{b}$  with  $b = |\mathbf{b}|$ . We assume  $\hat{\mathbf{b}}$  to be uniformly distributed on the unit hypersphere, and we calculate the probability that, for a given  $\mathbf{m}$ ,

$$\mathbb{P}(\inf_{n\in\mathbb{Z}}|\mathbf{b}\cdot\mathbf{m}-n|\leq\frac{x}{|\mathbf{m}|^{\tau}}).$$

We split  $\hat{\mathbf{b}} = \frac{a}{|\mathbf{m}|}\mathbf{m} + \hat{\mathbf{b}}_{\perp}$  and we evaluate this probability as

$$C_q \sum_n \int da (1-a^2) \frac{q-1}{2} \chi(|(ba|\mathbf{m}|-n)| \le \frac{x}{|\mathbf{m}|^{\tau}})$$
  
$$\le C_q \sum_n \int da (1-a^2) \frac{q-1}{2} \chi(|(a-\frac{n}{b|\mathbf{m}|}| \le \frac{C(q,\tau)x}{b|\mathbf{m}|^{\tau+1}})$$
  
$$\le b|\mathbf{m}| \frac{C(q,\tau)x}{b|\mathbf{m}|^{\tau+1}} = \frac{C(q,\tau)x}{|\mathbf{m}|^{\tau}}$$

and summing this over all of  $\mathbf{m} \in \mathbb{Z}^q$  yields the estimate, for some constant  $C'(q, \tau)$ ,

$$\mathbb{P}\left(\forall \mathbf{m} \in \mathbb{Z}^q : \inf_{n \in \mathbb{Z}} |\mathbf{b} \cdot \mathbf{m} - n| \ge \frac{x}{|\mathbf{m}|^{\tau}}\right) \le 1 - C'(q, \tau)x, \qquad \text{provided that } \tau > q$$

In particular, note that the magnitude  $b = |T\overline{\mathbf{J}}|$  did not influence our bound.