

Minimax Nonparametric Two-Sample Test under Smoothing

Xin Xing*, Zuofeng Shang[†], Pang Du[‡], Ping Ma[§],
Wenxuan Zhong,[§] Jun S. Liu[¶]

Abstract

We consider the problem of comparing probability densities between two groups. A new probabilistic tensor product smoothing spline framework is developed to model the joint density of two variables. Under such a framework, the probability density comparison is equivalent to testing the presence/absence of interactions. We propose a penalized likelihood ratio test for such interaction testing and show that the test statistic is asymptotically chi-square distributed under the null hypothesis. Furthermore, we derive a sharp minimax testing rate based on the Bernstein width for nonparametric two-sample tests and show that our proposed test statistics is minimax optimal. In addition, a data-adaptive tuning criterion is developed to choose the penalty parameter. Simulations and real applications demonstrate that the proposed test outperforms the conventional approaches under various scenarios.

Keywords: minimax optimality; nonparametric test; penalized likelihood ratio test; smoothing splines; two-sample test; Wilks' phenomenon.

*Department of Statistics, Virginia Tech

[†]Department of Mathematical Sciences, New Jersey Institute of Technology

[‡]Department of Statistics, Virginia Tech

[§]Department of Statistics, University of Georgia

[¶]Department of Statistics, Harvard University

1 Introduction

A fundamental problem in statistics is to test whether the probability densities underlying two groups of observed data are the same, which is called the two-sample test. It plays an essential role in different scientific fields ranging from modern biological sciences to deep learning. For instance, in metagenomics studies, comparing densities of specific microbial species (or strains) from different treatment groups helps researchers gain insights on the disease and treatments (Turnbaugh et al., 2009); in genomics, identifying differentially expressed genes between two groups or conditions is fundamental to many downstream analyses (Tang et al., 2009); in machine learning, the two-sample test is becoming an essential component in some deep learning algorithms (Li et al., 2017).

In these modern applications, the underlying distributions usually demonstrate complex patterns, including multi-modality and long-tails. Hence, it is often difficult to specify their distributional families. Classical normality-based tests such as the two-sample t-test (Anderson, 1958) and the Shapiro-Wilk test (Shapiro and Wilk, 1965) are generally inappropriate. Nonparametric approaches are more appealing due to their distribution-free feature. Examples include distance-based tests such as the Kolmogorov-Smirnov (KS) test (Darling, 1957), the Anderson-Darling test (Scholz and Stephens, 1987), and their variants; an alternative direction is using discretization (“slicing”) of continuous random variables. Jiang et al. (2015) proposed the dynamic slicing test (DSLICE), which penalizes the number of slices to regularize the test statistics. Recently, Gretton et al. (2007, 2012) proposed maximum mean discrepancy (MMD) two-sample tests via embedding the probability distribution into a reproducible kernel Hilbert space (RKHS). Eric et al. (2008) proposed the regularized MMD test by regularizing eigenvalues of the kernel matrix. In addition, a class of approaches based on kernel density estimation was proposed (Anderson et al., 1994; Cao and Van Keilegom, 2006; Martínez-Camblor et al., 2008; Zhan and Hart, 2014). One common challenge for MMD based and kernel density-based testing approaches is the choice of tuning parameters, e.g., kernel bandwidth or roughness penalty, since these parameters

sensitively affect the methods' power. Besides, they have some drawbacks when applied to data of long-tailed distributions: since the kernel bandwidth is fixed across the entire sample (Silverman, 1986), they tend to lack power to detect changes in tails. In many applications such as gene expression analyses, metagenomics, and economics, long-tailed distributions are widespread.

To overcome these limitations, we propose a likelihood-based test which can automatically adapt to densities with different shapes and further develop a data-adaptive tuning criteria to choose the penalization parameter. Let X be a continuous random vector and Z be a binary random variable that indicates the group information. Instead of directly comparing the two group densities, we characterize the dependence between X and Z through its log-transformed joint density $\eta(x, z)$ within a space $\mathcal{H}$. The key idea is to uniquely decompose the log-transformed joint density $\eta \in \mathcal{H}$ into the main effects η_X, η_Z and the interaction effect η_{XZ} through a novel probabilistic decomposition of $\mathcal{H}$ so that the magnitude of the interaction exactly quantifies the density difference between two groups. The two-sample test is thus equivalent to the interaction test

$$H_0 : \eta_{XZ}(x, z) = 0 \text{ vs. } H_1 : \eta_{XZ}(x, z) \neq 0. \quad (1.1)$$

We propose a penalized likelihood ratio (PLR) test by evaluating the penalized log-likelihood functional of η under H_0 and H_1 , and establish its null distribution as a chi-square distribution. Compared with distance-based testing methods, the proposed PLR test can be easily generalized to a k -sample test by letting $Z \in \{1, \dots, k\}$. We further propose a data-adaptive rule to select the tuning parameter to guarantee testing optimality. The PLR test makes full use of the distribution information and is sensitive to the density difference between the null and alternative hypotheses.

This work has three main contributions. First, we propose a probabilistic decomposition of the tensor product RKHS in Section 3. Existing references on functional decomposition without considering probabilistic measures (Gu, 2013; Wahba, 1990) mainly focus on estimation while leaving the hypothesis testing an open problem. Embedding the probability

measures of X and Z into the tensor product decomposition of $\mathcal{H}$, we can transform the problem of density comparison to the problem of significance test of the interaction between X and Z , which provides a foundation to establish the minimax testing principle (see Section 4). This new probabilistic decomposition framework can be generalized to a broader class of dependence tests, including higher order independence tests and conditional independence tests, by using the magnitudes of the decomposed terms to measure the corresponding dependency. Second, we establish the minimax lower bound for density comparison problems based on the Bernstein width (Pinkus, 2012). Existing minimax lower bounds of the testing rate are commonly derived based on Gaussian sequence models (Ingster, 1989, 1993; Wei and Wainwright, 2018) in a simple regression setting, thus cannot be adapted to density comparison. In contrast, our result can be easily generalized to a wide range of dependence testing problems. We further prove the PLR based two-sample test is minimax optimal. Third, we reveal an interesting connection between PLR and MMD test in Section 5. We show that the MMD test (with a particularly selected kernel) is exactly the squared norm of the gradient of the log-likelihood ratio. Compared with our proposed PLR test, the log-likelihood ratio without a penalty term does not enjoy the minimax optimality. Parallel to our work, Li and Yuan (2019) proposed a normalized MMD by appropriately choosing scaling parameters of the Gaussian kernel, and established its minimax property. Similar to the original MMD (Gretton et al., 2007), the approach in Li and Yuan (2019) is also based on the fixed kernel bandwidth, which reduces the sensitivity when the underlying densities are long-tailed. However, our proposed approach is based on the penalized likelihood estimators, which can automatically adapt to long-tailed distributions. As shown in various simulation and real data studies in Sections 6 and 7, our proposed test shows a higher power when the underlying densities have complex features such as long-tails and multi-modality.

The penalized likelihood-based estimation is widely used under nonparametric regression settings (Silverman, 1986; Eggermont and LaRiccia, 2001; Wood, 2011). There are

only a few exceptions of applying the likelihood principle to hypothesis testing in nonparametric models. [Fan et al. \(2001\)](#) proposed a generalized likelihood ratio test statistic based on a local polynomial estimator of the regression function. [Shang and Cheng \(2013\)](#) laid out a coherent theory of local and global inference for a smoothing spline regression model. However, these developments all focus on the likelihood-based inference under regression settings. To the best of our knowledge, our proposed PLR test is the first likelihood-based test under nonparametric density estimation framework with an optimality guarantee.

The rest of this paper is organized as follows. In [Section 2](#), we construct our proposed penalized likelihood ratio test. [Section 2](#) introduces the construction of the probabilistic decomposition of tensor product RHKS and main theoretical results, including the asymptotic distribution of the PLR test and its power performance. [Section 4](#) established the minimax lower bound of density comparisons. In [Section 5](#), we build the connection between our PLR test and the MMD test. In [Section 6](#), we demonstrate the finite sample performance of our test through simulation studies. [Section 7](#) is the analysis of two real-world examples using our test. [Section 8](#) contains some discussion. [Section A](#) is the appendix holding the proofs of the main results. Additional proofs for the lemmas are deferred to a supplement document.

2 Penalized Likelihood Ratio for Two-sample Test

The two-sample problem can be stated as follows. Suppose that we have n independent d -dimensional observations X_i 's and the associated labels Z_i 's, where Z_i is either 0 or 1 indicating that X_i is taken from either the population with a probability density function f_0 or another population with a probability density function f_1 . We aim to test whether f_0 and f_1 are the same. Other than a smoothness constraint, we will not impose any other constraints on the probability density functions f_0 and f_1 . For the convenience of presentation, instead of the marginal probability formulation, we consider an equivalent formulation in terms of conditional independence. That is, we have n i.i.d. observations

$\mathbf{Y}_i = (X_i, Z_i)$, $i = 1, \dots, n$, taken from a population $Y = (X, Z)$ with a joint probability density $f(x, z)$. Let $f_{X|Z=z}(x)$ be the conditional density of X given $Z = z$ for $z = 0, 1$. The two-sample problem is equivalent to testing whether X and Z are independent, or whether $f_{X|Z=0}(\cdot) = f_{X|Z=1}(\cdot)$, i.e.,

$$H_0 : f_{X|Z=0}(\cdot) = f_{X|Z=1}(\cdot) \quad v.s. \quad H_1 : f_{X|Z=0}(\cdot) \neq f_{X|Z=1}(\cdot) \quad (2.1)$$

We characterize the dependence between X and Z by their interaction with respect to their joint density, and show that testing the significance of such interaction is equivalent to the two-sample test. In order to characterize the interaction between X and Z , we introduce a probabilistic decomposition. Let $\eta(x, z) = \log(f(x, z))$ be the log-transformed joint density function. We define two averaging operators acting on the bivariate function $\eta(x, z)$. For any x , the operator $\mathcal{A}_x$ maps $\eta(x, z)$ to $\mathbb{E}_X \eta(X, z)$, a function in z ; and for any z , the operator $\mathcal{A}_z$ maps $\eta(x, z)$ to $\mathbb{E}_Z \eta(x, Z)$. The interaction term is then

$$\eta_{XZ}(x, z) = (\mathcal{I} - \mathcal{A}_x)(\mathcal{I} - \mathcal{A}_z)\eta(x, z) \equiv \eta(x, z) - (\mathcal{A}_x \eta)(z) - (\mathcal{A}_z \eta)(x) + \mathcal{A}_x \mathcal{A}_z \eta, \quad (2.2)$$

where $\mathcal{I}$ is the identity operator. Note that (2.2) is essentially derived from a functional ANOVA decomposition of $\eta(x, z)$ where $\mathcal{A}_x \mathcal{A}_z \eta$ is the constant, $(\mathcal{I} - \mathcal{A}_x) \mathcal{A}_z \eta$ and $(\mathcal{I} - \mathcal{A}_z) \mathcal{A}_x \eta$ are respectively the main effects of x and z , and $(\mathcal{I} - \mathcal{A}_x)(\mathcal{I} - \mathcal{A}_z) \eta$ is the interaction effect. A straightforward derivation shows that the two sample test is equivalent to testing whether η_{XZ} is zero or not (Proposition S.4 in the Appendix).

We assume that η is in a reproducing kernel Hilbert space (RKHS) $\mathcal{H}$ and let $\mathcal{H}_0 = \{\eta \in \mathcal{H} \mid \eta_{XZ} = 0\}$ be the subspace of $\mathcal{H}$ containing all the bivariate functions whose ANOVA decomposition has a zero interaction term. Based on Proposition S.4, the two-sample test problem in (2.1) is equivalent to testing

$$H_0 : \eta \in \mathcal{H}_0 \quad v.s. \quad H_1 : \eta \in \mathcal{H} \setminus \mathcal{H}_0. \quad (2.3)$$

Consider estimating η by the minimizer of the penalized likelihood

$$\ell_{n,\lambda}(\eta) = -\frac{1}{n} \sum_{i=1}^n \eta(x_i, z_i) + \sum_{z \in \{0,1\}} \int_{\mathcal{X}} e^{\eta(x,z)} dx + \frac{\lambda}{2} J(\eta), \quad (2.4)$$

where the first two sums from the negative log-likelihood representing the goodness-of-fit, $J(\cdot)$ is a quadratic functional enforcing a roughness penalty on η , and $\lambda > 0$ is a tuning parameter controlling the trade-off. Note that the integrals in (2.4) are to guarantee the unitary constraint of a probability density function (see Theorem 3.1 in Silverman (1982)).

We propose the following penalized likelihood ratio test statistic

$$PLR = 2 \left\{ \sup_{\eta \in \mathcal{H}_0} \ell_{n,\lambda}(\eta) - \sup_{\eta \in \mathcal{H}} \ell_{n,\lambda}(\eta) \right\}, \quad (2.5)$$

where the first and second terms are respectively the optimal penalized likelihoods under the reduced model $\mathcal{H}_0$ and the full model $\mathcal{H}$.

2.1 Penalized likelihood functional under the full model

Under the full model, the minimization of (2.4) is performed in $\mathcal{H}$. Let $\mathcal{H}^{(X)}$ be an RKHS of functions on the marginal domain $\mathbb{R}^d$ and $\mathcal{H}^{(Z)}$ be an RKHS of functions on $\{0, 1\}$. Then the full space $\mathcal{H} = \mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}$ is their tensor product and also an RKHS, where $\otimes$ denotes the tensor product of two linear spaces. Correspondingly, if $K^{(X)}$ and $K^{(Z)}$ are respectively the reproducing kernels (RKs) uniquely associated with the RKHS $\mathcal{H}^{(X)}$ and $\mathcal{H}^{(Z)}$, then the RK for $\mathcal{H}$ is simply the product of $K^{(X)}$ and $K^{(Z)}$, that is,

$$K(\mathbf{Y}_i, \mathbf{Y}_j) = K^{(X)}(X_i, X_j)K^{(Z)}(Z_i, Z_j). \quad (2.6)$$

One example for $\mathcal{H}^{(X)}$ is the m th order homogeneous Sobolev space $\{f \mid f^{(m)} \text{ is square integrable, } f^{(j)} \text{ is absolutely continuous and } f^j(0) = f^j(1) \text{ for } j = 0, 1, \dots, m-1\}$ associated with the kernel $K^{(X)}(X_i, X_j) = 1 + (-1)^{m-1}k_{2m}(X_i - X_j)$, where $k_{2m}(x)$ is the $2m$ -th order scaled Bernoulli polynomial (Abramowitz and Stegun, 1948). When $m = 2$, $k_4(x) = \frac{1}{24}((x - 0.5)^4 - 0.5(x - 0.5)^2 + \frac{7}{240})$ and the corresponding $K^{(X)}$ is known as the homogeneous cubic spline kernel. An example for the discrete kernel is $K(Z_i, Z_j) = \mathbb{1}_{\{Z_i=Z_j\}}$.

By the representer theorem (Kimeldorf and Wahba, 1971), the minimizer $\eta(\cdot)$ of (2.4) has the form

$$\eta(\cdot) = \sum_{i=1}^n K(\mathbf{Y}_i, \cdot) c_i = \zeta^T \mathbf{c}, \quad \forall \eta \in \mathcal{H} \quad (2.7)$$

where $\zeta^T = (K(\mathbf{Y}_1, \cdot), \dots, K(\mathbf{Y}_n, \cdot))$ is the vector of functions obtained from kernel K with its first argument fixed at $\mathbf{Y}_i$, and $\mathbf{c} = (c_1, \dots, c_n)$ is coefficient vector. This representation converts the infinite-dimensional minimization problem of (2.4) with respect to η to the finite-dimensional optimization problem with respect to the coefficient vector $\mathbf{c}$ by solving

$$\hat{\mathbf{c}} = \underset{\mathbf{c}}{\operatorname{argmin}} \left\{ -\frac{1}{n} \mathbf{1}_n^T Q \mathbf{c} + \int_{\mathcal{Y}} \exp\{\zeta^T \mathbf{c}\} dy + \frac{\lambda}{2} \mathbf{c}^T Q \mathbf{c} \right\}. \quad (2.8)$$

where $\mathbf{1}_n$ is a $n \times 1$ vector of ones, $Q \in R^{n \times n}$ is the empirical kernel matrix with its (i, j) -th entry being $Q_{ij} = K(Y_i, Y_j)$, and the second term is the same as the second term in (2.4) with summation and integration over (x, z) replaced by integration over y for the convenience of presentation. The objective function in (2.8) is strictly convex (Tapia and Thompson, 1978). Its optimization with respect to $\mathbf{c}$ can be performed via a standard convex optimization procedure such as the Newton-Raphson algorithm; see, e.g., Gu (2013) and Wang (2011). The integrals in (2.8) can be calculated by numerical integration (see 7.4.2 in Gu (2013) for details). When n is large, the representation (2.7) involves a large number of coefficients, which may lead to numerical instability. To tackle this, one may consider only a subsample of $\{\mathcal{Y}_i : i = 1, \dots, n\}$ to use in the presentation (Ma et al., 2015). In general, we denote by

$$\hat{\eta}_{n,\lambda} = \zeta^T \hat{\mathbf{c}} \quad (2.9)$$

the penalized maximum likelihood estimate under the full model.

2.2 Penalized likelihood functional under the reduced model

Under H_0 in (2.3), we denote $\hat{\eta}_{n,\lambda}^0$ to be the penalized likelihood estimator of η , that is,

$$\hat{\eta}_{n,\lambda}^0 = \underset{\eta \in \mathcal{H}_0}{\operatorname{argmin}} \ell_{n,\lambda}(\eta). \quad (2.10)$$

In Section 3.1, we show that $\mathcal{H}_0$ is also an RKHS equipped with kernel function $K^0(\cdot, \cdot)$, which enables us to use a similar reparameterization trick to solve the problem in (2.10). In the following, we show the expression of the kernel function K^0 .

$$K^0(Y_i, Y_j) = K_0^{(X)}(X_i, X_j) K_0^Z(Z_i, Z_j) + K_1^{(X)}(X_i, X_j) K_0^Z(Z_i, Z_j) + K_0^{(X)}(X_i, X_j) K_1^Z(Z_i, Z_j),$$

where $K_0^{\langle X \rangle}(X_i, X_j) = \mathbb{E}_X[K^{\langle X \rangle}(X, X_j)] + \mathbb{E}_X[K^{\langle X \rangle}(X_i, X)] - \mathbb{E}_{X, \tilde{X}}[K^{\langle X \rangle}(X, \tilde{X})]$, $K_1^{\langle X \rangle} = K^{\langle X \rangle} - K_0^{\langle X \rangle}$, $K_0^{\langle Z \rangle}(Z_i, Z_j) = \omega_{Z_i} + \omega_{Z_j} - \sum_{\ell=0}^1 \omega_\ell^2$, $K_1^{\langle Z \rangle} = K^{\langle Z \rangle} - K_0^{\langle Z \rangle}$, $\omega_0 = P(Z = 0)$, and $\omega_1 = P(Z = 1)$. The detailed derivation of K^0 depends on our proposed probabilistic decomposition of $\mathcal{H}$, and is deferred to Section 3.1.

Similar to (2.7), we apply the representation theorem and have

$$\eta(\cdot) = \sum_{i=1}^n K^0(\mathbf{Y}_i, \cdot) c_i = \zeta_0^T \mathbf{c}_0, \quad \forall \eta \in \mathcal{H}_0. \quad (2.11)$$

The penalized likelihood estimators can be obtained by first solving the quadratic program

$$\hat{\mathbf{c}}_0 = \underset{\mathbf{c}}{\operatorname{argmin}} \left\{ -\frac{1}{n} \mathbf{1}_n^T Q_0 \mathbf{c} + \int_{\mathcal{Y}} \exp\{\zeta_0^T \mathbf{c}\} + \frac{\lambda}{2} \mathbf{c}^T Q_0 \mathbf{c} \right\} \quad (2.12)$$

where the (i, j) -th entry of Q_0 is $K^0(\mathbf{Y}_i, \mathbf{Y}_j)$ and the (i, j) -th entry of Q_0 is $K^0(\mathbf{Y}_i, \mathbf{Y}_j)$. Numerically, We could express

$$\begin{aligned} Q_0 = & [(I_n - H)Q^{\langle X \rangle}(I_n - H)] \circ [(I_n - H)Q^{\langle Z \rangle}(I_n - H)] + [H Q^{\langle X \rangle} H] \circ [(I_n - H)Q^{\langle Z \rangle}(I_n - H)] \\ & + [(I_n - H)Q^{\langle X \rangle}(I_n - H)] \circ [H Q^{\langle Z \rangle} H] \end{aligned}$$

where $Q^{\langle X \rangle}$ is the empirical kernel matrix of $\mathcal{H}^{\langle X \rangle}$ with (i, j) -th entry $Q_{ij}^{\langle X \rangle} = K^{\langle X \rangle}(X_i, X_j)$, $Q^{\langle Z \rangle}$ is the empirical kernel matrix of $\mathcal{H}^{\langle Z \rangle}$ with (i, j) -th entry $Q_{ij}^{\langle Z \rangle} = K^{\langle Z \rangle}(Z_i, Z_j)$, and $H = I_n - \frac{1}{n} \mathbf{1}_n \mathbf{1}_n^T$ for I_n as a $n \times n$ identity matrix and $\mathbf{1}_n$ as a $n \times 1$ vector of ones. Then we solve the quadratic optimization similar to (2.8) and output the function estimates

$$\hat{\eta}_{n,\lambda}^0 = \zeta_0^T \hat{\mathbf{c}}^0. \quad (2.13)$$

2.3 Test statistics

Plugging the maximizers of penalized likelihood functional under full and reduced model into (2.5), we have the penalized likelihood ratio (PLR) statistic

$$PLR_{n,\lambda} = \ell_{n,\lambda}(\hat{\eta}_{n,\lambda}^0) - \ell_{n,\lambda}(\hat{\eta}_{n,\lambda}). \quad (2.14)$$

We will show in Section 3.2 that $PLR_{n,\lambda}$ is asymptotically χ^2 distributed under H_0 in the sense that $(2b_{n,\lambda})^{-1/2}(2PLR_{n,\lambda} - b_{n,\lambda}) \rightarrow N(0, 1)$ with $b_{n,\lambda}$ diverges for a wide range of λ . Also, the distribution of $PLR_{n,\lambda}$ is independent with the nuisance parameters in $\mathcal{H}_0$, which fulfills the Wilks' phenomenon.

For the nonparametric two-sample test, the parameter space under H_0 is infinite-dimensional as $n \rightarrow \infty$. The assumptions of the Neyman-Pearson Lemma cannot be satisfied. Thus the uniformly most powerful test may not exist in general. We evaluate the power performance by the minimax rate of testing, which is defined as the minimal distance between the null and the alternative hypotheses such that valid testing is possible (Ingster, 1989). For any generic 0-1 valued testing rule $\Phi = \Phi(\mathbf{Y}_1, \dots, \mathbf{Y}_n)$ and a separation rate $d_n > 0$, define the *total error* $\text{Err}(\Phi, d_n)$ of Φ under d_n as

$$\text{Err}(\Phi, d_n) = \mathbb{E}_{H_0} \{\Phi\} + \sup_{\|\eta_{XZ}\|_2 \geq d_n} \mathbb{E}_\eta \{1 - \Phi\}, \quad (2.15)$$

where $\mathbb{E}_{H_0} \{\cdot\}$ denotes the expectation under H_0 , d_n measures the distance between the null and the alternative hypotheses. The first and second terms on the right side of (2.15) represent type I and type II errors of Φ respectively. In Section 3, we show that the distinguishable rate of our proposed PLR test is related to tuning parameter λ and derive the optimal distinguishable rate by carefully selecting λ . A data-adaptive tuning method is developed for applications. In Section 4, we will use information theory to establish the minimum separation rate d_n for general testing rules, which extends the minimax testing principle pioneered in Ingster (1989) to density comparison.

3 Theoretical Properties of PLR Test

In this section, we first introduce the probabilistic decomposition of tensor produce RKHS, enabling us to construct kernel on the subspace $\mathcal{H}_0$. Such decomposition is also of independent interest for studying different kinds of dependence among random variables. We derive the null asymptotic distribution of our proposed test statistics and derive the opti-

mal power for the proposed test. Then we develop data-adaptive tuning method to choose the penalty parameter.

3.1 Probabilistic decomposition of the tensor product RKHS

We assume that bi-variate function $\eta(x, z)$ belong to a tensor product RKHS $\mathcal{H} = \mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}$, in which $\mathcal{H}^{(X)}$ and $\mathcal{H}^{(Z)}$ represent the marginal RKHS of X and Z respectively. We aim to decompose $\mathcal{H}$ into orthogonal subspaces with a hierarchical structure similar to the main effects and interactions in smoothing spline ANOVA (Wahba, 1990; Gu, 2013; Lin, 2000; Wang, 2011), while embedding the probabilistic distribution of X and Z into the decomposition. Such decomposition enables us to convert the two-sample test problem into testing whether the interaction presents or not. It includes two steps: decompose each marginal RKHS into mean and main effect; apply distributive law to expand the tensor product of marginal RKHS into a series of subspaces.

We first introduce the probabilistic tensor decomposition of the discrete domain $\mathcal{H}^{(Z)}$ via a probabilistic averaging operator. The kernel on $\mathcal{H}^{(Z)} := \{f : f \in \{0, 1\}\} = \mathbb{R}^2$ with the Euclidean inner product $(\langle \cdot, \cdot \rangle_{\mathcal{H}^{(Z)}})$ is $K^{(Z)}(z, \tilde{z}) = \mathbb{1}_{\{z=\tilde{z}\}}$. Consider a discrete probabilistic measure $\mathbb{P}$ on $\mathcal{Z} = \{0, 1\}$ such that $\mathbb{P}(Z = j) = \omega_j \geq 0$ with $\omega_0 + \omega_1 = 1$. Let $\boldsymbol{\omega} = (\omega_0, \omega_1)$, and define the probabilistic averaging operator as $\mathcal{A}_Z := f \rightarrow \mathbb{E}_Z f(Z) = \langle \boldsymbol{\omega}, f \rangle_{\mathcal{H}^{(Z)}}$. Since $\mathbb{E}_Z[K_Z^{(Z)}] = \boldsymbol{\omega}$, we can rewrite the probabilistic averaging operator as $\mathcal{A}_Z := f \rightarrow \mathbb{E}_Z f(Z) = \langle \mathbb{E}_Z[K_Z^{(Z)}], f \rangle_{\mathcal{H}^{(Z)}}$. Then $\mathbb{E}_Z[K_Z^{(Z)}]$ can be treated as a mean embedding of $\mathbb{P}$ in $\mathcal{H}^{(Z)}$. We further define the tensor sum decomposition of $\mathcal{H}^{(Z)}$ as

$$\mathcal{H}^{(Z)} = \mathcal{H}_0^{(Z)} \oplus \mathcal{H}_1^{(Z)} := \text{span}\{\mathbb{E}_Z K_Z^{(Z)}\} \oplus \{f \in \mathcal{H} : \mathbb{E}_Z\{f(Z)\} = 0\}, \quad (3.1)$$

where $\mathcal{H}_0^{(Z)}$ is the grand mean space, $\mathcal{H}_1^{(Z)}$ is the main effect space. Each subspace in (3.1) is an RKHS with their corresponding kernels stated in Lemma S.1. If more than two samples present, we can extend the decomposition in (3.1) to a general discrete domain where $Z \in \{0, \dots, a\}$ for $a \geq 1$ by considering $\mathcal{H}^{(Z)}$ as an a -dimensional Euclidean space.

Consider the continuous random variable $X \in \mathcal{X}$ and let $\mathbb{P}$ be a probability measure on $\mathcal{X}$. We suppose $\mathcal{H}^{(X)}$ is the m th order Soblev space, i.e.,

$$\mathcal{H}^{(X)} = \{f \in L^2(\mathbb{P}) \mid f^{(j)} \text{ is absolutely continuous and for } j = 0, 1, \dots, m-1, f^{(m)} \in L_2(\mathbb{P})\},$$

with inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}^{(X)}}$. The results also hold for its homogeneous subspace. Let $K^{(X)}$ be the corresponding kernel satisfying $\langle f, K_x^{(X)} \rangle_{\mathcal{H}^{(X)}} = f(x)$ for any $f \in \mathcal{H}^{(X)}$. Similarly, the probabilistic averaging operator is $\mathcal{A}_X := f \rightarrow \mathbb{E}_X f(X) = \mathbb{E}_X \langle K_X^{(X)}, f \rangle_{\mathcal{H}^{(X)}} = \langle \mathbb{E}_X K_X^{(X)}, f \rangle_{\mathcal{H}^{(X)}}$. $\mathbb{E}_X K_X^{(X)}$ has the same role as ω in the Euclidean space. Then, the tensor sum decomposition of a functional space is defined as

$$\mathcal{H}^{(X)} = \mathcal{H}_0^{(X)} \oplus \mathcal{H}_1^{(X)} := \text{span}\{\mathbb{E}_X K_X^{(X)}\} \oplus \{f \in \mathcal{H}^{(X)} : \mathcal{A}_X f = 0\}. \quad (3.2)$$

Analogously, we name $\mathcal{H}_0^{(X)}$ as the grand mean space and $\mathcal{H}_1^{(X)}$ as the main effect space. $\mathbb{E}_X K_X^{(X)}$ is known as the kernel mean embedding which is well established in the statistics literature (Berlinet and Thomas-Agnan, 2011). The construction of kernel functions for $\mathcal{H}_0^{(X)}$ and $\mathcal{H}_1^{(X)}$ are included in S.2.

We are now ready to consider the RKHS $\mathcal{H} = \mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}$ on the product domain $\mathcal{Y} = \mathcal{X} \times \mathcal{Z}$. Applying the distributive rule, the decomposition of $\mathcal{H}$ is written as

$$\mathcal{H} = (\mathcal{H}_0^{(X)} \oplus \mathcal{H}_1^{(X)}) \otimes (\mathcal{H}_0^{(Z)} \oplus \mathcal{H}_1^{(Z)}) \equiv \mathcal{H}_{00} \oplus \mathcal{H}_{10} \oplus \mathcal{H}_{01} \oplus \mathcal{H}_{11}, \quad (3.3)$$

where $\mathcal{H}_{ij} = \mathcal{H}_i^{(X)} \otimes \mathcal{H}_j^{(Z)}$ for $i = 0, 1$ and $j = 0, 1$. Analogous to the classic ANOVA, $\mathcal{H}_{10}$ and $\mathcal{H}_{01}$ are the RKHS's for the main effects, and $\mathcal{H}_{11}$ is the RKHS for the interaction. We call the decomposition of $\mathcal{H}$ in (3.3) as the *probabilistic decomposition* of the tensor product RKHS $\mathcal{H}$ since it embeds the probability measure of the random variable X and Z . Based on Theorems 2.6 in Gu (2013), we can construct the kernels K^{00}, K^{10}, K^{01} and K^{11} for the subspaces $\mathcal{H}_{00}, \mathcal{H}_{10}, \mathcal{H}_{01}$ and $\mathcal{H}_{11}$ accordingly (detailed construction is given in S.3).

3.2 Asymptotic distribution and Wilks' phenomenon

In this subsection, we present the asymptotic distribution of our PLR test (see Theorem 3.5). The proof relies on a technical lemma about the eigen-structures of $\mathcal{H}_0$ and $\mathcal{H}$; see

Lemma 3.1 below. For any $\eta, \tilde{\eta} \in \mathcal{H}$, define

$$\langle \eta, \tilde{\eta} \rangle = V(\eta, \tilde{\eta}) + \lambda J(\eta, \tilde{\eta}), \quad (3.4)$$

where $V(\eta, \tilde{\eta}) = \mathbb{E}_{\eta^*} \{\eta(\mathbf{Y})\tilde{\eta}(\mathbf{Y})\}$ with expectation taken under the true η^* , and J is a bilinear form corresponding to (2.4). It holds that $\mathcal{H}$ and $\mathcal{H}_0$, endowed with the inner product (3.4), are both RKHSs; see Lemma 3.2 in the Appendix. In the following lemma, we characterize the eigenvalues and eigenvectors of the Rayleigh quotient V/J .

Lemma 3.1. (a) *There exist a sequence of functions $\{\xi_p\}_{p=1}^\infty \subset \mathcal{H}$ and a sequence of nonnegative eigenvalues $\{\rho_p\}_{p=1}^\infty$ with $\rho_p \asymp p^{2m/d}$ such that*

$$V(\xi_p, \xi_{p'}) = \delta_{p,p'}, \quad J(\xi_p, \xi_{p'}) = \rho_p \delta_{p,p'}, \quad \text{for all } p, p' \geq 1, \quad (3.5)$$

and that any $\eta \in \mathcal{H}$ can be written as $\eta = \sum_{p=1}^\infty V(\eta, \xi_p) \xi_p$.

(b) *Moreover, there exists a proper subset $\{\rho_p^0, \xi_p^0\}_{p=1}^\infty$ of $\{\rho_p, \xi_p\}_{p=1}^\infty$ satisfying $\{\xi_p^0\}_{p=1}^\infty \subset \mathcal{H}_0$ and for any $\eta \in \mathcal{H}_0$, $\eta = \sum_{p=1}^\infty V(\eta, \xi_p^0) \xi_p^0$. Convergence of both series holds under (3.4).*

(c) $\rho_p^\perp \asymp p^{2m}$, where $\{\rho_p^\perp\}_{p=1}^\infty \subset \{\rho_p\}_{p=1}^\infty$ is a subset of eigenvalues corresponding to $\{\xi_p^\perp\}_{p=1}^\infty \equiv \{\xi_p\}_{p=1}^\infty \setminus \{\xi_p^0\}_{p=1}^\infty$. The set $\{\xi_p^\perp\}_{p=1}^\infty$ generates the orthogonal complement of $\mathcal{H}_0$ under the inner product (3.4).

Lemma 3.1 introduces an eigensystem that simultaneously diagonalizes the bilinear forms V and J . This eigensystem does not depend on the unknown null density, and only depends on the functional space $\mathcal{H}$. Moreover, $\mathcal{H}_0$ can be generated by a proper subset of the eigenfunctions, which is crucial for analyzing the likelihood ratios.

Let $\langle \cdot, \cdot \rangle_0$ denote the restriction of $\langle \cdot, \cdot \rangle$ on the subspace $\mathcal{H}_0$. Specifically, for any $\eta, \tilde{\eta} \in \mathcal{H}_0$, $\langle \eta, \tilde{\eta} \rangle_0 = \langle \eta, \tilde{\eta} \rangle$. Then $\mathcal{H}$ and $\mathcal{H}_0$ are both RKHS's endowed with these inner products.

Lemma 3.2. *$(\mathcal{H}, \langle \cdot, \cdot \rangle)$ and $(\mathcal{H}_0, \langle \cdot, \cdot \rangle_0)$ are both RKHS's with the corresponding inner products.*

Following Lemma 3.2, there exist reproducing kernel functions $\tilde{K}(\cdot, \cdot)$ and $\tilde{K}^0(\cdot, \cdot)$ defined on $\mathcal{Y} \times \mathcal{Y}$ satisfying, for any $\mathbf{y} \in \mathcal{Y}$, $\eta \in \mathcal{H}$, $\tilde{\eta} \in \mathcal{H}_0$:

$$\begin{aligned}\tilde{K}_{\mathbf{y}}(\cdot) &\equiv \tilde{K}(\mathbf{y}, \cdot) \in \mathcal{H}, & \tilde{K}_{\mathbf{y}}^0(\cdot) &\equiv \tilde{K}^0(\mathbf{y}, \cdot) \in \mathcal{H}_0, \\ \langle \tilde{K}_{\mathbf{y}}, \eta \rangle &= \eta(\mathbf{y}), & \langle \tilde{K}_{\mathbf{y}}^0, \tilde{\eta} \rangle_0 &= \tilde{\eta}(\mathbf{y}).\end{aligned}\tag{3.6}$$

We further introduce positive definite self-adjoint operators $W_\lambda : \mathcal{H} \rightarrow \mathcal{H}$ and $W_\lambda^0 : \mathcal{H}_0 \rightarrow \mathcal{H}_0$ such that

$$\begin{aligned}\langle W_\lambda \eta, \tilde{\eta} \rangle &= \lambda J(\eta, \tilde{\eta}) \text{ for all } \eta, \tilde{\eta} \in \mathcal{H}, \\ \langle W_\lambda^0 \eta, \tilde{\eta} \rangle_0 &= \lambda J_0(\eta, \tilde{\eta}) \text{ for all } \eta, \tilde{\eta} \in \mathcal{H}_0,\end{aligned}\tag{3.7}$$

where $J_0(\eta, \tilde{\eta}) = \theta_{01}^{-1} J_{01}(\eta, \tilde{\eta}) + \theta_{10}^{-1} J_{10}(\eta, \tilde{\eta})$ is the restriction of J over $\mathcal{H}_0$. By (3.7) we get $\langle \eta, \tilde{\eta} \rangle = V(\eta, \tilde{\eta}) + \langle W_\lambda \eta, \tilde{\eta} \rangle$, $\langle \eta, \tilde{\eta} \rangle_0 = V(\eta, \tilde{\eta}) + \langle W_\lambda^0 \eta, \tilde{\eta} \rangle_0$. In the following, we give the explicit expression of $\tilde{K}_y(\cdot)$ and $W_\lambda \xi_p(\cdot)$.

Proposition 3.3. *For any $\mathbf{y} \in \mathcal{Y}$ and $\eta \in \mathcal{H}$, we have*

$$\begin{aligned}\|\eta\|^2 &= \sum_{p=1}^{\infty} |V(\eta, \xi_p)|^2 (1 + \lambda \rho_p), \\ \tilde{K}_{\mathbf{y}}(\cdot) &= \sum_{p=1}^{\infty} \frac{\xi_p(\mathbf{y})}{1 + \lambda \rho_p} \xi_p(\cdot), & \tilde{K}_{\mathbf{y}}^0(\cdot) &= \sum_{p=1}^{\infty} \frac{\xi_p^0(\mathbf{y})}{1 + \lambda \rho_p^0} \xi_p^0(\cdot), \\ W_\lambda \xi_p(\cdot) &= \frac{\lambda \rho_p}{1 + \lambda \rho_p} \xi_p(\cdot), & W_\lambda^0 \xi_p^0(\cdot) &= \frac{\lambda \rho_p^0}{1 + \lambda \rho_p^0} \xi_p^0(\cdot).\end{aligned}$$

where $\{\rho_p^0, \xi_p^0\}_{p=1}^{\infty}$ and $\{\rho_p, \xi_p\}_{p=1}^{\infty}$ are eigensystems defined in Lemma 3.1.

As shown in Proposition 3.3, the eigenvalues for $\tilde{K}$ are $\{(1 + \lambda \rho_p)^{-1}\}_{p=1}^{\infty}$, having a slower decay rate due to the scaling by λ . $\tilde{K}$ can be viewed as a scaled kernel comparing with the product kernel $K^{\mathcal{H}} = K^{00} + K^{01} + K^{10} + K^{11}$ introduced in Lemma S.3. Note that

$$\text{trace}(\tilde{K}) = \sum_{p=1}^{\infty} (1 + \lambda \rho_p)^{-1} \asymp \lambda^{-d/(2m)}$$

is the effective dimension that measures the complexity of $\mathcal{H}$; see [Bartlett et al. \(2005\)](#).

Next, we will derive the null asymptotic distribution of the PLR statistics, which relies on the Taylor expansion of the PLR functional. First, we introduce the Frechét derivatives of the log-likelihood functional. Let D, D^2, D^3 be the first-, second- and third-order Frechét derivatives of $\ell_{n,\lambda}(\eta)$. Based on the above notations, these derivatives can be summarized as follows. Let $\mathbf{y} = (x, z)$. For any $\eta, \Delta\eta_1, \Delta\eta_2, \Delta\eta_3 \in \mathcal{H}$,

$$\begin{aligned} D\ell_{n,\lambda}(\eta)\Delta\eta_1 &= -\frac{1}{n} \sum_{i=1}^n \Delta\eta_1(\mathbf{Y}_i) + \int_{\mathcal{Y}} \Delta\eta_1(\mathbf{y}) e^{\eta(\mathbf{y})} d\mathbf{y} + \lambda J(\eta, \Delta\eta_1) \\ &= \left\langle -\frac{1}{n} \sum_{i=1}^n \tilde{K}_{\mathbf{Y}_i} + \mathbb{E}_{\eta} \tilde{K}_{\mathbf{Y}} + W_{\lambda} \eta, \Delta\eta_1 \right\rangle \\ &\equiv \langle S_{n,\lambda}(\eta), \Delta\eta_1 \rangle, \end{aligned} \quad (3.8)$$

$$D^2\ell_{n,\lambda}(\eta)\Delta\eta_1\Delta\eta_2 = \int_{\mathcal{Y}} \Delta\eta_1(\mathbf{y})\Delta\eta_2(\mathbf{y}) e^{\eta(\mathbf{y})} d\mathbf{y} + \lambda J(\Delta\eta_1, \Delta\eta_2), \quad (3.9)$$

$$D^3\ell_{n,\lambda}(\eta)\Delta\eta_1\Delta\eta_2\Delta\eta_3 = \int_{\mathcal{Y}} \Delta\eta_1(\mathbf{y})\Delta\eta_2(\mathbf{y})\Delta\eta_3(\mathbf{y}) e^{\eta(\mathbf{y})} d\mathbf{y}. \quad (3.10)$$

The second equality of (3.8) is due to the reproducing property (3.6) and that

$$\int_{\mathcal{Y}} \Delta\eta(\mathbf{y}) e^{\eta(\mathbf{y})} d\mathbf{y} = \mathbb{E}_{\eta} \Delta\eta_1(\mathbf{Y}) = \mathbb{E}_{\eta} \langle \tilde{K}_{\mathbf{Y}}, \Delta\eta_1 \rangle = \langle \mathbb{E}_{\eta} \tilde{K}_{\mathbf{Y}}, \Delta\eta_1 \rangle.$$

We denote $S_{n,\lambda}(\eta)$ as the score function of the log-likelihood functional $\ell_{n,\lambda}$. Similarly, we define $S_{n,\lambda}^0$ as the score function of the log-likelihood functional $\ell_{n,\lambda}^0$.

Then we have the following Taylor expansion of PLR functional. Let $g = \hat{\eta}_{n,\lambda}^0 - \hat{\eta}_{n,\lambda}$, we have

$$\begin{aligned} PLR_{n,\lambda} &= \ell_{n,\lambda}(\hat{\eta}_{n,\lambda}^0) - \ell_{n,\lambda}(\hat{\eta}_{n,\lambda}) \\ &= D\ell_{n,\lambda}(\hat{\eta}_{n,\lambda})g + \int_0^1 \int_0^1 s D^2\ell_{n,\lambda}(\hat{\eta}_{n,\lambda} + ss'g) g g ds ds' \\ &= \int_0^1 \int_0^1 s \{ D^2\ell_{n,\lambda}(\hat{\eta}_{n,\lambda} + ss'g) g g - D^2\ell_{n,\lambda}(\eta^*) g g \} ds ds' + \frac{1}{2} D^2\ell_{n,\lambda}(\eta^*) g g \\ &\equiv I_1 + I_2 \end{aligned} \quad (3.11)$$

where η^* is the underlying truth. In the proof of Theorem 3.5, we will show that I_2 is a leading term compared with I_1 . From (3.9), we have that $I_2 = \frac{1}{2}\|g\|^2 = \frac{1}{2}\|\hat{\eta}_{n,\lambda}^0 - \hat{\eta}_{n,\lambda}\|^2$. As we will see, the asymptotic distribution of $\|\hat{\eta}_{n,\lambda} - \hat{\eta}_{n,\lambda}^0\|^2$ relies on Bahadur representations of $\hat{\eta}_{n,\lambda}^0$ and $\hat{\eta}_{n,\lambda}$.

We further prove the following Bahadur representations for the difference of the two penalized likelihood estimators, by adapting an empirical processes technique in Shang and Cheng (2013). Lemma 3.4 is crucial for proving Theorem 3.5.

Lemma 3.4. *Suppose $h = \lambda^{\frac{d}{2m}}$ and $nh^2 \rightarrow \infty$. Then we have*

$$n^{1/2}\|\hat{\eta}_{n,\lambda} - \hat{\eta}_{n,\lambda}^0\| = n^{1/2}\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\| + o_P(1).$$

where $S_{n,\lambda}(\eta^*)$ and $S_{n,\lambda}^0(\eta^*)$ are the score functions for $\ell_{n,\lambda}$ and $\ell_{n,\lambda}^0$, respectively.

This lemma shows that the main term I_2 in Taylor's expansion of the PLR functional is determined by the norm of the difference between the score function of $\ell_{n,\lambda}$ and the score function of $\ell_{n,\lambda}^0$. Since the score functions have the explicit expression through Proposition 3.3, we can characterize the null asymptotic distribution of I_2 by the eigensystem introduced in Lemma 3.1.

Before stating our main theorem, we introduce an assumption which is commonly used in literature for deriving the rates of density estimates; see Theorem 9.3 of Gu (2013).

Assumption 1. *There exists a convex set $B \subset \mathcal{H}$ around η^* and a constant $c > 0$ such that, for any $\eta \in B$, $c\mathbb{E}_{\eta^*}\{\tilde{\eta}^2(\mathbf{Y})\} \leq \mathbb{E}_{\eta}\{\tilde{\eta}^2(\mathbf{Y})\}$. Furthermore, with the probability approaching one, $\hat{\eta}_{n,\lambda} \in B$; and under H_0 , with the probability approaching one, $\hat{\eta}_{n,\lambda}^0 \in B$.*

This condition is satisfied when the members of B have uniform upper and lower bounds on the domain $\mathcal{Y}$, as well as that $\hat{\eta}_{n,\lambda}$ and $\hat{\eta}_{n,\lambda}^0$ are stochastically bounded. The following theorem provides the asymptotic distribution for the PLR test statistic under Assumption 1.

Theorem 3.5. Suppose $m \geq 1$ and Assumption 1 holds. Let $h = \lambda^{\frac{d}{2m}}$ and $nh^{2m+d} = O(1)$, $nh^2 \rightarrow \infty$ as $n \rightarrow \infty$. Under H_0 , we have

$$\frac{2n \cdot PLR_{n,\lambda} - \theta_\lambda}{\sqrt{2}\sigma_\lambda} \xrightarrow{d} N(0, 1), \quad n \rightarrow \infty, \quad (3.12)$$

where $\theta_\lambda = \sum_{p=1}^{\infty} \frac{1}{1+\lambda\rho_p^\perp}$, $\sigma_\lambda^2 = \sum_{p=1}^{\infty} \frac{1}{(1+\lambda\rho_p^\perp)^2}$.

We notice that $h \asymp n^{-c}$ with $\frac{1}{2m+d} \leq c \leq \frac{1}{2}$ satisfies the rate conditions in Theorem 3.5, so the asymptotic distribution (3.12) holds under a wide-ranging choice of h . The quantities θ_λ and σ_λ solely depend on the eigenvalues $\rho_p^\perp$'s and λ . Based on (3.12), we propose the following decision rule $\Phi_{n,\lambda}$ at the significance level α :

$$\Phi_{n,\lambda}(\alpha) = \mathbb{1}(|2n \cdot PLR_{n,\lambda} - \theta_\lambda| \geq z_{1-\alpha/2}\sqrt{2}\sigma_\lambda) \quad (3.13)$$

where $\mathbb{1}(\cdot)$ is the indicator function, $z_{1-\alpha/2}$ is the $1 - \alpha/2$ quantile of the standard normal distribution. Hence, we reject H_0 at the significance level α if $\Phi_{n,\lambda} = 1$. Similar Wilks' phenomenon is also observed in the nonparametric/semiparametric regression framework (Fan et al., 2001; Shang and Cheng, 2013). Specifically, let $r_\lambda = \frac{\theta_\lambda}{\sigma_\lambda^2}$, then (3.12) implies that, as $n \rightarrow \infty$,

$$\frac{2nr_\lambda \cdot PLR_{n,\lambda} - r_\lambda\theta_\lambda}{\sqrt{2r_\lambda\theta_\lambda}} \xrightarrow{d} N(0, 1).$$

Therefore, $2nr_\lambda \cdot PLR_{n,\lambda}$ is asymptotically distributed as a χ^2 distribution with degrees of freedom $r_\lambda\theta_\lambda$. In practice, $\rho_p^\perp$'s can be estimated by the sample eigenvalues of the empirical kernel matrix, from which the quantities r_λ and θ_λ can be accurately approximated. Our numerical study in Sections 6 and 7 adopt such an approximation and the performance is satisfactory.

3.3 Power analysis and minimaxity

In this section, we investigate the power of PLR under local alternatives. Define the separation rate as

$$d_n := \sqrt{\lambda + \sigma_\lambda/n}. \quad (3.14)$$

The separation rate is used to measure the distance between the null and the alternative hypotheses. Theorem 3.6 shows that the power of PLR approaches one, provided that the norm of η_{XZ}^* , the interaction term in the probabilistic decomposition of η^* , has a norm bounded below by d_n . The squared separation rate d_n^2 consists of two components: λ representing the squared bias of the estimator, and σ_λ/n with the order of $n^{-1}h^{-1/2}$ representing the standard derivation of $PLR_{n,\lambda}$. Since σ_λ is decreasing with λ , the minimal separation rate for the PLR test is achieved by choosing appropriate λ such that $\lambda \asymp \sigma_\lambda/n$. Our result owes much to the analytic expression of independence (in terms of interactions) based on the proposed probabilistic tensor product decomposition framework.

Let P_{η^*} denote the probability measure induced under η^* , $\|\eta\|_{\text{sup}}$ the supremum norm over $\mathcal{Y}$, and $\|\eta\|_2 = \sqrt{V(\eta)}$.

Theorem 3.6. *Suppose Assumption 1 holds and let d_n be the separation rate defined in (3.14), $m > 3/2$, $\eta^* \in \mathcal{H}$ with $\|\eta_{XZ}^*\|_{\text{sup}} = o(1)$, $J(\eta_{XZ}^*) < \infty$, $\|\eta_{XZ}^*\|_2 \gtrsim d_n$. For any $\varepsilon \in (0, 1)$, there exists a positive N_ε such that, for any $n \geq N_\varepsilon$, $\mathbb{P}_{\eta^*}(\Phi_{n,\lambda}(\alpha) = 1) \geq 1 - \varepsilon$. When $\lambda \asymp \lambda^* \equiv n^{-4m/(4m+d)}$, d_n is upper bounded by $d_n^* \equiv n^{-2m/(4m+d)}$.*

Theorem 3.6 demonstrates that, when $\lambda \asymp \lambda^*$, PLR can successfully detect any local alternatives, provided that they separate from the null at least by d_n^* . In Section 4, we show that this upper bound is unimprovable by establishing the minimax lower bound of distinguishable rate for general two-sample test. It means that no test can successfully detect the local alternatives if they separate from the null by a rate faster than d_n^* . We claim that our PLR test is minimax optimal.

For any $\varepsilon \in (0, 1)$ and $\alpha \in (0, \varepsilon)$, Theorem 3.5 shows that $\mathbb{E}_{H_0}\{\Phi_{n,\lambda^*}(\alpha)\}$ tends to α ; Theorem 3.6 shows that $\mathbb{E}_{\eta^*}\{1 - \Phi_{n,\lambda^*}(\alpha)\} \leq \varepsilon - \alpha$, provided that $\|\eta_{XZ}^*\|_2 \geq C_{\varepsilon-\alpha}d_n^*$ for a large constant $C_{\varepsilon-\alpha}$. That means, asymptotically,

$$\text{Err}(\Phi_{n,\lambda^*}(\alpha), C_{\varepsilon-\alpha}d_n^*) \leq \varepsilon. \quad (3.15)$$

In other words, the total error of PLR is controlled by an arbitrary ε provided that the

null and local alternatives are separated by d_n^* .

3.4 Data-adaptive tuning parameter selection

Smoothing parameter selection plays an important role in nonparametric estimation. Classical methods including generalized cross-validation (GCV) (Craven and Wahba, 1976) and restricted maximum likelihood (REML) (Wood, 2011) provide data-adaptive estimate of the smoothing parameter. However, how to select the smoothing parameter in the nonparametric inference is still an open question. Here, we introduce a data-splitting method to select the smoothing parameter in our proposed PLR test. Here we use the first half of the data to select the smoothing parameter and the second half of the data to calculate the test statistics. Since we divide the data into two independent parts, the selecting event in the first half is independent with the PLR statistics calculated in the second half. When the sample size is small, data-splitting strategy will suffer from the power reduction. An interesting area for further work would be to establish the post-regularization inference which is recently studied in linear models like Lee et al. (2016).

In practice, how to choose the tuning parameter λ is essential to achieve the high power of the proposed test. Theorem 3.6 provides a theoretical guidance that optimal testing rate can be achieved by choosing λ^* to minimize the separation rate d_n defined in (3.14), i.e., satisfying the trade-off between the squared bias of the estimator and the standard deviation of the test statistic. Since d_n is related to the spectral of the population kernel which is usually unknown, we define a sample estimate of d_n by plugging in the empirical eigenvalue of kernel matrix as

$$\hat{d}_n := \sqrt{\lambda + \hat{\sigma}_\lambda}$$

where $\hat{\sigma}_\lambda^2 = \sum_{p=1}^n \frac{1}{(1+\lambda\hat{\rho}_p^2)^2}$ and $\hat{\rho}_p$, $p = 1, \dots, n$, is the empirical eigenvalue of the kernel matrix K^{11} with the ij th entry $K^{11}((x_i, z_i), (x_j, z_j))$. Since $\hat{\sigma}_\lambda$ is a decreasing function of λ , minimizing $\hat{d}_n$ with respect to λ is written as

$$\hat{\lambda}^* = \max \{ \lambda \mid \lambda < \hat{\sigma}_\lambda/n \}. \quad (3.16)$$

We call (3.16) as our data-adaptive criterion for choosing λ . Notice that $\hat{\lambda}^*$ depends on the eigenvalues of the kernel matrix, especially the first few leading eigenvalues. When the sample size is large, we can approximate $\hat{\sigma}_\lambda^2$ via the top eigenvalues, see Drineas and Mahoney (2005) for fast computation of the leading eigenvalues.

4 Minimax Lower Bound of the Distinguishable Rate

For any $\varepsilon \in (0, 1)$, define the minimax separation rate $d_n^\dagger(\varepsilon)$ as

$$d_n^\dagger(\varepsilon) = \inf\{d_n > 0 : \inf_{\Phi} \text{Err}(\Phi, d_n) \leq \varepsilon\}, \quad (4.1)$$

where the infimum in (4.1) is taken over all 0-1 valued testing rules based on samples $\mathbf{Y}_i$'s. Here we consider the local alternative by assuming $\|\eta\|_{\mathcal{H}} < 1/2$. And $d_n^\dagger(\varepsilon)$ characterizes the smallest separation between the null and local alternatives such that there exists a testing approach with a total error of at most ε . Next we establish a lower bound for $d_n^\dagger$, i.e., if d_n is smaller than a certain lower bound, there exists no test that can distinguish the alternative from the null.

We first introduce a geometric interpretation of the hypothesis testing (2.3). Geometrically, $\mathcal{E} = \{\eta \in \mathcal{H} : \|\eta\|_{\mathcal{H}} < 1/2\}$ is an ellipse with axis lengths equal to eigenvalues of $\mathcal{H}$ as shown in Figure 1. For any $\eta \in \mathcal{E}$, the projection of η on $\mathcal{E}_{11} := \mathcal{H}_{11} \cap \mathcal{E}$ is η_{XZ} where $\mathcal{H}_{11}$ is defined in (3.3). The magnitude of the interaction η_{XZ} can be qualified by $\|\eta_{XZ}\|_2$. The distinguishable rate d_n is the radius of the sphere centered at $\eta_{XZ} = 0$ in $\mathcal{E}_{11}$.

Intuitively, the testing will be harder when the projection of η on $\mathcal{H}_{11}$ is closer to the original point $\eta_{XZ} = 0$. We then introduce the Bernstein width in Pinkus (2012) to characterize the testing difficulty. For a compact set C , the Bernstein k -width is defined as

$$b_{k,2}(C) := \arg\max_{r \geq 0} \{\mathbb{B}_2^{k+1}(r) \subset C \cap S \text{ for some subspace } S \in S_{k+1}\} \quad (4.2)$$

where S_{k+1} denotes the set of all $k+1$ dimensional subspace, $\mathbb{B}_2^{k+1}(r)$ is a $(k+1)$ -dimensional L_2 -ball with radius r centered at $\eta_{XZ} = 0$ in $\mathcal{H}_{11}$. Based on the Bernstein width, we give




Figure 1: Geometric interpretation of the distinguishable rate of the testing H_0 .

an upper bound of the testing radius, i.e., for any η projected in the ball with radius less than the certain bound, the total error is larger than $1/2$.

Lemma 4.1. *For any $\eta \in \mathcal{H}$, we have*

$$\text{Err}(\Phi, d_n) \geq 1/2$$

for all

$$d_n \ll r_B(\delta^*) := \sup\{\delta \mid \delta \leq \frac{1}{2\sqrt{n}}(k_B(\delta))^{1/4}\}$$

where $k_B(\delta) := \arg\max_k \{b_{k-1,2}^2(\mathcal{H}_{11}) \geq \delta^2\}$ is the Bernstein lower critical dimension and $r_B(\delta^*)$ is called the Bernstein lower critical radius.

In Lemma 4.1, we show that when d_n is less than $r_B(\delta^*)$, there is no test that can distinguish the alternative from the null. In order to achieve a non-trivial power, we need d_n to be larger than the Bernstein lower critical radius $r_B(\delta^*)$. The critical radius $r_B(\delta^*)$ depends on the shape of the space $\mathcal{H}_{11}$. The lower bound of $k_B(\delta)$ depends on the decay rate of the eigenvalues for $\mathcal{H}_{11}$. According to the Lieb's law, the radius of k -dimensional ball that can be embedded into $\mathcal{H}_{11}$ is determined by k th largest eigenvalue. In Lemma 4.2, we characterize the lower bound of $k_B(\delta)$ by the largest k such that the k th largest eigenvalue is larger than δ^2 .

Lemma 4.2. *Let γ_k be the k th largest eigenvalue of $\mathcal{H}_{11}$. Then we have*

$$k_B(\delta) > \operatorname{argmax}_k \{\sqrt{\gamma_k} \geq \delta\} \quad (4.3)$$

Note that $\gamma_k \asymp k^{-2m/d}$, then $\operatorname{argmax}_k \{\sqrt{\gamma_k} \geq \delta\} \asymp \delta^{-d/m}$. Plug in the lower bound of $k_B(\delta)$ to Lemma 4.1, we achieve $r_B(\delta^*)$, which is the minimax lower bound for the distinguishable rate in the following theorem.

Theorem 4.3. *Suppose $\eta \in \mathcal{H}$. For any $\varepsilon \in (0, 1)$, the minimax distinguishable rate for the testing hypotheses (2.3) is $d_n^\dagger(\varepsilon) \gtrsim n^{-2m/(4m+d)}$.*

Theorem 4.3 provides a general guidance to justify a local minimax test for testing $\eta_{XZ} = 0$. The proof of Theorem 4.3 is presented in Appendix. Comparing $d_n^\dagger(\varepsilon)$ with d_n^* derived in Theorem 3.6, we proved that the PLR test is minimax.

5 Connection to Maximum Mean Discrepancy

We first briefly summarize the maximum mean discrepancy (MMD) proposed in Gretton et al. (2012). Given the kernel function $K^{(X)}$ on $\mathcal{H}^{(X)}$, denote the embedding that maps a probability distribution $f_{X|Z=z}$ into $\mathcal{H}^{(X)}$ by $\mu_z(\cdot) = \int_{\mathcal{X}} K^{(X)}(x, \cdot) f_{X|Z=z}(x) dx$, then the squared MMD between $f_{X|Z=0}$ and $f_{X|Z=1}$ is defined as the squared distance between embeddings of distributions to reproducing kernel Hilbert spaces (RKHS):

$$\begin{aligned} \text{MMD}_b^2(\mathcal{H}^{(X)}; f_{X|Z=0}, f_{X|Z=1}) &= \frac{1}{n_0^2} \sum_{\{i,j \mid Z_i=Z_j=0\}} K_1^{(X)}(X_i, X_j) \\ &\quad - \frac{2}{n_0 n_1} \sum_{\{i,j \mid Z_i \neq Z_j\}} K_1^{(X)}(X_i, X_j) + \frac{1}{n_1^2} \sum_{\{i,j \mid Z_i=Z_j=1\}} K_1^{(X)}(X_i, X_j); \end{aligned} \quad (5.1)$$

where $K_1^{(X)}(X_i, X_j)$ is introduced in Lemma S.2.

We next show that the MMD estimate is equivalent to the squared score function based on the likelihood functional without penalty. Let ℓ_n be the negative likelihood functional

defined as $\ell_n(\eta) = -\frac{1}{n} \sum_{i=1}^n \eta(\mathbf{Y}_i)$, and LR_n be the likelihood ratio functional defined as

$$LR_n(\eta) = \ell_n(\eta) - \ell_n(P_{\mathcal{H}_0}\eta) = -\frac{1}{n} \sum_{i=1}^n \{\eta(\mathbf{Y}_i) - P_{\mathcal{H}_0}\eta(\mathbf{Y}_i)\} = -\frac{1}{n} \sum_{i=1}^n \{\langle K_{\mathbf{Y}_i}^{\mathcal{H}}, \eta \rangle_{\mathcal{H}} - \langle K_{\mathbf{Y}_i}^{\mathcal{H}_0}, \eta \rangle_{\mathcal{H}}\} \quad (5.2)$$

where $P_{\mathcal{H}_0}$ is the projection operator from $\mathcal{H}$ to $\mathcal{H}_0$ and $K^{\mathcal{H}} = K^{00} + K^{01} + K^{10} + K^{11}$ is the kernel for $\mathcal{H}$ and $K^{\mathcal{H}_0} = K^{00} + K^{01} + K^{10}$ is the kernel for $\mathcal{H}_0$.

Now we calculate the Fréchet derivative of the likelihood ratio functional as the score function, i.e.,

$$DLR_n(\eta)\Delta\eta = \langle \frac{1}{n} \sum_{i=1}^n (K_{\mathbf{Y}_i}^{\mathcal{H}} - K_{\mathbf{Y}_i}^{\mathcal{H}_0}), \Delta\eta \rangle_{\mathcal{H}} = \langle \frac{1}{n} \sum_{i=1}^n K_{\mathbf{Y}_i}^{11}, \Delta\eta \rangle_{\mathcal{H}},$$

where K^{11} is the kernel for $\mathcal{H}_{11}$. We further define a score test statistics as the squared $\|\cdot\|_{\mathcal{H}}$ norm of the score function as follows

$$S_n^2 = \left\| \frac{1}{n} \sum_{i=1}^n K_{\mathbf{Y}_i}^{11} \right\|_{\mathcal{H}}^2 = \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n K^{11}(\mathbf{Y}_i, \mathbf{Y}_j), \quad (5.3)$$

where the second equality holds by the reproducing property. Recall that by Lemma S.1 the kernel on $\mathcal{H}_1^{(Z)}$ is $K_1^{(Z)}(Z_i, Z_j) = \mathbb{1}\{Z_i = Z_j\} - \omega_{Z_i} - \omega_{Z_j} + \sum_{l=1}^2 \omega_l^2$, and by Lemma S.2, the kernel on $\mathcal{H}_1^{(X)}$ is $K_1^{(X)}(X_i, X_j) = K(X_i, X_j) - \mathbb{E}_X[K(X, X_j)] - \mathbb{E}_{\tilde{X}}[K(X_i, \tilde{X})] + \mathbb{E}_{X, \tilde{X}}[K(X, \tilde{X})]$. Then we have $K^{11}(\mathbf{Y}_i, \mathbf{Y}_j) = K_1^{(Z)}(Z_i, Z_j)K_1^{(X)}(X_i, X_j)$ based on Lemma S.3. Let $\omega_0 = n_0/(n_0 + n_1)$ and $\omega_1 = n_1/(n_0 + n_1)$ where n_0 is the number of observations in group 0 and n_1 is the number of observations in group 1. Thus, the scaled score test statistic is equivalent to the MMD test statistic, i.e.,

$$\frac{4n_0n_1}{(n_0 + n_1)^2} S_n^2 = \text{MMD}_b^2(\mathcal{H}^{(X)}; f_{X|Z=0}, f_{X|Z=1}) \quad (5.4)$$

under the null hypothesis. When $n_0 = n_1$, i.e. the number of observations are equal in two groups, we have $S_n^2 = \text{MMD}_b^2(\mathcal{H}^{(X)}; f_{X|Z=0}, f_{X|Z=1})$.

The minimax optimality of the score test statistics S_n^2 based on the likelihood ratio is yet unknown. In previous Section 2, we established the minimax optimality of the PLR

test. We further show the difference between the MMD and our proposed PLR statistic. As shown in the proof of Theorem 3.5, the PLR test statistic has an asymptotic expression

$$PLR_{n,\lambda} \sim \|S_{n,\lambda}^0(\eta) - S_{n,\lambda}(\eta)\|^2 \sim \frac{1}{n} \left\| \sum_{i=1}^n \tilde{K}_{\mathbf{Y}_i}^1 \right\|^2, \quad (5.5)$$

where $S_{n,\lambda}$ and $S_{n,\lambda}^0$ are the score functions defined in (3.8) based on the penalized likelihood ratio functional, and $\tilde{K}_{\mathbf{Y}_i}^1 = \tilde{K}_{\mathbf{Y}_i} - \tilde{K}_{\mathbf{Y}_i}^0 = \sum_{p=1}^{\infty} \frac{\xi_p^\perp(\mathbf{Y}_i) \xi_p^\perp}{1 + \lambda \rho_p^\perp}$. Notice that $\tilde{K}^1$ can be viewed as a scaled version of the product kernel K^{11} by replacing the eigenvalues $\{\rho_p^\perp\}$ with $\{1 + \lambda \rho_p^\perp\}$. By choosing $\lambda = \lambda^*$, $\text{trace}(\tilde{K}^1) = \sum_{p=1}^{\infty} \frac{1}{1 + \lambda^* \rho_p^\perp} \asymp n^{2/(4m+d)}$ matches the lower bound of $k_B(d_n^\dagger)$ with $d_n^\dagger = n^{-2m/(4m+d)}$ as the minimax lower bound for the distinguishable rate in Lemma 4.2. In contrast, the MMD is based on kernel K^{11} without regularization, thus the optimality of the power performance cannot be guaranteed.

6 Simulation Study

In this section, we demonstrate the finite sample performance of the proposed test alongside its competitors through a simulation study. We choose the KS test and Anderson-Darling (AD) as two representers of the most popular CDF-based tests, the normalized MMD test (Li and Yuan, 2019) as a representer of kernel-based tests, the ELT (Cao and Van Keilegom, 2006) as a representer of density-based tests, and the dynamic slicing test (DSLICE) (Jiang et al., 2015) as a representer of discretization-based tests. We use the function `ad.test()` provided in the *kSamples* R package for the AD test, conduct the MMD test using the *dHSIC* R package with the default Gaussian kernel, and implement the ELT test using the code provided by the authors. For DSLICE, we follow the authors' suggestions by choosing for each sample size n a penalty parameter so that the rejection region that corresponds to the test statistic being zero is approximate of size $\alpha 100\%$. For our proposed PLR test, we choose the roughness parameter based on the data-adaptive tuning parameter selection criteria in section 3.4.

The samples $\mathbf{Y}_i = (X_i, Z_i)$, $i = 1, \dots, n$, were generated as follows. We first generated

$Z_i \stackrel{iid}{\sim} \text{Bernoulli}(0.5)$, with 0/1 representing the control/treatment group. Then X_i 's were independently generated from the conditional distribution $f_{X|Z}(x)$ in the following Settings 1 and 2. In each setting, we chose the averaged sample size n in each group as 125, 250, 375, 500, 625, 750, 875, 1000. Size and power were calculated as the proportions of rejection based on 1000 independent trials.

Setting 1: we consider the case that X in each group follows the Gaussian distribution with mean zero and a group-specific variance:

$$X \mid Z = z \sim N\left(0, (1 + \delta_1 \mathbb{1}_{z=1})^2\right),$$

where $\delta_1 = 0, 0.2, 0.3$.

Setting 2: We consider the uni-modal Gaussian distribution versus bi-modal Gaussian distribution:

$$X \mid Z = z \sim 0.5 * N\left(-\delta_2 \mathbb{1}_{z=1}, (1 + \delta_2^2 \mathbb{1}_{z=0})\right) + 0.5 * N\left(\delta_2 \mathbb{1}_{z=1}, (1 + \delta_2^2 \mathbb{1}_{z=0})\right)$$

where we set $\delta_2 = 0, 1, 1.2$.

Setting 3: X in the two groups follow Gaussian asymmetric mixture distributions, i.e.,

$$X \mid Z = z \sim 0.5N(2, 1) + 0.5N(-2, (1 - \delta_3 \mathbb{1}_{z=1})^2)$$

where $\delta_3 = 0, 0.3, 0.45$.


Setting 4: we consider symmetric mixtre distributions, i.e.,

$$X \mid Z = z \sim 0.5N(2, (1 - \delta_4 \mathbb{1}_{z=1})^2) + 0.5N(-2, (1 - \delta_4 \mathbb{1}_{z=1})^2)$$


where $\delta_4 = 0, 0.3, 0.6$.

In particular, $\delta_1 = 0$, $\delta_2 = 0$, $\delta_3 = 0$ or $\delta_4 = 0$ corresponds to the true H_0 which will be used to examine the size of the test statistics. Nonzero δ 's are corresponding to different level of heterogeneity between the two groups.


Figures 2 display the powers of the six tests. For Setting 1, Figure 2(a)(b) show that the powers of the PLR, MMD, ELT, AD, and DSLICE tests rapidly approach one when n




(a) Setting 1: $\delta_1 = 0.2$




(b) Setting 1: $\delta_1 = 0.3$




(c) Setting 2: $\delta_2 = 1$




(d) Setting 2: $\delta_2 = 1.2$




(e) Setting 3: $\delta_3 = 0.3$



(f) Setting 3: $\delta_3 = 0.45$



(g) Setting 4: $\delta_4 = 0.3$



(h) Setting 4: $\delta_4 = 0.6$

Figure 2: Power vs. sample size for PLR, KS, MMD, ELT, AD, and DSLICE.




Figure 3: Size vs. sample size for KS, MMD, ELT, AD, DSLICE and PLR tests.

or δ_1 increases. The power of the KS test increases slightly slower than the other five tests. DSLICE appears to be slightly less powerful than the other four tests, maybe because of its discrete nature and its challenges in choosing a proper penalization parameter in their penalized slicing approach. In Setting 2, as shown on 2(c)(d), the MMD and PLR test shows comparable power. PLR test has slightly higher power when the heterogeneity is higher. The distinguishable of DSLICE and ELT increases as δ_2 increases. AD and K-S show significantly lower power. For Setting 3, Figure 2(e)(f) shows again that the PLR test has the highest power. DSLICE performs quite well here, maybe due to its flexibility in slicing. In contrast, the powers of KS, MMD, ELT, AD are significantly lower than both PLR and DSLICE. In Setting 4, PLR and DSLICE shows the similar power in Figure 2(g)(h). The power of MMD, K-S and AD tests are significantly lower than the others. The results demonstrate that both PLR and DSLICE are more adaptive to differently shaped distributions than the other four methods, while PLR enjoys additional advantages than DSLICE when the underlying distribution is smooth.

Figure 3 displays the size of KS, MMD, ELT, AD, DSLICE and PLR tests. It can be seen that the sizes of the six tests are around to the nominal level 0.05 in Setting 1&2, confirming that all tests are asymptotically valid. In Setting 3&4, the size of the PLR test

is still asymptotically correct, and that for DSLICE is reasonably close; while the sizes of KS, MMD, and ELT are way below 0.05, showing that these three tests are too conservative in handling bimodal distributions.

7 Real Data Analysis

In this section, two real-world applications are provided to compare our PLR test with KS and MMD tests.

Metagenomic analysis of type II diabetes: The gut microbiota influences numerous biological functions throughout the body. Recent studies have indicated that gut microbiota plays an important role in many human diseases such as obesity and diabetes. The association between disease and gut microbial composition has been reported in many studies (Turnbaugh et al., 2009; Qin et al., 2012). Due to the rapid development of metagenomics, it is possible to study the DNA through environmental samples directly. Compared with traditional culture-based methods, metagenomics can study unculturable microorganisms. Recently, several metagenomic binning algorithms such as MetaGen (Xing et al., 2017) were proposed to estimate the abundance of microbial species with high accuracy. As observed in Turnbaugh et al. (2009), the microbial distributions demonstrate large cross-individual differences since there are many environmental factors, such as age and antibiotic usage, that could alter the distribution of gut microbiota. A powerful test that can detect such distributional differences would be useful in metagenomic analysis.

This study aims to detect whether the microbial species have different distributions between case and control groups. For a particular microbial species, let X_i be the log-transformed abundance for the i th individual, and let $Z_i = 1/0$ represent the case/control group. We applied the proposed PLR test to a metagenomic data set with 145 sequenced gut microbial DNA samples from 71 T2D patients (case group) and 74 individuals unaffected by T2D (control group) using Illumina Genome Analyzer and obtained 378.4 gigabase paired-end reads. We used MetaGen (Xing et al., 2017) to do the metagenomic binning in

which DNA fragments were clustered into species-level bins and estimated the abundance of 2450 identified species bins. We applied the KS, MMD, and PLR tests on 1005 species clusters with abundance larger than 1% of the averaged abundance in more than 50% of the total samples. The 1005 p-values were calculated by KS, MMD and PLR for each species. We adjusted the p-values by the Benjamini-Hochberg method (Benjamini and Hochberg, 1995). Through controlling the false discovery at 5%, we compared the identified species from the three methods in Figure 4(A). The PLR test identified 101 species, the KS test identified 4 species, and the MMD test identified 13 species. The species identified by PLR cover those by KS or MMD.




Figure 4: (A). A Venn diagram showing the numbers of species identified by PLR, KS and MMD. (B). Densities of log-transformed abundance for *Roseburia intestinalis* in case/control status. (C). Densities of log-transformed abundance for *Faecalibacterium prauitzii* in case/control status. (D). A Venn diagram showing the numbers of genes selected by PLR, KS and MMD. (E). Densities of gene expression levels from *TGFB2* in mutated/unmutated status. (F). Densities of gene expression levels from *DTX1* in mutated/unmutated status.

Moreover, we highlighted two species that were only identified by the PLR test in Figure 4(B-C). The densities of these two species are both bimodal in both the case and control groups. Figure 4(B) plots the conditional density of the log-transformed abundance of *Roseburia intestinalis*. The majority of the case group has a significantly low abundance.

In Figure 4(C), the other species, *Faecalibacterium prausnitzii* has lower abundance for a subgroup of patients in the case group. Both species are butyrate-producing bacteria which can exert profound immunometabolic effects, and thus are probiotic less abundant in T2D patients. Our finding is consistent with [Tilg and Moschen \(2014\)](#) who also observed that the two species' concentrations are lower in T2D subjects.

Gene expression of Chronic Lymphocytic Leukaemia: Chronic lymphocytic leukaemia (CLL), the most common leukaemia among adults in Western countries, is a heterogeneous disease with variable clinical presentation and evolution. Studies have shown that CLL patients with a mutated Immunoglobulin Heavy Chain Variable (IGHV) gene have a much more favorable outcome and low probability of developing progressive disease. In contrast, those with the unmutated IGHV gene are much more likely to develop the progressive disease and have shorter survival. The molecular changes leading to the pathogenesis of the disease are still poorly understood. To further investigate the role of the mutation status in IGHV gene, we aimed to test whether the distributions of the gene expressions are the same between the IGHV mutated and the IGHV unmutated patients.

This study considered a data set of 225 CLL patients in which 131 were IGHV mutated and 85 were IGHV unmutated. The Affymetrix technique measured the gene expressions in which proper quality control and normalization methods were performed ([Maura et al., 2015](#)). We used the Log2-transformed expression value extracted from the CEL files as the measurement of the expression level. For the i th subject, let X_i denote the expression level and Z_i denote the IGHV mutation status. In particular, $Z_i = 0$ denotes the unmutated status, and $Z_i = 1$ denotes the mutated status. We aimed to test $H_0 : f_{X|Z=0}(x) = f_{X|Z=1}(x)$, i.e. whether the gene expression level's conditional densities are the same between the two IGHV mutation status. Rejection of H_0 implies that the gene expression level distribution varies significantly across the mutation status.

We applied the PLR, KS, and MMD tests to the 18863 genes. We performed the Bonferroni correction on the p-values to control the false discovery rate less than 0.05.

The three methods selected 1071 genes, 275 genes and 412 genes, respectively. Results are summarized in a Venn diagram (Figure 4(D)) which demonstrates that the genes selected by PLR cover those selected by KS and MMD. There were 272 genes selected by all methods and 412 genes selected by both PLR and MMD. For instance, TGFB2 was missed by KS but discovered by PLR and MMD. In literature, it has been verified by real-time quantitative PCR (Bomben et al., 2007) that TGFB2 is down-regulated in IGHV mutated CLL cases compared with IGHV unmutated cases; see Figure 4(E) for a comparison of the conditional densities from both groups. So PLR and MMD tends to be more sensitive to select informative genes. There were 597 genes, including DTX1, uniquely selected by PLR. DTX1 is a well-established direct target of NOTCH1, which plays a significant role in a variety of developmental processes as well as in the pathogenesis of certain human cancers and genetic disorders (Fabbri et al., 2017); see Figure 4(F) for a comparison of the conditional densities, which indicates the expression levels of DTX1 are significantly different between two groups.

8 Discussion

We proposed a probabilistic decomposition approach for probability densities, and developed the penalized likelihood ratio (PLR) to compare probability densities between groups. As demonstrated in simulation studies, our method performs well under various families of density functions of different modalities. Notably, our test possesses the Wilks' phenomenon and testing minimaxity. Such results are not easy to derive for distance-based methods. Furthermore, the Wilks' phenomenon leads to an easy-to-execute testing rule that does not involve resampling. An additional natural extension is to test the independence or conditional independence between random variables. This can be carried out through a higher-order probabilistic decomposition of tensor product RKHS. A challenge for such an extension is to characterize the properties of the eigenvalues of the functional space spanned by interactions. We will explore this direction in future work.

References

- Abramowitz, M. and Stegun, I. A. (1948). *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55. US Government printing office.
- Anderson, N. H., Hall, P., Titterington, D. M., et al. (1994). Two-sample test statistics for measuring discrepancies between two multivariate probability density functions using kernel-based density estimates. *Journal of Multivariate Analysis*, 50(1):41–54.
- Anderson, T. W. (1958). *An introduction to multivariate statistical analysis*. Wiley, New York.
- Bartlett, P. L., Bousquet, O., and Mendelson, S. (2005). Local rademacher complexities. *The Annals of Statistics*, 33(4):1497–1537.
- Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the royal statistical society. Series B (Methodological)*, pages 289–300.
- Berlinet, A. and Thomas-Agnan, C. (2011). *Reproducing kernel Hilbert spaces in probability and statistics*. Springer Science & Business Media.
- Bomben, R., Dal Bo, M., Capello, D., Benedetti, D., Marconi, D., Zucchetto, A., Forconi, F., Maffei, R., Ghia, E. M., Laurenti, L., et al. (2007). Comprehensive characterization of ighv3-21-expressing b-cell chronic lymphocytic leukemia: an italian multicenter study. *Blood*, 109(7):2989–2998.
- Cao, R. and Van Keilegom, I. (2006). Empirical likelihood tests for two-sample problems via nonparametric density estimation. *Canadian Journal of Statistics*, 34(1):61–77.
- Craven, P. and Wahba, G. (1976). Smoothing noisy data with spline functions: Estimating the correct degree of smoothing by the method of generalized cross-validation. *Numerische Mathematik*, 31.
- Darling, D. A. (1957). The kolmogorov-smirnov, cramer-von mises tests. *The Annals of Mathematical Statistics*, 28(4):823–838.
- de Jong, P. (1987). A central limit theorem for generalized quadratic forms. *Probability Theory and Related Fields*, 75(2):261–277.
- Drineas, P. and Mahoney, M. W. (2005). On the nyström method for approximating a gram matrix for improved kernel-based learning. *Journal of Machine Learning Research*, 6(Dec):2153–2175.

- Eggermont, P. P. B. and LaRiccia, V. N. (2001). *Maximum penalized likelihood estimation*, volume I. Springer-Verlag, New York.
- Eric, M., Bach, F. R., and Harchaoui, Z. (2008). Testing for homogeneity with kernel fisher discriminant analysis. In *Advances in Neural Information Processing Systems*, pages 609–616.
- Fabbri, G., Holmes, A. B., Viganotti, M., Scuoppo, C., Belver, L., Herranz, D., Yan, X.-J., Kieso, Y., Rossi, D., Gaidano, G., et al. (2017). Common nonmutational notch1 activation in chronic lymphocytic leukemia. *Proceedings of the National Academy of Sciences*, 114(14):E2911–E2919.
- Fan, J., Zhang, C., and Zhang, J. (2001). Generalized likelihood ratio statistics and wilks phenomenon. *Annals of statistics*, 29:153–193.
- Gretton, A., Borgwardt, K., Rasch, M., Schölkopf, B., and Smola, A. J. (2007). A kernel method for the two-sample-problem. In *Advances in Neural Information Processing Systems*, pages 513–520.
- Gretton, A., Borgwardt, K. M., Rasch, M. J., Scholkopf, B., and Smola, A. (2012). A kernel two-sample test. *Journal of Machine Learning Research*, 13(Mar):723–773.
- Gu, C. (2013). *Smoothing spline ANOVA models*, volume 297. Springer Science & Business Media.
- Ingster, Y. I. (1987). Minimax testing of nonparametric hypotheses on a distribution density in the l.p metrics. *Theory of Probability & Its Applications*, 31(2):333–337.
- Ingster, Y. I. (1989). Asymptotic minimax testing of independence hypothesis. *Journal of Soviet Mathematics*, 44(4):466–476.
- Ingster, Y. I. (1993). Asymptotically minimax hypothesis testing for nonparametric alternatives. i, ii, iii. *Math. Methods Statist*, 2(2):85–114.
- Jiang, B., Ye, C., and Liu, J. S. (2015). Nonparametric k-sample tests via dynamic slicing. *Journal of the American Statistical Association*, 110(510):642–653.
- Kimeldorf, G. and Wahba, G. (1971). Some results on tchebycheffian spline functions. *Journal of mathematical analysis and applications*, 33(1):82–95.
- Lee, J. D., Sun, D. L., Sun, Y., Taylor, J. E., et al. (2016). Exact post-selection inference, with application to the lasso. *The Annals of Statistics*, 44(3):907–927.

- Li, C.-L., Chang, W.-C., Cheng, Y., Yang, Y., and Póczos, B. (2017). Mmd gan: Towards deeper understanding of moment matching network. In *Advances in Neural Information Processing Systems*, pages 2203–2213.
- Li, T. and Yuan, M. (2019). On the optimality of gaussian kernel based nonparametric tests against smooth alternatives. *arXiv preprint arXiv:1909.03302*.
- Lin, Y. (2000). Tensor product space anova models. *Annals of Statistics*, pages 734–755.
- Ma, P., Huang, J. Z., and Zhang, N. (2015). Efficient computation of smoothing splines via adaptive basis sampling. *Biometrika*, 102(3):631–645.
- Martínez-Camblor, P., De Una-Alvarez, J., and Corral, N. (2008). k-sample test based on the common area of kernel density estimators. *Journal of Statistical Planning and Inference*, 138(12):4006–4020.
- Maura, F., Cutrona, G., Mosca, L., Matis, S., Lionetti, M., Fabris, S., Agnelli, L., Colombo, M., Massucco, C., Ferracin, M., et al. (2015). Association between gene and mirna expression profiles and stereotyped subset# 4 b-cell receptor in chronic lymphocytic leukemia. *Leukemia & lymphoma*, 56(11):3150–3158.
- Pinkus, A. (2012). *N-widths in Approximation Theory*, volume 7. Springer Science & Business Media.
- Qin, J., Li, Y., Cai, Z., Li, S., Zhu, J., Zhang, F., Liang, S., Zhang, W., Guan, Y., Shen, D., et al. (2012). A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature*, 490(7418):55.
- Scholz, F. W. and Stephens, M. A. (1987). K-sample anderson–darling tests. *Journal of the American Statistical Association*, 82(399):918–924.
- Shang, Z. and Cheng, G. (2013). Local and global asymptotic inference in smoothing spline models. *The Annals of Statistics*, 41(5):2608–2638.
- Shapiro, S. S. and Wilk, M. B. (1965). An analysis of variance test for normality (complete samples). *Biometrika*, 52(3/4):591–611.
- Silverman, B. W. (1982). On the estimation of a probability density function by the maximum penalized likelihood method. *The Annals of Statistics*, pages 795–810.
- Silverman, B. W. (1986). *Density estimation for statistics and data analysis*, volume 26. CRC press.

- Tang, F., Barbacioru, C., Wang, Y., Nordman, E., Lee, C., Xu, N., Wang, X., Bodeau, J., Tuch, B. B., Siddiqui, A., et al. (2009). mrna-seq whole-transcriptome analysis of a single cell. *Nature methods*, 6(5):377–382.
- Tapia, R. and Thompson, J. (1978). *Nonparametric Probability Density Estimation*. Goucher College Series. Johns Hopkins University Press.
- Tilg, H. and Moschen, A. R. (2014). Microbiota and diabetes: an evolving relationship. *Gut*, 63(9):1513–1521.
- Turnbaugh, P. J., Hamady, M., Yatsunenko, T., Cantarel, B. L., Duncan, A., Ley, R. E., Sogin, M. L., Jones, W. J., Roe, B. A., Affourtit, J. P., et al. (2009). A core gut microbiome in obese and lean twins. *Nature*, 457(7228):480.
- Wahba, G. (1990). *Spline models for observational data*, volume 59. Siam.
- Wang, Y. (2011). *Smoothing splines: methods and applications*. CRC Press.
- Wei, Y. and Wainwright, M. J. (2018). The local geometry of testing in ellipses: Tight control via localized kolmogorov widths. *arXiv:1712.00711*.
- Weinberger, H. F. (1974). *Variational methods for eigenvalue approximation*. SIAM.
- Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(1):3–36.
- Xing, X., Liu, J. S., and Zhong, W. (2017). Metagen: reference-free learning with multiple metagenomic samples. *Genome Biology*, 18(1):187.
- Zhan, D. and Hart, J. (2014). Testing equality of a large number of densities. *Biometrika*, 101(2):449–464.

Supplementary Material

Minimax Nonparametric Two-Sample Test under Smoothing

Section A contains proofs of the main results in Theorem 3.5, 3.6 and 4.3. Proofs of Lemma 3.1, 3.2, 4.1, 4.2, and Proposition 3.3 as well as some auxiliary results, are also included.

Section B contains proofs of auxiliary Lemmas S.5-S.9.

Section C contains additional simulation studies on Beta and Beta mixture distribution.

A Proofs of the Main Results

- Section A.1 includes the notation table.
- Section A.2 includes the proofs of auxiliary lemmas in Section 3: Lemma S.4, Lemma 3.1, Lemma 3.2, Proposition 3.3, and Lemma 3.4.
- Section A.3 includes the proof of main Theorems: Theorem 3.5 and Theorem 3.6.
- Section A.4 includes preliminaries for the minimax lower bound, the proof of Lemma 4.1, Lemma 4.2, Theorem 4.3.

A.1 Notation table

We list the notations in the paper in Table 1.

X	d -dimensional continuous covariate
Z	discrete random variable for the group membership
Y	(X, Z)
$\eta(x, z)$	log-transformed joint density of X, Z
$\mathcal{H}$	tensor product RKHS
$\langle \cdot, \cdot \rangle_{\mathcal{H}}, \ \cdot\ _{\mathcal{H}}$	the inner product and norm under $\mathcal{H}$
$K(\cdot, \cdot)$	kernel function under the norm $\ \cdot\ _{\mathcal{H}}$
$\mathcal{H}^{(X)} = \mathcal{H}_0^{(X)} \oplus \mathcal{H}_1^{(X)}$	marginal RKHS of X
$\mathcal{H}^{(Z)} = \mathcal{H}_0^{(Z)} \oplus \mathcal{H}_1^{(Z)}$	marginal RKHS of Z
$K_i^{(X)}$	kernel function for $\mathcal{H}_i^{(X)}, i = 0, 1$
$K_i^{(Z)}$	kernel function for $\mathcal{H}_i^{(Z)}, i = 0, 1$
$\mathcal{H}_{ij}$	RKHS for intercept, main effects, interaction effect
K^{ij}	kernel function for $\mathcal{H}_{ij}$
$\mathcal{A}$	averaging operator
$\{\mu_i, \phi_i\}_{i=0}^{\infty}$	eigensystem for $\mathcal{H}^{(X)}$
$\{\nu_i, \psi_i\}_{i=0}^{\infty}$	eigensystem for $\mathcal{H}^{(Z)}$
$\ell_{n,\lambda}(\eta)$	negative penalized likelihood function
$\hat{\eta}_{n,\lambda}^0$	penalized likelihood estimator of η under H_0
$\hat{\eta}_{n,\lambda}$	penalized likelihood estimator of η in $\mathcal{H}$
$\langle \cdot, \cdot \rangle, \ \cdot\ $	embedded inner product and norm in $\mathcal{H}$
$\langle \cdot, \cdot \rangle_0, \ \cdot\ _0$	embedded inner product and norm in $\mathcal{H}_0$ under H_0
$V(\cdot, \cdot)$	L_2 inner product
$J(\cdot)$	penalty function
$\tilde{K}(\cdot, \cdot)$	kernel function equipped with $\ \cdot\ $ in $\mathcal{H}$
$\tilde{K}^0(\cdot, \cdot)$	kernel function equipped with $\ \cdot\ _0$ in $\mathcal{H}_0$ under H_0
$PLR_{n,\lambda}$	penalized likelihood ratio test statistic
$\ \cdot\ _{sup}$	the supremum norm
W_{λ}	self-adjoint operator satisfies $\langle W_{\lambda}\eta, \tilde{\eta} \rangle = \lambda J(\eta, \tilde{\eta})$
$\{\rho_p, \xi_p\}_{p=1}^{\infty}$	eigensystem that simultaneously diagonalizes V and J in $\mathcal{H}$
$\{\rho_p^0, \xi_p^0\}_{p=1}^{\infty}$	eigensystem that simultaneously diagonalizes V and J in $\mathcal{H}_0$
$\{\rho_p^{\perp}, \xi_p^{\perp}\}_{p=1}^{\infty}$	eigensystem generates the orthogonal complement of $\mathcal{H}_0$
$D\ell_{n,\lambda}, D^2\ell_{n,\lambda}, D^3\ell_{n,\lambda}$	first-, second-, third-order Frechét derivatives of $\ell_{n,\lambda}(\eta)$
$\Phi_{n,\lambda}(\alpha)$	decision rule at the significance level α
$d_n^{\dagger}(\varepsilon)$	minimax separation rate
$LR_n(\eta)$	likelihood ratio function
$\tilde{K}^1(\cdot, \cdot)$	$\tilde{K}(\cdot, \cdot) - \tilde{K}^0(\cdot, \cdot)$ 2

Table 1: A table that lists all useful notation and their meanings.

A.2 Proofs of Lemmas in Section 3

A.2.1 Some Auxiliary Lemmas

We first state some auxiliary lemmas in Lemma S.1, Lemma S.2, and Lemma S.3 to construct kernel functions of the RKHS, which lays the foundation to prove results in Section 3.

Lemma S.1. *For the RKHS $\mathcal{H}^{(Z)}$ on the discrete domain $\{0, 1\}$ with probability measure $\mathbb{P}(Z = z) = \omega_z$ for $z = 0, 1$, there corresponds a unique non-negative definite reproducing kernel $K^{(Z)}$. Based on the tensor sum decomposition $\mathcal{H}^{(Z)} = \mathcal{H}_0^{(Z)} \oplus \mathcal{H}_1^{(Z)}$ where $\mathcal{H}_0^{(Z)} = \{\mathbb{E}_Z[K_Z^{(Z)}]\}$ and $\mathcal{H}_1^{(Z)} = \{f \in \mathcal{H} : \mathbb{E}_Z(f(Z)) = 0\}$, we have that the kernel for $\mathcal{H}_0^{(Z)}$ is*

$$K_0^{(Z)}(z, \tilde{z}) = \omega_z + \omega_{\tilde{z}}$$

and the kernel for $\mathcal{H}_1^{(Z)}$ is

$$K_1^{(Z)}(z, \tilde{z}) = \mathbb{1}_{\{z=\tilde{z}\}} - \omega_z - \omega_{\tilde{z}}$$

where $\mathbb{1}$ is the indicator function.

Lemma S.2. *For the RKHS $\mathcal{H}^{(X)}$ on a continuous domain $\mathcal{X}$ with probability measure $\mathbb{P}$ equipped with inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}^{(X)}}$, there corresponds a unique nonnegative definite reproducing kernel $K^{(X)}$. Based on the tensor sum decomposition $\mathcal{H}^{(X)} = \mathcal{H}_0^{(X)} \oplus \mathcal{H}_1^{(X)}$ where $\mathcal{H}_0^{(X)} = \{\mathbb{E}_X K_X^{(X)}\}$ and $\mathcal{H}_1^{(X)} = \{f \in \mathcal{H} : \mathbb{E}_X(f(X)) = 0\}$, we have that the kernel for $\mathcal{H}_0^{(X)}$ is*

$$K_0^{(X)}(x, \tilde{x}) = \mathbb{E}_X[K(X, \tilde{x})] + \mathbb{E}_{\tilde{X}}[K(x, \tilde{X})] - \mathbb{E}_{X, \tilde{X}}[K(X, \tilde{X})], \quad (\text{A.1})$$

and the kernel for $\mathcal{H}_1^{(X)}$ is

$$\begin{aligned} K_1^{(X)}(x, \tilde{x}) &= \langle K_x^{(X)} - \mathbb{E}_X K_X^{(X)}, K_{\tilde{x}}^{(X)} - \mathbb{E}_{\tilde{X}} K_{\tilde{X}}^{(X)} \rangle_{\mathcal{H}^{(X)}} \\ &= K^{(X)}(x, \tilde{x}) - \mathbb{E}_X[K^{(X)}(X, y)] - \mathbb{E}_{\tilde{X}}[K^{(X)}(x, \tilde{X})] + \mathbb{E}_{X, \tilde{X}}[K^{(X)}(X, \tilde{X})]. \end{aligned}$$

Lemma S.3. *Suppose $K_i^{(X)}$ is the reproducing kernel of $\mathcal{H}_i^{(X)}$ on $\mathcal{X}$, and $K_j^{(Z)}$ is the reproducing kernel of $\mathcal{H}_j^{(Z)}$ on $\mathcal{Z}$ for $i = 0, 1$ and $j = 0, 1$. Then the reproducing kernels of $\mathcal{H}_i^{(X)} \otimes \mathcal{H}_j^{(Z)}$ on $\mathcal{Y} = \mathcal{X} \times \mathcal{Z}$ is $K^{ij}((x, z), (\tilde{x}, \tilde{z})) = K_i^{(X)}(x, \tilde{x}) K_j^{(Z)}(z, \tilde{z})$ with $x, \tilde{x} \in \mathcal{X}$ and $z, \tilde{z} \in \mathcal{Z}$.*

A.2.2 The equivalence between the two-sample test and the interaction test

In the following Proposition S.4, we show that the two-sample test is equivalent to testing whether the interaction η_{XZ} is 0 or not.

Proposition S.4. *Let η be the log-transformed density function of (X, Z) and η_{XZ} be the interaction term defined in (2.2), we have $\eta_{XZ} = 0$ if and only if $f_{X|Z=0}(\cdot) = f_{X|Z=1}(\cdot)$, where $f_{X|Z=z}(x)$ is the conditional density of X given $Z = z$.*

Proof. Write the log-transformed joint density as $\eta(x, z) = \eta_0 + \eta_X(x) + \eta_Z(z) + \eta_{XZ}(x, z)$ according to (3.3). if $\eta_{XZ} = 0$, then $f(x, z) \propto e^{\eta_X(x)} e^{\eta_Z(z)}$, and hence, X, Z are independent.

On the other hand, if X and Z are independent, then the joint density $f(x, z) = f_X(x)f_Z(z)$, where f_X, f_Z are the marginal densities of X and Z . Take log-transformations on both sides, i.e., $\eta(x, z) = \log(f(x, z)) = \log(f_X(x)) + \log(f_Z(z))$. By the decomposition (3.3), we have $\mathcal{A}_X \eta_{XZ} = 0$ and $\mathcal{A}_Z \eta_{XZ} = 0$. If we have $\eta_{XZ} \neq 0$, then $f(x, z)$ can not be factorized. Hence, we have $\eta_{XZ} = 0$

□

A.2.3 Proof of Lemma 3.1

Proof. We aim to construct the eigensystems on the marginal domain $\mathcal{H}^{(X)}$ and $\mathcal{H}^{(Z)}$, based on which the eigensystem on $\mathcal{H}$ will be constructed. First, we consider $\mathcal{X} = [0, 1]^d$. Recall the Sobolev norm $V_X(g_1, g_2) + J_X(g_1, g_2)$ on $\mathcal{H}^{(X)}$. Let $\mathbb{N}_0$ denote the set of non-negative integers. Following Shang and Cheng (2013), we choose the eigenvalues and eigenfunctions of $\mathcal{H}^{(X)}$ as the solution to the following systems of partial differential equations: for integer $k \in \mathbb{N}_0$ and $\alpha_1, \dots, \alpha_d \in \mathbb{N}_0$ satisfying $\alpha_1 + \dots + \alpha_d = m$,

$$(-1)^m \frac{\partial^m}{\partial^{\alpha_1} \dots \partial^{\alpha_d}} \phi_k(x_1, \dots, x_d) = \mu_k f_X(x_1, \dots, x_d) \phi_k(x_1, \dots, x_d) \quad (\text{A.2})$$

with boundary conditions: for any $l = m, \dots, 2m - 1$ and non-negative integers $\beta_1, \dots, \beta_d$ satisfying $\beta_1 + \dots + \beta_d = l$,

$$\frac{\partial^m}{\partial^{\alpha_1} \dots \partial^{\alpha_d}} \phi(x_1, \dots, x_d) = 0 \text{ for } (x_1, \dots, x_d) \in \partial[0, 1]^d,$$

where f_X is the marginal density of X , $\partial[0, 1]^d$ denotes the boundary of $[0, 1]^d$, μ_k 's are non-negative, non-decreasing and normalized so that $V_X(\phi_k, \phi_k) = 1$ for any $k \geq 0$. Simple integration by parts can show that the solutions to (A.2) satisfy $V_X(\phi_k, \phi_{k'}) = \delta_{kk'}$ and

$J_X(\phi_k, \phi_{k'}) = \mu_k \delta_{kk'}$. Meanwhile, the null space has dimension $M = \binom{m+d-1}{d}$, so one has $0 = \mu_0 = \mu_1 = \dots = \mu_{M-1} \leq \mu_M \leq \mu_{M+1} \leq \dots$ with $\mu_k \asymp k^{2m/d}$. Furthermore, one can actually choose $\phi_0 \equiv 1$. To see this, note that $\phi_0, \dots, \phi_{M-1}$ are basis of the null space of monomials on $[0, 1]^d$ with orders up to $m-1$. For $0 \leq k \leq M-1$, there exists $\mathbf{t} = (t_1, \dots, t_d) \in \mathbb{N}_0^d$ satisfying $|\mathbf{t}| \equiv \sum_{l=1}^d t_l < m$ such that one can write $\phi_k(x) \equiv \phi_{\mathbf{t}}(x) = \sum_{i=1}^M a_{i,k} x_1^{t_1} \dots x_d^{t_d}$. For $\mathbf{t}, \mathbf{t}' \in \mathbb{N}_0^d$ satisfying $0 \leq |\mathbf{t}|, |\mathbf{t}'| < m$, define $M_{\mathbf{t}\mathbf{t}'} = \int_{[0,1]^d} x_1^{t_1+t_1'} \dots x_d^{t_d+t_d'} f_X(x) dx$. Let $A_k = (a_{1,k}, \dots, a_{M,k})^T$ and $\mathbf{M} = [M_{\mathbf{t}\mathbf{t}'}]_{|\mathbf{t}|, |\mathbf{t}'|=0}^{m-1}$. Since $V_X(\phi_k, \phi_{k'}) = \delta_{kk'}$ for $k, k' = 1, \dots, M$, we have $A_k^T \mathbf{M} A_{k'} = \delta_{kk'}$. Purposely choose $A_1 = (1, 0, \dots, 0)^T$ and treat the rest $A_2, \dots, A_M$ as unknowns to be determined. This leaves us $M^2 - M$ unknown coefficients and $\frac{M^2+M}{2} - 1$ equations. Since $M^2 - M \geq \frac{M^2+M}{2} - 1$ for any positive integer M , there always exist A_k 's for $k = 2, \dots, M$ that satisfy $A_k^T \mathbf{M} A_{k'} = \delta_{kk'}$. This shows that we can choose $\phi_0 \equiv 1$ while maintaining the simultaneous diagonalization.

The space $\mathcal{H}^{(Z)}$ is an a -dimensional Euclidean space endowed with Euclidean norm. Let $\{\psi_l\}_{l=0}^{a-1}$ denote the orthonormal eigenvectors. The corresponding eigenvalues are $\nu_0 = \dots = \nu_{a-1} = 1$. To see this, note that the reproducing kernel is $R(z, z') = 1 (z = z')$, hence, $\langle R_z, \psi_l \rangle_Z = \psi_l(z)$. On the other hand, $R(z, z') = \sum_{l=0}^{a-1} \nu_l \psi_l(z) \psi_l(z')$, hence, $\langle R_z, \psi_l \rangle_Z = \psi_l(z) \nu_l$, leading to $\nu_l = 1$. For convenience, we choose ψ_0 as constant function, i.e., $\psi_0(z) \equiv 1/\sqrt{a}$ for $z = 1, \dots, a$.

Let $\|\cdot\|_{\mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}}$ denote the tensor product norm induced by $V_X(g_1, g_2) + J_X(g_1, g_2)$ on $\mathcal{H}^{(X)}$ and the Euclidean norm on $\mathcal{H}^{(Z)}$. The marginal basis for $\mathcal{H}^{(X)}$ and $\mathcal{H}^{(Z)}$ naturally provide a basis for the tensor space, i.e., $\{\phi_k \psi_l : k \geq 0, 0 \leq l \leq a-1\}$, that satisfy

$$\langle \phi_k \psi_l, \phi_{k'} \psi_{l'} \rangle_{\mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}} = (1 + \mu_k \nu_l) \delta_{kk'} \delta_{ll'}. \quad (\text{A.3})$$

The right hand side $\mu_k \nu_l$ of (A.3) is the eigenvalue corresponding to basis $\phi_k \psi_l$. Indeed, they form the eigenvalues of the Rayleigh quotient $\|\cdot\|_{L^2(X) \otimes L^2(Z)}^2 / \|\cdot\|_{\mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}}^2$ since ϕ_k and ψ_l are eigenvalues of the marginal Rayleigh quotients; see (Lin, 2000, Section 2.3). We arrange the eigenvalues $\{\mu_k \nu_l\}$ in an increasing order, and denote them as $\pi_1 \leq \pi_2 \leq \dots$, i.e., $\pi_{ra+s} = \mu_r$ for $r \geq 0$ and $1 \leq s \leq a$.

Consider the orthogonal decomposition $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$ in (3.4). By Weinberger (1974), we can use the Rayleigh quotient $V/(V+J)$ to produce $\xi_p^0 \in \mathcal{H}_0$ and $\xi_p^\perp \in \mathcal{H}_1$ with corresponding eigenvalues ρ_p^0 and $\rho_p^\perp$ that satisfy: $V(\xi_p^j, \xi_{p'}^j) = \delta_{pp'}$, $J(\xi_p^j, \xi_{p'}^j) = \rho_p^j \delta_{pp'}$, for $j = 0, \perp$. Let $\{\xi_p\}_{p=1}^\infty = \{\xi_p^0, \xi_p^\perp\}_{p=1}^\infty$ and $\{\rho_p\}_{p=1}^\infty = \{\rho_p^0, \rho_p^\perp\}_{p=1}^\infty$, where ρ_p are arranged in an increasing order. It is easy to verify that ξ_p 's are Rayleigh quotient eigenvalues of

$V/(V+J)$ over $\mathcal{H}$ as defined in (Weinberger, 1974, Section 2). We also have

$$V(\xi_p, \xi_{p'}) = \delta_{pp'}, \quad J(\xi_p, \xi_{p'}) = \rho_p \delta_{pp'}.$$

By (A.6), the Rayleigh quotients corresponding to $(\|\cdot\|_{L^2(X) \otimes L^2(Z)}, \|\cdot\|_{\mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}})$ and $(V, V+J)$ are equivalent. By the Mapping theorem (Weinberger, 1974, Section 3.3), there exist constants $c_1, c_2 > 0$ s.t.

$$\frac{c_1}{1 + \pi_p} \leq \frac{1}{1 + \rho_p} \leq \frac{c_2}{1 + \pi_p}, \quad p \geq 1. \quad (\text{A.4})$$

Following (A.4) we have $\rho_p \asymp \pi_p \asymp p^{2m/d}$. By Fourier expansion, we have $\eta = \sum_{p=1}^{\infty} V(\eta, \xi_p) \xi_p$.

When restricted on $\mathcal{H}_0$, the Rayleigh quotients corresponding to $(V, V+J)$ and $(\|\cdot\|_{L^2(X) \otimes L^2(Z)}, \|\cdot\|_{\mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}})$ are still equivalent. Similar to (A.4), by Mapping theorem,

$$\frac{c_1}{1 + \pi_p^0} \leq \frac{1}{1 + \rho_p^0} \leq \frac{c_2}{1 + \pi_p^0}, \quad p \geq 1. \quad (\text{A.5})$$

where $\{\pi_p^0\}_{p=1}^{\infty} = \{\mu_k, \nu_l : l = 0, \dots, a-1, k \geq 0\}$ are eigenvalues (with increasing order) corresponding to $\{\phi_k, \psi_l : l = 0, \dots, a-1, k \geq 0\}$. Specifically, $\pi_p^0 = \pi_p$ for $p = 1, \dots, a$, and $\pi_{a+s}^0 = \pi_{sa+1}$ for $s \geq 1$. Now remove $\{\pi_p^0\}_{p \geq 1}$ from $\{\pi_p\}_{p \geq 1}$ and denote the rest as $\{\pi_p^\perp\}_{p \geq 1}$. From (A.4) and (A.5), we have

$$\frac{c_1}{1 + \pi_p^\perp} \leq \frac{1}{1 + \rho_p^\perp} \leq \frac{c_2}{1 + \pi_p^\perp}, \quad p \geq 1.$$

Since $\nu_1 = \dots = \nu_{a-1} = 1$ which leads to $\pi_{r(a-1)+s}^\perp = \mu_{r+1}$ for $r \geq 0$ and $s = 1, \dots, a-1$, we have $\rho_p^\perp \asymp \pi_p^\perp \asymp \mu_{\lfloor p/(a-1) \rfloor} \asymp p^{2m/d}$. $\square$

A.2.4 Proof of Lemma 3.2

Proof. Following Gu (2013), $J(\cdot)$ is the roughness penalty, hence it is standard in the sense of Lin (2000). Following Lin (2000), the norm based on $\int_{\mathcal{Y}} \eta(x, z)^2 dx dz + J(\eta)$ is equivalent to $\|\cdot\|_{\mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}}$, where $\|\cdot\|_{\mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}}$ is the tensor product norm induced by the Sobolev norm $V_X(g_1, g_2) + J_X(g_1, g_2)$ on $\mathcal{H}^{(X)}$ and the Euclidean norm on $\mathcal{H}^{(Z)}$. Since $f(x, z)$ is bounded away from zero and infinity, there exist constants $0 < c_1 \leq c_2 < \infty$ such that, for any $\eta \in \mathcal{H}$,

$$c_1 \int_{\mathcal{Y}} \eta(x, z)^2 dx dz \leq V(\eta, \eta) \leq c_2 \int_{\mathcal{Y}} \eta(x, z)^2 dx dz. \quad (\text{A.6})$$

Therefore, $\|\cdot\|$ and $\|\cdot\|_{\mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}}$ are equivalent norms. Since $\mathcal{H}$ endowed with $\|\cdot\|_{\mathcal{H}^{(X)} \otimes \mathcal{H}^{(Z)}}$ is an RKHS, $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ is an RKHS. Since $\mathcal{H}_0$ is a closed subset of $\mathcal{H}$, and $\langle \cdot, \cdot \rangle_0$ is inherited from $\langle \cdot, \cdot \rangle$, we have that $(\mathcal{H}_0, \langle \cdot, \cdot \rangle_0)$ is also an RKHS. $\square$

A.2.5 Proof of Proposition 3.3

Proof. The proof of $\|\eta\|^2 = \sum_{p=1}^{\infty} |V(\eta, \xi_p)|^2 (1 + \lambda \rho_p)$ follows by (3.4) and the Fourier expansion of η : $\eta = \sum_{p=1}^{\infty} V(\eta, \xi_p) \xi_p$. For any $p' \geq 1$,

$$\langle \eta, \xi_{p'} \rangle = \left\langle \sum_{p=1}^{\infty} V(\eta, \xi_p) \xi_p, \xi_{p'} \right\rangle = V(\eta, \xi_{p'}) (1 + \lambda \rho_{p'}). \quad (\text{A.7})$$

By (A.7), $V(\tilde{K}_{\mathbf{y}}, \xi_p) = \frac{\langle \tilde{K}_{\mathbf{y}}, \xi_p \rangle}{1 + \lambda \rho_p} = \frac{\xi_p(\mathbf{y})}{1 + \lambda \rho_p}$. Hence $\tilde{K}_{\mathbf{y}}(\cdot) = \sum_{p=1}^{\infty} \frac{\xi_p(\mathbf{y})}{1 + \lambda \rho_p} \xi_p(\cdot)$ follows. Meanwhile, (A.7) implies that $V(W_{\lambda} \xi_p, \xi_{p'}) = \frac{\langle W_{\lambda} \xi_p, \xi_{p'} \rangle}{1 + \lambda \rho_{p'}} = \frac{\lambda \rho_p \delta_{p,p'}}{1 + \lambda \rho_p}$. Thus we have $W_{\lambda} \xi_p(\cdot) = \frac{\lambda \rho_p}{1 + \lambda \rho_p} \xi_p(\cdot)$.

By Lemma 3.1, any $\eta \in \mathcal{H}_0$ satisfies $\eta = \sum_{p=1}^{\infty} V(\eta, \xi_p^0) \xi_p^0$. Therefore, $V(\tilde{K}_{\mathbf{y}}^0, \xi_p^0) = \langle \tilde{K}_{\mathbf{y}}^0, \xi_p^0 \rangle / (1 + \lambda \rho_p^0)$. Hence, $\tilde{K}_{\mathbf{y}}^0(\cdot) = \sum_{p=1}^{\infty} \frac{\xi_p^0(\mathbf{y})}{1 + \lambda \rho_p^0} \xi_p^0(\cdot)$, and likewise, $W_{\lambda} \xi_p^0(\cdot) = \frac{\lambda \rho_p^0}{1 + \lambda \rho_p^0} \xi_p^0(\cdot)$. $\square$

A.2.6 Proof of Lemma 3.4

We first state and prove several preliminary lemmas. Define

$$h^{-1} = \sum_{p=1}^{\infty} \frac{1}{(1 + \lambda \rho_p)^2}, \quad h_0^{-1} = \sum_{p=1}^{\infty} \frac{1}{(1 + \lambda \rho_p^0)^2}. \quad (\text{A.8})$$

From Lemma 3.1, we have $\rho_p \asymp p^{2m/d}$ and $\rho_p^0 \asymp p^{2m/d}$. The following lemma provides an relation between h (or h_0) and λ .

Lemma S.5. $h \asymp \lambda^{d/2m}$ and $h_0 \asymp \lambda^{d/2m}$.

The following Lemma presents a relationship between the two norms $\|\cdot\|_{\text{sup}}$ and $\|\cdot\|$.

Lemma S.6. *There exists an absolute constant $c_m > 0$ s.t. $\|\eta\|_{\text{sup}} \leq c_m h^{-1/2} \|\eta\|$.*

Proofs of Lemmas S.5 and S.6 can be executed similar to Shang and Cheng (2013).

The following two lemmas characterize the convergence rates of $\hat{\eta}_{n,\lambda}$ and $\hat{\eta}_{n,\lambda}^0$ under H_0 .

Lemma S.7. *Assume $\lambda \rightarrow 0$ and H_0 . Then $\|\hat{\eta}_{n,\lambda}^0 - \eta^*\|_0 = O_P((nh_0)^{-1/2} + \lambda^{1/2})$ and $\|\hat{\eta}_{n,\lambda} - \eta^*\| = O_P((nh)^{-1/2} + \lambda^{1/2})$.*

Lemma S.7 can be proved based on a quadratic approximation method proposed by Gu (2013), i.e., apply (Gu, 2013, Section 9.2.2) to both $(\hat{\eta}_{n,\lambda}, \mathcal{H})$ and $(\hat{\eta}_{n,\lambda}^0, \mathcal{H}_0)$. The optimal rates for both estimators achieve at $h \asymp n^{-1/(2m+d)}$, $h_0 \asymp n^{-1/(2m+d)}$. Notice that $\|\cdot\|$ and $\|\cdot\|_0$ are equivalent under the null hypothesis for any $\eta \in \mathcal{H}_0$. Thus, in what follows, we will not distinguish the two norms for notation convenience. We also do not distinguish h and h_0 since they have the same order for achieving optimality.

Next, we prove Lemma 3.4 as follows.

Proof. Let $g = \hat{\eta}_{n,\lambda} - \eta^*$. By Taylor's expansion we have

$$S_{n,\lambda}(\hat{\eta}_{n,\lambda}) = S_{n,\lambda}(\eta^*) + DS_{n,\lambda}(\eta^*)g + \int_0^1 \int_0^1 s D^2 S_{n,\lambda}(\eta^* + ss'g) g g ds ds'.$$

By (A.16) and (6), one can check that $\langle DS_{n,\lambda}(\eta^*)g_1, g_2 \rangle = \langle g_1, g_2 \rangle$, and thus, $DS_{n,\lambda} = id$ is an identity operator. By the fact $S_{n,\lambda}(\hat{\eta}_{n,\lambda}) = 0$, we have

$$\|\hat{\eta}_{n,\lambda} - \eta^* - S_{n,\lambda}(\eta^*)\| = \left\| \int_0^1 \int_0^1 s D^2 S_{n,\lambda}(\eta^* + ss'g) g g ds ds' \right\|. \quad (\text{A.9})$$

By (A.17) we have $D^2 S_{n,\lambda}(\eta^* + ss'g) g g = \int_{\mathcal{Y}} g(\mathbf{y})^2 \tilde{K}_{\mathbf{y}} e^{\eta^*(\mathbf{y}) + ss'g(\mathbf{y})} d\mathbf{y}$. By Proposition A.1 and Lemma A.3, we have

$$\sup_{\mathbf{y} \in \mathcal{Y}} |g(\mathbf{y})|^2 \leq c_m h^{-1} \|g\|^2 = c_m h^{-1} O_P((nh)^{-1} + h^{2m}),$$

where h^{-1} is defined in (A.8). By (B.1), we have $\|\mathbb{E}_{\eta^*}\{\tilde{K}_{\mathbf{Y}}\}\| \leq c_m^{1/2} h^{-1/2}$. Thus, we have

$$\|D^2 S_{n,\lambda}(\eta^* + ss'g) g g\| = O(h^{-3/2}((nh)^{-1} + h^{2m})). \quad (\text{A.10})$$

Plugging (A.10) into (A.9), we finish the proof. $\square$

A.3 Proof of Theorem 3.5 and Theorem 3.6

A.3.1 Proof of Theorem 3.5

The proof of Theorem 3.5 is sketched as follows. By Lemma 3.4, $n^{1/2}\|\hat{\eta}_{n,\lambda}^0 - \hat{\eta}_{n,\lambda} - S_{n,\lambda}^0(\eta^*) + S_{n,\lambda}(\eta^*)\| = o_P(1)$. So we have the following

$$n^{1/2}\|\hat{\eta}_{n,\lambda} - \hat{\eta}_{n,\lambda}^0\| = n^{1/2}\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\| + o_P(1).$$

Thus we only focus on $n^{1/2}\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\|$. Moreover, the following expressions of $S_{n,\lambda}^0(\eta^*)$ and $S_{n,\lambda}(\eta^*)$ are reserved for future use:

$$S_{n,\lambda}(\eta^*) = -\frac{1}{n} \sum_{i=1}^n \tilde{K}_{\mathbf{Y}_i} + \mathbb{E}_{\eta^*} \tilde{K}_{\mathbf{Y}} + W_{\lambda} \eta^*, \quad (\text{A.11})$$

$$S_{n,\lambda}^0(\eta^*) = -\frac{1}{n} \sum_{i=1}^n \tilde{K}_{\mathbf{Y}_i}^0 + \mathbb{E}_{\eta^*} \tilde{K}_{\mathbf{Y}}^0 + W_{\lambda}^0 \eta^*. \quad (\text{A.12})$$

Proof of Theorem 3.5. Let us first analyze I_1 . Let $\tilde{g} = \hat{\eta}_{n,\lambda} + ss'g - \eta^*$, for any $0 \leq s, s' \leq 1$. By Lemma S.7, we have $\|\tilde{g}\| = O_P((nh)^{-1/2} + h^{m/d}) = o_P(1)$. Notice that

$$D^2 \ell_{n,\lambda}(\hat{\eta}_{n,\lambda} + ss'g)gg = D^2 \ell_{n,\lambda}(\tilde{g} + \eta^*)gg = \int_{\mathcal{Y}} g^2(\mathbf{y}) e^{\tilde{g}(\mathbf{y}) + \eta^*(\mathbf{y})} d\mathbf{y} + \lambda J(g, g), \quad (\text{A.13})$$

and

$$D^2 \ell_{n,\lambda}(\eta^*)gg = \int_{\mathcal{Y}} g^2(\mathbf{y}) e^{\eta^*(\mathbf{y})} d\mathbf{y} + \lambda J(g, g). \quad (\text{A.14})$$

Combining (A.13) and (A.14), we have

$$|D^2 \ell_{n,\lambda}(\hat{\eta}_{n,\lambda} + ss'g)gg - D^2 \ell_{n,\lambda}(\eta^*)gg| \leq \int_{\mathcal{Y}} g^2(\mathbf{y}) e^{\eta^*(\mathbf{y})} |e^{\tilde{g}(\mathbf{y})} - 1| d\mathbf{y}.$$

By Taylor expansion of $e^{\tilde{g}(\mathbf{y}) + \eta^*(\mathbf{y})}$ at $\eta^*(\mathbf{y})$ for any $\mathbf{y} \in \mathcal{Y}$, it trivially holds that $e^{\eta^*(\mathbf{y})} |e^{\tilde{g}(\mathbf{y})} - 1| = e^{\eta^*(\mathbf{y})} O(|\tilde{g}(\mathbf{y})|)$. Since $\sup_{\mathbf{y} \in \mathcal{Y}} |\tilde{g}(\mathbf{y})| \leq c_m h^{-1/2} \|\tilde{g}\|$ (Lemma S.6), and $h^{-1/2}((nh)^{-1} + \lambda)^{1/2} = o(1)$, we have

$$|I_1| = O_P(h^{-1/2}(\|\hat{\eta}_{n,\lambda} - \eta^*\| + \|g\|) \cdot \|g\|^2) = o_P(\|g\|^2). \quad (\text{A.15})$$

Let us then analyze I_2 . From (3.9), we have $D^2 \ell_{n,\lambda}(\eta^*)gg = \|g\|^2 = \|\hat{\eta}_{n,\lambda} - \hat{\eta}_{n,\lambda}^0\|^2$, which dominates I_1 , since $h^{-1/2}(\|\hat{\eta}_{n,\lambda} - \eta^*\| + \|g\|) = o_P(1)$. Next let us analyze $\|\hat{\eta}_{n,\lambda} - \hat{\eta}_{n,\lambda}^0\|^2$. By Lemma 3.4, we have

$$n^{1/2}\|\hat{\eta}_{n,\lambda}^0 - \hat{\eta}_{n,\lambda} - S_{n,\lambda}^0(\eta^*) + S_{n,\lambda}(\eta^*)\| = O_P(n^{1/2}h^{-3/2}((nh)^{-1} + h^{2m/d})) = o_P(1).$$

Thus we only need to focus on $n^{1/2}\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\|$. Recall $S_{n,\lambda}(\hat{\eta}_{n,\lambda}) = 0$ and $S_{n,\lambda}(\eta^*)$, $S_{n,\lambda}^0(\eta^*)$ have expressions (A.11), (A.12). For any $\mathbf{y} \in \mathcal{Y}$, define $\tilde{K}_{\mathbf{y}}^1 = \tilde{K}_{\mathbf{y}} - \tilde{K}_{\mathbf{y}}^0$ and $W_{\lambda}^1 = W_{\lambda} - W_{\lambda}^0$, then $S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*) = -\frac{1}{n} \sum_{i=1}^n \tilde{K}_{\mathbf{Y}_i}^1 + \mathbb{E} \tilde{K}_{\mathbf{Y}}^1 + W_{\lambda}^1 \eta^*$.

By Proposition 3.3, $\tilde{K}_Y^1$ can be expressed as a series of $\xi_p^\perp(\mathbf{y})$. Since $\xi_p^\perp \in \mathcal{H}_1$ and $\phi_0 \equiv 1 \in \mathcal{H}_0$, we have

$$\mathbb{E}_{\eta^*}\{\xi_p^\perp(\mathbf{y})\} = \mathbb{E}_{\eta^*}\{\xi_p^\perp(\mathbf{y})\phi_0(X)\} = V(\xi_p^\perp, \phi_0) = 0.$$

And so $\mathbb{E}_{\eta^*}\{\tilde{K}_Y^1\} = 0$. Therefore, $S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*) = -\frac{1}{n} \sum_{i=1}^n \tilde{K}_{Y_i}^1 + W_\lambda^1 \eta^*$. Then

$$\begin{aligned} n\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\|^2 &= n^{-1} \left\| \sum_{i=1}^n \tilde{K}_{Y_i}^1 \right\|^2 - 2 \sum_{i=1}^n \langle \tilde{K}_{Y_i}^1, W_\lambda^1 \eta^* \rangle + n\|W_\lambda^1 \eta^*\|^2 \\ &\equiv W_1 - 2W_2 + W_3. \end{aligned}$$

Since $\eta^* \in \mathcal{H}_0$, it follows by Lemma 3.1 that η^* is expanded by a series of ξ_p^0 . By Proposition 3.3, $W_\lambda \xi_p^0 \propto \xi_p^0$ which implies $W_\lambda \eta^* = W_\lambda^0 \eta^*$. And hence, $W_\lambda^1 \eta^* = W_\lambda \eta^* - W_\lambda^0 \eta^* = 0$ which yields that $W_2 = W_3 = 0$. Write $W_1 = n^{-1} \left\| \sum_{i=1}^n \tilde{K}_{Y_i}^1 \right\|^2 = n^{-1} \sum_{i=1}^n \|\tilde{K}_{Y_i}^1\|^2 + n^{-1}W(n)$, where $W(n) = \sum_{i \neq j} \tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_j)$.

Next let us consider the term $\sum_{i=1}^n \tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_i)$. Let $\mathbb{E}$ denote $\mathbb{E}_{\eta^*}$ unless otherwise indicated. Let $\theta(n) = \mathbb{E}\{\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_i)\}$. By Lemma S.6 we have $\mathbb{E}\{|\sum_{i=1}^n \{\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_i) - \theta(n)\}|^2\} \leq n\mathbb{E}\{\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_i)^2\} = O(nh^{-2})$, so

$$\sum_{i=1}^n [\tilde{K}(\mathbf{Y}_i, \mathbf{Y}_i) - \theta(n)] = O_p(n^{1/2}h^{-1}). \quad (\text{A.16})$$

Next, we derive the asymptotic distribution of $W(n)$. Define $W_{ij} = 2\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_j)$, then $W(n) = \sum_{1 \leq i < j \leq n} W_{ij}$. Let $\sigma(n)^2 = \text{Var}(W(n))$ and

$$G_I = \sum_{i < j} \mathbb{E}\{W_{ij}^4\},$$

$$G_{II} = \sum_{i < j < k} (\mathbb{E}\{W_{ij}^2 W_{ik}^2\} + \mathbb{E}\{W_{ji}^2 W_{jk}^2\} + \mathbb{E}\{W_{ki}^2 W_{kj}^2\}), \quad \text{and}$$

$$G_{IV} = \sum_{i < j < k < l} (\mathbb{E}\{W_{ij} W_{ik} W_{lj} W_{lk}\} + \mathbb{E}\{W_{ij} W_{il} W_{kj} W_{kl}\} + \mathbb{E}\{W_{ik} W_{il} W_{jk} W_{jl}\}).$$

By $\mathbb{E}\{\tilde{K}_Y^1\} = 0$ and direct examinations we have

$$\begin{aligned} \sigma^2(n) = \text{Var}(W(n)) &= \sum_{1 \leq i < j \leq n} \mathbb{E}\{(\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_j) - \mathbb{E}[\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_j)])^2\} \\ &= \sum_{1 \leq i < j \leq n} \mathbb{E}\{\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_j)^2\} \asymp n^2 h^{-1}. \end{aligned}$$

Since $\mathbb{E}\{W_{ij}^4\} = 16\mathbb{E}\{\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_j)^4\} = O(h^{-4})$, we have $G_I = O(n^2h^{-4})$. Obviously, $\mathbb{E}\{W_{ij}^2W_{ik}^2\} \leq \mathbb{E}\{W_{ij}^4\} = O(h^{-4})$, implying $G_{II} = O(n^3h^{-4})$. For pairwise different i, j, k, l , we have

$$\begin{aligned}\mathbb{E}\{W_{ij}W_{ik}W_{lj}W_{lk}\} &= 16\mathbb{E}\{\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_j)\tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_k)\tilde{K}^1(\mathbf{Y}_l, \mathbf{Y}_j)\tilde{K}^1(\mathbf{Y}_l, \mathbf{Y}_k)\} \\ &= \sum_{p=1}^{\infty} \frac{1}{(1 + \lambda\rho_p^\perp)^4} = O(h^{-1}),\end{aligned}$$

which leads to $G_{IV} = O(n^4h^{-1})$.

It follows by $h = o(1)$ and $(nh^2)^{-1} = o(1)$ that G_I , G_{II} and G_{IV} are of lower order than $\sigma(n)^4$. By Proposition 3.2 of [de Jong \(1987\)](#) we get that

$$\frac{W(n)}{\sigma(n)} \xrightarrow{d} N(0, 1). \quad (\text{A.17})$$

From (A.16) and (A.17), we get $\frac{1}{n} \sum_{i=1}^n \tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}_i)^2 = \theta(n) + o_P(1)$, which implies $n\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\|^2 = O_P(h^{-1} + n\lambda + h^{-1/2}) = O_P(h^{-1})$, and hence $n^{1/2}\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\| = O_P(h^{-1/2})$. Thus,

$$\begin{aligned}2n \cdot PLR_{n,\lambda} &= n\|\hat{\eta}_{n,\lambda} - \eta^*\|^2 + o_P(h^{-1/2}) \\ &= (n^{1/2}\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\| + o_P(1))^2 + o_P(h^{-1/2}) \\ &= n\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\|^2 + 2n^{1/2}\|S_{n,\lambda}^0(\eta^*) - S_{n,\lambda}(\eta^*)\| \cdot o_P(1) + o_P(h^{-1/2}) \\ &= n^{-1} \left\| \sum_{i=1}^n \tilde{K}_{\mathbf{Y}_i}^1 \right\|^2 + o_P(h^{-1/2}).\end{aligned} \quad (\text{A.18})$$

By (A.17), (A.18) and Slutsky's theorem, $\frac{2n \cdot PLR_{n,\lambda} - \theta(n)}{\sigma(n)/n} \xrightarrow{d} N(0, 1)$. Since $\theta_\lambda = \sum_{p=1}^{\infty} \frac{1}{1 + \lambda\rho_p^\perp}$, $\sigma_\lambda^2 = \sum_{p=1}^{\infty} \frac{1}{(1 + \lambda\rho_p^\perp)^2}$, we have $\theta(n) = \theta_\lambda$ and $\frac{\sigma(n)}{n} = \sqrt{\binom{n}{2} \mathbb{E}(W_{ij}^2)}/n = \sqrt{2}\sigma_\lambda$. $\square$

A.3.2 Proof of Theorem 3.6

Before proving Theorem 3.6, we provide some preliminary lemmas. For $\eta^* \in \mathcal{H}$, consider decomposition $\eta^* = \eta_0^* + \eta_{XZ}^*$ where η_0^* is the projection of η^* on $\mathcal{H}_0$. The following lemma says that, for general $\eta^* \in \mathcal{H}$, the restricted penalized likelihood estimator $\hat{\eta}_{n,\lambda}^0$ converges to η_0^* with rate of convergence provided.

Lemma S.8. *Suppose that Assumption 1 is satisfied. We have $\|\hat{\eta}_{n,\lambda}^0 - \eta_0^*\|_0 = O_P((nh)^{-1/2} + \lambda^{1/2})$.*

Parallel to Lemma 3.4, when $\eta^* \in \mathcal{H}$, we have the following result characterizing the higher order expansion of $\hat{\eta}_{n,\lambda}^0$.

Lemma S.9. *Suppose that $nh^2 \rightarrow \infty$. We have*

$$\|\hat{\eta}_{n,\lambda}^0 - \eta_0^* - S_{n,\lambda}^0(\eta_0^*)\|_0 = O_P(h^{-3/2}((nh)^{-1} + h^{2m/d})).$$

Proof of Theorem 3.6. Let $g = \hat{\eta}_{n,\lambda}^0 - \hat{\eta}_{n,\lambda}$. Recall the Taylor expansion (3.11):

$$\begin{aligned} PLR_{n,\lambda} &= \ell_{n,\lambda}(\hat{\eta}_{n,\lambda}^0) - \ell_{n,\lambda}(\hat{\eta}_{n,\lambda}) \\ &= \int_0^1 \int_0^1 s \{D^2 f(\hat{\eta}_{n,\lambda} + ss'g)gg - D^2 f(\eta^*)gg\} ds ds' + \frac{1}{2} D^2 f(\eta^*)gg \\ &= O_P((\|\hat{\eta}_{n,\lambda} - \eta^*\|_{\sup} + \|g\|_{\sup}) \cdot \|g\|^2) + \frac{1}{2} \|g\|^2, \end{aligned}$$

where the O_P term in the last equation follows from (A.15). By Lemmas S.6 and S.7, $\|\hat{\eta}_{n,\lambda} - \eta^*\|_{\sup} = o_P(1)$. By assumption $\|\eta_{XZ}^*\|_{\sup} \leq (\log n)^{-1} = o(1)$ and Lemma S.8, we have $\|g\|_{\sup} = \|\hat{\eta}_{n,\lambda}^0 - \eta_0^* + \eta^* - \hat{\eta}_{n,\lambda} - \eta_{XZ}^*\|_{\sup} = o_P(1)$. Hence, the O_P term in (A.19) is dominated by $\frac{1}{2}\|g\|^2$, for which we only focus on the latter. Combining the results of Lemmas 3.4 and S.9, we have

$$\begin{aligned} \|\hat{\eta}_{n,\lambda} - \eta^* - S_{n,\lambda}(\eta^*)\| &= O_P(h^{-2}((nh)^{-1} + h^{2m/d})), \\ \|\hat{\eta}_{n,\lambda}^0 - \eta_0^* - S_{n,\lambda}^0(\eta_0^*)\|_0 &= O_P(h^{-2}((nh)^{-1} + h^{2m/d})). \end{aligned}$$

Recalling $\eta^* - \eta_0^* = \eta_{XZ}^*$, we have $\|g\| = \|\eta_{XZ}^* + S_{n,\lambda}(\eta^*) - S_{n,\lambda}^0(\eta_0^*)\| + O_P(h^{-2}((nh)^{-1} + h^{2m/d}))$. In what follows, we focus on $\|\eta_{XZ}^* + S_{n,\lambda}(\eta^*) - S_{n,\lambda}^0(\eta_0^*)\|$. By definition of $S_{n,\lambda}(\eta^*)$, $S_{n,\lambda}^0(\eta_0^*)$ (see (3.8)) and direct calculations, it can be shown that

$$\begin{aligned} &\|\eta_{XZ}^* + S_{n,\lambda}(\eta^*) - S_{n,\lambda}^0(\eta_0^*)\|^2 \\ &= \left\| \frac{1}{n} \sum_{i=1}^n \tilde{K}_{\mathbf{Y}_i}^1 \right\|^2 + \|\eta_{XZ}^*\|^2 + \|\mathbb{E} \tilde{K}_{\mathbf{Y}} - \mathbb{E} \tilde{K}_{\mathbf{Y}}^0\|^2 + \|W_{\lambda}^1 \eta_{XZ}^*\|^2 \\ &\quad - \frac{2}{n} \sum_{i=1}^n \eta_{XZ}^*(\mathbf{Y}_i) + 2\mathbb{E} \eta_{XZ}^*(\mathbf{Y}) + 2\langle W_{\lambda}^1 \eta_{XZ}^*, \eta_{XZ}^* \rangle - \frac{2}{n} \sum_{i=1}^n \mathbb{E} \tilde{K}^1(\mathbf{Y}_i, \mathbf{Y}) \\ &\quad - \frac{2}{n} (W_{\lambda}^1 \eta_{XZ}^*)(\mathbf{Y}_i) + 2\mathbb{E} (W_{\lambda}^1 \eta_{XZ}^*)(\mathbf{Y}), \end{aligned}$$

where $\mathbb{E}$ denotes $\mathbb{E}_{\eta^*}$. Since $\mathbb{E}\{\tilde{K}_{\mathbf{Y}} - \tilde{K}_{\mathbf{Y}}^0\} = \mathbb{E}\tilde{K}_{\mathbf{Y}}^1 = 0$, we have

$$\begin{aligned} & \|\eta_{XZ}^* + S_{n,\lambda}(\eta^*) - S_{n,\lambda}^0(\eta_0^*)\|^2 \\ & \geq \left\| \frac{1}{n} \sum_{i=1}^n \tilde{K}_{\mathbf{Y}_i}^1 \right\|^2 + \|\eta_{XZ}^*\|^2 + \left[-\frac{2}{n} \sum_{i=1}^n \eta_{XZ}^*(\mathbf{Y}_i) + 2\mathbb{E}_{\eta^*} \eta_{XZ}^*(\mathbf{Y}) \right] + 2\langle W_{\lambda}^1 \eta_{XZ}^*, \eta_{XZ}^* \rangle \\ & \quad + \left[-\sum_{i=1}^n \frac{2}{n} (W_{\lambda}^1 \eta_{XZ}^*)(\mathbf{Y}_i) + 2\mathbb{E}_{\eta^*} (W_{\lambda}^1 \eta_{XZ}^*)(\mathbf{Y}) \right] \equiv V_1 + V_2 + V_3 + V_4 + V_5. \end{aligned}$$

$$\text{Since } \text{Var}(V_3) \leq \frac{4}{n} \mathbb{E}(\eta_{XZ}^*(\mathbf{Y}))^2 \leq \frac{4}{n} \|\eta_{XZ}^*\|^2,$$

$$V_3 = O_P(n^{-1/2}) \|\eta_{XZ}^*\|. \quad (\text{A.19})$$

By assumption $J(\eta_{XZ}^*, \eta_{XZ}^*) \leq C$, we have

$$V_4 = \lambda J(\eta_{XZ}^*, \eta_{XZ}^*) \leq C\lambda. \quad (\text{A.20})$$

Since $\text{Var}(V_5) \leq \mathbb{E}|(W_{\lambda} \eta_{XZ}^*)|^2 = V(W_{\lambda} \eta_{XZ}^*, W_{\lambda} \eta_{XZ}^*)$. By Proposition 3.3, we have

$$V(W_{\lambda} \eta_{XZ}^*, W_{\lambda} \eta_{XZ}^*) = \sum_{p=1}^{\infty} |V(\eta_{XZ}^*, \xi_p)|^2 \left(\frac{\lambda \rho_p}{1 + \lambda \rho_p} \right)^2 = o(\lambda),$$

where the last equality follows by $\sum_{p=1}^{\infty} |V(\eta_{XZ}^*, \xi_p)|^2 \rho_p < \infty$ and the dominated convergence theorem. Thus we have

$$V_5 = o_P(n^{-1/2} \lambda^{1/2}) \quad (\text{A.21})$$

Combining (A.19), (A.20) and (A.21) we have

$$\begin{aligned} & \frac{2n \cdot PLR_{n,\lambda} - \theta(n)}{\sigma(n)} \\ & \geq \frac{2n \cdot V_1 - \theta(n)}{\sigma(n)} + \frac{2n \cdot (V_2 + V_3 + V_4 + V_5)}{\sigma(n)} \\ & \geq O_P(1) + 2n\sigma^{-1}(n)(\|\eta_{XZ}^*\|^2 + O_P(n^{-1/2} \|\eta_{XZ}^*\|) + O(\lambda) + o_P(n^{-1/2} \lambda^{1/2})). \end{aligned}$$

For $C_{\varepsilon} > 0$ sufficiently large, let η_{XZ}^* satisfy $\|\eta_{XZ}^*\|^2 \geq C_{\varepsilon} n^{-1/2} \|\eta_{XZ}^*\|$, $\|\eta_{XZ}^*\|^2 \geq C_{\varepsilon} \lambda$, $nh^{1/2} \|\eta_{XZ}^*\|^2 \geq C_{\varepsilon}$, $n \|\eta_{XZ}^*\|^2 / \sigma(n) \geq C_{\varepsilon}$, which implies that with probability greater than $1 - \varepsilon$, $|\frac{2n \cdot PLR_{n,\lambda} - \theta(n)}{\sigma(n)}| \geq c_{\alpha}$ (i.e., $\Phi_{n,\lambda}(\alpha) = 1$), where c_{α} is the $1 - \alpha$ percentile of standard normal distribution. It can be seen that the above conditions on η_{XZ}^* are satisfied if $\|\eta_{XZ}^*\|^2 \geq C_{\varepsilon}(\lambda + (nh^{1/2})^{-1})$. The result follows immediately by the fact $\|\eta_{XZ}^*\|_2 \leq \|\eta_{XZ}^*\|$. Proof is completed. $\square$

A.4 Proofs of the Minimax Lower Bound in Section 4

A.4.1 Preliminaries for the minimax lower bound

Lemma S.10. *Let $\mathbb{P}_0$ be the probability measure under the null, and $\mathbb{P}_1$ be the probability with density in $\{\eta \mid \|\eta_{XZ}\|_{\mathcal{H}} < d_n\}$. We have*

$$\inf_{\phi_n} \text{Err}(\phi_n, d_n) \geq 1 - \delta(\sqrt{\delta + 4} - \delta),$$

where $\delta^2 = \mathbb{E}_{\mathbb{P}_0}(d\mathbb{P}_1/d\mathbb{P}_0 - 1)^2$.

Proof. The test is bounded below by $1 - \|\mathbb{P}_0 - \mathbb{P}_1\|_{TV}$, where $\|\cdot\|_{TV}$ is the total variation distance between $\mathbb{P}_0$ and $\mathbb{P}_1$. By the theorem in Ingster (1987), we have

$$\frac{1}{2}\|\mathbb{P}_0 - \mathbb{P}_1\|_{TV} \leq \delta(1 - \frac{1}{2}\|\mathbb{P}_0 - \mathbb{P}_1\|_{TV})^{1/2},$$

which directly implies the result. $\square$

A.4.2 Proof of Lemma 4.1

Proof. As show in Lemma S.10, we have

$$\inf_{\phi_n} \text{Err}(\phi_n, d_n) \geq 1 - \delta(\sqrt{\delta + 4} - \delta). \quad (\text{A.22})$$

Next we show that if $d_n^2 \leq \frac{\sqrt{k_B(d_n)}}{4n}$, we have that the last term in (A.22) is larger than 1/2. For simplicity, denote $k = k_B(d_n)$. For any $b = (b_1, \dots, b_k) \in \{-1, 1\}^k$, let $\theta_b = \frac{d_n}{\sqrt{k}} \sum_{i=1}^k b_i \mathbf{e}_i \in \mathbb{R}^N$, where $\mathbf{e}_i$ is the standard basis vector with i th coordinate as one. We assume b is uniformly distributed over $\{-1, 1\}^k$ so that θ_b is uniformly distributed over $\mathbb{Q} := \{\theta_b : b \in \{-1, 1\}^k\}$. Since $\mathbb{E}_{\mathbb{P}_0} e^{\eta_{XZ}^{\theta_b}} - 1 = 0$, we have $e^{\eta_{XZ}^{\theta_b}} - 1 \in \mathcal{H}_{11}$. Define

$$\exp(\eta_{XZ}^{\theta_b}) - 1 = \frac{d_n}{\sqrt{k}} \sum_{l=1}^k b_l \psi_l \phi_1, \quad (\text{A.23})$$

where $\{\psi_l \phi_1\}_{l=1}^k$ are basis function for $\mathcal{H}_{11}$. We denote $\mathbb{P}_1^{(n)}$ and $\mathbb{P}_0^{(n)}$ as the empirical measures under the alternative and null respectively. The ratio of densities of $\mathbb{P}_1^{(n)}$ and $\mathbb{P}_0^{(n)}$ is

$$\frac{d\mathbb{P}_1^{(n)}}{d\mathbb{P}_0^{(n)}} = \mathbb{E}_{\theta_b} \prod_{i=1}^n \exp(\eta_{XZ}^{\theta_b}(\mathbf{Y}_i)).$$

Then, we denote the empirical version of δ as δ_n which can be written as

$$\begin{aligned}
\delta_n^2 &= \mathbb{E}_{\mathbb{P}_0^{(n)}}(d\mathbb{P}_1^{(n)}/d\mathbb{P}_0^{(n)} - 1)^2 \\
&= \mathbb{E}_{\mathbb{P}_0^{(n)}}[\mathbb{E}_{\boldsymbol{\theta}_b} \prod_{i=1}^n \exp(\eta_{XZ}^{\boldsymbol{\theta}_b}(\mathbf{Y}_i))]^2 - 1 \\
&= \mathbb{E}_{\mathbb{P}_0^{(n)}}[\mathbb{E}_{\boldsymbol{\theta}_b} \prod_{i=1}^n \exp(\eta_{XZ}^{\boldsymbol{\theta}_b}(\mathbf{Y}_i))][\mathbb{E}_{\boldsymbol{\theta}_{b'}} \prod_{i=1}^n \exp(\eta_{XZ}^{\boldsymbol{\theta}_{b'}}(\mathbf{Y}_i))] - 1 \\
&= \mathbb{E}_{\boldsymbol{\theta}_b, \boldsymbol{\theta}_{b'}} \prod_{i=1}^n \mathbb{E}_{\mathbb{P}_0} \exp(\eta_{XZ}^{\boldsymbol{\theta}_b}(\mathbf{Y}_i)) \exp(\eta_{XZ}^{\boldsymbol{\theta}_{b'}}(\mathbf{Y}_i)) - 1 \\
&= \mathbb{E}_{\boldsymbol{\theta}_b, \boldsymbol{\theta}_{b'}} [\mathbb{E}_{\mathbb{P}_0} \exp(\eta_{XZ}^{\boldsymbol{\theta}_b}(\mathbf{Y})) \exp(\eta_{XZ}^{\boldsymbol{\theta}_{b'}}(\mathbf{Y}))]^n - 1.
\end{aligned}$$

Plugging (A.23) in, we have

$$\begin{aligned}
\delta_n + 1 &= \mathbb{E}_{\boldsymbol{\theta}_b} \mathbb{E}_{\boldsymbol{\theta}_{b'}} [\mathbb{E}_{\mathbb{P}_0} (1 + \frac{d_n}{\sqrt{k}} \sum_{l=1}^k b_l \psi_l \phi_1) (1 + \frac{d_n}{\sqrt{k}} \sum_{l=1}^k b'_l \psi_l \phi_1)]^n \\
&= \frac{1}{2^k} \sum_{b, b'} (1 + \frac{d_n^2}{k} b^T b')^n \\
&\leq \frac{1}{2^k} \sum_b \exp\{\frac{n d_n^2 b^T 1_k}{k}\} \\
&= \frac{1}{2^k} \sum_{i=0}^k \binom{k}{i} \exp\{\frac{n(k-2i)d_n^2}{k}\} \\
&= \frac{1}{2^k} (\exp\{\frac{n d_n^2}{k}\} + \exp\{-\frac{n d_n^2}{k}\})^k \\
&\stackrel{(i)}{\leq} (1 + \frac{n^2 d_n^4}{k^2})^k \\
&\stackrel{(ii)}{\leq} \exp\{\frac{n^2 d_n^4}{k}\},
\end{aligned}$$

where (i) is due to the fact that $\frac{1}{2}(\exp(x) + \exp(-x)) \leq 1 + x^2$ for $|x| \leq 1/2$ and (ii) is due to the fact $1 + x \leq e^x$. Thus for any $d_n^4 \leq \frac{k}{16n^2}$, we have

$$\inf_{\phi_n} \text{Err}(\phi_n, d_n) \geq 1 - \delta_n(\sqrt{\delta_n + 4} - \delta_n) \geq 1 - e^{1/16}(\sqrt{e^{1/16} + 4}) \geq 1/2.$$

For $d_n \lesssim k^{1/4}/\sqrt{n}$, we have

$$|\exp\{\eta_{XZ}^{\boldsymbol{\theta}_b}\} - 1| = \frac{d_n}{\sqrt{k}} \left| \sum_{l=1}^k b_l \psi_l \phi_1 \right| \lesssim \frac{k^{3/4}}{\sqrt{n}}.$$

Thus, there exists $c_1, c_2 > 0$ such that

$$c_1 |\eta_{XZ}^{\theta_b}(\mathbf{y})| < |\exp\{\eta_{XZ}^{\theta_b}\} - 1| < c_2 |\eta_{XZ}^{\theta_b}(\mathbf{y})|, \quad (\text{A.24})$$

which indicates that $\|\exp\{\eta_{XZ}^{\theta_b}\} - 1\|_2 \asymp \|\eta_{XZ}^{\theta_b}\|_2$. By the definition of $r_B(\delta^*)$, we have $\text{Err}(\phi_n, d_n) > 1/2$ for all $d_n \leq r_B(\delta^*)$. $\square$

A.4.3 Proof of Lemma 4.2

Proof. We show that $b_{k,2}(\mathcal{E}_{11})$ is bounded below by $\sqrt{\gamma_{k+1}}$. It is sufficient to show that $\mathcal{E}_{11}$ contains a l_2 ball centered at $\eta_{XZ} = 0$ with radius $\sqrt{\gamma_{k+1}}$. For any $v \in \mathcal{E}_{11}$ with $\|v\|_2 \leq \sqrt{\gamma_{k+1}}$, we have

$$b_{2,k} \leq \sum_{i=1}^{k+1} \frac{v_i^2}{\gamma_i} \stackrel{(ii)}{\leq} \frac{1}{\mu_{k+1}} \sum_{i=1}^{k+1} v_i^2$$

where the inequality (i) holds by set the $(k+1)$ -dimensional subspace spanned by the eigenvectors corresponding to the first $(k+1)$ largest eigenvalues; the inequality (ii) holds by the decreasing order of the eigenvalues, i.e., $\gamma_1 \geq \gamma_2 \geq \dots \gamma_{k+1}$.

Recall that the definition of the Bernstein lower critical dimension is $k_B(\delta) = \arg\max_k \{b_{k-1,2}^2(\mathcal{E}_{11}) \geq \delta^2\}$, we have

$$k_B(\delta) \geq \arg\max_k \{\sqrt{\gamma_k} \geq \delta\}.$$

$\square$

A.4.4 Proof of Theorem 4.3

Proof. By Lemme 4.1, we have

$$d_n \leq \sup\{\delta : k_B(\delta) \geq 16n^2\delta^4\}.$$

Then we plug in the lower bound of k_B in Lemma 4.2 and we have

$$d_n \leq \sup\{\delta : \arg\max_k \{\sqrt{\gamma_k} \geq \delta\} \geq 16n^2\delta^4\} \quad (\text{A.25})$$

The eigenvalues have polynomial decay rate i.e., $\gamma_k \asymp k^{-2m/d}$, and consequently, $\arg\max_k \{\sqrt{\gamma_k} \geq \delta\} \asymp \delta^{-d/m}$. Plugging this into (A.25), it is easy to see that the supremum on the right hand side has an order $n^{-\frac{2m}{4m+d}}$. Proof is thus completed. $\square$

B Proofs of the Auxillary Results

In this document, additional proofs of auxillary lemmas are included.

- Section B.1 includes the proof of Lemma S.5.
- Section B.2 includes the proof of Lemma S.6.
- Section B.3 includes the proof of Lemma S.7.
- Section B.4 includes the proof of Lemma S.8.
- Section B.5 includes the proof of Lemma S.9.

B.1 Proof of Lemma S.5

Since $\rho_p \asymp p^{2m/d}$, we have

$$h^{-1} = \sum_{p=0}^{\infty} \frac{1}{(1 + \lambda\rho_p)^2} \asymp \int_1^{\infty} \frac{1}{(1 + \lambda p^{2m/d})^2} = \int_{\lambda^{d/2m}}^{\infty} \frac{1}{(1 + x^{2m/d})^2} dx = O(\lambda^{-d/2m})$$

Thus we have $h \asymp \lambda^{d/2m}$. Similarly, $h_0 \asymp \lambda^{d/2m}$.

B.2 Proof of Lemma S.6

For any $\mathbf{y} \in \mathcal{Y}$ and $\eta \in \mathcal{H}$, we have $|\eta(\mathbf{y})| = |\langle \tilde{K}_{\mathbf{y}}, \eta \rangle| \leq \|\tilde{K}_{\mathbf{y}}\| \cdot \|\eta\|$. So it is sufficient to find the upper bound for $\|\tilde{K}_{\mathbf{y}}\|$. By Proposition A.1 and the boundedness of ξ_p 's, we have

$$\|\tilde{K}_{\mathbf{y}}\|^2 = \tilde{K}(\mathbf{y}, \mathbf{y}) = \sum_{p=1}^{\infty} \frac{|\xi_p(\mathbf{y})|^2}{1 + \lambda\rho_p} \leq c_m h^{-1} \quad (\text{B.1})$$

where $c_m > 0$ is a constant free of $\mathbf{y}$ and η .

B.3 Proof of Lemma S.7

The proof is rooted in Gu (2013). Consider the quadratic approximation of the integral $\int_{\mathcal{Y}} e^{\eta(\mathbf{y})} d\mathbf{y}$:

$$\int_{\mathcal{Y}} e^{\eta(\mathbf{y})} d\mathbf{y} \approx \int_{\mathcal{Y}} e^{\eta^*(\mathbf{y})} d\mathbf{y} + \int_{\mathcal{Y}} (\eta - \eta^*) e^{\eta^*(\mathbf{y})} d\mathbf{y} + \frac{1}{2} V(\eta - \eta^*, \eta - \eta^*). \quad (\text{B.2})$$

Dropping the terms that do not involve η , and plugging (B.2) into (4), $\ell_{n,\lambda}(\eta)$ has a quadratic approximation $q_{n,\lambda}(\eta)$:

$$q_{n,\lambda}(\eta) = -\frac{1}{n} \sum_{i=1}^n \eta(\mathbf{Y}_i) + \int_{\mathcal{Y}} \eta e^{\eta^*} d\mathbf{y} + \frac{1}{2} V(\eta - \eta^*, \eta - \eta^*) + \frac{1}{2} J(\eta, \eta). \quad (\text{B.3})$$

Consider the Fourier expansions of η and η^* :

$$\eta(x, z) = \sum_{k=1}^{\infty} \sum_{l=1}^a \beta_{kl} \phi_k(x) \psi_l(z), \quad \eta^*(x, z) = \sum_{k=1}^{\infty} \sum_{l=1}^a \beta_{kl}^* \phi_k(x) \psi_l(z).$$

Then, we have

$$\begin{aligned} q_{n,\lambda}(\eta) = & \sum_{k=1}^{\infty} \sum_{l=1}^a \left\{ -\beta_{kl} \left(\frac{1}{n} \sum_{i=1}^n \phi_k(x_i) \psi_l(z_i) - \mathbb{E}\{\phi_k(X) \psi_l(Z)\} \right) \right. \\ & \left. + \frac{1}{2} (\beta_{kl} - \beta_{kl}^*)^2 + \frac{\lambda}{2} \mu_k \nu_l \beta_{kl}^2 \right\}. \end{aligned} \quad (\text{B.4})$$

Write $\gamma_{kl} = n^{-1} \sum_{i=1}^n \phi_k(X_i) \psi_l(Z_i) - \mathbb{E}\{\phi_k(X) \psi_l(Z)\}$. Minimizing (B.4) with respect to β_{kl} 's, we get the optimizer:

$$\tilde{\beta}_{kl} = (\gamma_{kl} + \beta_{kl}^*) / (1 + \lambda \mu_k \nu_l), \quad k \geq 1, l = 1, \dots, a.$$

Then $\tilde{\eta} = \sum_{k=1}^{\infty} \sum_{l=1}^a \tilde{\beta}_{kl} \phi_k \psi_l$ becomes a linear approximation of $\hat{\eta}_{n,\lambda}$. By direct calculations we get that

$$V(\tilde{\eta} - \eta^*) = \sum_{k=1}^{\infty} \sum_{l=1}^a (\beta_{kl} - \beta_{kl}^*)^2, \quad \lambda J(\tilde{\eta} - \eta^*) = \sum_{i=1}^{\infty} \sum_{j=1}^a \lambda \mu_k \nu_l (\beta_{kl} - \beta_{kl}^*)^2.$$

Since $\mathbb{E}\gamma_{kl} = 0$ and $\mathbb{E}\gamma_{kl}^2 = 1/n$, we have

$$\begin{aligned}\mathbb{E}\{V(\tilde{\eta} - \eta^*)\} &= \sum_{i=1}^{\infty} \sum_{j=1}^a \frac{1}{(1 + \lambda\mu_k\nu_l)^2} + \lambda \sum_{i=1}^{\infty} \sum_{j=1}^a \frac{\lambda\mu_k\nu_l}{(1 + \lambda\mu_k\nu_l)^2} \mu_k\nu_l\beta_{kl}^*\beta_{kl}^* \\ \mathbb{E}\{\lambda J(\tilde{\eta} - \eta^*)\} &= \sum_{i=1}^{\infty} \sum_{j=1}^a \frac{1}{(1 + \lambda\mu_k\nu_l)^2} + \lambda \sum_{i=1}^{\infty} \sum_{j=1}^a \frac{(\lambda\mu_k\nu_l)^2}{(1 + \lambda\mu_k\nu_l)^2} \mu_k\nu_l\beta_{kl}^*\beta_{kl}^*\end{aligned}\tag{B.5}$$

By similar derivations in Lemma S.6, it can be verified that

$$\begin{aligned}\sum_{i=1}^{\infty} \sum_{j=1}^a \frac{1}{(1 + \lambda\mu_k\nu_l)^2} &= O(\lambda^{-1/2m}), \\ \sum_{i=1}^{\infty} \sum_{j=1}^a \frac{\lambda\mu_k\nu_l}{(1 + \lambda\mu_k\nu_l)^2} &= O(\lambda^{-1/2m}), \\ \sum_{i=1}^{\infty} \sum_{j=1}^a \frac{1}{(1 + \lambda\mu_k\nu_l)} &= O(\lambda^{-1/2m})\end{aligned}$$

Plugging into (B.5), we obtain that

$$\|\tilde{\eta} - \eta^*\|^2 = (V + \lambda J)(\tilde{\eta} - \eta^*) = O_p(n^{-1}\lambda^{-1/2m} + \lambda).\tag{B.6}$$

We now turn to the approximation error $\hat{\eta} - \tilde{\eta}$. We calculate the Fréchet derivative of the quadratic approximation in (B.3) as

$$Dq_{n,\lambda}(\eta)\Delta\eta = -\frac{1}{n} \sum_{i=1}^n \Delta\eta(\mathbf{Y}_i) + \int_{\mathcal{Y}} \Delta\eta e^{\eta^*} d\mathbf{y} + \lambda V(\eta - \eta^*, \Delta\eta) + \lambda J(\eta, \Delta\eta).\tag{B.7}$$

Since $Dq_{n,\lambda}(\tilde{\eta}) = 0$, setting $\Delta\eta = \hat{\eta}_{n,\lambda} - \tilde{\eta}$, (B.7) is equal to

$$-\frac{1}{n} \sum_{i=1}^n (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{Y}_i) + \int_{\mathcal{Y}} (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\eta^*(\mathbf{y})} d\mathbf{y} + V(\tilde{\eta} - \eta^*, \hat{\eta}_{n,\lambda} - \tilde{\eta}) + \lambda J(\tilde{\eta}, \hat{\eta}_{n,\lambda} - \tilde{\eta})\tag{B.8}$$

Since $D\ell_{n,\lambda}(\hat{\eta}_{n,\lambda}) = 0$, setting $\Delta\eta = \hat{\eta}_{n,\lambda} - \tilde{\eta}$ yields

$$D\ell_{n,\lambda}(\eta)\Delta\eta = -\frac{1}{n} \sum_{i=1}^n (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{Y}_i) + \int_{\mathcal{Y}} (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\hat{\eta}_{n,\lambda}(\mathbf{y})} d\mathbf{y} + \lambda J(\hat{\eta}_{n,\lambda}, \hat{\eta}_{n,\lambda} - \tilde{\eta}).$$

Combining (B.8) and (B.9), we have

$$\begin{aligned} & \int (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\hat{\eta}_{n,\lambda}(\mathbf{y})} d\mathbf{y} - \int_{\mathcal{Y}} (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\tilde{\eta}(\mathbf{y})} d\mathbf{y} + \lambda J(\hat{\eta}_{n,\lambda} - \tilde{\eta}) \\ &= V(\tilde{\eta} - \eta^*, \hat{\eta}_{n,\lambda} - \tilde{\eta}) + \int_{\mathcal{Y}} (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\eta^*(\mathbf{y})} d\mathbf{y} - \int_{\mathcal{Y}} (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\tilde{\eta}(\mathbf{y})} d\mathbf{y}. \end{aligned}$$

By Taylor expansion,

$$\int (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\tilde{\eta}(\mathbf{y})} d\mathbf{y} - \int_{\mathcal{Y}} (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\eta^*(\mathbf{y})} d\mathbf{y} = V(\hat{\eta}_{n,\lambda} - \tilde{\eta}, \tilde{\eta} - \eta^*)(1 + o_p(1)),$$

where the o_p term holds as $\lambda \rightarrow 0$ and $n\lambda^{1/2m} \rightarrow \infty$. Define

$$D(\alpha) = \int_{\mathcal{Y}} (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\hat{\eta}_{n,\lambda}(\mathbf{y}) + \alpha(\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y})} d\mathbf{y}.$$

It can be shown that $\dot{D}(\alpha) = V_{\tilde{\eta} + \alpha(\hat{\eta}_{n,\lambda} - \tilde{\eta})}(\hat{\eta}_{n,\lambda} - \tilde{\eta})$. By the mean value theorem,

$$\begin{aligned} & \int_{\mathcal{Y}} (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\hat{\eta}_{n,\lambda}(\mathbf{y})} d\mathbf{y} - \int_{\mathcal{Y}} (\hat{\eta}_{n,\lambda} - \tilde{\eta})(\mathbf{y}) e^{\tilde{\eta}(\mathbf{y})} d\mathbf{y} \\ &= D(1) - D(0) = \dot{D}(\alpha) = V_{\tilde{\eta} + \alpha(\hat{\eta}_{n,\lambda} - \tilde{\eta})}(\hat{\eta}_{n,\lambda} - \tilde{\eta}), \end{aligned}$$

for some $\alpha \in [0, 1]$. Then by Assumption 1, we have

$$c_1 V(\hat{\eta}_{n,\lambda} - \tilde{\eta}) + \lambda J(\hat{\eta}_{n,\lambda} - \tilde{\eta}) \leq o_p(V(\tilde{\eta} - \eta^*, \hat{\eta}_{n,\lambda} - \tilde{\eta})) = o_p(\{V(\hat{\eta}_{n,\lambda} - \tilde{\eta})V(\tilde{\eta} - \eta^*)\}^{1/2})$$

Combine with the estimation error (B.6), we have

$$\|\hat{\eta}_{n,\lambda} - \eta^*\|^2 = V(\hat{\eta}_{n,\lambda} - \eta^*) + \lambda J(\hat{\eta}_{n,\lambda} - \eta^*) = O_p(n^{-1}\lambda^{1/2m} + \lambda).$$

B.4 Proof of Lemma S.8

Suppose the η_0^* is the projection of η^* on $\mathcal{H}_0$. Define an index set $\mathcal{I}_0 = \{(k, l) | k = 1 \text{ or } l = 1\}$ corresponding to the basis, $\{\phi_k \psi_l | k = 1 \text{ or } l = 1\}$, of $\mathcal{H}_0$. When restricted to $\mathcal{H}_0$, the Fourier expansion of η^* is

$$\eta_0^*(x, z) = \sum_{(k, l) \in \mathcal{I}_0} \beta_{kl}^0 \phi_k(x) \psi_l(z).$$

Substituting the above η_0^* as well as its Fourier expansion into the proof of Lemma A.4, all results remain valid, provided the following truth:

$$\mathbb{E}\left\{\frac{1}{n}\sum_{i=1}^n\phi_k(X_i)\psi_l(Z_i)-\mathbb{E}_{\eta^*}(\phi_k\psi_l)\right\}^2=\frac{1}{n}$$

$$\mathbb{E}\left\{\frac{1}{n}\sum_{i=1}^n\phi_k(X_i)\psi_l(Z_i)\phi_{k'}(X_i)\psi_{l'}(Z_i)-\mathbb{E}_{\eta^*}(\phi_k\psi_l\phi_{k'}\psi_{l'})\right\}^2\leq\frac{c}{n},$$

where c is a positive constant. The existence of such c is guaranteed by the uniform boundedness of $\phi_k(x)$'s as proved by [Shang and Cheng \(2013\)](#). Let η_0^* be the projection of η^* on the subspace $\mathcal{H}_0$ and $g = \hat{\eta}_{n,\lambda}^0 - \eta_0^*$. Substituting η_0^* and $\hat{\eta}_{n,\lambda}^0$ into the proof of Lemma A.4, the results would follow.

B.5 Proof of Lemma S.9

Let η_0^* be the projection of η^* on the subspace $\mathcal{H}_0$ and $g = \hat{\eta}_{n,\lambda}^0 - \eta_0^*$. Substituting η_0^* and $\hat{\eta}_{n,\lambda}^0$ into the proof of Lemma 3.4, one can show the desired results.

C Additional Simulation with Beta and Beta Mixture

In this section, we consider the distribution with different shapes. Specifically, we considered the follow two settings:

Setting 5: the simple Beta distributions,

$$X \mid Z = z \sim \text{Beta}(2(1 + \delta_5 \mathbb{1}_{z=1}), 2(1 + \delta_5 \mathbb{1}_{z=1}))$$

where $\delta_5 = 0, 0.4, 0.6$.

Setting 6: Beta mixture distributions,

$$X \mid Z = z \sim 0.5\text{Beta}(2(1 + \delta_6 \mathbb{1}_{z=1}), 6(1 + \delta_6 \mathbb{1}_{z=1}))$$

$$+ 0.5\text{Beta}(6(1 + \delta_6 \mathbb{1}_{z=1}), 2(1 + \delta_6 \mathbb{1}_{z=1}))$$

where $\delta_6 = 0, 0.3, 0.45$. We calculated the size and power based 1000 independent trials.

Setting 5 corresponds to a Beta distribution while Setting 6 corresponds to a mixture of Beta distributions. With $\delta_5 = 0$ and $\delta_6 = 0$, we intended to examine the size of the test under the H_0 . The power of the testing methods were examined with positive δ_5 's and δ_6 's.




Figure 5: Size vs. sample size for KS, MMD, ELT, AD, DSLICE and PLR tests.

As shown in Figure 5(a), the empirical sizes of Setting 5 were all around 0.05 for the six test procedures when the density is a unimodal Beta distribution. Whereas, for setting 6, Figure 5(b) shows that the empirical sizes of KS, MMD, ELT, AD, and DSLICE tests were significantly lower than 0.05, while the sizes of PLR test were still around 0.05. This demonstrates that our PLR test is asymptotically correct for both unimodal and bimodal distributions.

Figure 6(a) and (b) examine the power of the three tests under Setting 5. In Setting 5, when $\delta_5 = 0.6$, the empirical powers of the MMD, AD and PLR test approached 1 as n increased. In contrast, the power of KS and ELT test were lower than 0.5 even when the averaged sample size in each group reaches 1000. DSLICE has power slightly over 0.5 when $\delta_5 = 0.6$ when $n = 1000$. In Setting 6, as shown in Figure 6(c)(d) the power of KS, MMD, and ELT test were below 0.2 even when the averaged sample size in each group is 1000. The power of AD and DSLICE is slightly over 0.5 when $n = 1000$ and $\delta_6 = 0.45$. In contrast, the power of PLR test approached 1 rapidly when δ_6 was 0.30 or 0.45. We conclude that the PLR test is still the most powerful among the four tests in all the considered settings,




Figure 6: Power vs. sample size for PLR, KS, MMD, ELT, AD, and DSLICE.

even when the data distribution is multimodal and non-Gaussian.