

# On algebras of $\Omega^n$ -finite and $\Omega^\infty$ -infinite representation type

Marcos Barrios · Gustavo Mata

the date of receipt and acceptance should be inserted later

**Abstract** In this article we characterize the subcategory  $\Omega^\infty(\text{mod}A)$  for algebras of  $\Omega^n$ -finite representation type. As a consequence, we characterize when a truncated path algebra is a Co-Gorenstein algebra in terms of its associated quiver. We also study the behaviour of Artin algebras of  $\Omega^\infty$ -infinite representation type. Finally, it is presented an example of a non Gorenstein algebra of  $\Omega^\infty$ -infinite representation type and an example of a finite dimensional algebra with infinite  $\phi$ -dimension.

**Keywords** Truncated path algebra, Co-Gorenstein algebra, Igusa-Todorov function.

**PACS** 16W50 · 16E30

**Mathematics Subject Classification (2010)** 16G10

## 1 Introduction

The Finitistic Dimension Conjecture (FDC), the Generalized Nakayama Conjecture (GNC) and the Nakayama Conjecture (NC) are examples of important conjectures in Representation Theory of Artin algebras and they still remain open. These conjectures are related in the following way:

If  $A$  is an Artin algebra,

- $A$  verifies FDC then  $A$  verifies GNC,
- $A$  verifies GNC then  $A$  verifies NC.

An algebra  $A$  is Co-Gorenstein if the subcategories  $\Omega^\infty(\text{mod}A)$  and  $\mathcal{G}\mathcal{P}(A)$  agree. In [11] the authors prove that there is a relation between NC and GNC and Co-Gorenstein algebras. This justifies the relevance of studying this family of algebras.

---

M. Barrios  
Universidad de la República, Montevideo, Uruguay  
E-mail: marcosb@fing.edu.uy G. Mata  
Universidad de la República, Montevideo, Uruguay  
E-mail: gmata@fing.edu.uy

In this article we characterize the subcategory  $\Omega^\infty(\text{mod}A)$  when  $A$  is of  $\Omega^n$ -finite representation type (Theorem 37). In particular, truncated path algebras are of  $\Omega^1$ -finite representation type (Theorem 5.11 of [3]). We use both theorems and determine, in terms of the geometry of the quiver, when a truncated path algebra is a Co-Gorenstein algebra.

In an attempt to prove FDC, K. Igusa and G. Todorov define, in [10], two functions from the objects of  $\text{mod}A$  to the natural numbers, which generalizes the notion of projective dimension. Nowadays, they are known as the Igusa-Todorov functions,  $\phi$  and  $\psi$ . In that article the authors prove, with the Igusa-Todorov functions, that if an Artin algebra  $A$  has  $\text{repdim}(A) \leq 3$  then  $A$  verifies the FDC.

More recently, it was conjectured in [7] that the associated dimensions to these functions are finite ( $\phi$ DC). This conjecture is also related with the FDC as follows: For  $A$  an Artin algebra, if  $A$  verifies  $\phi$ DC then  $A$  verifies FDC. In [13], the authors prove that every Artin algebra of  $\Omega^n$ -finite representation type and every Gorenstein algebra verifies  $\phi$ DC. Therefore, to find an example of an algebra that does not verify the  $\phi$ DC, it should be looked in the family of non Gorenstein algebras of  $\Omega^\infty$ -infinite representation type.

In this article we show that there are algebras of  $\Omega^\infty$ -infinite representation type that are not Gorenstein algebras (Examples 53 and 54). Finally we exhibit an example (Example 54) where  $\phi$ DC and FDC are not equivalent.

## 2 Preliminaries

From now on  $A$  will always be an Artin algebra. We denote by  $\text{mod}A$  the category of finitely generated right  $A$ -modules. The full subcategory of  $\text{mod}A$  whose objects are direct summands of finite sums of copies of  $M$  is denoted by  $\text{add}(M)$ . For a module  $M \in \text{mod}A$ ,  $\text{pd}(M)$  and  $\Omega(M)$  are the projective dimension and the syzygy of  $M$ , respectively. For an algebra  $A$ ,  $\text{gldim}(A)$  and  $\text{findim}(A)$  are the global and finitistic dimension of  $A$ , respectively.

We denote by  $\mathcal{P}(A)$  the full subcategory formed by projective modules and by  $\mathcal{GP}(A)$  the full subcategory formed by Gorenstein projective modules. Observe that  $\mathcal{P}(A) \subset \mathcal{GP}(A)$ . Recall that an algebra  $A$  is self-injective if and only if  $\mathcal{GP}(A) = \text{mod}A$ , that is, all modules are Gorenstein projective. On the other hand,  $A$  is called CM-free provided that  $\mathcal{GP}(A) = \mathcal{P}(A)$ , that is, all its finitely generated Gorenstein projective modules are projective. Recall that for an Artin algebra  $A$ , the set  $\text{ind } \mathcal{GP}(A)$  of isoclasses of indecomposable objects in the stable category  $\mathcal{GP}(A)$  is identified with the set of isoclasses of indecomposable non-projective Gorenstein-projective  $A$ -modules.

We denote by  $\mathcal{P}^{<\infty}(A)$  the full subcategory of  $\text{mod}A$  formed by the modules of finite projective dimension and by  $\mathcal{P}^\infty(A)$  the full subcategory of  $\text{mod}A$  formed by the modules of infinite projective dimension with no summand with finite projective dimension.

We denote by  $\text{add}(\Omega^n(\text{mod}A))$  the full subcategory of  $\text{mod}A$  formed by the direct sums of direct summands of modules in  $\Omega^n(\text{mod}A)$ . For  $n \in \mathbb{N}$ , we say that  $\Omega^n(\text{mod}A)$  is of **finite representation type** if there are only finitely many indecomposable modules  $M$ , up to isomorphism, such that  $M \in \text{add}(N)$  with  $N = \Omega^n(N')$ . In this case  $A$  is called of  **$\Omega^n$ -finite representation type**. If  $A$  is not of  $\Omega^n$ -finite representation type for any  $n \in \mathbb{N}$ , we say that  $A$  is  **$\Omega^\infty$ -infinite representation type**.

## 2.1 Quivers and path algebras

If  $Q = (Q_0, Q_1, s, t)$  is a finite connected quiver,  $\mathbb{k}Q$  denotes its associated path algebra. Given  $\rho$  a path in  $\mathbb{k}Q$ , we denote by  $l(\rho)$ ,  $s(\rho)$  and  $t(\rho)$  the length, start and target of  $\rho$ , respectively.  $C^n$  denotes the oriented cycle graph with  $n$  vertices. We denote by  $J$  the ideal generated by all the arrows from  $Q_1$ . Given a quiver  $Q$  with cycles, we denote by  $Q^\infty$  the full subquiver of  $Q$  formed by the vertices  $w$  with infinite paths starting at  $w$ . Let  $S$  be a set of paths in  $Q$ . A path  $p$  in  $S$  is **left-minimal** in  $S$  provided that there is no path  $q$  such that  $q \in S$  and  $p = qp'$  for some nontrivial path  $p'$ . Dually, one defines a **right-minimal** path in  $S$ . A full subquiver  $Q'$  of a quiver  $Q$  is called a **final subheart** (see[14]) if it is minimal in the family of full subquivers of  $Q$  which are closed by successors. We say a final subheart is **trivial** if  $Q'$  is a simple vertex.

**Remark 21** *If  $Q$  has a cycle and has not surces, then  $Q$  has a non trivial final subheart.*

Let  $A = \frac{\mathbb{k}Q}{I}$  be an algebra, if  $v \in Q_0$  we denote by  $S_v$  and  $I_v$  the simple  $A$ -module and the injective  $A$ -module associated to  $v$ , respectively. For  $M \in \text{mod } \frac{\mathbb{k}Q}{I}$ , we denote by  $\underline{\dim}(M)$  its dimension vector. A finite dimensional algebra  $A = \frac{\mathbb{k}Q}{I}$  is called a **monomial algebra** if  $I$  is generated by paths.

Let  $A = \frac{\mathbb{k}Q}{I}$  be a monomial algebra. For a nonzero nontrivial path  $p$ , we define  $L(p)$  to be the set of right-minimal paths in the set  $\{\text{nonzero paths } q \text{ such that } s(q) = t(p) \text{ and } qp = 0\}$  and  $R(p)$  the set of left-minimal paths in the set  $\{\text{nonzero paths } q \text{ such that } t(q) = s(p) \text{ and } pq = 0\}$ . We call a pair  $(p, q)$  of nonzero paths in  $A$  a **perfect pair** provided that the following conditions are satisfied:

- both of the nonzero paths  $p, q$  are nontrivial satisfying  $s(p) = t(q)$  and  $pq = 0$  in  $A$ ;
- if  $pq' = 0$  for a nonzero path  $q'$  with  $t(q') = s(p)$ , then  $q' = qq''$  for some path  $q''$ ; in other words,  $R(p) = \{q\}$ ;
- if  $p'q = 0$  for a nonzero path  $p'$  with  $s(p') = t(q)$ , then  $p' = p''p$  for some path  $p''$ ; in other words,  $L(q) = \{p\}$ .

Let  $A = \frac{\mathbb{k}Q}{I}$  be a monomial algebra. We call a nonzero path  $p$  in  $A$  a **perfect path**, provided that there exists a sequence:

$$p = p_1, p_2, \dots, p_n, p_{n+1} = p$$

of nonzero paths such that  $(p_i, p_{i+1})$  are perfect pairs for all  $1 \leq i \leq n$ . If the given nonzero paths  $p_i$  are pairwise distinct, we refer to the sequence  $p = p_1, p_2, \dots, p_n, p_{n+1} = p$  as a **relation-cycle** for  $p$ .

The next theorem (See Theorem 4.11 of [4]) shows how to compute every non projective Gorenstein projective module.

**Theorem 22** [4, Theorem 4.11] *Let  $A$  be a monomial algebra. Then there is a bijection 1 : 1*

$$\{\text{perfect paths in } A\} \rightarrow \text{ind } \underline{\mathcal{G}\mathcal{P}}(A)$$

*sending a perfect path  $p$  to the  $A$ -module  $Ap$ .*

The following theorem (see [15]) shows that monomial algebras are of  $\Omega^2$ -finite representation type algebras.

**Theorem 23** [15, Theorem 3.1] *Let  $A$  be a finite dimensional monomial algebra. If  $M$  is an  $A$ -module such that  $M \cong \Omega^t(N)$  for an  $A$ -module  $N$  and  $t \geq 2$ , then it is a direct sum of right ideals of  $A$  which are generated by paths with length greater or equal than 1. In particular, monomial algebras are of  $\Omega^2$ -finite representation type.*

## 2.2 Truncated path algebras

A monomial algebra  $A = \frac{\mathbb{k}Q}{J^k}$  is called a **truncated path algebra** if there is a  $k \in \mathbb{N}, k \geq 2$  such that  $I = J^k$ . If  $k = 2$  we say that  $A$  is a **radical square zero algebra**. Following [2], we use the below notation.

**Notation 24** For a truncated path algebra  $A$ , we denote by  $M_v^l(A)$  the ideal  $\rho A$ , where  $l(\rho) = l$ ,  $t(\rho) = v$  and  $M^l(A) = \bigoplus_{v \in Q_0} M_v^l(A)$ .

Given  $0 < l \leq k-1$ , it is easy to see that  $M_v^l(A)$  exists if and only if  $\text{id}_{\frac{\mathbb{k}Q}{J^2}} S_v \geq l$ . Also,  $M_v^l(A)$  is a projective  $A$ -module if and only if  $\text{pd}_{\frac{\mathbb{k}Q}{J^2}} S_v \leq k-l-1$ .

**Remark 25** If  $A = \frac{\mathbb{k}Q}{J^k}$  is a truncated path algebra, then

$$\Omega(M_v^l(A)) = \bigoplus_{\rho: \begin{cases} s(\rho)=v \\ l(\rho)=k-l \end{cases}} M_{t(\rho)}^{k-l}(A),$$

Proposition 2.6 gives a method to compute the global dimension for truncated path algebras as a function of the length of the longest path of their quivers. See Theorem 5 of [6] for a proof.

**Proposition 26** [6, Theorem 5]

Let  $A \cong \frac{\mathbb{k}Q}{J^k}$  be a truncated path algebra. Then the following statements hold.

1.  $\text{gldim}(A) < \infty$  if and only if  $Q$  has no oriented cycles.
2. If  $A$  has finite global dimension, then

$$\text{gldim}(A) = \begin{cases} 2\frac{l}{k} & \text{if } l \equiv 0 \pmod{k}, \\ 2\lfloor \frac{l}{k} \rfloor + 1 & \text{otherwise,} \end{cases}$$

where  $l$  is the length of the largest path of  $\mathbb{k}Q$ .

**Definition 27** Let  $A$  be an Artin algebra. We define  $\|\cdot\| : \text{mod}A \rightarrow \mathbb{N}$  a function such that  $\|M\| = k$  if  $M \cong (\bigoplus_{i=1}^k M_i) \oplus (\bigoplus_{j=1}^m N_j)$  is the decomposition into indecomposable modules where  $\text{pd}(M_i) = \infty \forall i \in \{1, \dots, k\}$  and  $\text{pd}(N_j) < \infty \forall j \in \{1, \dots, m\}$ .

**Remark 28** If  $M$  and  $N$  are  $A$ -modules with infinite projective dimension, we have that:

- $\|M \oplus N\| = \|M\| + \|N\|$ .
- $\|M^k\| = k\|M\|$  for  $k \in \mathbb{N}$ .
- $\|M\| \leq \|\Omega(M)\|$ .
- If  $\Omega^m(M) \cong M$  then  $\|\Omega(M)\| = \|M\|$ .

## 2.3 Igusa-Todorov functions

We recall the definition of the Igusa-Todorov  $\phi$  function and some basic properties. We also define the  $\phi$ -dimension of an algebra.

**Definition 29** Let  $K_0(A)$  be the abelian group generated by all symbols  $[M]$ , where  $M \in \text{mod}A$ , modulo the relations

1.  $[M] - [M'] - [M'']$  if  $M \cong M' \oplus M''$ ,
2.  $[P]$  for each projective.

Let  $\bar{\Omega} : K_0(A) \rightarrow K_0(A)$  be the group endomorphism induced by  $\Omega$ , and let  $K_i(A) = \bar{\Omega}(K_{i-1}(A)) = \dots = \bar{\Omega}^i(K_0(A))$ . If  $\mathcal{C}$  is subcategory of  $\text{mod}A$  closed by direct sums and direct summands such that  $\Omega(\mathcal{C}) \subset \mathcal{C}$ , then we define by  $K_0(\mathcal{C}) = \{[M] : M \in \mathcal{C}\}$ . Now, if  $M$  is a finitely generated  $A$ -module then  $\langle \text{add}M \rangle$  denotes the subgroup of  $K_0(A)$  generated by the classes of indecomposable summands of  $M$ .

**Definition 210** [10] *The (right) Igusa-Todorov function  $\phi$  of  $M \in \text{mod}A$  is defined as*

$$\phi_A(M) = \min \left\{ l : \bar{\Omega}_A|_{\bar{\Omega}_A^{l+s} \langle \text{add}M \rangle} \text{ is a monomorphism for all } s \in \mathbb{N} \right\}.$$

In case that there is no possible misinterpretation we denote by  $\phi$  the Igusa-Todorov function  $\phi_A$ .

**Proposition 211** ([10], [9]) *Given  $M, N \in \text{mod}A$ .*

1.  $\phi(M) = \text{pd}(M)$  if  $\text{pd}(M) < \infty$ .
2.  $\phi(M) = 0$  if  $M \in \text{ind}A$  and  $\text{pd}(M) = \infty$ .
3.  $\phi(M) \leq \phi(M \oplus N)$ .
4.  $\phi(M^k) = \phi(M)$  for  $k \in \mathbb{N}$ .
5. If  $M \in \text{mod}A$ , then  $\phi(M) \leq \phi(\Omega(M)) + 1$ .

*Proof* For the statements 1-4 see [10], and for 5 see [9].

**Definition 212** *Let  $A$  an Artin algebra, and  $\mathcal{C}$  a subcategory of  $\text{mod}A$ . The  $\phi$ -dimension of  $\mathcal{C}$  is defined as*

$$\phi \dim(\mathcal{C}) = \sup \{ \phi(M) : M \in \mathcal{C} \}.$$

*In particular, we denote by  $\phi \dim(A)$  the  $\phi$ -dimension of  $\text{mod}A$*

The following theorem (see [7]) shows that the finitude of the  $\phi$ -dimension is invariant by derived equivalences.

**Theorem 213** [7, Theorem 4.10] *Let  $A$  and  $B$  be Artin algebras, which are derived equivalent. Then,  $\phi \dim(A) < \infty$  if and only if  $\phi \dim(B) < \infty$ .*

### 3 Periodic modules over algebras of $\Omega^n$ -finite representation type

In this section we prove that the subcategory of  $\Omega$ -periodic  $A$ -modules and the subcategory  $\Omega^\infty(\text{mod}A)$  agree for an algebra  $A$  of  $\Omega^n$ -finite representation type.

**Definition 31** *For an algebra  $A$  we denote by  $\Omega^\infty(\text{mod}A)$  the full subcategory generated by the class of modules  $\{M \in \text{mod}A : 0 \rightarrow M \rightarrow P_0 \rightarrow P_1 \rightarrow \dots \text{ with } P_i \in \mathcal{P}(A) \forall i \in \mathbb{N}\}$ . We say that  $\Omega^\infty(\text{mod}A)$  is **trivial** if  $\mathcal{P}(A) = \Omega^\infty(\text{mod}A)$ .*

**Definition 32** *When an  $A$ -module  $M$  verifies  $\Omega^m(M) \cong M$  for  $m \in \mathbb{Z}^+$ , we say that  $M$  is  **$\Omega$ -periodic**.*

**Remark 33** For an algebra  $A$  we have the inclusions:

$$\mathcal{P}(A) \subset \mathcal{G}\mathcal{P}(A) \subset \Omega^\infty(\text{mod}A).$$

If, in addition, the algebra  $A$  has finite global dimension, then:

$$\mathcal{P}(A) = \mathcal{G}\mathcal{P}(A)$$

**Remark 34** If the algebra  $A$  has  $\Omega^\infty(\text{mod}A)$  trivial, then  $A$  is CM-free.

**Remark 35** If the algebra  $A$  has  $\text{findim}(A) < \infty$ , every non projective module  $M$  in  $\Omega^\infty(\text{mod}A)$  verifies  $\text{pd}(M) = \infty$ .

**Lemma 36** If  $A$  is of  $\Omega^n$ -finite representation type for  $n \in \mathbb{Z}^+$ , then the following assertions are equivalent:

- There is a nonprojective module  $M$  in  $\Omega^\infty(\text{mod}A)$ .
- There is an  $A$ -module  $N \neq \{0\}$  and  $n \in \mathbb{Z}^+$  such that  $\Omega^n(N) \cong N$ .

*Proof* Suppose there exist  $M \in \text{mod}A$  and  $n \in \mathbb{Z}^+$  such that  $\Omega^n(M) \cong M$ . Then there exist a exact sequence:

$$\zeta : 0 \rightarrow M \rightarrow P_n \rightarrow \dots \rightarrow P_1 \rightarrow P_0 \rightarrow M \rightarrow 0.$$

We can join infinite exact sequences  $\zeta$ , thus we conclude that  $M \in \Omega^\infty(\text{mod}A)$ .

Suppose that  $A$  is  $\Omega^L$ -finite representation type and that  $M = M_1 \oplus M_2 \in \Omega^\infty(\text{mod}A)$  with  $M_1 \in \mathcal{P}^\infty(A)$  no null and  $M_2 \in \mathcal{P}^{<\infty}(A)$ . Consider the set  $\mathcal{W}_M = \{N \in \mathcal{P}^\infty(A) \cap \text{add}(\Omega^L(\text{mod}A)) \text{ such that } \|N\| \leq \|M\|\}$ . Clearly  $\mathcal{W}_M$  is a finite non empty set ( $M_1 \in \mathcal{W}_M$ ), thus there exist  $N_0 \in \mathcal{W}_M$ ,  $f(k) = n_k : \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$  an increasing function, and  $N_{n_k} \in \mathcal{P}^{<\infty}(A)$  such that  $\Omega^{n_k}(N_0 \oplus N_{n_k}) \cong M$  for all  $k \in \mathbb{N}$ . We deduce that there are  $i, j \in \mathbb{Z}^+$  such that  $n_j - n_i > m = \text{findim}(A)$  and  $\Omega^{n_j - n_i}(N_0) \cong N_0 \oplus N'$  with  $N' \in \mathcal{P}^{<\infty}(A)$ . Finally we conclude that  $N_0 \oplus N'$  is a  $\Omega$ -periodic module.

In the proof of the proposition above we observe that  $M \cong \Omega^k(N)$  for  $N$  a periodic module and  $k > \text{findim}(A)$ . So we deduce the next result.

**Theorem 37** Let  $A$  be an syzygy finite Artin algebra modules. If  $M$  is a non projective module, then  $M \in \Omega^\infty(\text{mod}A)$  if and only if there exist  $m \in \mathbb{Z}^+$  such that  $\Omega^m(M) \cong M$ .

By Theorem 23, for a monomial algebra is valid:

**Corollary 38** If  $A$  is a monomial algebra, the following statements are equivalent:

- There is a nonprojective module  $M$  in  $\Omega^\infty(\text{mod}A)$ .
- There are an  $A$ -module  $N \neq \{0\}$  and  $n \in \mathbb{Z}^+$  such that  $\Omega^n(N) \cong N$ .

We deduce the following corollary from the proof of Lemma 36.

**Corollary 39** Let  $A$  be an Artin algebra, if  $M$  is  $\Omega$ -periodic  $A$ -module then  $M \in \Omega^\infty(\text{mod}A)$ .

The following example shows that Theorem 37 is not valid if the algebra is not syzygy finite.



The following result characterizes when a truncated path algebra is a Co-Gorenstein algebra.

**Theorem 44** *If  $A = \frac{\mathbb{k}Q}{jk}$  is a truncated path algebra, then are equivalent:*

1.  $A$  is Co-Gorenstein
2.  $Q$  is one of the following types:
  - i.  $Q$  is acyclic.
  - ii.  $Q = C^n$ .
  - iii.  $Q^\infty$  has no final subheart equal to  $C^i$  for  $i \in \mathbb{Z}^+$ .

*Proof* Any finite quiver  $Q$  belongs to a only one of the following families:


- i.  $Q$  acyclic
- ii.  $Q = C^n$
- iii.  $Q$  has cycles,  $Q \neq C^n$  and every final subheart of  $Q^\infty$  is not equal to  $C^i$  for  $i \in \mathbb{Z}^+$ .
- iv.  $Q$  has cycles,  $Q \neq C^n$  and there exist a final subheart of  $Q^\infty$  equal to  $C^i$  for  $i \in \mathbb{Z}^+$ .

claim: if  $Q$  is in the first three families, then  $A$  is Co-Gorenstein.

- i. If  $Q$  is acyclic, then  $\text{gldim} A < \infty$ . Thus  $A$  is a Gorenstein algebra.
- ii. If  $Q = C^n$ , then  $A$  is a selfinjective algebra. Thus  $A$  is 0-Gorenstein.
- iii. Suppose that  $Q$  has cycles, then  $Q^\infty$  has cycles and its final subhearts are not trivial. Suppose that every final subheart of  $Q^\infty$  is not equal to  $C^i$  for any  $i \in \mathbb{Z}^+$ , then for every module  $M_v^l(\frac{\mathbb{k}Q^\infty}{jk})$  there exist  $m_v^l \in \mathbb{Z}^+$  such that  $\|M_v^l(A)\| < \|\Omega^{m_v^l}(M_v^l(A))\|$ . Using the last fact and that every module in  $\Omega^\infty(\frac{\mathbb{k}Q^\infty}{jk})$  must be a direct sum of modules  $M_v^l(\frac{\mathbb{k}Q^\infty}{jk})$  and a projective module we obtain the equality  $\Omega^\infty(\frac{\mathbb{k}Q^\infty}{jk}) = \mathcal{P}(\frac{\mathbb{k}Q^\infty}{jk})$ . By Corollary 38 we have that there are not  $\frac{\mathbb{k}Q^\infty}{jk}$ -periodic modules. By Proposition 43 we have that there are no  $A$ -periodic modules. Finally, again by Corollary 38, we have that  $\Omega^\infty(\text{mod} A) = \mathcal{P}(A)$ .


Finally consider  $Q$  a quiver such that  $Q^\infty$  has a final subheart equal to  $C^h$  for  $h \in \mathbb{Z}^+$ . Then there are two possible cases.

- Every final subheart of  $Q^\infty$  of type  $C^h$  can be extended to a subquiver  $\bar{Q}$  of  $Q$  as below





- There is a final subheart  $Q^\infty$  of type  $C^h$  that can not be extended as the previous case but can be extended to a subquiver  $\tilde{Q}$  of  $Q$  as below



In the second case, it is clear that there is a path  $\rho : i \rightarrow j$  with  $i, j \in \{1, \dots, h\}$  such that the right ideal  $\rho A$  is  $A$ -periodic but  $\rho$  is not a perfect path in  $Q$  because there are two paths  $\gamma_1$  and  $\gamma_2$  such that  $\rho \gamma_1 \in J^k$  and  $\rho \gamma_2 \in J^k$ . Therefore  $\rho A$  is a  $A$ -periodic non Co-Gorenstein module, and we deduce that  $A$  is not Co-Gorenstein.

In the first case, it is clear that there is a path  $\rho : i \rightarrow j$  with  $i, j \in \{1, \dots, h\}$  such that the right ideal  $\rho \frac{\mathbb{k}Q^\infty}{J^k}$  is  $\frac{\mathbb{k}Q^\infty}{J^k}$ -periodic but  $\rho$  is not a perfect path in  $Q$  because there are two paths  $\lambda_1$  and  $\lambda_2$  such that  $\lambda_1 \rho \in J^k$  and  $\lambda_2 \rho \in J^k$ . By Proposition 43, there is a  $A$ -module  $M \in \mathcal{P}^{<\infty}(A)$  such that  $\rho A \oplus M$  is  $A$ -periodic and it is not Gorenstein projective, thus we deduce that  $A$  is not Co-Gorenstein.

**Remark 45** Let  $A$  be an Artin algebra that verifies:

- $A$  is a truncated path algebra,
- $A$  is a Co-Gorenstein algebra, and
- $A$  is not a Gorenstein algebra,

then  $A$  is CM-free.

In [5] the authors prove that radical square zero algebras are either self-injective or CM-free. The example below shows that the behaviour of monomial algebras is more complicated. It shows that there are Co-Gorenstein monomial algebras that are not Gorenstein nor CM-free.

**Example 46** Consider the algebra  $A = \frac{\mathbb{k}Q}{I}$ , where

$$Q = 1 \begin{array}{c} \xrightarrow{\alpha} \\ \xleftarrow{\beta} \end{array} 2 \circlearrowright \gamma \text{ and } I = \langle \alpha\beta, \gamma^2 \rangle$$

Observe that  $\text{pd}(I_1 \oplus I_2) = \infty$ , hence  $A$  is not a Gorenstein algebra. On the other hand, the module  $A\gamma$  verifies  $\Omega(A\gamma) \cong A\gamma$ , and it is the only indecomposable periodic non projective module. In fact  $A\gamma$  is the only indecomposable non projective Gorenstein projective module by Theorem 22. We conclude that  $A$  is a Co-Gorenstein algebra and it is not CM-free.

## 5 $\Omega^\infty$ -infinite representation type and infinite $\phi$ -dimension

### 5.1 Igusa-Todorov function

The following lemma is a generalization of Proposition 3.9 of [14]. (For a proof see the cited proposition)

**Lemma 51** *Let  $A$  be an Artin algebra. If  $\mathcal{C}$  is a full subcategory of  $\text{mod}A$  closed by direct sums and direct summands and  $\Omega(\mathcal{C}) \subset \mathcal{C}$ , then*

$$\phi \dim(\mathcal{C}) = \min\{l : \bar{\Omega}|_{\Omega^l(\mathcal{K}_0(\mathcal{C}))} \text{ is injective}\}.$$

**Theorem 52** *Let  $Q$  be a quiver and  $C = \frac{\mathbb{k}Q}{I}$ . Suppose there exist two disjoint full subquivers of  $Q$ ,  $\Gamma$  and  $\bar{\Gamma}$  such that*

- $Q_0 = \Gamma_0 \cup \bar{\Gamma}_0$
- There is at least one arrow from  $\Gamma_0$  to  $\bar{\Gamma}_0$ , and there are no arrows of  $Q$  from  $\bar{\Gamma}_0$  to  $\Gamma_0$ .
- If  $\alpha \in Q_1$  with  $s(\alpha) \in \Gamma_0$  and  $t(\alpha) \in \bar{\Gamma}_0$  then  $\alpha\beta = 0$  for every arrow  $\beta$ .

Then  $\phi \dim(C) \leq \phi \dim(A) + \phi \dim(B) + 1$  where  $A = \frac{\mathbb{k}\Gamma}{I \cap \mathbb{k}\Gamma}$  and  $B = \frac{\mathbb{k}\bar{\Gamma}}{I \cap \mathbb{k}\bar{\Gamma}}$ .

*Proof* We denote by  $\partial A = \{v \in \Gamma_0 : \exists \alpha \in Q_1 \text{ with } s(\alpha) = v \text{ and } t(\alpha) \in \bar{\Gamma}_0\}$  and by  $\mathcal{P}_{\partial A}$  the full subcategory of  $\text{mod}A$  generated by the projective  $A$ -modules  $P_v$  with  $v \in \partial A$ . First note that if  $M \in \text{mod}C$  and

$$0 \rightarrow K \rightarrow P \rightarrow M \rightarrow 0$$

is an exact sequence in  $\text{mod}C$  with  $P$  a projective  $C$ -module, then  $K \cong M_1 \oplus M_2 \oplus M_3 \oplus Q$  where  $M_1 \in \Omega_A(\text{mod}A)$ ,  $M_2 \in \mathcal{P}_{\partial A}$ ,  $M_3 \in \text{mod}B$  and  $Q$  is a projective  $C$ -module.

Claim  $\phi(K) \leq \phi \dim(A) + \phi \dim(B)$

It is clear that

- $\Omega_C(M_1) = \Omega_A(M_1) \oplus N_1 \oplus N_2$  with  $N_1 \in \mathcal{P}_{\partial A}$  and  $N_2 \in \text{mod}B$ ,
- $\Omega_C(M_2) \in \text{mod}B$  and
- $\Omega_C(M_3) = \Omega_B(M_3)$ .

If  $\mathcal{D} = \Omega_A(\text{mod}A) \oplus \mathcal{P}_{\partial A} \oplus \text{mod}B$ , then  $K_0(\mathcal{D}) = K_1(A) \times K_0(\mathcal{P}_{\partial A}) \times K_0(B)$  and  $\bar{\Omega}_C$  restricted to  $K_0(\mathcal{D})$  is


$$\begin{pmatrix} \bar{\Omega}_A & 0 & 0 \\ T_1 & 0 & 0 \\ T_2 & T_3 & \bar{\Omega}_B \end{pmatrix}$$

Suppose that  $\phi \dim(C) \geq k$ , then, by Lemma 51, there exist  $k' \geq k$  and  $x = (x_1, x_2, x_3) \in K_0(\mathcal{D})$  such that  $\bar{\Omega}_C^{k'}(x) = 0$  and  $\bar{\Omega}_C^{k'-1}(x) \neq 0$ . Since  $\bar{\Omega}_A^i(x_1) = \Pi_{K_1(A)}(\bar{\Omega}_C^i(x))$  then  $\bar{\Omega}_A^{k'}(x_1) = 0$ . We deduce that  $\bar{\Omega}_A^l(x_1) = 0 \forall l \geq \phi \dim A - 1$  because  $x_1 \in K_1(A)$ , then  $\bar{\Omega}_C^{\phi \dim A - 1}(x) \in \{0\} \times K_0(\mathcal{P}_{\partial A}) \times K_0(B)$ . Hence  $k' \leq \phi \dim(A) + \phi \dim(B)$ . Finally, by Proposition 211 follows the thesis.

5.2 Examples

In [12] is showed that some Gorenstein algebras are of  $\Omega^\infty$ -infinite representation type (see example 4.1), hence this algebra has finite  $\phi$ -dimension (see Theorem 4.7 of [13]). Now we give an example of a non Gorenstein algebra of  $\Omega^\infty$ -infinite representation type, but its  $\phi$ -dimension is finite.

**Example 53** Let  $\mathbb{k}$  be an infinite field with characteristic different to 2. Consider the algebra  $A = \frac{\mathbb{k}Q}{I}$ , where  $Q$  is the quiver below:




and  $I = \langle \beta_1\alpha_1 - 2\beta_2\alpha_2, \alpha_1\beta_1 - \alpha_2\beta_2, \delta^2, \gamma\delta, \beta_1\gamma - \beta_2\gamma, \alpha_1\beta_1\gamma, \alpha_2\beta_1\gamma, \alpha_1\beta_2\gamma, \alpha_i\beta_j\alpha_k, \beta_i\alpha_j\beta_k \text{ with } i, j, k \in \{1, 2\} \rangle$ .

With the same computation from Example 310 we conclude that  $A$  is of  $\Omega^\infty$ -infinite representation type. It can be seen that the injective module  $I(3)$  has infinite projective dimension, hence  $A$  is not a Gorenstein algebra. Finally by Theorem 52 we conclude that  $\phi\dim(A) = 1$ .

In [13] was proved that  $\Omega^n$ -finite representation type algebras has finite  $\phi$ -dimension, therefore if we want to find an example of an algebra with infinite  $\phi$ -dimension, it must be of  $\Omega^\infty$ -infinite representation type.


The following example shows an radical cube zero algebra with infinite  $\phi$ -dimension. It is a counterexample of the conjecture stated in [7], however its finitistic dimension is finite (see [16]). Since  $\phi DC$  is not valid, then Theorem 213 in particular implies that the infinite  $\phi$ -dimensions are preserved by derived equivalences. The example also shows that the subcategory  $\Omega^\infty(\text{mod}A)$  can be trivial although the algebra  $A$  is of  $\Omega^\infty$ -infinite representation type.

**Example 54** Let  $A = \frac{\mathbb{k}Q}{I}$  be an algebra where  $Q$  is



and  $I = \langle \alpha_{i+1}\alpha_i - \bar{\alpha}_{i+1}\bar{\alpha}_i, \beta_{i+1}\beta_i - \bar{\beta}_{i+1}\bar{\beta}_i, \alpha_{i+1}\bar{\alpha}_i, \bar{\alpha}_{i+1}\alpha_i, \beta_{i+1}\bar{\beta}_i, \bar{\beta}_{i+1}\beta_i, J^3 \rangle$

It is clear that  $\text{pd}(S_i) = \infty$  for every  $i \in \mathbb{Z}_4$ . Let  $M_n^{\alpha_1}$  and  $M_n^{\beta_1}$ , with  $n \in \mathbb{Z}^+$ , be the indecomposable  $A$ -modules defined by



where the linear transformations  $i_m : \mathbb{k}^n \rightarrow \mathbb{k}^{3n+1}$ , with  $m \in \{1, 2, 3, 4\}$ , verifies:

- $i_1(e_j) = f_j \forall j \in \{1, \dots, n\}$ .
- $i_2(e_j) = f_{n+j} \forall j \in \{1, \dots, n\}$ .
- $i_3(e_j) = f_{n+j+1} \forall j \in \{1, \dots, n\}$ .
- $i_4(e_j) = f_{2n+j+1} \forall j \in \{1, \dots, n\}$ .

where  $\{e_1 \dots e_n\}$  and  $\{f_1, \dots, f_{3n+1}\}$  are the canonical bases of  $\mathbb{k}^n$  and  $\mathbb{k}^{3n+1}$  respectively.

In an analogous way we define  $M_n^{\alpha_2}, M_n^{\alpha_3}, M_n^{\alpha_4}$  and  $M_n^{\beta_2}, M_n^{\beta_3}, M_n^{\beta_4}$ . Then, the following assertions are valid for  $i \in \mathbb{Z}_4$ .

- $M_n^{\alpha_i} \not\cong M_n^{\beta_i}$  if  $n \geq 2$  and
- $M_1^{\alpha_i} \cong M_1^{\beta_i}$ .

On the other hand we compute the syzygies of the previous modules for  $n \geq 2$  and  $i \in \mathbb{Z}_4$  and we obtain

- $\Omega(M_n^{\alpha_i}) = M_{n-1}^{\alpha_{i+1}} \oplus S(i+2)^{7n+2}$ ,
- $\Omega(M_n^{\beta_i}) = M_{n-1}^{\beta_{i+1}} \oplus S(i+2)^{7n+2}$ .

And for  $n = 1$ , we have  $\Omega(M_1^{\alpha_i}) = \Omega(M_1^{\beta_i}) = S(i+2)^{10}$ .

By the previous facts we have  $\phi(M_n^{\alpha_i} \oplus M_n^{\beta_i}) = n - 1$  for any  $n \geq 2$ , then we conclude that  $\phi \dim(A) = \infty$ .

**Remark 55** Let  $A$  be the algebra defined in Example 54, then every  $A$ -module  $M \in \Omega^1(\text{mod}A)$  verifies  $\dim(\text{Top}(\Omega(M))) \geq \dim(\text{Top}(M)) + 1$ .

Consider an indecomposable  $A$ -module  $M \in \Omega^1(\text{mod}A)$ . We can suppose that  $\underline{\dim}(M) = (\alpha, \beta, 0, 0)$ , where  $1 \leq \beta \leq 4\alpha - 1$ . Then  $\underline{\dim}(\Omega(M)) = (0, 4\alpha - \beta, 10\alpha, 0)$ .

If  $4\alpha - \beta \geq \alpha + 1$ , we have  $\dim(\text{Top}(\Omega(M))) \geq \dim(\text{Top}(M)) + 1$ . Now suppose that  $4\alpha - \beta \leq \alpha$ , then  $(S_3)^{6n}$  is a direct summand of  $\Omega(M)$  therefore  $\dim(\text{Top}(\Omega(M))) \geq 6\dim(\text{Top}(M))$ .

As a consequence of the above computation we conclude that  $\Omega^\infty(\text{mod}A) = \mathcal{P}(A)$ . Therefore  $A$  is a CM-free Co-Gorenstein algebra.

## References

1. Apostolos Beligiannis. *The homological theory of contravariantly finite subcategories: Auslander-Buchweitz contexts, Gorenstein categories and (co-)stabilization*. Comm. Algebra, **28** (10) pp. 4547-4596 (2000).
2. Marcos Barrios, Gustavo Mata, Gustavo Rama. *Igusa-Todorov  $\phi$  function for truncated path algebras*, Algebras and Representation Theory (2019) <https://doi.org/10.1007/s10468-019-09883-7>
3. E. Babson, B. Huisgen-Zimmermann, and R. Thomas, *Generic representation theory of quivers with relations*, J. Algebra **322** (6), pp. 1877-1918 (2009)
4. Xiao-Wu Chen, Dawei Shen, Guodong Zhou. *The Gorenstein-projective modules over a monomial algebra*, Proceedings of the Royal Society of Edinburgh Section A: Mathematics **148** (6), pp. 1115-1134 (2018).
5. Xiao-Wu Chen, *Algebras with radical square zero are either self-injective or CM-free*, Proc. Am. Math. Soc. **140** (1) pp. 93-98 (2012).
6. A. Dugas, B. Huisgen-Zimmermann and J. Learned, *Truncated path algebras are homologically transparent. Part I*, Models, Modules and Abelian Groups (R. Göbel and B. Goldsmith, eds.), de Gruyter, Berlin, pp. 445-461 (2008).
7. S. Fernandes, M. Lanzilotta, O. Mendoza, *The Phi-dimension: A new homological measure*, Algebras and representation theory **17** (5), pp. 463-476 (2014).
8. M. Hoshino, *Algebras of finite self-injective dimension*, Proc. Amer. Math. Soc. **112** (3), pp. 619-622 (1991).
9. F. Huard, M. Lanzilotta, O. Mendoza. *An approach to the finitistic dimension conjecture*, J. Algebra **319** (9), pp. 3916-3934 (2008).
10. K. Igusa, G. Todorov, *On finitistic global dimension conjecture for artin algebras*, Representations of algebras and related topics, Fields Inst. Commun. **45**, American Mathematical Society, Providence, RI, pp. 201-204 (2005).
11. Sondre Kvamme, René Marczinzik, *Co-Gorenstein algebras*, applied categorical structures **27** (3), pp. 277-287 (2019).
12. Gustavo Mata, *Igusa-Todorov function on path rings*, Bol. Soc. Paran. Mat. To appear.
13. M. Lanzilotta, G. Mata, *Igusa-Todorov functions for Artin algebras*, Journal of pure and applied algebra, **222** (1), pp. 202-212 (2018).
14. M. Lanzilotta, E. Marcos, G. Mata, *Igusa-Todorov functions for radical square zero algebras*, J. Algebra **487**, pp. 357-385 (2017).
15. B. Zimmermann, *Predicting syzygies over monomial relations algebras*, Manuscripta Math. **70** (1), pp. 157-182 (1991).
16. B. Zimmermann, *Bounds on finitistic and global dimension for artinian rings with vanishing radical cube*, J. Algebra **161** pp. 47-68 (1993).
17. P. Zhang. *A brief introduction to Gorenstein projective modules*, Notes <https://www.math.uni-bielefeld.de/~sek/sem/abs/zhangpu4.pdf>