

# Anomalous Behavior of Magnetic Susceptibility Obtained by Quench Experiments in Isolated Quantum Systems

Yuuya Chiba,<sup>1,2,\*</sup> Kenichi Asano,<sup>3,†</sup> and Akira Shimizu<sup>1,2,‡</sup>

<sup>1</sup>*Komaba Institute for Science, The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8902, Japan*

<sup>2</sup>*Department of Basic Science, The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8902, Japan*

<sup>3</sup>*Center for Education in Liberal Arts and Sciences,  
Osaka University, Toyonaka, Osaka 560-0043, Japan*

(Dated: January 9, 2021)

We examine how the magnetic susceptibility obtained by the quench experiment on isolated quantum systems is related to the isothermal and adiabatic susceptibilities defined in thermodynamics. Under the conditions similar to the eigenstate thermalization hypothesis, together with some additional natural ones, we prove that for translationally invariant systems the quench susceptibility as a function of wave vector  $\mathbf{k}$  is discontinuous at  $\mathbf{k} = \mathbf{0}$ . Moreover, its values at  $\mathbf{k} = \mathbf{0}$  and the  $\mathbf{k} \rightarrow \mathbf{0}$  limit coincide with the adiabatic and the isothermal susceptibilities, respectively. We give numerical predictions on how these particular behaviors can be observed in experiments on the  $XYZ$  spin chain with tunable parameters, and how they deviate when the conditions are not fully satisfied.

*Introduction.*— Ultracold atoms [1, 2] and molecules [3–5] in optical lattices offer nearly ideal playgrounds for studying quantum many-body systems experimentally. Various model systems [6–21] are realized on the optical lattices with various geometry [22–27] and with tunable physical parameters [2, 28–31]. Furthermore, one can isolate the systems from the environments over a reasonably long period, which enables the direct observation of the dynamics of isolated quantum systems induced by suddenly changing a physical parameter [32–38]. After this so-called quench, the system often relaxes to a steady state, where the expectation values of local observables become almost time independent [21, 39–44]. The nature of such a steady state has been discussed in terms of the eigenstate thermalization hypothesis (ETH) [45–56]. For example, if the “strong” ETH is satisfied, the steady state is an equilibrium state [45–51].

In this Letter, we study the susceptibility obtained by the *quench* experiment, and explore whether or not it coincides with a thermodynamic susceptibility. This problem is highly nontrivial since there are two kinds of thermodynamic susceptibilities, the *isothermal* and the *adiabatic* ones, which take different values. In other words, it is not even clear which thermodynamic susceptibilities should be compared with the quench one. Furthermore, the wave number dependences of these susceptibilities make the problem more nontrivial, as we will reveal in this paper.

To be concrete, we consider the magnetic susceptibility of a quantum spin system. Suppose that the initial equilibrium state is in a uniform “offset” magnetic field  $h$ , and a weak extra magnetic field of wave number  $\mathbf{k}$  is suddenly applied. The *quench susceptibility*  $\chi^{\text{qch}}(\mathbf{k})$  is defined as the rate of magnetization change induced by such a quench. We explore its relation to the isothermal and the adiabatic thermodynamic susceptibilities,  $\chi^T(\mathbf{k})$  and  $\chi^S(\mathbf{k})$ , in the case where  $\chi^T(\mathbf{0}) > \chi^S(\mathbf{0})$ , which occurs when  $h \neq 0$ .

We reveal that  $\chi^{\text{qch}}(\mathbf{k})$  is *discontinuous* at  $\mathbf{k} = \mathbf{0}$  as a function of  $\mathbf{k}$ . Because of this discontinuity, *both* thermodynamic susceptibilities are obtained from the quench one, as  $\chi^{\text{qch}}(\mathbf{0}) = \chi^S(\mathbf{0})$  and  $\lim_{\mathbf{k} \rightarrow \mathbf{0}} \chi^{\text{qch}}(\mathbf{k}) = \chi^T(\mathbf{0})$ . The proof requires the conditions similar to the ETH, which hold when the dynamics of the system is complicated enough, as well as the natural conditions that are satisfied except at a phase transition point.

Furthermore, we numerically demonstrate how such anomalous behaviors should be observed in experiments on an isolated quantum spin system when it is nonintegrable. We also predict how the deviation from these behaviors is observed when the physical parameters of the system are tuned so that it becomes integrable.

*Setup.*— We deal with a quantum spin-1/2 system on a  $d$ -dimensional cubic lattice  $\Omega_N$  with linear size  $L$  and  $N = L^d$  spins. The periodic boundary conditions and the invariance under the discrete spatial translations are assumed for the prequench Hamiltonian  $\hat{H}(h)$ , where  $h$  denotes the uniform offset magnetic field. The density matrix of the initial state is chosen as the canonical Gibbs one,  $\hat{\rho}_{\text{ini}} = e^{-\beta \hat{H}(h)} / Z$  [57].

We are interested in the *quantum quench process* where the additional magnetic field  $\Delta h(\mathbf{r})$ , with wave number  $\mathbf{k}$  and small magnitude  $\Delta h_{\mathbf{k}}$ , is suddenly applied at  $t = 0$ . At  $t > 0$ , the isolated system obeys the Schrödinger dynamics of the postquench Hamiltonian,  $\hat{H}(h) - \sum_{\mathbf{r} \in \Omega_N} \hat{\sigma}_{\mathbf{r}}^z \Delta h(\mathbf{r})$ , where  $\hat{\sigma}_{\mathbf{r}}^\alpha$  ( $\alpha = x, y, z$ ) is the Pauli operator on site  $\mathbf{r} \in \Omega_N$ . For simplicity, we assume that  $\Delta h(\mathbf{r})$  is parallel to the offset field  $h$ , pointing in the  $z$  direction. While the previous works regarding the quantum quench focused only on the final state, we here study the *quench susceptibility*,

$$\chi_N^{\text{qch}}(\mathbf{k}) := \lim_{T \rightarrow \infty} \lim_{\Delta h_{\mathbf{k}} \rightarrow 0} \frac{\text{Tr}[\hat{\rho}(t) \hat{m}_{\mathbf{k}}] - \text{Tr}[\hat{\rho}_{\text{ini}} \hat{m}_{\mathbf{k}}]}{\Delta h_{\mathbf{k}}}, \quad (1)$$

which quantifies the *difference* of the expectation val-

ues of the  $\mathbf{k}$  component of magnetization,  $\hat{m}_{\mathbf{k}} = (1/N) \sum_{\mathbf{r} \in \Omega_N} e^{-i\mathbf{k} \cdot \mathbf{r}} \hat{\sigma}_{\mathbf{r}}^z$ , between the final and the initial states. Here,  $\hat{\rho}(t)$  is the density matrix at time  $t$ , and  $\overline{f(t)}^{\mathcal{T}}$  denotes the time average of  $f(t)$  over  $0 \leq t \leq \mathcal{T}$ .

For comparison, we consider the isothermal and the adiabatic thermodynamic susceptibilities,  $\chi_N^T(\mathbf{k})$  and  $\chi_N^S(\mathbf{k})$ , which are defined via the quasistatic processes with constant temperature and entropy, respectively. At  $\mathbf{k} = \mathbf{0}$ , they satisfy

$$\chi_N^S(\mathbf{0}) = \chi_N^T(\mathbf{0}) - \frac{T}{c_h} \left[ \left( \frac{\partial m_{\mathbf{0}}}{\partial T} \right)_h \right]^2, \quad (2)$$

where  $c_h$  is the specific heat at constant magnetic field and  $T = 1/\beta$  is the temperature [61]. We assume  $0 < T < +\infty$ , and exclude phase transition points where  $c_h$  diverges as  $N \rightarrow \infty$  and the case where  $(\partial m_{\mathbf{0}}/\partial T)_h$  vanishes, which is indeed unlikely for  $h \neq 0$ . This leads to the most interesting situation where the two susceptibilities take different values even in the thermodynamic limit,

$$\chi_{\infty}^S(\mathbf{0}) < \chi_{\infty}^T(\mathbf{0}), \quad (3)$$

where  $\chi_{\infty}^{\bullet}(\mathbf{k}) := \lim_{N \rightarrow \infty} \chi_N^{\bullet}(\mathbf{k})$ .

*Main results.*— Our main results are summarized as follows.

(i) The  $\mathbf{k} = \mathbf{0}$  value of the quench susceptibility agrees with that of the adiabatic one:

$$\chi_{\infty}^{\text{qch}}(\mathbf{0}) = \chi_{\infty}^S(\mathbf{0}), \quad (4)$$

if and only if condition (8), which is similar to but different from the ordinary ETH, is satisfied. Although the quench increases entropy, this equality implies it is irrelevant to  $\chi_{\infty}^{\text{qch}}(\mathbf{0})$  [61]. By contrast, the quench induces relevant changes in energy and temperature [61], which results in  $\chi_{\infty}^{\text{qch}}(\mathbf{0}) < \chi_{\infty}^T(\mathbf{0})$ .

(ii) The  $\mathbf{k} \neq \mathbf{0}$  value of the quench susceptibility agrees with those of the adiabatic and the isothermal ones [65],

$$\chi_{\infty}^{\text{qch}}(\mathbf{k}) = \chi_{\infty}^S(\mathbf{k}) = \chi_{\infty}^T(\mathbf{k}) \text{ for all } \mathbf{k} \neq \mathbf{0}, \quad (5)$$

if and only if condition (10), which is similar to but weaker than the ordinary “off-diagonal” ETH [44, 49–51], is satisfied.

(iii) The isothermal susceptibility,  $\chi_{\infty}^T(\mathbf{k})$ , is *uniformly* continuous as a function of  $\mathbf{k}$  [66] under two conditions (12) and (13) regarding the spatial spin-spin correlation function, both of which are fulfilled in normal systems.

(iv) When the conditions for (ii) and (iii), [namely, Eqs. (10), (12) and (13)] are all satisfied,

$$\lim_{\mathbf{k} \rightarrow \mathbf{0}} \chi_{\infty}^{\text{qch}}(\mathbf{k}) = \lim_{\mathbf{k} \rightarrow \mathbf{0}} \chi_{\infty}^T(\mathbf{k}) = \chi_{\infty}^T(\mathbf{0}). \quad (6)$$

This also shows that  $\chi_{\infty}^{\text{qch}}(\mathbf{k})$  is discontinuous at  $\mathbf{k} = \mathbf{0}$  because  $\chi_{\infty}^{\text{qch}}(\mathbf{0}) < \chi_{\infty}^T(\mathbf{0})$  as seen from the thermodynamic inequality (3) [69] and the general relation [61],

$$\chi_N^{\text{qch}}(\mathbf{0}) \leq \chi_N^S(\mathbf{0}). \quad (7)$$

(v) These results can be confirmed by a series of experiments in the isolated quantum systems, e.g., ultracold atoms, which simulate the  $XYZ$  spin chain. We predict the dependence of the above susceptibilities on  $\mathbf{k}$ ,  $N$ , and the exchange coupling parameters,  $J_x, J_y, J_z$ .

*Condition for (i).*— We introduce  $\hat{m}_{\mathbf{k}}^0 := \overline{e^{i\hat{H}(h)t} \hat{m}_{\mathbf{k}} e^{-i\hat{H}(h)t}}^{\mathcal{T}}$ , which is the energy-diagonal part of  $\hat{m}_{\mathbf{k}}$  [61]. Let  $|\nu\rangle$  be the simultaneous eigenstate of  $\hat{H}(h)$ , the translation operators, and  $\hat{m}_{\mathbf{k}=\mathbf{0}}^0$ , with eigenenergy  $E_{\nu}$  and crystal momentum  $\mathbf{K}_{\nu}$ . We also introduce  $\delta\hat{\sigma}_{\mathbf{r}}^z = \hat{\sigma}_{\mathbf{r}}^z - \text{Tr}[\hat{\rho}_{\text{ini}} \hat{\sigma}_{\mathbf{r}}^z]$  and  $\delta E_{\nu} = E_{\nu} - \text{Tr}[\hat{\rho}_{\text{ini}} \hat{H}(h)]$ . Then, we obtain the necessary and sufficient condition for (i) in the following form [61]: For *almost all*  $|\nu\rangle$  in a *narrow* energy region  $|\delta E_{\nu}| \lesssim T\sqrt{c_h N}$ , the diagonal elements  $\langle \nu | \delta\hat{\sigma}_{\mathbf{0}}^z | \nu \rangle$  are related almost linearly with  $\delta E_{\nu}$  as

$$\langle \nu | \delta\hat{\sigma}_{\mathbf{0}}^z | \nu \rangle = C \delta E_{\nu}/N + o(1/\sqrt{N}), \quad (8)$$

where  $C = \mathcal{O}(1)$  is some constant independent of  $\nu$  [70]. This is similar to but different from the ordinary two forms of ETH in the following points. The ordinary strong ETH [48–51] requires more stringently that *all*  $\langle \nu | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle$  behave like a smooth function of  $E_{\nu}/N$ , which is often satisfied in nonintegrable systems [71]. Since a smooth function of  $E_{\nu}/N$  can be regarded as linear within the narrow region  $|\delta E_{\nu}| \lesssim T\sqrt{c_h N}$ , any system satisfying the strong ETH also satisfies condition (8). By contrast, the ordinary weak ETH [53–55] requires only that  $\langle \nu | \delta\hat{\sigma}_{\mathbf{0}}^z | \nu \rangle = o(1)$  for almost all  $\nu$  in the same energy region. For this reason, some models that satisfy the ordinary weak ETH do not satisfy Eq. (8), as will be demonstrated shortly.

*Demonstration of (i).*— We now demonstrate how result (i) can be observed in experiments on the  $XYZ$  spin chain, which has the prequench Hamiltonian,

$$\hat{H}(h) = - \sum_{j=0}^{N-1} \sum_{\alpha=x,y,z} J_{\alpha} \hat{\sigma}_j^{\alpha} \hat{\sigma}_{j+1}^{\alpha} - \sum_{j=0}^{N-1} h \hat{\sigma}_j^z, \quad (9)$$

with periodic boundary condition,  $\hat{\sigma}_N = \hat{\sigma}_0$ . Since spin systems [14–21] and a 1D ring [22, 23] can be separately realized in ultracold atoms and molecules, we expect this model can also be realized experimentally. This model alone covers three different classes of systems, (a)  $XYZ$ , (b)  $XXZ$  ( $J_x = J_y \neq J_z$ ), and (c)  $XY$  ( $J_z = 0$ ) models, by tuning the parameters  $J_{\alpha}$ . We here predict the behaviors of the susceptibilities by means of the numerical diagonalization for (a) and (b), and the analytic evaluation for (c), respectively.

Figure 1(a) shows the  $N$  dependence of the  $k = 0$  components  $\chi_N^{\text{qch}}(0)$ ,  $\chi_N^T(0)$ , and  $\chi_N^S(0)$  in the  $XYZ$  model [72]. Since the model has no local conserved quantity for  $h \neq 0$  [73], it is expected that the condition (8) is fulfilled, so that Eq. (4) holds. In fact, Fig. 1(a) shows




FIG. 1. Size- $N$  dependence of  $\chi_N^{\text{qch}}(0)$ ,  $\chi_N^T(0)$ , and  $\chi_N^S(0)$  of the (a) XYZ, (b) XXZ, and (c) XY models. We take (a)  $(J_x - J_y, J_z) = (1.2, 1.0)$ , (b)  $(0.0, 1.0)$ , and (c)  $(1.2, 0.0)$ , for fixed values of  $J_x + J_y = 0.6$ ,  $h = 0.8$ ,  $\beta = 0.15$ . Inset of (a):  $\chi_N^S(0) - \chi_N^{\text{qch}}(0)$  in the logarithmic scale. Solid lines in (c):  $\chi_\infty^{\text{qch}}(0)$ ,  $\chi_\infty^T(0)$ , and  $\chi_\infty^S(0)$ .

that  $\chi_N^{\text{qch}}(0)$  approaches  $\chi_N^S(0)$  as  $N$  increases. Their difference decreases nearly exponentially, as shown in the inset, where the function  $0.083 e^{-0.193N}$  is also plotted as a guide to the eye. Both of them remain far off from  $\chi_N^T(0)$ .

Contrastingly, Eq. (4) does *not* hold for the XXZ or the XY models, as shown in Figs. 1(b) and 1(c), respectively. In these two cases, there exist some local conserved quantities that result in the violation of Eq. (8) and its equivalent (4). In other words, they do not satisfy Eq. (8) because of its integrability [74], while they *do satisfy* the ordinary “weak” ETH [53–56]. It should be noted that our results (a)-(c) are consistent with inequalities (3) and (7).

*Conditions for (ii).*— As is proved in Ref. [61], Eq. (5) holds if and only if *almost all*  $|\nu\rangle$  in a *narrow* energy region  $|\delta E_\nu| \lesssim T\sqrt{c_n N}$  satisfy

$$\sum_{\nu'} \delta_{E_\nu, E_{\nu'}} \delta_{\mathbf{K}_\nu, \mathbf{K}_{\nu'} + \mathbf{k}} |\langle \nu' | \hat{\sigma}_0^z | \nu \rangle|^2 = o(1/N) \quad \text{for all } \mathbf{k} \neq \mathbf{0}. \quad (10)$$

This is similar to the “off-diagonal ETH” [44, 49–52], except for the following points. First, the off-diagonal ETH requires that *all* off-diagonal elements of *all* local operators tend to vanish as  $N \rightarrow \infty$ . By contrast, Eq. (10) refers only to a *particular* spin operator  $\hat{\sigma}_0^z$  and to the off-diagonal elements between *specific* pairs of states such that

$$E_\nu = E_{\nu'} \quad \text{and} \quad \mathbf{K}_\nu = \mathbf{K}_{\nu'} + \mathbf{k}. \quad (11)$$

Furthermore, it requires not all such off-diagonal elements but *most* of them tend to vanish. Second, the ordinary off-diagonal ETH [44, 49–51] requires exponentially fast decay of all the off-diagonal elements, which

is not necessarily satisfied in integrable models. By contrast, Eq. (10) is a weaker condition [61] that can be satisfied even in integrable models, as we will demonstrate shortly for the XY model.

It is noteworthy that if we impose Eqs. (8) and (10) not only on a *particular* spin operator  $\hat{\sigma}_0^z$  but also on all other local operators, we obtain a *new necessary condition for thermalization*, which is also a sufficient condition as long as the quench parameter  $\Delta h_{\mathbf{k}}$  is small.

*Conditions for (iii).*— We introduce the canonical spin-spin correlation function [61, 75] as  $\phi_N^T(\mathbf{r}) := \beta \langle \delta \hat{\sigma}_0^z; \delta \hat{\sigma}_\tau^z \rangle_{\text{ini}}$ . Then, we can show [61] that  $\chi_\infty^T(\mathbf{k})$  is *uniformly continuous on the whole region* (including  $\mathbf{k} = \mathbf{0}$ ), if  $\phi_\infty^T(\mathbf{r})$  decays fast enough such that

$$\lim_{N \rightarrow \infty} \sum_{\mathbf{r} \in \Omega_N} |\phi_\infty^T(\mathbf{r})| < \infty \quad (12)$$

and if finite-size effects are small such that

$$\lim_{N \rightarrow \infty} \sum_{\mathbf{r} \in \Omega_N} |\phi_N^T(\mathbf{r}) - \phi_\infty^T(\mathbf{r})| = 0. \quad (13)$$

Since we exclude phase transition points, condition (12) is expected to be satisfied in most systems. Moreover, it seems normal that the condition (13) holds, since the canonical ensemble well emulates a subsystem in an infinite system [76, 77].

If conditions (10), (12) and (13) are all fulfilled, Eq. (6) follows from results (ii) and (iii). It also follows that  $\chi_\infty^{\text{qch}}(\mathbf{k})$  is discontinuous at  $\mathbf{k} = \mathbf{0}$ , as discussed in (iv).

*Demonstrations of (ii)-(iv).*— The discontinuity of  $\chi_\infty^{\text{qch}}(\mathbf{k})$  may seem counterintuitive, but can be verified experimentally by adopting the isolated system representing Eq. (9). The observed susceptibility should follow the following results of the numerical simulation.

Figure 2 shows the  $k$  dependence of  $\chi_N^{\text{qch}}(k)$ ,  $\chi_N^T(k)$ , and  $\chi_N^S(k)$  in the (a) XYZ, (b) XXZ, and (c) XY models. Recalling that the condition (10) is weaker than the ordinary off-diagonal ETH [44, 49–51], we expect that it is fulfilled in all these models. In fact, our data show that Eq. (5),  $\chi_\infty^{\text{qch}}(k) = \chi_\infty^S(k) = \chi_\infty^T(k)$  for all  $k \neq 0$ , holds in each model. We also find that  $\chi_N^T(k) - \chi_N^{\text{qch}}(k)$  for  $k \neq 0$  scales as  $\Theta(1/N)$  in (c). This is because the off-diagonal elements  $|\langle \nu' | \hat{\sigma}_0^z | \nu \rangle|$  that satisfy Eq. (11) decay not exponentially but algebraically as  $\Theta(1/N)$  for the XY model.

The conditions (12) and (13) are the natural ones that will also be satisfied in all these models. In fact, Figs. 2(a)-2(c) indicate Eq. (6),  $\lim_{k \rightarrow 0} \chi_\infty^{\text{qch}}(k) = \chi_\infty^T(0)$ , holds and hence  $\chi_\infty^{\text{qch}}(k)$  is discontinuous at  $k = 0$  while  $\chi_\infty^T(k)$  is uniformly continuous.

For the parameters presented here, Eqs. (5) and (6) hold in all three cases, while Eq. (4) only in the XYZ one. By further varying  $J_x$  and  $J_y$ , we can also construct a model for which *none* of Eqs. (4)-(6) holds [61]. In such a case, the condition (10) is violated, while the conditions (12) and (13) are still fulfilled.




FIG. 2.  $k$  dependence of  $\chi_N^{\text{qch}}(k)$ ,  $\chi_N^T(k)$ , and  $\chi_N^S(k)$  in (a) XYZ, (b) XXZ, and (c) XY models, with the same parameters as in Fig. 1. We take (a), (b)  $N = 12-17$  and  $k = 2\pi n_k/N$ , and (c)  $N = 2^n$  with  $n = 3-9$  and  $k = 2\pi n_k/8$ , with  $n_k = 0-4$ . Solid line in (c):  $\chi_N^{\text{qch}}(k) [= \chi_N^S(k) = \chi_N^T(k)]$  for  $k \neq 0$ , whereas the dashed line shows its discontinuous jump to  $\chi_\infty^{\text{qch}}(0)$ .

*Discussion on discontinuity.* — The discontinuity of  $\chi_\infty^{\text{qch}}(\mathbf{k})$  at  $\mathbf{k} = \mathbf{0}$  seems nontrivial. When results (i) and (ii) hold, this discontinuity is related to that of  $\chi_\infty^S(\mathbf{k})$ . To help understand the former, we here explain the latter discontinuity intuitively [66]. We also explain the continuity of  $\chi_\infty^T(\mathbf{k})$ .

Suppose that a huge system is enclosed by an adiabatic wall. We consider a thermodynamic process in which  $\Delta h(\mathbf{r})$  is applied quasistatically. If  $\Delta h(\mathbf{r})$  is localized and uniform in a subsystem with  $N = L^d$  sites, one obtains the adiabatic susceptibility of wave number  $|\mathbf{k}| \sim 1/L$ . Since the total system size is huge, it is well approximated by  $\chi_\infty^S(\mathbf{k})$ . On the other hand, this thermodynamic process can also be regarded as an isothermal process for the subsystem because the rest of the system works as a heat reservoir. According to this picture, one obtains  $\chi_N^T(\mathbf{0})$ . Since the two pictures have to give the same results,  $\chi_\infty^S(\mathbf{k}) = \chi_N^T(\mathbf{0})$  for  $|\mathbf{k}| \sim 1/L$ . By increasing  $N$ , we obtain  $\lim_{\mathbf{k} \rightarrow \mathbf{0}} \chi_\infty^S(\mathbf{k}) = \chi_\infty^T(\mathbf{0})$ . Comparing this with inequality (3), we can see that  $\chi_\infty^S(\mathbf{k})$  is discontinuous at  $\mathbf{k} = \mathbf{0}$ . By contrast, we can argue similarly the case where the adiabatic wall is replaced with a heat reservoir. Then we have  $\lim_{\mathbf{k} \rightarrow \mathbf{0}} \chi_\infty^T(\mathbf{k}) = \chi_\infty^T(\mathbf{0})$ , which shows that  $\chi_\infty^T(\mathbf{k})$  is continuous at  $\mathbf{k} = \mathbf{0}$ .

*Relation to Kubo formula.* — Since the Schrödinger dynamics is assumed, our results are applicable to experiments on isolated quantum systems. Moreover, since many formulas of physics were derived assuming the Schrödinger dynamics, our results contribute also to foundations of such formulas. As an example of the latter, we finally discuss the susceptibility obtained by the Kubo formula [78],  $\chi_N^{\text{Kubo}}(\mathbf{k}, \omega + i\varepsilon)$ . Here,  $\omega$  is the frequency and  $\varepsilon$  is an infinitesimal positive number. While we have defined  $\chi_N^{\text{qch}}$  through a sudden quench of  $\Delta h(\mathbf{r})$ , Kubo derived  $\chi_N^{\text{Kubo}}$  assuming that  $\Delta h(\mathbf{r})$  is switched on

gradually over a long timescale  $\sim 1/\varepsilon$ .

It is generally believed that the  $\varepsilon \rightarrow +0$  limit of  $\chi_N^{\text{Kubo}}$  should be taken *after* the  $N \rightarrow \infty$  limit [79–83]. However, some works took the  $\varepsilon \rightarrow +0$  limit keeping  $N$  finite [84–86]. For the latter limit, we can show [61]

$$\lim_{\varepsilon \rightarrow +0} \chi_N^{\text{Kubo}}(\mathbf{k}, 0 + i\varepsilon) = \chi_N^{\text{qch}}(\mathbf{k}) \quad \text{for all } N, \quad (14)$$

although the left-hand side and the right-hand side correspond to the slow and fast processes, respectively, which would result in different final states. Therefore, all the statements (i)-(iv) for  $\chi_\infty^{\text{qch}}(\mathbf{k})$  hold also for  $\lim_{N \rightarrow \infty} \lim_{\varepsilon \rightarrow +0} \chi_N^{\text{Kubo}}(\mathbf{k}, 0 + i\varepsilon)$  [87]. Moreover, the previous results on  $\lim_{\varepsilon \rightarrow +0} \chi_N^{\text{Kubo}}(\mathbf{0}, 0 + i\varepsilon)$  [84–86] can be understood more precisely using (i) [61]. However, it is noteworthy that  $\chi_N^{\text{Kubo}}$  is hard to measure in experiments in contrast to  $\chi_N^{\text{qch}}$ , since the system cannot be isolated for the infinitely long timescale.

In conclusion, we have revealed the anomalous natures of the quench susceptibility, demonstrating together that experimental verifications are feasible enough.

We thank Y. Yoneta, A. Noguchi and Y. Kato for discussions, and A. Ikeda, C. Hotta, R. Hamazaki and K. Saito for helpful comments. This work was supported by JSPS KAKENHI Grants No. JP19H01810, No. JP15H05700, and No. JP17K05497.

\* chiba@as.c.u-tokyo.ac.jp  
 † asano@celas.osaka-u.ac.jp  
 ‡ shmz@as.c.u-tokyo.ac.jp

- [1] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, A. Sen, and U. Sen, Ultracold atomic gases in optical lattices: mimicking condensed matter physics and beyond, *Adv. Phys.* **56**, 243 (2007).
- [2] I. Bloch, J. Dalibard, and W. Zwerger, Many-body physics with ultracold gases, *Rev. Mod. Phys.* **80**, 885 (2008).
- [3] A. Micheli, G. K. Brennen, and P. Zoller, A toolbox for lattice-spin models with polar molecules, *Nat. Phys.* **2**, 341 (2006).
- [4] C. Chin, V. V. Flambaum, and M. G. Kozlov, Ultracold molecules: new probes on the variation of fundamental constants, *New J. Phys.* **11**, 055048 (2009).
- [5] L. D. Carr, D. DeMille, R. V. Krems, and J. Ye, Cold and ultracold molecules: science, technology and applications, *New J. Phys.* **11**, 055049 (2009).
- [6] D. Jaksch, C. Bruder, J. I. Cirac, C. W. Gardiner, and P. Zoller, Cold Bosonic Atoms in Optical Lattices, *Phys. Rev. Lett.* **81**, 3108 (1998).
- [7] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, Quantum phase transition from a superfluid to a Mott insulator in a gas of ultracold atoms, *Nature (London)* **415**, 39 (2002).
- [8] T. Stöferle, H. Moritz, C. Schori, M. Köhl, and T. Esslinger, Transition from a Strongly Interacting 1D

- Superfluid to a Mott Insulator, *Phys. Rev. Lett.* **92**, 130403 (2004).
- [9] I. B. Spielman, W. D. Phillips, and J. V. Porto, Mott-Insulator Transition in a Two-Dimensional Atomic Bose Gas, *Phys. Rev. Lett.* **98**, 080404 (2007).
- [10] I. B. Spielman, W. D. Phillips, and J. V. Porto, Condensate Fraction in a 2D Bose Gas Measured across the Mott-Insulator Transition, *Phys. Rev. Lett.* **100**, 120402 (2008).
- [11] M. Köhl, H. Moritz, T. Stöferle, K. Günter, and T. Esslinger, Fermionic Atoms in a Three Dimensional Optical Lattice: Observing Fermi Surfaces, Dynamics, and Interactions, *Phys. Rev. Lett.* **94**, 080403 (2005).
- [12] M. F. Parsons, A. Mazurenko, C. S. Chiu, G. Ji, D. Greif, and M. Greiner, Site-resolved measurement of the spin-correlation function in the Fermi-Hubbard model, *Science* **353**, 1253 (2016).
- [13] T. Esslinger, Fermi-Hubbard Physics with Atoms in an Optical Lattice, *Annu. Rev. Condens. Matter Phys.* **1**, 129 (2010).
- [14] M. L. Wall, K. Maeda, and L. D. Carr, Realizing unconventional quantum magnetism with symmetric top molecules, *New J. Phys.* **17**, 025001 (2015).
- [15] G. Pelegrí, J. Mompart, V. Ahufinger, and A. J. Daley, Quantum magnetism with ultracold bosons carrying orbital angular momentum, *Phys. Rev. A* **100**, 023615 (2019).
- [16] J. Simon, W. S. Bakr, R. Ma, M. E. Tai, P. M. Preiss, and M. Greiner, Quantum simulation of antiferromagnetic spin chains in an optical lattice, *Nature (London)* **472**, 307 (2011).
- [17] T. Fukuhara, P. Schauß, M. Endres, S. Hild, M. Cheneau, I. Bloch, and C. Gross, Microscopic observation of magnon bound states and their dynamics, *Nature (London)* **502**, 76 (2013).
- [18] T. Fukuhara, S. Hild, J. Zeiher, P. Schauß, I. Bloch, M. Endres, and C. Gross, Spatially Resolved Detection of a Spin-Entanglement Wave in a Bose-Hubbard Chain, *Phys. Rev. Lett.* **115**, 035302 (2015).
- [19] B. Yan, S. A. Moses, B. Gadway, J. P. Covey, K. R. Hazzard, A. M. Rey, D. S. Jin, and J. Ye, Observation of dipolar spin-exchange interactions with lattice-confined polar molecules, *Nature (London)* **501**, 521 (2013).
- [20] K. R. Hazzard, B. Gadway, M. Foss-Feig, B. Yan, S. A. Moses, J. P. Covey, N. Y. Yao, M. D. Lukin, J. Ye, D. S. Jin, and A. M. Rey, Many-Body Dynamics of Dipolar Molecules in an Optical Lattice, *Phys. Rev. Lett.* **113**, 195302 (2014).
- [21] A. P. Orioli, A. Signoles, H. Wildhagen, G. Günter, J. Berges, S. Whitlock, and M. Weidemüller, Relaxation of an Isolated Dipolar-Interacting Rydberg Quantum Spin System, *Phys. Rev. Lett.* **120**, 063601 (2018).
- [22] L. Amico, A. Osterloh, and F. Cataliotti, Quantum Many Particle Systems in Ring-Shaped Optical Lattices, *Phys. Rev. Lett.* **95**, 063201 (2005).
- [23] K. Henderson, C. Ryu, C. MacCormick, and M. G. Boshier, Experimental demonstration of painting arbitrary and dynamic potentials for Bose-Einstein condensates, *New J. Phys.* **11**, 043030 (2009).
- [24] G. Wirth, M. Ölschläger, and A. Hemmerich, Evidence for orbital superfluidity in the P-band of a bipartite optical square lattice, *Nat. Phys.* **7**, 147 (2011).
- [25] P. Soltan-Panahi, J. Struck, P. Hauke, A. Bick, W. Plenkers, G. Meineke, C. Becker, P. Windpassinger, M. Lewenstein, and K. Sengstock, Multi-component quantum gases in spin-dependent hexagonal lattices, *Nat. Phys.* **7**, 434 (2011).
- [26] L. Tarruell, D. Greif, T. Uehlinger, G. Jotzu, and T. Esslinger, Creating, moving and merging Dirac points with a Fermi gas in a tunable honeycomb lattice, *Nature (London)* **483**, 302 (2012).
- [27] G. B. Jo, J. Guzman, C. K. Thomas, P. Hosur, A. Vishwanath, and D. M. Stamper-Kurn, Ultracold Atoms in a Tunable Optical Kagome Lattice, *Phys. Rev. Lett.* **108**, 045305 (2012).
- [28] H. Feshbach, Unified theory of nuclear reactions, *Ann. Phys. (N.Y.)* **5**, 357 (1958).
- [29] U. Fano, Effects of Configuration Interaction on Intensities and Phase Shifts, *Phys. Rev.* **124**, 1866 (1961).
- [30] E. Tiesinga, B. J. Verhaar, and H. T. C. Stoof, Threshold and resonance phenomena in ultracold ground-state collisions, *Phys. Rev. A* **47**, 4114 (1993).
- [31] H. T. Stoof, M. Houbiers, C. A. Sackett, and R. G. Hulet, Superfluidity of Spin-Polarized  ${}^6\text{Li}$ , *Phys. Rev. Lett.* **76**, 10 (1996).
- [32] M. Greiner, O. Mandel, T. W. Hänsch, and I. Bloch, Collapse and revival of the matter wave field of a Bose-Einstein condensate, *Nature (London)* **419**, 51 (2002).
- [33] L. E. Sadler, J. M. Higbie, S. R. Leslie, M. Vengalattore, and D. M. Stamper-Kurn, Spontaneous symmetry breaking in a quenched ferromagnetic spinor Bose-Einstein condensate, *Nature (London)* **443**, 312 (2006).
- [34] F. Meinert, M. J. Mark, E. Kirilov, K. Lauber, P. Weinmann, A. J. Daley, and H. C. Nägerl, Quantum Quench in an Atomic One-Dimensional Ising Chain, *Phys. Rev. Lett.* **111**, 053003 (2013).
- [35] G. Lamporesi, S. Donadello, S. Serafini, F. Dalfovo, and G. Ferrari, Spontaneous creation of Kibble-Zurek solitons in a Bose-Einstein condensate, *Nat. Phys.* **9**, 656 (2013).
- [36] C. L. Hung, V. Gurarie, and C. Chin, From Cosmology to Cold Atoms: Observation of Sakharov Oscillations in a Quenched Atomic Superfluid, *Science* **341**, 1213 (2013).
- [37] T. Fukuhara, A. Kantian, M. Endres, M. Cheneau, P. Schauß, S. Hild, D. Bellem, U. Schollwöck, T. Gimarchi, C. Gross, I. Bloch, and S. Kuhr, Quantum dynamics of a mobile spin impurity, *Nat. Phys.* **9**, 235 (2013).
- [38] S. Hild, T. Fukuhara, P. Schauß, J. Zeiher, M. Knap, E. Demler, I. Bloch, and C. Gross, Far-from-Equilibrium Spin Transport in Heisenberg Quantum Magnets, *Phys. Rev. Lett.* **113**, 147205 (2014).
- [39] S. Trotzky, Y. A. Chen, A. Flesch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, Probing the relaxation towards equilibrium in an isolated strongly correlated one-dimensional Bose gas, *Nat. Phys.* **8**, 325 (2012).
- [40] T. Kinoshita, T. Wenger, and D. S. Weiss, A quantum Newton's cradle, *Nature (London)* **440**, 900 (2006).
- [41] M. Gring, M. Kuhnert, T. Langen, T. Kitagawa, B. Rauer, M. Schreitl, I. Mazets, D. A. Smith, E. Demler, and J. Schmiedmayer, Relaxation and Prethermalization in an Isolated Quantum System, *Science* **337**, 1318 (2012).
- [42] H. Tasaki, From Quantum Dynamics to the Canonical Distribution: General Picture and a Rigorous Example, *Phys. Rev. Lett.* **80**, 1373 (1998).
- [43] P. Reimann, Foundation of Statistical Mechanics under Experimentally Realistic Conditions, *Phys. Rev. Lett.*

- 101**, 190403 (2008).
- [44] T. Mori, T. N. Ikeda, E. Kaminishi, and M. Ueda, Thermalization and prethermalization in isolated quantum systems: a theoretical overview, *J. Phys. B* **51**, 112001 (2018).
- [45] J. von Neumann, Proof of the ergodic theorem and the H-theorem in quantum mechanics, *Eur. Phys. J. H* **35**, 201 (2010).
- [46] J. M. Deutsch, Quantum statistical mechanics in a closed system, *Phys. Rev. A* **43**, 2046 (1991).
- [47] M. Srednicki, Chaos and quantum thermalization, *Phys. Rev. E* **50**, 888 (1994).
- [48] M. Rigol, V. Dunjko, and M. Olshanii, Thermalization and its mechanism for generic isolated quantum systems, *Nature (London)* **452**, 854 (2008).
- [49] M. Srednicki, The approach to thermal equilibrium in quantized chaotic systems, *J. Phys. A* **32**, 1163 (1999).
- [50] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, *Adv. Phys.* **65**, 239 (2016).
- [51] F. Anza, C. Gogolin, and M. Huber, Eigenstate Thermalization for Degenerate Observables, *Phys. Rev. Lett.* **120**, 150603 (2018).
- [52] S. Goldstein, D. A. Huse, J. L. Lebowitz, and R. Tumulka, Macroscopic and microscopic thermal equilibrium, *Ann. Phys. (Berlin)* **529**, 1600301 (2017).
- [53] G. Biroli, C. Kollath, and A. M. Läuchli, Effect of Rare Fluctuations on the Thermalization of Isolated Quantum Systems, *Phys. Rev. Lett.* **105**, 250401 (2010).
- [54] E. Iyoda, K. Kaneko, and T. Sagawa, Fluctuation Theorem for Many-Body Pure Quantum States, *Phys. Rev. Lett.* **119**, 100601 (2017).
- [55] T. Mori, Weak eigenstate thermalization with large deviation bound, arXiv:1609.09776.
- [56] T. Kuwahara and K. Saito, Ensemble equivalence and eigenstate thermalization from clustering of correlation, arXiv:1905.01886.
- [57] The initial state can be replaced with an appropriate pure quantum state that approximates the state in Ref. [58], which gives the same results as the Gibbs state for both equilibrium [58] and dynamical [59, 60] properties.
- [58] S. Sugiura and A. Shimizu, Canonical Thermal Pure Quantum State, *Phys. Rev. Lett.* **111**, 010401 (2013).
- [59] A. Shimizu and K. Fujikura, Quantum violation of fluctuation-dissipation theorem, *J. Stat. Mech.* (2017) 024004.
- [60] H. Endo, C. Hotta, and A. Shimizu, From Linear to Non-linear Responses of Thermal Pure Quantum States, *Phys. Rev. Lett.* **121**, 220601 (2018).
- [61] See Supplemental Material for detailed calculations, which includes Refs. [62–64].
- [62] P. Mazur, Non-ergodicity of phase functions in certain systems, *Physica* **43**, 533 (1969).
- [63] W. Beugeling, R. Moessner, and M. Haque, Finite-size scaling of eigenstate thermalization, *Phys. Rev. E* **89**, 042112 (2014).
- [64] R. Steinigeweg, A. Khodja, H. Niemeyer, C. Gogolin, and J. Gemmer, Pushing the Limits of the Eigenstate Thermalization Hypothesis towards Mesoscopic Quantum Systems, *Phys. Rev. Lett.* **112**, 130403 (2014).
- [65] The equality,  $\chi_N^S(\mathbf{k}) = \chi_N^T(\mathbf{k})$  for all  $\mathbf{k} \neq \mathbf{0}$  and for all  $N$ , can be proved only from the translation invariance [61].
- [66] See Refs. [67, 68] for related discussions.
- [67] J. L. Lebowitz, J. K. Percus, and L. Verlet, Ensemble Dependence of Fluctuations with Application to Machine Computations, *Phys. Rev.* **153**, 250 (1967).
- [68] J.-P. Hansen and I. R. McDonald, *Theory of Simple Liquids*, 4th ed. (Academic Press, Oxford, 2013).
- [69] If  $h = 0$ , so that  $m_{\mathbf{0}} = 0$ , we have  $\chi_{\infty}^S(\mathbf{0}) = \chi_{\infty}^T(\mathbf{0})$  unlike (3). Even in such a case, Eq. (6) shows that  $\chi_{\infty}^{\text{ch}}(\mathbf{k})$  is discontinuous at  $\mathbf{k} = \mathbf{0}$  unless the condition for (i) is satisfied.
- [70] Note that condition (8) can be satisfied even when  $C = 0$ , although  $C$  do not vanish in our setting where  $h \neq 0$  [61].
- [71] H. Kim, T. N. Ikeda, and D. A. Huse, Testing whether all eigenstates obey the eigenstate thermalization hypothesis, *Phys. Rev. E* **90**, 052105 (2014).
- [72] Here, we take  $J_y$  negative in (a) because we found that the  $o(1/\sqrt{N})$  term of Eq. (8) is smaller for larger  $J_x - J_y$ .
- [73] N. Shiraishi, Proof of the absence of local conserved quantities in the XYZ chain with a magnetic field, *Europhys. Lett.* **128**, 17002 (2019).
- [74] V. Alba, Eigenstate thermalization hypothesis and integrability in quantum spin chains, *Phys. Rev. B* **91**, 155123 (2015).
- [75] R. Kubo, M. Toda, and N. Hashitsume, *Statistical Physics II*, 2nd ed. (Springer, Berlin, 1991).
- [76] M. Hyuga, S. Sugiura, K. Sakai, and A. Shimizu, Thermal pure quantum states of many-particle systems, *Phys. Rev. B* **90**, 121110 (2014).
- [77] D. Iyer, M. Srednicki, and M. Rigol, Optimization of finite-size errors in finite-temperature calculations of unordered phases, *Phys. Rev. E* **91**, 062142 (2015).
- [78] R. Kubo, Statistical-Mechanical Theory of Irreversible Processes. I. General Theory and Simple Applications to Magnetic and Conduction Problems, *J. Phys. Soc. Jpn.* **12**, 570 (1957).
- [79] D. Pines and P. Nozieres, *The Theory of Quantum Liquids, Vol I: Normal Fermi Liquids* (W. A. Benjamin, New York, 1966).
- [80] G. Giuliani and G. Vignale, *Quantum Theory of the Electron Liquid* (Cambridge University Press, Cambridge, England, 2005).
- [81] D.N.Zubarev, *Nonequilibrium Statistical Thermodynamics* (Consultants Bureau, New York, 1974).
- [82] D. Zubarev, V. Morozov, and G. Röpke, *Statistical Mechanics of Nonequilibrium Processes, Volume 1: Basic Concepts, Kinetic Theory* (Akademie-Verlag, Berlin, 1996).
- [83] D. Zubarev, V. Morozov, and G. Röpke, *Statistical Mechanics of Nonequilibrium Processes, Volume 2: Relaxation and Hydrodynamic Processes* (Akademie-Verlag, Berlin, 1997).
- [84] H. Falk, Lower Bound for the Isothermal Magnetic Susceptibility, *Phys. Rev.* **165**, 602 (1968).
- [85] R. M. Wilcox, Bounds for the Isothermal, Adiabatic, and Isolated Static Susceptibility Tensors, *Phys. Rev.* **174**, 624 (1968).
- [86] M. Suzuki, Ergodicity, constants of motion, and bounds for susceptibilities, *Physica* **51**, 277 (1971).
- [87] For the former order of limits ( $\varepsilon \rightarrow +0$  after  $N \rightarrow \infty$ ), the conventional wisdom [79, 80] is that  $\lim_{\mathbf{k} \rightarrow \mathbf{0}} \lim_{\varepsilon \rightarrow +0} \lim_{N \rightarrow \infty} \chi_N^{\text{Kubo}}(\mathbf{k}, 0 + i\varepsilon) = \chi_{\infty}^T(\mathbf{0})$ , which however does not always hold. Our results (ii)-(iv) suggest the condition for the validity of this wisdom, although we have not yet proved  $\lim_{\varepsilon \rightarrow +0} \lim_{N \rightarrow \infty} \chi_N^{\text{Kubo}}(\mathbf{k}, 0 + i\varepsilon) =$

$$\chi_{\infty}^{\text{qch}}(\mathbf{k}).$$

# Anomalous Behavior of Magnetic Susceptibility Obtained by Quench Experiments in Isolated Quantum Systems: *Supplemental Material*

Yuuya Chiba\* and Akira Shimizu†

*Komaba Institute for Science, The University of Tokyo,  
3-8-1 Komaba, Meguro, Tokyo 153-8902, Japan and*

*Department of Basic Science, The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8902, Japan*

Kenichi Asano‡

*Center for Education in Liberal Arts and Sciences,  
Osaka University, Toyonaka, Osaka 560-0043, Japan*

(Dated: March 14, 2020)

## A. Quench susceptibility

We deal with a quantum spin system on a  $d$ -dimensional hypercubic lattice  $\Omega_N$  with linear size  $L$  and  $N = |\Omega_N| = L^d$  spins centered at  $\mathbf{r} = \mathbf{0}$ . The unit of length is taken as the lattice constant. We consider a quantum quench process where the weak additional field  $\Delta h(\mathbf{r})$ , with wave number  $\mathbf{k}$  and magnitude  $\Delta h_{\mathbf{k}}$ , is applied suddenly at  $t = 0$  and after that the expectation value of  $\hat{\sigma}_{\mathbf{r}}^z$  evolves in time as

$$\langle \hat{\sigma}_{\mathbf{r}}^z \rangle^{\text{qch}}(t) = \langle \hat{\sigma}_{\mathbf{r}}^z \rangle_{\text{ini}} + \sum_{\mathbf{r}' \in \Omega_N} \phi_N^{\text{qch}}(\mathbf{r} - \mathbf{r}'; t) \Delta h(\mathbf{r}') + \mathcal{O}(\Delta h_{\mathbf{k}}^2), \quad (\text{S1})$$

where  $\langle \bullet \rangle_{\text{ini}} = \text{Tr}[\hat{\rho}_{\text{ini}} \bullet]$ . Here,  $\phi_N^{\text{qch}}(\mathbf{r}; t) = \beta \langle \delta \hat{\sigma}_{\mathbf{0}}^z; \delta \hat{\sigma}_{\mathbf{r}}^z(t) \rangle_{\text{ini}} - \beta \langle \delta \hat{\sigma}_{\mathbf{0}}^z; \delta \hat{\sigma}_{\mathbf{r}}^z(t) \rangle_{\text{ini}}$  is a periodic function of  $\mathbf{r}$  with period  $L$ , where  $\hat{X}(t) = e^{i\hat{H}(h)t} \hat{X} e^{-i\hat{H}(h)t}$  is the Heisenberg operator and  $\langle \hat{X}; \hat{Y} \rangle_{\text{ini}} = \frac{1}{\beta} \int_0^\beta du \langle e^{u\hat{H}(h)} \hat{X}^\dagger e^{-u\hat{H}(h)} \hat{Y} \rangle_{\text{ini}}$  is the canonical correlation. Then, the response of  $\hat{m}_{\mathbf{k}}$  at time  $t$  reads  $\Delta \langle \hat{m}_{\mathbf{k}} \rangle^{\text{qch}}(t) = \langle \hat{m}_{\mathbf{k}} \rangle^{\text{qch}}(t) - \langle \hat{m}_{\mathbf{k}} \rangle_{\text{ini}} = \chi_N^{\text{qch}}(\mathbf{k}; t) \Delta h_{\mathbf{k}} + \mathcal{O}(\Delta h_{\mathbf{k}}^2)$ , where

$$\chi_N^{\text{qch}}(\mathbf{k}; t) = \sum_{\mathbf{r} \in \Omega_N} e^{-i\mathbf{k} \cdot \mathbf{r}} \phi_N^{\text{qch}}(\mathbf{r}; t) = \beta N \langle \delta \hat{m}_{\mathbf{k}}; \delta \hat{m}_{\mathbf{k}} \rangle_{\text{ini}} - \beta N \langle \delta \hat{m}_{\mathbf{k}}; \delta \hat{m}_{\mathbf{k}}(t) \rangle_{\text{ini}}. \quad (\text{S2})$$

Since we are only interested in the relaxed value of  $\Delta \langle \hat{m}_{\mathbf{k}} \rangle^{\text{qch}}(t)$ , we define the quench susceptibility  $\chi_N^{\text{qch}}(\mathbf{k})$  as the long time average of  $\chi_N^{\text{qch}}(\mathbf{k}; t)$ ,

$$\chi_N^{\text{qch}}(\mathbf{k}) = \lim_{\mathcal{T} \rightarrow \infty} \overline{\chi_N^{\text{qch}}(\mathbf{k}; t)}_{\mathcal{T}} = \beta N \langle \delta \hat{m}_{\mathbf{k}}; \delta \hat{m}_{\mathbf{k}} \rangle_{\text{ini}} - \beta N \langle \delta \hat{m}_{\mathbf{k}}^0; \delta \hat{m}_{\mathbf{k}}^0 \rangle_{\text{ini}}. \quad (\text{S3})$$

Here the energy diagonal part of an operator  $\hat{X}$  is given as  $\hat{X}^0 = \lim_{\mathcal{T} \rightarrow \infty} \overline{\hat{X}(t)}_{\mathcal{T}} = \sum_{\nu, \nu'} \delta_{E_\nu, E_{\nu'}} |\nu\rangle \langle \nu | \hat{X} | \nu' \rangle \langle \nu' |$ .

Figures S1(a) and S1(b) show the time dependence of  $\chi_N^{\text{qch}}(0; t)$  and  $\chi_N^{\text{qch}}(\pi/2; t)$  in 1D  $XYZ$  model, respectively. For  $t \gtrsim 5$ , i.e., after the transient regime,  $\chi_N^{\text{qch}}(\mathbf{k}; t)$  fluctuates in time around the quench susceptibility  $\chi_N^{\text{qch}}(\mathbf{k})$ , which is shown by the solid line. When the system size  $N$  is increased as 8, 12 and 16, this time fluctuation gets small. Therefore, if  $\chi_N^{\text{qch}}(\mathbf{k}; t)$  is measured after the transient regime, the measured value of  $\chi_N^{\text{qch}}(\mathbf{k}; t)$  will be close to  $\chi_N^{\text{qch}}(\mathbf{k})$ .

In the definition of  $\chi_N^{\text{qch}}(\mathbf{k})$ ,  $N$  is kept finite until the two limits  $\Delta h_{\mathbf{k}} \rightarrow 0$  and  $\mathcal{T} \rightarrow \infty$  are taken. This is reasonable in the sense that experiments on isolated quantum systems are performed only for finite  $N$ , such as  $N \simeq 30$  [1].

## B. Thermodynamic susceptibilities

We consider the isothermal quasistatic process in which the weak additional field is applied gradually and the final state of the system is the canonical Gibbs one,  $\hat{\rho}_{\text{fin}}^T \propto \exp(-\beta(\hat{H}(h) - \sum_{\mathbf{r} \in \Omega_N} \hat{\sigma}_{\mathbf{r}}^z \Delta h(\mathbf{r})))$ , with the same inverse

\* chiba@as.c.u-tokyo.ac.jp

† shmz@as.c.u-tokyo.ac.jp

‡ asano@celas.osaka-u.ac.jp






FIG. S1. Time dependence of (a)  $\chi_N^{\text{qch}}(0; t)$  and (b)  $\chi_N^{\text{qch}}(\pi/2; t)$  in XYZ model, with the parameters,  $J_x + J_y = 0.6$ ,  $J_x - J_y = 1.2$ ,  $J_z = 1.0$ ,  $h = 0.8$ , and  $\beta = 0.15$ . We take  $N = 8, 12, 16$ . The solid lines in (a) and (b) show  $\chi_N^{\text{qch}}(0)$  and  $\chi_N^{\text{qch}}(\pi/2)$  for each  $N$ , respectively. As the system size  $N$  is increased, the time fluctuation of  $\chi_N^{\text{qch}}(k; t)$  from its time average  $\chi_N^{\text{qch}}(k)$  gets small in both (a) and (b).

temperature as the initial one. Then, the expectation value of  $\hat{\sigma}_{\mathbf{r}}^z$  changes by

$$\Delta \langle \hat{\sigma}_{\mathbf{r}}^z \rangle^T = \text{Tr}[\hat{\rho}_{\text{fin}}^T \hat{\sigma}_{\mathbf{r}}^z] - \langle \hat{\sigma}_{\mathbf{r}}^z \rangle_{\text{ini}} = \sum_{\mathbf{r}' \in \Omega_N} \phi_N^T(\mathbf{r} - \mathbf{r}') \Delta h(\mathbf{r}') + \mathcal{O}(\Delta h_{\mathbf{k}}^2), \quad (\text{S4})$$

where  $\phi_N^T(\mathbf{r}) = \beta \langle \delta \hat{\sigma}_{\mathbf{0}}^z; \delta \hat{\sigma}_{\mathbf{r}}^z \rangle_{\text{ini}}$  is defined as a periodic function of  $\mathbf{r}$  in the same way as  $\phi_N^{\text{qch}}(\mathbf{r}; t)$ . From Eq. (S4), the response of  $\hat{m}_{\mathbf{k}}$  is given as  $\Delta \langle \hat{m}_{\mathbf{k}} \rangle^T = \text{Tr}[\hat{\rho}_{\text{fin}}^T \hat{m}_{\mathbf{k}}] - \langle \hat{m}_{\mathbf{k}} \rangle_{\text{ini}} = \chi_N^T(\mathbf{k}) \Delta h_{\mathbf{k}} + \mathcal{O}(\Delta h_{\mathbf{k}}^2)$ , where

$$\chi_N^T(\mathbf{k}) = \sum_{\mathbf{r} \in \Omega_N} e^{-i\mathbf{k} \cdot \mathbf{r}} \phi_N^T(\mathbf{r}) = \beta N \langle \delta \hat{m}_{\mathbf{k}}; \delta \hat{m}_{\mathbf{k}} \rangle_{\text{ini}} \quad (\text{S5})$$

is the isothermal susceptibility.

We also consider the adiabatic quasistatic process in which the weak additional field is applied gradually and the final state of the system is the canonical Gibbs one  $\hat{\rho}_{\text{fin}}^S \propto \exp(-\beta_{\text{fin}}^S (\hat{H}(h) - \sum_{\mathbf{r} \in \Omega_N} \hat{\sigma}_{\mathbf{r}}^z \Delta h(\mathbf{r})))$  with the same entropy as the initial one,  $-\text{Tr}[\hat{\rho}_{\text{fin}}^S \log \hat{\rho}_{\text{fin}}^S]/N = -\text{Tr}[\hat{\rho}_{\text{ini}} \log \hat{\rho}_{\text{ini}}]/N$ . From this condition, the final inverse temperature  $\beta_{\text{fin}}^S$  is determined as

$$\beta_{\text{fin}}^S = \beta + \sum_{\mathbf{r} \in \Omega_N} \beta \frac{\langle \delta \hat{H}(h) \delta \hat{\sigma}_{\mathbf{r}}^z \rangle_{\text{ini}}}{\langle \delta \hat{H}(h)^2 \rangle_{\text{ini}}} \Delta h(\mathbf{r}) + \mathcal{O}(\Delta h_{\mathbf{k}}^2). \quad (\text{S6})$$

The change of the expectation value of  $\hat{\sigma}_{\mathbf{r}}^z$ ,  $\Delta \langle \hat{\sigma}_{\mathbf{r}}^z \rangle^S = \text{Tr}[\hat{\rho}_{\text{fin}}^S \hat{\sigma}_{\mathbf{r}}^z] - \langle \hat{\sigma}_{\mathbf{r}}^z \rangle_{\text{ini}}$ , is given as

$$\Delta \langle \hat{\sigma}_{\mathbf{r}}^z \rangle^S = \Delta \langle \hat{\sigma}_{\mathbf{r}}^z \rangle^T - (\beta_{\text{fin}}^S - \beta) \langle \delta \hat{H}(h) \delta \hat{\sigma}_{\mathbf{r}}^z \rangle_{\text{ini}} + \mathcal{O}(\Delta h_{\mathbf{k}}^2) = \sum_{\mathbf{r}' \in \Omega_N} \phi_N^S(\mathbf{r} - \mathbf{r}') \Delta h(\mathbf{r}') + \mathcal{O}(\Delta h_{\mathbf{k}}^2), \quad (\text{S7})$$

where

$$\phi_N^S(\mathbf{r}) = \phi_N^T(\mathbf{r}) - \beta \frac{\langle \delta \hat{H}(h) \delta \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{ini}}^2}{\langle \delta \hat{H}(h)^2 \rangle_{\text{ini}}}. \quad (\text{S8})$$

Then, the response of  $\hat{m}_{\mathbf{k}}$  is also given as  $\Delta \langle \hat{m}_{\mathbf{k}} \rangle^S = \text{Tr}[\hat{\rho}_{\text{fin}}^S \hat{m}_{\mathbf{k}}] - \langle \hat{m}_{\mathbf{k}} \rangle_{\text{ini}} = \chi_N^S(\mathbf{k}) \Delta h_{\mathbf{k}} + \mathcal{O}(\Delta h_{\mathbf{k}}^2)$ , where

$$\chi_N^S(\mathbf{k}) = \sum_{\mathbf{r} \in \Omega_N} e^{-i\mathbf{k} \cdot \mathbf{r}} \phi_N^S(\mathbf{r}) = \chi_N^T(\mathbf{k}) - \beta N \frac{|\langle \delta \hat{H}(h) \delta \hat{m}_{\mathbf{k}} \rangle_{\text{ini}}|^2}{\langle \delta \hat{H}(h)^2 \rangle_{\text{ini}}} \quad (\text{S9})$$

is the adiabatic susceptibility.

### C. Relations between the susceptibilities

From Eq. (S9), we have  $\chi_N^S(\mathbf{k}) = \chi_N^T(\mathbf{k}) - \frac{T}{c_h} |(\partial m_{\mathbf{k}} / \partial T)_h|^2$ , where  $c_h = \beta^2 \langle \delta \hat{H}(h)^2 \rangle_{\text{ini}} / N$  is the specific heat at constant magnetic field and  $(\partial m_{\mathbf{k}} / \partial T)_h = -\beta^2 \langle \delta \hat{H}(h) \delta \hat{m}_{\mathbf{k}} \rangle_{\text{ini}}$ . In contrast to  $\mathbf{k} = \mathbf{0}$  component,  $(\partial m_{\mathbf{k}} / \partial T)_h = 0$  hold for all  $\mathbf{k} \neq \mathbf{0}$  because of the translation invariance of  $\hat{H}(h)$ , yielding

$$\chi_N^S(\mathbf{k}) = \chi_N^T(\mathbf{k}) \quad \text{for all } \mathbf{k} \neq \mathbf{0}. \quad (\text{S10})$$

Comparing Eqs. (S3) and (S9), we have

$$\chi_N^S(\mathbf{0}) - \chi_N^{\text{qch}}(\mathbf{0}) = \beta N \langle \delta \hat{m}_{\mathbf{k}=\mathbf{0}}^0; \delta \hat{m}_{\mathbf{k}=\mathbf{0}}^0 \rangle_{\text{ini}} - \beta N \frac{|\langle \delta \hat{H}(h) \delta \hat{m}_{\mathbf{k}=\mathbf{0}} \rangle_{\text{ini}}|^2}{\langle \delta \hat{H}(h)^2 \rangle_{\text{ini}}} \quad (\text{S11})$$

$$= \beta N \left( \sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} \langle \nu | \delta \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle^2 \right) - \beta N \left( \sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} \delta E_{\nu} \langle \nu | \delta \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle \right)^2 / \left( \sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} \delta E_{\nu}^2 \right) \geq 0 \quad (\text{S12})$$

from the Cauchy-Schwarz inequality. Here  $\langle \nu' | \hat{m}_{\mathbf{k}=\mathbf{0}}^0 | \nu \rangle = \delta_{\nu, \nu'} \langle \nu | \hat{m}_{\mathbf{k}=\mathbf{0}}^0 | \nu \rangle = \delta_{\nu, \nu'} \langle \nu | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle$  holds, since  $|\nu\rangle$  is the simultaneous eigenstate of  $\hat{H}(h)$ , translation operators, and  $\hat{m}_{\mathbf{k}=\mathbf{0}}^0$ . This yields the general relation (7) [2–4]. The equality for finite  $N$  holds if and only if  $\langle \nu | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle = C \delta E_{\nu}/N$  for all  $\nu$ , where  $C$  is some constant independent of  $\nu$ . This is not satisfied in almost all systems. In the thermodynamic limit  $N \rightarrow \infty$ , the condition for the equality is relaxed as follows.

Result (i) : From Eq. (S12), the necessary and sufficient condition for Eq. (4) is given as

$$\lim_{N \rightarrow \infty} \sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} N \left| \langle \nu | \delta \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle - \left( \sum_{\nu'} \frac{e^{-\beta E_{\nu'}}}{Z} \delta E_{\nu'} \langle \nu' | \delta \hat{\sigma}_{\mathbf{0}}^z | \nu' \rangle \right) \delta E_{\nu} / \left( \sum_{\nu'} \frac{e^{-\beta E_{\nu'}}}{Z} \delta E_{\nu'}^2 \right) \right|^2 = 0. \quad (\text{S13})$$

This condition can be rephrased as Eq. (8), where the constant  $C$  is given by

$$C = N \left( \sum_{\nu'} \frac{e^{-\beta E_{\nu'}}}{Z} \delta E_{\nu'} \langle \nu' | \delta \hat{\sigma}_{\mathbf{0}}^z | \nu' \rangle \right) / \left( \sum_{\nu'} \frac{e^{-\beta E_{\nu'}}}{Z} \delta E_{\nu'}^2 \right) + o(1), \quad (\text{S14})$$

which does not vanish in our case for the reasons explained below Eq. (2) of the main text.

We can relate condition (8) with the ordinary ETH more directly. Let us introduce the microcanonical average over the energy shell  $(E - \delta, E]$  as  $\langle \bullet \rangle_{\text{mc}}(E/N)$  and the number of states in  $(E - \delta, E]$  as  $W(E/N)$ , assuming that the energy width  $\delta$  can be taken as  $\delta_N = \Theta(1/N^{1+\alpha})$ , where  $\alpha$  is some small positive number. Then we can evaluate  $\langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{ini}}$  as

$$\langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{ini}} = \sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} \langle \nu | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle = \frac{\int de \exp(N(s_N(e) - \beta e)) \langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{mc}}(e)}{\int de \exp(N(s_N(e) - \beta e))} + \mathcal{O}(\delta_N), \quad (\text{S15})$$

where  $s_N(e) = \log W(e)/N$ . Except at a phase transition point, we can use the saddle point method and obtain

$$\langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{ini}} = \langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{mc}}(e^*) + \mathcal{O}(1/N), \quad (\text{S16})$$

where  $e^*$  is determined by  $s'_N(e^*) = \frac{ds_N}{de}(e^*) = \beta$ . In the same way,

$$\langle \hat{H}(h) \rangle_{\text{ini}}/N = e^* + \mathcal{O}(1/N), \quad (\text{S17})$$

$$\sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} \delta E_{\nu}^2 / N = 1/|s''_N(e^*)| + o(1), \quad (\text{S18})$$

$$\sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} \delta E_{\nu} \langle \nu | \delta \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle = \frac{d\langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{mc}}}{de}(e^*) / |s''_N(e^*)| + o(1), \quad (\text{S19})$$

$$N \sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} \langle \nu | \delta \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle^2 = N \sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} |\langle \nu | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle - \langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{mc}}(E_{\nu}/N)|^2 + \left( \frac{d\langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{mc}}}{de}(e^*) \right)^2 / |s''_N(e^*)| + o(1) \quad (\text{S20})$$

can be shown. From Eqs. (S18), (S19), and (S20), the following result holds.

Result (i') : Eq. (4) or its equivalent condition (8) holds if and only if

$$N \sum_{\nu} \frac{e^{-\beta E_{\nu}}}{Z} |\langle \nu | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle - \langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{mc}}(E_{\nu}/N)|^2 = o(1), \quad (\text{S21})$$

which is similar to the weak ETH [5–7] in that it requires almost all  $\langle \nu | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle$  should be close to  $\langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{mc}}(E_{\nu}/N)$ . Condition (S21) will be satisfied in nonintegrable systems, where  $\langle \nu | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle$  is often exponentially close to  $\langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{mc}}(E_{\nu}/N)$  [8, 9]. Note that, there are some integrable models which satisfy the ordinary weak ETH [5, 6, 10] but do not satisfy condition (S21). This fact can be confirmed by the violation of its equivalent Eq. (4),  $\chi_{\infty}^{\text{qch}}(\mathbf{0}) = \chi_{\infty}^S(\mathbf{0})$ . (See main text.) Indeed, condition (S21) is more stringent than the ordinary weak ETH [5, 6, 10] in that condition (S21) requires

$|\langle \nu | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle - \langle \hat{\sigma}_{\mathbf{0}}^z \rangle_{\text{mc}}(E_\nu/N)|^2$  to be typically  $o(1/N)$ , while the ordinary weak ETH [5, 6, 10] allows this quantity to be larger than  $\Theta(1/N)$ . Here, functions of  $N$ ,  $f_N$  and  $g_N$ , satisfy  $g_N = \Theta(f_N)$ , if there are positive constants  $0 < c_1 \leq c_2 < \infty$  such that  $c_1 f_N \leq g_N \leq c_2 f_N$  holds for sufficiently large  $N$ .

Eqs. (S3) and (S5) give a relation between  $\mathbf{k} \neq \mathbf{0}$  components,

$$\chi_N^T(\mathbf{k}) - \chi_N^{\text{qch}}(\mathbf{k}) = \beta N \langle \delta \hat{m}_{\mathbf{k}}^0; \delta \hat{m}_{\mathbf{k}}^0 \rangle = \beta N \sum_{\nu} \frac{e^{-\beta E_\nu}}{Z} \sum_{\nu'} \delta_{E_\nu, E_{\nu'}} \delta_{\mathbf{K}_\nu, \mathbf{K}_{\nu'} + \mathbf{k}} |\langle \nu' | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle|^2, \quad (\text{S22})$$

where the crystal momentum  $\mathbf{K}_\nu$  is defined so that the eigenvalue of  $\mathbf{r}$  sites translation operator is written as  $e^{-i\mathbf{K}_\nu \cdot \mathbf{r}}$  and we used  $|\langle \nu' | \hat{m}_{\mathbf{k}} | \nu \rangle| = \delta_{\mathbf{K}_\nu, \mathbf{K}_{\nu'} + \mathbf{k}} |\langle \nu' | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle|$ . Therefore Eqs. (S10) and (S22) yield the following.

Result (ii) : Eq. (5) holds if and only if the off-diagonal elements are small so that

$$\sum_{\nu} \frac{e^{-\beta E_\nu}}{Z} N \sum_{\nu'} \delta_{E_\nu, E_{\nu'}} \delta_{\mathbf{K}_\nu, \mathbf{K}_{\nu'} + \mathbf{k}} |\langle \nu' | \hat{\sigma}_{\mathbf{0}}^z | \nu \rangle|^2 = o(1) \quad \text{for all } \mathbf{k} \neq \mathbf{0}. \quad (\text{S23})$$

This condition can be rephrased as Eq. (10), which is weaker than the ordinary off-diagonal ETH [11–14] as explained below using  $XY$  model.

#### D. Analysis of the quench process using thermodynamics

In this section, we assume that thermalization occurs after the quench process, where the small uniform magnetic field  $\Delta h_{\mathbf{0}}$  is applied. From this assumption, the state of the system, which evolves from the initial equilibrium state, relaxes to another equilibrium state. Since the expectation value of the postquench Hamiltonian  $\hat{H}(h + \Delta h_{\mathbf{0}})$  does not change before and after the quench, the initial and the final equilibrium states satisfy

$$e + \Delta e = \langle \hat{H}(h + \Delta h_{\mathbf{0}}) \rangle_{\text{fin}}/N = \langle \hat{H}(h + \Delta h_{\mathbf{0}}) \rangle_{\text{ini}}/N = \langle \hat{H}(h) \rangle_{\text{ini}}/N - \langle \hat{m}_{\mathbf{0}} \rangle_{\text{ini}} \Delta h_{\mathbf{0}} = e - m_{\mathbf{0}} \Delta h_{\mathbf{0}}, \quad (\text{S24})$$

where  $e$  and  $e + \Delta e$  are the initial and the final equilibrium values of the energy per site and  $m_{\mathbf{0}}$  is the initial equilibrium value of the  $\mathbf{k} = \mathbf{0}$  component of the magnetization. From Eq. (S24), the change of the entropy is

$$\Delta s = \frac{\partial s}{\partial e}(e, h) \Delta e + \frac{\partial s}{\partial h}(e, h) \Delta h_{\mathbf{0}} + \mathcal{O}((\Delta h_{\mathbf{0}})^2) = \beta \Delta e + \beta m_{\mathbf{0}} \Delta h_{\mathbf{0}} + \mathcal{O}((\Delta h_{\mathbf{0}})^2) = \mathcal{O}((\Delta h_{\mathbf{0}})^2), \quad (\text{S25})$$

where  $\beta$  is the initial inverse temperature. Eq. (S25) is consistent with Eq. (4). Note that the change of the energy per site  $\Delta e = -m_{\mathbf{0}} \Delta h_{\mathbf{0}}$  and the change of the inverse temperature

$$\Delta \beta = \frac{\beta}{c_h} \left( \frac{\partial m_{\mathbf{0}}}{\partial T} \right)_h \Delta h_{\mathbf{0}} + \mathcal{O}((\Delta h_{\mathbf{0}})^2) \quad (\text{S26})$$

are  $\mathcal{O}(\Delta h_{\mathbf{0}})$  because  $h \neq 0$ . This results in  $\chi_{\infty}^{\text{qch}}(\mathbf{0}) < \chi_{\infty}^T(\mathbf{0})$ .

#### E. Proof of (iii)

From condition (12), we can define  $\chi^{\text{inf}}(\mathbf{k}) = \lim_{N \rightarrow \infty} \sum_{\mathbf{r} \in \Omega_N} e^{-i\mathbf{k} \cdot \mathbf{r}} \phi_{\infty}^T(\mathbf{r})$ , which is uniformly continuous in  $\mathbf{k}$  by the property of Fourier transform. From Eq. (S5),

$$|\chi_N^T(\mathbf{k}) - \chi^{\text{inf}}(\mathbf{k})| \leq \left| \sum_{\mathbf{r} \in \Omega_N} e^{-i\mathbf{k} \cdot \mathbf{r}} (\phi_N^T(\mathbf{r}) - \phi_{\infty}^T(\mathbf{r})) \right| + \left| \lim_{N' \rightarrow \infty} \sum_{\mathbf{r} \in \Omega_{N'} \setminus \Omega_N} e^{-i\mathbf{k} \cdot \mathbf{r}} \phi_{\infty}^T(\mathbf{r}) \right| \quad (\text{S27})$$

$$\leq \sum_{\mathbf{r} \in \Omega_N} |\phi_N^T(\mathbf{r}) - \phi_{\infty}^T(\mathbf{r})| + \lim_{N' \rightarrow \infty} \sum_{\mathbf{r} \in \Omega_{N'} \setminus \Omega_N} |\phi_{\infty}^T(\mathbf{r})|. \quad (\text{S28})$$

In the  $N \rightarrow \infty$  limit, the first term and the second term of Eq. (S28) converges to 0 from condition (13) and (12), respectively. As a result,  $\chi_N^T(\mathbf{k})$  converges to  $\chi^{\text{inf}}(\mathbf{k})$  in the  $N \rightarrow \infty$  limit,  $\chi_{\infty}^T(\mathbf{k}) = \chi^{\text{inf}}(\mathbf{k})$  for all  $\mathbf{k}$ , which implies that  $\chi_{\infty}^T(\mathbf{k})$  is also uniformly continuous in  $\mathbf{k}$ .  $\square$

Note that condition (13) is essential for the uniform continuity of  $\chi_{\infty}^T(\mathbf{k})$ . Since  $\phi_{\infty}^S(\mathbf{r}) = \phi_{\infty}^T(\mathbf{r})$  for all  $\mathbf{r}$  follows from Eq. (S8), condition (12) holds also for  $\phi^S$ . However condition (13) does not hold for  $\phi^S$ :

$$\lim_{N \rightarrow \infty} \sum_{\mathbf{r} \in \Omega_N} |\phi_N^S(\mathbf{r}) - \phi_{\infty}^S(\mathbf{r})| = \lim_{N \rightarrow \infty} \sum_{\mathbf{r} \in \Omega_N} |\phi_N^S(\mathbf{r}) - \phi_N^T(\mathbf{r}) + \phi_N^T(\mathbf{r}) - \phi_{\infty}^T(\mathbf{r})| = \chi_{\infty}^T(\mathbf{0}) - \chi_{\infty}^S(\mathbf{0}) > 0, \quad (\text{S29})$$




FIG. S2. Verification of conditions for (iii) in  $XYZ$  model, with the same parameters as in Fig. S1. We investigate the  $N$  dependence of (a)  $D_N$ , the sum of  $|\phi_{N_{\max}}^T(\mathbf{r})|$  over all  $\mathbf{r} \in \Omega_N$ , and (b)  $F_N$ , the sum of  $|\phi_N^T(\mathbf{r}) - \phi_{N_{\max}}^T(\mathbf{r})|$  over all  $\mathbf{r} \in \Omega_N$ . We take  $N_{\max} = 16$ .

which is consistent with the discontinuity of  $\chi_\infty^S(\mathbf{k})$  at  $\mathbf{k} = \mathbf{0}$ .

In Fig. S2, we verify the conditions for (iii), (a)  $\phi_N^T(\mathbf{r})$  decays fast enough and (b) finite-size effects of  $\phi_N^T(\mathbf{r})$  are small, in  $XYZ$  model. To this end, we introduce two quantities, (a)  $D_N = \sum_{\mathbf{r} \in \Omega_N} |\phi_{N_{\max}}^T(\mathbf{r})|$  and (b)  $F_N = \sum_{\mathbf{r} \in \Omega_N} |\phi_N^T(\mathbf{r}) - \phi_{N_{\max}}^T(\mathbf{r})|$ , where  $N_{\max}$  is taken as large as possible. Fig. S2 (a) shows  $N$  dependence of  $D_N$  in  $XYZ$  model. As  $N$  increases,  $D_N$  is saturated, suggesting that condition (12) holds. Fig. S2 (b) shows  $N$  dependence of  $F_N$  in the same system. As  $N$  increases,  $F_N$  decreases, suggesting that condition (13) holds.

### F. Analytic solutions in 1D $XY$ model

We here describe the analytic solutions  $\chi_\infty^{\text{qch}}(k)$ ,  $\chi_\infty^S(k)$ , and  $\chi_\infty^T(k)$  in 1D  $XY$  model and verify whether the above relations hold or not in this model. By defining  $J_s = J_x + J_y$ ,  $J_a = J_x - J_y$ , and  $\varepsilon_k = \sqrt{(J_s \cos k + h)^2 + J_a^2 \sin^2 k}$ , we can write the results as follows.

For the  $k = 0$  components, we have

$$\chi_\infty^{\text{qch}}(0) = \frac{1}{2\pi} \int_0^{2\pi} dk' \frac{J_a^2 \sin^2 k' \tanh \beta \varepsilon_{k'}}{\varepsilon_{k'}^2 \varepsilon_{k'}}, \quad (\text{S30})$$

$$\chi_\infty^T(0) = \chi_\infty^{\text{qch}}(0) + \frac{\beta}{2\pi} \int_0^{2\pi} dk' \frac{(J_s \cos k' + h)^2}{\varepsilon_{k'}^2} \frac{1}{\cosh^2 \beta \varepsilon_{k'}}, \quad (\text{S31})$$

$$\chi_\infty^S(0) = \chi_\infty^T(0) - \left( \frac{\beta}{2\pi} \int_0^{2\pi} dk' \frac{J_s \cos k' + h}{\cosh^2 \beta \varepsilon_{k'}} \right)^2 / \left( \frac{\beta}{2\pi} \int_0^{2\pi} dk' \frac{\varepsilon_{k'}^2}{\cosh^2 \beta \varepsilon_{k'}} \right) < \chi_\infty^T(0). \quad (\text{S32})$$

From Eqs. (S30) and (S32), Eq. (4) is violated except at  $J_s = J_a = 0$  [free spin model] or  $|J_s| = |J_a| = |h|$  [a special case of the transverse field Ising model]. Therefore, condition (8) does not hold, whereas the ordinary weak ETH [5, 6, 10] is satisfied in this model.

For the  $k \neq 0$  components, Eq. (5) is satisfied as

$$\begin{aligned} \chi_\infty^T(k) &= \chi_\infty^S(k) = \chi_\infty^{\text{qch}}(k) \\ &= \frac{1}{2\pi} \int_0^{2\pi} dk' \frac{\varepsilon_{k'} \varepsilon_{k'+k} - (J_s \cos k' + h)(J_s \cos(k' + k) + h) + J_a^2 \sin k' \sin(k' + k)}{2\varepsilon_{k'} \varepsilon_{k'+k}} \\ &\quad \times \frac{\tanh \beta \varepsilon_{k'} + \tanh \beta \varepsilon_{k'+k}}{\varepsilon_{k'} + \varepsilon_{k'+k}} \\ &+ \frac{1}{2\pi} \int_0^{2\pi} dk' \frac{\sinh \beta(\varepsilon_{k'} - \varepsilon_{k'+k})}{\varepsilon_{k'} - \varepsilon_{k'+k}} \frac{1}{\cosh \beta \varepsilon_{k'} \cosh \beta \varepsilon_{k'+k}} \\ &\quad \times \frac{\varepsilon_{k'} \varepsilon_{k'+k} + (J_s \cos k' + h)(J_s \cos(k' + k) + h) - J_a^2 \sin k' \sin(k' + k)}{2\varepsilon_{k'} \varepsilon_{k'+k}}. \end{aligned} \quad (\text{S33})$$

This indicates condition (S23) is satisfied in this model. From Eqs. (S31) and (S33), Eq. (6) holds and  $\chi_\infty^T(k)$  is uniformly continuous in  $k$ , while  $\chi_\infty^{\text{qch}}(k)$  is discontinuous at  $k = 0$ . Moreover, for  $k \neq 0$ ,  $\chi_N^T(k) - \chi_N^{\text{qch}}(k)$  scales as

$$\chi_N^T(k) - \chi_N^{\text{qch}}(k) = \frac{\beta}{2N} \left( \frac{1}{\cosh^2 \beta \varepsilon_{k/2}} + \frac{1}{\cosh^2 \beta \varepsilon_{\pi-k/2}} \right) + \exp(-\Theta(N)) = \Theta(1/N), \quad (\text{S34})$$

because some off-diagonal elements  $|\langle \nu' | \hat{\sigma}_j^z | \nu \rangle|$  that are appeared in Eq. (S23) scale as  $\Theta(1/N)$ . That indicates the ordinary off-diagonal ETH [11–14], which requires exponentially fast decay of all off-diagonal elements, is not satisfied in this model.

### G. Additional demonstrations of (ii)-(iv)

Although condition (S23) is weaker than the ordinary off-diagonal ETH [11–14] as mentioned above, there are some models which do not satisfy it such as the longitudinal field Ising model ( $J_x = J_y = 0$ ). Fig. S3 (a) shows  $k$  dependence of  $\chi_N^{\text{qch}}(k)$ ,  $\chi_N^T(k)$ , and  $\chi_N^S(k)$  in this model. Since  $\hat{m}_k$  is conserved,  $\chi_N^{\text{qch}}(k) = 0$  holds, while  $\chi_N^T(k)$  and  $\chi_N^S(k) > 0$  for all  $k$ , resulting in the violation of Eq. (5) or equivalent condition (S23). In contrast, Fig. S3 (b) shows how the susceptibilities behave when a small nonintegrability ( $J_x + J_y = 0.1$ ,  $J_x - J_y = 0.2$ ) is added to this system. For the  $k = 0$  component,  $\chi_N^{\text{qch}}(0)$  does not approach  $\chi_N^S(0)$  as  $N$  is increased, and Eq. (4) is not satisfied as in (a). On the other hand, the  $k \neq 0$  components  $\chi_N^{\text{qch}}(k)$  differs dramatically from those of (a), and Eq. (5) seems satisfied in (b).

These results suggest that Eq. (5) is easily satisfied as in (b), while we need more nonintegrability for Eq. (4). Reflecting these facts,  $\chi_\infty^{\text{qch}}(k)$  is discontinuous at  $k = 0$  in only (b), while  $\chi_\infty^T(k)$  is uniformly continuous in both (a) and (b).




FIG. S3.  $k$  dependence of  $\chi_N^{\text{qch}}(k)$ ,  $\chi_N^T(k)$ , and  $\chi_N^S(k)$  in (a) longitudinal field Ising model ( $J_x = J_y = 0$ ) and (b) XYZ model with small  $J_x$  and  $J_y$  ( $J_x + J_y = 0.1$ ,  $J_x - J_y = 0.2$ ).  $J_z = 1.0$ ,  $h = 0.8$ , and  $\beta = 0.15$  are fixed. We take  $N = 12-14$  and  $k = 2\pi n_k/N$  with  $n_k \in \mathbb{Z}$ .

### H. Relation to Kubo formula

The susceptibility obtained by Kubo formula [15, 16] is given as

$$\chi_N^{\text{Kubo}}(\mathbf{k}, \omega + i\varepsilon) = \int_0^\infty dt e^{i\omega t - \varepsilon t} \frac{N}{i} \langle [\hat{m}_{\mathbf{k}}(t), -\hat{m}_{\mathbf{k}}^\dagger] \rangle_{\text{ini}} \quad (\text{S35})$$

$$= \chi_N^T(\mathbf{k}) + (i\omega - \varepsilon) \int_0^\infty dt e^{i\omega t - \varepsilon t} \beta N \langle \delta \hat{m}_{\mathbf{k}}; \delta \hat{m}_{\mathbf{k}}(t) \rangle_{\text{ini}}, \quad (\text{S36})$$

where  $[\hat{X}, \hat{Y}] = \hat{X}\hat{Y} - \hat{Y}\hat{X}$  is the commutator,  $\omega$  is the angular frequency, and  $\varepsilon$  is a small positive number.

From Eqs. (S36) and (S3), the following holds for all  $N$  and for all  $\mathbf{k}$ ,

$$\lim_{\varepsilon \rightarrow +0} \chi_N^{\text{Kubo}}(\mathbf{k}, 0 + i\varepsilon) = \chi_N^T(\mathbf{k}) - \beta N \langle \delta \hat{m}_{\mathbf{k}}^0; \delta \hat{m}_{\mathbf{k}}^0 \rangle_{\text{ini}} = \chi_N^{\text{qch}}(\mathbf{k}). \quad (\text{S37})$$

- 
- [1] S. Trotzky, Y. A. Chen, A. Flesch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, Probing the relaxation towards equilibrium in an isolated strongly correlated one-dimensional Bose gas, *Nat. Phys.* **8**, 325 (2012).
  - [2] R. M. Wilcox, Bounds for the Isothermal, Adiabatic, and Isolated Static Susceptibility Tensors, *Phys. Rev.* **174**, 624 (1968).
  - [3] P. Mazur, Non-ergodicity of phase functions in certain systems, *Physica* **43**, 533 (1969).
  - [4] M. Suzuki, Ergodicity, constants of motion, and bounds for susceptibilities, *Physica* **51**, 277 (1971).
  - [5] G. Biroli, C. Kollath, and A. M. Läuchli, Effect of Rare Fluctuations on the Thermalization of Isolated Quantum Systems, *Phys. Rev. Lett.* **105**, 250401 (2010).
  - [6] E. Iyoda, K. Kaneko, and T. Sagawa, Fluctuation Theorem for Many-Body Pure Quantum States, *Phys. Rev. Lett.* **119**, 100601 (2017).
  - [7] T. Mori, Weak eigenstate thermalization with large deviation bound, [arXiv:1609.09776](https://arxiv.org/abs/1609.09776).
  - [8] W. Beugeling, R. Moessner, and M. Haque, Finite-size scaling of eigenstate thermalization, *Phys. Rev. E* **89**, 042112 (2014).
  - [9] R. Steinigeweg, A. Khodja, H. Niemeyer, C. Gogolin, and J. Gemmer, Pushing the Limits of the Eigenstate Thermalization Hypothesis towards Mesoscopic Quantum Systems, *Phys. Rev. Lett.* **112**, 130403 (2014).
  - [10] T. Kuwahara and K. Saito, Ensemble equivalence and eigenstate thermalization from clustering of correlation, [arXiv:1905.01886](https://arxiv.org/abs/1905.01886).
  - [11] M. Srednicki, The approach to thermal equilibrium in quantized chaotic systems, *J. Phys. A* **32**, 1163 (1999).
  - [12] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, *Adv. Phys.* **65**, 239 (2016).
  - [13] T. Mori, T. N. Ikeda, E. Kaminishi, and M. Ueda, Thermalization and prethermalization in isolated quantum systems: a theoretical overview, *J. Phys. B* **51**, 112001 (2018).
  - [14] F. Anza, C. Gogolin, and M. Huber, Eigenstate Thermalization for Degenerate Observables, *Phys. Rev. Lett.* **120**, 150603 (2018).
  - [15] R. Kubo, Statistical-Mechanical Theory of Irreversible Processes. I. General Theory and Simple Applications to Magnetic and Conduction Problems, *J. Phys. Soc. Jpn.* **12**, 570 (1957).
  - [16] R. Kubo, M. Toda, and N. Hashitsume, *Statistical Physics II*, 2nd ed. (Springer, Berlin, 1991).