

CLASSIFICATION OF COSILTING MODULES IN TYPE \tilde{A}

KARIN BAUR AND ROSANNA LAKING

ABSTRACT. Torsion pairs in the category of finitely presented modules over a noetherian ring can be parametrised by the class of cosilting modules. In this paper, we characterise such modules in terms of their indecomposable summands, providing a new approach to the classification of torsion pairs. In particular, we classify cosilting modules over cluster-tilted algebras of type \tilde{A} . We do this by using a geometric model for finite- and infinite-dimensional modules over such algebras.

Cosilting modules were first introduced by Breaz and Pop ([17]) as a dual notion to silting modules ([3]). These modules are a common generalisation of (large) cotilting modules and support τ^{-1} -tilting modules over finite-dimensional algebras. The next theorem establishes a close relationship between cosilting modules in the module category $\text{Mod}(R)$ and the torsion pairs in the subcategory $\text{mod}(R)$ of noetherian modules when R is left noetherian. It follows immediately from the combination of two known results in the literature. The notation used in the bijection is defined in the notation section at the end of the introduction.

Theorem ([2, Cor. 3.9], [24, Lem. 4.4]). *For a left noetherian ring R , the assignment sending a cosilting module C to the torsion pair $({}^{\perp_0}C \cap \text{mod}(R), \text{Cogen}(C) \cap \text{mod}(R))$ induces a bijection between the following sets.*

- (1) *The set of equivalence classes of cosilting modules C in $\text{Mod}(R)$.*
- (2) *The set of torsion pairs $(\mathcal{X}, \mathcal{Y})$ in $\text{mod}(R)$.*

Torsion pairs play a fundamental role in localisation theory (see, for example, [15]) and also provide a direct connection between the module category and t-structures in the derived category ([33]). Tilting theory has played a major role in the study of torsion pairs in module categories, but this often only yields a restricted class of torsion pairs. For example, under the above bijection, the cotilting modules correspond to the torsion pairs $(\mathcal{X}, \mathcal{Y})$ such that \mathcal{Y} generates $\text{mod}(R)$ ([18, Thm. A]) and, if R is a finite-dimensional algebra, then the support τ^{-1} -tilting modules correspond to the torsion pairs $(\mathcal{X}, \mathcal{Y})$ such that \mathcal{Y} is functorially finite ([1, Thm. 2.15]). More recently the lattice of all torsion pairs in $\text{mod}(R)$ has been investigated in the case where R is a finite-dimensional algebra via the study of bricks in $\text{mod}(R)$ ([26]).

The above theorem suggests a different approach to the classification of torsion pairs in $\text{mod}(R)$: classify equivalence classes of cosilting modules in $\text{Mod}(R)$. With a view to following this approach, we show in Theorem 1.13 that equivalence classes of cotilting modules over left artinian rings are parametrised by maximal sets of pairwise Ext-orthogonal indecomposable pure-injective modules of injective dimension less than or equal to 1. This yields Corollary 1.17, which is a new characterisation of cosilting modules in the same setting.

From this starting point we consider the case of cosilting modules over cluster-tilted algebras of type \tilde{A} . This family of algebras was introduced by Buan, Marsh and Reiten ([19]) in the context of cluster theory and has been extensively studied since. Such algebras can be characterised as surface algebras over the annulus ([6]) and, in particular, all indecomposable

2010 *Mathematics Subject Classification.* 16G10, 16G20, 16S90, 16E05.

Key words and phrases. cosilting, pure-injective, cluster-tilted, torsion pair, geometric model, gentle algebra.

The first author was supported by FWF grants P30549-N26 and DK1230. She is supported by a Royal Society Wolfson Research Merit Award. Currently, she is on leave from the University of Graz. The second author was supported by the Max Planck Institute for Mathematics and also by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement No. 797281.

finite-dimensional modules are known to be string or band modules. Moreover, it is possible to express all finite-dimensional string modules as arcs between marked points on the surface.


In the same way, we express the infinite-dimensional (indecomposable pure-injective) string modules as *asymptotic arcs* in the surface, starting at a marked point and converging towards the unique non-contractible closed curve (see [8, 9]). The remaining infinite-dimensional indecomposable pure-injective modules, which were classified in [47], consist of band modules parametrised by $K^* \times \{\infty, -\infty\}$, as well as the unique generic module. We represent this family of band modules, together with the finite-dimensional band modules, by the non-contractible closed curve in the surface. Using this geometric model of the pure-injective modules, we establish the following classification of the cosilting modules up to equivalence. We use the term *arc* to refer to either a finite or asymptotic arc up to homotopy.

Main Theorem (Theorem 2.36 and Remark 2.23). *Let $A(\Gamma)$ be a cluster-tilted K -algebra of type \tilde{A} where Γ is a triangulation of the annulus S with marked points M . There are the following bijective correspondences:*

$$\begin{aligned} \mathcal{A}(S, M) \times \mathcal{P}(K^*) &\xleftrightarrow{1-1} \text{Cosilt-}A(\Gamma) \\ \mathcal{T}(S, M) &\xleftrightarrow{1-1} \text{cosilt-}A(\Gamma) \end{aligned}$$

Where $\mathcal{A}(S, M)$ is the set of maximal collections of non-crossing arcs in (S, M) that contain at least one asymptotic arc; $\mathcal{P}(K^*)$ is the powerset of K^* ; $\mathcal{T}(S, M)$ is the set of maximal collections of non-crossing arcs in (S, M) that contain no asymptotic arcs; $\text{Cosilt-}A(\Gamma)$ is the set of infinite-dimensional cosilting $A(\Gamma)$ -modules up to equivalence; and $\text{cosilt-}A(\Gamma)$ is the set of finite-dimensional cosilting $A(\Gamma)$ -modules up to equivalence.

The diagram below depicts an element of $\mathcal{A}(S, M)$ where (S, M) is as in Figure 2. By the Main Theorem, each subset of K^* determines an infinite-dimensional cosilting module with the indecomposable pure-injective modules corresponding to the arcs $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5$ occurring as direct summands.



This classification encompasses some known results, including the classification of cotilting modules over the path algebra of a quiver of type \tilde{A} ([18]) and the classification of torsion pairs in a tube category in terms of asymptotic arcs ([8]); see Remark 2.37. We also note that there is a significant body of existing work on the classification of torsion pairs in cluster categories via geometric models [32, 44, 34, 35, 36, 55].

We end this introduction with a summary of the contents of the article. The first section is dedicated to giving a characterisation of cosilting modules over a left artinian ring R in terms of certain classes of indecomposable pure-injective modules (Corollary 1.17). To do this, we first introduce the fundamental notions from the theory of purity (Section 1.1) and then provide a characterisation of cotilting R -modules (Section 1.2). Section 2 of the paper focuses on cluster-tilted algebras of type \tilde{A} . In Section 2.1, we introduce the algebras and the geometric model. In Section 2.2 we provide several preliminary results that will lead us to the classification. In particular, we describe the indecomposable pure-injective modules (Section 2.2.1); determine their injective dimensions (Section 2.2.2); and describe extensions between them (Section 2.2.3). The main result of Section 2.2.3 is Theorem 2.16, which establishes a correspondence between certain crossings of arcs in the surface and extensions between the corresponding modules. A

large portion of the proof of the theorem makes use of the combinatorial description of the derived category of $A(\Gamma)$; we postpone these arguments until Appendix A. The final section (Section 2.3) consists of the proof of Theorem 2.36.

Notation. Throughout R denotes an associative unital ring. The category of left R -modules is denoted by $\text{Mod}(R)$ and the full subcategory of finitely presented left R -modules is denoted by $\text{mod}(R)$.

Let N be a left R -module and I a set. We use the notation N^I to denote the direct product of copies of N indexed by I . We use the notation $N^{(I)}$ to denote the direct sum of copies of N indexed by I .

For a set \mathcal{N} of modules we use the notation $\text{Prod}(\mathcal{N})$ to denote the set of direct summands of direct products of copies of objects in \mathcal{N} . If $\mathcal{N} = \{N\}$ then we will write $\text{Prod}(N)$ for $\text{Prod}(\mathcal{N})$.

We denote the set of submodules of modules in $\text{Prod}(\mathcal{N})$ (respectively in $\text{Prod}(N)$) by $\text{Cogen}(\mathcal{N})$ (respectively $\text{Cogen}(N)$) for a set \mathcal{N} of modules or a module N . For a module N , define the sets ${}^{\perp_1}N := \{M \in \text{Mod}(R) \mid \text{Ext}_R^1(M, N) = 0\}$ and ${}^{\perp_0}N := \{M \in \text{Mod}(R) \mid \text{Hom}_R(M, N) = 0\}$.

1. COSILTING MODULES OVER ARTINIAN RINGS.

1.1. Background on purity in module categories. In this section we briefly define some key concepts from the theory of purity in module categories. Consider the category $\mathcal{F}(R)$ of additive functors from the category $\text{mod}(R^{\text{op}})$ of finitely presented right R -modules to the category Ab of abelian groups. There exists a fully faithful functor $t: \text{Mod}(R) \rightarrow \mathcal{F}(R)$ where, for each left R -modules M , we define the functor $t(M): \text{mod}(R^{\text{op}}) \rightarrow \text{Ab}$ which takes $N \mapsto N \otimes_R M$ and $f \mapsto f \otimes_R 1_M$.

Definition 1.1. An exact sequence $0 \rightarrow X \xrightarrow{f} Y \xrightarrow{g} Z \rightarrow 0$ in $\text{Mod}(R)$ is called **pure-exact** if its image $0 \rightarrow t(X) \rightarrow t(Y) \rightarrow t(Z) \rightarrow 0$ is an exact sequence in $\mathcal{F}(R)$. In this case, the morphism f is called a **pure monomorphism** and g is called a **pure epimorphism**. A module N is called **pure-injective** if every pure monomorphism $N \rightarrow Y$ splits. Let $\text{Pinj}(R)$ denote the full subcategory of pure-injective modules.

It turns out that the functor $t: \text{Mod}(R) \rightarrow \mathcal{F}(R)$ restricts to an equivalence of categories $t: \text{Pinj}(R) \xrightarrow{\sim} \text{Inj}(\mathcal{F}(R))$ where $\text{Inj}(\mathcal{F}(R))$ is the full subcategory of injective objects in $\mathcal{F}(R)$ (see, for example, [40, Thm. B.16]).

Definition 1.2. We say that a module N in $\text{Pinj}(R)$ has **no superdecomposable part** if every nonzero direct summand of N has a nonzero indecomposable direct summand.

If N has no superdecomposable part, then there exists a set \mathcal{N} of indecomposable pure-injective modules such that $N \cong \text{PE}(\bigoplus_{M \in \mathcal{N}} M)$ where $\text{PE}(-)$ denotes the **pure-injective envelope** of $\bigoplus_{M \in \mathcal{N}} M$ in $\text{Mod}(R)$ i.e. there exists a pure-monomorphism $\bigoplus_{M \in \mathcal{N}} M \rightarrow N$ such that $t(\bigoplus_{M \in \mathcal{N}} M) \rightarrow t(N)$ is an injective envelope of $t(\bigoplus_{M \in \mathcal{N}} M)$ in $\mathcal{F}(R)$ (see [28], cf. [45, Thm. 4.A14] for more details).

Remark 1.3. If

$$\text{PE}\left(\bigoplus_{M \in \mathcal{N}} M\right) \cong \text{PE}\left(\bigoplus_{L \in \mathcal{L}} L\right)$$

for sets \mathcal{L} and \mathcal{N} of indecomposable pure-injective modules, then there is a bijection $\sigma: \mathcal{N} \rightarrow \mathcal{L}$ such that $M \cong \sigma(M)$ for every M in \mathcal{N} (again, see [45, Thm. 4.A14]). It follows that (up to reordering and isomorphism) \mathcal{N} is the set of indecomposable direct summands of N . For a pure-injective module N with no superdecomposable part, we will use the notation \mathcal{N}_N to denote this set. Note that the elements of \mathcal{N}_N are necessarily pure-injective.

Definition 1.4. An additive functor from $\text{Mod}(R)$ to Ab is called a **coherent functor** if there exists a morphism $f: M \rightarrow N$ in $\text{mod}(R)$ such that

$$\text{Hom}_R(N, L) \xrightarrow{\text{Hom}_R(f, L)} \text{Hom}_R(M, L) \rightarrow F(L) \rightarrow 0$$

is an exact sequence of abelian groups for every $L \in \text{Mod}(R)$.

Definition 1.5. A full subcategory \mathcal{D} of $\text{Mod}(R)$ is **definable** if there exists a set $\{F_i\}_{i \in I}$ of coherent functors such that

$$\mathcal{D} = \{M \in \text{Mod}(R) \mid F_i(M) = 0 \text{ for all } i \in I\}.$$

In [25], Crawley-Boevey characterises definable subcategories of $\text{Mod}(R)$ in terms of closure conditions. He shows that a full subcategory \mathcal{D} of $\text{Mod}(R)$ is definable if and only if \mathcal{D} is closed under direct limits, direct products and pure submodules i.e. if $X \rightarrow Y$ is a pure monomorphism with $Y \in \mathcal{D}$, then $X \in \mathcal{D}$.

1.2. Cotilting modules over artinian rings. In this section we characterise cotilting modules over artinian rings in terms of their indecomposable direct summands. First we recall the definition of a cotilting module in $\text{Mod}(R)$. Note that we are only considering cotilting modules of injective dimension ≤ 1 (so-called 1-cotilting modules).

Definition 1.6. A module C in $\text{Mod}(R)$ is called a **cotilting module** if $\text{Cogen}(C) = {}^{\perp 1}C$. If C_1 and C_2 are cotilting modules, then we say that C_1 is **equivalent** to C_2 if $\text{Prod}(C_1) = \text{Prod}(C_2)$.

We will make use of the following properties of cotilting modules.

Theorem 1.7 ([13, Thm. 2.8, Prop. 3.2]). *Every cotilting module over R is a pure-injective module and the full subcategory $\text{Cogen}(C) = {}^{\perp 1}C$ is definable.*

Next we prove a structure theorem for cotilting modules over an artinian ring. We require the following definition that can be found in [51, Sec. 3.4] (referred to as a 1-rigid system). For tame hereditary artin algebras, similar ideas can be found in [18].

Definition 1.8. Let \mathcal{N} be a non-empty set of indecomposable pure-injective modules. Then \mathcal{N} is called a **rigid system** if the following conditions are satisfied.

- (1) $\text{id}_R(M) \leq 1$ for all $M \in \mathcal{N}$.
- (2) $\text{Ext}_R^1(M, N) = 0$ for all $M, N \in \mathcal{N}$.

We refer to a rigid system \mathcal{N} as **maximal** if the following property is satisfied: if there exists a rigid system \mathcal{L} such that $\text{Prod}(\mathcal{N}) \subseteq \text{Prod}(\mathcal{L})$, then $\text{Prod}(\mathcal{N}) = \text{Prod}(\mathcal{L})$. Two maximal rigid systems \mathcal{N} and \mathcal{L} are **equivalent** if $\text{Prod}(\mathcal{N}) = \text{Prod}(\mathcal{L})$.

We will also need the following two lemmas. The first is implicit in the proof of [51, Thm. 3.7] and the second can be found in [18, Cor. 2.3]. We include the proofs for completeness. See Remark 1.3 for the definition of \mathcal{N}_N .

Lemma 1.9. *Let R be a left noetherian ring. For a pure-injective module N with no superdecomposable part, the following statements are equivalent.*

- (1) $\text{id}_R(N) \leq 1$.
- (2) $\text{id}_R(M) \leq 1$ for all $M \in \mathcal{N}_N$.

Proof. Since R is left noetherian, we may apply Baer's criterion and conclude that the class of modules of injective dimension ≤ 1 is a definable subcategory since the functor $\text{Ext}_R^1(R/I, -)$ is a coherent functor for every ideal I . Then, since definable subcategories are closed under direct summands, direct sums and pure-injective envelopes, the statement follows from the structure of N described in Section 1.1. \square

Lemma 1.10. *Let R be a left artinian ring. For a pure-injective module N with no superdecomposable part and $\text{id}_R(N) \leq 1$, the following statements are equivalent.*

- (1) $\text{Ext}_R^1(N, N) = 0$.

- (2) $\text{Ext}_R^1(N^I, N) = 0$ for all sets I .
(3) $\text{Ext}_R^1(M', M) = 0$ for all $M', M \in \mathcal{N}_N$.

Proof. The argument for **(2)** \Rightarrow **(3)** is straight forward. To show **(1)** \Rightarrow **(2)** we apply [18, Cor. 1.10], which tells us that ${}^{\perp_1}N$ is closed under products.

Finally we prove **(3)** \Rightarrow **(1)**. Since N has no superdecomposable part, we may write $N \cong \text{PE}\left(\bigoplus_{M \in \mathcal{N}_N} M\right)$. Moreover, N is a direct summand of $K := \prod_{M \in \mathcal{N}_N} M$ and it therefore suffices to show that $\text{Ext}_R^1(K, K) = 0$. By [18, Cor. 1.10] and Lemma 1.9, we have that ${}^{\perp_1}M$ is closed under products for each $M \in \mathcal{N}_N$. Applying our assumption, we therefore have that $\text{Ext}_R^1(K, M) = 0$. But then we have that $\text{Ext}_R^1(K, K) \cong \prod_{M \in \mathcal{N}_N} \text{Ext}_R^1(K, M) = 0$. \square

Remark 1.11. Let \mathcal{N} be a set of indecomposable pure-injective modules and consider the modules $N := \text{PE}\left(\bigoplus_{M \in \mathcal{N}} M\right)$ and $L := \prod_{M \in \mathcal{N}} M$. Then N is a direct summand of L and so we have the following containments $\text{Prod}(L) = \text{Prod}(\mathcal{N}) \subseteq \text{Prod}(N) \subseteq \text{Prod}(L)$. That is, we have $\text{Prod}(L) = \text{Prod}(\mathcal{N}) = \text{Prod}(N)$.

In Theorem 1.13 below, we prove that cotilting modulevs over artinian rings are determined by their indecomposable summands. A slightly weaker version of this theorem can be found in [51, Thm. 3.7]. Trlifaj produces a similar correspondence between cotilting *classes* and (not necessarily maximal) rigid systems. Our stronger version requires the following lemma, which follows immediately from [43, Thm. 3.3]. A cotilting module is of **cofinite type** if there is a set \mathcal{S} of finitely generated right R -modules such that

$$\text{Cogen}(C) = \{X \in \text{Mod}(R) \mid \text{Tor}_1^R(S, X) = 0 \text{ for all } S \in \mathcal{S}\}.$$

Lemma 1.12. *Let R be a ring and let C be a cotilting module of cofinite type. Then there is a direct summand D of C that has no superdecomposable part and is cotilting. In particular, if R is left noetherian, then every cotilting module is equivalent to one with no superdecomposable part.*

Proof. Let C be a cotilting module of cofinite type. By [31, Thm. 15.18], there exists a tilting right R -module such that $C \cong \text{Hom}_{\mathbb{Z}}(T, \mathbb{Q}/\mathbb{Z})$. It follows from [43, Thm. 3.3] that there exists a set of \mathcal{N} of indecomposable direct summands of C such that $\text{Prod}(C) = \text{Prod}(\mathcal{N})$. Then, by Remark 1.11, we have that $\text{PE}\left(\bigoplus_{N \in \mathcal{N}} N\right)$ is a cotilting module equivalent to C . The last statement follows from [4, Thm. 2.10]. \square

Theorem 1.13. *Let R be a left artinian ring. The assignments*

$$C \mapsto \mathcal{N}_C \quad \text{and} \quad \mathcal{N} \mapsto \prod_{M \in \mathcal{N}} M$$

are mutually inverse bijections between the set of cotilting modules C in $\text{Mod}(R)$ (up to equivalence) and the set of maximal rigid systems \mathcal{N} in $\text{Mod}(R)$ (up to equivalence).

Proof. Let C be a cotilting module. By Lemma 1.12, we may assume that C has no superdecomposable part. By Lemma 1.10 and Lemma 1.9 we have that \mathcal{N}_C is a rigid system so it remains to show that \mathcal{N}_C is maximal. Suppose \mathcal{N} is a rigid system with $\text{Prod}(\mathcal{N}_C) \subseteq \text{Prod}(\mathcal{N})$. Then let $N := \text{PE}\left(\bigoplus_{M \in \mathcal{N}} M\right)$ and note that, since $\bigoplus_{M \in \mathcal{N}_C} M$ is a direct summand of $\bigoplus_{M \in \mathcal{N}} M$, it follows that C is a direct summand of N . We may then argue as in the proof of (1) \Rightarrow (2) in [18, Prop. 3.1] to obtain that $\text{Prod}(C) = \text{Prod}(N)$ and hence that $\text{Prod}(\mathcal{N}_C) = \text{Prod}(\mathcal{N})$ by Remark 1.11.

Let \mathcal{N} be a maximal rigid system and consider $C := \text{PE}\left(\bigoplus_{M \in \mathcal{N}} M\right)$. Then $\mathcal{N} = \mathcal{N}_C$ by Remark 1.3 and so we have that $\text{Ext}_R^1(C, C) = 0$ and $\text{id}_R(C) \leq 1$ by Lemma 1.10 and Lemma 1.9. By [18, Cor. 1.12], there exists a pure-injective module N such that $C \oplus N$ is cotilting. Note that $\mathcal{N}_{C \oplus N} = \mathcal{N}_C \cup \mathcal{N}_N = \mathcal{N} \cup \mathcal{N}_N$ and so, by applying Lemma 1.12, we can ensure that N has no superdecomposable part. As $\mathcal{N} \cup \mathcal{N}_N$ is a rigid system containing \mathcal{N} , we have

$\text{Prod}(\prod_{M \in \mathcal{N}} M) = \text{Prod}(C) = \text{Prod}(\mathcal{N}) = \text{Prod}(\mathcal{N} \cup \mathcal{N}_N) = \text{Prod}(N \oplus C)$ by the maximality of \mathcal{N} and Remark 1.11. This implies that

$$\text{Cogen}(\prod_{M \in \mathcal{N}} M) = \text{Cogen}(N \oplus C) = {}^{\perp_1}(N \oplus C) = {}^{\perp_1}\left(\prod_{M \in \mathcal{N}} M\right)$$

and so $\prod_{M \in \mathcal{N}} M$ is a cotilting module.

The fact that the assignments are mutually inverse bijections follows immediately from Remark 1.11. \square

1.3. Cosilting modules over artinian rings. Cosilting modules over a ring R , introduced by Breaz and Pop [17], are a generalisation of cotilting modules over R . In fact, the cotilting modules are exactly the faithful cosilting modules.

Definition 1.14. A module C over R is a **cosilting module** if there exists an injective copresentation $0 \rightarrow C \rightarrow E_0 \xrightarrow{\sigma} E_1$ such that $\text{Cogen}(C) = \{M \in \text{Mod}(R) \mid \text{Hom}_R(M, \sigma) \text{ is surjective}\}$. Two cosilting modules C_1 and C_2 are said to be **equivalent** if $\text{Prod}(C_1) = \text{Prod}(C_2)$.

In the later sections we will make use of the following useful characterisation of cosilting modules. For this precise formulation, see [2, Thm. 3.6].

Proposition 1.15 ([56]). *Let R be a ring. An R -module C is a cosilting module if and only if C satisfies the following two conditions:*

- (1) C is a cotilting module over $R/\text{Ann}(C)$ where $\text{Ann}(C) := \{r \in R \mid r.C = 0\}$.
- (2) $\text{Cogen}(C)$ is a torsion-free class in $\text{Mod}(R)$.

Combining this characterisation with Theorem 1.7, we obtain the following corollary.

Corollary 1.16 ([56], [17]). *Every cosilting module C over R is a pure-injective module and $\text{Cogen}(C)$ is a definable subcategory.*

If R is a left artinian ring, then R/I is left artinian for any ideal I in R . Thus we may combine the results of Section 1.2 with Proposition 1.15 to obtain the following characterisation of cosilting modules over an artinian ring.

Corollary 1.17. *Let R be a left artinian ring. Then a pure-injective module C is a cosilting module if and only if C satisfies the following conditions:*

- (1) *There exists a maximal rigid system \mathcal{N} in $\text{Mod}(R/\text{Ann}(C))$ such that $\text{Prod}(C) = \text{Prod}(\mathcal{N})$.*
- (2) $\text{Cogen}(C)$ is a torsion-free class in $\text{Mod}(R)$.

2. COSILTING MODULES OVER CLUSTER-TILTED ALGEBRAS.

Throughout this section we fix an algebraically closed field K . This assumption is needed in order to appeal to the classification of indecomposable pure-injective modules over a domestic string algebra used in Theorem 2.11.

2.1. Cluster-tilted algebras of type \tilde{A} . Cluster-tilted algebras were introduced by Buan, Marsh and Reiten ([19]) as endomorphism algebras of cluster-tilting objects in cluster categories: If \mathcal{C}_Q is the cluster category of an acyclic quiver Q and T a cluster-tilting object in \mathcal{C}_Q , the endomorphism ring $\text{End}_{\mathcal{C}_Q}(T)$ is a **cluster-tilted algebra** of type Q . We will use the characterisation of cluster-tilted algebras through surface triangulations: Assem et al. show in [6, Theorem 1.1] that the cluster-tilted algebras of type \tilde{A} are the gentle algebras arising from triangulations of an annulus.

Let S be an oriented surface and M a nonempty finite set of points in the boundary of S such that each boundary component has at least one marked point. An **arc** in S is a non self-intersecting curve $\gamma: [0, 1] \rightarrow S$ with endpoints in M whose interior is disjoint from the boundary of S and which does not cut out a monogon. The arc is a **boundary arc** if it cuts out a digon, it is **internal** otherwise. Arcs are considered up to homotopy fixing endpoints. A




FIGURE 1. Annulus with universal cover $U_{3,2}$




FIGURE 2. Triangulation in the cover, in the annulus and its quiver

triangulation of a S (and M) is a maximal collection of pairwise non-crossing internal arcs. For details we refer to [29]. It is known that in the case of an annulus, the number of internal arcs of any triangulation is equal to the number of marked points on the two boundary components, [29, Proposition 2.10]. Let Γ be a triangulation of an annulus. Then its quiver $Q = Q(\Gamma)$ has as vertices the set $Q_0 := \{1, 2, \dots, m\}$ indexed by the set $\{\gamma_1, \dots, \gamma_m\}$ of arcs in Γ . The set of arrows Q_1 is defined to consist of arrows defined as follows: for any two internal arcs γ_i, γ_j in Γ bounding a common triangle we draw an arrow $a: i \rightarrow j$ if j is clockwise from i (using the orientation of S). As usual, the quiver Q comes with assignments $s, t: Q_1 \rightarrow Q_0$ such that $s(a) = i$ and $t(a) = j$.

A triangle Δ in Γ is **internal** if all its edges are internal arcs. Every internal triangle gives rise to an oriented cycle $a_\Delta b_\Delta c_\Delta$ in Q (up to cyclic permutation). Then the algebra $A(\Gamma)$ is defined to be the quotient $KQ/I(\Gamma)$ where $I(\Gamma)$ is the ideal generated by the paths $a_\Delta b_\Delta, b_\Delta c_\Delta, c_\Delta a_\Delta$ for all triangles Δ in Γ . This is the (noncompleted) Jacobian algebra of Q with the potential arising from all the cycles of Γ [27, 41].

Since we are working with cluster-tilted algebras of type \tilde{A} , we will from now on we will fix the surface (S, M) to be an annulus with p respectively q marked points on the two boundary components, $p, q > 0$. We will also fix a triangulation Γ and the algebra $A(\Gamma)$. Unless otherwise stated, the notation Q will refer to the quiver $Q(\Gamma)$ i.e. $A(\Gamma) \cong KQ/I(\Gamma)$.

2.1.1. Geometric model. As well as allowing us to define $A(\Gamma)$, the surface (S, M) provides us with a geometric model of the module category. Internal arcs will simply be called arcs. Arcs with both endpoints at the same boundary are called **peripheral** arcs, arcs connecting the two boundaries are called **bridging**. It is customary to draw an annulus and its triangulation on its universal cover $U = U_{p,q}$. This is a horizontal strip in the plane, with a projection map to the annulus. We will label marked points on the boundary components by integers, indicating the upper boundary with a prime. We arrange the marked points so that $0'$ is above 0 in the universal cover. In every copy of the fundamental domain, there are p marked points on the lower boundary and q marked points on the upper boundary. See Figure 1 for an example with $p = 3$ and $q = 2$.

As an example, we consider a triangulation of an annulus with three marked points on the outer boundary and 2 marked points on the inner boundary. See Figure 2. The associated quiver is on the right.

2.2. Towards maximal rigid systems. In Section 2.3 we classify the cosilting $A(\Gamma)$ -modules using Corollary 1.17 i.e. we identify the pure-injective modules C such that the indecomposable summands of C form a maximal rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(C))$ and $\text{Cogen}(C)$ is a torsion-free class in $\text{Mod}(A(\Gamma))$.

Recall that the elements of a maximal rigid system are pairwise Ext-orthogonal indecomposable pure-injective modules that have injective dimension less than or equal to 1. In this section, we set out some preliminary results that will allow us to identify maximal rigid systems in the subsequent section. In particular, in Section 2.2.1 we consider indecomposable pure-injective modules; in Section 2.2.2 we consider conditions for injective dimension less than or equal to 1; and in Section 2.2.3 we consider extensions between indecomposable pure-injective modules.

2.2.1. Indecomposable pure-injective modules. Here we describe indecomposable finite-dimensional modules for $A(\Gamma)$ via arcs in the annulus as introduced in [10], see also [53, 8], where we allow marked points on both boundaries as in [12] and arbitrary orientation of Q .

We will extend this description of $A(\Gamma)$ -modules to include the indecomposable infinite-dimensional pure-injective modules, which were classified in [47].

Finite dimensional string modules. By a **finite arc** in (S, M) we mean an arc α in S whose endpoints $\alpha(0), \alpha(1)$ are contained in M and all other points in α are contained in the interior of S .

Let $\alpha: [0, 1] \rightarrow S$ be a (finite) arc in (S, M) that is not homotopic to any arc in Γ . We assume that α is chosen in its homotopy class so that, if α intersects an arc γ in Γ , then α and γ intersect transversally and the number of intersections is minimal.

We will define a module $M(\alpha)$ for every finite arc α and in order to do this we must first define strings. For every arrow $a \in Q_1$, we introduce a formal inverse a^{-1} and we extend the assignments s and t so that $s(a^{-1}) = t(a)$ and $t(a^{-1}) = s(a)$. If a^{-1} is the inverse of an arrow a , then its formal inverse $(a^{-1})^{-1}$ is defined to be a . An arrow a will be referred to as a **direct letter** and its inverse a^{-1} will be referred to as an **inverse letter**. For each vertex $i \in Q_0$, we introduce a **trivial letter** or **trivial string**, which we denote e_i . We define $e_i^{-1} = e_i$. We will refer to the elements of the set

$$Q_1 \cup \{a^{-1} \mid a \in Q_1\} \cup \{e_i \mid i \in Q_0\}$$

as **letters**. For any sequence of letters $p_n \dots p_1 p_0$ we define the inverse sequence $(p_n \dots p_1 p_0)^{-1}$ to be $p_0^{-1} p_1^{-1} \dots p_n^{-1}$.

Orienting α from $\alpha(0)$ to $\alpha(1)$, suppose γ_{i_0} is the first arc of Γ that intersects α , that γ_{i_1} is the second arc of Γ that intersects α , and so on. In this way we obtain a sequence (i_0, i_1, \dots, i_n) of vertices in Q , which uniquely defines a sequence of letters u_α . This sequence is

$$u_\alpha := \begin{cases} a_n a_{n-1} \dots a_1 & \text{if } n > 0 \\ e_{i_0} & \text{if } n = 0 \end{cases}$$

where $a_j: i_{j-1} \rightarrow i_j$ for all $1 \leq j \leq n$. We will call such a sequence a **finite string** of length n . We also extend the definition of s and t to strings so that $s(u_\alpha) = i_0$ and $t(u_\alpha) = i_n$.

Example 2.1. As an example, consider algebra $A(\Gamma)$ where Γ is the triangulation as in Figure 2. Let α be the arc in Figure 3. Then the string of α is $h^{-1}dgfe$.

Remark 2.2. Strings were first introduced in [20] and used to parametrise the string modules over an arbitrary string algebra. The above definition of string coincides with this original definition by [6, Prop. 4.2].

Let α be an arc in (S, M) with associated string u_α as above. We define the **string module** $M(\alpha)$ as a representation in the following way.

For each $i \in Q_0$, let $\mathcal{I}_i := \{k \mid i_k = i \text{ for } 0 \leq k \leq n\}$ and define

$$M(\alpha)_i := K^{(\mathcal{I}_i)}.$$




FIGURE 3. Finite arc in the triangulated cover and in the triangulated annulus




FIGURE 4. Asymptotic arcs in the triangulated cover and in the triangulated annulus

For each arrow $a: i \rightarrow j$ in Q , let

$$M(\alpha)_a: K^{(\mathcal{I}_i)} \rightarrow K^{(\mathcal{I}_j)}$$

be the matrix with (k, l) th entry ($k \in \mathcal{I}_i, l \in \mathcal{I}_j$) given by 1_K if either $l = k + 1$ and $a_l = a$ or if $k = l + 1$ and $a_k = a^{-1}$. All other entries of $M(\alpha)_a$ are defined to be zero.

Remark 2.3. It follows directly from the definitions that $u_\alpha = u_\delta^{-1}$ if and only if $M(\alpha) \cong M(\delta)$. So for our purposes, it will not matter which orientation we pick for a finite arc.

Infinite-dimensional string modules. In order to define infinite-dimensional string modules, we will consider ‘infinite’ arcs with only one endpoint. A curve $\alpha: [0, 1) \rightarrow S$ is called an **asymptotic arc** in (S, M) if the endpoint $\alpha(0)$ is in M , the interior of α is contained in the interior of S and α spirals (clockwise or anticlockwise) around the inner boundary component of S . For an example, see the asymptotic arc α in Figure 4.

As with the finite arcs, we consider asymptotic arcs up to homotopy and we assume that α is chosen in its homotopy class so that, if α intersects an arc γ in Γ , then α and γ intersect transversally and the number of intersections is minimal.

From α we define a sequence of vertices in Q as we did in the finite case. We obtain an infinite sequence $(i_0, i_1, i_2, \dots, i_n, \dots)$, which uniquely determines a sequence of letters $u_\alpha = \dots a_n \dots a_2 a_1$ such that $a_j: i_{j-1} \rightarrow i_j$ for all $j > 0$. We call such a sequence an **N-string**. We also define $s(u_\alpha)$ to be i_0 .

Example 2.4. As an example, we consider the algebra $A(\Gamma)$ for the triangulated annulus from Figure 2. Let α be the arc in Figure 4, starting at 1 on the lower (outer) boundary. Its string is

$$\dots c^{-1} g f c^{-1} g f e$$

Let α' be the asymptotic arc starting at $0'$ on the upper (inner) boundary. Its string is

$$\dots gfc^{-1}gfc^{-1}$$

Remark 2.5. Such \mathbb{N} -strings were first considered in the context of string algebras by Ringel in [48]. Ringel defines an \mathbb{N} -word to be an infinite sequence of letters $\dots a_3a_2a_1$ such that every truncation $a_n \dots a_2a_1$ is a finite string. In our setting \mathbb{N} -strings coincide with \mathbb{N} -words. Indeed, if we consider $u_\alpha = \dots a_3a_2a_1$, then, for every $n \in \mathbb{N}$, we can construct an arc α_n such that $u_{\alpha_n} = a_n \dots a_2a_1$ as follows. Let δ be the segment of α between $\alpha(0)$ and the point p where α intersects γ_{i_n} . We may then define α_n to be the concatenation of δ with an arc connecting p with the marked point opposite γ_{i_n} .

Let α be an asymptotic arc in (S, M) and u_α the corresponding \mathbb{N} -string. We will define an infinite-dimensional indecomposable pure-injective module $M(\alpha)$. Let i_p be the first bridging arc crossed by α , then the \mathbb{N} -string $\dots a_{p+3}a_{p+2}a_{p+1}$ is the **maximal periodic part** of u_α . Let $n \in \mathbb{N}$ be minimal such that $i_{p+n} = i_p$. If a_{p+n} is direct, then α is said to be **expanding**. If a_{p+n} is inverse, then α is said to be **contracting**.

Example 2.6. We consider the algebra $A(\Gamma)$ for the triangulated annulus from Figure 2. The \mathbb{N} -string of the arc α in Figure 4 is contracting. The \mathbb{N} -string of the arc α' is expanding.

As before we may define a **string module** $M(\alpha)$. For each $i \in Q_0$, let $\mathcal{I}_i := \{k \mid i_k = i \text{ for } k \geq 0\}$. From this data, we define two modules. Firstly let

$$M^\Pi(\alpha)_i := K^{\mathcal{I}_i}$$

and secondly let

$$M^\oplus(\alpha)_i := K^{(\mathcal{I}_i)}.$$

The action of Λ is defined in the same way in both cases. For each arrow $a: i \rightarrow j$ in Q , let

$$M^*(\alpha)_a: M^*(\alpha)_i \rightarrow M^*(\alpha)_j$$

(with $*$ = \prod, \oplus) be the morphism determined by the following components. For $k \in \mathcal{I}_i$ and $l \in \mathcal{I}_j$, take the component map to be 1_K if either $l = k + 1$ and $a_l = a$ or if $k = l + 1$ and $a_k = a^{-1}$. Otherwise take the (k, l) th component of $M^*(\alpha)_a$ to be zero.

If α is expanding, then define $M(\alpha) := M^\Pi(\alpha)$ and if α is contracting, then define $M(\alpha) := M^\oplus(\alpha)$.

Band modules. Let β be the unique simple closed arc lying in the interior of (S, M) (considered up to homotopy). As before, we choose a representative in the homotopy class such that β crosses all arcs $\gamma \in \Gamma$ transversally and the number of crossings between each γ and β is minimal. We orient β in the same way as the boundary: in the universal cover, β is oriented from left to right, in the annulus anticlockwise.

Choose an arc γ_{i_0} in Γ that is crossed by β such that, if γ_{i_1} is the next arc in Γ crossed by β in a clockwise direction and $\gamma_{i_{-1}}$ is the next arc crossed by β in an anticlockwise direction, then the letter $a_0: i_{-1} \rightarrow i_0$ is direct and the letter $a_1: i_0 \rightarrow i_1$ is inverse.¹ Following β in a clockwise direction, let γ_{i_2} be the next arc in Γ crossed by β after γ_{i_1} , let γ_{i_3} be the next arc in Γ crossed by β , and so on. After carrying out a similar procedure in the anticlockwise direction, the arc β uniquely determines a sequence $(\dots i_{-1}, i_0, i_1, \dots)$ of vertices in Q . This sequence then uniquely determines a sequence of letters $u_\beta = \dots a_1a_0a_{-1} \dots$ such that $a_j: i_{j-1} \rightarrow i_j$ for all $j \in \mathbb{Z}$. We will refer to u_β as the **periodic \mathbb{Z} -string** and, for formal reasons, we will say that the **maximal periodic part** of u_β is u_β . Let $s > 0$ be minimal such that $i_s = i_0$. Then we call the sequence $b := a_s \dots a_1$ of letters the **band** corresponding to β .

Remark 2.7. Consider the band $b := a_s \dots a_1$. Let J be the ideal in $A(\Gamma)$ generated by the idempotent elements corresponding to the set V consisting of the peripheral arcs in Γ and let $B := A(\Gamma)/J$. The algebra B is the surface algebra given by the surface obtained by cutting

¹This can always be done, as the bridging arcs in the triangulation change orientation at least twice, i.e. they form both a figure N and a \mathcal{N} where the first and last of the three arcs may coincide.




FIGURE 5. Arc β in the cover and in the annulus

along all of the arcs $\gamma_j \in V$. So B is given by a triangulation of the annulus consisting of only bridging arcs, i.e. $B \cong K\tilde{A}_{s-1}$.

Example 2.8. Consider the algebra $A(\Gamma)$ for the triangulated annulus from Figure 2. The sequence for β as in Figure 5 is $(\dots, 4, 2, 1, 4, 2, 1, 4, \dots)$ with periodic \mathbb{Z} -string $\dots fc^{-1}gfc^{-1}g\dots$. We have $i_0 = 1$, $s = 3$ and $b = c^{-1}gf$.

As before, we define a module $M^\oplus(\beta)$ associated to u_β . For each $i \in Q_0$, let $\mathcal{I}_i := \{k \mid i_k = i \text{ for } k \in \mathbb{Z}\}$ and define $M^\oplus(\beta)_i := K^{\mathcal{I}_i}$. For each arrow $a: i \rightarrow j$ in Q , let $M^\oplus(\beta)_a: M^\oplus(\beta)_i \rightarrow M^\oplus(\beta)_j$ be the morphism defined exactly as in the definition of the infinite-dimensional string modules.

It is clear from the construction of $M^\oplus(\beta)$ that the K -basis vectors of $M^\oplus(\beta)$ are canonically parametrised with the integers: $\{\dots v_{-1}, v_0, v_1, \dots\}$. Let $\Phi: M^\oplus(\beta) \rightarrow M^\oplus(\beta)$ be such that $v_i \mapsto v_{i+s}$ for all $i \in \mathbb{Z}$. It can easily be checked that $\Phi \in \text{End}_A(M^\oplus(\beta))$ and that Φ^{-1} is the endomorphism such that $v_i \mapsto v_{i-s}$ for all $i \in \mathbb{Z}$. This endomorphism equips $M^\oplus(\beta)$ with a $(A(\Gamma), R)$ -bimodule structure where $R := K[\Phi, \Phi^{-1}]$. We consider the functor

$$M^\oplus(\beta) \otimes_R (-): \text{Mod}(R) \rightarrow \text{Mod}(A(\Gamma)).$$

An $A(\Gamma)$ -module of the form $N \cong M^\oplus(\beta) \otimes_R L$ is an indecomposable pure-injective module if L is an indecomposable pure-injective R -module (see, for example, [48]).

Remark 2.9. The following is a complete list of the indecomposable pure-injective modules over R .

- (1) Finitely-generated module $N(\lambda, n)$ i.e. $K[\Phi]/\mathfrak{p}_\lambda^n$ where $\mathfrak{p}_\lambda = (\Phi - \lambda)$, for each $n \geq 1$ and $\lambda \in K^*$.
- (2) Prüfer module $N(\lambda, \infty)$ for each $\lambda \in K^*$ i.e. the direct limit $\varinjlim_{n \geq 1} N(\lambda, n)$.
- (3) \mathfrak{p}_λ -adic completion $N(\lambda, -\infty)$ of $K[\Phi]$ for each $\lambda \in K^*$ i.e. the inverse limit $\varprojlim_{n \geq 1} N(\lambda, n)$.
- (4) Field of fractions $K(\Phi)$.

The above list is obtained from the classification of indecomposable pure-injective modules over a Dedekind domain in [57] by observing that the category of R -modules can be realised as a definable subcategory of the category of $K[\Phi]$ -modules. We also use that, since K is algebraically closed, all the maximal ideals of $K[\Phi]$ are of the form \mathfrak{p}_λ for $\lambda \in K$.

From the list of Remark 2.9, we obtain the following indecomposable pure-injective $A(\Gamma)$ -modules. We will refer to them as **band modules**.

- (1) Finite-dimensional band module $M(\lambda, n) := M^\oplus(\beta) \otimes_R N(\lambda, n)$ for each $\lambda \in K^*$ and $n \in \mathbb{N}$.
- (2) Prüfer module $M(\lambda, \infty) := M^\oplus(\beta) \otimes_R N(\lambda, \infty)$ for each $\lambda \in K^*$.
- (3) Adic module $M(\lambda, -\infty) := M^\oplus(\beta) \otimes_R N(\lambda, -\infty)$ for each $\lambda \in K^*$.
- (4) Generic module $G := M^\oplus(\beta) \otimes_R K(\Phi)$.

Remark 2.10. The definition of the band modules depends on the choice of arc γ_{i_0} in the construction. However, the set of modules defined by different choices of γ_{i_0} are the same up to isomorphism. See [48] for details.

Classification of indecomposable pure-injective $A(\Gamma)$ -modules. The algebras $A(\Gamma)$ are domestic string algebras, therefore we may apply the following classification result.

Theorem 2.11 ([47]). *Let $A(\Gamma)$ be the algebra associated to a triangulated Γ of the annulus (S, M) . Then the following is a complete list of indecomposable pure-injective $A(\Gamma)$ -modules.*

- (1) *The string module $M(\alpha)$ for each α (up to homotopy) such that α is a finite or asymptotic arc in (S, M) that is not homotopic to any arc in Γ .*
- (2) *The band module $M(\lambda, n)$ for every $\lambda \in K^*$ and every $n \in \mathbb{N} \cup \{\infty, -\infty\}$.*
- (3) *The generic module G .*

The term **arc in (S, M)** is used to refer to finite arcs, asymptotic arcs or β . For a string module $M(\alpha)$, we say that α is the **associated arc** and u_α is the **associated string**. For a band module $M(n, \lambda)$ ($n \in \mathbb{N} \cup \{\infty, -\infty\}, \lambda \in K^*$) or G , we say that β is the **associated arc** and u_β is the **associated string**.

We will take the convention that, for any $\gamma \in \Gamma$, the module $M(\gamma)$ is defined to be the zero module.

2.2.2. Injective dimension of indecomposable pure-injective modules. In this subsection we consider the injective dimension of the indecomposable pure-injective modules introduced in the previous subsection.

Injective dimension of string modules. According to Corollary 1.17, we must show that certain modules have injective dimension less than or equal to 1 over a quotient of $A(\Gamma)$. It turns out that this quotient is still a gentle algebra (see Corollary 2.27). In this section we therefore consider the injective dimension of string modules over an arbitrary gentle algebra $A \cong KQ/I$.

In [48] Ringel defines finite strings, infinite strings and bands over any string algebra (and hence any gentle algebra). It is also shown in [48] that in this context infinite- and finite-dimensional string and band modules are always indecomposable pure-injective modules.

Several authors have computed syzygies and hence projective resolutions of finite-dimensional string and band modules (see, for example, [21, 52, 11]). Syzygies of infinite-dimensional string modules have also been considered in [37]. Since we are considering injective dimension rather than projective dimension, we include a full proof of the next lemma. However everything is completely dual to the papers cited here.

Let w be a finite string. We can write $w = p_0^{-1}q_0p_1^{-1}q_1 \dots p_n^{-1}q_n$ where p_i, q_i are paths in (Q, I) for all $0 \leq i \leq n$ and the length of p_i (resp. q_j) is greater than or equal to 1 when $i > 0$ (resp. $j < n$). Similarly we can write any \mathbb{N} -string as $w = \dots q_1^{-1}p_1q_0^{-1}p_0$. Note that this partition into direct and inverse paths is uniquely determined.

Lemma 2.12. *Let $A = KQ/I$ be a gentle algebra. Let $w = q_n^{-1}p_n \dots q_1^{-1}p_1q_0^{-1}p_0$ (respectively $w = \dots q_1^{-1}p_1q_0^{-1}p_0$) be a finite string (respectively an \mathbb{N} -string).*

Then $\text{id}_A(M(w)) \leq 1$ if and only if the following conditions are satisfied.

- (1) *If there is an arrow l_0 such that p_0l_0 is a path in (Q, I) , then there is no arrow a such that $l_0a \in I$.*
- (2) *If w is a finite string and there is an arrow l_n such that q_nl_n is a path in (Q, I) , then there is no arrow a such that $l_na \in I$.*

Proof. We begin by proving the proposition for a finite string w . The injective envelope of $M(w)$ is $\bigoplus_{i=0}^n M(b_i^{-1}q_i^{-1}p_ia_i)$ where a_i is the longest path in Q such that p_ia_i is not contained in I and b_i is the longest path in Q such that q_ib_i is not contained in I . The first cosyzygy of $M(w)$ is isomorphic to $\bigoplus_{i=1}^n M(b_{i-1}^{-1}a_i) \oplus M(a'_0) \oplus M(b'_n)$ where $a_0 = l_0a'_0$ and $b_n = l_nb'_n$ for arrows l_0, l_n if they exist and zero otherwise.

The module $M(a_i^{-1}b_{i-1})$ is injective for all $1 \leq i \leq n$. So $\text{id}_A(M) \leq 1$ if and only if $M(a'_0)$ and $M(b'_n)$ are injective if and only if conditions (1) and (2) are satisfied.

Now consider an \mathbb{N} -string w . If w is expanding then the injective envelope of $M(w)$ is given by $\prod_{i \geq 0} M(b_i^{-1} q_i^{-1} p_i a_i)$ and if w is contracting then the injective envelope of $M(w)$ is given by $\bigoplus_{i \geq 0} M(b_i^{-1} q_i^{-1} p_i a_i)$. In both cases a_i and b_i are defined as in the finite case.

Then, as in the finite case, the first cosyzygy is the direct sum of an injective module and a serial module $M(a'_0)$ where a'_0 is either 0 or $a_0 = l_0 a'_0$ for an arrow l_0 . Finally we observe that $M(a'_0)$ is injective if and only if condition (1) is satisfied as required. \square

Injective dimension of band modules. It suffices for us to consider the injective dimension of band modules over $A(\Gamma)$. Consider the quotient algebra $B := A(\Gamma)/J$ from Remark 2.7 and let $f: A(\Gamma) \rightarrow B$ be the corresponding ring epimorphism.

Remark 2.13. The image of the restriction functor $f^*: \text{Mod}(B) \rightarrow \text{Mod}(A)$ contains all the band modules and is closed under extensions because J is an idempotent ideal (see, for example, [39, Cor. 2.2]). Moreover, it is a bireflective and therefore definable subcategory.


Lemma 2.14. *If N is a band module over $A(\Gamma)$, then $\text{id}_{A(\Gamma)}(N) \leq 1$ and $\text{pd}_{A(\Gamma)}(N) \leq 1$.*

Proof. By [21, Cor. 2.13] finite-dimensional band modules have projective dimension equal to 1. It follows from [30] that the algebra is 1-Iwanga-Gorenstein, and so the finite-dimensional band modules also have injective dimension equal to 1 (see [38, Thm. 5]).

Recall that, by Baer's criterion, the full subcategory with objects in the class \mathcal{F} of modules of injective dimension less than or equal to 1 is definable. Similarly, the full subcategory with objects in the class \mathcal{P} of modules of projective dimension less than or equal to 1 is definable (see, for example, [42, Lem. 5]). Let \mathcal{D} be the image of the restriction functor $f^*: \text{Mod}(B) \rightarrow \text{Mod}(A(\Gamma))$. Thus $\mathcal{F} \cap \mathcal{D}$ (resp. $\mathcal{P} \cap \mathcal{D}$) is a definable subcategory and it contains all the finite-dimensional band modules. By [49, Thm. A], the remaining band modules are in also in $\mathcal{F} \cap \mathcal{D}$ (resp. $\mathcal{P} \cap \mathcal{D}$). \square

2.2.3. Extensions between indecomposable pure-injective modules. In this section, specifically in Theorem 2.16, we prove that there is a very explicit description of the extensions between the indecomposable pure-injective $A(\Gamma)$ -modules in terms of crossings of the corresponding associated arcs.

Definition 2.15. Let N and M be string or band modules for $A(\Gamma)$ and let α, δ be their respective associated arcs in (S, M) . Then we say that α and δ **cross in a 3-cycle** if we have one of the following local configuration with $\gamma_i, \gamma_j, \gamma_k$ bridging arcs in the triangulation.



Similarly, we will say that α and δ **do not cross outside of 3-cycles** if the only crossings between α and δ occur as in the above diagrams.

The left-hand picture in Definition 2.15 may be expressed in terms of the associated strings by saying we have that α and δ cross in a 3-cycle if there exists a 3-cycle $i \xrightarrow{d} j$ (so necessarily

$$\begin{array}{ccc} & d & \\ i & \xrightarrow{\quad} & j \\ & a \swarrow \downarrow c & \\ & & k \end{array}$$

$cd, ac, da \in I(\Gamma)$) such that u_α contains the letter a or a^{-1} and we have that $s(u_\delta) = j$ (assuming without loss of generality that $\delta(0)$ is shown in the diagram). An analogous description can be given for the the right-hand picture.

In Theorem 2.16 we will show that all crossings between arcs that are not in a 3-cycle indicate the existence of extensions between the associated modules.

Theorem 2.16. *Let M and N be string or band modules over $A(\Gamma)$ and let α and δ be their respective associated arcs. If M and N are not both band modules, then $\text{Ext}_{A(\Gamma)}^1(M, N) = 0 = \text{Ext}_{A(\Gamma)}^1(N, M)$ if and only if α and δ do not cross outside of 3-cycles.*

Proof. The case where M and N are both finite-dimensional string modules is covered in [23]. We consider the case where M or N is the generic module; all other cases are addressed in Appendix A since the arguments require the derived category. Throughout the proof we will denote the group $\text{Ext}_{A(\Gamma)}^1(X, Y)$ by $\text{Ext}^1(X, Y)$ for any pair X, Y of $A(\Gamma)$ -modules.

First suppose that M is the generic module and N is a string module. Then the associated arc of M is $\alpha = \beta$. By [49, Prop. 4], there exists an endomorphism ϕ of an adic module $M(\lambda, -\infty)$ such that $M(\lambda, -\infty) \xrightarrow{\phi} M(\lambda, -\infty) \xrightarrow{\phi} \dots$ is a direct system of monomorphisms and the direct limit L is isomorphic to $M^{(J)}$ for some set J (where $M^{(J)}$ denotes the direct sum of copies of M indexed by J). This induces an inverse system

$$\text{Ext}^1(M(\lambda, -\infty), N) \xleftarrow{\text{Ext}^1(\phi, N)} \text{Ext}^1(M(\lambda, -\infty), N) \xleftarrow{\text{Ext}^1(\phi, N)} \dots$$

Denote its inverse limit by L' . It then follows that $\text{Ext}^1(M, N) = 0$ if and only if $0 = \text{Ext}^1(M, N)^J \cong \text{Ext}^1(M^{(J)}, N) \cong \text{Ext}^1(L, N) \cong L'$ where the final isomorphism follows from [7, Prop. I.10.1]. By Appendix A, we have that $\text{Ext}^1(M(\lambda, -\infty), N) = 0$ and if and only if β and δ do not cross outside of 3-cycles. Thus, it suffices to show that $L' = 0$ if and only if $\text{Ext}^1(M(\lambda, -\infty), N) = 0$.

Clearly, if $\text{Ext}^1(M(\lambda, -\infty), N) = 0$, then $L' = 0$. We will therefore prove the claim by showing that $\dim_k(\text{Ext}^1(M(\lambda, -\infty), N)) \leq \dim_k(L')$. The inverse system consists of epimorphisms because $\text{coker } \phi$ is a band module and so $\text{pd}_{A(\Gamma)}(\text{coker } \phi) \leq 1$ by Lemma 2.14. Moreover, we can express the universal map $L' \rightarrow \text{Ext}^1(M(\lambda, -\infty), N)$ as an inverse limit of epimorphisms of the form

$$\text{Ext}^1(M(\lambda, -\infty), N) \xrightarrow{\text{Ext}^1(\phi^n, N)} \text{Ext}^1(M(\lambda, -\infty), N)$$

and so it is an epimorphism (see, for example, [50, Lem. 2.2]). This proves that indeed $\dim_k(\text{Ext}^1(M(\lambda, -\infty), N)) \leq \dim_k(L')$.

Now suppose that N is the generic module and M is a string module. Then the associated arc of N is $\delta = \beta$. By [49, Prop. 3], N is a direct summand of $M(\lambda, \infty)^K$ for any Prüfer module $M(\lambda, \infty)$ and any infinite set K . If α and β only cross in 3-cycles, then, by Appendix A, we have that $0 = \text{Ext}^1(M, M(\lambda, \infty))^K \cong \text{Ext}^1(M, M(\lambda, \infty)^K)$ and so $\text{Ext}^1(M, N) = 0$.

Conversely, suppose α and β cross outside of a 3-cycle. Then α must be a bridging arc and $\text{Ext}(M, X) \neq 0$ for any finite-dimensional band module X . By [49, Prop. 5], for any infinite set \mathcal{X} of finite-dimensional band modules from pairwise different homogeneous tubes, there exists a set H such that $0 \rightarrow \bigoplus_{X \in \mathcal{X}} X \rightarrow \prod_{X \in \mathcal{X}} X \rightarrow N^{(H)} \rightarrow 0$ is an exact sequence. In fact, this is a pure-exact sequence (see, for example, [46, Lem. 2.1.20]) and so, combining this with the fact that, by Lemma 2.14, we have $\text{id}_{A(\Gamma)}(\bigoplus_{X \in \mathcal{X}} X) \leq 1$, we have an exact sequence

$$0 \rightarrow \text{Ext}^1(M, \bigoplus_{X \in \mathcal{X}} X) \rightarrow \text{Ext}^1(M, \prod_{X \in \mathcal{X}} X) \rightarrow \text{Ext}^1(M, N^{(H)}) \rightarrow 0.$$

Finally, since M is finite-dimensional, $\text{Ext}^1(M, -)$ commutes with direct sums and direct products. It therefore follows that $\text{Ext}^1(M, N^{(H)}) \cong \text{Ext}^1(M, N)^{(H)} \neq 0$ and hence $\text{Ext}^1(M, N)$ is non-zero. \square

Next we consider the case where both indecomposable pure-injective modules are band modules. Since finite-dimensional band modules have self extensions, we only need to consider extensions between non-isomorphic infinite-dimensional band modules.

Proposition 2.17. *If M and N are both band modules over $A(\Gamma)$, then the following statements hold.*

- (1) Suppose M and N are both infinite dimensional. Then $\text{Ext}_{A(\Gamma)}^1(M, N) \neq 0$ if and only if $M \cong M(\lambda, \infty)$ and $N \cong M(\lambda, -\infty)$ for $\lambda \in K^*$.
- (2) If $M \cong M(\lambda, n)$ for $n \in \mathbb{N}$, then $\text{Ext}_{A(\Gamma)}^1(M, M) \neq 0$.

Proof. By Remark 2.13, we have $\text{Ext}_{A(\Gamma)}^1(M, N) \cong \text{Ext}_B^1(M, N)$ for any pair of indecomposable band modules where $B \cong K\tilde{A}_{s-1}$. The first statement then follows from [18, Lemma 2.7] and the second statement is well-known. \square

Middle terms of extensions. In the case of string and 1-dimensional band modules, it is possible to identify the middle terms of the extensions in $\text{Mod}(A(\Gamma))$. In fact, Proposition 2.20 gives a complete list of the extensions between string and 1-dimensional band modules that correspond to crossings of the associated arcs in (S, M) .

Definition 2.18. Consider the unique (periodic) \mathbb{Z} -string u_β . Consider the module $M^\Pi(\beta)$ that is defined exactly as $M^\oplus(\beta)$ but taking a direct product of copies of K instead of a direct sum. In particular, we have

$$M^\Pi(\beta) \cong \prod_{i \in \mathbb{Z}} K_i$$

(as vector spaces) where $K_i \cong K$ for each $i \in \mathbb{Z}$. Define $M^-(\beta)$ to be the submodule of $M^\Pi(\beta)$ such that $(\lambda_i)_{i \in \mathbb{Z}}$ is in $M^-(\beta) \subseteq M^\Pi(\beta)$ if and only if $\lambda_i = 0$ for $i \ll 0$. Define $M^+(\beta)$ to be the submodule of $M^\Pi(\beta)$ such that $(\lambda_i)_{i \in \mathbb{Z}}$ is in $M^+(\beta) \subseteq M^\Pi(\beta)$ if and only if $\lambda_i = 0$ for $i \gg 0$.

Definition 2.19. Let $A(\Gamma)$ be the algebra associated to a triangulated Γ of the annulus (S, M) and let α and δ be arcs in (S, M) .

In the statements of Proposition 2.20, v_{Le} , $d^{-1}v_R$, w_{Lc}^{-1} and aw_R are (possibly trivial) strings such that a, c, d, e are direct letters whenever they are contained in a non-trivial string. Moreover m is a (possibly trivial) finite string. We define

$$*_\delta = \begin{cases} \prod & \text{if } \delta \text{ is an expanding asymptotic arc;} \\ \oplus & \text{if } \delta \text{ is a contracting asymptotic arc;} \\ \emptyset & \text{if } \delta \text{ is a finite arc.} \end{cases}$$

$$*_\alpha = \begin{cases} \prod & \text{if } \alpha \text{ is an expanding asymptotic arc;} \\ \oplus & \text{if } \alpha \text{ is a contracting asymptotic arc;} \\ \emptyset & \text{if } \alpha \text{ is a finite arc.} \end{cases}$$

and the notation $M^\emptyset(\alpha)$ will mean $M(\alpha)$ for any finite arc α .

Proposition 2.20. Let $A(\Gamma)$ be the algebra associated to a triangulated Γ of the annulus (S, M) and let α and δ be arcs in (S, M) .

- (1) Suppose $u_\delta = v_{Lem}d^{-1}v_R$ and $u_\alpha = w_{Lc}^{-1}maw_R$ are such that v_{Le} and w_{Lc}^{-1} are not both trivial and $d^{-1}v_R$ and aw_R are not both trivial. Then there exists a non-split extension

$$0 \rightarrow M(\delta) \rightarrow M^{*\delta}(\varepsilon_1) \oplus M^{*\alpha}(\varepsilon_2) \rightarrow M(\alpha) \rightarrow 0$$

where $u_{\varepsilon_1} = v_{Lem}aw_R$ and $u_{\varepsilon_2} = w_{Lc}^{-1}md^{-1}v_R$. We call such an extension an **overlap extension** between string modules.

- (2) Suppose there exists $l \geq 1$ such that $u_\alpha = v_{Lem}d^{-1}v_R$ and $b^l = w_{Lc}^{-1}maw_R$. If we choose l to be minimal, then there exists a non-split overlap extension

$$0 \rightarrow M(\alpha) \rightarrow M(\varepsilon) \rightarrow M(\lambda, 1) \rightarrow 0$$

for all $\lambda \in K^*$ where

- (i) if $l = 1$ then $u_\varepsilon = v_{Lem}aw_Rw_{Lc}^{-1}md^{-1}v_R$; and
- (ii) if $l > 1$ then $u_\varepsilon = v_{Lem}m_2bm_1d^{-1}v_R$ such that $m = m_2m_1$ and $b = m_1aw_R$.

We call such an extension an **overlap extension** from a band module to a string module.

- (3) Suppose there exists $l \geq 1$ such that $b^l = v_L e m d^{-1} v_R$ and $u_\delta = w_L c^{-1} m a w_R$. If we choose l to be minimal, then there exists a non-split overlap extension

$$0 \rightarrow M(\lambda, 1) \rightarrow M(\varepsilon) \rightarrow M(\delta) \rightarrow 0$$

for all $\lambda \in K^*$ where

- (i) if $l = 1$ then $u_\varepsilon = w_L c^{-1} m d^{-1} v_R v_L e m a w_R$; and
- (ii) if $l > 1$ then $u_\varepsilon = w_L c^{-1} m_2 b m_1 a w_R$ such that $m = m_2 m_1$ and $b = m_1 d^{-1} v_R$.

We call such an extension an **overlap extension** from a string module to a band module.

- (4) Suppose there exists an arrow $a \in Q_1$ such that either $u_\alpha a^{-1} u_\delta^{-1}$ or $u_\delta a u_\alpha^{-1}$ is a finite string, an \mathbb{N} -string or u_β and denote this string by u_ε . Then there exists a non-split extension

$$0 \rightarrow M(\delta) \rightarrow M^*(\varepsilon) \rightarrow M(\alpha) \rightarrow 0$$

where

$$* = \begin{cases} + & \text{if } u_\varepsilon = u_\alpha a^{-1} u_\delta^{-1} = u_\beta; \\ - & \text{if } u_\varepsilon = u_\delta a u_\alpha^{-1} = u_\beta; \\ *_\alpha & \text{if } u_\varepsilon = u_\alpha a^{-1} u_\delta^{-1} \neq u_\beta; \\ *_\delta & \text{if } u_\varepsilon = u_\delta a u_\alpha^{-1} \neq u_\beta. \end{cases}$$


We call such an extension an **arrow extension**.

We refer to arrow and overlap extensions as **standard extensions**.


Proof. In [21], the authors prove this result for finite-dimensional modules. The extensions between infinite-dimensional modules are completely analogous. \square

- Remark 2.21.** (1) In cases (2) and (3) the arcs α and δ must be finite arcs since otherwise the above conditions on the strings u_α and u_δ cannot be satisfied.
(2) In case (4) α must be contracting and δ must be expanding.

Example 2.22. The first three pictures illustrate case 1 of Proposition 2.20.



The second three pictures illustrate case 4.



2.3. The classification of cosilting modules. In this section will classify the cosilting modules over $A(\Gamma)$ for a triangulation Γ of an annulus (S, M) , with quiver $Q = Q(\Gamma)$ and $A(\Gamma) = KQ/I(\Gamma)$. We begin by introducing the $A(\Gamma)$ -modules $M(t)$ that we will show are cosilting.

Let T be a collection of non-crossing arcs in (S, M) . We will call T **strictly asymptotic** if it contains an asymptotic arc. If either

- (i) $t = (T, P_1, P_2)$ where T is strictly asymptotic and P_1 and P_2 are disjoint subsets of K^* or
- (ii) $t = T$ where T is not strictly asymptotic,

then we call t a **partial asymptotic triangulation**. If T is a maximal collection of non-crossing asymptotic arcs and $K^* = P_1 \cup P_2$ (when t is strictly asymptotic), then t is called an **asymptotic triangulation**.

Remark 2.23. Note that the strictly asymptotic triangulations are in bijection with the set $\mathcal{A}(S, M) \times \mathcal{P}(K^*)$ defined in the Main Theorem in the introduction. Similarly the asymptotic triangulations that are not strictly asymptotic are in bijection with the set $\mathcal{T}(S, M)$.

Let t be a partial asymptotic triangulation, then we define a set of indecomposable pure-injective objects as follows. If t is strictly asymptotic, then let

$$\mathcal{N}_t := \{M(\alpha) \mid \alpha \in T\} \cup \{M(\lambda, \infty) \mid \lambda \in P_1\} \cup \{M(\lambda, -\infty) \mid \lambda \in P_2\} \cup \{G\}.$$

If t is not strictly asymptotic, then let

$$\mathcal{N}_t := \{M(\alpha) \mid \alpha \in T\}.$$

For each partial asymptotic triangulation t , let

$$M(t) := \prod_{N \in \mathcal{N}_t} N.$$

2.3.1. Annihilators of asymptotic triangulations. We will consider partial asymptotic triangulations t and consider the associated $A(\Gamma)$ -module $M(t)$. In order to show that $M(t)$ is a cosilting module over $A(\Gamma)$ whenever t is an asymptotic triangulation, we will make use of the characterisation of cosilting modules over left artinian rings given in Corollary 1.17. The first step will be to understand the annihilator ideal $\text{Ann}(M(t))$. The following observation holds for an arbitrary gentle algebra; see [48] for definitions of (possibly infinite-dimensional) string and band modules.

Observation 2.24. *Let $B = KQ/I$ be an arbitrary gentle algebra. Then the following hold.*

- (1) *If M is a string module corresponding to a string u , then $\text{Ann}(M)$ has a K -linear basis consisting of the paths p in (Q, I) such that neither p nor p^{-1} is contained in u .*

- (2) If N is a band module corresponding to a band b , then $\text{Ann}(N)$ has a K -linear basis consisting of the paths q in (Q, I) such that neither q nor q^{-1} is contained in \mathbb{Z} -string ${}_{\infty}b_{\infty}$.

For the rest of the section, let $Q = Q(\Gamma)$ and $I = I(\Gamma)$ for Γ a triangulation of an annulus, with string algebra $A(\Gamma) = KQ/I$ and let t be a (partial) asymptotic triangulation.

Lemma 2.25. *The annihilator ideal $\text{Ann}(M(t))$ in $A(\Gamma)$ has a K -linear basis consisting of all paths p satisfying the following conditions.*

- (1) Neither p nor p^{-1} occur as a substring of u_{α} for any $\alpha \in T$.
(2) If t is strictly asymptotic, then neither p nor p^{-1} occur as a substring of u_{β} .

Proof. The lemma follows immediately from the above observation since

$$\text{Ann}(M(t)) = \bigcap_{M \in \mathcal{N}_t} \text{Ann}(M).$$

□

Given Lemma 2.25, we will use $\text{Ann}(t)$ to denote the annihilator ideal associated to the module $M(t)$.

Lemma 2.26. *Let t be a partial asymptotic triangulation of (S, M) and let $p = p_n \dots p_0$ be a path of length $n + 1$ in $(Q, I(\Gamma))$. Then $p \in \text{Ann}(t)$ if and only if $p_i \in \text{Ann}(t)$ for some $i \in \{0, \dots, n\}$.*


Proof. By definition, either $t = T$ or $t = (T, P_1, P_2)$. Suppose p is contained in $\text{Ann}(t)$ but p_0 is not. By Lemma 2.25 there is some $\alpha \in T$ such that u_{α} contains p_0 or p_0^{-1} as a substring or T is strictly asymptotic and u_{β} contains p_0 or p_0^{-1} as a substring. Any arc γ such that u_{γ} contains $p_n \dots p_1$ or $p_1^{-1} \dots p_n^{-1}$ as a substring but does not contain p or p^{-1} as a substring will necessarily cross α and so cannot be contained in T . Thus $p_n \dots p_1 \in \text{Ann}(t)$ by Lemma 2.25. By induction on the length of p we have that $p \in \text{Ann}(t)$ implies $p_i \in \text{Ann}(t)$ for some $0 \leq i \leq n$. The converse is immediate. □

Corollary 2.27. *The algebra $A(\Gamma)/\text{Ann}(t)$ is a gentle algebra.*

Proof. By Lemma 2.26, the algebra $A(\Gamma)/\text{Ann}(t)$ is obtained from $A(\Gamma)$ by removing arrows and vertices and so the axioms of a gentle algebra will still be satisfied by $A(\Gamma)/\text{Ann}(t)$. □

Next we consider the annihilator of an asymptotic triangulation t . In this case we can describe the annihilator in more detail. In particular, $\text{Ann}(t)$ will contain arrows satisfying the following definition with respect to some $\alpha \in T$.

Definition 2.28. Let a be an arrow in Q with $s(a) = i$ and $t(a) = j$. We say that an arc α **crosses a in a 3-cycle** if i and j are edges of an internal triangle in Γ and either $s(u_{\alpha}) = k$ or $t(u_{\alpha}) = k$ where k is the remaining edge in the triangle, see figure below.



Note that this means that α and the arc corresponding to a cross in a 3-cycle, cf. Definition 2.15.

Lemma 2.29. *Let t be an asymptotic triangulation with corresponding T . Then the following statements hold.*

- (1) Let $a : i \rightarrow j$ be an arrow in Q . Then a is contained in $\text{Ann}(t)$ if and only if one of the following conditions hold.
(a) One of γ_i or γ_j is contained in T .

- (b) *There exists some $\alpha \in T$ such that α crosses a in a 3-cycle.*
(2) *Let i be a vertex in Q . Then e_i is contained in $\text{Ann}(t)$ if and only if $\gamma_i \in T \cap \Gamma$.*

Proof. (1) If either of the conditions hold then, it follows from Lemma 2.25, that $a \in \text{Ann}(t)$. We will therefore show the converse implication.

So let $a \in \text{Ann}(t)$. First suppose that γ_i and γ_j are edges of a triangle such that the remaining edge is a boundary component and suppose γ_i is not contained in T . As T is a triangulation, there must be some arc $\alpha \in T$ that crosses γ_i . By Lemma 2.25, neither a nor a^{-1} occur as a substring of u_α and so $s(u_\alpha) = i$ or $t(u_\alpha) = i$. Any arc γ that crosses γ_j such that u_γ does not contain a or a^{-1} as a substring will cross α and so cannot be contained in T . So γ_j is compatible with all arcs in T and we must therefore have that $\gamma_j \in T$.

Next suppose that γ_i and γ_j are edges of an internal triangle in Γ with remaining edge γ_k . If γ_i is not contained in T , then there exists an arc $\alpha \in T$ that crosses γ_i . Since u_α cannot contain a or a^{-1} as a substring, either $s(u_\alpha) = i$ or $t(u_\alpha) = i$ or α crosses γ_k immediately after γ_i . In the first situation, it follows that γ_j must be contained in $\text{Ann}(t)$. In the second situation, there must be an arc γ in T that crosses a in a 3-cycle and then agrees with α until its endpoint. As before, this arc is compatible with all of T and so is in T .

(2) The statement follows immediately from the fact that γ_i is crossed by an arc in T if and only if γ_i is not contained in T . \square

We now associate to any asymptotic triangulation t a new quiver Q_t as follows:

- I. If γ_i is contained in $\Gamma \cap T$, remove the vertex i from Q and all arrows incident with i .
- II. If a is an arrow in Q that is crossed in a 3-cycle by some $\alpha \in T$, then remove a from Q .

Let I_t be the admissible ideal in KQ_t generated by the generating relations in $I(\Gamma)$ that are supported on Q_t .

Corollary 2.30. *Let t be an asymptotic triangulation with quiver Q_t and ideal I_t . Then $A(\Gamma)/\text{Ann}(t)$ is isomorphic to the algebra KQ_t/I_t .*


2.3.2. Cosilting modules are partial asymptotic triangulations. Recall from Lemma 1.12 and Proposition 1.15 that every cosilting module C is equivalent to a cosilting module with no superdecomposable part. As in Remark 1.3, we will use the notation \mathcal{N}_C to denote the set of indecomposable pure-injective summands of C up to isomorphism.

Proposition 2.31. *Let C be a cosilting module over $A(\Gamma)$. There exists a partial asymptotic triangulation t such that $\text{Prod}(\mathcal{N}_C) = \text{Prod}(\mathcal{N}_t)$.*

Proof. By Theorem 1.13 and Proposition 1.15, the set \mathcal{N}_C is a maximal rigid system in the category $\text{Mod}(A(\Gamma)/\text{Ann}(C))$. We consider $\text{Mod}(A(\Gamma)/\text{Ann}(C))$ as a full subcategory of $\text{Mod}(A(\Gamma))$.


Let $T := \{\alpha \mid M(\alpha) \in \mathcal{N}_C\}$. We will show that T is a collection of non-crossing arcs in (S, M) . Let α and δ be contained in T . By Proposition 1.15, we have that $\text{Cogen}(C) \subseteq \text{Mod}(A(\Gamma)/\text{Ann}(C))$ is closed under extensions. It follows that $\text{Ext}_{A(\Gamma)/\text{Ann}(C)}^1(M(\alpha), M(\delta)) = 0$ if and only if $\text{Ext}_{A(\Gamma)}^1(M(\alpha), M(\delta)) = 0$. Similarly $\text{Ext}_{A(\Gamma)/\text{Ann}(C)}^1(M(\delta), M(\alpha)) = 0$ if and only if $\text{Ext}_{A(\Gamma)}^1(M(\delta), M(\alpha)) = 0$. By Theorem 2.16 the arcs α and δ cannot cross outside of 3-cycles or else this would contradict that \mathcal{N}_C is a maximal rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(C))$.

Suppose that α and δ cross in a 3-cycle. Without loss of generality, suppose we have the following local configuration.



Then we must have that $c \in \text{Ann}(C)$ because otherwise $\text{id}_{A(\Gamma)/\text{Ann}(C)}(M(\delta)) > 1$ by Lemma 2.12, which contradicts the fact that \mathcal{N}_C is a maximal rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(C))$. Let

ε be the arc that crosses d in a 3-cycle and follows α until its endpoint (if possible) or until ε meets an inverse letter. That is, we have the following local configuration.



Then $M(\varepsilon)$ is a finite-dimensional submodule of $M(\alpha)$ and hence $M(\varepsilon) \in \text{Cogen}(C)$. However, there is an arrow extension $0 \rightarrow M(\delta) \rightarrow M(\nu) \rightarrow M(\varepsilon) \rightarrow 0$ where $u_\nu = u_\delta c u_\varepsilon$ in $\text{Mod}(A(\Gamma))$ and $M(\nu)$ is not contained in $\text{Cogen}(C) \subseteq \text{Mod}(A(\Gamma)/\text{Ann}(C))$ because $c \in \text{Ann}(C)$. This contradicts that $\text{Cogen}(C)$ is closed under extensions and so α and δ do not cross.


If T is strictly asymptotic, then define $t := (T, P_1, P_2)$ where $P_1 := \{\lambda \in K^* \mid M(\lambda, \infty) \in \mathcal{N}_C\}$ and $P_2 := \{\lambda \in K^* \mid M(\lambda, -\infty) \in \mathcal{N}_C\}$. Note that $P_1 \cap P_2 = \emptyset$ by Theorem 2.16 and the same reasoning as above. If T is not strictly asymptotic, then define $t := T$. We have shown that t is a partial asymptotic triangulation. Clearly \mathcal{N}_C is a rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(C))$ and $\mathcal{N}_C \subseteq \mathcal{N}_t$ (with a proper inclusion exactly when G is not contained in \mathcal{N}_C). Since \mathcal{N}_C is maximal, we may conclude that $\text{Prod}(\mathcal{N}_C) = \text{Prod}(\mathcal{N}_t)$. \square

2.3.3. Asymptotic triangulations are cosilting modules. Let t be an asymptotic triangulation of (S, M) . In this section we show that $M(t)$ is a cosilting module over $A(\Gamma)$. By Corollary 1.17, it is enough to show that \mathcal{N}_t is a maximal rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(t))$ and that $\text{Cogen}(M(t))$ is a torsion-free class.

Proposition 2.32. *Let t be an asymptotic triangulation. The set \mathcal{N}_t is a maximal rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(t))$ and $M(t)$ is a cotilting $A(\Gamma)/\text{Ann}(t)$ -module.*

Proof. First note that, by Theorem 2.16, we have that $\text{Ext}_{A(\Gamma)}^1(M, N) = 0$ for all $N, M \in \mathcal{N}_t$ and so it follows that also $\text{Ext}_{A(\Gamma)/\text{Ann}(t)}^1(M, N) = 0$ for all $N, M \in \mathcal{N}_t$.

Next we show that $\text{id}_{A(\Gamma)/\text{Ann}(t)}(M) \leq 1$ for all $M \in \mathcal{N}_t$. If M is a band module then $\text{id}_{A(\Gamma)/\text{Ann}(t)}(M) \leq \text{id}_{A(\Gamma)}(M) \leq 1$ by Lemma 2.14. Suppose $M \cong M(\alpha)$ for some arc α . We show that the conditions of Lemma 2.12 holds for u_α . Suppose that the endpoint of α is contained in a triangle with edges γ_i, μ and ν such that ν follows μ in the anticlockwise direction.



Without loss of generality assume that $s(u_\alpha) = i$. If all of γ_i, μ and ν are arcs in Γ , say $\mu = \gamma_j$ and $\nu = \gamma_k$, then there are arrows $a: j \rightarrow k$, $c: k \rightarrow i$ and $d: i \rightarrow j$ in Q . Moreover, the arrow c is the unique arrow such that $u_\alpha c$ is a string and the arrow a is the unique arrow such that $ca \in I(\Gamma)$. However, the arc α crosses a in a 3-cycle so by Corollary 2.30 there is no arrow d in Q_t such that $cd \in I_t$. Thus condition (i) of Lemma 2.12 holds. In a similar way, if either or both of μ and ν are boundary components, then, even if there exists an arrow c such that $u_\alpha c$ is a string, there is no arrow a such that $ca \in I_t$. If α is finite, we may run the same argument at the other endpoint of α and we have that $\text{id}_{A(\Gamma)/\text{Ann}(t)}(M) \leq 1$ by Proposition 2.12.

We have shown that \mathcal{N}_t is a rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(t))$. Finally we must show that it is maximal. Let \mathcal{L} be a rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(t))$ such that $\text{Prod}(\mathcal{N}_t) \subseteq \text{Prod}(\mathcal{L})$.

Without loss of generality, we may assume that $M(t)$ is a direct summand of the direct product L of the objects in \mathcal{L} i.e. we have that $L \cong M(t) \oplus X$ for some module X . Now, by Lemma 1.10, we must have that $\text{Ext}_{A(\Gamma)/\text{Ann}(t)}^1(L, L) = 0$. It follows that $\text{Ext}_{A(\Gamma)/\text{Ann}(t)}^1(Y, M(t)) = 0 = \text{Ext}_{A(\Gamma)/\text{Ann}(t)}^1(M(t), Y)$ for all indecomposable summands Y of X . Suppose there is an indecomposable summand Y of X such that $\text{Ext}_{A(\Gamma)}^1(Y, M(t)) \neq 0$ or $\text{Ext}_{A(\Gamma)}^1(M(t), Y) \neq 0$. By Lemma A.10, there exists an arrow or an overlap extension between Y and some $N \in \mathcal{N}_t$. By Lemma 2.29, the middle term of any overlap extension between Y and M will be contained in $\text{Mod}(A(\Gamma)/\text{Ann}(t))$. Therefore the only possibility is the following. There is some $\delta \in T$ and a arc ε such that $Y \cong M(\varepsilon)$ such that there is an arrow extension between $M(\delta)$ and $M(\varepsilon)$ with the corresponding arrow a contained in $\text{Ann}(t)$. However, by Lemma 2.29, this implies that there is a arc $\alpha \in T$ such that α crosses a in a 3-cycle. But then α crosses δ , which is a contradiction.


So then all indecomposable summands of X must be contained in \mathcal{N}_t because any other indecomposable pure-injective module will have a non-trivial extension with $M(t)$. By [47, Thm. 5.6] there are no superdecomposable pure-injective modules over $A/\text{Ann}(t)$ and so, by Remark 1.11, we have shown that $\text{Prod}(\mathcal{L}) = \text{Prod}(\mathcal{N}_t)$.

By Theorem 1.13, we also have that $M(t)$ is a cotilting module over $A(\Gamma)/\text{Ann}(t)$. \square

Proposition 2.33. *If t is an asymptotic triangulation, then $M(t)$ is a cosilting $A(\Gamma)$ -module.*

Proof. By Corollary 1.17 and Proposition 2.32, it remains to show that $\text{Cogen}(M(t))$ is a torsion-free class in $\text{Mod}(A(\Gamma))$. We know that $\text{Cogen}(M(t))$ is a torsion-free class in $\text{Mod}(A(\Gamma)/\text{Ann}(t))$ because $M(t)$ is a $A(\Gamma)/\text{Ann}(t)$ -cotilting module by Proposition 2.32. In particular, if M, N are in $\text{Cogen}(M(t))$ such that $0 \rightarrow M \rightarrow L \rightarrow N \rightarrow 0$ is a short exact sequence in $\text{Mod}(A(\Gamma)/\text{Ann}(t))$, then $L \in \text{Cogen}(M(t))$. As we saw in the proof of Proposition 2.32, any overlap extension in $\text{Mod}(A(\Gamma)/\text{Ann}(t))$ is also an overlap extension in $\text{Mod}(A(\Gamma))$. We must rule out the possibility that there is an arrow extension $0 \rightarrow M \rightarrow L \rightarrow N \rightarrow 0$ in $\text{Mod}(A(\Gamma))$ with $L \notin \text{Cogen}(M(t))$.

Suppose there is an arrow extension between M and N . Without loss of generality we may assume that $M \cong M(\alpha)$ and $N \cong M(\delta)$ for arcs α and δ and that there is an arrow $a \in \text{Ann}(t)$ such that $u_\delta a^{-1} u_\alpha$ is a string. By Lemma 2.29, this means that there is $\varepsilon \in T$ such that ε crosses a in a 3-cycle. We must therefore have the following local configuration.



The arrows c and d cannot be in $\text{Ann}(t)$ because any arc crossing them in a 3-cycle would cross ε . Thus $\text{Ext}_{A(\Gamma)/\text{Ann}(t)}^1(M(\alpha), M(\varepsilon)) \neq 0$ since $u_\alpha c^{-1} u_\varepsilon$ is a string in (Q_t, I_t) . But this contradicts that $\text{Cogen}(M(t)) = {}^\perp M(t)$ in $\text{Mod}(A(\Gamma)/\text{Ann}(t))$. \square

2.3.4. The classification.

Lemma 2.34. *If t is a partial asymptotic triangulation, with corresponding T , then the following statements hold.*


- (1) For $i \in Q$, $e_i \in \text{Ann}(t)$ if and only if $\gamma_i \in \Gamma$ does not cross any arc in T .
- (2) Let a be an arrow in Q such that $e_{s(a)}$ and $e_{t(a)}$ are not contained in $\text{Ann}(t)$. Then $a \in \text{Ann}(t)$ if and only if one of the following statements hold.
 - (i) There is an arc $\alpha \in T$ that crosses a in a 3-cycle.
 - (ii) There is an arc α in (S, M) that crosses a in a 3-cycle such that $\text{Ann}(t) = \text{Ann}(t \cup \{\alpha\})$. Moreover, the arc α does not cross any arc in T .

Proof. (1) This is clear from Observation 2.24.

(2) (\Leftarrow) In both case (i) and (ii), we must have $a \in \text{Ann}(t)$ because there cannot be any arc $\delta \in T$ such that u_δ contains a or a^{-1} as a substring since such a δ would cross α .

(\Rightarrow) Suppose $a \in \text{Ann}(t)$ and there is no arc in T that crosses a in a 3-cycle. First we will argue that there is no arc δ in T such that $s(u_\delta) = s(a)$ or $s(u_\delta) = t(a)$. If there is a $\delta \in T$ such that $s(u_\delta) = t(a)$, then $e_{s(a)} \in \text{Ann}(t)$ because any other arc η that crosses $\gamma_{s(a)}$ must either cross δ or u_η must contain a or a^{-1} as a substring. Such an η cannot be contained in T . A similar argument yields that, if there is a $\delta \in T$ such that $s(u_\delta) = s(a)$, then $e_{t(a)} \in \text{Ann}(t)$. We have assumed that $e_{s(a)}$ and $e_{t(a)}$ are not contained in $\text{Ann}(t)$ so there is no arc δ in T such that $s(u_\delta) = s(a)$ or $s(u_\delta) = t(a)$.

It follows that there exists an arc α that crosses a in a 3-cycle and does not cross any arcs in T . Finally we must show that we can choose α such that $\text{Ann}(t) = \text{Ann}(t \cup \{\alpha\})$. We have that $\text{Ann}(t \cup \{\alpha\}) \subset \text{Ann}(t)$, so we must show that u_α does not contain any (trivial or non-trivial) letters that are not already contained in u_δ for some $\delta \in T$. Let $\gamma_k \in \Gamma$ be such that $\gamma_{s(a)}, \gamma_{t(a)}, \gamma_k$ form a triangle. The conclusion of the first part of this proof tells us that we have some arcs δ_1 and δ_2 in T such that δ_1 crosses $\gamma_{s(a)}$ followed by γ_k and δ_2 crosses $\gamma_{t(a)}$ followed by γ_k . Choose α to cross a in a 3-cycle and then follow δ_1 to its endpoint.



□

Lemma 2.35. *Let t be a partial asymptotic triangulation. There exists an asymptotic triangulation s such that $\mathcal{N}_t \subseteq \mathcal{N}_s$ and $\text{Ann}(t) = \text{Ann}(s)$.*

Proof. By definition, we have that either $t = T$ or $t = (T, P_1, P_2)$. For the second case, let $Q_1 := P_1$ and let $Q_2 := K^* \setminus P_1$. In the following proof we will define sets S_0, \dots, S_{n+1} of arcs in (S, M) for some $n \in \mathbb{N}$, all containing T , from which we will construct s .

We fix the following notation: for each $0 \leq i \leq n+1$, let

$$s_i := \begin{cases} S_i & \text{if } t = T; \\ (S_i, Q_1, Q_2) & \text{if } t = (T, P_1, P_2) \end{cases}$$

where the sets S_i are defined as follows. Let $S_0 := \{\gamma_i \in \Gamma \mid \gamma_i \text{ does not cross any } \alpha \in T\} \cup T$. Clearly $\text{Ann}(s_0) = \text{Ann}(t)$. Let a_1, \dots, a_n be the arrows in $\text{Ann}(t)$ that are not crossed in a 3-cycle by any arc in T and such that $s(a_i), t(a_i)$ are not contained in $\text{Ann}(t)$. Then, for $1 \leq i \leq n$, let $S_i := S_0 \cup \{\alpha_1, \dots, \alpha_i\}$ where α_i is an arc such that α_i crosses a_i in a 3-cycle and $\text{Ann}(s_i) = \text{Ann}(s_{i-1})$. Each α_i exists by Lemma 2.34.

Let $S := S_{n+1}$ be any completion of S_n to a full asymptotic triangulation of (S, M) and let $s := s_{n+1}$. Clearly $\mathcal{N}_t \subseteq \mathcal{N}_s$. We finish the proof by arguing that $\text{Ann}(s) = \text{Ann}(t)$. Since $M(t)$ is a direct summand of $M(s)$, we have that $\text{Ann}(s) \subseteq \text{Ann}(t)$. By Lemma 2.26, in order to show that $\text{Ann}(t) \subseteq \text{Ann}(s)$, it is sufficient to show that every vertex and every arrow contained in $\text{Ann}(t)$ is also contained in $\text{Ann}(s)$.

Suppose e_i is a vertex in $\text{Ann}(t)$. Then either $\gamma_i \in T \cap \Gamma \subseteq S \cap \Gamma$ or γ_i is not crossed by any arc in T . In the latter case we have that $\gamma_i \in S_0 \subseteq S \cap \Gamma$. By Lemma 2.29, both cases imply that $e_i \in \text{Ann}(s)$.

Suppose a is an arrow contained in $\text{Ann}(t)$. By the above argument, we may assume that both $e_{s(a)}$ and $e_{t(a)}$ are not in $\text{Ann}(t)$. Then either a is crossed in a 3-cycle by some $\alpha \in T \subseteq S$

or $a = a_j$ for some $1 \leq j \leq n$. In the latter case, we have that a is crossed in a 3-cycle by $\alpha_j \in S_j \subseteq S$. By Lemma 2.29, we have that $a \in \text{Ann}(s)$. \square

By Remark 2.23, the following theorem yields the Main Theorem of the introduction.

Theorem 2.36. *Every cosilting $A(\Gamma)$ -module C is equivalent to a module $M(t)$ where t is a uniquely determined asymptotic triangulation.*

Proof. Suppose C is a cosilting module then C is a cotilting module over $A(\Gamma)/\text{Ann}(C)$ and by Theorem 1.13, we have that \mathcal{N}_C is a maximal rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(C))$. Then, by Proposition 2.31, there is a partial asymptotic triangulation such that $\mathcal{N}_C = \mathcal{N}_t$ and so $\text{Ann}(C) = \text{Ann}(t)$. By Lemma 2.35, there exists an asymptotic triangulation s such that $\text{Ann}(s) = \text{Ann}(t)$ and $\mathcal{N}_C = \mathcal{N}_t \subseteq \mathcal{N}_s$. By Proposition 2.33, we have that $M(s)$ is cosilting and so \mathcal{N}_s is a maximal rigid system in $\text{Mod}(A(\Gamma)/\text{Ann}(C))$ such that $\text{Prod}(\mathcal{N}_C) \subseteq \text{Prod}(\mathcal{N}_s)$. Therefore, by the maximality of \mathcal{N}_C , we have that $\text{Prod}(\mathcal{N}_C) = \text{Prod}(\mathcal{N}_s)$ and so C is equivalent to $M(s)$.

Suppose there is some other asymptotic triangulation s such that $M(s)$ is equivalent to C . Then $\text{Ann}(s) = \text{Ann}(t)$ and $\text{Prod}(\mathcal{N}_s) = \text{Prod}(\mathcal{N}_t)$. As $M(s) \in \text{Prod}(\mathcal{N}_t)$, there exists a module X such that $M(s) \oplus X \cong \prod_{i \in I} M_i$ with $M_i \in \mathcal{N}_t$. Without loss of generality, assume that a copy of every module in \mathcal{N}_t occurs as a summand of $X \oplus M(s)$. By Lemma 1.10, we have that $\text{Ext}_{A(\Gamma)/\text{Ann}(C)}^1(\prod_{i \in I} M_i, \prod_{i \in I} M_i) = 0$ and hence, for every $N \in \mathcal{N}_t$, we have $\text{Ext}_{A(\Gamma)/\text{Ann}(C)}^1(N, M) = 0 = \text{Ext}_{A(\Gamma)/\text{Ann}(C)}^1(M, N)$ for all $M \in \mathcal{N}_s$. Let δ be the arc corresponding to N and first suppose $N \cong M(\delta)$. As we argued in Proposition 2.32, it follows that, for every arc $\alpha \in S$, if δ and α cross, then they cross in a 3-cycle or in an arrow $a \in \text{Ann}(s) = \text{Ann}(t)$. If α and δ cross in a 3-cycle, then $\text{Ann}(s) \neq \text{Ann}(t)$, by Lemma 2.29, which is a contradiction so α and δ can not cross in a 3-cycle.

If α and δ cross in an arrow $a \in \text{Ann}(t) = \text{Ann}(s)$, then there are arcs $\varepsilon, \varepsilon'$ (with $\varepsilon \in S$ and $\varepsilon' \in T$) that cross a in a 3-cycle, by Lemma 2.29. But then ε crosses α and ε' crosses δ , which is not possible. Thus δ does not cross any arc $\alpha \in S$ and so it must be that δ is in S . If N is a band module then the above argument implies that s is strictly asymptotic, since otherwise there would be a bridging arc that crosses β . Again Ext-orthogonality implies $P_1 = Q_1$ and $P_2 = Q_2$. We may argue in the same, exchanging the roles of s and t . Therefore we have shown $t = s$. \square

Remark 2.37. From this theorem we recover the classification of cotilting modules over $\tilde{K}\tilde{A}_n$ given in [18] by choosing Γ to consist only of bridging arcs. The cotilting modules are then given by the asymptotic triangulations t such that $\Gamma \cap T = \emptyset$.

Such a choice of Γ also yields the description of the torsion classes in a tube category given in [8]. To see this, one should note that the arcs involving only one boundary component gives rise to a geometric model of one of the two (possibly non-homogeneous) tubes in $\text{mod}(A(\Gamma))$ that contain string modules.

APPENDIX APPENDIX A EXTENSIONS AND ASYMPTOTIC ARCS.

Throughout this appendix we will follow the definitions and notation set up in Section 2. In particular, we fix S to be the annulus and M to be a set of marked points in the boundary of S . We will then consider the K -algebra $A(\Gamma)$ (defined as in Subsection 2.1) where Γ is a triangulation of (S, M) . As in Section 2, we assume that K is algebraically closed.

The main result will be the following.

Theorem A.1. *Let M and N be string or band modules in $\text{Mod}(A(\Gamma))$ and let α and δ be their respective associated arcs. If M and N are not both band modules, then the following statements are equivalent.*

- (1) *The K -vector spaces $\text{Ext}_{A(\Gamma)}^1(M, N)$ and $\text{Ext}_{A(\Gamma)}^1(N, M)$ are both zero.*
- (2) *The arcs α and δ do not cross outside of 3-cycles.*

We will approach this theorem in stages. First we will describe explicitly the form of extensions between indecomposable pure-injective $A(\Gamma)$ -modules. Our approach is similar to [5, 21, 22]. We will then connect these extensions to crossings of arcs in (S, M) (outside of 3-cycles). In this first section our approach is similar to [23].

A.1 Extensions in $\text{Mod}(A(\Gamma))$ and morphisms in the homotopy category. Let M and N be string or band modules over $A(\Gamma)$ and let α and δ be their respective associated arcs. Assume that M and N are not both band modules. We will identify M and N with their projective resolutions, denoted P_M^\bullet and P_N^\bullet respectively, and make use of the following well-known isomorphisms (see, for example, [54, Cor. 10.4.7, Cor. 10.7.5]).

$$\begin{aligned}\text{Ext}_{A(\Gamma)}^1(M, N) &\cong \text{Hom}_{D(A(\Gamma))}(M[0], N[1]) \\ &\cong \text{Hom}_{K(A(\Gamma))}(P_M^\bullet, P_N^\bullet[1]).\end{aligned}$$

where $D(A(\Gamma))$ is the unbounded derived category of $\text{Mod}(A(\Gamma))$ and $K(A(\Gamma))$ is the homotopy category of (co)chain complexes in $\text{Mod}(A(\Gamma))$.

In [14] it is shown that every indecomposable complex in $D^b(\text{mod}(A(\Gamma)))$ is isomorphic to a complex of finite-dimensional projective modules that is either a string complex or a band complex associated to a homotopy string or a homotopy band. We will see that P_N^\bullet and P_M^\bullet also have the form of a string or a band complex, however the projective modules may be infinite-dimensional.

Next we define what we mean by a homotopy string in this setting. Observe that any finite string may be uniquely expressed as a sequence $p_n q_n^{-1} \dots p_1 q_1^{-1} p_0 q_0^{-1}$ where p_i and q_j are paths of length ≥ 1 for $0 \leq i < n$ and $0 < j \leq n$ and p_n and q_0 are possibly trivial paths. Similarly, any \mathbb{N} -string may be uniquely expressed as a sequence $\dots p_1 q_1^{-1} p_0 q_0^{-1}$. In the following definition, all strings will be represented in this way.

- Definition A.2.** (1) A sequence of arrows $\dots a_3 a_2 a_1 a_0$ is called a **antipath** if $a_{i+1} a_i \in I(\Gamma)$ for all $i \geq 0$.
- (2) Let $u_\alpha = \dots p_1 q_1^{-1} p_0 q_0^{-1}$ be an \mathbb{N} -string (respectively let $u_\alpha = p_n q_n^{-1} \dots p_1 q_1^{-1} p_0 q_0^{-1}$ be a finite string).
- (a) If there exists an arrow a_0 such that $u_\alpha a_0^{-1}$ is a string, then define $h(u_\alpha)$ to be the sequence $u_\alpha a_0^{-1} a_1^{-1} a_2^{-1} \dots$ where $\dots a_2 a_1 a_0$ is the longest possible antipath starting at a_0 .
- (b) If there is no such arrow a_0 , then define $h(u_\alpha)$ to be the sequence $\dots p_1 q_1^{-1} p_0$ (respectively $p_n q_n^{-1} \dots p_1 q_1^{-1} p_0$).
- (3) Let u_α be an \mathbb{N} -string, then the **homotopy string** τ_α **associated to** u_α is defined to be $h(u_\alpha)$.
- (4) Let u_α be a finite string, then the **homotopy string** τ_α **associated to** u_α is defined to be $h(h(u_\alpha^{-1})^{-1})$.

From this data we can define string complexes, which are essentially infinite-dimensional versions of the string complexes defined in [14]. For a vertex i in Q , we will denote the corresponding indecomposable projective module by $P(i)$. Moreover, we will make use of the K -basis of $\text{Hom}_{A(\Gamma)}(P(i), P(j))$ given by paths $p: j \rightarrow i$. We will denote these basis elements by the paths p .

As with string infinite-dimensional string modules, we must consider the expanding and contracting cases separately.

- Definition A.3.** (1) Suppose $u_\alpha = \dots p_1 q_1^{-1} p_0 q_0^{-1}$ is a contracting \mathbb{N} -string and let τ_α be the homotopy string associated to u_α . If $h(u_\alpha)$ fell into case (b), then we take $P(t(a_k))$, $a_0 q_0$ and a_{k+1} to be zero for $k \geq 0$.

Define the **string complex** P_α^\bullet as follows.

$$P_\alpha^0 := \bigoplus_{i \geq 0} P(s(p_i)), \quad P_\alpha^{-1} := P(t(a_0)) \oplus \left(\bigoplus_{i \geq 0} P(t(p_i)) \right), \quad P_\alpha^{-k} := P(t(a_{k-1})), \quad P_\alpha^{-l} := 0$$

for $k > 1$ and $l < 0$. Also let

$$d_\alpha^{-1} := (A_{rs})_{r>0, s>0}, \quad d_\alpha^{-k} := a_{k-1}, \quad d_\alpha^{-l} := 0$$

for $k > 1$ and $l < 1$ where

$$A_{rs} := \begin{cases} a_0 q_0 & \text{if } r = 1, s = 1, \\ q_i & \text{if } r = i + 1, s = i + 1 \text{ and } i > 0, \\ p_i & \text{if } r = i + 1, s = i + 2 \text{ and } i \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

- (2) Suppose $u_\alpha = \dots p_1 q_1^{-1} p_0 q_0^{-1}$ is a expanding \mathbb{N} -string and τ_α is the homotopy string associated to u_α . Define the **string complex** P_α^\bullet in the same way as in the contracting case but replacing any direct sums with direct products.
- (3) Suppose u_α is a finite string and let τ_α be the associated homotopy string. If $h(u_\alpha^{-1})$ fell into case **(a)**, then denote the antipath by $\dots a_2 a_1 a_0$. If $h(h(u_\alpha^{-1})^{-1})$ fell into case **(a)**, then denote the antipath by $\dots c_2 c_1 c_0$. If $h(u_\alpha^{-1})$ fell into case **(b)**, then we take $P(t(a_k))$, $a_0 p_0$ and a_{k+1} to be zero for $k \geq 0$. If $h(h(u_\alpha^{-1})^{-1})$ fell into case **(b)**, then we take $P(t(c_k))$, $c_0 p_n$ and c_{k+1} to be zero for $k \geq 0$.

Define the **string complex** P_α^\bullet as follows.

$$P_\alpha^0 := \bigoplus_{i=0}^n P(s(p_i)), \quad P_\alpha^{-1} := P(t(a_0)) \oplus \left(\bigoplus_{i=0}^n P(t(p_i)) \right) \oplus P(t(c_0)),$$

$$P_\alpha^{-k} := P(t(a_{k-1})) \oplus P(t(c_{k-1})), \quad P_\alpha^{-l} := 0$$

for $k > 1$ and $l < 0$. Also let

$$d_\alpha^{-1} := (A_{rs})_{0 < r \leq n+1, 0 < s \leq n+2}, \quad d_\alpha^{-k} := \begin{pmatrix} a_{k-1} & 0 \\ 0 & c_{k-1} \end{pmatrix}, \quad d_\alpha^{-l} := 0$$

for $k > 1$ and $l < 1$ where

$$A_{rs} := \begin{cases} a_0 q_0 & \text{if } r = 1, s = 1, \\ q_i & \text{if } r = i + 1, s = i + 1 \text{ and } 0 < i \leq n - 1, \\ p_i & \text{if } r = i + 1, s = i + 2 \text{ and } 0 < i \leq n - 1, \\ c_0 p_n & \text{if } r = n + 1, s = n + 2, \\ 0 & \text{otherwise.} \end{cases}$$

We can view the definition of P_α^\bullet for an \mathbb{N} -string schematically as follows.

$$\begin{array}{ccccc} & & \vdots & & \\ & & \xrightarrow{q_2} & & \\ & & \searrow^{p_1 q_1} & & \\ & & \xrightarrow{p_0} & & \\ & & \searrow^{a_0 q_0} & & \\ \dots & \xrightarrow{a_2} & P(t(a_1)) & \xrightarrow{a_1} & P(t(a_0)) & \xrightarrow{a_0 q_0} & P(s(q_0)) \end{array}$$

Similarly, we can view the definition of P_α^\bullet for a finite string schematically as follows.

$$\begin{array}{ccccccc}
 \dots & \xrightarrow{c_2} & P(t(c_2)) & \xrightarrow{c_1} & P(t(c_0)) & & \\
 & & & & & \searrow^{c_0 p_n} & \\
 & & & & P(t(p_{n-1})) & \xrightarrow{q_n} & P(s(p_n)) \\
 & & & & & \vdots & \\
 & & & & P(t(p_1)) & & \\
 & & & & & \searrow^{p_1} & \\
 & & & & P(t(p_0)) & \xrightarrow{q_1} & P(s(p_1)) \\
 & & & & & \searrow^{p_0} & \\
 \dots & \xrightarrow{a_2} & P(t(a_1)) & \xrightarrow{a_1} & P(t(a_0)) & \xrightarrow{a_0 q_0} & P(s(q_0))
 \end{array}$$

As in [5, 21, 22], we will make use of unfolded diagrams i.e. we represent the complexes P_α^\bullet in the following ways.

The figure consists of two horizontal diagrams illustrating a sequence of points and intervals. The top diagram shows a sequence of points (dots) with intervals labeled $p_2, q_2, p_1, q_1, p_0, a_0q_0, a_1, a_2$. The points are labeled with integers above them: 0, -1, 0, -1, 0, -1, -2. The bottom diagram shows a sequence of points with intervals labeled $c_1, c_0p_n, q_n, p_{n-1}, p_0, a_0q_0, a_1$. The points are labeled with integers above them: -2, -1, 0, -1, 0, -1, 0, -2. The diagrams illustrate the construction of a sequence of points and intervals, with the top diagram showing the initial construction and the bottom diagram showing the construction after a shift.

Recall that u_β is the \mathbb{Z} -string associated to the closed arc β . We will refer to u_β as the **homotopy string associate to β** and we will denote u_β by τ_β when we are considering it as a homotopy string. Next we will use a string complex $P_{u_\beta}^\bullet$ to define generalised band complexes. Let $R := K[\Phi, \Phi^{-1}]$ and recall the classification of indecomposable pure-injective modules given in Remark 2.9.

Definition A.4. Let $u_\beta = \dots p_1 q_1^{-1} p_0 q_0^{-1} p_{-1} q_{-1}^{-1} \dots$ be the periodic \mathbb{Z} -string. Define the complex $P_{u_\beta}^\bullet$ to be

$$P_{u_\beta}^{-1} := \bigoplus_{i \in \mathbb{Z}} P(t(p_i)), \quad P_{u_\beta}^0 := \bigoplus_{i \in \mathbb{Z}} P(s(p_i)), \quad P_{u_\beta}^k := 0$$

for all $k \neq 0, -1$. Also let $d_{u_\beta}^{-1} := (A_{rs})_{r,s \in \mathbb{Z}}$ and $d_{u_\beta}^k := 0$ for all $k \neq -1$ where

$$A_{rs} := \begin{cases} q_i & \text{if } r = i + 1, s = i + 1, \\ p_i & \text{if } r = i + 1, s = i + 2, \\ 0 & \text{otherwise.} \end{cases}$$

We can view $P_{u_\beta}^\bullet$ as a complex of $(A(\Gamma), R)$ -bimodules (see, for example, [16, Def. 1.3.45]). For each $\lambda \in K^*$ and each $n \in \mathbb{N} \cup \{\infty, -\infty\}$, define the **band complex** $B_{\lambda,n}^\bullet$ to be the complex obtained by applying $-\otimes_R N_{\lambda,n}$ to $P_{u_\beta}^\bullet$.

Remark A.5. (1) The complex $B_{\lambda,n}^\bullet$ is a complex of projective $A(\Gamma)$ -modules (see, for example, [16, Lem. 1.3.47]).

(2) Since K is algebraically closed, the complex $B_{\lambda,n}^\bullet$ is a band complex as defined in [14] when $n \in \mathbb{N}$.

We also associate an unfolded diagram to the complexes $B_{\lambda,1}^\bullet$. The unfolded diagrams of the other band complexes will be considered later (see ...). Since u_β is periodic, we have that there exist non-trivial paths m_0, \dots, m_t and n_0, \dots, n_t such that $u_\beta = \dots m_t^{-1} n_0 m_0^{-1} n_1 \dots n_t m_t^{-1} n_0 \dots$. The complex $B_{\lambda,1}^\bullet$ is then represented as follows.

$$\dots \xrightarrow{n_{t-2}} \overset{0}{\bullet} \xleftarrow{m_{t-1}} \overset{-1}{\bullet} \xrightarrow{n_{t-1}} \overset{0}{\bullet} \xleftarrow{\lambda m_t} \overset{-1}{\bullet} \xrightarrow{n_0} \overset{0}{\bullet} \xleftarrow{m_0} \overset{-1}{\bullet} \xrightarrow{n_1} \overset{0}{\bullet} \xleftarrow{m_1} \dots$$

The aim of the preceding definitions was to identify the projective resolutions of the indecomposable pure-injective modules (excluding G). The following lemma confirms that this is the case.

Lemma A.6. (1) Let u_α be a finite or an \mathbb{N} -string and let α be the associated homotopy string. The complex P_α^\bullet is quasi-isomorphic to $M(\alpha)[0]$.
(2) Let $\lambda \in K^*$ and let $n \in \mathbb{N} \cup \{\infty, -\infty\}$. The complex $B_{\lambda,n}$ is quasi-isomorphic to $M(\lambda, n)[0]$.

Proof. (1) Compute cohomology as in [21, Cor. 2.12].

(2) By computing cohomology of $P_{u_\beta}^\bullet$, we have that the following sequence is a projective resolution of $M^\oplus(\beta)$.

$$0 \rightarrow P_{u_\beta}^{-1} \rightarrow P_{u_\beta}^0 \rightarrow M^\oplus(\beta) \rightarrow 0.$$

Moreover, the functor $- \otimes_R N_{\lambda,n}$ is right exact, so

$$0 \rightarrow B_{\lambda,n}^{-1} \rightarrow B_{\lambda,n}^0 \rightarrow M(\lambda, n) \rightarrow 0$$

is right exact. The module $M^\oplus(\beta)$ is free as an R -module, so $\text{Tor}_1^R(M^\oplus(\beta), N_{\lambda,n}) = 0$, therefore the sequence is exact. We have shown that this sequence is a projective resolution of $M(\lambda, n)$ and so in particular, the complex $B_{\lambda,n}^\bullet$ is quasi-isomorphic to $M(\lambda, n)[0]$. \square

Let M and N be string modules or band modules with $n = 1$, such that M and N are not both band modules. Let α and δ be the associated arcs in (S, M) . Also, denote the projective resolutions of M and N by P_M^\bullet and P_N^\bullet respectively; note that these are given in Lemma A.6. Next we will establish when there exists a non-zero morphism $P_M^\bullet \rightarrow P_N^\bullet[1]$ by comparing the homotopy strings τ_α and τ_δ .

We refer to [5, Sec. 3] for the definition of **(singleton) single, double, graph and quasi-graph maps** between P_M^\bullet and $P_N^\bullet[1]$. These definitions extend easily to include projective resolutions of infinite-dimensional string modules.

Remark A.7. In [5] a quasi-graph map “from P_M^\bullet to P_N^\bullet ” determines a non-trivial homotopy class of morphisms $P_M^\bullet \rightarrow P_N^\bullet[1]$ that does not contain zero. We use the notation $P_M^\bullet \rightsquigarrow P_N^\bullet[1]$ to indicate that there is such a quasi-graph map.

For there to be a graph map $P_M^\bullet \rightarrow P_N^\bullet[1]$ or quasi-graph map $P_M^\bullet \rightsquigarrow P_N^\bullet[1]$, there must be a (maximal) common substring shared by τ_α and τ_δ ; we stress that this common substring ρ must be finite unless ρ contains an infinite length antipath. We will call quasi-graph maps $P_M^\bullet \rightsquigarrow P_N^\bullet[1]$, graph maps, singleton single and singleton double maps $P_M^\bullet \rightarrow P_N^\bullet[1]$ the **standard maps** from P_M^\bullet to $P_N^\bullet[1]$.

If we apply the main theorem of [5] to our setting, then we obtain the following proposition. Since the proof of Proposition A.9 will extend the arguments given in [5], we include a step-by-step sketch of the proof in order to refer to it later.

Proposition A.8 ([5, Thm. 3.15]). *Let M and N be finite-dimensional string modules or band modules with $n = 1$, such that M and N are not both band modules, and let P_M^\bullet and P_N^\bullet be the projective resolutions of M and N respectively. The standard maps $P_M^\bullet \rightarrow P_N^\bullet[1]$ form a K -basis for $\text{Hom}_{K(A(\Gamma))}(P_M^\bullet, P_N^\bullet[1])$.*

Proof. **Step 1::** First it is shown that the graph, single and double maps form a basis for $\text{Hom}_{C(A(\Gamma))}(P_M^\bullet, P_N^\bullet[1])$ where $C(A(\Gamma))$ is the category of chain complexes in $\text{Mod}(A(\Gamma))$. This is done by showing that the morphisms are linearly independent ([5, Lem. 4.3, Cor. 4.4]) and that any non-zero component of a morphism $f: P_M^\bullet \rightarrow P_N^\bullet[1]$ determines a graph, single or double map that is a summand of f ([5, Lem. 4.2]). Since P_M^\bullet and P_N^\bullet are bounded complexes of finite-dimensional modules, this is enough to prove that the graph, single and double maps also span $\text{Hom}_{C(A(\Gamma))}(P_M^\bullet, P_N^\bullet[1])$ (see [5, Prop. 4.1]).

Step 2:: In order to consider morphisms in $K(A(\Gamma))$, the next step is to consider the class $\mathcal{H}(f)$ of single, double or graph maps g such that f is homotopy equivalent to μg for some $0 \neq \mu \in K$ (see [5, Def. 4.6]). It is shown in [5, Prop. 4.8] that if $f: P_M^\bullet \rightarrow P_N^\bullet[1]$ is a single or double map such that $\mathcal{H}(f) \neq \{f\}$, then there is a quasi-graph map $P_M^\bullet \rightsquigarrow P_N^\bullet[1]$. The idea of the proof is to produce an algorithm that is determined by a given single or double map and produces the corresponding quasi-graph map. Since P_M^\bullet and P_N^\bullet are bounded complexes of finite-dimensional modules, the algorithm terminates after finitely many steps.

Step 3:: The final step is to show that if f is a graph map or a singleton single or double map, then $\mathcal{H}(f) = \{f\}$. This is done in [5, Lem. 4.12, Lem. 4.12]. □

Next we allow M and N to be infinite-dimensional string modules. The standard maps no longer necessarily form a basis, however we do obtain the following proposition.

Proposition A.9. *Let M and N be string modules or band modules with $n = 1$, such that M and N are not both band modules, and let P_M^\bullet and P_N^\bullet be the projective resolutions of M and N respectively. The following statements are equivalent.*

- (1) *The K -vector space $\text{Hom}_{K(A(\Gamma))}(P_M^\bullet, P_N^\bullet[1])$ is zero.*
- (2) *There exists a standard map $P_M^\bullet \rightarrow P_N^\bullet[1]$.*

Proof. In the following proof we will denote the arcs associated to M and N by α and δ respectively.


The idea of the proof is to extend to arguments of [5] to include projective resolutions of infinite-dimensional string modules. The proof is divided into three steps to reflect the three steps used to prove [5, Thm. 3.15] (see Proposition A.8). For each step it is necessary to apply the corresponding arguments in [5] and then extend them in the way that is described below.

Step 1:: First we show that the Proposition A.9 holds in the category $C(A(\Gamma))$. The arguments used to obtain [5, Lem. 4.2, Lem. 4.3, Cor. 4.4] may also be applied to projective resolutions of infinite-dimensional string modules. This yields that the graph, single and double maps $P_M^\bullet \rightarrow P_N^\bullet[1]$ are linearly independent in $C(A(\Gamma))$. By applying the same argument as the proof of [5, Prop. 4.1], we achieve Step 1 by ruling out the case where g is a ‘graph map’ induced by an infinite common substring that is not an antipath. Such a situation cannot arise in our setup since the degrees of any infinite common substring would not be compatible. Thus the proposition holds in the category $C(A(\Gamma))$.

Step 2:: If neither of M and N are finite-dimensional string modules and $f: P_M^\bullet \rightarrow P_N^\bullet[1]$ is a single or double map such that $\mathcal{H}(f) \neq \{f\}$, then the algorithm described in [5, Prop. 4.8] works in the same way but may not terminate. If the algorithm stops, then there is a quasi-graph map $g: P_N^\bullet \rightsquigarrow P_M^\bullet[1]$.

We show that if the algorithm does not terminate, then the morphism f is null-homotopic. By the same argument as [21, Lem. 3.5], the maximal common substring produced by the algorithm is supported in cohomological degrees -1 and 0 only. Then the algorithm described in the proof of [5, Prop. 4.8] yields the following two extra cases:


Case 1:



where the top unfolded diagram is the unfolded diagram of τ_α and the bottom one is the unfolded diagram of $\tau_{\alpha'}$. Moreover, there is no path p from $t(q'_m)$ to $t(q_n)$ such that

$q_n = pq'_m$. This is only possible if either δ is an asymptotic arc with $m = 0$ and q'_0 is trivial or there is a non-trivial path q such that $qq_n = q'_m$.

Case 2:



where, again, the top unfolded diagram is the unfolded diagram of τ_α and the bottom one is the unfolded diagram of τ_δ , and there is no path p from $s(p'_{m-1})$ to $s(p_{n-1})$ such that $p_{n-1}p = p'_{m-1}$. This is only possible if α is an asymptotic arc and $n = 0$ or there exists a non-trivial path q such that $p_{n-1} = p'_{m-1}q$.

If α and δ are both expanding asymptotic arcs or both contracting asymptotic arcs, then both cases 1 and 2 yield classes of null-homotopic maps.

Suppose that α and δ are not both expanding asymptotic arcs nor both contracting asymptotic arcs. We will show that we must have that α is a contracting asymptotic arc or δ is an expanding asymptotic arc; either scenario yields a class of null-homotopic maps. Note that the maximal common substring of τ_δ and τ_α must be the maximal periodic part of τ_α or τ_δ , otherwise they would be both contracting or both expanding asymptotic arcs. We will deal with case 1, case 2 is similar.

First we assume we have case 1 and suppose that $m \neq 0$. The maximal common substring of τ_δ and τ_α is $w := \dots q_{n+2}^{-1}p_{n+1}q_{n+1}^{-1}p_nq_n^{-1} = \dots (q'_{m+2})^{-1}p'_{m+1}(q'_{m+1})^{-1}p'_mq_n^{-1}$. If w is the maximal periodic part of τ_α , then it is also the maximal periodic part of u_α and α is a contracting asymptotic arc. If w is the maximal periodic part of τ_δ , then it is also the maximal periodic part of u_δ and δ is an expanding asymptotic arc. Now suppose we are in case 1 with δ an asymptotic arc and $m = 0$. We cannot have that $u_\delta = \dots p'_1(q'_1)^{-1}p'_0$ since, by Definition A.2, the existence of the last letter of q_n would imply that $\tau_\delta \neq u_\delta$. It must therefore be the case that $u_\delta = \dots p'_1(q'_1)^{-1}p'_0(q'_0)^{-1}$ where q'_0 is a path and there is no arrow a_0 such that $a_0q'_0$ is a string. That is, we must have that $q'_0 = rq_n$ for some (possibly trivial) path r . It follows that the maximal common substring of u_α and u_δ is $w := \dots p'_1(q'_1)^{-1}p'_0q_n^{-1} = \dots p_{n+1}q_{n+1}^{-1}p_nq_n^{-1}$. If r is non-trivial then we have already dealt with this case. If r is trivial, then u_δ is the maximal periodic substring of itself and must be expanding.

Step 3:: The arguments given in [5, Lem. 4.12, Lem. 4.12] also yield that if f is a graph map or a singleton single or double map, then $\mathcal{H}(f) = \{f\}$ if M and/or N is an infinite-dimensional string module.

□

A.2 Extensions in $\text{Mod}(A(\Gamma))$ and standard maps. Let M and N be string or band modules with $n = 1$ over $A(\Gamma)$ and let α and δ be their respective associated arcs. Assume that M and N are not both band modules. Let

$$\Phi: \text{Hom}_{K(A(\Gamma))}(P_M^\bullet, P_N^\bullet[1]) \rightarrow \text{Ext}_{A(\Gamma)}(M, N)$$

be the isomorphism described at the beginning of the appendix.

In [21], the authors show that if M and N are finite-dimensional, then Φ induces a bijection between the standard maps in $\text{Hom}_{K(A(\Gamma))}(P_M^\bullet, P_N^\bullet[1])$ and the standard extensions in $\text{Ext}_{A(\Gamma)}(M, N)$, which are defined in Proposition 2.20. The arguments used in [21, 22] do not depend on M and N being finite-dimensional and so we directly obtain the following lemma.

Lemma A.10. *Let M and N be string or band modules with $n = 1$ over $A(\Gamma)$ and let α and δ be their respective associated arcs. Assume that M and N are not both band modules. Then Φ*


induces a bijection as follows.

$$\left\{ \text{Standard maps in } \text{Hom}_{K(A(\Gamma))}(P_M^\bullet, P_N^\bullet[1]) \right\} \xleftrightarrow{1-1} \left\{ \text{Standard extensions in } \text{Ext}_{A(\Gamma)}^1(M, N) \right\}$$


In particular, there exists a standard extension $0 \rightarrow N \rightarrow E \rightarrow M \rightarrow 0$ in $\text{Mod}(A(\Gamma))$ if and only if $\text{Ext}_{A(\Gamma)}^1(M, N) \neq 0$.

A.3 Extensions in $\text{Mod}(A(\Gamma))$ and crossings of arcs. As before, let M and N be string or band modules with $n = 1$ such that M and N are not both band modules. We will denote the arc associated to M by α and the arc associated to N by δ . The next proposition extends some of the results of [23] to include infinite-dimensional string modules and band modules with $n = 1$.

The following observations are similar to the proof of [23, Prop. 3.3]. They essentially follow from the definitions. There exists an overlap extension $0 \rightarrow N \rightarrow E \rightarrow M \rightarrow 0$ if and only if we have the following local configuration.



There is an arrow extension $0 \rightarrow N \rightarrow E \rightarrow M \rightarrow 0$ if and only if we have the following local configuration.



The only remaining possible crossing between α and δ (up to homotopy and reordering of α and δ) corresponds to α and δ crossing in a 3-cycle.


Lemma A.11. *Let M and N be string or band modules with $n = 1$ such that M and N are not both band modules. Denote the arcs associated to M and N by α and δ respectively. The following statements are equivalent.*

- (1) *The K -vector spaces $\text{Ext}_{A(\Gamma)}^1(M, N)$ and $\text{Ext}_{A(\Gamma)}^1(N, M)$ are both zero.*
- (2) *The arcs α and δ do not cross outside of 3-cycles.*


Proof. By Lemma A.10, we have that the groups $\text{Ext}_{A(\Gamma)}^1(M, N)$ and $\text{Ext}_{A(\Gamma)}^1(N, M)$ are both trivial if and only if there are no standard extensions in $\text{Ext}_{A(\Gamma)}^1(M, N)$ or $\text{Ext}_{A(\Gamma)}^1(N, M)$. By the observations preceding this lemma, this arises exactly when α and δ do not cross outside of 3-cycles. \square

A.4 Band modules with $n \neq 1$. In this section we will show that, for a finite or asymptotic arc α , a scalar $\lambda \in K^*$ and $n \in \mathbb{N} \cup \{\infty, -\infty\}$, we have that the homomorphisms in $K(A(\Gamma))$ between the string complex P_α^\bullet and a band complex $B_{\lambda, n}^\bullet$ are determined by the homomorphisms between P_α^\bullet and $B_{\lambda, 1}^\bullet$. In order to do this we will follow the arguments of [5, Sec. 5].

Since K is algebraically closed, we associate the following unfolded diagram to the complexes $B_{\lambda, n}^\bullet$ with $n \in \mathbb{N}$ as in [5, Sec. 5].



In the same way we associate the following unfolded diagrams to $B_{\lambda,\infty}^\bullet$ and $B_{\lambda,-\infty}^\bullet$ respectively.



Definition A.12 (cf. [5, Def. 5.2]). Consider a finite or asymptotic arc α in (S, M) and let $\lambda \in K^*$, $n \in \mathbb{N} \cup \{\infty, -\infty\}$.

A morphism f in $\text{Hom}_{K(A(\Gamma))}(P_\alpha^\bullet, B_{\lambda,n}^\bullet[1])$ is said to be **lifted from the morphism g in $\text{Hom}_{K(A(\Gamma))}(P_\alpha^\bullet, B_{\lambda,1}^\bullet[1])$** if there exists an $m \in \mathbb{N}$ such that the components of f that have codomain in layer m of $B_{\lambda,n}^\bullet[1]$ are exactly the same as the components of g and f is the minimal such morphism.

A morphism f in $\text{Hom}_{K(A(\Gamma))}(B_{\lambda,n}^\bullet, P_\alpha^\bullet[1])$ is defined to be **lifted from the morphism g in $\text{Hom}_{K(A(\Gamma))}(B_{\lambda,1}^\bullet, P_\alpha^\bullet[1])$** analogously.

The following lemma is adapted from [5, Lem. 5.3] and the argument applies easily to the infinite-dimensional case.

Lemma A.13. *Let α be a finite or asymptotic arc in (S, M) and let $\lambda \in K^*$ and $n \in \mathbb{N} \cup \{\infty, -\infty\}$. Then the following statements hold.*

- (1) *If $f \neq 0$ is a morphism in $\text{Hom}_{K(A(\Gamma))}(P_\alpha^\bullet, B_{\lambda,n}^\bullet[1])$, then there exists a morphism g in $\text{Hom}_{K(A(\Gamma))}(P_\alpha^\bullet, B_{\lambda,n}^\bullet[1])$ and $\mu \in K^*$ such that $f = \mu g + f'$ and g is lifted from a non-zero standard map in $\text{Hom}_{K(A(\Gamma))}(P_\alpha^\bullet, B_{\lambda,1}^\bullet[1])$.*
- (2) *If $f \neq 0$ is a morphism in $\text{Hom}_{K(A(\Gamma))}(B_{\lambda,n}^\bullet, P_\alpha^\bullet[1])$, then there exists a morphism g in $\text{Hom}_{K(A(\Gamma))}(B_{\lambda,n}^\bullet, P_\alpha^\bullet[1])$ and $\mu \in K^*$ such that $f = \mu g + f'$ and g is lifted from a non-zero standard map in $\text{Hom}_{K(A(\Gamma))}(B_{\lambda,1}^\bullet, P_\alpha^\bullet[1])$.*

Proposition A.14. *Let α be a finite or asymptotic arc in (S, M) and let $\lambda \in K^*$ and $n \in \mathbb{N} \cup \{\infty, -\infty\}$. Then the following statements hold.*

- (1) $\text{Ext}_{A(\Gamma)}^1(M(\alpha), M_{\lambda,n}) = 0$ if and only if $\text{Ext}_{A(\Gamma)}^1(M\alpha, M_{\lambda,1}) = 0$.
- (2) $\text{Ext}_{A(\Gamma)}^1(M_{\lambda,n}, M(\alpha)) = 0$ if and only if $\text{Ext}_{A(\Gamma)}^1(M_{\lambda,1}, M(\alpha)) = 0$.

Proof. We prove (1), the proof of (2) is similar. By Proposition A.9, we have that the K -vector space $\text{Hom}_{K(A(\Gamma))}(P_\alpha^\bullet, B_{\lambda,1}^\bullet[1])$ is zero if and only if there are no standard maps in from P_α^\bullet to $B_{\lambda,1}^\bullet[1]$. By Lemma A.13, we also have that the K -vector space $\text{Hom}_{K(A(\Gamma))}(P_\alpha^\bullet, B_{\lambda,n}^\bullet[1])$ is zero if and only if there are no standard maps from P_α^\bullet to $B_{\lambda,1}^\bullet[1]$. The proposition follows immediately after observing the isomorphisms $\text{Ext}_{A(\Gamma)}^1(M, N) \cong \text{Hom}_{K(A(\Gamma))}(P_M^\bullet, P_N^\bullet[1])$ and $\text{Ext}_{A(\Gamma)}^1(N, M) \cong \text{Hom}_{K(A(\Gamma))}(P_N^\bullet, P_M^\bullet[1])$ given in Section A.1. \square

A.5 Proof of Theorem A.1. Following Sections A.1 to A.4, we are now ready to give a proof of the main theorem of the appendix.

Proof of Theorem A.1. If neither of M and N is isomorphic to $M_{\lambda,n}$ with $n \neq 1$, then the theorem holds by Lemma A.11. Without loss of generality, assume that $M \cong M_{\lambda,n}$ for $n \neq 1$. By Proposition A.14, we have that $\text{Ext}_{A(\Gamma)}^1(M, N) = 0 = \text{Ext}_{A(\Gamma)}^1(N, M)$ exactly when $\text{Ext}_{A(\Gamma)}^1(M_{\lambda,1}, N) = 0 = \text{Ext}_{A(\Gamma)}^1(N, M_{\lambda,1})$. Therefore we may apply Lemma A.11 again to conclude that this is the case exactly when $\alpha = \beta$ and δ do not cross outside of 3-cycles. \square

REFERENCES

- [1] Takahide Adachi, Osamu Iyama, and Idun Reiten. τ -tilting theory. *Compos. Math.*, 150(3):415–452, 2014.
- [2] Lidia Angeleri Hügel. On the abundance of silting modules. In *Surveys in representation theory of algebras*, volume 716 of *Contemp. Math.*, pages 1–23. Amer. Math. Soc., Providence, RI, 2018.
- [3] Lidia Angeleri Hügel, Frederik Marks, and Jorge Vitória. Silting modules. *Int. Math. Res. Not. IMRN*, 2016(4):1251–1284, 2016.
- [4] Lidia Angeleri Hügel, David Pospíšil, Jan Šťovíček, and Jan Trlifaj. Tilting, cotilting, and spectra of commutative Noetherian rings. *Trans. Amer. Math. Soc.*, 366(7):3487–3517, 2014.
- [5] Kirstin K. Arnesen, Rosanna Laking, and David Pauksztello. Morphisms between indecomposable complexes in the bounded derived category of a gentle algebra. *J. Algebra*, 467:1–46, 2016.
- [6] Ibrahim Assem, Thomas Brüstle, Gabrielle Charbonneau-Jodoin, and Pierre-Guy Plamondon. Gentle algebras arising from surface triangulations. *Algebra Number Theory*, 4(2):201–229, 2010.
- [7] Maurice Auslander. Functors and morphisms determined by objects. In *Representation theory of algebras (Proc. Conf., Temple Univ., Philadelphia, Pa., 1976)*, pages 1–244. Lecture Notes in Pure Appl. Math., Vol. 37. Dekker, New York, 1978.
- [8] Karin Baur, Aslak B. Buan, and Robert J. Marsh. Torsion pairs and rigid objects in tubes. *Algebr. Represent. Theory*, 17(2):565–591, 2014.
- [9] Karin Baur and Grégoire Dupont. Compactifying exchange graphs I: Annuli and tubes. *Ann. Comb.*, 18(3):383–396, 2014.
- [10] Karin Baur and Robert J. Marsh. A geometric model of tube categories. *J. Algebra*, 362:178–191, 2012.
- [11] Karin Baur and Sibylle Schroll. Higher extensions for gentle algebras. *arXiv:1906.05257*, 2019.
- [12] Karin Baur and Hermund André Torkildsen. A geometric interpretation of categories of type \tilde{A}_n and of morphisms in the infinite radical. *Algebras and Representation Theory*, 2019.
- [13] Silvana Bazzoni. Cotilting modules are pure-injective. *Proc. Amer. Math. Soc.*, 131(12):3665–3672, 2003.
- [14] Viktor Bekkert and Héctor. A. Merklen. Indecomposables in derived categories of gentle algebras. *Algebr. Represent. Theory*, 6(3):285–302, 2003.
- [15] Apostolos Beligiannis and Idun Reiten. Homological and homotopical aspects of torsion theories. *Mem. Amer. Math. Soc.*, 188(883):viii+207, 2007.
- [16] Raphael Bennett-Tennenhaus. *Functorial Filtrations for Semiperfect generalisations of Gentle Algebras*. PhD thesis, University of Leeds, 2017.
- [17] Simion Breaz and Flaviu Pop. Cosilting modules. *Algebr. Represent. Theory*, 20(5):1305–1321, 2017.
- [18] Aslak B. Buan and Henning Krause. Cotilting modules over tame hereditary algebras. *Pacific J. Math.*, 211(1):41–59, 2003.
- [19] Aslak B. Buan, Robert J. Marsh, and Idun Reiten. Cluster-tilted algebras. *Trans. Amer. Math. Soc.*, 359(1):323–332, 2007.
- [20] Michael C. R. Butler and Claus M. Ringel. Auslander-Reiten sequences with few middle terms and applications to string algebras. *Comm. Algebra*, 15(1-2):145–179, 1987.
- [21] İlke Çanakçı, David Pauksztello, and Sibylle Schroll. On extensions for gentle algebras. *arXiv:1707.06934v2*, 2017.
- [22] İlke Çanakçı, David Pauksztello, and Sibylle Schroll. Mapping cones in the bounded derived category of a gentle algebra. *J. Algebra*, 530:163–194, 2019.
- [23] İlke Çanakçı and Sibylle Schroll. Extensions in Jacobian algebras and cluster categories of marked surfaces. *Adv. Math.*, 313:1–49, 2017.
- [24] William Crawley-Boevey. Locally finitely presented additive categories. *Comm. Algebra*, 22(5):1641–1674, 1994.
- [25] William Crawley-Boevey. Infinite-dimensional modules in the representation theory of finite-dimensional algebras. In *Algebras and modules, I (Trondheim, 1996)*, volume 23 of *CMS Conf. Proc.*, pages 29–54. Amer. Math. Soc., Providence, RI, 1998.
- [26] Laurent Demonet, Osamu Iyama, Nathan Reading, Idun Reiten, and Hugh Thomas. Lattice theory of torsion classes. *arXiv:1711.01785*, Preprint, 2017.
- [27] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky. Quivers with potentials and their representations. I. Mutations. *Selecta Math. (N.S.)*, 14(1):59–119, 2008.
- [28] Edward R. Fisher. *Abelian structures. I*. Springer, Berlin, 1977.

- [29] Sergey Fomin, Michael Shapiro, and Dylan Thurston. Cluster algebras and triangulated surfaces. I. Cluster complexes. *Acta Math.*, 201(1):83–146, 2008.
- [30] Christof Geiß and Idun Reiten. Gentle algebras are Gorenstein. In *Representations of algebras and related topics*, volume 45 of *Fields Inst. Commun.*, pages 129–133. Amer. Math. Soc., Providence, RI, 2005.
- [31] Rüdiger Göbel and Jan Trlifaj. *Approximations and endomorphism algebras of modules*, volume 41 of *De Gruyter Expositions in Mathematics*. Walter de Gruyter GmbH & Co. KG, Berlin, 2006.
- [32] Sira Gratz, Thorsten Holm, and Peter Jørgensen. Cluster tilting subcategories and torsion pairs in Igusa-Todorov cluster categories of Dynkin type A_∞ . *Math. Z.*, 292(1-2):33–56, 2019.
- [33] Dieter Happel, Idun Reiten, and Sverre O. Smalø. Tilting in abelian categories and quasitilted algebras. *Mem. Amer. Math. Soc.*, 120(575):viii+ 88, 1996.
- [34] Thorsten Holm, Peter Jørgensen, and Martin Rubey. Ptolemy diagrams and torsion pairs in the cluster category of Dynkin type A_n . *J. Algebraic Combin.*, 34(3):507–523, 2011.
- [35] Thorsten Holm, Peter Jørgensen, and Martin Rubey. Ptolemy diagrams and torsion pairs in the cluster categories of Dynkin type D . *Adv. in Appl. Math.*, 51(5):583–605, 2013.
- [36] Thorsten Holm, Peter Jørgensen, and Martin Rubey. Torsion pairs in cluster tubes. *J. Algebraic Combin.*, 39(3):587–605, 2014.
- [37] Birge Huisgen-Zimmermann and Sverre O. Smalø. A homological bridge between finite- and infinite-dimensional representations of algebras. *Algebr. Represent. Theory*, 1(2):169–188, 1998.
- [38] Yasuo Iwanaga. On rings with finite self-injective dimension. *Comm. Algebra*, 7(4):393–414, 1979.
- [39] J. P. Jans. Some aspects of torsion. *Pacific J. Math.*, 15:1249–1259, 1965.
- [40] Christian U. Jensen and Helmut Lenzing. Model theory and representations of algebras. In *Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979)*, volume 832 of *Lecture Notes in Math.*, pages 302–310. Springer, Berlin, 1980.
- [41] Bernhard Keller. Calabi-Yau triangulated categories. In *Trends in representation theory of algebras and related topics*, EMS Ser. Congr. Rep., pages 467–489. Eur. Math. Soc., Zürich, 2008.
- [42] Henning Krause. Finitistic dimension and Ziegler spectrum. *Proc. Amer. Math. Soc.*, 126(4):983–987, 1998.
- [43] Akeel Ramadan Mehdi and Mike Prest. Almost dual pairs and definable classes of modules. *Comm. Algebra*, 43(4):1387–1397, 2015.
- [44] Puiman Ng. A characterisation of torsion theories in the cluster category of dynkin type a_∞ . *arXiv:1005.4364*, 2010.
- [45] Mike Prest. *Model theory and modules*, volume 130 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1988.
- [46] Mike Prest. *Purity, spectra and localisation*, volume 121 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2009.
- [47] Gena Puninski and Mike Prest. Ringel’s conjecture for domestic string algebras. *Math. Z.*, 282(1-2):61–77, 2016.
- [48] Claus M. Ringel. Some algebraically compact modules. I. In *Abelian groups and modules (Padova, 1994)*, volume 343 of *Math. Appl.*, pages 419–439. Kluwer Acad. Publ., Dordrecht, 1995.
- [49] Claus M. Ringel. The Ziegler spectrum of a tame hereditary algebra. *Colloq. Math.*, 76(1):105–115, 1998.
- [50] Jan Trlifaj. Ext and inverse limits. *Illinois J. Math.*, 47(1-2):529–538, 2003. Special issue in honor of Reinhold Baer (1902–1979).
- [51] Jan Trlifaj. Infinite dimensional tilting modules and cotorsion pairs. In *Handbook of tilting theory*, volume 332 of *London Math. Soc. Lecture Note Ser.*, pages 279–321. Cambridge Univ. Press, Cambridge, 2007.
- [52] Burkhard Wald and Josef Waschbüsch. Tame biserial algebras. *J. Algebra*, 95(2):480–500, 1985.
- [53] Matthias Warkentin. Fadenmoduln über \widetilde{A}_n und Cluster-Kombinatorik. *Diploma Thesis*, Universität Bonn, <http://nbn-resolving.de/urn:nbn:de:bsz:ch1-qucosa-94793>, 2008.
- [54] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.
- [55] Jie Zhang, Yu Zhou, and Bin Zhu. Cotorsion pairs in the cluster category of a marked surface. *J. Algebra*, 391:209–226, 2013.
- [56] Peiyu Zhang and Jiaqun Wei. Cosilting complexes and AIR-cotilting modules. *J. Algebra*, 491:1–31, 2017.
- [57] Martin Ziegler. Model theory of modules. *Ann. Pure Appl. Logic*, 26(2):149–213, 1984.

KARIN BAUR, SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS, LS2 9JT, UK
E-mail address: pmtkb@leeds.ac.uk
URL: <http://www1.maths.leeds.ac.uk/~pmtkb/>

ROSANNA LAKING, DIPARTIMENTO DI INFORMATICA - SETTORE DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI VERONA, STRADA LE GRAZIE 15 - CA’ VIGNAL, I-37134 VERONA, ITALY
E-mail address: rosanna.laking@univr.it
URL: <http://profs.scienze.univr.it/laking/>