

Elementary function representations for the moments of the Meyer-König and Zeller operators

Ulrich Abel

*Technische Hochschule Mittelhessen
Fachbereich MND
Wilhelm-Leuschner-Straße 13, 61169 Friedberg,
Germany
Ulrich.Abel@mnd.thm.de*

Abstract.

We calculate the moments of the Meyer-König and Zeller operators in terms of elementary functions and polylogarithms.

Mathematics Subject Classification (2010): 33C05, 33E99, 41A36.

Keywords: Moments of Meyer-König and Zeller operators, elementary functions, polylogarithm.

1 Introduction

The operators of Meyer-König and Zeller [6] in the slight modification of Cheney and Sharma [4], defined by

$$(M_n f)(x) = (1-x)^{n+1} \sum_{\nu=0}^{\infty} \binom{\nu+n}{\nu} f\left(\frac{\nu}{\nu+n}\right) x^{\nu} \quad (0 \leq x < 1), \quad (1)$$

(also called Bernstein power series, or briefly MKZ operators) were the object of several investigations in approximation theory.

Let $e_r(x) = x^r$ denote the monomials. The moments of the MKZ operators are given by $(M_n e_r)(x)$. They are starting point for several studies of their approximation properties.

In 1984, J. A. H. Alkemade [3] found a representation of the second moment

$$(M_n e_2)(x) = x^2 + \frac{x(1-x)^2}{n+1} {}_2F_1(1, 2; n+2; x) \quad (2)$$

in terms of hypergeometric functions. The way of deriving this expression is based upon a differential equation which is satisfied by the right-hand side of (1). Alkemade wrote: “For the MKZ operators, an explicit expression for $(M_n e_2)(x)$ does not yet occur in the literature.” [3, Page 262]. Gavrea and Ivan [5, Theorem 2.2, Eq. (2.3)] observed that the second moment is an elementary function and found an explicit representation of $(M_n e_2)(x)$ in terms of the logarithm and rational functions.

Entering ${}_2F_1(1, 2; n+2; x)$ into a computer algebra software (Wolfram Mathematica Version 9), for small values of the integer variable n (up to $n = 9$), one obtains explicit expressions of the form

$$\sum_{k=1}^n \frac{a_k}{x^k} + \log(1-x) \sum_{k=2}^{n+1} \frac{b_k}{x^k}.$$

The command **Simplify** reveals the second sum to be a multiple of $(1-x)^{n+1} x^{-n-1}$. This is by no means surprising:

$${}_2F_1(1, 2; n+2; x) = (n+1)! \sum_{k=0}^{\infty} \frac{(k+1)!}{(n+k+1)!} x^k.$$

Now $\sum_{k=0}^{\infty} \frac{(k+1)!}{(n+k+1)!} x^k$ is a n -th antiderivative of $\sum_{k=0}^{\infty} x^{k+1} = \frac{1}{1-x} - 1$. Therefore, it can be expected that ${}_2F_1(1, 2; n+2; x)$ has a representation in terms of logarithmic functions $\log(1-x)$ and rational functions.

The purpose of this note is to provide a short and direct proof of this fact. It follows by a more general representation for arbitrary MKZ moments in terms of polylogarithmic functions. The proof uses only the definition (1) of the MKZ operators. We mention that an integral representation for $(M_n e_r)(x)$ can be found in [1]. It is useful when studying the asymptotic properties of the MKZ operators $(M_n f)(x)$ for smooth functions f as n tends to infinity (see [2]).

2 Results

For $s = 0, 1, 2, \dots$, put $g_s(x) = (1-x)^s$. Obviously, $M_n g_0 = e_0$. For $s, n \in \mathbb{N}$, we obtain

$$\begin{aligned} (M_n g_s)(x) &= (1-x)^{n+1} \sum_{\nu=0}^{\infty} \binom{\nu+n}{\nu} \left(\frac{n}{\nu+n} \right)^s x^\nu \\ &= \frac{n^s}{n!} (1-x)^{n+1} \sum_{\nu=0}^{\infty} (\nu+n) \cdots (\nu+2)(\nu+1) \frac{x^\nu}{(\nu+n)^s} \\ &= \frac{n^s}{n!} (1-x)^{n+1} \left(\sum_{\nu=0}^{\infty} \frac{x^{\nu+n}}{(\nu+n)^s} \right)^{(n)} \end{aligned}$$

$$= \frac{n^s}{n!} (1-x)^{n+1} \text{Li}_s^{(n)}(x),$$

where

$$\text{Li}_s(x) = \sum_{\nu=1}^{\infty} \frac{x^\nu}{\nu^s} \quad (|x| < 1)$$

is the polylogarithm. Using $e_r = \sum_{s=0}^r (-1)^s \binom{r}{s} g_s$ we obtain the following result.

Theorem 1 *For $r, n \in \mathbb{N}$ and $|x| < 1$, the moments of the MKZ operators have the representation*

$$(M_n e_r)(x) = 1 + \frac{1}{n!} (1-x)^{n+1} \sum_{s=1}^r (-1)^s \binom{r}{s} n^s \text{Li}_s^{(n)}(x).$$

Applying the well-known (elementary) relation $(d/dx)\text{Li}_s(x) = x^{-1}\text{Li}_{s-1}(x)$, for $s \geq 2$, one can reduce the terms $\text{Li}_s^{(n)}(x)$ to certain derivatives of $\text{Li}_1(x)$. Note that $\text{Li}_1(x) = -\log(1-x)$ is an elementary function. As an immediate consequence all moments of the MKZ operators can be represented in terms of polylogarithms and rational functions. In particular, we have, for $n \in \mathbb{N}$,

$$\begin{aligned} \text{Li}_1^{(n)}(x) &= \frac{(n-1)!}{(1-x)^n}, \\ \text{Li}_2^{(n)}(x) &= (x^{-1}\text{Li}_1(x))^{(n-1)} = (-1)^n \frac{(n-1)!}{x^n} \log(1-x) \\ &\quad + \sum_{k=1}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} \frac{(n-k-1)!}{x^{n-k}} \frac{(k-1)!}{(1-x)^k} \\ &= (-1)^n \frac{(n-1)!}{x^n} \log(1-x) + \frac{(n-1)!}{x^n} \sum_{k=1}^{n-1} (-1)^{n-1-k} \frac{1}{k} \left(\frac{x}{1-x} \right)^k. \end{aligned}$$

In the special case $r = 1$ we recover the well-known fact that MKZ operators preserve linear functions:

$$(M_n e_1)(x) = 1 + \frac{1}{n!} (1-x)^{n+1} \text{Li}_1^{(n)}(x) = 1 - (1-x) = x.$$

In the special case $r = 2$ we obtain

$$(M_n e_2)(x) = 1 + \frac{1}{n!} (1-x)^{n+1} \left(-rn \text{Li}_1^{(n)}(x) + n^2 \text{Li}_2^{(n)}(x) \right)$$

A short calculation leads to the following formula which is comparable to [5, Theorem 2.2, Eq. (2.3)].

Corollary 2 *For $n \in \mathbb{N}$ and $|x| < 1$, the second moment of the MKZ operators has the representation*

$$\begin{aligned} (M_n e_2)(x) &= (-1)^n \frac{n(1-x)^{n+1}}{x^n} \log(1-x) \\ &\quad + 2x - 1 + n(1-x) \sum_{k=1}^{n-1} (-1)^{n-1-k} \frac{1}{k} \left(\frac{1-x}{x} \right)^{n-k}. \end{aligned}$$

References

- [1] Ulrich Abel, The moments for the Meyer-König and Zeller operators, *J. Approx. Theory* **82** (1995), 352–361.
- [2] Ulrich Abel, The complete asymptotic expansion for the Meyer-König and Zeller operators, *J. Math. Anal. Appl.* **208** (1997), 109–119.
- [3] J. A. H. Alkemade, The second moment for the Meyer-König and Zeller operators, *J. Approx. Theory* **40** (1984), 261–273.
- [4] E. W. Cheney and A. Sharma, Bernstein power series, *Canad. J. Math.* **16** (1964), 241–253.
- [5] Ioan Gavrea and Mircea Ivan, An elementary function representation of the second-order moment of the Meyer-König and Zeller operators, *Mediterr. J. Math.* **15** (2018), No. 1, Paper No. 20, 8 p.
- [6] W. Meyer-König and K. Zeller, Bernsteinsche Potenzreihen, *Studia Math.* **19** (1960), 89–94.