

INFINITE DIMENSIONAL SYSTEMS OF PARTICLES WITH INTERACTIONS GIVEN BY DUNKL OPERATORS

ANDREI VELICU

ABSTRACT. Firstly we consider a finite dimensional Markov semigroup generated by Dunkl laplacian with drift terms. Using gradient bounds we show that for small coefficients this semigroup has an invariant measure. We then extend this analysis to an infinite dimensional semigroup on $(\mathbb{R}^N)^{\mathbb{Z}^d}$ which we construct using gradient bounds, and finally we study the existence of invariant measures and ergodicity properties.

1. INTRODUCTION

In this paper we study infinite dimensional models of interacting particle systems generated by Dunkl operators. More precisely, we study the Markov semigroup on the infinite dimensional space $(\mathbb{R}^N)^{\mathbb{Z}^d}$ generated by

$$\mathcal{L} := \sum_{l \in \mathbb{Z}^d} (L^{(l)} + L_1^{(l)}).$$

The operators $L^{(l)}$ and $L_1^{(l)}$ are defined by

$$L^{(l)} = \Delta_k^{(l)} + b^{(l)} \cdot \nabla_k^{(l)},$$

and

$$L_1^{(l)} = e^{(l)} \cdot \nabla_k^{(l)},$$

where $\Delta_k^{(l)}$ and $\nabla_k^{(l)}$ are the usual Dunkl laplacian and gradient on $\mathbb{R}^N$ which act only on the l component of $\omega = (\omega_l)_{l \in \mathbb{Z}^d} \in (\mathbb{R}^N)^{\mathbb{Z}^d}$. The functions $b^{(l)} : (\mathbb{R}^N)^{\mathbb{Z}^d} \rightarrow \mathbb{R}^N$ depend only on the ω_l component, while $e^{(l)} : (\mathbb{R}^N)^{\mathbb{Z}^d} \rightarrow \mathbb{R}^N$ depend on the components ω_j with $|j - l| < R$ for a finite R called the range of interactions. In other words, the operators $L^{(l)}$ are operators that only depend on the ω_l component and so they define finite dimensional models. In the absence of the interaction terms $e^{(l)}$, these evolve independently of each other.

Dunkl operators are differential-difference operators defined in terms of a root system. They were introduced in [4] to study special functions associated to root systems, but have since been studied extensively and have found applications outside harmonic analysis. One typical application is in mathematical physics where they were used to study quantum many body problems associated to Calogero-Moser-Sutherland models; see [12] for more details. In probability theory, Dunkl operators have been used to define Markov processes with remarkable properties; for an overview of probabilistic aspects of Dunkl theory see [5].

2010 *Mathematics Subject Classification.* 60J35, 37A25, 47D07, 60K35, 82C22.

Key words and phrases. Infinite dimensional Markov semigroups, Gradient bounds, Ergodicity.

To tackle the problem introduced above, we first study the finite dimensional case of a single operator $L^{(l)}$ acting on $\mathbb{R}^N$, for which a priori the existence of an invariant measure is unknown. To begin with, we prove gradient bounds for the resulting semigroup. An essential trick here is to consider a symmetrised gradient which takes into account the reflections associated to the root system defining Dunkl operators. This allows us to deal with the resulting reflection terms and to obtain appropriate bounds. Using these bounds we then prove the existence of an invariant measure for the semigroup in the case of small coefficients k_α .

This analysis is then extended to an infinite dimensional setting, by introducing interaction terms. In this case, corresponding to the description above, the construction of the semigroup is not obvious and it requires an approximation procedure. This requires gradient bounds similar to the finite dimensional case, but which are technically more involved. The bounds obtained allow us to prove that the semigroup has an invariant measure and study its ergodicity properties.

Dissipative dynamics for infinite systems of interacting particles has been studied a long time; see the classic references [9] and [6] for more details. For a study of Markov semigroups in infinite dimensional settings see [15] in the case of elliptic operators, [3] for subelliptic operators, [13, 14] for Levy-type operators and [8] for operators of hypocoercive type.

Similar analysis for Dunkl-type operators in the infinite dimensional setting but in a simplified context corresponding to the very particular case of a root system A_1 , was studied in [16]. In this paper we extend these results to the case of general root systems in all dimensions.

The structure of the paper is as follows. In section 2 we introduce Dunkl operators and basic results that will be used later in the paper. In section 3 we deal with the finite dimensional case, and in section 3.2 we present the infinite dimensional case.

2. PRELIMINARIES

In this section we present a very quick introduction to Dunkl theory. For more details see the survey papers [10] and [1].

We define a root system to be a finite set $R \subset \mathbb{R}^N \setminus \{0\}$ such that $R \cap \alpha\mathbb{R} = \{-\alpha, \alpha\}$ and $\sigma_\alpha(R) = R$ for all $\alpha \in R$. Here σ_α is the reflection in the hyperplane orthogonal to the root α , i.e.,

$$\sigma_\alpha x = x - 2 \frac{\langle \alpha, x \rangle}{\langle \alpha, \alpha \rangle} \alpha.$$

The group generated by all the reflections σ_α for $\alpha \in R$ is a finite group, and we denote it by G .

An invariant function is a map $k : R \rightarrow [0, \infty)$, denoted $\alpha \mapsto k_\alpha$, such that $k(\alpha) = k(g\alpha)$ for all $g \in G$ and all $\alpha \in R$. It is possible to write the root system R as a disjoint union $R = R_+ \cup (-R_+)$ such that R_+ and $-R_+$ lie on different sides of a hyperplane through the origin; in this case, we call R_+ a positive subsystem. This decomposition is not unique, but the particular choice of positive subsystem does not make a difference in the definitions below because of the G -invariance of the coefficients k .

From now on we fix a root system in $\mathbb{R}^N$ with positive subsystem R_+ , and a multiplicity function k . We also assume without loss of generality that $|\alpha|^2 = 2$ for all $\alpha \in R$. For

$i = 1, \dots, N$ we define the Dunkl operator on $C^1(\mathbb{R}^N)$ by

$$T_i f(x) = \partial_i f(x) + \sum_{\alpha \in R_+} k_\alpha \alpha_i \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle}.$$

An important result, due to Dunkl [4], is that Dunkl operators commute, i.e.,

$$T_i T_j = T_j T_i \quad \forall 1 \leq i, j \leq N.$$

We will denote by $\nabla_k = (T_1, \dots, T_N)$ the Dunkl gradient, and $\Delta_k = \sum_{i=1}^N T_i^2$ will denote the Dunkl laplacian. Note that for $k = 0$ Dunkl operators reduce to partial derivatives, and $\nabla_0 = \nabla$ and $\Delta_0 = \Delta$ are the usual gradient and laplacian. The Dunkl laplacian can be expressed in terms of the usual gradient and laplacian using the following formula:

$$(2.1) \quad \Delta_k f(x) = \Delta f(x) + 2 \sum_{\alpha \in R_+} k_\alpha \left[\frac{\langle \nabla f(x), \alpha \rangle}{\langle \alpha, x \rangle} - \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle^2} \right].$$

The following representation formula

$$(2.2) \quad \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle} = \int_0^1 \alpha \cdot \nabla f(x - t\langle \alpha, x \rangle \alpha) dt,$$

which holds for all roots $\alpha \in R$, shows that Dunkl operators leave invariant classical spaces of functions, for example $\mathcal{S}(\mathbb{R}^N)$, the space of Schwarz functions, or $C_c^\infty(\mathbb{R}^N)$, the space of smooth compactly supported functions.

The natural spaces to study Dunkl operators are the weighted $L^p(\mu_k)$, where $d\mu_k = w_k(x) dx$ and the weight function is defined as

$$w_k(x) = \prod_{\alpha \in R_+} |\langle \alpha, x \rangle|^{2k_\alpha}.$$

This is a homogeneous function of degree

$$\gamma := \sum_{\alpha \in R_+} k_\alpha.$$

We will simply write $\|\cdot\|_p$ for the norm in $L^p(\mu_k)$. With respect to this weighted measure we have the integration by parts formula

$$\int_{\mathbb{R}^N} T_i(f)g d\mu_k = - \int_{\mathbb{R}^N} f T_i(g) d\mu_k.$$

One of the main differences between Dunkl operators and classical partial derivatives is that the Leibniz rule does not hold in general. Instead, we have the following.

Lemma 2.1. *If one of the functions f, g is G -invariant, then we have the Leibniz rule*

$$T_i(fg) = f T_i g + g T_i f.$$

In general, we have

$$T_i(fg)(x) = T_i f(x)g(x) + f(x)T_i g(x) - \sum_{\alpha \in R_+} k_\alpha \alpha_i \frac{(f(x) - f(\sigma_\alpha x))(g(x) - g(\sigma_\alpha x))}{\langle \alpha, x \rangle}.$$

The theory of Dunkl operators is further enriched by the existence of an intertwining operator, which connects Dunkl operators to partial derivatives, and by the construction of the Dunkl kernel, which acts as a generalisation of the classical exponential function. Using these tools, it is then possible to define a Dunkl transform, which generalises the classical Fourier transform, with which it shares many important properties. The methods in this paper do not make any use of these notions, so we will not go into further details here; a more complete account can be found in the review papers recommended at the beginning of this section.

3. THE FINITE DIMENSIONAL CASE

We consider the operator

$$(3.1) \quad L := \sum_{i=1}^N (T_i^2 + b_i T_i) = \Delta_k + b \cdot \nabla_k,$$

where each $b_i \in C^1(\mathbb{R}^N)$ for $i = 1, \dots, N$. We will assume throughout this section that L generates a Markov semigroup on $C_b(\mathbb{R}^N)$, the space of bounded continuous functions on $\mathbb{R}^N$. Let $P_t := e^{tL}$ be the semigroup generated by L and denote for brevity $f_t = P_t f$. The carré-du-champ operator associated with the operator L is given by

$$\Gamma_L(f) := \frac{1}{2} (L(f^2) - 2fLf).$$

This should not be confused with the carré-du-champ operator associated with the Dunkl laplacian

$$\Gamma(f) := \frac{1}{2} (\Delta_k(f^2) - 2f\Delta_k f),$$

which can be computed using the following lemma.

Lemma 3.1. *We have*

$$\Gamma(f) = |\nabla f|^2 + \sum_{\alpha \in R_+} k_\alpha \left(\frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle} \right)^2.$$

Proof. Using (2.1), we have

$$\begin{aligned} \Delta_k(f^2) &= \Delta(f^2) + 2 \sum_{\alpha \in R_+} k_\alpha \left(\frac{\langle \nabla(f^2), \alpha \rangle}{\langle \alpha, x \rangle} - \frac{f^2(x) - f^2(\sigma_\alpha x)}{\langle \alpha, x \rangle^2} \right) \\ &= 2f\Delta_k f + 2|\nabla f|^2 + 4f(x) \sum_{\alpha \in R_+} k_\alpha \left(\frac{\langle \nabla f, \alpha \rangle}{\langle \alpha, x \rangle} - \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle^2} \right) \\ &\quad + 2 \sum_{\alpha \in R_+} k_\alpha \left(\frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle} \right)^2 \\ &= 2f\Delta_k f + 2|\nabla f|^2 + 2 \sum_{\alpha \in R_+} k_\alpha \left(\frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle} \right)^2. \end{aligned}$$

The expression for $\Gamma(f)$ then follows immediately from this and the definition. $\square$

In order to avoid long expressions, in this paper we use the (non-standard) notation

$$A_\alpha f(x) := \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle}$$

for the α -dependent difference part in the definition of the Dunkl operators.

Example 3.2. The main example throughout this section is the linear case corresponding to $b(x) = -cx$ for some $c > 0$. This case is related to the generalised Ornstein-Uhlenbeck semigroup studied in [11], where it was shown that the operator

$$L_{OU} := \Delta_k - cx \cdot \nabla f$$

generates a Markov semigroup on $C_b(\mathbb{R}^N)$.

We note that our operator is related to L_{OU} via

$$Lf = L_{OU}f - c \sum_{\alpha \in R_+} k_\alpha (f - f \circ \sigma_\alpha).$$

Since the linear operator $f \mapsto -c \sum_{\alpha \in R_+} k_\alpha (f - f \circ \sigma_\alpha)$ is clearly bounded on $C_b(\mathbb{R}^N)$, stan-

dard results from general semigroup theory imply that L generates a contraction semigroup $(P_t)_{t \geq 0}$ (the perturbation of a contraction semigroup generator by a bounded operator generates a contraction semigroup, see for example [2, Theorem 3.1]).

It is easy to see that the assumption (3.2) that we impose below on the drifts b will assure the positive maximum principle holds and thus $(P_t)_{t \geq 0}$ is indeed positivity-preserving, so a Markov semigroup (see [7, Theorem 19.11]).

Moreover, adding suitably small perturbations to the linear drifts $b(x) = -cx$ should produce further Markov semigroups with generators of the desired form (3.1).

Remark. We expect that the class of functions b for which the operator L generates a Markov semigroup is much larger but this is the subject of a different investigation.

3.1. Gradient bounds. The main result of this section is a bound on the symmetrised gradient form

$$\tilde{\Gamma}(f)(x) := \sum_{g \in G} |\nabla_k f(gx)|^2.$$

This form of the gradient has the advantage that it is G -invariant, so we do not have to deal with the reflected terms that will appear in the computations below.

Theorem 3.3. *Let $b_i \in C^1(\mathbb{R}^N)$ such that $\nabla(b_i)$ is bounded for all $i = 1, \dots, N$ and which satisfy the assumptions*

$$(3.2) \quad \frac{2}{\langle \alpha, x \rangle^2} - \langle b(x), \alpha \rangle \frac{1}{\langle \alpha, x \rangle} \geq 0 \quad \forall x \in \mathbb{R}^N, \forall \alpha \in R_+$$

and

$$(3.3) \quad b(gx) = gb(x) \quad \forall g \in G, \forall x \in \mathbb{R}^N.$$

Consider the operator

$$L = \Delta_k + b(x) \cdot \nabla_k$$

which is assumed to generate a Markov semigroup $(P_t)_{t \geq 0}$ on $C_b(\mathbb{R}^N)$.

We then have the bound

$$(3.4) \quad \tilde{\Gamma}(P_t f) \leq e^{2\eta t} P_t(\tilde{\Gamma}(f)) \quad \forall t > 0,$$

where

$$\eta := \max_i \sup_{x \in \mathbb{R}^N} \partial_i(b_i)(x) + (N-1) \max_{i \neq j} \|\partial_j(b_i)\|_\infty + \sqrt{2}\gamma \max_{\alpha \in R_+} \|A_\alpha(b)\|_\infty.$$

Proof. Firstly, let $g \in G$ and we compute

$$\begin{aligned} \frac{d}{ds} P_{t-s}(|\nabla_k f_s|^2 \circ g) &= \sum_{j=1}^N \frac{d}{ds} P_{t-s}((T_j f_s)^2 \circ g) \\ &= \sum_{j=1}^N P_{t-s}(-L((T_j f_s) \circ g)^2) + 2(T_j f_s) \circ g \cdot (T_j L f_s) \circ g \\ &= P_{t-s} \left(-2 \sum_{j=1}^N \Gamma_L((T_j f_s) \circ g) + I(f_s) \circ g + 2 \sum_{j=1}^N (T_j f_s) \circ g \cdot J_g(T_j f_s) \right), \end{aligned}$$

where

$$I(h) := 2\nabla_k h \cdot [\nabla_k, L]h = 2 \sum_{i=1}^N \nabla_k h \cdot [\nabla_k, b_i T_i]h$$

and

$$J_g(h) := (Lh) \circ g - L(h \circ g).$$

Here and everywhere in the below the notation $[A, B]$ stands for the commutator of the operators A and B , i.e., $[A, B] := AB - BA$.

Step 1: compute $\Gamma_L(h)$. We have

$$\begin{aligned} \Gamma_L(h) &= \frac{1}{2} \left(\Delta_k(h^2) - 2h\Delta_k h + \sum_{i=1}^N b_i [T_i(h^2) - 2hT_i h] \right) \\ &= \Gamma(h) + \frac{1}{2} \sum_{i=1}^N b_i [T_i(h^2) - 2hT_i h]. \end{aligned}$$

Using Lemmas 3.1 and 2.1, we have

$$\begin{aligned} \Gamma_L(h)(x) &= |\nabla h(x)|^2 + \sum_{\alpha \in R_+} k_\alpha \left(\frac{h(x) - h(\sigma_\alpha x)}{\langle \alpha, x \rangle} \right)^2 - \frac{1}{2} \sum_{i=1}^N \sum_{\alpha \in R_+} k_\alpha \alpha_i b_i(x) \frac{(h(x) - h(\sigma_\alpha x))^2}{\langle \alpha, x \rangle} \\ &= |\nabla h(x)|^2 + \frac{1}{2} \sum_{\alpha \in R_+} k_\alpha (h(x) - h(\sigma_\alpha x))^2 \left(\frac{2}{\langle \alpha, x \rangle^2} - \langle \alpha, b(x) \rangle \frac{1}{\langle \alpha, x \rangle} \right). \end{aligned}$$

Thus, assumption (3.2) assures that $\Gamma_L(h) \geq 0$.

Step 2: compute $J_g(h)$. It will be useful here to see $g \in G$ as a matrix, and recall that $gg^T = g^T g = I$. A simple computation shows that

$$(3.5) \quad \nabla(h \circ g) = g^T(\nabla h) \circ g.$$

Also, note that

$$g\sigma_\alpha g^T(x) = x - \langle \alpha, g^T x \rangle g\alpha = \sigma_{g\alpha}(x),$$

so

$$A_\alpha(h \circ g)(x) = \frac{h(gx) - h(g\sigma_\alpha x)}{\langle \alpha, x \rangle} = \frac{h(gx) - h(\sigma_{g\alpha}(gx))}{\langle g\alpha, gx \rangle} = A_{g\alpha}(h)(gx).$$

Thus, since by G -invariance we have $k_\alpha = k_{g\alpha}$, we obtain

$$(3.6) \quad \sum_{\alpha \in R_+} k_\alpha \alpha A_\alpha(h \circ g) = g^T \sum_{\alpha \in R_+} k_{g\alpha} g\alpha A_{g\alpha}(h) \circ g = g^T \sum_{\alpha \in R_+} k_\alpha \alpha A_\alpha(h) \circ g,$$

where in the last step we simply used a change of variables. Finally, from (3.5) and (3.6), we obtain

$$(3.7) \quad \nabla_k(h \circ g) = g^T \nabla_k(h) \circ g.$$

Using (3.7), we can now compute

$$\Delta_k(h \circ g) = \Delta_k(h) \circ g,$$

and also

$$J_g(h)(x) = b(gx) \cdot (\nabla_k h) \circ g - b(x) \cdot g^T (\nabla_k h) \circ g = (b(gx) - gb(x)) \cdot (\nabla_k h) \circ g = 0,$$

by our assumption (3.3).

Step 3: estimate $I(h)$. We first note that, using the commutativity of Dunkl operators,

$$[T_j, b_i T_i] = [T_j, b_i] T_i + b_i [T_j, T_i] = [T_j, b_i] T_i,$$

and

$$\begin{aligned} [T_j, b_i] T_i h(x) &= T_j(b_i T_i h)(x) - b_i(x) T_j T_i h(x) \\ &= T_j(b_i)(x) T_i h(x) - \sum_{\alpha \in R_+} k_\alpha \alpha_j \frac{(b_i(x) - b_i(\sigma_\alpha x))(T_i h(x) - T_i h(\sigma_\alpha x))}{\langle \alpha, x \rangle}. \end{aligned}$$

Thus we have

$$\begin{aligned} I(h) &= 2 \sum_{i,j=1}^N T_j h [T_j, b_i] T_i h \\ &= 2 \sum_{i,j=1}^N \left(T_j(b_i) T_j h T_i h - \sum_{\alpha \in R_+} k_\alpha \alpha_j A_\alpha(b_i) T_j h (T_i h - (T_i h) \circ \sigma_\alpha) \right) \\ &= 2 \sum_{i,j=1}^N \left(\partial_j(b_i) T_j h T_i h + \sum_{\alpha \in R_+} k_\alpha \alpha_j A_\alpha(b_i) T_j h (T_i h) \circ \sigma_\alpha \right) \\ &= 2 \sum_{i=1}^N \partial_i(b_i) |T_i h|^2 + 2 \sum_{i \neq j} \partial_j(b_i) T_i h T_j h + 2 \sum_{i,j=1}^N \sum_{\alpha \in R_+} k_\alpha \alpha_j A_\alpha(b_i) T_j h (T_i h) \circ \sigma_\alpha. \end{aligned}$$

Here, in the last step, we simply isolated the terms containing second powers $|T_i h|^2$, which will be necessary in obtaining gradient bounds. To obtain similar second powers from the

mixed terms, we use the basic inequality $2xy \leq x^2 + y^2$. This gives, for the second sum,

$$\begin{aligned} 2 \sum_{i \neq j} \partial_j(b_i) T_i h T_j h &\leq \sum_{i \neq j} \|\partial_j(b_i)\|_\infty [(T_i h)^2 + (T_j h)^2] \\ &\leq 2(N-1) \max_{i \neq j} \|\partial_j(b_i)\|_\infty |\nabla_k h|^2, \end{aligned}$$

and for the second sum, recalling that $|\alpha|^2 = 2$, we obtain

$$\begin{aligned} 2 \sum_{i,j=1}^N \sum_{\alpha \in R_+} k_\alpha \alpha_j A_\alpha(b_i) T_j h (T_i h) \circ \sigma_\alpha &= 2 \sum_{\alpha \in R_+} k_\alpha \langle \alpha, \nabla_k h \rangle \langle A_\alpha(b), (\nabla_k h) \circ \sigma_\alpha \rangle \\ &\leq \sqrt{2} \max_{\alpha \in R_+} \|A_\alpha(b)\|_\infty \sum_{\alpha \in R_+} k_\alpha [|\nabla_k h|^2 + |(\nabla_k h) \circ \sigma_\alpha|^2] \end{aligned}$$

Step 4: gradient bounds. From the previous three steps, summing up the computations over all $g \in G$, we obtain

$$\begin{aligned} \frac{d}{ds} P_{t-s}(\tilde{\Gamma}(f_s)) &\leq \sum_{g \in G} P_{t-s}(I(f_s) \circ g) \\ &\leq 2 \left(\max_i \sup_{x \in \mathbb{R}^N} \partial_i(b_i)(x) + (N-1) \max_{i \neq j} \|\partial_j(b_i)\|_\infty + \sqrt{2} \gamma \max_{\alpha \in R_+} \|A_\alpha(b)\|_\infty \right) P_{t-s}(\tilde{\Gamma}(f_s)). \end{aligned}$$

So we have proved that

$$\frac{d}{ds} P_{t-s}(\tilde{\Gamma}(f_s)) \leq 2\eta P_{t-s}(\tilde{\Gamma}(f_s)),$$

where

$$\eta := \max_i \sup_{x \in \mathbb{R}^N} \partial_i(b_i)(x) + (N-1) \max_{i \neq j} \|\partial_j(b_i)\|_\infty + \sqrt{2} \gamma \max_{\alpha \in R_+} \|A_\alpha(b)\|_\infty.$$

Integrating this inequality, we obtain

$$\tilde{\Gamma}(P_t f) \leq e^{2\eta t} P_t(\tilde{\Gamma}(f)),$$

as required. $\square$

Example 3.4. If $b(x) = -cx$, for a constant $c > 0$, then it is clear that it satisfies the assumption (3.2) of the Theorem. Moreover, $\partial_i(b_j) = -c\delta_{ij}$ and

$$A_\alpha(b_i)(x) = c \frac{-x_i + (x - \langle \alpha, x \rangle \alpha)_i}{\langle \alpha, x \rangle} = -c\alpha_i,$$

so

$$\eta = -c + 0 + \sqrt{2} \gamma \max_{\alpha} |c\alpha| = -c + 2c\gamma.$$

Thus, if $\gamma < \frac{1}{2}$, the inequality (3.4) proved above is in fact a coercive bound.

Remark. It is possible to extend the previous example of linear drifts by adding small perturbations and such that the conditions of the previous Theorem still hold.

3.2. Invariant measures. The strategy for proving existence of invariant measure is to use general results from probability theory such as the Prokhorov and Krylov-Bogoliubov theorems. A useful tool in proving the tightness conditions in these theorems is that of a Lyapunov function.

Definition 3.5. We say that a smooth function $\rho : \mathbb{R}^N \rightarrow [0, \infty]$ is a **Lyapunov function** for L if it satisfies the conditions:

- (i) $\rho^{-1}([0, \infty)) \neq \emptyset$;
- (ii) for any $M > 0$, the level set $\{x \in \mathbb{R}^N : \rho(x) \leq M\}$ is compact;
- (iii) there exist positive constants C_1, C_2 , with $C_2 \neq 0$, such that

$$(3.8) \quad L\rho(x) \leq C_1 - C_2\rho(x)$$

holds for all $x \in \mathbb{R}^N$ for which $\rho(x) \neq \infty$.

We then have the following easy consequence.

Lemma 3.6. *Let ρ be a Lyapunov function. Then, for any $x \in \mathbb{R}^N$ for which $\rho(x) \neq \infty$, $P_t\rho(x)$ is bounded in t , i.e., there exists a constant $C > 0$ such that*

$$P_t\rho(x) \leq C \quad \text{for all } t > 0.$$

Here, $P_t\rho(x)$ should be understood as the limit

$$P_t\rho(x) = \lim_{a \rightarrow \infty} P_t(\rho_a)(x),$$

where $\rho_a(x) = \min\{\rho(x), a\}$.

Proof. Note that by (3.8) we have

$$L\rho_a(y) \leq C_1 - C_2\rho_a(y)$$

holds for all $y \in \mathbb{R}^N$ if $a > \frac{C_1}{C_2}$. Since $\rho_a \in C_b(\mathbb{R}^N)$, then $P_t(\rho_a)$ is well defined and, moreover, we have

$$\frac{d}{dt}P_t(\rho_a) = P_t(L\rho_a) \leq C_1 - C_2P_t(\rho_a).$$

Integrating this inequality, we obtain

$$P_t(\rho_a)(x) \leq e^{-C_2t}\rho_a(x) + \frac{C_1}{C_2}(1 - e^{-C_2t}) \leq \rho_a(x) + \frac{C_1}{C_2}.$$

Taking $a \rightarrow \infty$ we have

$$P_t(\rho) \leq \rho(x) + \frac{C_1}{C_2}.$$

Thus, the conclusion follows by taking $C = \rho(x) + \frac{C_1}{C_2}$ (recall that x is fixed). $\square$

Let us now construct a Lyapunov function as a cutoff of the euclidean distance. To this end, let $\chi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a smooth function such that

$$\chi(t) = \begin{cases} 0 & \text{if } t \leq 1 \\ 1 & \text{if } t \geq 2. \end{cases}$$

Consider then the function

$$(3.9) \quad \rho(x) = |x|\chi(|x|).$$

Then clearly ρ satisfies the conditions (i) and (ii) of Definition 3.5. In order to satisfy assumption (3.8) as well, we need to find suitable bounds on ρ . We first compute, using (2.1),

$$\begin{aligned}
\Delta_k \rho(x) &= \Delta \rho(x) + 2 \sum_{\alpha \in R_+} k_\alpha \frac{\langle \nabla \rho(x), \alpha \rangle}{\langle \alpha, x \rangle} \\
&= \sum_{i=1}^N \partial_i \left(\frac{x_i}{|x|} \chi(|x|) + x_i \chi'(|x|) \right) + 2\gamma \left(\frac{1}{|x|} \chi(|x|) + \chi'(|x|) \right) \\
&= \sum_{i=1}^N \left(\frac{1}{|x|} \chi(|x|) - \frac{x_i^2}{|x|^3} \chi(|x|) + \frac{x_i^2}{|x|^2} \chi'(|x|) + \chi'(|x|) + \frac{x_i^2}{|x|} \chi''(|x|) \right) \\
&\quad + 2\gamma \left(\frac{1}{|x|} \chi(|x|) + \chi'(|x|) \right) \\
&= (N + 2\gamma - 1) \frac{1}{|x|} \chi(|x|) + (N + 2\gamma + 1) \chi'(|x|) + |x| \chi''(|x|),
\end{aligned}$$

and we notice that, from the choice of χ , there exists a constant $C_1 \geq 0$ such that

$$\Delta_k \rho \leq C_1.$$

Thus, we have

$$\begin{aligned}
L\rho(x) &\leq C_1 + \sum_{i=1}^N b_i(x) \partial_i \rho(x) \\
&= C_1 + \langle x, b(x) \rangle \left(\frac{1}{|x|} \chi(|x|) + \chi'(|x|) \right) \\
&\leq C_2 + \frac{\langle x, b(x) \rangle}{|x|^2} \rho(x),
\end{aligned}$$

for a constant $C_2 \geq 0$. To summarise, we have proved the following Lemma.

Lemma 3.7. *Assume that there exists a constant $C > 0$ such that*

$$\frac{\langle x, b(x) \rangle}{|x|^2} \leq -C \quad \text{for all } x \in \mathbb{R}^N.$$

Then, the function ρ defined by (3.9) is a Lyapunov function.

Example 3.8. In the case of linear drifts $b(x) = -cx$ with $c > 0$, we have

$$\frac{\langle x, b(x) \rangle}{|x|^2} = -c,$$

so the function ρ does satisfy the assumption of the Lemma, so it is a Lyapunov function.

We are now ready to prove the main result of this section.

Theorem 3.9. *Let $\gamma < \frac{1}{2}$. Assume that there exists $\eta < 0$ such that the following inequality holds*

$$\tilde{\Gamma}(P_t f) \leq e^{2\eta t} P_t(\tilde{\Gamma}(f)).$$

Moreover, assume that there exists a Lyapunov function ρ .

Then, there exists a sequence $(P_{t_l})_{l \geq 0}$ and a probability measure ν such that for all f bounded and continuous, and for all $x \in \mathbb{R}^N$ such that $\rho(x) \neq \infty$, we have

$$P_{t_l} f(x) \rightarrow \int f d\nu \quad \text{as } l \rightarrow \infty.$$

Additionally, the measure ν is invariant for the Markov semigroup P_t .

Proof. Fix $x \in \rho^{-1}([0, \infty))$ and define the probability measures

$$p_t^x(A) := P_t(\mathbb{1}_A)(x).$$

Step 1: $(p_t^x)_{t \geq 0}$ is a tight family. By Markov's inequality we have, for any $M > 0$,

$$p_t^x(\{\rho \geq M\}) \leq \frac{1}{M} \int \rho dp_t^x = \frac{1}{M} P_t \rho(x) \leq \frac{C}{M},$$

where in the last step we used Lemma 3.6. This implies that

$$(3.10) \quad p_t^x(\{\rho \leq M\}) \geq 1 - \frac{C}{M}.$$

Since the set $\{\rho \leq M\}$ is compact, and $M > 0$ is arbitrary, this shows that the family $(p_t^x)_{t \geq 0}$ is tight. By Prokhorov's theorem, there exists a sequence $(t_l)_{l \geq 0}$ such that $(p_{t_l}^x)_{l \geq 0}$ converges weakly to a probability measure, say ν .

Step 2: $p_t^y \Rightarrow \nu$ for all $y \in \rho^{-1}([0, \infty))$.

Let $\gamma_{x,y}(s) = x + s(y - x)$. We first compute

$$\begin{aligned} |P_{t_l} f(y) - P_{t_l} f(x)| &= \left| \int_0^1 \nabla(P_{t_l} f)(\gamma_{x,y}(s)) \cdot (y - x) ds \right| \\ &\leq |x - y| \int_0^1 |\nabla(P_{t_l} f)(\gamma_{x,y}(s))| ds \\ &= |x - y| \int_0^1 \left| \nabla_k(P_{t_l} f)(\gamma_{x,y}(s)) - \sum_{\alpha \in R_+} k_\alpha \alpha \frac{P_{t_l} f(\gamma_{x,y}(s)) - P_{t_l} f(\sigma_\alpha(\gamma_{x,y}(s)))}{\langle \alpha, \gamma_{x,y}(s) \rangle} \right| ds \\ &\leq |x - y| \int_0^1 |\nabla_k(P_{t_l} f)(\gamma_{x,y}(s))| ds \\ &\quad + \sqrt{2}|x - y| \sum_{\alpha \in R_+} k_\alpha \int_0^1 \left| \frac{P_{t_l} f(\gamma_{x,y}(s)) - P_{t_l} f(\sigma_\alpha(\gamma_{x,y}(s)))}{\langle \alpha, \gamma_{x,y}(s) \rangle} \right| ds. \end{aligned}$$

From our assumption, we have

$$(3.11) \quad |\nabla_k(P_t f)| \leq \sqrt{\tilde{\Gamma}(P_t f)} \leq e^{\eta t} \left\| \tilde{\Gamma}(f) \right\|_\infty^{1/2},$$

so the above becomes

$$(3.12) \quad \begin{aligned} |P_{t_l} f(y) - P_{t_l} f(x)| &\leq |x - y| e^{\eta t_l} \left\| \tilde{\Gamma}(f) \right\|_\infty^{1/2} \\ &\quad + \sqrt{2}|x - y| \sum_{\alpha \in R_+} k_\alpha \int_0^1 \left| \frac{P_{t_l} f(\gamma_{x,y}(s)) - P_{t_l} f(\sigma_\alpha(\gamma_{x,y}(s)))}{\langle \alpha, \gamma_{x,y}(s) \rangle} \right| ds. \end{aligned}$$

Without loss of generality we can assume that $x = 0$ and we will use an iteration argument applying (3.12) repeatedly. We first have

$$(3.13) \quad \begin{aligned} |P_{t_l} f(y) - P_{t_l} f(0)| &\leq |y| e^{\eta t_l} \left\| \tilde{\Gamma}(f) \right\|_{\infty}^{1/2} \\ &\quad + \sqrt{2} |y| \sum_{\alpha \in R_+} \frac{k_{\alpha}}{|\langle \alpha, y \rangle|} \int_0^1 \frac{1}{s} |P_{t_l} f(sy) - P_{t_l} f(\sigma_{\alpha}(sy))| ds. \end{aligned}$$

Applying (3.12) again, noting that $|sy - \sigma_{\alpha}(sy)| = s\sqrt{2}|\langle \alpha, y \rangle|$, we have

$$\begin{aligned} |P_{t_l} f(y) - P_{t_l} f(0)| &\leq |y| e^{\eta t_l} \left\| \tilde{\Gamma}(f) \right\|_{\infty}^{1/2} \left(1 + 2 \sum_{\alpha \in R_+} k_{\alpha} \right) \\ &\quad + 2^{3/2} |y| \sum_{\alpha \in R_+} \sum_{\beta \in R_+} k_{\alpha} k_{\beta} \int_0^1 \int_0^1 \left| \frac{P_{t_l} f(\gamma_{\sigma_{\alpha}(sy), sy}(u)) - P_{t_l} f(\sigma_{\beta}(\gamma_{\sigma_{\alpha}(sy), sy}(u)))}{\langle \beta, \gamma_{\sigma_{\alpha}(sy), sy}(u) \rangle} \right| du ds. \end{aligned}$$

The next step is already too difficult to write down, but we can see that we obtain

$$\begin{aligned} |P_{t_l} f(y) - P_{t_l} f(0)| &\leq |y| e^{\eta t_l} \left\| \tilde{\Gamma}(f) \right\|_{\infty}^{1/2} \sum_{i=0}^{\infty} \left[2^i \sum_{\alpha_1 \in R_+} \sum_{\alpha_2 \in R_+} \cdots \sum_{\alpha_i \in R_+} k_{\alpha_1} k_{\alpha_2} \cdots k_{\alpha_i} \right] \\ &= |y| e^{\eta t_l} \left\| \tilde{\Gamma}(f) \right\|_{\infty}^{1/2} \sum_{i=0}^{\infty} (2\gamma)^i. \end{aligned}$$

Therefore, if $\gamma < \frac{1}{2}$, the infinite sum converges and we have

$$(3.14) \quad |P_{t_l} f(y) - P_{t_l} f(0)| \leq C |y| e^{\eta t_l} \left\| \tilde{\Gamma}(f) \right\|_{\infty}^{1/2}.$$

Since $\eta < 0$, this shows that $p_{t_l}^y \Rightarrow \nu$, as required.

Step 3: ν is an invariant measure for P_t . We will now apply the Krylov-Bogoliubov theorem. We have, using (3.10),

$$\mu_T^x(\{\rho \leq M\}) = \frac{1}{T} \int_0^T p_t^x(\{\rho \leq M\}) dt \geq 1 - \frac{C}{M},$$

so the family $(\mu_T^x)_{T>0}$ is tight. By the Krylov-Bogoliubov theorem, there exists an invariant measure for P_t , say μ .

Fix a bounded and continuous function f and let $\epsilon > 0$. Then, since $p_{t_l}^x \Rightarrow \nu$, there exists $l_1 > 0$ such that

$$|p_{t_l}^x(f) - \nu(f)| < \frac{\epsilon}{2} \quad \text{for all } l \geq l_1.$$

On the other hand, since μ is an invariant measure, we have

$$|p_{t_l}^x(f) - \mu(f)| \leq \int |P_{t_l} f(x) - P_{t_l} f(y)| d\mu(y) \leq C \left\| \tilde{\Gamma}(f) \right\|_{\infty}^{1/2} e^{\eta t_l} \int |x - y| d\mu(y),$$

where, in the second step, we used (3.14). Since $\eta < 0$, there exists $k_2 > 0$ such that

$$|p_{t_l}^x(f) - \mu(f)| < \frac{\epsilon}{2} \quad \text{for all } l > l_2.$$

Thus, for all $l > \max\{l_1, l_2\}$, we have

$$|\mu(f) - \nu(f)| \leq |p_{t_l}^x(f) - \mu(f)| + |p_{t_l}^x(f) - \nu(f)| < \epsilon,$$

which shows that $\mu = \nu$. This finishes the proof. $\square$

Remark. We note that we needed to impose conditions for the coefficients k_α to be small both in order to obtain $\eta < 0$, as well as in the proof of existence of invariant measure. It would be interesting to find out whether this is just a shortfall of our chosen method, or whether this is a deeper characteristic of Dunkl operators.

Example 3.10. If $b(x) = -cx$, for some $c > 0$, then an invariant measure for the resulting semigroup is given by $d\nu(x) = e^{-c|x|^2/2} d\mu_k(x)$. Indeed, we have

$$\begin{aligned} \int_{\mathbb{R}^N} Lf d\nu &= \int_{\mathbb{R}^N} (\Delta_k f - cx \cdot \nabla_k f) e^{-c|x|^2/2} d\mu_k \\ &= - \int_{\mathbb{R}^N} \left(\nabla \left(-\frac{c|x|^2}{2} \right) - cx \right) \cdot \nabla_k f d\nu = 0. \end{aligned}$$

4. THE INFINITE DIMENSIONAL CASE

In this section we work over the infinite dimensional space $\Omega = (\mathbb{R}^N)^{\mathbb{Z}^d}$. An element of this set will be denoted by $\omega = (\omega_l)_{l \in \mathbb{Z}^d}$, where each $\omega_l \in \mathbb{R}^N$. On the lattice $\mathbb{Z}^d$ we define the distance $|l - j| = d(l, j) := \sum_{i=1}^d |l_i - j_i|$. A cylinder function on Ω is a smooth function

$f : \Omega \rightarrow \mathbb{R}$ that only depends the components ω_l for $l \in \Lambda$ and $\Lambda \subset \mathbb{Z}^d$ is a finite set. The smallest such subset $\Lambda \subset \mathbb{Z}^d$ will be denoted $\Lambda(f)$. It is known that the set of cylinder functions on Ω is dense in $\mathcal{C}(\Omega)$, see for example [6].

4.1. Construction of the semigroup. We want to define an infinite dimensional semigroup on $(\mathbb{R}^N)^{\mathbb{Z}^d}$ with generator

$$\mathcal{L} = \sum_{l \in \mathbb{Z}^d} L^{(l)} + \sum_{l \in \mathbb{Z}^d} e^{(l)} \cdot \nabla_k^{(l)},$$

where

$$L^{(l)} = \Delta_k^{(l)} + b^{(l)} \cdot \nabla_k^{(l)}$$

is a copy of the operator studied in the previous section, acting only on the l component of $\omega \in (\mathbb{R}^N)^{\mathbb{Z}^d}$. Here $b^{(l)} : \Omega \rightarrow \mathbb{R}^N$ is a function that only depends on the ω_l component. In addition, $e^{(l)} : \Omega \rightarrow \mathbb{R}^N$ is a C^1 function which depends only on ω_j for $|j - l| < R$ for some fixed $R > 0$ (we say that $e^{(l)}$ has finite range of interaction), and such that $e^{(l)}$ and all its first degree derivatives are uniformly bounded. As in the previous section, we assume firstly that for any $l \in \mathbb{Z}^d$ we have

$$(4.1) \quad \frac{2}{\langle \alpha, \omega_l \rangle^2} - \langle \alpha, b^{(l)}(\omega) \rangle \frac{1}{\langle \alpha, \omega_l \rangle} \geq 0 \quad \forall \omega \in \Omega, \forall \alpha \in R_+$$

and

$$(4.2) \quad \frac{2}{\langle \alpha, \omega_l \rangle^2} - \langle \alpha, b^{(l)}(\omega) + e^{(l)}(\omega) \rangle \frac{1}{\langle \alpha, \omega_l \rangle} \geq 0 \quad \forall \omega \in \Omega, \forall \alpha \in R_+.$$

Secondly, we also assume that for all $l \in \mathbb{Z}^d$ we have

$$(4.3) \quad b^{(l)} \circ g^{(l)} = g^{(l)} b^{(l)} \quad \text{and} \quad e^{(l)} \circ g^{(l)} = g^{(l)} e^{(l)},$$

and

$$(4.4) \quad e^{(\bar{l})} \circ g^{(l)} = e^{(\bar{l})} \quad \forall \bar{l} \neq l.$$

These two sets of assumptions mirror the conditions (3.2) and (3.3), respectively, of the finite dimensional case.

The operators $L^{(l)}$ are commutative and, as in the previous section, each generates a finite dimensional semigroup. Thus, their sum $\sum_{l \in \mathbb{Z}^d} L^{(l)}$ also generates a semigroup. In this case, the dynamics is given by infinitely many copies of the same model acting independently of each other. Introducing the terms corresponding to the functions $e^{(l)}$ makes the dynamics more interesting as it allows for interactions between the individual diffusions.

In order to define this semigroup formally, we first consider the truncated operator for some finite $\Lambda \subset \mathbb{Z}^d$,

$$\mathcal{L}_\Lambda := \sum_{l \in \mathbb{Z}^d} L^{(l)} + \sum_{l \in \Lambda} e^{(l)} \cdot \nabla_k^{(l)}.$$

This operator is well defined on the space of cylinder functions on Ω and we assume that it generates a Markov semigroup $(P_t^\Lambda)_{t \geq 0}$. Since Λ is finite and by the finite range of interactions assumption, then in fact only the diffusions corresponding to the points of $\Lambda^R := \{l \in \mathbb{Z}^d : d(l, \Lambda) < R\}$ interact with each other. Thus, we can split $\mathcal{L}_\Lambda$ into two components

$$\mathcal{L}_\Lambda = \sum_{l \in \mathbb{Z}^d \setminus \Lambda^R} L^{(l)} + \left(\sum_{l \in \Lambda^R} L^{(l)} + \sum_{l \in \Lambda} e^{(l)} \cdot \nabla_k^{(l)} \right).$$

As discussed above, the operators $L^{(l)}$ are commutative, so the first part generates a semigroup of diffusions that do not interact with each other, while the second part is essentially finite dimensional and it can be dealt with as in the previous section.

Define the carré-du-champ operator

$$\Gamma_\Lambda(f) := \frac{1}{2} (\mathcal{L}_\Lambda(f^2) - 2f \mathcal{L}_\Lambda f).$$

Using computations similar to the finite dimensional case, we have

$$\begin{aligned} \Gamma_\Lambda(f)(\omega) &= \sum_{l \in \mathbb{Z}^d} |\nabla^{(l)} f(\omega)|^2 + \frac{1}{2} \sum_{l \in \mathbb{Z}^d \setminus \Lambda} k_\alpha A_\alpha^{(l)}(f)^2(\omega) \left[2 - \langle \alpha, b^{(l)}(\omega) \rangle \langle \alpha, \omega \rangle \right] \\ &\quad + \frac{1}{2} \sum_{l \in \Lambda} k_\alpha A_\alpha^{(l)}(f)^2(\omega) \left[2 - \langle \alpha, b^{(l)}(\omega) + e^{(l)}(\omega) \rangle \langle \alpha, \omega \rangle \right]. \end{aligned}$$

Here, in keeping with the notation above, $\nabla^{(l)}$ is the usual gradient acting only on the ω_l component of $\omega \in \Omega$, and $A_\alpha^{(l)}$ is the operator A_α acting on the same component ω_l . From the assumptions (4.1) and (4.2) we have

$$\Gamma_\Lambda(f) \geq 0.$$

We want to define the infinite semigroup for any cylindrical function f as the limit in the uniform norm

$$P_t f = \lim_{\Lambda \rightarrow \mathbb{Z}^d} P_t^\Lambda f.$$

In order to prove that this limit exists, it is enough to show that for each increasing sequence $(\Lambda_n)_{n \in \mathbb{N}}$ of finite subsets of $\mathbb{Z}^d$, the sequence $(P_t^{\Lambda_n} f)_{n \in \mathbb{N}}$ is Cauchy. In other words, fixing a cylindrical function f , for finite $\Lambda \subset \Lambda' \subset \mathbb{Z}^d$, we want to estimate the norm

$$\left\| P_t^{\Lambda'} f - P_t^{\Lambda} f \right\|_{\infty}.$$

We assume also that Λ is large enough so that $\Lambda(f) \subset \Lambda$. We construct a sequence $\Lambda_0, \dots, \Lambda_n$, where $n = |\Lambda'| - |\Lambda|$, such that $\Lambda_0 = \Lambda$, $\Lambda_n = \Lambda'$ and for each $j = 0, \dots, n-1$, $\Lambda_{j+1} \setminus \Lambda_j = \{h_j\}$. Then, we have

$$(4.5) \quad \left\| P_t^{\Lambda'} f - P_t^{\Lambda} f \right\|_{\infty} \leq \sum_{j=0}^{n-1} \left\| P_t^{\Lambda_{j+1}} f - P_t^{\Lambda_j} f \right\|_{\infty}.$$

But we can compute

$$P_t^{\Lambda_{j+1}} f - P_t^{\Lambda_j} f = \int_0^t \frac{d}{ds} \left[P_{t-s}^{\Lambda_j} (P_s^{\Lambda_{j+1}} f) \right] ds = \int_0^t P_{t-s}^{\Lambda_j} (\mathcal{L}_{\Lambda_{j+1}} - \mathcal{L}_{\Lambda_j}) P_s^{\Lambda_{j+1}} f ds,$$

and since $P_t^{\Lambda_j}$ is a Markov semigroup we can estimate

$$|P_t^{\Lambda_{j+1}} f - P_t^{\Lambda_j} f| \leq \left\| e^{(h_j)} \right\|_{\infty} \int_0^t P_{t-s}^{\Lambda_j} \left(\left| \nabla_k^{(h_j)} (P_s^{\Lambda_{j+1}} f) \right| \right) ds$$

Similarly to the previous section, we now introduce the symmetrised gradient form

$$(4.6) \quad \tilde{\Gamma}^{(l)}(h)(\omega) = \sum_{g \in G} |\nabla_k^{(l)} h(g^{(l)} \omega)|^2,$$

defined for all $l \in \mathbb{Z}^d$, and where $g^{(l)}$ is the element g of the reflection group G acting on the $\omega_l \in \mathbb{R}^N$ coordinate of $\omega \in \Omega$. Using this definition, we have

$$\left| \nabla_k^{(h_j)} (P_s^{\Lambda_{j+1}} f) \right|^2 \leq \tilde{\Gamma}^{(h_j)}(P_s^{\Lambda_{j+1}} f),$$

and hence, from the above we deduce

$$(4.7) \quad \left\| P_t^{\Lambda_{j+1}} f - P_t^{\Lambda_j} f \right\|_{\infty} \leq \left\| e^{(h_j)} \right\|_{\infty} \int_0^t \left\| \tilde{\Gamma}^{(h_j)}(P_s^{\Lambda_{j+1}} f) \right\|_{\infty}^{1/2} ds.$$

Thus, we need to obtain bounds on the quantity $\left\| \tilde{\Gamma}^{(h_j)}(P_s^{\Lambda_{j+1}} f) \right\|_{\infty}$. This is done in the following Theorem.

Theorem 4.1. *Let Λ be a finite subset of $\mathbb{Z}^d$ and consider the Markov semigroup $(P_t^{\Lambda})_{t \geq 0}$ generated by the operator*

$$\mathcal{L}_{\Lambda} := \sum_{l \in \mathbb{Z}^d} L^{(l)} + \sum_{l \in \Lambda} e^{(l)} \cdot \nabla_k^{(l)}$$

and denote $f_t := P_t^{\Lambda} f$. Assume that

$$\eta := \max_i \sup_{\omega \in \Omega} \partial_i^{(l)}(b_i^{(l)})(\omega) + (N-1) \max_{m \neq i} \left\| \partial_m^{(l)}(b_i) \right\|_{\infty} + \sqrt{2}\gamma \max_{\alpha \in R_+} \left\| A_{\alpha}^{(l)}(b^{(l)}) \right\|_{\infty} < 0$$

and

$$\zeta := \sum_{l \in \mathbb{Z}^d} \left\| e^{(l)} \right\|_{\infty} < \infty.$$

Let f be a cylinder function such that $\Lambda(f) \subset \Lambda$ and let $l \notin \Lambda(f)$. Let also

$$N_l = \left\lfloor \frac{d(l, \Lambda(f))}{R} \right\rfloor + 1,$$

where the square brackets indicate integer part. Then, for any $\sigma > 0$ there exists $\tau \geq 1$ large enough such that if $N_l \geq \tau s$, and we have the bound

$$(4.8) \quad \left\| \tilde{\Gamma}^{(l)}(f_s) \right\|_{\infty} \leq e^{-2N_l\sigma - 2s\sigma} \sum_{j \in \mathbb{Z}^d} \left\| \tilde{\Gamma}^{(j)}(f) \right\|_{\infty}.$$

Proof. Fix some $i = 1, \dots, N$. We first look at

$$(4.9) \quad \begin{aligned} \partial_u P_{s-u}^{\Lambda}(|T_i^{(l)} f_u|^2) &= P_{s-u}^{\Lambda}(-\mathcal{L}_{\Lambda}(|T_i^{(l)} f_u|^2) + 2T_i^{(l)} f_u \cdot T_i^{(l)} \mathcal{L}_{\Lambda} f_u) \\ &= P_{s-u}^{\Lambda}(-2\Gamma_{\Lambda}(T_i^{(l)} f_u) + 2T_i^{(l)} f_u \cdot [T_i^{(l)}, \mathcal{L}_{\Lambda}] f_u) \\ &\leq P_{s-u}^{\Lambda}(2T_i^{(l)} f_u \cdot [T_i^{(l)}, \mathcal{L}_{\Lambda}] f_u), \end{aligned}$$

where we used the positivity of Γ_{Λ} .

We can check directly that for distinct $l, l' \in \mathbb{Z}^d$, the operators $T_i^{(l)}$ and $T_j^{(l')}$ commute for any $i, j = 1, \dots, N$. The same holds for $l = l'$, but here we need to use the fact that Dunkl operators commute. As a consequence (recall also that $b^{(j)}$ depends only on the ω_j component), we have $[T_i^{(l)}, L^{(j)}] = 0$ for all $j \neq l$. Thus, we can simplify the commutator

$$\begin{aligned} [T_i^{(l)}, \mathcal{L}_{\Lambda}] &= [T_i^{(l)}, L^{(l)}] + \sum_{j \in \Lambda} [T_i^{(l)}, e^{(j)} \cdot \nabla_k^{(j)}] \\ &= [T_i^{(l)}, L^{(l)}] + \sum_{j \in \Lambda} [T_i^{(l)}, e^{(j)}] \cdot \nabla_k^{(j)}, \end{aligned}$$

where in the last step we used again the commutativity of $T_i^{(l)}$ with $T_u^{(j)}$. So we next need to compute $[T_i^{(l)}, e^{(j)}]$. We have

$$\begin{aligned} [T_i^{(l)}, e^{(j)}]h &= \partial_i^{(l)}(he^{(j)}) + \sum_{\alpha \in R_+} k_{\alpha} \alpha_i A_{\alpha}^{(l)}(he^{(j)}) - e^{(j)} \partial_i^{(l)}h - \sum_{\alpha \in R_+} k_{\alpha} \alpha_i A_{\alpha}^{(l)}(h)e^{(j)} \\ &= h \partial_i^{(l)}(e^{(j)}) - \sum_{\alpha \in R_+} k_{\alpha} \alpha_i [A_{\alpha}^{(l)}(h)e^{(j)} - A_{\alpha}^{(l)}(he^{(j)})]. \end{aligned}$$

But

$$\begin{aligned} A_{\alpha}^{(l)}(h)e^{(j)} - A_{\alpha}^{(l)}(he^{(j)}) &= \frac{1}{\langle \alpha, \omega_l \rangle} \left[e^{(j)}(\omega)(h(\omega) - h(\sigma_{\alpha}^{(l)}\omega)) - (h(\omega)e^{(j)}(\omega) - h(\sigma_{\alpha}^{(l)}\omega)e^{(j)}(\sigma_{\alpha}^{(l)}\omega)) \right] \\ &= -h(\sigma_{\alpha}^{(l)}\omega)A_{\alpha}^{(l)}(e^{(j)}). \end{aligned}$$

Thus, we have

$$[T_i^{(l)}, e^{(j)}]h = h \partial_i^{(l)}(e^{(j)}) + \sum_{\alpha \in R_+} k_{\alpha} \alpha_i h(\sigma_{\alpha}^{(l)}\omega)A_{\alpha}^{(l)}(e^{(j)}).$$

Therefore, we have obtained

(4.10)

$$[T_i^{(l)}, \mathcal{L}_\Lambda]h = [T_i^{(l)}, L^{(l)}] + \sum_{j \in \Lambda} \left[\partial_i^{(l)}(e^{(j)}) \cdot \nabla_k^{(j)} h + \sum_{\alpha \in R_+} k_\alpha \alpha_i A_\alpha^{(l)}(e^{(j)}) \cdot \nabla_k^{(j)} h(\sigma_\alpha^{(l)} \omega) \right].$$

Recall the definition of the symmetrised gradient form given in (4.6). As in the finite case, and due to the assumptions (4.3) and (4.4), the computations in (4.9) and (4.10) can be extended to more difficult terms, involving compositions with reflections $g^{(l)}$. For simplicity, we do not include all the details here, but the method is the same as before. Thus, we obtain

$$\begin{aligned} & \frac{d}{du} P_{s-u}^\Lambda(\tilde{\Gamma}^{(l)}(f_u)) \\ (4.11) \quad & \leq \sum_{g \in G} \sum_{i=1}^N P_{s-u}^\Lambda \left(2T_i^{(l)} f_u(g^{(l)} \omega) [T_i^{(l)}, L^{(l)}] f_u(g^{(l)} \omega) \right) \\ (4.12) \quad & + \sum_{g \in G} \sum_{i=1}^N \sum_{m=1}^N \sum_{j \in \Lambda} 2P_{s-u}^\Lambda \left[\left(\partial_i^{(l)}(e_m^{(j)}) T_m^{(j)} f_u T_i^{(l)} f_u \right) \circ g^{(l)} \right] \\ (4.13) \quad & + \sum_{g \in G} \sum_{i=1}^N \sum_{m=1}^N \sum_{j \in \Lambda} \sum_{\alpha \in R_+} 2k_\alpha \alpha_i P_{s-u}^\Lambda \left[\left(A_\alpha^{(l)}(e_m^{(j)}) (T_m^{(j)} f_u) \circ \sigma_\alpha^{(l)} T_i^{(l)} f_u \right) \circ g^{(l)} \right]. \end{aligned}$$

The term in (4.11) is essentially the finite dimensional case discussed in Theorem 3.3 since all operators involved only act on the l component. We thus have

$$\sum_{g \in G} \sum_{i=1}^N P_{s-u}^\Lambda \left(2T_i^{(l)} f_u(g^{(l)} \omega) [T_i^{(l)}, L^{(l)}] f_u(g^{(l)} \omega) \right) \leq 2\eta_l P_{s-u}^\Lambda(\tilde{\Gamma}^{(l)}(f_u)),$$

where

$$\eta_l := \max_i \sup_{\omega \in \Omega} \partial_i^{(l)}(b_i^{(l)})(\omega) + (N-1) \max_{m \neq i} \left\| \partial_m^{(l)}(b_i) \right\|_\infty + \sqrt{2}\gamma \max_{\alpha \in R_+} \left\| A_\alpha^{(l)}(b^{(l)}) \right\|_\infty.$$

In order to estimate the next two terms, let

$$E_{l,j} := \max \left\{ \max_{i,m} \left\| \partial_i^{(l)}(e_m^{(j)}) \right\|_\infty, \max_{i,m,\alpha} \left\| A_\alpha^{(l)}(e_m^{(j)}) \right\|_\infty \right\} \leq \sqrt{2} \max_{i,m} \left\| \partial_i^{(l)}(e_m^{(j)}) \right\|_\infty,$$

where the second inequality follows from (2.2). Clearly $E_{l,j} \geq 0$ for all $l, j \in \mathbb{Z}^d$, but note also that $E_{l,j} = 0$ if $d(l, j) \geq R$ since $e^{(l)}$ only depends on the components ω_j with $d(i, j) < R$. The strategy in estimating the remaining terms (4.12) and (4.13) is to use the inequality

$2xy \leq \epsilon x^2 + \frac{1}{\epsilon} y^2$ to separate the products into sums of squares. We first estimate

$$\begin{aligned}
& \sum_{g \in G} \sum_{i=1}^N \sum_{m=1}^N \sum_{j \in \Lambda} 2P_{s-u}^\Lambda \left[\left(\partial_i^{(l)}(e_m^{(j)}) T_m^{(j)} f_u T_i^{(l)} f_u \right) \circ g^{(l)} \right] \\
& \leq N \sum_{g \in G} \sum_{j \in \Lambda} E_{l,j} P_{s-u}^\Lambda \left[\epsilon |\nabla_k^{(l)} f_u|^2 \circ g^{(l)} + \frac{1}{\epsilon} |\nabla_k^{(j)} f_u|^2 \circ g^{(l)} \right] \\
& = \epsilon N \left(\sum_{j \in \Lambda} E_{l,j} \right) P_{s-u}^\Lambda(\tilde{\Gamma}^{(l)}(f_u)) + \frac{1}{\epsilon} N \sum_{g \in G} \sum_{j \in \Lambda} E_{l,j} P_{s-u}^\Lambda \left[|\nabla_k^{(j)} f_u|^2 \circ g^{(l)} \right] \\
& \leq \epsilon N \left(\sum_{j \in \Lambda} E_{l,j} \right) P_{s-u}^\Lambda(\tilde{\Gamma}^{(l)}(f_u)) + \frac{1}{\epsilon} N \sum_{g \in G} \sum_{j \in \Lambda} E_{l,j} P_{s-u}^\Lambda \left[(\tilde{\Gamma}^{(j)} f_u) \circ g^{(l)} \right],
\end{aligned}$$

and similarly

$$\begin{aligned}
& \sum_{g \in G} \sum_{i=1}^N \sum_{m=1}^N \sum_{j \in \Lambda} \sum_{\alpha \in R_+} 2k_\alpha \alpha_i P_{s-u}^\Lambda \left[\left(A_\alpha^{(l)}(e_m^{(j)}) (T_m^{(j)} f_u) \circ \sigma_\alpha^{(l)} T_i^{(l)} f_u \right) \circ g^{(l)} \right] \\
& \leq \sum_{g \in G} \sum_{j \in \Lambda} \sum_{\alpha \in R_+} \sqrt{2} N k_\alpha E_{l,j} P_{s-u}^\Lambda \left[\epsilon |\nabla_k^{(l)} f_u|^2 \circ g^{(l)} + \frac{1}{\epsilon} |\nabla_k^{(j)} f_u|^2 \circ (\sigma_\alpha^{(l)} \circ g^{(l)}) \right] \\
& = \sqrt{2} N \gamma \epsilon \left(\sum_{j \in \Lambda} E_{l,j} \right) P_{s-u}^\Lambda(\tilde{\Gamma}^{(l)}(f_u)) + \sqrt{2} N \gamma \frac{1}{\epsilon} \sum_{g \in G} \sum_{j \in \Lambda} E_{l,j} P_{s-u}^\Lambda \left[|\nabla_k^{(j)} f_u|^2 \circ g^{(l)} \right] \\
& \leq \sqrt{2} N \gamma \epsilon \left(\sum_{j \in \Lambda} E_{l,j} \right) P_{s-u}^\Lambda(\tilde{\Gamma}^{(l)}(f_u)) + \sqrt{2} N \gamma \frac{1}{\epsilon} \sum_{g \in G} \sum_{j \in \Lambda} E_{l,j} P_{s-u}^\Lambda \left[\tilde{\Gamma}^{(j)}(f_u) \circ g^{(l)} \right].
\end{aligned}$$

Combining all these results, we have obtained that

$$(4.14) \quad \frac{d}{du} P_{s-u}^\Lambda(\tilde{\Gamma}^{(l)}(f_u)) \leq 2\tilde{\eta}_l P_{s-u}^\Lambda(\tilde{\Gamma}^{(l)}(f_u)) + \sum_{g \in G} \sum_{j \in \Lambda} \tilde{E}_{l,j} P_{s-u}^\Lambda \left[\tilde{\Gamma}^{(j)}(f_u) \circ g^{(l)} \right],$$

where

$$(4.15) \quad \tilde{\eta}_l := \eta_l + \epsilon \frac{N}{2} (1 + \sqrt{2}\gamma) \left(\sum_{j \in \Lambda} E_{l,j} \right),$$

and

$$\tilde{E}_{l,j} := \frac{1}{\epsilon} N (1 + \sqrt{2}\gamma) E_{l,j}.$$

We choose $\epsilon > 0$ such that $\tilde{\eta}_l < 0$; this is possible because by our assumption $\eta_l < 0$. Multiplying inequality (4.14) by $e^{-2\tilde{\eta}_l u}$, and integrating with respect to u from 0 to s , we obtain

$$\tilde{\Gamma}^{(l)}(f_s) \leq e^{2\tilde{\eta}_l s} P_s^\Lambda(\tilde{\Gamma}^{(l)} f) + \sum_{g \in G} \sum_{j \in \Lambda} \tilde{E}_{l,j} \int_0^s e^{2\tilde{\eta}_l(s-u)} P_{s-u}^\Lambda \left[\tilde{\Gamma}^{(j)}(f_u) \circ g^{(l)} \right] du.$$

Taking supremum norm over this inequality, and since $\tilde{\eta}_l < 0$, we have

$$(4.16) \quad \left\| \tilde{\Gamma}^{(l)}(f_s) \right\|_\infty \leq \left\| \tilde{\Gamma}^{(l)} f \right\|_\infty + |G| \sum_{j \in \Lambda} \tilde{E}_{l,j} \int_0^s \left\| \tilde{\Gamma}^{(j)}(f_u) \right\|_\infty du.$$

Up until this point we have not used the assumption that $l \notin \Lambda(f)$ so this inequality holds for general $l \in \mathbb{Z}^d$. Since $l \notin \Lambda(f)$, then f does not depend on the ω_l coordinate and so the first term on the right hand side of inequality (4.16) vanishes.

We can improve inequality (4.16) by applying itself iteratively to the term under the integral. At the first step we obtain

$$\begin{aligned} \left\| \tilde{\Gamma}^{(l)}(f_s) \right\|_{\infty} &\leq |G| \sum_{j \in \Lambda} \tilde{E}_{l,j} \int_0^s \left[\left\| \tilde{\Gamma}^{(j)} f \right\|_{\infty} + |G| \sum_{j' \in \Lambda} \tilde{E}_{j,j'} \int_0^u \left\| \tilde{\Gamma}^{(j')}(f_v) \right\|_{\infty} dv \right] du \\ &\leq s |G| \sum_{j \in \Lambda} \tilde{E}_{l,j} \left\| \tilde{\Gamma}^{(j)} f \right\|_{\infty} + |G|^2 \sum_{j \in \Lambda} \sum_{j' \in \Lambda} \tilde{E}_{l,j} \tilde{E}_{j,j'} \int_0^s \int_0^u \left\| \tilde{\Gamma}^{(j')}(f_u) \right\|_{\infty} dv du. \end{aligned}$$

Note that if $d(l, \Lambda(f)) \geq R$ and $d(j, l) < R$, then $j \notin \Lambda(f)$, so the first sum vanishes again. In fact, we need to apply this method iteratively $N_l = \left\lceil \frac{d(l, \Lambda(f))}{R} \right\rceil + 1$ times in order to get a non-trivial contribution. After infinitely many iterations, we obtain

$$\begin{aligned} \left\| \tilde{\Gamma}^{(l)}(f_s) \right\|_{\infty} &\leq \left(\frac{(sC_l)^{N_l}}{N_l!} + \frac{(sC_l)^{N_l+1}}{(N_l+1)!} + \dots \right) \sum_{j \in \mathbb{Z}^d} \left\| \tilde{\Gamma}^{(j)}(f) \right\|_{\infty} \\ &\leq \frac{(sC_l)^{N_l}}{(N_l)!} e^{sC_l} \sum_{j \in \mathbb{Z}^d} \left\| \tilde{\Gamma}^{(j)}(f) \right\|_{\infty}, \end{aligned}$$

where

$$(4.17) \quad C_l = |G| \sum_{j \in \Lambda} \tilde{E}_{l,j}.$$

Note that the factorials appearing in the denominators above are a result of computing nested integrals.

Let us estimate the constant appearing in this inequality. Using the fact that for any natural number n we have $n! > \left(\frac{n}{e}\right)^n$ (which follows from the expansion of e^n), we obtain

$$\left\| \tilde{\Gamma}^{(l)}(f_s) \right\|_{\infty} \leq e^{sC_l + N_l(\log sC_l - \log N_l + 1)} \sum_{j \in \mathbb{Z}^d} \left\| \tilde{\Gamma}^{(j)}(f) \right\|_{\infty}.$$

Now fix $\sigma > 0$. There exists $\tau \geq 0$ large enough such that

$$(4.18) \quad \log \frac{C_l}{\tau} + \frac{C_l}{\tau} + 1 \leq -4\sigma.$$

If, in addition, $N_l \geq \tau s$, then

$$sC_l + N_l(\log sC_l - \log N_l + 1) \leq N_l \left(\frac{C_l}{\tau} + \log \frac{C_l}{\tau} + 1 \right) \leq -4\sigma N_l \leq -2N_l\sigma - 2s\sigma.$$

Finally, this gives

$$(4.19) \quad \left\| \tilde{\Gamma}^{(l)}(f_s) \right\|_{\infty} \leq e^{-2N_l\sigma - 2s\sigma} \sum_{j \in \mathbb{Z}^d} \left\| \tilde{\Gamma}^{(j)}(f) \right\|_{\infty},$$

as required. $\square$

Now that we have proved the required bounds on the symmetrised gradient, we can go back to the definition of the infinite dimensional semigroup. Fix $\sigma > 0$ and let $\tilde{N} =$

$\left\lceil \frac{d(\mathbb{Z}^d \setminus \Lambda, \Lambda(f))}{R} \right\rceil + 1$. By definition, we have $N_{h_j} \geq \tilde{N}$ for all $j = 0, \dots, n-1$. From (4.5) and (4.7), with the help of Theorem 4.1, we obtain

$$(4.20) \quad \begin{aligned} \|P_t^{\Lambda'} f - P_t^\Lambda f\|_\infty &\leq \left(\sum_{l \in \mathbb{Z}^d} \|\tilde{\Gamma}^{(l)}(f)\|_\infty \right)^{1/2} \sum_{j=0}^{n-1} \|e^{(h_j)}\|_\infty \int_0^t e^{-\sigma N_{h_j} - \sigma s} ds \\ &\leq \zeta e^{-\sigma \tilde{N}} \frac{1 - e^{\sigma t}}{\sigma} \left(\sum_{l \in \mathbb{Z}^d} \|\tilde{\Gamma}^{(l)}(f)\|_\infty \right)^{1/2}, \end{aligned}$$

which holds for all $\tilde{N} \geq \tau t$, where $\tau \geq 1$ is fixed. Recall that $\zeta = \sum_{l \in \mathbb{Z}^d} \|e^{(l)}\|_\infty$ was assumed to be finite.

This concludes the proof that for each increasing sequence $(\Lambda_n)_{n \geq 1}$ of finite subsets of $\mathbb{Z}^d$, and for each cylindrical function f , the sequence $(P_t^{\Lambda_n} f)_n$ is Cauchy. Therefore, we have proved the following Theorem.

Theorem 4.2. *Consider the operator*

$$\mathcal{L} := \sum_{l \in \mathbb{Z}^d} (\Delta_k^{(l)} + b^{(l)} \cdot \nabla_k^{(l)}) + \sum_{l \in \mathbb{Z}^d} e^{(l)} \cdot \nabla_k^{(l)}.$$

Assume that the coefficients $b^{(l)}$ and $e^{(l)}$ are uniformly bounded such that

$$\eta_l := \max_i \sup_{\omega \in \Omega} \partial_i^{(l)}(b_i^{(l)})(\omega) + (N-1) \max_{m \neq i} \|\partial_m^{(l)}(b_i)\|_\infty + \sqrt{2}\gamma \max_{\alpha \in R_+} \|A_\alpha^{(l)}(b^{(l)})\|_\infty < 0$$

and

$$\zeta := \sum_{l \in \mathbb{Z}^d} \|e^{(l)}\|_\infty < \infty.$$

Then, for any cylinder function f , the following limit exists in the uniform norm

$$P_t f := \lim_{\Lambda \rightarrow \mathbb{Z}^d} P_t^\Lambda f,$$

and it defines a Markov semigroup $(P_t)_{t \geq 0}$ on $C_b(\Omega)$, with infinitesimal generator given by $\mathcal{L}$.

4.2. Invariant measure. In this section we discuss the existence of an invariant measure for the infinite dimensional semigroup that we have just defined. The strategy is the same as in section 3.2, and we start by defining a Lyapunov function ρ that is suitable to our new setting.

Let $(a_l)_{l \in \mathbb{Z}^d} \subset (0, \infty)$ be an absolutely convergent series, i.e.,

$$\sum_{l \in \mathbb{Z}^d} a_l < \infty.$$

We assume that for each $l \in \mathbb{Z}^d$ there exists a function $\rho_l : \Omega \rightarrow (0, \infty)$, which depends only on the ω_l coordinate, and such that $\omega_l \mapsto \rho_l(\omega)$ is a Lyapunov function for the operator $L^{(l)}$ on $\mathbb{R}^N$. Moreover, assume that there exist constants $C_1 \geq 0$ and $C_2 > 0$ such that

$$(4.21) \quad L^{(l)} \rho_l + e^{(l)} \cdot \nabla_k^{(l)} \rho_l \leq C_1 - C_2 \rho_l \quad \forall l \in \mathbb{Z}^d.$$

For each $0 < r \leq \infty$, define the set

$$\Omega_r := \left\{ \omega \in \Omega : \sum_{l \in \mathbb{Z}^d} a_l \rho_l(\omega) < r \right\},$$

and consider the function $\rho : \Omega_\infty \rightarrow (0, \infty)$ given by

$$\rho(\omega) := \sum_{l \in \mathbb{Z}^d} a_l \rho_l(\omega).$$

Using condition (4.21) and the fact that ρ_l only depends on the ω_l coordinate, we have

$$(4.22) \quad \mathcal{L}\rho = \sum_{l \in \mathbb{Z}^d} a_l (L^{(l)} \rho_l + e^{(l)} \cdot \nabla_k^{(l)} \rho_l) \leq \sum_{l \in \mathbb{Z}^d} a_l (C_1 - C_2 \rho_l) = C_1 \sum_{l \in \mathbb{Z}^d} a_l - C_2 \rho.$$

Then ρ is a Lyapunov function for $\mathcal{L}$.

Remark. We could take, as in section 3.2, $\rho_l(\omega) = |\omega_l| \chi(|\omega_l|)$ for a cut-off function χ . If $b^{(l)}$ satisfies

$$\frac{\langle \omega_l, b^{(l)}(\omega) \rangle}{|\omega_l|^2} \leq -C \quad \forall \omega_l \in \mathbb{R}^N,$$

and if the interaction coefficients $e^{(l)}$ are sufficiently small, then more generally ρ_l satisfies the assumption (4.21).

Remark. We can relax assumption (4.21) to

$$L^{(l)} \rho_l \leq C_1 - C_2 \rho_l,$$

and

$$e^{(l)} \cdot \nabla_l^{(l)} \rho_l \leq C_3 - \sum_{j \in \mathbb{Z}^d} \epsilon_{l,j} \rho_j,$$

for suitably small positive numbers $\epsilon_{l,j}$. With a little more effort, the proof below can be changed to accommodate this case.

We are ready to prove the main result of this section.

Theorem 4.3. *Let ρ be defined as above, and let $(P_t)_{t \geq 0}$ be the Markov semigroup with generator $\mathcal{L}$ defined in Theorem 4.2. Then, for any $\omega \in \Omega_\infty$ there exists a subsequence $(P_{t_j})_{j \geq 0}$, and a probability measure ν_ω such that for all $f \in C(\Omega)$ bounded cylinder functions, we have*

$$P_{t_j} f(\omega) \rightarrow \int f d\nu_\omega \quad \text{as } j \rightarrow \infty.$$

Furthermore, $\nu_\omega(\Omega_\infty) = 1$.

Proof. Fix $\omega \in \Omega_\infty$ and define

$$p_t^\omega(A) := P_t(\mathbb{1}_A)(\omega).$$

Using the same approach as in Lemma 3.6, from (4.22) we can show that there exists a $C > 0$ such that

$$(4.23) \quad P_t \rho(\omega) \leq C \quad \text{for all } t > 0.$$

As in the proof of Theorem 3.9, using (4.23) and Prokhorov's theorem, we deduce that there exists a subsequence $(p_{t_j}^\omega)_{j \geq 0}$ which converges weakly to some probability measure ν_ω .

Moreover, using Markov's inequality we have for any $r > 0$ that

$$0 \leq 1 - \nu_\omega(\Omega_r) \leq \frac{1}{r} \int \rho d\nu_\omega = \frac{1}{r} \sup_{l \in \mathbb{N}} P_{t_l} \rho(\omega) \leq \frac{C}{r},$$

where in the last step we used (4.23). Taking $r \rightarrow \infty$, we obtain $\nu_\omega(\Omega_\infty) = 1$. $\square$

For any $\epsilon > 0$ for which we have $\sum_{l \in \mathbb{Z}^d} (1 + |l|)^{-\epsilon} < \infty$, define also the set

$$\tilde{\Omega}_\epsilon := \left\{ \omega \in \Omega : \sum_{l \in \mathbb{Z}^d} \frac{|\omega_l|}{(1 + |l|)^\epsilon} < \infty \right\}.$$

Note that for ρ_l defined as in the previous remark, and with $a_l = (1 + |l|)^{-\epsilon}$, then $\tilde{\Omega}_\epsilon = \Omega_\infty$.

Remark. From the proof below it will be clear that instead of the weights $(1 + |l|)^{-\epsilon}$ we could take more general weights in the definition of $\tilde{\Omega}_\epsilon$, even of exponential decay.

Theorem 4.4. *Assume that $\gamma < \frac{1}{2}$. Let $(P_t)_{t \geq 0}$ be the Markov semigroup defined in Theorem 4.2 and assume in addition that*

$$\tilde{\eta} := \sup_{l \in \mathbb{Z}^d} \tilde{\eta}_l < 0,$$

and

$$\tilde{C} := \sup_{l \in \mathbb{Z}^d} C_l \leq -2\tilde{\eta},$$

where the constants $\tilde{\eta}_l$ and C_l were defined in (4.15) and (4.17), respectively.

For any bounded cylinder function f and any $\omega, \omega' \in \tilde{\Omega}_\epsilon$, there exists $t_0 > 0$ and a constant $C(f, \omega, \omega') < \infty$, such that

$$|P_t f(\omega) - P_t f(\omega')| \leq C(f, \omega, \omega') e^{-ct},$$

where $c > 0$ is independent of f, t, ω, ω' .

Proof. Fix a cylinder function f and $\omega, \omega' \in \tilde{\Omega}_\epsilon$. Fix $\Lambda \subset \mathbb{Z}^d$ finite and such that $\Lambda(f) \subset \Lambda$. We then have

$$(4.24) \quad \begin{aligned} & |P_t f(\omega) - P_{t_j} f(\omega')| \\ & \leq |P_t f(\omega) - P_t^\Lambda f(\omega)| + |P_t^\Lambda f(\omega) - P_t^\Lambda f(\omega')| + |P_t^\Lambda f(\omega') - P_t f(\omega')|. \end{aligned}$$

Using the computations in (4.20), there exists $r > 0$ such that if $\Lambda = B_{rt}(0) = \{l \in \mathbb{Z}^d : |l| < rt\}$, and letting $\Lambda' \rightarrow \infty$, then we have

$$(4.25) \quad \|P_t f - P_t^\Lambda f\|_\infty \leq C_1(f) e^{-c_1 t},$$

which holds for all $t \geq t_0$, where t_0 is chosen such that $\Lambda(f) \subset \Lambda$. This deals with the first and the third terms on the right hand side of the inequality (4.24) above. We are thus only left to study the middle term.

If ω and ω' differ by just one coordinate, say $\omega_j = \omega'_j$ for all $j \neq l$, then let $\gamma_{\omega_l, \omega'_l}(s) = \omega_l + s(\omega'_l - \omega_l)$. We can then compute

$$\begin{aligned} |P_t^\Lambda f(\omega) - P_t^\Lambda f(\omega')| &= \left| \int_0^1 \nabla^{(l)}(P_t^\Lambda f)(\gamma_{\omega_l, \omega'_l}(s)) \cdot (\omega'_l - \omega_l) ds \right| \\ &\leq |\omega'_l - \omega_l| \int_0^1 |\nabla^{(l)}(P_t^\Lambda f)(\gamma_{\omega_l, \omega'_l}(s))| ds \\ &\leq |\omega'_l - \omega_l| \left\| \tilde{\Gamma}^{(l)}(P_t^\Lambda f) \right\|_\infty^{1/2} \\ &\quad + \sqrt{2} |\omega'_l - \omega_l| \sum_{\alpha \in R_+} k_\alpha \int_0^1 \left| \frac{P_t^\Lambda f(\gamma_{\omega_l, \omega'_l}(s)) - P_t^\Lambda f(\sigma_\alpha(\gamma_{\omega_l, \omega'_l}(s)))}{\langle \alpha, \gamma_{\omega_l, \omega'_l}(s) \rangle} \right|. \end{aligned}$$

Using the same method as in Step 2 in the proof of Theorem 3.9, since $\gamma < \frac{1}{2}$, then we obtain

$$(4.26) \quad |P_t^\Lambda f(\omega) - P_t^\Lambda f(\omega')| \leq C_2 |\omega_l - \omega'_l| \left\| \tilde{\Gamma}^{(l)}(P_t^\Lambda f) \right\|_\infty^{1/2}.$$

Going back to the proof of (4.16), note that we have in fact

$$\left\| \tilde{\Gamma}^{(l)}(f_t) \right\|_\infty \leq e^{2\tilde{\eta}t} \left\| \tilde{\Gamma}^{(l)} f \right\|_\infty + |G| \sum_{j \in \Lambda} \tilde{E}_{l,j} \int_0^t \left\| \tilde{\Gamma}^{(j)}(f_u) \right\|_\infty du.$$

Using the same iteration procedure as above, this implies that

$$(4.27) \quad \left\| \tilde{\Gamma}^{(l)}(f_t) \right\|_\infty \leq e^{2\tilde{\eta}t + tC_l} \sum_{j \in \mathbb{Z}^d} \left\| \tilde{\Gamma}^{(j)}(f) \right\|_\infty.$$

By our assumption $2\tilde{\eta} + C_l \leq 2\tilde{\eta} + \tilde{C} < 0$, so this inequality provides the sufficient decay required in (4.26).

We now turn to the case of general $\omega, \omega' \in \tilde{\Omega}_\epsilon$. First, we note that if $\Lambda(f) \subset \Lambda$, then by (4.16) we have $\Lambda(P_t^\Lambda f) \subset \Lambda^R := \{l \in \mathbb{Z}^d : d(l, \Lambda) < R\}$, where R is the range of interaction. But Λ^R is a finite set, say $|\Lambda^R| = n$, and let $l_1, \dots, l_n$ be its distinct elements. We construct the sequence $\omega^0 = \omega$, $\omega_j^{i+1} = \omega_j^i$ for all $j \neq l_i$, and $\omega_{l_i}^{i+1} = \omega'_{l_i}$. In other words, each two consecutive terms of the sequence differ in just one coordinate, l_i , and $\omega_j^n = \omega'_j$ for all $j \in \Lambda^R$. Thus, by the observation above, $P_t^\Lambda(\omega^n) = P_t^\Lambda(\omega')$, and so we have

$$|P_t^\Lambda f(\omega) - P_t^\Lambda f(\omega')| \leq \sum_{i=0}^{n-1} |P_t^\Lambda f(\omega^{i+1}) - P_t^\Lambda f(\omega^i)|.$$

We estimate each term in this sum using the same method as above. More precisely, using (4.26) and (4.27), we have

$$(4.28) \quad \begin{aligned} \sum_{i=0}^{n-1} |P_t^\Lambda f(\omega^{i+1}) - P_t^\Lambda f(\omega^i)| &\leq C_2 \sum_{i=0}^{n-1} |\omega_{l_i} - \omega'_{l_i}| e^{(\tilde{\eta} + \tilde{C}/2)t} \left(\sum_{j \in \mathbb{Z}^d} \left\| \tilde{\Gamma}^{(j)}(f) \right\|_\infty \right)^{1/2} \\ &\leq C_3(f) e^{(\tilde{\eta} + \tilde{C}/2)t} \sum_{l \in B_{R+tR}(0)} \frac{|\omega_l| + |\omega'_l|}{(1 + |l|)^\epsilon} (1 + |l|)^\epsilon \\ &\leq C_3(f) (C_\omega + C_{\omega'}) e^{(\tilde{\eta} + \tilde{C}/2)t} (1 + |R + rt|)^\epsilon \end{aligned}$$

where $C_\omega = \sum_{l \in \mathbb{Z}^d} \frac{|\omega_l|}{(1 + |l|)^\epsilon}$ and $C_{\omega'} = \sum_{l \in \mathbb{Z}^d} \frac{|\omega'_l|}{(1 + |l|)^\epsilon}$ which are both finite since $\omega, \omega' \in \tilde{\Omega}_\epsilon$.

From (4.25) and (4.28) we obtain the decay we require. $\square$

Acknowledgements. The author wishes to thank Boguslaw Zegarlinski for introducing him to the problem and for useful advice. Financial support from EPSRC is also gratefully acknowledged.

REFERENCES

- [1] J.-Ph. Anker. An introduction to Dunkl theory and its analytic aspects. In *Analytic, algebraic and geometric aspects of differential equations*, Trends Math., pages 3–58. Birkhäuser/Springer, Cham, 2017.
- [2] E. B. Davies. *One-parameter semigroups*, volume 15 of *London Mathematical Society Monographs*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York, 1980.

- [3] F. Dragoni, V. Kontis, and B. Zegarliński. Ergodicity of Markov semigroups with Hörmander type generators in infinite dimensions. *Potential Anal.*, 37(3):199–227, 2012.
- [4] C. F. Dunkl. Differential-difference operators associated to reflection groups. *Trans. Amer. Math. Soc.*, 311(1):167–183, 1989.
- [5] P. Graczyk, M. Rösler, and M. Yor, editors. *Harmonic and stochastic analysis of Dunkl processes*. Travaux en cour. Herman, Paris, 2008.
- [6] A. Guionnet and B. Zegarliński. Lectures on logarithmic Sobolev inequalities. In *Séminaire de Probabilités, XXXVI*, volume 1801 of *Lecture Notes in Math.*, pages 1–134. Springer, Berlin, 2003.
- [7] O. Kallenberg. *Foundations of modern probability*. Probability and its Applications (New York). Springer-Verlag, New York, second edition, 2002.
- [8] V. Kontis, M. Ottobre, and B. Zegarliński. Markov semigroups with hypocoercive-type generator in infinite dimensions: ergodicity and smoothing. *J. Funct. Anal.*, 270(9):3173–3223, 2016.
- [9] T. M. Liggett. *Interacting particle systems*, volume 276 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1985.
- [10] M. Rösler. Dunkl operators: theory and applications. In *Orthogonal polynomials and special functions (Leuven, 2002)*, volume 1817 of *Lecture Notes in Math.*, pages 93–135. Springer, Berlin, 2003.
- [11] M. Rösler and M. Voit. Markov processes related with Dunkl operators. *Adv. in Appl. Math.*, 21(4):575–643, 1998.
- [12] J. F. van Diejen and L. Vinet, editors. *Calogero-Moser-Sutherland models*, CRM Series in Mathematical Physics. Springer-Verlag, New York, 2000.
- [13] L. Xu and B. Zegarliński. Ergodicity of the finite and infinite dimensional α -stable systems. *Stoch. Anal. Appl.*, 27(4):797–824, 2009.
- [14] L. Xu and B. Zegarliński. Existence and exponential mixing of infinite white α -stable systems with unbounded interactions. *Electron. J. Probab.*, 15:no. 65, 1994–2018, 2010.
- [15] B. Zegarliński. The strong decay to equilibrium for the stochastic dynamics of unbounded spin systems on a lattice. *Comm. Math. Phys.*, 175(2):401–432, 1996.
- [16] B. Zegarliński. Linear and nonlinear dissipative dynamics. *Rep. Math. Phys.*, 77(3):377–397, 2016.

ANDREI VELICU, DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, HUXLEY BUILDING, 180 QUEEN’S GATE, LONDON SW7 2AZ, UNITED KINGDOM *and* INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UNIVERSITÉ PAUL SABATIER, 118 ROUTE DE NARBONNE, 31062 TOULOUSE, FRANCE
E-mail address: `andrei.velicu@math.univ-toulouse.fr`