ON GEOMETRY OF NUMBERS AND UNIFORM RATIONAL APPROXIMATION TO THE VERONESE CURVE

JOHANNES SCHLEISCHITZ

ABSTRACT. Consider the classical problem of rational simultaneous approximation to a point in \mathbb{R}^n . The optimal lower bound on the gap between the induced ordinary and uniform approximation exponents has been established by Marnat and Moshchevitin in 2018. Recently Nguyen, Poels and Roy provided information on the best approximating rational vectors to points where the gap is close to this minimal value. Combining the latter result with parametric geometry of numbers, we effectively bound the dual linear form exponents in the described situation. As an application, we slightly improve the upper bound for the classical exponent of uniform Diophantine approximation $\hat{\lambda}_n(\xi)$, for even $n \geq 4$. Unfortunately our improvements are small, for n = 4 only in the fifth decimal digit. However, the underlying method in principle can be improved with more effort to provide better bounds. We indeed establish reasonably stronger results for numbers which almost satisfy equality in the estimate by Marnat and Moshchevitin. We conclude with consequences on the classical problem of approximation to real numbers by algebraic numbers/integers of uniformly bounded degree.

Keywords: exponents of Diophantine approximation, regular graph, parametric geometry of numbers Math Subject Classification 2010: 11J13, 11J82

1. Consequences of a recent Theorem in geometry of numbers

Classical topics in Diophantine approximation are to study simultaneous rational approximation to points $\underline{\xi} = (\xi_1, \ldots, \xi_n) \in \mathbb{R}^n$ and to find small values of linear forms in $\underline{\xi}$ with integer coefficients. This naturally leads to investigation of classical exponents of Diophantine approximation. Denote by $\omega(\underline{\xi})$ the (possibly infinite) supremum of ω such that

(1)
$$Y_{\underline{x}} := \max_{1 \le i \le n} |x\xi_i - y_i| \le x^{-\omega}$$

has infinitely many integer vector solutions $\underline{x} = (x, y_1, \ldots, y_n) \in \mathbb{Z}^{n+1}$. This is usually referred to as the ordinary exponent of simultaneous approximation. Similarly, let the (possibly infinite) uniform exponent of simultaneous approximation denoted by $\widehat{\omega}(\underline{\xi})$ be the supremum of ω such that

(2)
$$0 < x \le X, \qquad Y_x \le X^{-\omega}$$

Middle East Technical University, Northern Cyprus Campus, Kalkanli, Güzelyurt jschleischitz@outlook.com, johannes@metu.edu.tr.

has a solution $\underline{x} \in \mathbb{Z}^{n+1}$ for all large X. For linear forms, let $\omega^*(\underline{\xi})$ and $\widehat{\omega}^*(\underline{\xi})$ respectively be the (possibly infinite) supremum of ω^* such that

$$0 < \max_{1 \le j \le n} |a_j| \le X, \qquad |a_0 + a_1\xi_1 + \dots + a_n\xi_n| \le X^{-\omega^*}$$

has a solution in integers a_j for certain arbitrarily large X and all large X, respectively. Variants of Dirichlet's Box principle yield the lower bounds

(3)
$$\omega(\underline{\xi}) \ge \widehat{\omega}(\underline{\xi}) \ge \frac{1}{n}, \qquad \omega^*(\underline{\xi}) \ge \widehat{\omega}^*(\underline{\xi}) \ge n$$

for any $\underline{\xi} \in \mathbb{R}^n$. Finding refined relations between these exponents is an important topic in Diophantine approximation. Assume in the sequel that $\underline{\xi}$ is linearly independent over \mathbb{Q} together with {1}. Marnat and Moshchevitin [16] recently proved a remarkable (sharp) improvement of the trivial inequalities $\omega(\underline{\xi}) \geq \widehat{\omega}(\underline{\xi})$ and $\omega^*(\underline{\xi}) \geq \widehat{\omega}^*(\underline{\xi})$ that had been conjectured by Schmidt and Summerer [30]. The first can be stated as

(4)
$$\widehat{\omega}(\underline{\xi}) + \frac{\widehat{\omega}(\underline{\xi})^2}{\omega(\underline{\xi})} + \dots + \frac{\widehat{\omega}(\underline{\xi})^n}{\omega(\underline{\xi})^{n-1}} \le 1,$$

where we mean that $\widehat{\omega}(\underline{\xi}) = 1$ implies $\omega(\underline{\xi}) = \infty$. The sharpness of the estimate had previously been settled by Roy who constructed the equality case for any possible parameter pair in [22]. We further want to refer to his reasonably more general, deep results in [23]. We do not require the dual linear form estimate in this paper, but rather want to refer to [29], [10] for a weaker inequality that had been established previously, and the PhD thesis of Rivard-Cooke [19] for a simplified proof of (4) and a conjectural generalization. Equality in (4) is obtained in a case that Schmidt and Summerer in [30] refer to as the regular graph. We omit the geometrical motivation behind it, but point out that in this scenario the logarithms of the numbers x realizing the exponent $\omega(\underline{\xi})$ in (1) (for appropriate y_i) form almost a geometric sequence with ratio $\omega(\underline{\xi})/\widehat{\omega}(\underline{\xi})$. The recent preprint by Nguyen, Poels and Roy [17] provides a clearer picture by essentially showing that if $\omega(\underline{\xi}), \widehat{\omega}(\underline{\xi})$ almost satisfy identity in the estimate (4), then similar properties must still be satisfied. Indeed this information is provided in a compact form in Theorem 1.4 from [17], which we rephrase below upon omitting some subtle additional information not required here. With $Y_{\underline{x}}$ defined in (1), for a parameter X > 1 we derive the quantity

$$\mathscr{L}_{\xi}(X) = \min\{Y_{\underline{x}} : 1 \le x \le X\},\$$

where the minimum is taken over all integer vectors $\underline{x} = (x, y_1, \ldots, y_n)$ with first coordinate positive and not exceeding X. The induced sequence of vectors \underline{x} realizing the minimum for some X are sometimes referred to as best approximations or minimal points.

Theorem 1.1 (Nguyen, Poels, Roy). Let $n \ge 1$ be an integer. Let $\underline{\xi} \in \mathbb{R}^n$ with \mathbb{Q} -linearly independent coordinates together with $\{1\}$. Suppose that there exist positive real numbers a, b, α, β such that for all large enough X we have

(5)
$$bX^{-\beta} \le \mathscr{L}_{\underline{\xi}}(X) \le aX^{-\alpha}.$$

Then $\alpha \leq \beta$ and

(6)
$$\epsilon = \epsilon_{\alpha,\beta} := 1 - \left(\alpha + \frac{\alpha^2}{\beta} + \dots + \frac{\alpha^n}{\beta^{n-1}}\right) \ge 0.$$

If

(7)
$$\epsilon \leq \frac{1}{4n} \left(\frac{\alpha}{\beta}\right)^n \min\{\alpha, \beta - \alpha\},$$

then there exists an unbounded sequence of integer vectors $\underline{x}_j = (x_j, y_{j,1}, \dots, y_{j,n})$ with the properties

- $|\alpha \log x_{j+1} \beta \log x_j| \le C + 4\epsilon (\beta/\alpha)^n \log x_{j+1}$ $|\log Y_{\underline{x}_j} + \beta \log x_j| \le C + 4\epsilon (\beta/\alpha)^2 \log x_j$

- the vectors $\underline{x}_j, \underline{x}_{j+1}, \dots, \underline{x}_{j+n}$ are linearly independent There is no vector $\underline{x} = (x, y_1, \dots, y_n)$ so that $1 \le x \le x_j$ and $Y_{\underline{x}} < Y_{\underline{x}_j}$

Clearly assumption (5) implies

$$\alpha \le \widehat{\omega}(\underline{\xi}) \le \omega(\underline{\xi}) \le \beta,$$

and conversely any $\underline{\xi}$ satisfies (5) for any $\alpha < \widehat{\omega}(\underline{\xi})$ and $\beta > \omega(\underline{\xi})$ with suitable a, b. We will thus occasionally identify α and β with $\widehat{\omega}(\xi)$ and $\omega(\xi)$, respectively. Then $\epsilon = 0$ corresponds to equality in (4). In the situation of Theorem 1.1, we use the description of best approximating vectors to provide estimates on the dual exponents $\omega^*(\xi), \widehat{\omega}^*(\xi)$.

Theorem 1.2. Let $n \ge 1$ be an integer. Assume $\underline{\xi} = (\xi_1, \ldots, \xi_n)$ is \mathbb{Q} -linearly independent coordinates together with $\{1\}$ and satisfies condition (5) of Theorem 1.1 with given a, b, α, β . Assume ϵ as in (6) satisfies (7), and let

$$\phi := \frac{4\epsilon\beta^{n-1}}{\alpha^n}, \qquad \rho := \frac{4\epsilon\beta^2}{\alpha^2}$$

Then we have

(8)
$$\widehat{\omega}^*(\underline{\xi}) \ge \frac{(\beta - \rho)S}{(\frac{\alpha}{\beta} - \phi)^{-n} + (\beta - \rho)(1 - S)}, \qquad S := \sum_{j=1}^n (\frac{\alpha}{\beta} + \phi)^{1-j}.$$

Moreover

(9)
$$\omega^*(\underline{\xi}) \ge \frac{\rho^2 - \beta^2 - (\beta + \rho)^2 T}{\rho - \beta + (\beta + \rho)^2 T}, \qquad T := \sum_{j=1}^{n-1} (\frac{\alpha}{\beta} + \phi)^j.$$

Conversely we have the upper bounds

(10)
$$\widehat{\omega}^*(\underline{\xi}) \le (\beta - \rho)^{-1} (\frac{\alpha}{\beta} - \phi)^{-n},$$

and

(11)
$$\omega^*(\underline{\xi}) \le (\beta - \rho)^{-1} (\frac{\alpha}{\beta} - \phi)^{-n-1}.$$

All bounds in Theorem 1.2 can in principle be improved as will be indicated in its proof, thereby implying better bounds in Theorem 2.2 below as well. However we do not attempt to optimize the method as it would lead to a significantly more cumbersome proof. Theorem 1.2 becomes interesting for $n \geq 3$. For n = 1 clearly $\omega^*(\xi) = \omega(\xi)$ and a short argument by Khintchine [12] shows $\widehat{\omega}(\xi) = \widehat{\omega}^*(\xi) = 1$ for any $\xi \in \mathbb{R} \setminus \mathbb{Q}$. For n = 2,

as soon as $1, \xi_1, \xi_2$ are linearly independent over \mathbb{Q} , we have Jarník's identity [11] given as $\widehat{\omega}^*(\xi_1, \xi_2) = (1 - \widehat{\omega}(\xi_1, \xi_2))^{-1}$ and the sharp estimates

$$\frac{\omega(\xi_1,\xi_2) + \widehat{\omega}(\xi_1,\xi_2)}{1 - \widehat{\omega}(\xi_1,\xi_2)} \le \omega^*(\xi_1,\xi_2) \le \frac{\omega(\xi_1,\xi_2)}{\widehat{\omega}(\xi_1,\xi_2) - \omega(\xi_1,\xi_2) + \omega(\xi_1,\xi_2)\widehat{\omega}(\xi_1,\xi_2)}$$

due to Laurent [14], where the right inequality requires a positive right hand side (which is true upon equality in (4)) and otherwise $\omega^*(\xi_1, \xi_2) = \infty$ is possible. For additional information in the case n = 2, we refer to Roy [24] and Schmidt and Summerer [31]. Concretely, our classical exponents can be interpreted as extremal values of the first and last successive minimum of some lattice point problem from the geometry of numbers. The papers [24, 31] provide additional information on the missing second successive minimum, not reflected by $\omega, \hat{\omega}, \omega^*, \hat{\omega}^*$. See also the introduction of Section 3 for the correspondence geometry of numbers. For larger n, sharp estimates linking $\omega(\underline{\xi})$ and $\omega^*(\underline{\xi})$ on the one hand and $\hat{\omega}(\underline{\xi})$ and $\hat{\omega}^*(\underline{\xi})$ on the other hand are due to Khintchine (see Theorem B5 in [2]) and German [8] respectively, however they do not take into account the additional information in Theorem 1.2 on another exponent. As a consequence, they do not provide any upper bound for $\omega^*(\underline{\xi})$ (or $\hat{\omega}^*(\underline{\xi})$) if $\omega(\underline{\xi}) \geq 1/(n-1)$ (or $\hat{\omega}(\underline{\xi}) \geq 1/(n-1)$), and indeed these exponents may take the value $+\infty$ when no further restriction is imposed. See however [4] for refinements of Khintchine's estimates that contain triples $(\omega, \hat{\omega}, \omega^*)$ resp. $(\omega, \omega^*, \hat{\omega}^*)$ of classical exponents.

We also should point out that Theorem 1.2 is mainly of interest when ϵ in (6) is very small, when we increase ϵ the inequalities (8), (9) quickly become worse than the bounds in (3). Consequently, the numerical improvements in Theorem 2.2 below will be small.

A consequence of Theorem 1.2 is that equality in (4) is sufficient for all classical exponents to attain the values as in the corresponding regular graph, and by continuity reasons they cannot differ much from them if the error in (4) is sufficiently small.

Corollary 1.3. Assume $\underline{\xi} = (\xi_1, \ldots, \xi_n)$ induces equality in (4). Then

$$\widehat{\omega}^*(\underline{\xi}) = \frac{\omega(\underline{\xi})^{n-1}}{\widehat{\omega}(\underline{\xi})^n}, \qquad \omega^*(\underline{\xi}) = \frac{\omega(\underline{\xi})^n}{\widehat{\omega}(\underline{\xi})^{n+1}}$$

Moreover, for any $\varepsilon > 0$ and c < 1, if we identify $\alpha = \widehat{\omega}(\underline{\xi})$ and $\beta = \omega(\underline{\xi})$, upon $\alpha \leq c$ there exists $\delta = \delta(n, c) > 0$ such that the estimate $\epsilon = \epsilon_{\alpha,\beta} < \delta$ implies

$$\frac{\omega(\underline{\xi})^{n-1}}{\widehat{\omega}(\underline{\xi})^n} - \varepsilon \le \widehat{\omega}^*(\underline{\xi}) \le \frac{\omega(\underline{\xi})^{n-1}}{\widehat{\omega}(\underline{\xi})^n} + \varepsilon, \qquad \frac{\omega(\underline{\xi})^n}{\widehat{\omega}(\underline{\xi})^{n+1}} - \varepsilon \le \omega^*(\underline{\xi}) \le \frac{\omega(\underline{\xi})^n}{\widehat{\omega}(\underline{\xi})^{n+1}} + \varepsilon.$$

Unfortunately δ depends in a sensitive way on ϵ (that is, on α, β), concordant with the strong dependence of the bounds in Theorem 1.2 on ϵ .

2. An application to the Veronese curve

We consider $\underline{\xi} = (\xi, \xi^2, \dots, \xi^n)$ on the Veronese curve, where we assume ξ is transcendental. As customary we write

(12)
$$\lambda_n(\xi) = \omega(\underline{\xi}), \quad \widehat{\lambda}_n(\xi) = \widehat{\omega}(\underline{\xi}), \quad w_n(\xi) = \omega^*(\underline{\xi}), \quad \widehat{w}_n(\xi) = \widehat{\omega}^*(\underline{\xi}),$$

5

for the intensely studied classical exponents of Diophantine approximation (note: the classical definition of $w_n(\xi)$, $\hat{w}_n(\xi)$ differs if ξ is algebraic of degree $\leq n$, by excluding vanishing polynomial evaluations at ξ). For n = 1 we just have $\lambda_1(\xi) = \omega(\xi) = \omega(\xi)$, and similarly for the other exponents. The exponents $w_n(\xi)$ date back to Mahler [15], others have been defined in [3]. For n = 1 we have $\hat{\lambda}_1(\xi) = 1$ for any irrational ξ , as already noticed in Section 1. The consequence $\hat{\omega}(\underline{\xi}) \leq 1$ for any $\underline{\xi} \notin \mathbb{Q}^n$ agrees with (4). Without restriction to the Veronese curve the uniform exponent $\hat{\omega}(\underline{\xi})$ attains the maximum value 1 for certain $\underline{\xi} \in \mathbb{R}^n$ with \mathbb{Q} -linearly independent coordinates with $\{1\}$ no matter how large n is (see Poels [18] for general classes of manifolds containing points with this property, and further references). On the other hand, on the Veronese curve the exponent turns out to be always significantly smaller. Only for n = 2 the optimal bound is known, given as

(13)
$$\widehat{\lambda}_2(\xi) \le \frac{\sqrt{5}-1}{2} = 0.6180\dots$$

The inequality was found by Davenport and Schmidt [7], the optimality is due to Roy [20]. For even n the following estimates are from [27].

Theorem 2.1 (Schleischitz). Let $n \ge 2$ be an even integer. For any transcendental real number ξ we have $\widehat{\lambda}_n(\xi) \le \tau_n$ where τ_n is the solution of

$$(\frac{n}{2})^n t^{n+1} - (\frac{n}{2} + 1)t + 1 = 0$$

in the interval $(\frac{2}{n+2}, \frac{2}{n})$.

The bound τ_n is obtained as the solution for identity in (4) for $\hat{\lambda}_n(\xi)$ when $\lambda_n(\xi) = 2/n$. For n = 2 we confirm the optimal estimate (13). Other numerical bounds are

(14)
$$\widehat{\lambda}_4(\xi) \le 0.370635\ldots, \quad \widehat{\lambda}_6(\xi) \le 0.268186\ldots, \quad \widehat{\lambda}_{20}(\xi) \le 0.092803\ldots$$

For large n, the bound is of order

(15)
$$\tau_n = \frac{2}{n} - \frac{\chi}{n^2} + o(n^{-2}), \qquad n \to \infty,$$

where $\chi = 3.18...$ can be expressed as a zero of some power series, see [27]. For sake of completeness, we remark that for the case of odd n the bound $\hat{\lambda}_n(\xi) \leq 2/(n+1)$ was established by Laurent [13], an improvement for n = 3 is due to Roy [21]. Application of (8) from our new Theorem 1.2 leads to a small improvement that can be stated in the following way.

Theorem 2.2. Let $n \ge 4$ be an even integer and ξ be any transcendental real number. For $\alpha \in [1/n, 1]$ define

$$\epsilon_{\alpha} = 1 - \alpha - \frac{n}{2}\alpha^2 - \dots - \left(\frac{n}{2}\right)^{n-1}\alpha^n, \qquad S_{\alpha} = \sum_{j=1}^n \left(\frac{n\alpha}{2} + \frac{2^{n+1}\epsilon_{\alpha}}{n^{n-1}\alpha^n}\right)^{1-j}.$$

Then $\widehat{\lambda}_n(\xi) \leq \sigma_n$ with σ_n the solution of the implicit equation

(16)
$$\frac{\left(\frac{2}{n} - \frac{16\epsilon_{\alpha}}{n^{2}\alpha^{2}}\right)S_{\alpha}}{\left(\frac{n\alpha}{2} - \frac{2^{n+1}\epsilon_{\alpha}}{n^{n}\alpha^{n}}\right)^{-n} + \left(\frac{2}{n} - \frac{16\epsilon_{\alpha}}{n^{2}\alpha^{2}}\right)(1 - S_{\alpha})} = \mu_{n} := \max\{2n - 2, w(n)\},$$

for α in the interval (0,1), where w(n) is the solution of

$$\frac{(n-1)w}{w-n} - w + 1 = \left(\frac{n-1}{w-n}\right)^n$$

in the interval [n, 2n-1). Precisely for $n \ge 10$ we get $\mu_n = 2n-2$.

The left hand side in (16) is (8) with $\beta = 2/n$. It can be checked and follows from the proof that $\sigma_n < \tau_n$ for $n \ge 4$. However, the quantities differ only by a very small amount, for example we obtain the numerical bounds

$$\widehat{\lambda}_4(\xi) \le 0.370629\dots, \qquad \widehat{\lambda}_6(\xi) \le 0.268183\dots,$$

that may be compared with (14). The asymptotics (15) remain unaffected and the improvement occurs in a lower order term. Improvements can be made via improving the estimate (8) in Theorem 1.2. Another source of improvement would be better bounds for the exponent $\hat{w}_n(\xi)$, related to μ_n in the theorem. On the other hand, increasing the bound for ϵ in (7) would not lead to stronger bounds when combined with Theorem 1.2 in its present form.

The sensitive dependence of (8), (9), (10), (11) on $\epsilon = \epsilon_{\alpha,\beta}$ is the key problem why the improvement compared to Theorem 2.1 is small. We wonder about estimates in the optimal case $\epsilon = 0$. This is partly motivated by the fact that for n = 2 identity in (13) is attained (only) for numbers ξ where (ξ, ξ^2) induces a regular graph $\epsilon = 0$, see [20]. We also include a result for the case where the difference from the regular graph is very small, derived by continuity reasoning. Mind the difference between ϵ and ε therein.

Theorem 2.3. Let $n \ge 4$ be even and let $\xi \in \mathbb{R}$. First assume $\underline{\xi} = (\xi, \xi^2, \dots, \xi^n)$ induces equality in (4). If $n \in \{4, 6, 8\}$, then

(17)
$$\widehat{\lambda}_4(\xi) < 0.3588, \quad \widehat{\lambda}_6(\xi) < 0.2540, \quad \widehat{\lambda}_8(\xi) < 0.1968,$$

and as $n \to \infty$ we have the asymptotical bound

(18)
$$\widehat{\lambda}_n(\xi) \le \frac{\Theta + o(1)}{n}, \qquad n \to \infty,$$

where $\Theta = 1.7564...$ is the solution to $e^t/t = 2\sqrt{e}$ with t > 1.

When we drop equality assumption in (4), we still have the following continuity result. For $n \ge n_0$ and every $\varepsilon > 0$, there exists $\delta_n > 0$ such that if $\alpha = \widehat{\lambda}_n(\xi) \in [1/n, 1]$ and $\beta = \lambda_n(\xi) \in [1/n, \infty]$ are linked by $\epsilon = \epsilon_{\alpha,\beta} < \delta_n$, then we have

(19)
$$\widehat{\lambda}_n(\xi) \le \frac{\Theta + \varepsilon}{n}.$$

Unfortunately, similar to Corollary 1.3 the bound δ_n for the conclusion (19) is very small when ε is small. On the other hand, we remark that for n = 8 the bound in (17) is already smaller than the left interval end point 2/(n+2) = 0.2 from Theorem 2.1.

We derive a corollary on approximation to real numbers by algebraic integers. For an algebraic integer α denote by $H(\alpha)$ its height, i.e. the naive height (maximum modulus among its coefficients) of its irreducible minimal polynomial over $\mathbb{Z}[T]$. By a well-known argument that originates in work of Davenport and Schmidt [7], from Theorem 2.2 and Theorem 2.3 we infer

Corollary 2.4. Let $n \ge 4$ be an even integer, ξ a real transcendental number and $\varepsilon > 0$. Then for σ_n as in Theorem 2.2 the inequality

(20)
$$|\xi - \alpha| < H(\alpha)^{-\frac{1}{\sigma_n} - 1 + \varepsilon}$$

has infinitely many solutions in real algebraic integers α of degree at most n + 1 (or real algebraic numbers of degree precisely n). If $\underline{\xi} = (\xi, \xi^2, \dots, \xi^n)$ satisfies equality in (4), then for $n \ge n_0$ and Θ as in Theorem 2.3 the estimate

(21)
$$|\xi - \alpha| < H(\alpha)^{-\frac{n}{\Theta} - 1 + \varepsilon}$$

has infinitely many solutions α as above.

The deduction for algebraic integers of degree n + 1 is immediate from [7, Lemma 1]. For the implication on algebraic numbers of precise degree n, see Bugeaud and Teulié [6]. An unconditional proof of (21) would be highly desirable, which would improve on the currently best known bounds of order -n/2 - O(1) from [6, 7] for the exponent in (20) regarding both approximation by algebraic integers of degree n+1 and algebraic numbers of exact degree n. However, we remark that concerning approximation by algebraic numbers of degree at most n, a slightly stronger bound with exponent smaller than $-n/\sqrt{3} - 1 = -n/1.73... - 1$ has recently been settled in [1].

3. Proof of Theorem 1.2

We first introduce parametric geometry of numbers that our proof is based on. We follow the introductory paper of Schmidt and Summerer [28]. Let $n \ge 1$ an integer and $\underline{\xi} = (\xi_1, \ldots, \xi_n) \in \mathbb{R}^n$ be fixed. For a parameter Q > 1 and every $1 \le j \le n+1$, let $\psi_j(Q)$ be the infimum of ν for which the system

$$|x| \le Q^{1+\nu}, \qquad Y_x \le Q^{-1/n+\nu}$$

has j linearly independent integral solution vectors $\underline{x} = (x, y_1, \dots, y_n)$, with $Y_{\underline{x}}$ as in (1). Let $q = \log Q$ and define the functions $L_j(q)$ from $\psi_j(Q)$ via

$$\psi_j(Q) = \frac{L_j(q)}{q}, \qquad 1 \le j \le n+1.$$

It can be checked that $L_j(q)$ are piecewise linear with slopes among $\{-1, 1/n\}$. In fact, any $L_j(q)$ is locally induced by a function

(22)
$$L_{\underline{x}}(q) = \max\{\log x - q, \ \log Y_{\underline{x}} + \frac{q}{n}\}.$$

JOHANNES SCHLEISCHITZ

more precisely we may write

(23)
$$L_j(q) = \min \max_{1 \le i \le j} L_{\underline{x}_i}(q),$$

with minimum taken over all sets of j linearly independent vectors $\underline{x}_1, \ldots, \underline{x}_j \in \mathbb{Z}^{n+1}$. We infer

$$-1 \le \underline{\psi}_j \le \overline{\psi}_j \le \frac{1}{n}, \qquad 1 \le j \le n+1,$$

where we have put

$$\underline{\psi}_j = \liminf_{Q \to \infty} \psi_j(Q) = \liminf_{q \to \infty} \frac{L_j(q)}{q}, \qquad \overline{\psi}_j = \limsup_{Q \to \infty} \psi_j(Q) = \limsup_{q \to \infty} \frac{L_j(q)}{q},$$

Another important property of the functions highlighted in [28] is

(24)
$$\left|\sum_{j=1}^{n+1} L_j(q)\right| = O(1)$$

This reflects Minkowski's Second Convex Body Theorem. In particular, on long intervals on average one function $L_j(q)$ will decay with slope -1 whereas the remaining functions rise with slope 1/n. The next result linking the quantities $\underline{\psi}_{n+1}, \overline{\psi}_{n+1}$ with classical exponents originates in German's paper [9], see also Schmidt and Summerer [28].

Proposition 3.1. We have

(25)
$$(1+\widehat{\omega}^*(\underline{\xi})^{-1})(1+\underline{\psi}_{n+1}) = \frac{n+1}{n}.$$

and

(26)
$$(1 + \omega^*(\underline{\xi})^{-1})(1 + \overline{\psi}_{n+1}) = \frac{n+1}{n}.$$

Proof. Define $\omega_{n+1}(\underline{\xi})$ and $\widehat{\omega}_{n+1}(\underline{\xi})$ as the supremum of ω such that (2) has n+1 linearly independent solution vectors (x, y_1, \ldots, y_n) for arbitrarily large X and all large X, respectively. An application of Mahler's Theorem on Polar Convex Bodies implies $\widehat{\omega}^*(\underline{\xi})^{-1} = \omega_{n+1}(\underline{\xi})$ and similarly $\omega^*(\underline{\xi})^{-1} = \widehat{\omega}_{n+1}(\underline{\xi})$, as observed in [9, Corollary 8.5]. With these identifications, the identities just become a special case of [9, Proposition 7.1] (see also [28, Theorem 1.4] for a special case whose proof can be readily extended).

The following direct consequence of Theorem 1.1 is just formulated for convenience.

Proposition 3.2. Let $\underline{\xi}$ satisfy the hypothesis of Theorem 1.1 and consider the induced sequence $(\underline{x}_j)_{j\geq 1}$ with $\underline{x}_j = (x_j, y_{j,1}, \ldots, y_{j,n})$ from its claim. Then for $\ell \geq 0$ any fixed integer as $j \to \infty$ we have

(27)
$$\left(\left(\frac{\alpha}{\beta}-\phi\right)^{\ell}+o(1)\right)\log x_{j+\ell} \leq \log x_j \leq \left(\left(\frac{\alpha}{\beta}+\phi\right)^{\ell}+o(1)\right)\log x_{j+\ell}, \quad j\geq 1,$$

where

$$\phi := \frac{4\epsilon\beta^{n-1}}{\alpha^n}.$$

Moreover, again as $j \to \infty$ we have

(28) $(-\beta - \rho + o(1)) \log x_j \leq \log Y_j \leq (-\beta + \rho + o(1)) \log x_j, \qquad j \geq 1,$ where $4\epsilon\beta^2$

$$\rho := \frac{4\epsilon\beta^2}{\alpha^2}$$

Proof. The first property of Theorem 1.1 can be formulated

$$\log x_j = \left(\frac{\alpha}{\beta} + \phi_j + o(1)\right) \log x_{j+1}, \qquad j \to \infty,$$

with some $|\phi_j| \leq \phi$. The first claim (27) follows. The second claim (28) follows directly from the second property of Theorem 1.1.

In the proof, in accordance with notation in (1) we write

$$Y_j = \max_{1 \le i \le n} |x_j \xi_i - y_{j,i}|, \qquad j \ge 1,$$

when $\underline{x}_j = (x_j, y_{j,1}, \dots, y_{j,n})$ are the vectors from Theorem 1.1.

Proof of Theorem 1.2. We start with (8), which turns out to be the most tedious calculation. In view of (25), we have to provide lower bounds for $\underline{\psi}_{n+1}$. For $k \ge 1$, let q_k denote the minimum of $L_{\underline{x}_k}$. It obviously follows from Theorem 1.1 that $q_k < q_{k+1}$ for all large k. By the last property of Theorem 1.1 each q_k is a local minimum of L_1 , however they may not capture all such minima.

We estimate the first n functions $L_1(q), L_2(q), \ldots, L_n(q)$ from above in intervals $q \in I_k := [q_k, q_{k+1})$. Since $[q_m, \infty)$ is the disjoint union of these intervals I_k over $k \ge m$ and by (24)

(29)
$$\psi_{n+1}(e^q) = \frac{L_{n+1}(q)}{q} \ge -\sum_{j=1}^n \frac{L_j(q)}{q} - O(q^{-1}), \qquad q > 0,$$

this will lead to the desired lower bound for $\underline{\psi}_{n+1}$.

Notice now that the sets of vectors

$$\{\underline{x}_{k-n+1}, \underline{x}_{k-n+2}, \dots, \underline{x}_k\}, \qquad \{\underline{x}_{k-n+2}, \underline{x}_{k-n+3}, \dots, \underline{x}_{k+1}\}$$

are both linearly independent by the third claim of Theorem 1.1. By (23) we infer that for any $q \in [q_k, q_{k+1})$ every function $L_j(q)$ for $1 \leq j \leq n$ is bounded above by $L_{\underline{x}_g}(q)$ for some $k - n + 1 \leq g \leq k + 1$ (g depends on j and q). Concretely, we have

(30)
$$\sum_{j=1}^{n} \frac{L_j(q)}{q} \le \min\left\{\sum_{j=1}^{n} \frac{L_{\underline{x}_{k-n+j}}(q)}{q}, \sum_{j=2}^{n+1} \frac{L_{\underline{x}_{k-n+j}}(q)}{q}\right\}, \qquad q \in [q_k, q_{k+1}).$$

We have to estimate the right hand side in dependence of q. Let r_k be the intersection point of $L_{\underline{x}_k}$ and $L_{\underline{x}_{k+1}}$. The values r_k are closely connected to the local maxima of L_1 , but do not necessarily coincide with them. We have $q_k < r_k < q_{k+1}$. Keep in mind that by assumption $L_{\underline{x}_k}(r_k) = L_{\underline{x}_{k+1}}(r_k)$. First, we observe that we can restrict to intervals $[q_k, r_k]$. Indeed, the function $L_{\underline{x}_{k+1}}$ decreases with slope -1 in $(r_k, q_{k+1}]$, and since the others have slope at most 1/n the sum of all slopes is at most $-1+(n-1)\cdot n^{-1} = -1/n < 0$ within $(r_k, q_{k+1}]$. Since the expression in (30) cannot exceed -1/n + o(1) as $q \to \infty$ by (24) and $L_{n+1}(q) \leq q/n$, thus the sum decays in $(r_k, q_{k+1}]$. Hence the worst (largest) bound in (30) and thus the worst (smallest) bound in (29) is attained at the left endpoint r_k . Thus indeed we can restrict to $[q_k, r_k]$.

Next we further split the interval $[q_k, r_k]$ into two intervals $[q_k, s_k)$ and $[s_k, r_k]$ where s_k is the first coordinate of the point where $L_{\underline{x}_{k-n+1}}(q)$ meets $L_{\underline{x}_{k+1}}(q)$, by (22) that is the solution for q of

$$\log x_{k+1} - q = \log Y_{k-n+1} + \frac{q}{n},$$

which yields

(31)
$$s_k = \frac{n}{n+1} (\log x_{k+1} - \log Y_{k-n+1}).$$

In the interval $q \in [q_k, s_k]$ we estimate the left expression in (30), that is

(32)
$$\sum_{j=1}^{n} \frac{L_{\underline{x}_{k-n+j}}(q)}{q}$$

Since every $L_{\underline{x}_i}(q)$ reaches its minimum at q_i and the sequence $(q_i)_{i\geq 1}$ is clearly increasing, in this interval all involved functions increase with slope 1/n. Thus

$$L_{\underline{x}_i}(q_k) = \log Y_i + \frac{q_k}{n}, \qquad 1 \le i \le k,$$

and the sum (32) reaches its maximum at the right end point $q = s_k$ and we conclude

$$\max_{q \in [q_k, s_k]} \sum_{j=1}^n \frac{L_{\underline{x}_{k-n+j}}(q)}{q} \le \frac{\sum_{j=1}^n L_{\underline{x}_{k-n+j}}(s_k)}{s_k} \le \frac{\sum_{j=1}^n (\log Y_{k-n+j} + \frac{s_k}{n})}{s_k} = 1 + \frac{\sum_{j=1}^n \log Y_{k-n+j}}{s_k}.$$

Thus by (31) we infer

(33)
$$\max_{q \in [q_k, s_k]} \sum_{j=1}^n \frac{L_{x_{k-n+j}}(q)}{q} \le 1 + \frac{n+1}{n} \cdot \frac{\sum_{j=1}^n \log Y_{k-n+j}}{\log x_{k+1} - \log Y_{k-n+1}}$$

Finally, in the interval $q \in [s_k, r_k]$ we estimate the right expression in (30), that is

(34)
$$\sum_{j=2}^{n+1} \frac{L_{\underline{x}_{k-n+j}}(q)}{q}$$

Since in this interval $L_{\underline{x}_{k+1}}(q)$ decays with slope -1, again the sum of slopes in this interval is at most $-1 + (n-1) \cdot n^{-1} = -1/n < 0$. Thus the most disadvantageous case of a maximum in (30) is attained at the left endpoint s_k , which leads to the same bound as in (33) again.

This bound (33) remains to be estimated, which we perform via Theorem 1.1. We first readily verify that by $x_{k+1} > 1$ and $Y_i < 1$ the expression is increasing in all involved variables $x_{k+1}, Y_{k-n+1}, \ldots, Y_k$. Thus we have to find upper bounds for each variable.

By (30) and (28) applied to j from j = k - n + 1 up to j = k, we obtain (35)

$$\max_{q \in [q_k, q_{k+1}]} \sum_{j=1}^n \frac{L_j(q)}{q} \le 1 + \frac{n+1}{n} \cdot \frac{\sum_{j=1}^n (-\beta + \rho + o(1)) \log x_{k-n+j}}{\log x_{k+1} - (-\beta + \rho + o(1)) \log x_{k-n+1}}, \qquad k \to \infty.$$

Observe that $\rho < \beta$ as this is equivalent to $\alpha^2/\beta > 4\epsilon$, but since $n \ge 2$ and $0 < \alpha < \beta$ we compute

(36)
$$\frac{\alpha^2}{\beta} = \beta \left(\frac{\alpha}{\beta}\right)^2 \ge \beta \left(\frac{\alpha}{\beta}\right)^n \ge (\beta - \alpha) \left(\frac{\alpha}{\beta}\right)^n > \frac{\left(\frac{\alpha}{\beta}\right)^n (\beta - \alpha)}{n} \ge 4\epsilon,$$

where the most right inequality follows from (7). Hence we verify that this time the expression (35) is decreasing in x_{k-n+2}, \ldots, x_k but increasing in x_{k+1} . For x_{k-n+1} the situation is unclear, depending on the sign of $\log x_{k+1} - (\beta - \rho) \sum_{j=2}^{n} \log x_{k-n+j}$. Thus we want to find lower bounds for x_{k-n+2}, \ldots, x_k and upper bounds for x_{k+1} in terms of x_{k-n+1} .

It can be checked similar to (36) that $\alpha/\beta - \phi > 0$. By (27) we can estimate

$$\log x_{k-n+j} \ge \frac{\log x_{k-n+1}}{(\frac{\alpha}{\beta} + \phi)^{j-1} + o(1)}, \ 1 \le j \le n, \qquad \log x_{k+1} \le \frac{\log x_{k-n+1}}{(\frac{\alpha}{\beta} - \phi)^n + o(1)}$$

We remark that these are crude estimates that can be improved by a refined analysis of the interplay of the possible quotients of the occurring $\log x_j$, however calculations become rather cumbersome. Similar situations will occur more often below when we apply Proposition 3.1. Inserting these bounds in (35) we divide numerator and denominator by x_{n-k+1} to conclude

$$\max_{q \in [q_k, q_{k+1}]} \sum_{j=1}^n \frac{L_j(q)}{q} \le 1 + \frac{n+1}{n} \cdot \frac{\sum_{j=1}^n (-\beta + \rho + o(1)) \frac{1}{(\frac{\alpha}{\beta} + \phi)^{j-1} + o(1)}}{\frac{1}{(\frac{\alpha}{\beta} - \phi)^n + o(1)} + \beta - \rho - o(1)}$$
$$= 1 + \frac{n+1}{n} (-\beta + \rho) \cdot \frac{\sum_{j=1}^n \frac{1}{(\frac{\alpha}{\beta} + \phi)^{j-1}}}{\frac{1}{(\frac{\alpha}{\beta} - \phi)^n} + \beta - \rho} + o(1), \qquad k \to \infty.$$

Now we finally use (29) to conclude

$$\underline{\psi}_{n+1} = \liminf_{q \to \infty} \frac{L_{n+1}(q)}{q} = \liminf_{k \to \infty} \min_{q \in [q_k, q_{k+1})} \frac{L_{n+1}(q)}{q} \ge -1 + \frac{n+1}{n} (\beta - \rho) \cdot \frac{\sum_{j=1}^n \frac{1}{(\frac{\alpha}{\beta} + \phi)^{j-1}}}{\frac{1}{(\frac{\alpha}{\beta} - \phi)^n} + \beta - \rho}$$

Using (25) leads to the stated bound (8) after some rearrangements.

We turn to the lower bound (9) for $\omega^*(\underline{\xi})$. Here we look at the values $q = q_k$ where the functions $L_{\underline{x}_k}(q)$ attain their minimum values. Again since $(q_i)_{i\geq 1}$ increases, all

$$L_{\underline{x}_{k-n+1}}(q), \ L_{\underline{x}_{k-n+2}}(q), \ \dots, \ L_{\underline{x}_{k-1}}(q)$$

increase with slope 1/n at $q = q_k$. Hence

$$L_{\underline{x}_{k-n+j}}(q_k) = \log Y_{k-n+j} + \frac{q_k}{n}, \qquad 1 \le j \le n,$$

where for j = n we can use that representation as well as there is equality in the expressions of (22) for $L_{\underline{x}_k}(q)$ at $q = q_k$. From the linear dependence of $L_{\underline{x}_{k-n+1}}, L_{\underline{x}_{k-n+2}}, \ldots, L_{\underline{x}_{k-1}}$ by Theorem 1.1 and (29), if we let $Q_k = e^{q_k}$ we infer

$$\psi_{n+1}(Q_k) = \frac{L_{n+1}(q_k)}{q_k} \ge -\sum_{j=1}^n \frac{L_{\underline{x}_{k-n+j}}(q_k)}{q_k} - O(q_k^{-1}) = -\frac{\sum_{j=1}^n \log Y_{k-n+j}}{q_k} - 1 - O(q_k^{-1}).$$

Now q_k is by (22) given as the solution of

$$\log x_k - q_k = \log Y_k + \frac{q_k}{n}$$

hence

$$q_k = \frac{n}{n+1} \cdot (\log x_k - \log Y_k).$$

Plugging in yields

$$\psi_{n+1}(Q_k) \ge \frac{n+1}{n} \cdot \frac{\sum_{j=1}^n \log Y_{k-n+j}}{\log Y_k - \log x_k} - 1 - O(q_k^{-1}).$$

We see that the right hand side expression is decreasing in x_k and in $Y_{k-n+1}, \ldots, Y_{k-1}$, the situation is unclear for Y_k depending on the sign of $\log x_k + \sum_{j=1}^{n-1} \log Y_j$. Thus we look for lower bounds for the other variables in terms of Y_k . By (28) we have

$$\psi_{n+1}(Q_k) \ge \frac{n+1}{n} \cdot \frac{\log Y_k - (\beta + \rho + o(1)) \sum_{j=1}^{n-1} \log x_{k-n+j}}{\log Y_k (1 + \frac{1}{\beta + \rho}) + o(1)} - 1, \qquad k \to \infty.$$

Now this expression is increasing in Y_k so again by (28) we can estimate

$$\psi_{n+1}(Q_k) \ge \frac{n+1}{n} \cdot \frac{(\rho - \beta + o(1))\log x_k - (\beta + \rho + o(1))\sum_{j=1}^{n-1}\log x_{k-n+j}}{(\rho - \beta + o(1))(1 + \frac{1}{\beta + \rho})\log x_k} - 1.$$

Since $\rho - \beta < 0$, the right hand side is increasing in $x_{k-n+1}, \ldots, x_{k-1}$. Thus by (27) we can replace the sum in the numerator expression by T + o(1) with

$$T := \sum_{j=1}^{n-1} (\frac{\alpha}{\beta} + \phi)^j,$$

without making the expression larger. Dividing denominator and numerator by $\log x_k$ and dropping the negligible lower order terms yields

$$\overline{\psi}_{n+1} \ge \limsup_{k \to \infty} \psi_{n+1}(Q_k) \ge \frac{n+1}{n} \cdot \frac{(\rho - \beta) - (\beta + \rho)T}{(\rho - \beta)(1 + \frac{1}{\beta + \rho})} - 1$$

Inserting in (26) gives (9).

We turn to the upper bounds. Here we have to find upper bounds for $\overline{\psi}_{n+1}$ and $\underline{\psi}_{n+1}$, respectively. For the uniform exponent we choose $q = u_k$ as the points where the graphs of $L_{\underline{x}_{k-n}}(q)$ and $L_{\underline{x}_k}(q)$ meet, that is the solution of

$$L_{\underline{x}_{k-n}}(q) = L_{\underline{x}_k}(q).$$

Since at this position clearly $L_{\underline{x}_{k-n}}(q)$ increases whereas $L_{\underline{x}_k}(q)$ decreases, from (22) we obtain

(37)
$$u_k = \frac{n}{n+1} (\log x_k - \log Y_{k-n}).$$

Again since $\underline{x}_{k-n}, \underline{x}_{k-n+1}, \ldots, \underline{x}_k$ are linearly independent, by (23) we have to bound each of the corresponding functions $L_{\underline{x}_j}, k-n \leq j \leq k$, at position u_k . It is easy to see that $L_{\underline{x}_k}(u_k) = L_{\underline{x}_{k-n}}(u_k)$ is the maximum among the values, since the rising part of the graph of $L_{\underline{x}_{k-n}}(q)$ will intersect the falling part of each of $L_{\underline{x}_{k-n+1}}(q), \ldots, L_{\underline{x}_{k-1}}(q)$ before (i.e. at smaller q values) it eventually meets $L_{\underline{x}_k}(q)$ at $q = u_k$. Moreover by (22) the function value is given as $L_{\underline{x}_k}(u_k) = \log x_k - u_k$. Hence we estimate

$$\underline{\psi}_{n+1} \le \liminf_{k \to \infty} \frac{L_{\underline{x}_k}(u_k)}{u_k} \le \liminf_{k \to \infty} \frac{\log x_k - u_k}{u_k} = \liminf_{k \to \infty} \frac{\log x_k}{u_k} - 1.$$

Using the representation (37) of u_k yields

$$\underline{\psi}_{n+1} \le \liminf_{k \to \infty} \frac{n+1}{n} \cdot \frac{\log x_k}{\log x_k - \log Y_{k-n}} - 1$$

We check that the expression increases in Y_{k-n} . Hence we can use (28) to estimate

$$\underline{\psi}_{n+1} \le \liminf_{k \to \infty} \frac{n+1}{n} \cdot \frac{\log x_k}{\log x_k - (-\beta + \rho) \log x_{k-n}} - 1$$

Again since $\beta > \rho$ we see the right hand side decays in x_{k-n} , application of (27) with $\ell = n$ and yields

(38)
$$\underline{\psi}_{n+1} \le \frac{n+1}{n} \cdot \frac{1}{1 + (\beta - \rho)(\frac{\alpha}{\beta} - \phi)^n} - 1.$$

Plugging this into (25) we get (10).

Finally we show (11). For this we estimate $\overline{\psi}_{n+1}$ from above to apply (26). Recall we defined u_k as the value where $L_{\underline{x}_{k-n}}$ meets $L_{\underline{x}_k}$. Consider the interval $[u_k, u_{k+1})$. We have seen above that the last successive minimum function at u_k is at most

$$L_{n+1}(u_k) \le L_{\underline{x}_k}(u_k) = L_{\underline{x}_{k-n}}(u_k) = \frac{u_k}{n} + \log Y_{k-n}.$$

Now let p_k be the point where $L_{\underline{x}_{k-n}}$ meets $L_{\underline{x}_{k+1}}$. Clearly $p_k > u_k$. Similarly as in (37) above we get

(39)
$$p_k = \frac{n}{n+1} (\log x_{k+1} - \log Y_{k-n}).$$

Comparison with (37) for index k + 1 gives $p_k < u_{k+1}$ since $(Y_j)_{j\geq 1}$ decreases. We split $[u_k, u_{k+1})$ into $[u_k, p_k)$ and $[p_k, u_{k+1})$. It is readily verified that for $q \in [u_k, p_k)$ we have

$$L_{\underline{x}_{k-n}}(q) > L_{\underline{x}_{k-n+1}}(q) > \dots > L_{\underline{x}_k}(q)$$

and since these vectors are linearly independent, we have

(40)
$$L_{n+1}(q) \le L_{\underline{x}_{k-n}}(q), \qquad q \in [q_k, p_k)$$

In the other partial interval $q \in [p_k, u_{k+1})$ we similarly see that

$$L_{\underline{x}_{k+1}}(q) > L_{\underline{x}_{k-n+1}}(q) > L_{\underline{x}_{k-n+2}}(q) > \dots > L_{\underline{x}_{k}}(q)$$

and again by the linear independence of the n + 1 functions $L_{\underline{x}_{k-n+1}}, L_{\underline{x}_{k-n+2}}, \ldots, L_{\underline{x}_{k+1}}$ by Theorem 1.1 and the definition of u_{k+1} we conclude

$$L_{n+1}(q) \le L_{\underline{x}_{k+1}}(q), \qquad q \in [p_k, u_{k+1}).$$

Since $L_{\underline{x}_{k+1}}$ which decays with slope -1 in the latter interval $[p_k, u_{k+1})$, the corresponding values $L_{\underline{x}_{k+1}}(q)/q$ decrease in this interval. Thus we only need to take into account $[u_k, p_k]$ when looking for upper bounds for $\overline{\psi}_{n+1}$. Moreover, by (40) and since $L_{\underline{x}_{k-n}}(q)$ increases in this interval with slope 1/n, the quantity $L_{\underline{x}_{k-n}}(q)/q$ increases in $[u_k, p_k]$, it suffices to consider the right end point $q = p_k$. From (39) and since $[u_m, \infty)$ is the disjoint union of the intervals $[u_k, u_{k+1})$ over $k \geq m$, observing the similarity between (37) and (39), a very similar argument as for (38) above yields

$$\overline{\psi}_{n+1} \le \limsup_{k \to \infty} \frac{L_{\underline{x}_{k+1}}(p_k)}{p_k} \le \frac{n+1}{n} \cdot \frac{1}{1 + (\beta - \rho)(\frac{\alpha}{\beta} - \phi)^{n+1}} - 1.$$

Again using (26) we obtain the bound (11).

We deduce the corollary.

Proof of Corollary 1.3. We identify $\alpha = \widehat{\omega}(\underline{\xi})$ and $\beta = \omega(\underline{\xi})$. Assume equality in (4). Then $\epsilon = \rho = \phi = 0$ and the bounds of Theorem 1.2 simplify reasonably, in fact (10) and (11) directly become β^{n-1}/α^n and β^n/α^{n+1} , respectively. To identify the lower bounds (8), (9) with these respective values, some rearrangements and explicit use of equality in (4) is required, we do not carry it out. We describe how to infer the second claim by a continuity argument. Let $\varepsilon > 0$. First fix $\alpha \in [1/n, 1)$. Let β_0 be the solution to equality in (4). Then by continuity of the bounds in Theorem 1.2 in α, β , there is some $\tilde{\delta} > 0$ that depends on n, ε, α such that $\beta \in (\beta_0 - \tilde{\delta}, \beta_0 + \tilde{\delta})$ implies that the bound expressions differ from the respective values $\omega(\underline{\xi})^{n-1}/\widehat{\omega}(\underline{\xi})^n$ and $\omega(\underline{\xi})^n/\widehat{\omega}(\underline{\xi})^{n+1}$ of the case $\beta = \beta_0$ by less than ε . By implicit function theorem, for given ϵ we have that the solution β to (7) depends continuously on α , and for $\epsilon = 0$ it becomes β_0 . Hence the same claim holds if $\epsilon = \epsilon_{\alpha,\beta}$ in (6) is smaller than some modified $\delta > 0$. Since we restrict α to the compact interval [1, c] and (a lower bound for) δ depends continuously on α , we may choose $\delta > 0$ independently of α .

4. Proof of Theorems 2.2, 2.3

The proof of Theorem 2.2 is based on a case distinction $\lambda_n(\xi) > \frac{2}{n}$ and $\lambda_n(\xi) \le \frac{2}{n}$. The first case is dealt with by the following result from [27, Section 4].

Theorem 4.1 (Schleischitz). Let $n \ge 2$ be an even integer. Assume ξ is transcendental real and satisfies

0

$\lambda_n(\xi)$	$>\frac{2}{n}$.
$\widehat{\lambda}_n(\xi) \leq$	$\frac{2}{n+2}$

Then

It can be verified that the bound 2/(n+2) is smaller than σ_n in Theorem 2.2 (thus clearly also smaller than τ_n in Theorem 2.1).

In the latter case $\lambda_n(\xi) \leq \frac{2}{n}$ we use Theorem 1.2. We point out that the slightly weaker bounds in our Theorem 2.1 obtained in [27] followed from combining Theorem 4.1 with (4), corresponding to $\epsilon = 0$ in (6). The key observation for the improvement is that when $\beta = 2/n$ and $\epsilon = 0$, then the bound for the exponent $\widehat{w}_n(\xi)$ we obtain from (8) is larger than settled upper bounds for this exponent rephrased in Theorem 4.2 below. Thus by continuity we expect that if β is not too much smaller than 2/n and ϵ sufficiently small, we still get a contradiction to (8). Hereby we want to recall the identifications $\alpha = \widehat{\omega}(\xi) = \widehat{\lambda}_n(\xi), \ \beta = \omega(\xi) = \lambda_n(\xi)$ and $\widehat{w}_n(\xi) = \widehat{\omega}^*(\xi)$.

We turn to the upper bound for $\widehat{w}_n(\xi)$ indicated above. Indeed, in contrast to general points in \mathbb{R}^n where $\widehat{\omega}^*(\underline{\xi}) = \infty$ may occur (if $n \ge 2$), for points on the Veronese curve the uniform exponent is bounded in terms of n. The following currently best known bound is a consequence of Bugeaud and Schleischitz [5] when incorporating the linear form result (dual to (4) in some sense) from [16], already observed in [26].

Theorem 4.2 (Bugeaud, Schleischitz). For any $n \ge 1$ and transcendental real ξ we have $\widehat{w}_n(\xi) \le \mu_n$ where μ_n is as defined in Theorem 2.2.

For sake of completeness we mention that the paper [5] in turn improved slightly on a bound by Davenport and Schmidt [7]. All aforementioned proofs are based on some variation (see [7, Lemma 8]) of the Liouville inequality that provides effective lower bounds for the distance between two distinct algebraic numbers in terms of their degrees and heights. Recall (12) for the following proof.

Proof of Theorem 2.2. We may assume $\lambda_n(\xi) \leq 2/n$ by Theorem 4.1. Observe that by (4) this implies an upper bound for $\hat{\lambda}_n(\xi)$ that is just slightly larger than σ_n . Denote this inferred bound by Ψ_n and write $I_n := (\sigma_n, \Psi_n)$.

Now assume on the contrary that $\widehat{\lambda}_n(\xi) > \sigma_n$ for some ξ . Then $\widehat{\lambda}_n(\xi) \in I_n$, and again from (4) we obtain a lower bound for $\lambda_n(\xi)$ just slightly smaller than 2/n. Denote the bound by Φ_n and the resulting range for $\lambda_n(\xi)$ by $J_n := (\Phi_n, 2/n]$. For given parameters α, β write

$$W_{\alpha,\beta} := \frac{(\beta - \frac{4\epsilon\beta^2}{\alpha^2})S}{(\frac{\alpha}{\beta} - \frac{4\beta^{n-1}\epsilon}{\alpha^n})^{-n} + (\beta - \frac{4\epsilon\beta^2}{\alpha^2})(1-S)}$$

where $\epsilon = \epsilon_{\alpha,\beta}$ is defined in (6), and $S = S_{\alpha,\beta}$ in (8). Then in particular $W_{\alpha} := W_{\alpha,2/n}$ denotes the left hand side in (16). By construction $\epsilon_{\alpha} = \epsilon_{\alpha,2/n}$ and $S_{\alpha} = S_{\alpha,2/n}$ and σ_n is the solution for α to equality $W_{\alpha} = \mu_n$, thus

(41)
$$W_{\sigma_n,2/n} = \mu_n$$

A brief calculation verifies that $\epsilon = \epsilon_{\alpha,\beta}$ from (6) satisfies (7) when

(42)
$$\alpha \in I_n, \qquad \beta \in J_n \cup (\frac{2}{n}, \frac{2}{n} + \varepsilon) = (\Phi_n, \frac{2}{n} + \varepsilon) =: K_n$$

for some small $\varepsilon = \varepsilon(n) > 0$ (independent of α).

Next observe that by the strict inequality $\widehat{\lambda}_n(\xi) > \sigma_n$, we have that the hypothesis (5) of Theorem 1.1 holds for every pair (α, β) with $\alpha \in (\sigma_n, \widehat{\lambda}_n(\xi)) \subseteq I_n$ and $\beta > 2/n$, and suitable a, b. Thus, hypothesis (5) holds in particular for $\alpha \in I_n$ and $\beta \in (2/n, 2/n + \varepsilon) \subseteq K_n$. Hence we can apply Theorem 1.2 for any

(43)
$$\alpha \in (\sigma_n, \widehat{\lambda}_n(\xi)) \subseteq I_n, \qquad \beta \in (\frac{2}{n}, \frac{2}{n} + \varepsilon) \subseteq K_n,$$

and (8) yields for any pair α, β as in (43) the inequality

(44)
$$\widehat{w}_n(\xi) = \widehat{\omega}^*(\underline{\xi}) \ge W_{\alpha,\beta}.$$

A short calculation further shows that when $\beta \in K_n$ is fixed, the expression $W_{\alpha,\beta}$ increases as α increases in I_n . Thus

$$W_{\alpha,\beta} > W_{\sigma_n,\beta}$$

with strict inequality because $\alpha > \sigma_n$ strictly. By continuity of $W_{\alpha,\beta}$ in the second argument, for any fixed $\alpha > \sigma_n$ we still have

(45)
$$W_{\alpha,\beta} > W_{\sigma_n,2/n}$$

if β is sufficiently close to 2/n (alternatively one can start with $\mu_n + \varepsilon$ for arbitrarily small $\varepsilon > 0$ in the right hand side of (16), and use continuity to derive the contradiction below). Thus for any pair α, β as in (43), combining (41), (44), (45) upon making ε smaller if necessary we conclude

$$\widehat{w}_n(\xi) = \widehat{\omega}^*(\underline{\xi}) \ge W_{\alpha,\beta} > W_{\sigma_n,2/n} = \mu_n$$

This contradicts Theorem 4.2. Thus we cannot have $\widehat{\lambda}_n(\xi) > \sigma_n$.

Finally we prove Theorem 2.3 with a similar method.

Proof of Theorem 2.3. We proceed as in the proof of Theorem 2.2. Let us first assume $\epsilon = 0$ for some α, β , i.e. there is equality in (4) and we are in the situation of the regular graph. Then $\alpha = \hat{\lambda}_n(\xi)$ and $\beta = \lambda_n(\xi)$ and by Corollary 1.3, we have identity

(46)
$$\widehat{w}_n(\xi) = \widehat{\omega}^*(\underline{\xi}) = \frac{\omega(\underline{\xi})^{n-1}}{\widehat{\omega}(\underline{\xi})^n} = \frac{\lambda_n(\xi)^{n-1}}{\widehat{\lambda}_n(\xi)^n}.$$

It is easily checked that upon equality in (4) the expressions are increasing as functions in $\alpha = \hat{\lambda}_n(\xi)$. By Theorem 4.2, the exponent $\hat{\lambda}_n(\xi)$ is thus bounded by the solution to

(47)
$$\mu_n = \frac{\lambda_n(\xi)^{n-1}}{\widehat{\lambda}_n(\xi)^n},$$

with the exponents $\lambda_n(\xi)$ and $\hat{\lambda}_n(\xi)$ linked by an identity in (4). For $n \in \{4, 6, 8\}$ we derive the stated numerical bounds (17) with some computation. For large n, we have $\mu_n = 2n - 2 < 2n$ and with some analysis of the regular graph the claimed asymptotics (18) can be derived. We give some more details. We use identity [25, (31)] which,

upon identifying $\widehat{\omega}_{n+1}(\underline{\xi})^{-1} = \omega(\underline{\xi})$ by [9, Corollary 8.5] (already used in the proof of Proposition 3.1), can be written

(48)
$$(1+\omega^*(\underline{\xi})) \cdot \left(1+\frac{1}{\omega^*(\underline{\xi})}\right)^n = \left(1+\frac{1}{\omega(\underline{\xi})}\right) \cdot (1+\omega(\underline{\xi}))^n.$$

Moreover by (46) and (47) we have $\hat{\omega}^*(\underline{\xi})/n = \mu_n/n = 2 - o(1)$ and then also $\omega^*(\underline{\xi})/n = 2 + o(1)$ as can be derived from identity [25, (33)], we see that the left hand side in (48) is of order $(2\sqrt{e} + o(1))n$ as $n \to \infty$. Thus so is the right hand side and we readily conclude $n\omega(\underline{\xi}) = \Theta + o(1)$. Finally the smaller quantity $n\hat{\omega}(\underline{\xi}) = n\hat{\lambda}_n(\xi)$ will be asymptotically of the same order as $n \to \infty$ (see (30) in [25]), thus (18) follows.

Finally (19) follows by a similar continuity argument as in Corollary 1.3. First assume $\alpha \in [1/n, 1)$ is fixed. By Corollary 1.3 and its proof, the expression $\widehat{w}_n(\xi) = \widehat{\omega}^*(\underline{\xi})$ depends continuously on ϵ if $\beta = \beta(\epsilon)$ is such that there is identity in (7). Moreover for $\epsilon = 0$ and $\alpha = \widehat{\lambda}_n(\xi)$ larger than claimed, we get a contradiction $\widehat{w}_n(\xi) > \mu_n$ as we have proved above. Thus for $\epsilon \in [0, \delta_n]$ with $\delta_n > 0$ small enough in dependence of n, α, ε , in case of larger $\alpha = \widehat{\lambda}_n(\xi)$ we still obtain the same contradiction $\widehat{w}_n(\xi) > \mu_n$. Finally, again as we can restrict $\alpha = \widehat{\lambda}_n(\xi)$ to a compact interval like [1/n, 1/2], we can choose δ_n uniformly in α , depending only on n and ε .

The author thanks the referee for the careful reading and for providing references, as well as Nikolay Moshchevitin for pointing out an inaccuracy in the original proof of Theorem 1.2.

References

- D. Badziahin, J. Schleischitz. An improved bound in Wirsing's problem. Trans. Amer. Math. Soc. 374 (2021), no. 3, 1847–1861.
- [2] Y. Bugeaud. Approximation by algebraic numbers. Cambridge Tracts in Mathematics, 160. Cambridge University Press, Cambridge, 2004.
- [3] Y. Bugeaud, M. Laurent. Exponents of Diophantine Approximation and Sturmian Continued Fractions. Ann. Inst. Fourier (Grenoble) 55 (2005), 773–804.
- [4] Y. Bugeaud, M. Laurent. On transfer inequalities in Diophantine approximation. II. Math. Z. 265 (2010), no. 2, 249–262.
- [5] Y. Bugeaud, J. Schleischitz. On uniform approximation to real numbers. Acta Arith. 175 (2016), 255–268.
- [6] Y. Bugeaud, O. Teulié. Approximation d'un nombre réel par des nombres algébriques de degré donné. (French) [Approximation of a real number by algebraic numbers of a given degree]. Acta Arith. 93 (2000), no. 1, 77–86.
- [7] H. Davenport, W. M. Schmidt. Approximation to real numbers by algebraic integers. Acta Arith. 15 (1969), 393–416.
- [8] O. German, On Diophantine exponents and Khintchine's transference principle, Mosc. J. Comb. Number Theory 2, (2012), 22–51.
- [9] O. German. Intermediate Diophantine exponents and parametric geometry of numbers. Acta Arith. 154 (2012), no. 1, 79–101.
- [10] O. German, N.G. Moshchevitin. A simple proof of Schmidt-Summerer's inequality. Monatsh. Math. 170 (2013), no. 3-4, 361–370.

JOHANNES SCHLEISCHITZ

- [11] V. Jarník. Zum Khintchineschen Übertragungssatz. Acad. Sci. URSS., Fil. Géorgienne, Trav. Inst. Math. Tbilissi 3 (1938), 193–216.
- [12] Y. A. Khintchine, Uber eine Klasse linearer Diophantischer Approximationen. Rendiconti Palermo 50 (1926), 170–195.
- [13] M. Laurent. Simultaneous rational approximation to the successive powers of a real number. Indag. Math. (N.S.) 14 (2003), no. 1, 45–53.
- [14] M. Laurent. Exponents of Diophantine approximation in dimension two. Canad. J. Math. 61 (2009), no. 1, 165–189.
- [15] K. Mahler. Zur Approximation der Exponentialfunktionen und des Logarithmus I,II. J. reine angew. Math. 166 (1932), 118–150.
- [16] A. Marnat, N. Moshchevitin. An optimal bound for the ratio between ordinary and uniform exponents of Diophantine approximation. *Mathematika* 66 (2020), no. 3, 818–854.
- [17] N. A. V. Nguyen, A. Poels, D. Roy. A transference principle for simultaneous Diophantine approximation. J. Théor. Nombres Bordeaux 32 (2020), no. 2, 387–402.
- [18] A. Poels. A class of maximally singular sets for rational approximation. Int. J. Number Theory 16 (2020), no. 9, 2005–2012.
- [19] M. Rivard-Cooke. Parametric Geometry of Numbers. PhD thesis, University of Ottawa, 2019; https://ruor.uottawa.ca/handle/10393/38871.
- [20] D. Roy. Approximation to real numbers by cubic algebraic integers I. Proc. London Math. Soc. (3) 88 (2004), no. 1, 42–62.
- [21] D. Roy. On simultaneous rational approximations to a real number, its square, and its cube. Acta Arith. 133 (2008), no. 2, 185–197.
- [22] D. Roy. Construction of points realizing the regular systems of Wolfgang Schmidt and Leonhard Summerer. J. Théor. Nombres Bordeaux 27 (2015), no. 2, 591–603.
- [23] D. Roy. On Schmidt and Summerer parametric geometry of numbers. Ann. of Math. (2) 182 (2015), no. 2, 739–786.
- [24] D. Roy. On the topology of Diophantine approximation spectra. Compos. Math. 153 (2017), no. 7, 1512–1546.
- [25] J. Schleischitz. Some notes on the regular graph defined by Schmidt and Summerer and uniform approximation. JP J. Algebra, Number Theory Appl. 39 (2017), no. 2, 115–150.
- [26] J. Schleischitz. Uniform approximation and best approximation polynomials. Acta Arith. 185 (2018), no. 3, 249–274.
- [27] J. Schleischitz. An equivalence principle between polynomial and simultaneous Diophantine approximation. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 21 (2020), 1063–1085.
- [28] W.M. Schmidt, L. Summerer. Parametric geometry of numbers and applications. Acta Arith. 140 (2009), no. 1, 67–91.
- [29] W.M. Schmidt, L. Summerer. Diophantine approximation and parametric geometry of numbers. Monatsh. Math. 169 (2013), 51–104.
- [30] W.M. Schmidt, L. Summerer. Simultaneous approximation to three numbers. Mosc. J. Comb. Number Theory 3 (2013), no. 1, 84–107.
- [31] W.M. Schmidt, L. Summerer. Simultaneous approximation to two reals: bounds for the second successive minimum. *Mathematika* 63 (2017), no. 3, 1136–1151.