

# Traversable Wormholes in $f(R, T)$ Gravity

Ambuj Kumar Mishra<sup>1</sup>, Umesh Kumar Sharma<sup>2</sup>, Vipin Chandra Dubey<sup>3</sup>, Anirudh Pradhan<sup>4</sup>

<sup>1,2,3,4</sup>Department of Mathematics, Institute of Applied Sciences and Humanities, GLA University, Mathura-281 406, Uttar Pradesh, India

<sup>1</sup>E-mail: ambuj\_math@rediffmail.com

<sup>2</sup>E-mail: sharma.umesh@gla.ac.in

<sup>3</sup>E-mail: vipin.dubey@gla.ac.in

<sup>4</sup>E-mail: pradhan.anirudh@gmail.com

## Abstract

In the present article, models of traversable wormholes within the  $f(R, T)$  modified gravity theory are investigated. We have presented some wormhole models, developed from various hypothesis for the substance of their matter, i.e. various relationships with their components of pressure (lateral and radial). The solutions found for the shape functions of the wormholes produced complies with the required metric conditions. The suitability of solution is examined by exploring null, strong and dominant energy conditions. It is surmised that the normal matter in the throat may pursue the energy conditions yet the gravitational field exuding from the adjusted gravity hypothesis support the appearance of the non-standard geometries of wormholes.

Keywords: Traversable wormholes,  $f(R, T)$  gravity, Energy conditions, Shape functions

PACS number: 04.50.kd, 04.20.Gz, 04.20.q

## 1 Introduction

The exact solutions for wormholes (WH) were obtained connecting two distinct regions of asymptotically flat space time in general relativity [1, 2], also as two distinct anti-de Sitter or asymptotically de-Sitter regions [3]. Although, wormholes still have not been observed [4] but attempts to achieve has been proposed [5–13]. Theoretically wormholes were predicted a long time ago. Wormholes solutions were first proposed by Einstein and Rosen [14] with event horizon. Many decades later, Morris and Thorne have suggested that wormholes can be traversable, which violates the energy conditions, if filled with exotic matter [15].

In wormholes Physics, the presence of throat is the fundamental feature which satisfy flaring-out conditions [16]. Other theories have been also suggested wormhole solutions [17–26] and [27] for reviews. NEC for matter is also violated in these works. In fact, the wormhole throat can be threaded by the normal matter in the framework of modified

gravity theories and these nonstandard wormhole geometries are supported by the higher order curvature term [16], which can be termed as gravitational fluid. Theoretically, wormholes throats can be formulated in absence of exotic matter [28]. Wormholes were constructed by non-minimal coupling in [29,30]. In modified gravity alone, the fundamental fields sustained the wormhole throats using generic modified gravities [31]. Such solutions are also obtained in [32–38]. Rosa et al. [16] found wormholes solutions in which the null energy conditions (NEC) was obeyed by the matter not only at throat but everywhere in a hybrid generalized metric-Palatini gravity, with action  $f(R, R)$ .

Indeed, examination of wormhole solutions in various modified theories is a noteworthy and important point in Theoretical Physics. The General Relativity (GR) could be adjusted in various perspectives. By the introduction of gravitational action of the  $f(R, T)$  gravity in Ricci scalar  $R$  and the trace of energy momentum tensor  $T$ , T. Harko and fellow researchers [39] streamlined the  $f(R)$  theories [40–45] by replacing the function  $f(R)$  with a random function  $f(R, T)$ . This  $f(R, T)$  theory tested in Astrophysics of compact objects [46–49], Thermodynamics [50, 51] and cosmology [52–63].

The work has been done on the cases of  $WH$  geometry in  $f(R, T)$  gravity, where the redshift function  $\phi$  is not dependent on either time or spatial coordinate [64,65]. The spherically symmetric and static wormholes are considered in  $f(R, T) = f(R) + \Lambda T$  gravity with different fluids [66]. The geometry (non-commutative) of wormholes with respect to Lorentzian and Gaussian distributions of string theory is proposed as well as found the numerical and exact solutions in modified  $f(R, T)$  gravity [67]. An analytical approach is used to get the wormhole solutions in  $f(R, T)$  gravity [68]. An exponential shape function is defined for wormholes and a comparison is made between  $f(R, T)$  and  $f(R)$  gravity for the rationality of energy conditions [69]. Wormholes are studied for two types of varying Chaplygin gas and found the violation of dominated and null energy conditions in  $f(R, T)$  gravity [70]. In [71], an exponential  $f(R, T)$  gravity model has been considered and obtained solutions for wormhole. The charged wormhole solutions are investigated in  $f(R, T)$  gravity satisfying the energy conditions [72]. Using Noether symmetry approach, wormhole solutions were investigated for non dust and dust distributions in  $f(R, T)$  gravity [73]. Considering the different types of energy density, traversable wormhole solutions investigated in  $f(R, T)$  gravity [74]. Wormhole solutions have been studied by various researchers in modified gravity theories in various set-ups [75–79].

The fact that wormholes matter content is signified by an anisotropic fluid and  $T$ - dependence of the  $f(R, T)$  theory, which may be related to imperfect fluid forms the strong motivation behind working with the wormholes in  $f(R, T)$  gravity [68,80]. Recently, traversable wormhole are investigated for two particular types of shape functions and obtained wormhole structures filled with phantom fluid [81]. Further, traversable wormholes investigated the in  $f(R, T)$  gravity by taking a new form of  $f(R, T) = R + 2\ln T$  function to minimize the appearance of exotic matter near the throat of the wormhole [82].

Hence, in this paper, we have investigated the traversable wormholes by assuming functional form of  $f(R, T) = R + \Lambda T$  of  $f(R, T)$  gravity, originally proposed by the authors [39] considering the shape function given in [80], which has not been studied already.

The paper is organized as follows: in Section II, a brief review is presented for the  $f(R, T)$  theory. In Section III, the wormhole metric and its conditions are described. In Section IV, we presented the field equation solution for the wormhole metric in the  $f(R, T)$  gravity. In Section V, we presented models of the wormhole. Our results are discussed in Section VI.

## 2 The basic formalism of $f(R, T)$ gravity

The theory of  $f(R, T)$  gravity is given by the following action [39]

$$S = \frac{1}{16\pi} \int d^4x f(R, T) \sqrt{-g} + \int \mathcal{L}_m d^4x \sqrt{-g}, \quad (1)$$

where  $R$  is Ricci scalar and  $T$  is trace of the energy momentum tensor (EMT) in an arbitrary function  $f(R, T)$ ,  $g$  is the metric determinant and matter Lagrangian density is  $\mathcal{L}_m$ . These are connected to the EMT [83].

$$T_{ij} = -\frac{2}{\sqrt{-g}} \left[ \frac{\partial(\sqrt{-g}\mathcal{L}_m)}{\partial g^{ij}} - \frac{\partial}{\partial x^k} \frac{\partial(\sqrt{-g}\mathcal{L}_m)}{\partial(\frac{\partial g^{ij}}{\partial x^k})} \right]. \quad (2)$$

Moreover, units is taken such that  $c = 1 = G$ .

Following as [39,84], we consider  $\mathcal{L}_m$  relies only on  $g_{ij}$  (metric component) and not depends on its differential coefficients, such that we can get

$$T_{ij} = -\frac{\partial \mathcal{L}_m}{\partial g^{ij}} + g_{ij} \mathcal{L}_m. \quad (3)$$

In the Eq. (1), the action  $S$  is varying with  $g_{ij}$  which provides the  $f(R, T)$  field equations [39] as

$$\begin{aligned} f_R(R, T) \left( R_{ij} - \frac{1}{3} R g_{ij} + \frac{1}{6} f(R, T) g_{ij} \right) \\ = 8\pi G \left( T_{ij} - \frac{1}{2} T g_{ij} \right) - f_T(R, T) \left( T_{ij} - \frac{1}{3} T g_{ij} \right) \\ - f_T(R, T) \left( \theta_{ij} - \frac{1}{3} \theta g_{ij} \right) + \nabla_i \nabla_j f_R(R, T) \end{aligned} \quad (4)$$

with  $f_R(R, T) = \frac{\partial f(R, T)}{\partial R}$ ,  $f_T(R, T) = \frac{\partial f(R, T)}{\partial T}$  and

$$\theta_{ij} = g^{ij} \frac{\partial T_{ij}}{\partial g^{ij}}. \quad (5)$$

Such theories of gravity  $f(R, T)$  (as well as  $f(R, \mathcal{L}_m)$ ) can explain in their formalism a non-minimal matter-geometry coupling. It suggests that research particles in the field (gravitational) are not going to obey geodesic lines in such theories. The geometry and matter coupling prompts an additional force one behaves perpendicular to the four-velocity on the particles.

Interestingly, an additional force relies on the Lagrangian structure of the matter [85]. It has been found in [86], by taking the total pressure of  $\mathcal{L}_m = p$  with  $p$ , the additional force disappear. However, more normal structures of  $\mathcal{L}_m$  such as  $\mathcal{L}_m = -\rho$  with  $\rho$  represents

the density of energy are more common apparently and do not give the vanishing of the additional force.

Assuming the Lagrangian matter in this paper as  $\mathcal{L}_m = -\rho$ . It is therefore possible to write an equation (5) as

$$\theta_{ij} = -2T_{ij} - \rho g_{ij}. \quad (6)$$

Let us suppose the function  $f(R, T) = 2f(T) + R$ , where  $f(T)$  is a random function of  $T$ . The field equations (4) and (5) of  $f(R, T)$  gravity takes the form

$$R_{ij} - \frac{1}{2}Rg_{ij} = 8\pi T_{ij} + 2f^t(T)T_{ij} + [2\rho f^t(T) + f(T)]g_{ij}, \quad (7)$$

where  $R_{ij}$  is Ricci tensor and  $f^t(T) = \frac{df(T)}{dT}$ . By taking  $f(T) = \lambda T$ , with a constant  $\lambda$ , we can rewrite the above as

$$G_{ij} = (8\pi + 2\lambda)T_{ij} + \lambda(2\rho + T)g_{ij}, \quad (8)$$

where  $G_{ij}$  is usual Einstein tensor. Such a  $f(R, T)$  functional type assumption was originally proposed in [39].

### 3 Wormhole Metric and Its Conditions

In Schwarzschild coordinate  $(t, r, \theta, \phi)$ , the spherical symmetric wormhole metric is [87, 88]

$$ds^2 = -U(r)dt^2 + \frac{dr^2}{V} + r^2d\Omega^2, \quad (9)$$

where  $\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$  and  $V = 1 - \frac{b(r)}{r}$ . The  $U(r)$  that denotes redshift function and the  $b(r)$  that denotes shape function given in Eq. (9) will follow the conditions mentioned below [87, 88]

1. The coordinate  $r$  named as radial co-ordinate, ranges  $r_0 \leq r \leq \infty$ , where  $r_0$  refers to the radius of the throat.
2. The shape function  $(b(r))$  fulfills the condition at the throat

$$b(r_0) = r_0, \quad (10)$$

and for the out of throat i.e. for  $r > r_0$

$$0 < 1 - \frac{b(r)}{r}. \quad (11)$$

3. The flaring out condition must be satisfied by shape function  $b(r)$  at the throat, i.e.

$$b'(r_0) < 1, \quad (12)$$

where superscript  $(')$  represent derivative with respect to  $r$

4. The limit required for asymptotically flatness of the space-time geometry

$$\frac{b(r)}{r} \rightarrow 0 \quad \text{as} \quad |r| \rightarrow \infty. \quad (13)$$




Figure 1: Nature of  $b(r)$ , throat condition  $b(r) < r$ , flaring out condition  $b'(r) < 1$  and asymptotically flatness  $\lim_{r \rightarrow \infty} \frac{b(r)}{r} = 0$  for  $r_0 = 1$ .

5. At the throat  $r_0$ , redshift function  $U(r)$  must be non-vanishing and finite.

To achieve the anti-de Sitter and de Sitter asymptotic behavior, we can take  $U(r) = \text{constant}$ . Since the re-scaled time coordinate can absorb a constant redshift, we take  $U(r) = 1$ , such as [89, 90].

## 4 Solutions of Field Equations for Wormholes in $f(R, T)$ Gravity.

Considering that the description of matter is described by an anisotropic fluid of the form

$$T_j^i = \text{diag} \{-\rho, p_r, p_l, p_l\}. \quad (14)$$


Where  $\rho = \rho(r), p_r = p_r(r), p_l = p_l(r)$  are energy density, radial pressure and transverse (measured orthogonal to radial directions) pressure, respectively and the trace  $T$  of the Eq. (14) can be obtained as  $T = -\rho + p_r + 2p_l$ .

The field equation components from Eq. (8) for the metric given by Eq. (9) with Eq. (14) are obtained as :


$$\frac{b'}{r^2} = (8\pi + \lambda)\rho - \lambda(p_r + 2p_l), \quad (15)$$

$$-\frac{b}{r^3} = \lambda\rho + (8\pi + 3\lambda)p_r + 2\lambda p_l, \quad (16)$$

$$\frac{b - b'r}{2r^3} = \lambda\rho + \lambda p_r + (8\pi + 4\lambda)p_l, \quad (17)$$




(a)




(b)

Figure 2: (a) NEC,  $\rho + p_r$ , (b) NEC,  $\rho + p_t$ .



(a)



(b)

Figure 3: (a) SEC,  $\rho + p_r + 2p_t$ , (b) DEC,  $\rho - p_r$ .




Figure 4: (a) DEC,  $\rho - p_l$ , (b) DEC,  $p_l - p_r$ .

the field equations given by Eqs. (15), (16) and(17), yields the following solutions

$$\rho = \frac{b'}{r^2(8\pi + 2\lambda)}, \quad (18)$$

$$p_r = -\frac{b}{r^3(8\pi + 2\lambda)}, \quad (19)$$

$$p_l = \frac{b - b'r}{2r^3(8\pi + 2\lambda)}. \quad (20)$$

## 5 Models of Wormholes

We shall investigate models of wormholes for two different shape functions in this section.

### 5.1 Shape Function $b(r) = \frac{r_0 \log(r+1)}{\log(r_0+1)}$

For our first model we take radial pressure  $p_r$  and energy density  $\rho$  related as

$$p_r = \omega \rho, \quad (21)$$

where equation of state parameter  $\omega$  in terms of  $p_r$  is called radial state parameter. The  $f(R, T)$  gravity model is considered with the shape function [81]

$$b(r) = \frac{r_0 \log(r + 1)}{\log(r_0 + 1)}. \quad (22)$$

For this shape function, wormholes solutions are obtained. The energy density, radial pressure, tangential pressure can be obtained by using field equations (18)- (20) as

$$\rho = \frac{r_0}{(1 + r) \ln(1 + r_0) r^2 (8\pi + 2\lambda)}, \quad (23)$$

$$p_r = \frac{r_0 \ln(1+r)}{\ln(r_0+1) r^3 (8\pi+2\lambda)}, \quad (24)$$

$$p_l = 1/2 \left( \frac{r_0 \ln(1+r)}{\ln(r_0+1)} - \frac{r_0 r}{(1+r) \ln(1+r_0)} \right) r^{-3} (8\pi+2\lambda)^{-1}, \quad (25)$$

$$\rho + p_r = \frac{1}{2} \frac{r_0 (r + r \ln(1+r) + \ln(1+r))}{(1+r) \ln(r_0+1) r^3 (4\pi+\lambda)}, \quad (26)$$

$$\rho + p_l = \frac{1}{4} \frac{r_0 (r + r \ln(1+r) + \ln(r+1))}{(1+r) \ln(r_0+1) r^3 (4\pi+\lambda)}, \quad (27)$$

$$\rho + p_r + 2p_l = \frac{r_0 \ln(1+r)}{\ln(r_0+1) r^3 (4\pi+\lambda)}, \quad (28)$$

$$\rho - |p_r| = \frac{r_0}{(1+r) \ln(r_0+1) r^2 (8\pi+2\lambda)} - \left| \frac{r_0 \ln(1+r)}{\ln(r_0+1) r^3 (8\pi+2\lambda)} \right|, \quad (29)$$

$$\rho - |p_l| = \frac{r_0}{(1+r) \ln(r_0+1) r^2 (8\pi+2\lambda)} - \frac{1}{2} \left| \left( \frac{r_0 \ln(1+r)}{\ln(r_0+1)} - \frac{r r_0}{(1+r) \ln(r_0+1)} \right) r^{-3} (8\pi+2\lambda)^{-1} \right|, \quad (30)$$

$$p_l - p_r = -\frac{1}{4} \frac{r_0 (r + \ln(1+r) r + \ln(1+r))}{(1+r) \ln(r_0+1) r^3 (4\pi+\lambda)}. \quad (31)$$

We can see directly from Fig. 1, that all the conditions i.e. fundamental needs of shape functions, such as throat condition, flaring out condition and typical asymptotically flatness condition are satisfied for WH geometry. It can be observed from Eqs. (26, 27) and Figs. 2(a) and 2(b) the validation region of null energy condition (NEC) i.e.  $\rho + p_r \geq 0$  and  $\rho + p_l \geq 0$ . This is the weakest restriction and just represents the attractive nature of gravity.

The strong energy condition (SEC) derives from the attractive nature of gravity and its shape arises directly from the analysis of a spherically symmetrical metric in the GR system. From (28) the SEC is plotted in Fig. 3(a). The SEC  $\rho + p_r + 2p_l$  is found to be positive, decreasing and tending towards zero with the increment of  $r$ .

The dominant energy condition (DEC) limited the rate of the transfer of energy to the speed of light. DEC's are obtained in Eqs. (29, 30). DEC is plotted in Figs 3(b) and 4(a). one can observe that violation of  $\rho \geq |p_r|$  for present model. We also observe from Fig. 4(b), that anisotropy parameter is negative with change in  $r$ , which signifies that the geometry is of attractive nature.

## 5.2 Shape Function $b(r) = r_0 \left(\frac{r}{r_0}\right)^\gamma$ , $0 < \gamma < 1$ .

In this model, the wormholes solutions are obtained by using the shape function  $b(r) = r_0 \left(\frac{r}{r_0}\right)^\gamma$ ,  $0 < \gamma < 1$  (see in [81]). In this case energy density  $\rho$ , radial pressure  $p_r$ , tangential pressure  $p_l$  are obtained by using field equations (18)-(20)

$$\rho = r_0 \left(\frac{r}{r_0}\right)^\gamma \gamma r^{-3} (8\pi+2\lambda)^{-1} \quad (32)$$

$$p_r = r_0 \left(\frac{r}{r_0}\right)^\gamma r^{-3} (8\pi+2\lambda)^{-1} \quad (33)$$






Figure 5: Nature of  $b(r)$ , throat condition  $b(r) < r$ , flaring out condition  $b'(r) < 1$  and asymptotically flatness  $\lim_{r \rightarrow \infty} \frac{b(r)}{r} = 0$  for  $r_0 = 1$ .

$$p_l = 1/2 \left( r_0 \left( \frac{r}{r_0} \right)^\gamma - r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma \right) r^{-3} (8\pi + 2\lambda)^{-1} \quad (34)$$

$$\rho + p_r = r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma r^{-3} (8\pi + 2\lambda)^{-1} + r_0 \left( \frac{r}{r_0} \right)^\gamma r^{-3} (8\pi + 2\lambda)^{-1} \quad (35)$$

$$\rho - p_l = r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma r^{-3} (8\pi + 2\lambda)^{-1} + 1/2 \left( r_0 \left( \frac{r}{r_0} \right)^\gamma - r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma \right) r^{-3} (8\pi + 2\lambda)^{-1} \quad (36)$$


$$\begin{aligned} \rho + p_r + 2p_l &= r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma r^{-3} (8\pi + 2\lambda)^{-1} + r_0 \left( \frac{r}{r_0} \right)^\gamma r^{-3} (8\pi + 2\lambda)^{-1} \\ &\quad + \left( r_0 \left( \frac{r}{r_0} \right)^\gamma - r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma \right) r^{-3} (8\pi + 2\lambda)^{-1} \end{aligned} \quad (37)$$

$$\rho - |p_r| = r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma r^{-3} (8\pi + 2\lambda)^{-1} - \left| r_0 \left( \frac{r}{r_0} \right)^\gamma r^{-3} (8\pi + 2\lambda)^{-1} \right| \quad (38)$$


$$\rho - |p_l| = r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma r^{-3} (8\pi + 2\lambda)^{-1} - 1/2 \left| \left( r_0 \left( \frac{r}{r_0} \right)^\gamma - r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma \right) r^{-3} (8\pi + 2\lambda)^{-1} \right| \quad (39)$$

$$p_l - p_r = 1/2 \left( r_0 \left( \frac{r}{r_0} \right)^\gamma - r_0 \left( \frac{r}{r_0} \right)^\gamma \gamma \right) r^{-3} (8\pi + 2\lambda)^{-1} - r_0 \left( \frac{r}{r_0} \right)^\gamma r^{-3} (8\pi + 2\lambda)^{-1} \quad (40)$$

We can also see from Fig. 5 that all the conditions mentioned earlier for WH geometry are satisfied for second model. Further, in Figs 6(a) and 6(b) we plot the validity region of NEC ( $\rho > o$ ). The validity of SEC can also be seen in Fig. 7(a). One can observe from Fig. 7(b) that violation of DEC ( $\rho > |p_r|$ ).  $\rho - p_l$  is positive as  $r$  increases shown in Fig. 8(a). It can be seen from Fig. 8(b), that anisotropy parameter is negative with change in  $r$ , which signifies that the geometry is of attractive nature.




(a)




(b)

Figure 6: (a) NEC,  $\rho + p_r$ , (b) NEC,  $\rho + p_l$ .




(a)




(b)

Figure 7: (a) SEC,  $\rho + p_r + 2p_l$ , (b) DEC,  $\rho - p_r$ .



(a)



(b)

Figure 8: (a) DEC,  $\rho - p_l$ , (b) DEC,  $p_l - p_r$ .

## 6 Discussion

Modified gravitational theories have received considerable attention over the last few decades as a possible alternative to GR. The paramount feature of wormholes geometries are non-vacuum solutions of Einstein's field equations. According to this theory they are filled with a matter which is different from the normal matter and is known as exotic matter. Several researchers found exotic matter to be very useful in examining whether different modified gravity models were responsible for violating energy conditions via the effective energy momentum tensors while the usual matter obey these conditions. [87,88]. In the present paper we explored different models of static wormhole solutions with two different shape functions  $b(r) = \frac{r_0 \log(r+1)}{\log(r_0+1)}$  and  $b(r) = r_0 \left(\frac{r}{r_0}\right)^\gamma$ ,  $0 < \gamma < 1$  in the theory of  $f(R, T)$  gravity. One can observe from Fig. 1 and Fig. 5 that both the shape functions satisfies throat condition, flaring out condition and typical asymptotically flatness condition for WH geometry.

Furthermore, redshift function  $U(r)$  is assumed to be constant, it indicates that a theoretical traveler's tidal gravitational force is null. In both cases we found that the geometries of the wormhole can exist even if the NEC is not violated by the usual matter i.e. it is not the exotic matter that threads the WH, but these are the extra curvature ingredients that sustained the wormhole in a modified gravity context [88]. We analyzed the nature of SEC and WEC in the context of  $f(R, T)$  gravity with two different types of shape functions to investigate this study. We emphasized the details of static wormhole geometry with variable shape function and constant redshift.

In both the models, it is observed that, for appropriate values of the parameters, we have obtain a wormhole solution with valid NEC and SEC, while DEC is also satisfied in terms of  $p_l$  and only DEC is violated in terms of  $p_r$ . The presence of these solutions is in compliance with the understanding that traversable wormholes enabled by additional fundamental gravitational fields can occur without the need for exotic matter. On the other hand, the rather

realistic construction needed to build a complete space-time wormhole solution may suggest that in this class of theories there are not many such solutions around.

## Acknowledgments

The authors are thankful to GLA University, Mathura, India for providing support and facility to carry out this research work. The authors are grateful to Ayan Banerjee, University of KwaZulu Natal, South Africa, for fruitful discussion and suggestions during his visit to GLA University.

## References

- [1] M. S. Morris and K. S. Thorne, “Wormholes in space-time and their use for interstellar travel: A tool for teaching general relativity,” *Am. J. Phys.* **56**, 395 (1988).
- [2] M. Visser, *Lorentzian wormholes: From Einstein to Hawking* (Springer-Verlag, New York, 1996).
- [3] J. P. S. Lemos, F. S. N. Lobo and S. Quinet de Oliveira, “Morris-Thorne wormholes with a cosmological constant,” *Phys. Rev. D* **68**, 064004 (2003).
- [4] P. Jordan, “The present state of Dirac’s cosmological hypothesis,” *Z. Phys.* **157**, 112 (1959).
- [5] N. Tsukamoto, “Retrolensing by a wormhole at deflection angles  $\pi$  and  $3\pi$ ,” *Phys. Rev. D* **95**, no. 8, 084021 (2017)
- [6] N. Tsukamoto, “Strong deflection limit analysis and gravitational lensing of an Ellis wormhole,” *Phys. Rev. D* **94**, no. 12, 124001 (2016).
- [7] M. Zhou, A. Cardenas-Avendano, C. Bambi, B. Kleihaus and J. Kunz, “Search for astrophysical rotating Ellis wormholes with X-ray reflection spectroscopy,” *Phys. Rev. D* **94**, no. 2, 024036 (2016).
- [8] F. Rahaman, P. K. F. Kuhfittig, S. Ray and N. Islam, “Possible existence of wormholes in the galactic halo region,” *Eur. Phys. J. C* **74**, 2750 (2014).
- [9] P. K. F. Kuhfittig, “Gravitational lensing of wormholes in the galactic halo region,” *Eur. Phys. J. C* **74**, no. 99, 2818 (2014).
- [10] C. Bambi, “Can the supermassive objects at the centers of galaxies be traversable wormholes? The first test of strong gravity for mm/sub-mm very long baseline interferometry facilities,” *Phys. Rev. D* **87**, 107501 (2013)
- [11] Z. Li and C. Bambi, “Distinguishing black holes and wormholes with orbiting hot spots,” *Phys. Rev. D* **90**, 024071 (2014).
- [12] K. K. Nandi, Y. Z. Zhang and A. V. Zakharov, “Gravitational lensing by wormholes,” *Phys. Rev. D* **74**, 024020 (2006).

- [13] T. Harko, Z. Kovacs and F. S. N. Lobo, “Thin accretion disks in stationary axisymmetric wormhole spacetimes,” *Phys. Rev. D* **79**, 064001 (2009).
- [14] A. Einstein and N. Rosen, “The particle problem in the general theory of relativity,” *Phys. Rev.* **48**, 73 (1935).
- [15] M. S. Morris and K. S. Thorne, “Wormholes in space-time and their use for interstellar travel: A tool for teaching general relativity,” *Am. J. Phys.* **56**, 395 (1988).
- [16] J. L. Rosa, J. P. S. Lemos and F. S. N. Lobo, “Wormholes in generalized hybrid metric-Palatini gravity obeying the matter null energy condition everywhere,” *Phys. Rev. D* **98**, no. 6, 064054 (2018).
- [17] A. G. Agnese and M. La Camera, “Wormholes in the Brans-Dicke theory of gravitation,” *Phys. Rev. D* **51**, 2011 (1995).
- [18] K. K. Nandi, B. Bhattacharjee, S. M. K. Alam and J. Evans, “Brans-Dicke wormholes in the Jordan and Einstein frames,” *Phys. Rev. D* **57**, 823 (1998).
- [19] K. A. Bronnikov and S. W. Kim, “Possible wormholes in a brane world,” *Phys. Rev. D* **67**, 064027 (2003).
- [20] M. La Camera, “Wormhole solutions in the Randall-Sundrum scenario,” *Phys. Lett. B* **573**, 27 (2003).
- [21] F. S. N. Lobo, Exotic solutions in general relativity: Traversable wormholes and warp drive spacetimes, in *Classical and Quantum Gravity Research*, 1-78, (2008), Nova Sci. Pub. ISBN 978-1-60456-366-5; arXiv:0710.4474 [gr-qc].
- [22] R. Garattini and F. S. N. Lobo, “Self sustained phantom wormholes in semi-classical gravity,” *Class. Quant. Grav.* **24**, 2401 (2007).
- [23] F. S. N. Lobo, “General class of wormhole geometries in conformal Weyl gravity,” *Class. Quant. Grav.* **25**, 175006 (2008).
- [24] R. Garattini and F. S. N. Lobo, “Self-sustained traversable wormholes in noncommutative geometry,” *Phys. Lett. B* **671**, 146 (2009).
- [25] F. S. N. Lobo and M. A. Oliveira, “General class of vacuum Brans-Dicke wormholes,” *Phys. Rev. D* **81**, 067501 (2010).
- [26] R. Garattini and F. S. N. Lobo, “Self-sustained wormholes in modified dispersion relations,” *Phys. Rev. D* **85**, 024043 (2012).
- [27] F. S. N. Lobo (editor), “Wormholes, Warp Drives and Energy Conditions, Fundam”. *Theor. Phys.* 189, (Springer International Publishing, 2017).
- [28] F. S. N. Lobo and M. A. Oliveira, “Wormhole geometries in  $f(R)$  modified theories of gravity,” *Phys. Rev. D* **80**, 104012 (2009).
- [29] N. M. Garcia and F. S. N. Lobo, “Wormhole geometries supported by a nonminimal curvature-matter coupling,” *Phys. Rev. D* **82**, 104018 (2010).

- [30] N. Montelongo Garcia and F. S. N. Lobo, “Nonminimal curvature-matter coupled wormholes with matter satisfying the null energy condition,” *Class. Quant. Grav.* **28**, 085018 (2011).
- [31] T. Harko, F. S. N. Lobo, M. K. Mak and S. V. Sushkov, “Modified-gravity wormholes without exotic matter,” *Phys. Rev. D* **87**, no. 6, 067504 (2013).
- [32] B. Bhawal and S. Kar, “Lorentzian wormholes in Einstein-Gauss-Bonnet theory,” *Phys. Rev. D* **46**, 2464 (1992).
- [33] G. Dotti, J. Oliva and R. Troncoso, “Static wormhole solution for higher-dimensional gravity in vacuum,” *Phys. Rev. D* **75**, 024002 (2007).
- [34] M. R. Mehdizadeh, M. Kord Zangeneh and F. S. N. Lobo, “Einstein-Gauss-Bonnet traversable wormholes satisfying the weak energy condition,” *Phys. Rev. D* **91**, no. 8, 084004 (2015).
- [35] F. S. N. Lobo, “A General class of braneworld wormholes,” *Phys. Rev. D* **75**, 064027 (2007).
- [36] S. K. Sahu, S. G. Ganebo and G. G. Weldemariam, “Kaluza-Klein Tilted Cosmological Model in Lyra Geometry,” *Iran. J. Sci. Technol. A* **42**, no. 3, 1451 (2018).
- [37] L. A. Anchordoqui, S. E. Perez Bergliaffa and D. F. Torres, “Brans-Dicke wormholes in nonvacuum space-time,” *Phys. Rev. D* **55**, 5226 (1997).
- [38] S. Capozziello, T. Harko, T. S. Koivisto, F. S. N. Lobo and G. J. Olmo, “Wormholes supported by hybrid metric-Palatini gravity,” *Phys. Rev. D* **86**, 127504 (2012).
- [39] T. Harko, F. S. N. Lobo, S. Nojiri and S. D. Odintsov, “ $f(R, T)$  gravity,” *Phys. Rev. D* **84**, 024020 (2011) [arXiv:1104.2669 [gr-qc]].
- [40] A. A. Starobinsky, “A New Type of Isotropic Cosmological Models Without Singularity,” *Phys. Lett.* **91B**, 99 (1980).
- [41] O. Bertolami, C. G. Boehmer, T. Harko and F. S. N. Lobo, “Extra force in  $f(R)$  modified theories of gravity,” *Phys. Rev. D* **75**, 104016 (2007).
- [42] S. Nojiri, S. D. Odintsov, “Introduction to modified gravity and gravitational alternative for dark energy,” *Int. J. Geom. Methods Mod. Phys.* **4**, 115 (2007).
- [43] O. Bertolami, P. Frazao and J. Paramos, “Accelerated expansion from a non-minimal gravitational coupling to matter,” *Phys. Rev. D* **81**, 104046 (2010).
- [44] K. Bamba, S. Capozziello, S. Nojiri and S. D. Odintsov, “Dark energy cosmology: the equivalent description via different theoretical models and cosmography tests,” *Astrophys. Space Sci.* **342**, 155 (2012).
- [45] S. Nojiri, S. D. Odintsov and M. Sami, “Dark energy cosmology from higher-order, string-inspired gravity and its reconstruction,” *Phys. Rev. D* **74**, 046004 (2006).

- [46] P. H. R. S. Moraes, J. D. V. Arbail and M. Malheiro, “Stellar equilibrium configurations of compact stars in  $f(R, T)$  gravity,” JCAP **1606**, 005 (2016).
- [47] M. Zubair, G. Abbas and I. Noureen, “Possible formation of compact stars in  $f(R, T)$  gravity,” Astrophys. Space Sci. **361**, no. 1, 8 (2016).
- [48] M. F. Shamir, “Locally rotationally symmetric Bianchi type I cosmology in  $f(R, T)$  gravity,” Eur. Phys. J. C **75**, no. 8, 354 (2015).
- [49] M. Zubair and I. Noureen, “Evolution of axially symmetric anisotropic sources in  $f(R, T)$  gravity,” Eur. Phys. J. C **75**, no. 6, 265 (2015).
- [50] D. Momeni, P. H. R. S. Moraes and R. Myrzakulov, “Generalized second law of thermodynamics in  $f(R, T)$  theory of gravity,” Astrophys. Space Sci. **361**, no. 7, 228 (2016).
- [51] T. Harko, “Thermodynamic interpretation of the generalized gravity models with geometry - matter coupling,” Phys. Rev. D **90**, no. 4, 044067 (2014).
- [52] P. H. R. S. Moraes, G. Ribeiro and R. A. C. Correa, “A transition from a decelerated to an accelerated phase of the universe expansion from the simplest non-trivial polynomial function of  $T$  in the  $f(R, T)$  formalism,” Astrophys. Space Sci. **361**, no. 7, 227 (2016).
- [53] P. H. R. S. Moraes and R. A. C. Correa, “Braneworld cosmology in  $f(R, T)$  gravity,” Astrophys. Space Sci. **361**, no. 3, 91 (2016).
- [54] R. A. C. Correa and P. H. R. S. Moraes, “Configurational entropy in  $f(R, T)$  brane models,” Eur. Phys. J. C **76**, no. 2, 100 (2016).
- [55] P. H. R. S. Moraes and J. R. L. Santos, “A complete cosmological scenario from  $f(R, T^\phi)$  gravity theory,” Eur. Phys. J. C **76**, 60 (2016).
- [56] P. H. R. S. Moraes, “Cosmological solutions from Induced Matter Model applied to 5D  $f(R, T)$  gravity and the shrinking of the extra coordinate,” Eur. Phys. J. C **75**, no. 4, 168 (2015).
- [57] F. G. Alvarenga, A. de la Cruz-Dombriz, M. J. S. Houndjo, M. E. Rodrigues and D. Sez-Gmez, “Dynamics of scalar perturbations in  $f(R, T)$  gravity,” Phys. Rev. D **87**, no. 10, 103526 (2013).
- [58] R. Myrzakulov, “FRW Cosmology in  $F(R, T)$  gravity,” Eur. Phys. J. C **72**, 2203 (2012).
- [59] U. K. Sharma and A. Pradhan, “Cosmology in modified  $f(R, T)$ -gravity theory in a variant  $\Lambda(T)$  scenario-revisited,” Int. J. Geom. Meth. Mod. Phys. **15**, no. 01, 1850014 (2017).
- [60] R. Nagpal, S. K. J. Pacif, J. K. Singh, K. Bamba and A. Beesham, “Analysis with observational constraints in  $\Lambda$ -cosmology in  $f(R, T)$  gravity,” Eur. Phys. J. C **78**, no. 11, 946 (2018).
- [61] P. S. Debnath, “Bulk viscous cosmological model in  $f(R, T)$  theory of gravity” arXiv preprint arXiv:1907.02238 (2019).

- [62] N. Ahmed and S. Z. Alamri, “A stable flat universe with variable cosmological constant in  $f(R, T)$  gravity,” Res. Astron. Astrophys. **18**, no. 10, 123 (2018).
- [63] U. K. Sharma, R. Zia, A. Pradhan and A. Beesham, “Stability of LRS Bianchi type-I cosmological models in  $f(R, T)$ -gravity,” Res. Astron. Astrophys. **19**, no. 4, 055 (2019).
- [64] T. Azizi, “Wormhole Geometries In  $f(R, T)$  Gravity,” Int. J. Theor. Phys. **52**, 3486 (2013).
- [65] M. Zubair, S. Waheed and Y. Ahmad, “Static spherically symmetric wormholes in  $f(R, T)$  gravity,” Eur. Phys. J. C **76**, no. 8, 444 (2016).
- [66] M. Zubair, S. Waheed and Y. Ahmad, “Static spherically symmetric wormholes in  $f(R, T)$  gravity,” Eur. Phys. J. C **76**, no. 8, 444 (2016).
- [67] M. Zubair, G. Mustafa, S. Waheed and G. Abbas, “Existence of stable wormholes on a non-commutative-geometric background in modified gravity,” Eur. Phys. J. C **77**, no. 10, 680 (2017).
- [68] P. H. R. S. Moraes, R. A. C. Correa and R. V. Lobato, “Analytical general solutions for static wormholes in  $f(R, T)$  gravity” J. Cosmol. Astropart. Phys. **2017** (2017).
- [69] G. C. Samanta, N. Godani and K. Bamba, “Traversable Wormholes with Exponential Shape Function in Modified Gravity and in General Relativity: A Comparative Study” arXiv:1811.06834v1 [gr-qc] (2018).
- [70] E. Elizalde and M. Khurshudyan, “Wormhole formation in  $f(R, T)$  gravity: Varying Chaplygin gas and barotropic fluid,” Phys. Rev. D **98** 123525 (2018).
- [71] M. Z. Bhatti, Z. Yousaf and M. Ilyas, “Existence of wormhole solutions and energy conditions in  $f(R, T)$  gravity,” J. Astrophys. Astr. **39** 69 (2018).
- [72] P. H. R. S. Moraes, W. de Paula and R. A. C. Correa, “Charged wormholes in  $f(R, T)$  extended theory of gravity,” Int. J. Mod. Phys. D **28**, no. 08, 1950098 (2019).
- [73] M. Sharif and I. Nawazish, “Viable wormhole solutions and Noether symmetry in  $f(R, T)$  gravity,” Annals Phys. **400**, 37 (2019).
- [74] E. Elizalde and M. Khurshudyan, “Wormholes with  $\rho(R, R')$  matter in  $f(R, T)$  gravity,” Phys. Rev. D **99**, no. 2, 024051 (2019).
- [75] P. H. R. S. Moraes, P. K. Sahoo, “Wormholes in exponential  $f(R, T)$  gravity,” arXiv preprint arXiv:1903.03421 (2019).
- [76] P. Sahoo, A. Kirschner, P. K. Sahoo, “Phantom fluid wormhole in  $f(R, T)$  gravity,” arXiv preprint arXiv:1906.04048 (2019).
- [77] A. Banerjee, K. Singh, M. K. Jasim, F. Rahaman, “Traversable wormholes in  $f(R, T)$  gravity with conformal motions,” arXiv preprint arXiv:1908.04754 (2019).
- [78] M. Amir, A. Banerjee, S. D. Maharaj, “Shadow of charged wormholes in EinsteinMaxwell-dilaton theory,” Annals Phys. **400** 198-207 (2019).



- [79] A. Banerjee, P. H. R. S. Moraes, R. A. C. Correa, G. Ribeiro, “Wormholes in Randall-Sundrum braneworld,” arXiv preprint arXiv:1904.10310 (2019).
- [80] P. H. R. S. Moraes, P. K. Sahoo, “Modeling wormholes in  $f(R, T)$  gravity,” Phys. Rev. D, **96** (4), 044038 (2017).
- [81] N. Godani and G. C. Samanta, “Traversable Wormholes and Energy Conditions with Two Different Shape Functions in  $f(R)$  Gravity,” Int. J. Mod. Phys. D **28**, no. 02, 1950039 (2018).
- [82] N. Godani, G. C. Samanta, “Static Traversable Wormholes in  $f(R, T) = R + 2\ln T$  Gravity,” Chin. J. Phys. **62**, 161-171 (2019).
- [83] L.D Landau and E.M. Lifshitz, ”The Classical Theory of Fields” (Oxford, UK, 1998).
- [84] T. Harko and F. S. N. Lobo, “ $f(R, L_m)$  gravity,” Eur. Phys. J. C **70**, 373 (2010).
- [85] T. Harko and F. S. N. Lobo, “Generalized curvature-matter couplings in modified gravity,” Galaxies **2**, no. 3, 410 (2014).
- [86] O. Bertolami, F. S. N. Lobo and J. Paramos, “Non-minimum coupling of perfect fluids to curvature,” Phys. Rev. D **78** (2008) 064036.
- [87] M. S. Morris and K. S. Thorne, “Wormholes in space-time and their use for interstellar travel: A tool for teaching general relativity,” Am. J. Phys. **56**, 395 (1988).
- [88] M. Visser, ”Lorentzian wormholes: From Einstein to Hawking,” (AIP Press, New York, 1995).
- [89] M. Cataldo, P. Meza and P. Minning, “N-dimensional static and evolving Lorentzian wormholes with cosmological constant,” Phys. Rev. D **83**, 044050 (2011).
- [90] F. Rahaman, M. Kalam, M. Sarker, A. Ghosh and B. Raychaudhuri, “Wormhole with varying cosmological constant,” Gen. Rel. Grav. **39**, 145 (2007).