

# Regularity of the solution to fractional diffusion, advection, reaction equations

V.J. Ervin\*

January 24, 2022

## Abstract

In this report we investigate the regularity of the solution to the fractional diffusion, advection, reaction equation on a bounded domain in  $\mathbb{R}^1$ . The analysis is performed in the weighted Sobolev spaces,  $H_{(a,b)}^s(I)$ . Three different characterizations of  $H_{(a,b)}^s(I)$  are presented, together with needed embedding theorems for these spaces. The analysis shows that the regularity of the solution is bounded by the endpoint behavior of the solution, which is determined by the parameters  $\alpha$  and  $r$  defining the fractional diffusion operator. Additionally, the analysis shows that for a sufficiently smooth right hand side function, the regularity of the solution to fractional diffusion reaction equation is lower than that of the fractional diffusion equation. Also, the regularity of the solution to fractional diffusion advection reaction equation is two orders lower than that of the fractional diffusion reaction equation.

**Key words.** Fractional diffusion equation, regularity, weighted Sobolev spaces

**AMS Mathematics subject classifications.** 35R11, 35B65, 46E35

## 1 Introduction

Of interest in this report is the regularity of the solution of the fractional diffusion equation

$$\mathcal{L}_r^\alpha u(x) := -(rD^\alpha + (1-r)D^{\alpha*})u(x) = f(x), \quad x \in I, \quad (1.1)$$

$$\text{subject to } u(0) = u(1) = 0, \quad (1.2)$$

and the regularity of the solution of the fractional diffusion, advection, reaction equation

$$\mathcal{L}_r^\alpha u(x) + b(x)Du(x) + c(x)u(x) = f(x), \quad x \in I, \quad (1.3)$$

$$\text{subject to } u(0) = u(1) = 0, \quad (1.4)$$

where  $I := (0, 1)$ ,  $1 < \alpha < 2$ ,  $0 \leq r \leq 1$ ,  $c(x) - \frac{1}{2}Db(x) \geq 0$ ,  $D$  denotes the usual derivative operator,  $D^\alpha$  the  $\alpha$ -order left fractional derivative operator, and  $D^{\alpha*}$  the  $\alpha$ -order right fractional

---

\*School of Mathematical and Statistical Sciences, Clemson University, Clemson, South Carolina 29634-0975, USA.  
email: [vjervin@clemson.edu](mailto:vjervin@clemson.edu).

derivative operator, defined by:

$$D^\alpha u(x) := D \frac{1}{\Gamma(2-\alpha)} \int_0^x \frac{1}{(x-s)^{\alpha-1}} Du(s) ds, \quad (1.5)$$

$$D^{\alpha*} u(x) := D \frac{1}{\Gamma(2-\alpha)} \int_x^1 \frac{1}{(s-x)^{\alpha-1}} Du(s) ds. \quad (1.6)$$

The regularity of the solution to a differential equation plays a fundamental role in designing optimal approximation schemes for the solution.

In recent years fractional order differential equations have received increased attention due to their application in the modeling of physical phenomena such as in contaminant transport in ground water flow [7, 12], viscoelasticity [26], image processing [5, 10, 16, 17], turbulent flow [25, 34], and chaotic dynamics [37].

The diffusion operator,  $\mathcal{L}_r^\alpha$ , arises in a random walk process in which the jumps have an unbounded variance (Lévy process) [7, 30].

For the case  $r = 1/2$ ,  $\mathcal{L}_{1/2}^\alpha$  represents the *integral* fractional Laplacian operator [2]. The existence, uniqueness and regularity of the solution to the fractional Laplacian equation has been investigated by a number of authors, in  $\mathbb{R}^1$  see [3], in  $\mathbb{R}^{n \geq 2}$  [4, 11, 19, 31]. Recently the regularity results for the fractional Laplacian equation was extended by Hao and Zhang in [22, 38] to the fractional Laplacian equation with a constant advection and reaction term (i.e. (1.3),(1.4), for  $r = 1/2$ ,  $b(x) = b$ ,  $c(x) = c$ ,  $b, c \in \mathbb{R}$ ).

Fewer results on the regularity of the solution to the general fractional diffusion, advection, reaction equation have been established. In [14] Ervin and Roop established existence and uniqueness of solution,  $u \in H_0^{\alpha/2}(\mathbf{I})$ , for  $f \in H^{-\alpha/2}(\mathbf{I})$ . More recently in [13, 23] precise regularity results were obtained for the solution of (1.1),(1.2) for  $f \in H_{(a,b)}^s(\mathbf{I})$ , where  $H_{(a,b)}^s(\mathbf{I})$  denotes an appropriated weighted Sobolev space. In [21] Hao, Guang and Zhang obtained regularity estimates for the solution of (1.3) for  $b(x) = 0$ ,  $c(x) = c$ . Their numerical experiments indicated that their regularity estimates were not optimal.

In this article we present the general regularity results for (1.3),(1.4), in appropriately weighted Sobolev spaces. The analysis establishes that the presence of a reaction term (i.e.  $c(x) \neq 0$ ) limits the regularity of the solution, regardless of the smoothness of the right hand side function,  $f(x)$ . This reduction in regularity is greater (by a factor of 2) when an advective term (i.e.  $b(x) \neq 0$ ) appears in (1.3). This behavior of the solution is in sharp contrast to that for the integer order ( $\alpha = 2$ ) diffusion, advection, reaction equation. In that case, assuming  $b(x)$  and  $c(x)$  are sufficiently regular, for the right hand side function  $f \in H^s(\mathbf{I})$  the solution lies in  $H^{s+2}(\mathbf{I})$ .

The results we present herein extend those in [13] for the fractional diffusion equation, and those in [22] for the fractional Laplacian equation with a constant advection and reaction term. The proofs given are significantly different than those used in [21, 22].

The analysis of (1.3),(1.4) is most appropriately performed in weighted Sobolev spaces (due to the singular behavior of the solution at the endpoints). There are different ways to define the weighted Sobolev spaces: (i) using interpolation (Section 3), (ii) using an appropriate basis (Section 4), (iii) using an explicit definition for the fractional order norms (Section 6). Each of these representations have their advantages and are used in the analysis.

This paper is organized as follows. In the next section we present some preliminary definitions and results. Sections 3, 4, and 6 present the three different (but equivalent) definitions of the weighted Sobolev spaces. Section 4 also establishes some useful properties of the space  $H_{(a,b)}^s(\mathbb{I})$ . For example, which  $H_{(a,b)}^s(\mathbb{I})$  space the function  $f(x) = x^\mu$  lies in, and for which  $H_{(a,b)}^s(\mathbb{I})$  we have the embedding  $H_{(a,b)}^s(\mathbb{I}) \subset C^k(\mathbb{I})$ . The regularity of the solution to the fractional diffusion problem (1.1),(1.2), is discussed in Section 5. The theorems needed for determining which  $H_{(b,a)}^t(\mathbb{I})$  space  $(1-x)^a x^b \phi(x)$  lies in, when  $\phi(x) \in H_{(a,b)}^s(\mathbb{I})$ , are established in Section 6. Section 7 then discusses the regularity of solutions to (1.3),(1.4).

In the last section we relate the regularity results obtained in weighted Sobolev spaces to the usual (unweighted) Sobolev spaces.

## 2 Notation and Properties

Jacobi polynomial have an important connection with fractional order diffusion equations [3, 13, 28, 27]. We briefly review their definition and some of their important properties [1, 35].

**Usual Jacobi Polynomials,  $P_n^{(a,b)}(t)$ , on  $(-1, 1)$ .**

Definition:  $P_n^{(a,b)}(t) := \sum_{m=0}^n p_{n,m} (t-1)^{(n-m)}(t+1)^m$ , where

$$p_{n,m} := \frac{1}{2^n} \binom{n+a}{m} \binom{n+b}{n-m}. \quad (2.1)$$

Orthogonality:

$$\int_{-1}^1 (1-t)^a (1+t)^b P_j^{(a,b)}(t) P_k^{(a,b)}(t) dt = \begin{cases} 0, & k \neq j \\ |||P_j^{(a,b)}|||^2, & k = j \end{cases},$$

$$\text{where } |||P_j^{(a,b)}||| = \left( \frac{2^{(a+b+1)}}{(2j+a+b+1)} \frac{\Gamma(j+a+1) \Gamma(j+b+1)}{\Gamma(j+1) \Gamma(j+a+b+1)} \right)^{1/2}. \quad (2.2)$$

In order to transform the domain of the family of Jacobi polynomials to  $[0, 1]$ , let  $t \rightarrow 2x - 1$  and introduce  $G_n^{(a,b)}(x) = P_n^{(a,b)}(t(x))$ . From (2.2),

$$\begin{aligned} \int_{-1}^1 (1-t)^a (1+t)^b P_j^{(a,b)}(t) P_k^{(a,b)}(t) dt &= \int_0^1 2^a (1-x)^a 2^b x^b P_j^{(a,b)}(2x-1) P_k^{(a,b)}(2x-1) 2 dx \\ &= 2^{a+b+1} \int_0^1 (1-x)^a x^b G_j^{(a,b)}(x) G_k^{(a,b)}(x) dx \\ &= \begin{cases} 0, & k \neq j, \\ 2^{a+b+1} |||G_j^{(a,b)}|||^2, & k = j. \end{cases} \end{aligned}$$

$$\text{where } |||G_j^{(a,b)}||| = \left( \frac{1}{(2j+a+b+1)} \frac{\Gamma(j+a+1) \Gamma(j+b+1)}{\Gamma(j+1) \Gamma(j+a+b+1)} \right)^{1/2}. \quad (2.3)$$

$$\text{Note that } |||G_j^{(a,b)}||| = |||G_j^{(b,a)}|||. \quad (2.4)$$

From [27, equation (2.19)] we have that

$$\frac{d^k}{dt^k} P_n^{(a,b)}(t) = \frac{\Gamma(n+k+a+b+1)}{2^k \Gamma(n+a+b+1)} P_{n-k}^{(a+k,b+k)}(t). \quad (2.5)$$

Hence,

$$\frac{d^k}{dx^k} G_n^{(a,b)}(x) = \frac{\Gamma(n+k+a+b+1)}{\Gamma(n+a+b+1)} G_{n-k}^{(a+k,b+k)}(x). \quad (2.6)$$

Also, from [27, equation (2.15)],

$$\frac{d^k}{dt^k} \left\{ (1-t)^{a+k} (1+t)^{b+k} P_{n-k}^{(a+k,b+k)}(t) \right\} = \frac{(-1)^k 2^k n!}{(n-k)!} (1-t)^a (1+t)^b P_n^{(a,b)}(t), \quad n \geq k \geq 0, \quad (2.7)$$

from which it follows that

$$\frac{d^k}{dx^k} \left\{ (1-x)^{a+k} x^{b+k} G_{n-k}^{(a+k,b+k)}(x) \right\} = \frac{(-1)^k n!}{(n-k)!} (1-x)^a x^b G_n^{(a,b)}(x). \quad (2.8)$$

For compactness of notation we introduce

$$\rho^{(a,b)} = \rho^{(a,b)}(x) := (1-x)^a x^b. \quad (2.9)$$

We let  $\mathbb{N}_0 := \mathbb{N} \cup 0$  and use  $y_n \sim n^p$  to denote that there exists constants  $c$  and  $C > 0$  such that, as  $n \rightarrow \infty$ ,  $c n^p \leq |y_n| \leq C n^p$ . Additionally, we use  $a \lesssim b$  to denote that there exists a constant  $C$  such that  $a \leq C b$ .

For  $s \in \mathbb{R}$ ,  $\lfloor s \rfloor$  is used to denote the largest integer that is less than or equal to  $s$ , and  $\lceil s \rceil$  is used to denote the smallest integer that is greater than or equal to  $s$ .

Note, from Stirling's formula we have that

$$\lim_{n \rightarrow \infty} \frac{\Gamma(n+\sigma)}{\Gamma(n) n^\sigma} = 1, \quad \text{for } \sigma \in \mathbb{R}. \quad (2.10)$$

**Definition: Condition A**

For  $\alpha$  and  $r$  given, satisfying  $1 < \alpha < 2$ ,  $0 \leq r \leq 1$ , let  $\beta$  be determined by  $\alpha - 1 \leq \beta \leq 1$  and

$$r = \frac{\sin(\pi\beta)}{\sin(\pi(\alpha-\beta)) + \sin(\pi\beta)}. \quad (2.11)$$

Furthermore, introduce the constant  $c_*^*$  defined by

$$c_*^* = \frac{\sin(\pi\alpha)}{\sin(\pi(\alpha-\beta)) + \sin(\pi\beta)}. \quad (2.12)$$

**Function space**  $L_\omega^2(\mathbf{I})$ .

For  $\omega(x) > 0$ ,  $x \in (0, 1)$ , let

$$L_\omega^2(\mathbf{I}) := \{f(x) : \int_0^1 \omega(x) f(x)^2 dx < \infty\}. \quad (2.13)$$

Associated with  $L_\omega^2(0, 1)$  is the inner product,  $\langle \cdot, \cdot \rangle_\omega$ , and norm,  $\|\cdot\|_\omega$ , defined by

$$\langle f, g \rangle_\omega := \int_0^1 \omega(x) f(x) g(x) dx, \quad \text{and} \quad \|f\|_\omega := (\langle f, f \rangle_\omega)^{1/2}.$$

The set of orthogonal polynomials  $\{G_j^{(a,b)}\}_{j=0}^\infty$  form an orthogonal basis for  $L_{\rho^{(a,b)}}^2(\mathbf{I})$ .

### 3 Weighted Sobolev Space defined by interpolation

Following Babuška and Guo [6], and Guo and Wang [20],  $n \in \mathbb{N}_0$  define the weighted Sobolev spaces

$$H_{\rho^{(a,b)}}^n(\mathbf{I}) := \left\{ v : \sum_{j=0}^n \|D^j v\|_{\rho^{(a+j,b+j)}}^2 < \infty \right\}, \quad (3.1)$$

with associated norm  $\|v\|_{n,\rho^{(a,b)}} := \left( \sum_{j=0}^n \|D^j v\|_{\rho^{(a+j,b+j)}}^2 \right)^{1/2}$ .

Definition (3.1) is extended to  $s \in \mathbb{R}^+$  using the  $K$ - method of interpolation. For  $s < 0$  the spaces are defined by (weighted)  $L^2$  duality.

Babuška and Guo used the  $H_{\rho^{(a,b)}}^s(\mathbf{I})$  spaces in establishing the optimal convergence properties of the  $p$ - version of the finite element method. They also related the definition (3.1) to the decay property of the coefficients of the Jacobi polynomials of the expansion of the function  $v$ .

In [20] Guo and Wang derived approximation properties of Jacobi polynomials for functions in the weighted Sobolev spaces (3.1).

### 4 Weighted Sobolev Space defined by Jacobi coefficients

In this section we define function spaces in terms of the decay property of the Jacobi coefficients of their member functions. We then show that these spaces agree with the weighted Sobolev spaces defined using the  $K$ - method of interpolation.

This presentation parallels the work of Acosta, Borthagaray, Bruno and Maas in [3] who investigated the regularity and approximation of the 1-d fractional Laplacian equation. The 1-d fractional Laplacian operator they considered is a special case of the fractional diffusion operator,  $\mathcal{L}_r^\alpha(\cdot)$ , for the parameter  $r = 1/2$ . In [3] their analysis focused on the coefficients of the Gegenbauer polynomials, which are a special case of the Jacobi polynomials where the weight parameters are equal.

We begin by relating the decay rate of a function's Jacobi polynomial coefficient to its regularity. We then introduce the weighted Sobolev space  $H_{(a,b)}^s(\mathbf{I})$  and show that it corresponds to the space  $H_{\rho^{(a,b)}}^s(\mathbf{I})$ . We conclude the section with a corollary describing the precise solvability of (1.1),(1.2), an embedding theorem relating  $H_{(a,b)}^s(\mathbf{I})$  to  $C^k(\mathbf{I})$ , and a corollary stating a suitable sufficient condition for the solution of (1.1),(1.2) to be continuous.

Introduce  $\tilde{G}_j^{(a,b)}(x) = G_j^{(a,b)}(x)/\|G_j^{(a,b)}\|$  as an orthonormal basis for  $L_{\rho^{(a,b)}}^2(\mathbf{I})$ .

Then, given  $v \in L_{\rho^{(a,b)}}^2(\mathbf{I})$ , we have

$$v(x) = \sum_{j=0}^{\infty} v_j \tilde{G}_j^{(a,b)}(x), \quad (4.1)$$

which converges in  $L_{\rho^{(a,b)}}^2(\mathbf{I})$ , where

$$v_j = \int_0^1 \rho^{(a,b)}(x) v(x) \tilde{G}_j^{(a,b)}(x) dx. \quad (4.2)$$

Using (2.6) and (4.1) we could conjecture (by differentiating (4.1)) that for sufficiently smooth  $v(x)$

$$\begin{aligned} v^{(k)}(x) &= \sum_{j=k}^{\infty} \frac{\|G_{j-k}^{(a+k,b+k)}\|}{\|G_j^{(a,b)}\|} \frac{\Gamma(j+k+a+b+1)}{\Gamma(j+a+b+1)} v_j \tilde{G}_{j-k}^{(a+k,b+k)}(x) \\ &:= \sum_{j=k}^{\infty} v_{j-k}^{(k)} \tilde{G}_{j-k}^{(a+k,b+k)}(x), \end{aligned} \quad (4.3)$$

where

$$v_{j-k}^{(k)} = \frac{\|G_{j-k}^{(a+k,b+k)}\|}{\|G_j^{(a,b)}\|} \frac{\Gamma(j+k+a+b+1)}{\Gamma(j+a+b+1)} v_j. \quad (4.4)$$

**Remark:** The terms  $v_j$  and  $v_{j-k}^{(k)}$  denote real numbers, whereas  $v^{(k)}(x)$  is used to represent the  $k^{th}$  derivative of  $v(x)$ . (From (4.3),  $v_{j-k}^{(k)}$ 's are the Jacobi coefficients for  $v^{(k)}(x)$ .)

We investigate (4.4) more rigorously using the following lemma.

**Lemma 4.1** (See [3, Lemma 4.2] *Integration by parts*)

Let  $k \in \mathbb{N}$ ,  $k \geq 2$ , and let  $v \in C^{k-2}[0, 1]$  such that for the decomposition of  $[0, 1] = \cup_{i=1}^n [x_i, x_{i+1}]$  ( $0 = x_1 < x_2 < \dots < x_{n+1} = 1$ ), and for functions  $\hat{v}_i \in C^k[x_i, x_{i+1}]$ , we have  $v(x) = \hat{v}_i(x)$  for  $x \in (x_i, x_{i+1})$ , and  $1 \leq i \leq n$ . Then for  $j \geq k$  the  $(a, b)$  Jacobi coefficient defined by (4.2) satisfies

$$\begin{aligned} v_j &= B_j^k \int_0^1 \rho^{(a+k,b+k)}(x) \tilde{G}_{j-k}^{(a+k,b+k)}(x) v^{(k)}(x) dx \\ &\quad - B_j^k \sum_{l=1}^n \rho^{(a+k,b+k)}(x) \tilde{G}_{j-k}^{(a+k,b+k)}(x) v_l^{(k-1)}(x) \Big|_{x=x_l}^{x_{l+1}}, \end{aligned} \quad (4.5)$$

$$\text{where } B_j^k = \frac{\Gamma(j-k+1)}{\Gamma(j+1)} \frac{\|G_{j-k}^{(a+k,b+k)}\|}{\|G_j^{(a,b)}\|}. \quad (4.6)$$

**Proof:** From (2.8),

$$\int \rho^{(a,b)}(x) G_n^{(a,b)}(x) dx = \frac{-(n-1)!}{n!} \rho^{(a+1,b+1)}(x) G_{n-1}^{(a+1,b+1)}(x) = \frac{-1}{n} \rho^{(a+1,b+1)}(x) G_{n-1}^{(a+1,b+1)}(x).$$

Hence,

$$\int \rho^{(a,b)}(x) \tilde{G}_n^{(a,b)}(x) dx = \frac{-1}{n} \frac{|||G_{n-1}^{(a+1,b+1)}|||}{|||G_n^{(a,b)}|||} \rho^{(a+1,b+1)}(x) \tilde{G}_{n-1}^{(a+1,b+1)}(x).$$

Beginning with (4.2) and using integration by parts,

$$\begin{aligned} v_j &= \sum_{l=1}^n \int_{x_l}^{x_{l+1}} \hat{v}_l(x) \rho^{(a,b)}(x) \tilde{G}_j^{(a,b)}(x) dx \\ &= \sum_{l=1}^n \int_{x_l}^{x_{l+1}} \frac{1}{j} \frac{|||G_{j-1}^{(a+1,b+1)}|||}{|||G_j^{(a,b)}|||} \rho^{(a+1,b+1)}(x) \tilde{G}_{j-1}^{(a+1,b+1)}(x) \hat{v}_l^{(1)}(x) dx \\ &\quad - \frac{1}{j} \frac{|||G_{j-1}^{(a+1,b+1)}|||}{|||G_j^{(a,b)}|||} \rho^{(a+1,b+1)}(x) \tilde{G}_{j-1}^{(a+1,b+1)}(x) \hat{v}_l(x) \Big|_{x=x_l}^{x_{l+1}} \\ &= \sum_{l=1}^n \int_{x_l}^{x_{l+1}} \frac{1}{j} \frac{|||G_{j-1}^{(a+1,b+1)}|||}{|||G_j^{(a,b)}|||} \rho^{(a+1,b+1)}(x) \tilde{G}_{j-1}^{(a+1,b+1)}(x) \hat{v}_l^{(1)}(x) dx. \end{aligned} \quad (4.7)$$

Repeated use of integration by parts, and using that

$$\frac{1}{j} \frac{1}{j-1} \cdots \frac{1}{j-k+1} = \frac{\Gamma(j-k+1)}{\Gamma(j+1)},$$

(4.7) becomes

$$\begin{aligned} v_j &= \sum_{l=1}^n \int_{x_l}^{x_{l+1}} \frac{\Gamma(j-k+1)}{\Gamma(j+1)} \frac{|||G_{j-k}^{(a+k,b+k)}|||}{|||G_j^{(a,b)}|||} \rho^{(a+k,b+k)}(x) \tilde{G}_{j-k}^{(a+k,b+k)}(x) \hat{v}_l^{(k)}(x) dx \\ &\quad - \frac{\Gamma(j-k+1)}{\Gamma(j+1)} \frac{|||G_{j-k}^{(a+k,b+k)}|||}{|||G_j^{(a,b)}|||} \rho^{(a+k,b+k)}(x) \tilde{G}_{j-k}^{(a+k,b+k)}(x) \hat{v}_l^{(k-1)}(x) \Big|_{x=x_l}^{x_{l+1}} \\ &= \frac{\Gamma(j-k+1)}{\Gamma(j+1)} \frac{|||G_{j-k}^{(a+k,b+k)}|||}{|||G_j^{(a,b)}|||} \int_0^1 \rho^{(a+k,b+k)}(x) \tilde{G}_{j-k}^{(a+k,b+k)}(x) v^{(k)}(x) dx \\ &\quad - \frac{\Gamma(j-k+1)}{\Gamma(j+1)} \frac{|||G_{j-k}^{(a+k,b+k)}|||}{|||G_j^{(a,b)}|||} \sum_{l=1}^n \rho^{(a+k,b+k)}(x) \tilde{G}_{j-k}^{(a+k,b+k)}(x) \hat{v}_l^{(k-1)}(x) \Big|_{x=x_l}^{x_{l+1}}, \end{aligned} \quad (4.8)$$

which corresponds with (4.5). ■

Note, for  $v \in C^k[0,1]$  we have that the  $j^{th}$  Jacobi polynomial coefficient  $v_j^{(k)}$  is given by

$$v_j^{(k)} = \int_0^1 \rho^{(a+k,b+k)}(x) \tilde{G}_j^{(a+k,b+k)}(x) dx,$$

and from (4.8)

$$v_{j-k}^{(k)} = \frac{\Gamma(j+1)}{\Gamma(j-k+1)} \frac{|||G_j^{(a,b)}|||}{|||G_{j-k}^{(a+k,b+k)}|||} v_j, \quad \text{for } j \geq k. \quad (4.9)$$

Using (2.2), a simple calculation establishes the equivalence of (4.9) and (4.4).

In order to investigate how the decay rate of the Jacobi coefficients relate to the regularity of the function we establish the following lemma.

**Lemma 4.2** *There exists constants  $C_1$  and  $C_2$  such that*

$$C_1 j^{-k} \leq \frac{\Gamma(j-k+1)}{\Gamma(j+1)} \frac{|||G_{j-k}^{(a+k,b+k)}|||}{|||G_j^{(a,b)}|||} \leq C_2 j^{-k}. \quad (4.10)$$

**Proof:** We have that

$$\frac{\Gamma(j-k+1)}{\Gamma(j+1)} = \frac{\Gamma(j-k+1)}{j(j-1)\dots(j-k+1)\Gamma(j-k+1)} = \frac{1}{j(j-1)\dots(j-k+1)} \sim j^{-k}. \quad (4.11)$$

Next consider

$$\begin{aligned} \frac{|||G_{j-k}^{(a+k,b+k)}|||^2}{|||G_j^{(a,b)}|||^2} &= \frac{(2j+a+b+1)}{1} \frac{\Gamma(j+1)\Gamma(j+a+b+1)}{\Gamma(j+a+1)\Gamma(j+b+1)} \\ &\quad \cdot \frac{1}{(2j-2k+a+k+b+k+1)} \frac{\Gamma(j-k+a+k+1)\Gamma(j-k+b+k+1)}{\Gamma(j-k+1)\Gamma(j-k+a+k+b+k+1)} \\ &= \frac{\Gamma(j+1)\Gamma(j+a+b+1)}{\Gamma(j+1-k)\Gamma(j+a+b+1+k)} \\ &\sim (j+1)^k (j+a+b+1)^{-k} \sim \left( \frac{j+1}{j+1+a+b} \right)^k \\ &\sim 1. \end{aligned} \quad (4.12)$$

From (4.11) and (4.12), (4.10) follows. ■

**Corollary 4.1** *(See [3, Corollary 4.3]) Let  $k \in \mathbb{N}$  and  $v$  satisfy the hypothesis of Lemma 4.1. Then the Jacobi coefficients in (4.5) are quantities of order  $O(j^{-k})$  as  $j \rightarrow \infty$ , i.e.,*

$$|v_j| < C j^{-k}, \quad (4.13)$$

for a constant  $C$  that depends on  $v$  and  $k$ .

**Remark:** In words, this corollary states that if a function is  $C^k(\mathbb{I})$ , with the possible exception of a few points, then its  $j^{\text{th}}$  Jacobi coefficient must decay as  $C j^{-k}$ , for some constant  $C$ .



**Proof:** In order to establish the estimate, for the first term in (4.5) note that

$$\begin{aligned}
& \left| \int_0^1 \rho^{(a+k, b+k)}(x) \tilde{G}_{j-k}^{(a+k, b+k)}(x) v^{(k)}(x) dx \right| \\
& \leq \left( \int_0^1 \rho^{(a+k, b+k)}(x) (\tilde{G}_{j-k}^{(a+k, b+k)}(x))^2 dx \right)^{1/2} \left( \int_0^1 \rho^{(a+k, b+k)}(x) (v^{(k)}(x))^2 dx \right)^{1/2} \\
& \leq \|v^{(k)}\|_{L^\infty} \left( \int_0^1 \rho^{(a+k, b+k)}(x) dx \right)^{1/2} \\
& = C \|v^{(k)}\|_{L^\infty}.
\end{aligned} \tag{4.14}$$

To bound the second term in (4.5) consider Darboux's formula [35, pg. 196 (8.21.10)]

$$P_n^{(a,b)}(\cos \theta) = n^{-1/2} k(\theta) \cos(N\theta + \gamma) + O(n^{-3/2}), \tag{4.15}$$

$$\begin{aligned}
& \text{where } k(\theta) = \pi^{-1/2} (\sin(\theta/2))^{-a-1/2} (\cos(\theta/2))^{-b-1/2}, \quad N = n + (a+b+1)/2, \\
& \gamma = -(a+1/2)\pi/2, \quad 0 < \theta < \pi.
\end{aligned} \tag{4.16}$$

The bound for the error term holds uniformly in the interval  $[\epsilon, \pi - \epsilon]$ .

Using the trig. identities

$$\cos(\theta/2) = \frac{1}{\sqrt{2}} (1 + \cos(\theta))^{1/2}, \quad \text{and} \quad \sin(\theta/2) = \frac{1}{\sqrt{2}} (1 - \cos(\theta))^{1/2},$$

and the substitution  $x = \cos(\theta)$ , for  $k(\theta)$  we have

$$\begin{aligned}
k(x) &= \pi^{-1/2} 2^{-\frac{1}{2}(-a-1/2)} (1-x)^{\frac{1}{2}(-a-1/2)} 2^{-\frac{1}{2}(-b-1/2)} (1+x)^{\frac{1}{2}(-b-1/2)} \\
&= \pi^{-1/2} 2^{\frac{1}{2}(a+b+1)} (1-x)^{-\frac{1}{2}(a+1/2)} (1+x)^{-\frac{1}{2}(b+1/2)}.
\end{aligned}$$

Therefore, for  $P_n^{(a,b)}(x)$  we have

$$P_n^{(a,b)}(x) = n^{-1/2} \left( \pi^{-1/2} 2^{\frac{1}{2}(a+b+1)} (1-x)^{-\frac{1}{2}(a+1/2)} (1+x)^{-\frac{1}{2}(b+1/2)} \right) \cos(N \operatorname{Arccos}(x) + \gamma) + O(n^{-3/2}). \tag{4.17}$$

Using (4.17), we have for  $(x \rightarrow 2t - 1)$   $G_n^{(a,b)}(t)$ ,

$$G_n^{(a,b)}(t) = n^{-1/2} \left( \pi^{-1/2} (1-t)^{-\frac{1}{2}(a+1/2)} t^{-\frac{1}{2}(b+1/2)} \right) \cos(N \operatorname{Arccos}(2t-1) + \gamma) + O(n^{-3/2}).$$

Now,

$$\begin{aligned}
& \rho^{(a+k, b+k)}(x) G_n^{(a+k, b+k)}(x) = \\
& n^{-1/2} \left( \pi^{-1/2} (1-x)^{\frac{1}{2}(a+k-1)} x^{\frac{1}{2}(b+k-1)} \right) \cos(N \operatorname{Arccos}(2x-1) + \gamma) + O(n^{-3/2}) \\
& \text{i.e., } |\rho^{(a+k, b+k)}(x) G_n^{(a+k, b+k)}(x)| \leq C n^{-1/2},
\end{aligned} \tag{4.18}$$

as  $\left(\pi^{-1/2}(1-x)^{\frac{1}{2}(a+k-1)}x^{\frac{1}{2}(b+k-1)}\right)\cos(N\text{Arccos}(2x-1)+\gamma)$  is bounded for  $x \in [x_2, x_n]$  and on this subinterval the error estimate from Darboux's formula is uniform. (Note that we are not interested in the boundary pieces in the Integration by Parts formula at  $x_1$  and  $x_{n+1}$ , as at these points the boundary pieces are zero because of the  $\rho^{(a+k, b+k)}(x)$  term).

From (2.3) and (2.10) we have

$$\begin{aligned} \|G_j^{(a,b)}\|^2 &= \frac{1}{(2j+a+b+1)} \frac{\Gamma(j+a+1)\Gamma(j+b+1)}{\Gamma(j+1)\Gamma(j+a+b+1)} \\ &\sim j^{-1}(j+1)^a(j+b+1)^{-a} \sim j^{-1}. \end{aligned} \quad (4.19)$$

Combining (4.18) and (4.19) we obtain

$$|\rho^{(a+k, b+k)}(x) \tilde{G}_n^{(a+k, b+k)}(a_l)| \leq C \text{ for } l = 1, 2, \dots, n+1. \quad (4.20)$$

Using (4.14) and (4.20), together with Lemma 4.2 estimate (4.13) follows. ■

Next we introduce the  $(a, b)$ -weighted Sobolev spaces.

**Definition 4.1** Let  $s, a, b \in \mathbb{R}$ ,  $s \geq 0$ ,  $a, b > -1$ ,  $L_{(a,b)}^2(\mathbf{I}) := L_{\rho^{(a,b)}}^2(\mathbf{I})$ , and  $v_j$  be given by (4.2). Then, we define

$$H_{(a,b)}^s(\mathbf{I}) := \{v \in L_{\rho^{(a,b)}}^2(\mathbf{I}) : \sum_{j=0}^{\infty} (1+j^2)^s v_j^2 < \infty\} \quad (4.21)$$

as the  $(a, b)$ -weighted Sobolev space of order  $s$ .

**Lemma 4.3** (See Lemma 4.7 in [3]) Let  $s, a, b \in \mathbb{R}$ ,  $s \geq 0$ ,  $a, b > -1$ . Then the space  $H_{(a,b)}^s(\mathbf{I})$  endowed with the inner product

$$\langle v, w \rangle_{(a,b)}^s := \sum_{j=0}^{\infty} (1+j^2)^s v_j w_j, \text{ and associated norm } \|v\|_{s,(a,b)} := \left( \sum_{j=0}^{\infty} (1+j^2)^s v_j^2 \right)^{1/2} \quad (4.22)$$

is a Hilbert space.

**Proof:** The proof is completely analogously to that given in [24, Theorem 8.2]. ■

**Remarks.**

1. It follows immediately from the definition that for any function  $v \in H_{(a,b)}^s(\mathbf{I})$  its expansion (4.1) is convergent in  $H_{(a,b)}^s(\mathbf{I})$ .
2. From Parseval's identity, it follows that  $H_{(a,b)}^0(\mathbf{I})$  and  $L_{\rho^{(a,b)}}^2(\mathbf{I})$  coincide. Further, we have the dense compact embedding  $H_{(a,b)}^t(\mathbf{I}) \subset H_{(a,b)}^s(\mathbf{I})$  whenever  $t < s$ . (The density of the embedding follows 1. and that all polynomials are contained in  $H_{(a,b)}^s(\mathbf{I})$  for every  $s$ . The compactness follows as in [24, Theorem 8.3]).

**Definition 4.2** Let  $s, a, b \in \mathbb{R}$ ,  $s > 0$ ,  $a, b > -1$ . We denote by  $H_{(a,b)}^{-s}(\mathbb{I})$  the space of bounded linear functionals on  $H_{(a,b)}^s(\mathbb{I})$ .

**Lemma 4.4** For  $s > 0$ ,  $a, b > -1$  the space  $H_{(a,b)}^{-s}(\mathbb{I})$  can be equivalently characterized as the set

$$\mathcal{H}_{(a,b)}^{-s}(\mathbb{I}) := \left\{ f : f(x) = \sum_{j=0}^{\infty} f_j \tilde{G}_j^{(a,b)}(x), \text{ where } \sum_{j=0}^{\infty} (1+j^2)^{-s} f_j^2 < \infty \right\}. \quad (4.23)$$

Additionally, for  $F \in H_{(a,b)}^{-s}(\mathbb{I})$ , with representation  $F = \sum_{j=0}^{\infty} f_j \tilde{G}_j^{(a,b)}(x) := f(x)$

$$\|F\|_{H_{(a,b)}^{-s}} = \sup_{g \in H_{(a,b)}^s(\mathbb{I})} \frac{|F(g)|}{\|g\|_{s,(a,b)}} = \left( \sum_{j=0}^{\infty} (1+j^2)^{-s} f_j^2 \right)^{1/2} := \|f\|_{-s,(a,b)}. \quad (4.24)$$

**Proof:** To establish the characterization we demonstrate that there is an isometry between  $H_{(a,b)}^{-s}(\mathbb{I})$  and  $\mathcal{H}_{(a,b)}^{-s}(\mathbb{I})$ .

Let  $g(x) = \sum_{j=0}^{\infty} g_j \tilde{G}_j^{(a,b)}(x) \in H_{(a,b)}^s(\mathbb{I})$ .

Consider  $F(\cdot) \in H_{(a,b)}^{-s}(\mathbb{I})$ . Then, using the Riesz Representation Theorem, there exists a unique  $h(x) = \sum_{j=0}^{\infty} h_j \tilde{G}_j^{(a,b)}(x) \in H_{(a,b)}^s(\mathbb{I})$ , such that

$$\begin{aligned} F(g) &= \langle h, g \rangle_{(a,b)}^s = \sum_{j=0}^{\infty} (1+j^2)^s h_j g_j \\ &= \langle f, g \rangle_{(a,b)}^0, \end{aligned}$$

where  $f(x) = \sum_{j=0}^{\infty} f_j \tilde{G}_j^{(a,b)}(x)$ , for  $f_j = (1+j^2)^s h_j$ .

Note that as

$$\begin{aligned} \sum_{j=0}^{\infty} (1+j^2)^{-s} f_j^2 &= \sum_{j=0}^{\infty} (1+j^2)^{-s} (1+j^2)^{2s} h_j^2 \\ &= \sum_{j=0}^{\infty} (1+j^2)^s h_j^2 = \|h\|_{s,(a,b)}^2 < \infty, \end{aligned}$$

then  $f \in \mathcal{H}_{(a,b)}^{-s}(\mathbb{I})$ , and from (4.24),  $\|f\|_{-s,(a,b)} = \|h\|_{s,(a,b)}$ .

Next,

$$\begin{aligned}
\|F\|_{H_{(a,b)}^{-s}} &= \sup_{g \in H_{(a,b)}^s(\mathbb{I})} \frac{|F(g)|}{\|g\|_{s,(a,b)}} = \sup_{g \in H_{(a,b)}^s(\mathbb{I})} \frac{|\langle f, g \rangle_{(a,b)}^0|}{\|g\|_{s,(a,b)}} \\
&= \sup_{g \in H_{(a,b)}^s(\mathbb{I})} \frac{|\sum_{j=0}^{\infty} f_j g_j|}{\|g\|_{s,(a,b)}} = \sup_{g \in H_{(a,b)}^s(\mathbb{I})} \frac{|\sum_{j=0}^{\infty} (1+j^2)^{-s/2} f_j (1+j^2)^{s/2} g_j|}{\|g\|_{s,(a,b)}} \\
&\leq \sup_{g \in H_{(a,b)}^s(\mathbb{I})} \frac{\left(\sum_{j=0}^{\infty} (1+j^2)^{-s} f_j^2\right)^{1/2} \left(\sum_{j=0}^{\infty} (1+j^2)^s g_j^2\right)^{1/2}}{\|g\|_{s,(a,b)}} = \sup_{g \in H_{(a,b)}^s(\mathbb{I})} \frac{\|f\|_{-s,(a,b)} \|g\|_{s,(a,b)}}{\|g\|_{s,(a,b)}} \\
&= \|f\|_{-s,(a,b)}.
\end{aligned}$$

Additionally, as

$$\frac{|F(h)|}{\|h\|_{s,(a,b)}} = \frac{|\langle f, h \rangle_{(a,b)}^0|}{\|h\|_{s,(a,b)}} = \frac{|\sum_{j=0}^{\infty} f_j h_j|}{\|f\|_{-s,(a,b)}} = \frac{|\sum_{j=0}^{\infty} f_j (1+j^2)^{-s} f_j|}{\|f\|_{-s,(a,b)}} = \|f\|_{-s,(a,b)},$$

then it follows that  $\|F\|_{H_{(a,b)}^{-s}} = \|f\|_{-s,(a,b)}$ .

Finally, for  $f \in \mathcal{H}_{(a,b)}^{-s}(\mathbb{I})$  and any  $g \in H_{(a,b)}^s(\mathbb{I})$ ,

$$\langle f, g \rangle_{(a,b)}^0 = \sum_{j=0}^{\infty} f_j g_j = \sum_{j=0}^{\infty} (1+j^2)^{-s/2} f_j (1+j^2)^{s/2} g_j \leq \|f\|_{-s,(a,b)} \|g\|_{s,(a,b)}.$$

Thus,  $f$  defines a bounded linear functional on  $H_{(a,b)}^s(\mathbb{I})$ , i.e.,  $f \in H_{(a,b)}^{-s}(\mathbb{I})$ . ■

With the structure of the  $H_{(a,b)}^s(\mathbb{I})$  spaces it is straight forward to show that  $D$  is a bounded mapping from  $H_{(a,b)}^s(\mathbb{I})$  onto  $H_{(a+1,b+1)}^{s-1}(\mathbb{I})$ .

**Lemma 4.5** *For  $s, a, b \in \mathbb{R}$ ,  $a, b > -1$ . the differential operator  $D$  is a bounded mapping from  $H_{(a,b)}^s(\mathbb{I})$  onto  $H_{(a+1,b+1)}^{s-1}(\mathbb{I})$ .*

**Proof:** Let  $f \in H_{(a,b)}^s(\mathbb{I})$ . Then,  $f(x)$  is expressible as

$$f(x) = \sum_{j=0}^{\infty} c_j \tilde{G}_j^{(a,b)}(x) \quad \text{where} \quad \sum_{j=0}^{\infty} (1+j^2)^s c_j^2 < \infty.$$

Using (2.6),

$$\begin{aligned}
Df(x) &= \sum_{j=1}^{\infty} c_j \frac{1}{\|\tilde{G}_j^{(a,b)}\|} \frac{\Gamma(j+a+b+2)}{\Gamma(j+a+b+1)} G_{j-1}^{(a+1,b+1)}(x) \\
&= \sum_{j=1}^{\infty} c_j \frac{\|G_{j-1}^{(a+1,b+1)}\|}{\|\tilde{G}_j^{(a,b)}\|} (j+a+b+1) \tilde{G}_{j-1}^{(a+1,b+1)}(x).
\end{aligned} \tag{4.25}$$

From (4.12) with  $k = 1$ , we have that there exists a constant  $C > 0$  such that for  $j \geq 1$

$$c_j \frac{|||G_{j-1}^{(a+1, b+1)}|||}{|||G_j^{(a, b)}|||} (j + a + b + 1) \leq C j c_j.$$

Thus,

$$\|Df\|_{s-1, (a+1, b+1)}^2 = \sum_{j=0}^{\infty} (1+j^2)^{s-1} C (j+1)^2 b_{j+1}^2 \leq C \sum_{j=0}^{\infty} (1+j^2)^s c_j^2 = C \|f\|_{s, (a, b)}^2.$$

To establish that the mapping is onto, note that from (4.25) for  $g(x) = \sum_{j=0}^{\infty} d_j \tilde{G}_j^{(a+1, b+1)}(x) \in H_{(a+1, b+1)}^{s-1}(\mathbb{I})$ , the function  $f(x) = \sum_{j=0}^{\infty} d_j \frac{1}{j+a+b+2} \frac{|||G_{j+1}^{(a, b)}|||}{|||G_j^{(a+1, b+1)}|||} \tilde{G}_{j+1}^{(a, b)}(x) \in H_{(a, b)}^s(\mathbb{I})$ , and  $Df(x) = g(x)$ . ■

#### 4.1 Equivalence of the spaces $H_{\rho(a, b)}^s(\mathbb{I})$ and $H_{(a, b)}^s(\mathbb{I})$

In this section we show that the spaces  $H_{\rho(a, b)}^s(\mathbb{I})$  and  $H_{(a, b)}^s(\mathbb{I})$  are equivalent. In the first lemma we show equivalence of the space for  $s \in \mathbb{N}_0$ . Then, in the second lemma we establish equivalence of the spaces for  $s \in \mathbb{R} \setminus \mathbb{N}_0$ .

**Lemma 4.6** *For  $s \in \mathbb{N}_0$  the spaces  $H_{(a, b)}^s(\mathbb{I})$  and  $H_{\rho(a, b)}^s(\mathbb{I})$  coincide, and their corresponding norms are equivalent.*

**Proof:** For  $s = m \in \mathbb{N}$ , using Parseval's equality

$$\begin{aligned} \|v\|_{m, (a, b)}^2 &= \sum_{k=0}^m \|v^{(k)}\|_{L_{\rho(a, b)}^2}^2 = \sum_{k=0}^m \sum_{j=0}^{\infty} |v_j^{(k)}|^2 \\ &= \sum_{k=0}^m \sum_{j=k}^{\infty} |v_{j-k}^{(k)}|^2 \\ &= \sum_{k=0}^m \sum_{j=k}^{\infty} \left( \frac{|||G_{j-k}^{(a+k, b+k)}|||}{|||G_j^{(a, b)}|||} \frac{\Gamma(j+k+a+b+1)}{\Gamma(j+a+b+1)} v_j \right)^2 \quad (\text{using (4.4)}) \\ &\sim \sum_{k=0}^m \sum_{j=k}^{\infty} j^{2k} v_j^2 \quad (\text{using (4.10), and notation } 0^0 = 1) \\ &\sim \sum_{j=0}^{\infty} (1 + j^2 + \dots + j^{2m}) v_j^2. \end{aligned}$$

Noting that

$$(1 + j^2 + \dots + j^{2m}) \leq (1 + j^2)^m \leq \binom{m}{\lfloor m/2 \rfloor} (1 + j^2 + \dots + j^{2m}),$$

where  $\lfloor \cdot \rfloor$  denotes the integer part, we obtain

$$\|v\|_{m,\rho^{(a,b)}}^2 \sim \sum_{j=k}^{\infty} (1+j^2)^m v_j^2 = \|v\|_{m,(a,b)}^2.$$

■

Recall that for  $m, n \in \mathbb{N}$ ,  $m < s < n$ ,  $\theta$  satisfying  $s = (1 - \theta)m + \theta n$ , with

$$K(t, u) := \inf_{v \in H_{\rho^{(a,b)}}^n} \left( \|u - v\|_{m,\rho^{(a,b)}} + \|v\|_{n,\rho^{(a,b)}} \right), \quad (4.26)$$

$$\|u\|_{s,\rho^{(a,b)}} := \|u\|_{[H_{\rho^{(a,b)}}^m, H_{\rho^{(a,b)}}^n]_{\theta,2}} := \left( \int_0^\infty t^{-2\theta} K(t, u)^2 \frac{dt}{t} \right)^{1/2}.$$

**Theorem 4.1** *The spaces  $H_{(a,b)}^s(\mathbb{I})$  and  $H_{\rho^{(a,b)}}^s(\mathbb{I})$  coincide, and their corresponding norms are equivalent.*

**Proof:** (This part of the proof is similar to an argument used in [33]. See also [6, Theorem 2.1].) Let  $m = n - 1$ ,  $n$ ,  $s$  and  $\theta$  be as described above, and assume that  $u \in H_{\rho^{(a,b)}}^{n-1}(\mathbb{I})$ . Using  $\frac{1}{\sqrt{2}}(c + d) \leq (c^2 + d^2)^{1/2} \leq c + d$ , and  $H_{\rho^{(a,b)}}^j(\mathbb{I}) = H_{(a,b)}^j(\mathbb{I})$  for  $j \in \mathbb{N}_0$ , we have that

$$K(t, u) \sim \tilde{K}(t, u) = \inf_{v \in H_{(a,b)}^n} \left( \|u - v\|_{n-1,(a,b)}^2 + t^2 \|v\|_{n,(a,b)}^2 \right)^{1/2}.$$

For  $u = \sum_{k=0}^\infty u_k \tilde{G}_k^{(a,b)}(x)$  and  $v = \sum_{k=0}^\infty v_k \tilde{G}_k^{(a,b)}(x)$ ,

$$\|u - v\|_{n-1,(a,b)}^2 = \sum_{k=0}^\infty (1 + k^2)^{n-1} (u_k - v_k)^2, \quad \text{and} \quad \|v\|_{n,(a,b)}^2 = \sum_{k=0}^\infty (1 + k^2)^n v_k^2.$$

Then,

$$\begin{aligned} \tilde{K}(t, u)^2 &= \inf_{\{(1+k^2)^{n/2}v_k\}_{k=0}^\infty \in l^2} \sum_{k=0}^\infty (1 + k^2)^{n-1} (u_k - v_k)^2 + t^2 (1 + k^2)^n v_k^2 \\ &\geq \sum_{k=0}^\infty \inf_{v_k \in \mathbb{R}} \left( (1 + k^2)^{n-1} (u_k - v_k)^2 + t^2 (1 + k^2)^n v_k^2 \right). \end{aligned} \quad (4.27)$$

Each term in the summation in (4.27) is minimized for

$$v_k = \frac{u_k}{1 + t^2(1 + k^2)}. \quad (4.28)$$

For these  $v_k$  we have that

$$\begin{aligned} \sum_{k=1}^\infty (1 + k^2)^n v_k^2 &= \sum_{k=1}^\infty \frac{u_k^2 (1 + k^2)^n}{(1 + t^2(1 + k^2))^2} \\ &= t^{-4} \sum_{k=1}^\infty \frac{u_k^2 t^4 (1 + k^2)^n}{(1 + t^2(1 + k^2))^2} \leq t^{-4} \sum_{k=1}^\infty u_k^2 (1 + k^2)^{n-2} \\ &\leq t^{-4} \|u\|_{n-1,(a,b)}^2, \end{aligned}$$

which implies that  $(1 + k^2)^{n/2} v_k \in l^2$ . Hence,

$$\tilde{K}(t, u)^2 = \sum_{k=0}^{\infty} \frac{t^2(1 + k^2)^n}{(1 + t^2(1 + k^2))} u_k^2.$$

Then,

$$\begin{aligned} \|u\|_{s, \rho^{(a,b)}} &\sim \int_0^\infty t^{-2\theta} \tilde{K}(t, u)^2 \frac{dt}{t} = \int_0^\infty \sum_{k=0}^{\infty} t^{-2\theta} \frac{t^2(1 + k^2)^n}{(1 + t^2(1 + k^2))} u_k^2 \frac{dt}{t} \\ &= \sum_{k=0}^{\infty} \int_0^\infty t^{-2\theta} \frac{t^2(1 + k^2)^n}{(1 + t^2(1 + k^2))} u_k^2 \frac{dt}{t}. \end{aligned}$$

Letting  $\tau = (1 + k^2)^{1/2} t \Leftrightarrow t = (1 + k^2)^{-1/2} \tau \Rightarrow dt = (1 + k^2)^{-1/2} d\tau$ ,

$$\begin{aligned} \|u\|_{s, \rho^{(a,b)}}^2 &\sim \sum_{k=0}^{\infty} \int_0^\infty \tau^{-2\theta} (1 + k^2)^\theta \frac{\tau^2 (1 + k^2)^{n-1}}{(1 + \tau^2)} u_k^2 \frac{d\tau}{\tau} \\ &= \sum_{k=0}^{\infty} (1 + k^2)^{n-1+\theta} u_k^2 \int_0^\infty \frac{\tau^{1-2\theta}}{1 + \tau^2} d\tau \\ &= C_\theta^2 \sum_{k=0}^{\infty} (1 + k^2)^s u_k^2 = C_\theta^2 \|u\|_{s, (a,b)}^2, \end{aligned} \tag{4.29}$$

where  $C_\theta := \left( \int_0^\infty \frac{\tau^{1-2\theta}}{1 + \tau^2} d\tau \right)^{1/2}$ .

Thus, combining Lemma 4.6 and (4.29) we have for  $s \geq 0$  the spaces  $H_{(a,b)}^s(\mathbf{I})$  and  $H_{\rho^{(a,b)}}^s(\mathbf{I})$  coincide, and their corresponding norms are equivalent. It also then follows that for  $s < 0$  the corresponding (dual) spaces are equivalent. ■

We have the following straight forward lemma.

**Lemma 4.7** *For  $-1 < a_1 < a$ ,  $-1 < b_1 < b$ ,  $s \geq 0$ , we have that  $H_{(a_1, b_1)}^s(\mathbf{I}) \subset H_{(a,b)}^s(\mathbf{I})$ . Additionally, for  $s > 0$ ,  $H_{(a,b)}^{-s}(\mathbf{I}) \subset H_{(a_1, b_1)}^{-s}(\mathbf{I})$ .*

**Proof:**

For  $s \in \mathbb{N}_0$  the fact that for  $x \in \mathbf{I}$ ,  $(1 - x)^{a_1} x^{b_1} \leq (1 - x)^a x^b$ , together with definition (3.1) and the equivalence of  $H_{\rho^{(a,b)}}^s(\mathbf{I})$  and  $H_{(a,b)}^s(\mathbf{I})$  establishes that  $H_{(a_1, b_1)}^s(\mathbf{I}) \subset H_{(a,b)}^s(\mathbf{I})$ .

Next, for  $m = n - 1$ ,  $n$ ,  $s$  and  $\theta$  as defined in (4.26), consider

$$\begin{aligned} K_{(a_1, b_1)}(t, u) &:= \inf_{v \in H_{(a_1, b_1)}^n} (\|u - v\|_{n-1, (a_1, b_1)} + t \|v\|_{n, (a_1, b_1)}) \\ &\geq \inf_{v \in H_{(a_1, b_1)}^n} (\|u - v\|_{n-1, (a,b)} + t \|v\|_{n, (a,b)}) \\ &\geq \inf_{v \in H_{(a,b)}^n} (\|u - v\|_{n-1, (a,b)} + t \|v\|_{n, (a,b)}) \quad (\text{as } H_{(a_1, b_1)}^n(\mathbf{I}) \subset H_{(a,b)}^n(\mathbf{I})) \\ &:= K_{(a,b)}(t, u). \end{aligned}$$

Hence  $\|u\|_{s,(a,b)} \leq \|u\|_{s,(a_1,b_1)}$ , from which it follows that  $H_{(a_1,b_1)}^s(\mathbf{I}) \subset H_{(a,b)}^s(\mathbf{I})$ .

That  $H_{(a,b)}^{-s}(\mathbf{I}) \subset H_{(a_1,b_1)}^{-s}(\mathbf{I})$  follows from the definition of the dual space and that  $H_{(a_1,b_1)}^s(\mathbf{I}) \subset H_{(a,b)}^s(\mathbf{I})$  for  $s > 0$ . ■

We further illustrate the structure of the  $H_{(a,b)}^s(\mathbf{I})$  spaces with the following lemma, whose proof is motivated by the proof of Theorem 2.5 in [6].

**Lemma 4.8** *Let  $u(x) = x^\mu$ . Then,  $u \in H_{(a,b)}^s(\mathbf{I})$  for  $s < 2\mu + b + 1$ .*

**Proof:** Let  $\chi(x) \in C^\infty[0, \infty)$  denote the cutoff function satisfying

$$\chi(x) = \begin{cases} 1 & \text{for } 0 < x \leq 1/4 \\ 0 & \text{for } x \geq 3/4 \end{cases},$$

and let  $\chi_\delta(x) := \chi(\frac{x}{\delta})$ , for  $\delta > 0$ . Note that

$$\chi_\delta(x) = \begin{cases} 1 & \text{for } 0 < x \leq \delta/4 \\ 0 & \text{for } x \geq 3\delta/4 \end{cases}, \quad \text{and} \quad \frac{d^m}{dx^m} \chi_\delta(x) = \begin{cases} 0 & \text{for } 0 < x < \delta/4 \\ 0 & \text{for } x > 3\delta/4 \end{cases} \quad \text{for } m \in \mathbb{N}.$$

For  $\delta$  to be determined, let  $u = v + w$  where  $v(x) = \chi_\delta(x) u(x)$  and  $w(x) = (1 - \chi_\delta(x)) u(x)$ . We have that

$$\left| \frac{d^m v(x)}{dx^m} \right| \leq C \sum_{j=0}^m \delta^{-(m-j)} x^{\mu-j}, \quad \text{and is zero for } x > 3\delta/4.$$

Thus,

$$\begin{aligned} \int_{\mathbf{I}} (1-x)^{a+m} x^{b+m} \left| \frac{d^m v(x)}{dx^m} \right|^2 dx &\leq C \int_0^{3\delta/4} \sum_{j=0}^m \delta^{-2(m-j)} x^{2\mu-2j+b+m} dx \\ &\leq C \sum_{j=0}^m \delta^{-2(m-j)} \delta^{2\mu-2j+b+m+1}, \quad \text{provided } 2\mu - 2j + b + m > -1, \\ &\leq C \delta^{2\mu+b+1-m}, \end{aligned}$$

which implies that, for  $m < 2\mu + b + 1$ ,  $v \in H_{\rho(a,b)}^m(\mathbf{I})$  and

$$\|v\|_{m, \rho(a,b)}^2 \leq C \delta^{2\mu+b+1-m}. \quad (4.30)$$

Next, consider  $w(x)$ .

$$\left| \frac{d^m w(x)}{dx^m} \right| \leq C \left( (1 - \chi_\delta(x)) x^{\mu-m} + \sum_{j=0}^{m-1} x^{\mu-j} \frac{d^{m-j}}{dx^{m-j}} (1 - \chi_\delta(x)) \right). \quad (4.31)$$



The first term on the RHS of (4.31) vanishes for  $x < \delta/4$ , and the second term vanishes for  $x < \delta/4$  and  $x > 3\delta/4$ . Using this,

$$\begin{aligned} \int_I (1-x)^{a+m} x^{b+m} \left| \frac{d^m w(x)}{dx^m} \right|^2 dx &\leq C \left( \int_{\delta/4}^1 x^{b+m} x^{2\mu-2m} dx + \int_{\delta/4}^{3\delta/4} x^{b+m} \sum_{j=0}^{m-1} x^{2\mu-2j} \delta^{-2(m-j)} dx \right) \\ &\leq C \left( \int_{\delta/4}^1 x^{2\mu+b-m} dx + \sum_{j=0}^{m-1} \delta^{-2m+2j} \int_{\delta/4}^{3\delta/4} x^{2\mu+b+m-2j} dx \right) \\ &\leq \begin{cases} C (1 + \delta^{2\mu+b-m+1}) & \text{if } 2\mu + b - m \neq -1 \\ C (1 + |\log \delta|) & \text{if } 2\mu + b - m = -1 \end{cases} . \end{aligned} \quad (4.32)$$

Hence, for  $n > 2\mu + b + 1$ ,  $w \in H_{\rho(a,b)}^n(I)$  and

$$\|w\|_{n,\rho(a,b)}^2 \leq C \delta^{2\mu+b+1-n} . \quad (4.33)$$

(**Remark:** For  $n > 2\mu + b + 1$  the exponent of  $\delta$  in (4.33) is negative, so the '1' term in (4.32) is bounded by the  $\delta$  term.)

We have from (4.30) and (4.33)

$$K(t, u) = \inf_{u=u_1+u_2} \left( \|u_1\|_{H_{(a,b)}^m} + t \|u_2\|_{H_{(a,b)}^n} \right) \quad (4.34)$$

$$\begin{aligned} &\leq \|v\|_{m,\rho(a,b)} + t \|w\|_{n,\rho(a,b)} \\ &\leq C \left( \delta^{(2\mu+b+1-m)/2} + t \delta^{(2\mu+b+1-n)/2} \right) . \end{aligned} \quad (4.35)$$

Setting  $\delta = t^{2/(n-m)}$  leads to  $K(t, u) \leq C t^{(2\mu+b+1-m)/(n-m)}$ .

Recall that

$$\|u\|_{[H_{(a,b)}^m, H_{(a,b)}^n]_{\theta,2}}^2 = \int_0^\infty t^{-2\theta} (K(t, u))^2 \frac{dt}{t} . \quad (4.36)$$

The larger the value of  $\theta$  ( $0 < \theta < 1$ ) in (4.36) such that the integral is finite, the “nicer” (i.e., more regular) is the function  $u$ . Hence from (4.36), we are interested in the integrand about  $t = 0$ . We have trivially that for  $u_1 = u$ ,  $u_2 = 0$  in (4.34) that  $K(t, u) \leq \|u\|_{H_{(a,b)}^m} \leq C$ . Hence it follows that

$$K(t, u) \leq \begin{cases} C t^{(2\mu+b+1-m)/(n-m)} & \text{for } 0 < t < 1 \\ C & \text{for } t \geq 1 \end{cases} .$$

Using (4.36),

$$\begin{aligned} \|u\|_{[H_{(a,b)}^m, H_{(a,b)}^n]_{\theta,2}}^2 &\leq \int_0^1 C t^{-2\theta-1+2(2\mu+b+1-m)/(n-m)} dt + \int_1^\infty C t^{-2\theta-1} dt < \infty , \\ &\text{if } \theta < (2\mu + b + 1 - m)/(n - m) . \end{aligned} \quad (4.37)$$

For  $s = (1 - \theta)m + \theta n = m + \theta(n - m)$ , then  $s < 2\mu + b + 1$  using (4.37) .

Hence we can conclude that  $u(x) = x^\mu \in H_{(a,b)}^s(I)$  for  $s < 2\mu + b + 1$ . ■

We now present a result which connects the weighted Sobolev spaces with the spaces of continuous functions.

**Theorem 4.2** (See [3, Theorem 4.14]) Let  $a, b > -1$ ,  $k \in \mathbb{N}_0$  and  $s > k + 1 + \max\{a + k, b + k, -1/2\}$ . Then we have a continuous embedding  $H_{(a,b)}^s(\mathbf{I}) \subset C^k(\mathbf{I})$  of  $H_{(a,b)}^s(\mathbf{I})$  into the Banach space  $C^k(\mathbf{I})$  with the norm  $\|v\|_{C^k} = \|v\|_\infty + \|v^{(k)}\|_\infty$ .

**Proof:** We begin by establishing that  $v(x) \in C(\mathbf{I})$ . From the representation (4.1), consider the sequence of partial sums  $\{v_n(x)\}_{n=0}^\infty$ , where, using (2.3),

$$v_n(x) = \sum_{j=0}^n v_j \tilde{G}_j^{(a,b)}(x) = \sum_{j=0}^n v_j \left( (2j + a + b + 1) \frac{\Gamma(j+1) \Gamma(j+a+b+1)}{\Gamma(j+a+1) \Gamma(j+b+1)} \right)^{1/2} G_j^{(a,b)}(x).$$

From [35, Theorem 7.32.1], for  $x \in \mathbf{I}$ ,  $|G_n^{(a,b)}(x)| \sim n^q$ , where  $q = \max\{a, b, -1/2\}$ . Using this, together with (4.19), for  $n > m$

$$\begin{aligned} |v_n(x) - v_m(x)| &= \left| \sum_{j=m+1}^n v_j \left( (2j + a + b + 1) \frac{\Gamma(j+1) \Gamma(j+a+b+1)}{\Gamma(j+a+1) \Gamma(j+b+1)} \right)^{1/2} G_j^{(a,b)}(x) \right| \\ &\leq C \sum_{j=m+1}^n |v_j| j^{1/2} j^{\max\{a, b, -\frac{1}{2}\}} \\ &= C \sum_{j=m+1}^n (1 + j^2)^{-s/2} j^{\frac{1}{2} + \max\{a, b, -\frac{1}{2}\}} (1 + j^2)^{s/2} |v_j| \\ &\leq C \left( \sum_{j=m+1}^n (1 + j^2)^{-s} j^{1 + 2 \max\{a, b, -\frac{1}{2}\}} \right)^{1/2} \left( \sum_{j=m+1}^n (1 + j^2)^s v_j^2 \right)^{1/2} \\ &\leq C \left( \sum_{j=m+1}^n j^{1 - 2s + 2 \max\{a, b, -\frac{1}{2}\}} \right)^{1/2} \|v\|_{s, (a,b)}. \end{aligned} \quad (4.38)$$

Thus, for  $1 - 2s + 2 \max\{a, b, -\frac{1}{2}\} < -1$ , i.e., for  $s > 1 + \max\{a, b, -\frac{1}{2}\}$ ,  $\{v_n(x)\}_{n=0}^\infty$  converges uniformly on  $\mathbf{I}$ . As the limit function of a sequence of uniformly convergent, continuous functions is also continuous, it follows that  $v(x) \in C(\mathbf{I})$ .

To obtain the continuity of  $v^{(l)}(x)$ , for  $1 \leq l \leq k$ , we proceed in a similar manner. Using (2.6),  $m \geq l$

$$|v_n^{(l)}(x) - v_m^{(l)}(x)| = \left| \sum_{j=m+1}^n v_j \left( (2j + a + b + 1) \frac{\Gamma(j+1) \Gamma(j+a+b+1)}{\Gamma(j+a+1) \Gamma(j+b+1)} \right)^{1/2} \frac{\Gamma(j+l+a+b+1)}{\Gamma(j+a+b+1)} G_{j-l}^{(a+l, b+l)}(x) \right|.$$

Using

$$\frac{\Gamma(j+l+a+b+1)}{\Gamma(j+a+b+1)} \sim j^l \quad \text{for } j \text{ large,}$$

$$\begin{aligned}
|v_n^{(l)}(x) - v_m^{(l)}(x)| &\leq C \sum_{j=l}^n |v_j| j^{1/2} j^l (j-l)^{\max\{a+l, b+l, -1/2\}} \\
&= C \left| \sum_{j=m+1}^n (1+j^2)^{-s/2} j^{l+\frac{1}{2}+\max\{a+l, b+l, -1/2\}} (1+j^2)^{s/2} |v_j| \right| \\
&\leq C \left( \sum_{j=m+1}^n j^{2l+1-2s+2\max\{a+l, b+l, -1/2\}} \right)^{1/2} \|v\|_{s,(a,b)}. \tag{4.39}
\end{aligned}$$

Thus for  $2l+1-2s+2\max\{a+l, b+l, -1/2\} < -1$ , i.e., for  $s > l+1+\max\{a+l, b+l, -1/2\}$ ,  $\{v_n^{(l)}(x)\}_{n=0}^\infty$  converges uniformly on  $I$ . Furthermore, as  $\{v_n(x)\}_{n=0}^\infty$  converges to  $v(x)$ , then  $\{v_n^{(l)}(x)\}_{n=0}^\infty$  converges to  $v^{(l)}(x) \in C(I)$  (see [32, Theorem 7.17]).

A straight forward modification of the argument used in (4.38) and (4.39) can be used to establish that, for  $0 \leq l \leq k$ ,  $\|v^{(l)}\|_\infty \leq C \|v\|_{H_{(a,b)}^s}$ , where the precise value for  $C$  depends upon  $l$ . ■

## 5 Regularity of the solution to the fractional diffusion equation

Now we can give sharp regularity results for the fractional diffusion equation (1.1),(1.2).

**Theorem 5.1** *For  $\mathcal{L}_r^\alpha(\cdot)$  defined for  $1 < \alpha < 2$ ,  $0 \leq r \leq 1$ , let  $\beta$  be determined by **Condition A**. Then the mapping  $\mathcal{L}_r^\alpha(\cdot) : \rho^{(\alpha-\beta, \beta)}(x) \otimes H_{(\alpha-\beta, \beta)}^{s+\alpha}(I) \rightarrow H_{(\beta, \alpha-\beta)}^s(I)$  is bijective, continuous, and has a continuous inverse.*

**Proof:** From [13, 23],

$$\mathcal{L}_r^\alpha \left( \rho^{(\alpha-\beta, \beta)} \tilde{G}_k^{(\alpha-\beta, \beta)} \right) (x) = \lambda_k \tilde{G}_k^{(\beta, \alpha-\beta)}(x), \quad \text{where } \lambda_k = -c_*^* \frac{\Gamma(k+1+\alpha)}{\Gamma(k+1)}, \quad k = 0, 1, 2, \dots,$$

and  $c_*^*$  given by (2.12).

Using Stirling's formula,

$$\begin{aligned}
\lambda_k^2 &= (c_*^*)^2 \frac{\Gamma(k+1+\alpha)^2}{\Gamma(k+1)^2} \sim (k+1)^{2\alpha} \quad \text{as } k \rightarrow \infty, \\
\text{i.e., } c(1+k^2)^\alpha &\leq \lambda_k^2 \leq C(1+k^2)^\alpha. \tag{5.1}
\end{aligned}$$

Let  $\phi(x) = \sum_{k=0}^\infty \phi_k \tilde{G}_k^{(\alpha-\beta, \beta)}(x) \in H_{(\alpha-\beta, \beta)}^{s+\alpha}(I)$ , and  $\phi_N(x) = \sum_{k=0}^N \phi_k \tilde{G}_k^{(\alpha-\beta, \beta)}(x)$ . Then  $\{\phi_n\}_{n=1}^\infty$  is a Cauchy sequence in  $H_{(\alpha-\beta, \beta)}^{s+\alpha}(I)$ , with  $\lim_{n \rightarrow \infty} \phi_n = \phi$ .

Consider,

$$f_N(x) = \mathcal{L}_\alpha^r \left( \rho^{(\alpha-\beta, \beta)} \phi_N \right) (x) = \sum_{k=0}^N \lambda_k \phi_k \tilde{G}_k^{(\beta, \alpha-\beta)}(x) \in H_{(\beta, \alpha-\beta)}^s(I).$$

Then using (5.1),

$$\begin{aligned}\|f_N - f_M\|_{s,(\beta, \alpha-\beta)}^2 &= \sum_{k=M}^N (1+k^2)^s (\lambda_k \phi_k)^2 \leq C \sum_{k=M}^N (1+k^2)^s (1+k^2)^\alpha \phi_k^2 \\ &= C \|\phi_N - \phi_M\|_{s+\alpha, (\alpha-\beta, \beta)}^2.\end{aligned}$$

Thus  $\{f_n\}_{n=1}^\infty$  is a Cauchy sequence in  $H_{(\beta, \alpha-\beta)}^s(\mathbf{I})$ . It then follows that  $f := \lim_{n \rightarrow \infty} f_n$  satisfies

$$\begin{aligned}f &= \mathcal{L}_r^\alpha \left( \rho^{(\alpha-\beta, \beta)} \phi \right) (x) = \mathcal{L}_r^\alpha \left( \rho^{(\alpha-\beta, \beta)} \sum_{k=0}^\infty \phi_k \tilde{G}_k^{(\alpha-\beta, \beta)} \right) (x) \\ &= \sum_{k=0}^\infty \phi_k \mathcal{L}_r^\alpha \left( \rho^{(\alpha-\beta, \beta)} \tilde{G}_k^{(\alpha-\beta, \beta)} \right) (x) \\ &= \sum_{k=0}^\infty \lambda_k \phi_k \tilde{G}_k^{(\beta, \alpha-\beta)} (x).\end{aligned}$$

Also, using (5.1),

$$\|f\|_{s,(\beta, \alpha-\beta)}^2 \leq \sum_{k=0}^\infty (1+k^2)^s \lambda_k^2 \phi_k^2 \leq C \sum_{k=0}^\infty (1+k^2)^{s+\alpha} \phi_k^2 = C \|\phi\|_{s+\alpha, (\alpha-\beta, \beta)}^2. \quad (5.2)$$

Hence  $\mathcal{L}_r^\alpha(\cdot)$  is a one-to-one, continuous mapping from  $\rho^{(\alpha-\beta, \beta)}(x) \otimes H_{(\alpha-\beta, \beta)}^{s+\alpha}(\mathbf{I})$  into  $H_{(\beta, \alpha-\beta)}^s(\mathbf{I})$ .

Next, for  $f(x) = \sum_{k=0}^\infty f_k \tilde{G}_k^{(\beta, \alpha-\beta)}(x) \in H_{(\beta, \alpha-\beta)}^s(\mathbf{I})$ , let  $\phi(x) = \sum_{k=0}^\infty \frac{1}{\lambda_k} f_k \tilde{G}_k^{(\alpha-\beta, \beta)}(x)$ . Note that  $\mathcal{L}_r^\alpha(\rho^{(\alpha-\beta, \beta)} \phi)(x) = f(x)$ , and using a similar argument to that in (5.2),  $\|\phi\|_{s+\alpha, (\alpha-\beta, \beta)} \leq C \|f\|_{s, (\beta, \alpha-\beta)}$ , from which the stated result then follows. ■

Using this theorem we obtain the following sharp regularity result for (1.1),(1.2).

**Corollary 5.1** *For  $f \in H_{(\beta, \alpha-\beta)}^s(\mathbf{I})$  there exists a unique solution  $u$  to (1.1),(1.2), which can be expressed as  $u(x) = \rho^{(\alpha-\beta, \beta)}(x)\phi(x)$ , where  $\phi(x) \in H_{(\alpha-\beta, \beta)}^{s+\alpha}(\mathbf{I})$ , with  $\|\phi\|_{s+\alpha, (\alpha-\beta, \beta)} \leq C \|f\|_{s, (\beta, \alpha-\beta)}$ .* ■

Combining Theorem 4.2 and Corollary 5.1 we obtain a sufficient condition for the solution of (1.1),(1.2) to be continuous on  $\mathbf{I}$ .

**Corollary 5.2** *Let  $1 < \alpha < 2$ ,  $0 \leq r \leq 1$ , and  $\beta$  be determined by **Condition A**. Then for  $f \in H_{(\beta, \alpha-\beta)}^s(\mathbf{I})$  where*

$$s > (1 - \alpha) + \max\{\alpha - \beta, \beta\}, \quad (5.3)$$

*the solution of (1.1),(1.2) is continuous on  $\mathbf{I}$ .*

■

**Note:** For  $r = 0$  ( $r = 1$ ) Corollary 5.2 implies for  $f \in H_{(1, \alpha-1)}^s(\mathbf{I})$  ( $f \in H_{(\alpha-1, 1)}^s(\mathbf{I})$ ), where  $s > 2 - \alpha$ , that  $u \in C(\mathbf{I})$ .

For  $r = 1/2$  Corollary 5.2 implies for  $f \in H_{(\alpha/2, \alpha/2)}^s(\mathbf{I})$ , where  $s > 1 - \alpha/2$ , that  $u \in C(\mathbf{I})$ .

## 6 Weighted Sobolev Space defined using a Sobolev-Slobodeckij type semi-norm

To obtain the regularity results for (1.3), (1.4) in the following section we use the regularity result from (1.1), (1.2) and a boot-strapping argument. To apply the boot-strapping argument we need a precise estimate for  $t$  such that if  $\phi(x) \in H_{(\alpha-\beta, \beta)}^r(\mathbf{I})$ , then  $(1-x)^{\alpha-\beta} x^\beta \phi(x) \in H_{(\beta, \alpha-\beta)}^t(\mathbf{I})$ .

In Sections 3 and 4 we gave two characterizations of the weighted Sobolev space  $H_{(a,b)}^s(\mathbf{I})$ . Neither of these definitions are easily to apply in order to determine the value for  $t$  in order that  $(1-x)^{\alpha-\beta} x^\beta \phi(x) \in H_{(\beta, \alpha-\beta)}^t(\mathbf{I})$ . Therefore, following the work of Bernardi, Dauge, and Maday [9], and Fdez-Manin, and Munoz-Sola [15] we introduce a further characterization of  $H_{(a,b)}^s(\mathbf{I})$ .

**Remark:** The work in [9, 15] is presented for a single weighted space, i.e.,  $H_{(a,a)}^s(\mathbf{I})$ . However, it extends in a straight forward manner to  $H_{(a,b)}^s(\mathbf{I})$  spaces.

### 6.1 Equivalent definition for $H_{(a,b)}^s(\mathbf{I})$

For  $s > 0$ , let  $s = \lfloor s \rfloor + r$ , where  $0 < r < 1$ . Let the semi-norm,  $|\cdot|_{H_{(a,b)}^s(\mathbf{I})}$ , and norm,  $\|\cdot\|_{H_{(a,b)}^s(\mathbf{I})}$  be defined as

$$|f|_{H_{(a,b)}^s(\mathbf{I})}^2 := \iint_{\tilde{\Lambda}} (1-x)^{a+s} x^{b+s} \frac{|D^{\lfloor s \rfloor} f(x) - D^{\lfloor s \rfloor} f(y)|^2}{|x-y|^{1+2(s-\lfloor s \rfloor)}} dy dx$$

$$\text{and } \|f\|_{H_{(a,b)}^s(\mathbf{I})}^2 := \begin{cases} \sum_{j=0}^s \|D^j f\|_{L_{(a+j, b+j)}^2(\mathbf{I})}^2, & \text{for } s \in \mathbb{N}_0 \\ \sum_{j=0}^{\lfloor s \rfloor} \|D^j f\|_{L_{(a+j, b+j)}^2(\mathbf{I})}^2 + |f|_{H_{(a,b)}^s(\mathbf{I})}^2, & \text{for } s \in \mathbb{R}^+ \setminus \mathbb{N}_0 \end{cases},$$

where (see Figure 6.1)

$$\tilde{\Lambda} := \left\{ (x, y) : \frac{2}{3}x < y < \frac{3}{2}x, 0 < x < \frac{1}{2} \right\} \cup \left\{ (x, y) : \frac{3}{2}x - \frac{1}{2} < y < \frac{2}{3}x + \frac{1}{3}, 1/2 \leq x < 1 \right\}. \quad (6.1)$$

**Lemma 6.1** [15] *Let  $s \geq 0$  with  $s \neq 1 + a$  if  $a \in (-1, 0)$  and  $s \neq 1 + b$  if  $b \in (-1, 0)$ . Then,  $\{f : f \text{ is measurable and } \|f\|_{H_{(a,b)}^s(\mathbf{I})} < \infty\} = H_{(a,b)}^s(\mathbf{I})$ .*

**Proof:** From [15] it follows that  $\{f : f \text{ is measurable and } \|f\|_{H_{(a,b)}^s(\mathbf{I})} < \infty\} = H_{\rho(a,b)}^s(\mathbf{I})$ . The equality of the spaces  $H_{\rho(a,b)}^s(\mathbf{I})$  and  $H_{(a,b)}^s(\mathbf{I})$  was established in Section 4.

■




Figure 6.1: Domain  $\tilde{\Lambda}$ .




Figure 6.2: Domain  $\Lambda^* = \Lambda \cup \Lambda_1$ .

The  $H_{(a,b)}^s(\mathbf{I})$  space a function  $f$  lies in is determined by its behavior at: (i) the left endpoint ( $x = 0$ ), (ii) the right endpoint ( $x = 1$ ), and (iii) away from the endpoints. In order to separate the consideration of the endpoint behaviors, following [9] we introduce the following function space  $H_{(\gamma)}^s(\mathbf{J})$ . Let  $\mathbf{J} := (0, 3/4)$ , and

$$\begin{aligned} \Lambda^* &:= \left\{ (x, y) : \frac{2}{3}x < y < \frac{3}{2}x, 0 < x < \frac{1}{2} \right\} \cup \left\{ (x, y) : \frac{3}{2}x - \frac{1}{2} < y < \frac{2}{3}x + \frac{1}{3}, 1/2 \leq x < 3/4 \right\} \\ &:= \Lambda \cup \Lambda_1 \quad (\text{see Figure 6.2}). \end{aligned}$$

Introduce the semi-norm and norm

$$|f|_{H_{(\gamma)}^s(\mathbf{J})}^2 := \iint_{\Lambda} x^{\gamma+s} \frac{|D^{\lfloor s \rfloor} f(x) - D^{\lfloor s \rfloor} f(y)|^2}{|x - y|^{1+2(s-\lfloor s \rfloor)}} dy dx + \iint_{\Lambda_1} x^{\gamma+s} \frac{|D^{\lfloor s \rfloor} f(x) - D^{\lfloor s \rfloor} f(y)|^2}{|x - y|^{1+2(s-\lfloor s \rfloor)}} dy dx \quad (6.2)$$

$$:= |f|_{H_{(\gamma)}^s(\Lambda)}^2 + |f|_{H_{(\gamma)}^s(\Lambda_1)}^2, \quad (6.3)$$

$$(6.4)$$

$$\text{and } \|f\|_{H_{(\gamma)}^s(\mathbf{J})}^2 := \begin{cases} \sum_{j=0}^s \|D^j f\|_{L_{(\gamma+j)}^2(\mathbf{J})}^2, & \text{for } s \in \mathbb{N}_0 \\ \sum_{j=0}^{\lfloor s \rfloor} \|D^j f\|_{L_{(\gamma+j)}^2(\mathbf{J})}^2 + |f|_{H_{(\gamma)}^s(\mathbf{J})}^2, & \text{for } s \in \mathbb{R}^+ \setminus \mathbb{N}_0 \end{cases}, \quad (6.5)$$

$$(6.6)$$

$$\text{where } \|g\|_{L_{(\gamma)}^2(\mathbf{J})}^2 := \int_{\mathbf{J}} x^{\gamma} g^2(x) dx.$$

Then,  $H_{(\gamma)}^s(\mathbf{J}) := \{f : f \text{ is measurable and } \|f\|_{H_{(\gamma)}^s(\mathbf{J})} < \infty\}$ .

**Note:** A function  $f(x)$  is in  $H_{(a,b)}^s(\mathbf{I})$  if and only if  $f(\frac{3}{4}x) \in H_{(b)}^s(\mathbf{J})$  and  $f(\frac{3}{4}(1-x)) \in H_{(a)}^s(\mathbf{J})$ .

The following lemma and discussion allows us to further focus our analysis for determining the regularity of  $(1-x)^{\alpha-\beta} x^{\beta} \phi(x)$ .

**Lemma 6.2** *Let  $s \geq 0$ ,  $\psi \in H_{(\gamma)}^s(\mathbf{J})$ , and  $g \in C^{\lceil s \rceil}(\mathbf{J})$ . Then*

$$\|g\psi\|_{H_{(\gamma)}^s(\mathbf{J})}^2 \lesssim \|g\|_{C^{\lceil s \rceil}(\mathbf{J})}^2 \|\psi\|_{H_{(\gamma)}^s(\mathbf{J})}^2.$$

**Proof:** For  $s = 0$ ,

$$\|g\psi\|_{H_{(\gamma)}^0(\mathbf{J})}^2 = \|g\psi\|_{L_{(\gamma)}^2(\mathbf{J})}^2 \leq \|g\|_{L^\infty(\mathbf{J})}^2 \|\psi\|_{L_{(\gamma)}^2(\mathbf{J})}^2. \quad (6.7)$$

Then, for  $s = 1$ ,

$$\begin{aligned} \|g\psi\|_{H_{(\gamma)}^1(\mathbf{J})}^2 &\leq \|g\psi\|_{L_{(\gamma)}^2(\mathbf{J})}^2 + \|D(g\psi)\|_{L_{(\gamma+1)}^2(\mathbf{J})}^2 \\ &\lesssim \|g\|_{L^\infty(\mathbf{J})}^2 \|\psi\|_{L_{(\gamma)}^2(\mathbf{J})}^2 + \|\psi Dg\|_{L_{(\gamma+1)}^2(\mathbf{J})}^2 + \|g D\psi\|_{L_{(\gamma+1)}^2(\mathbf{J})}^2 \\ &\lesssim \|g\|_{L^\infty(\mathbf{J})}^2 \|\psi\|_{L_{(\gamma)}^2(\mathbf{J})}^2 + \|Dg\|_{L^\infty(\mathbf{J})}^2 \|\psi\|_{L_{(\gamma)}^2(\mathbf{J})}^2 + \|g\|_{L^\infty(\mathbf{J})}^2 \|D\psi\|_{L_{(\gamma+1)}^2(\mathbf{J})}^2 \\ &\lesssim \|g\|_{C^1(\mathbf{J})}^2 \|\psi\|_{H_{(\gamma)}^1(\mathbf{J})}^2. \end{aligned} \quad (6.8)$$

Next, for  $0 < s < 1$ , consider the mapping  $\mathcal{F} : H_{(\gamma)}^s(\mathbf{J}) \rightarrow H_{(\gamma)}^s(\mathbf{J})$ , defined by  $\mathcal{F}(\psi) := g\psi$ . From (6.7) and (6.8) we have that  $\mathcal{F}$  is a bounded mapping for  $s = 0$  and  $s = 1$ . As  $H_{(\gamma)}^s(\mathbf{J})$  is a family of interpolation spaces, it follows that  $\mathcal{F}$  is a bounded mapping for  $0 < s < 1$ , with  $\|\mathcal{F}\| \lesssim \|g\|_{C^1(\mathbf{J})}$ .

For  $s > 1$  the above argument extends in a straight forward manner. ■

Consider  $h(x) = g(x)\psi(x)$ , where  $\psi(x) \in H_{(\mu)}^s(\mathbf{J})$  and  $g(x) = \begin{cases} x^\alpha, & \frac{1}{2} \leq x \leq \frac{3}{4} \\ g_{ext}(x), & 0 \leq x \leq \frac{1}{2} \end{cases}$ , for  $g_{ext}(x)$  a  $C^{\lceil s \rceil}(0, \frac{1}{2})$  extension of  $x^\alpha$  satisfying  $\|g\|_{C^{\lceil s \rceil}(\mathbf{J})} \leq \|x^\alpha\|_{C^{\lceil s \rceil}[\frac{1}{2}, \frac{3}{4}]}$ .

Note that for  $t \leq s$ ,  $t \notin \mathbb{N}$ ,

$$\begin{aligned} \|x^\alpha \psi(x)\|_{H_{(\sigma)}^t(\mathbf{J})}^2 &\lesssim \sum_{j=0}^{\lfloor t \rfloor} \|D^j(x^\alpha \psi(x))\|_{L_{(\sigma+j)}^2(\mathbf{J})}^2 + \iint_{\Lambda} x^{\sigma+t} \frac{|D^{\lfloor t \rfloor}(x^\alpha \psi(x)) - D^{\lfloor t \rfloor}(y^\alpha \psi(y))|^2}{|x - y|^{1+2(t-\lfloor t \rfloor)}} dy dx \\ &\quad + \iint_{\Lambda_1} x^{\mu+t} \frac{|D^{\lfloor t \rfloor}(x^\alpha \psi(x)) - D^{\lfloor t \rfloor}(y^\alpha \psi(y))|^2}{|x - y|^{1+2(t-\lfloor t \rfloor)}} dy dx \\ &\lesssim \sum_{j=0}^{\lfloor t \rfloor} \|D^j(x^\alpha \psi(x))\|_{L_{(\sigma+j)}^2(\mathbf{J})}^2 + \iint_{\Lambda} x^{\sigma+t} \frac{|D^{\lfloor t \rfloor}(x^\alpha \psi(x)) - D^{\lfloor t \rfloor}(y^\alpha \psi(y))|^2}{|x - y|^{1+2(t-\lfloor t \rfloor)}} dy dx \\ &\quad + \|h\|_{H_{(\mu)}^t(\mathbf{J})}^2 \\ &\lesssim \sum_{j=0}^{\lfloor t \rfloor} \|D^j(x^\alpha \psi(x))\|_{L_{(\sigma+j)}^2(\mathbf{J})}^2 + \iint_{\Lambda} x^{\sigma+t} \frac{|D^{\lfloor t \rfloor}(x^\alpha \psi(x)) - D^{\lfloor t \rfloor}(y^\alpha \psi(y))|^2}{|x - y|^{1+2(t-\lfloor t \rfloor)}} dy dx \\ &\quad + \|\psi\|_{H_{(\mu)}^s(\mathbf{J})}^2 \quad (\text{using Lemma 6.2}). \end{aligned}$$

Hence for the analysis of the regularity of  $f(x) = (1-x)^{\alpha-\beta} x^\beta \phi(x)$  we can restrict our attention on the analysis of the semi-norm to  $|f|_{H_{(\sigma)}^t(\Lambda)}$ .

## 6.2 Regularity of $x^p\psi(x)$

In this section we establish the value of  $t$  and  $\sigma$  such that  $x^p\psi(x) \in H_{(\sigma)}^t(J)$  for  $\psi(x) \in H_{(\mu)}^s(J)$ . The general result is given in Theorem 6.4. There are two key terms which arise in the proof of Theorem 6.4. The analysis for one of these terms follows similarly to a term which occurs for the case of  $s$  between 0 and 1, discussed in Theorem 6.2. The other of these terms arises for the case of  $s$  between 1 and 2, discussed in Theorem 6.3. We begin this section with an embedding theorem which is used in the proofs of the subsequent theorems in this section.

**Theorem 6.1** *For  $s \geq 0$ ,  $\gamma - s > -1$ , then  $H_{(\gamma)}^s(J) \subset L_{(\gamma-s)}^2(J)$ .*

**Proof:** Firstly we consider the case for  $0 \leq s \leq 1$ .

For  $s = 0$  we have that

$$H_{(\gamma)}^s(J) = H_{(\gamma)}^0(J) = L_{(\gamma)}^2(J). \quad (6.9)$$

For  $s = 1$ , consider  $\phi \in L_{(\gamma-s)}^2(J) = L_{(\gamma-1)}^2(J)$ . Then, using Hardy's inequality [8, Lemma 3.2],

$$\begin{aligned} \|\phi\|_{L_{(\gamma-1)}^2}^2 &= \int_J x^{\gamma-1} (\phi(x))^2 dx \\ &\lesssim \int_J x^{\gamma+1} (\phi'(x))^2 dx + \int_J x^{\gamma+1} (\phi(x))^2 dx \\ &\leq \int_J x^{\gamma+1} (\phi'(x))^2 dx + \int_J x^{\gamma} (\phi(x))^2 dx \\ &= \|\phi\|_{H_{(\gamma)}^1}^2. \end{aligned} \quad (6.10)$$

From (6.9) and (6.10) the identity operator  $\mathbb{I}$  mapping from  $H_{(\gamma)}^s(J) \rightarrow L_{(\gamma-s)}^2(J)$ ,  $s = 0, 1$  is a bounded operator.

Additionally, the spaces  $L_{(\sigma)}^2(J)$  are interpolation spaces [36, Lemma 23.1].

Hence, for  $\theta = (1 - \theta)0 + \theta 1$ , using the fact that  $H_{(\gamma)}^{\theta}(J)$  and  $L_{(\theta)}^2(J)$  are interpolation spaces, it follows that

$$H_{(\gamma)}^{\theta}(J) \xrightarrow{\mathbb{I}} L_{((1-\theta)\gamma + \theta(\gamma-1))}^2(J) = L_{(\gamma-\theta)}^2(J)$$

is bounded. Thus, if  $u \in H_{(\gamma)}^{\theta}(J)$  then  $u \in L_{(\gamma-\theta)}^2(J)$  with  $\|u\|_{L_{(\gamma-\theta)}^2} \leq C \|u\|_{H_{(\gamma)}^{\theta}}$ .

Next, for  $1 \leq s \leq 2$ , consider  $s = 2$ . For  $\phi \in L_{(\gamma-s)}^2(J) = L_{(\gamma-2)}^2(J)$ , again using Hardy's inequality (and that  $\gamma - s > -1$ ),

$$\begin{aligned} \|\phi\|_{L_{(\gamma-2)}^2}^2 &= \int_J x^{\gamma-2} (\phi(x))^2 dx \\ &\lesssim \int_J x^{\gamma} (\phi'(x))^2 dx + \int_J x^{\gamma} (\phi(x))^2 dx \\ &\lesssim \int_J x^{\gamma+2} (\phi''(x))^2 dx + \int_J x^{\gamma+2} (\phi'(x))^2 dx + \int_J x^{\gamma} (\phi(x))^2 dx \\ &\leq \int_J x^{\gamma+2} (\phi''(x))^2 dx + \int_J x^{\gamma+1} (\phi'(x))^2 dx + \int_J x^{\gamma} (\phi(x))^2 dx \\ &= \|\phi\|_{H_{(\gamma)}^2}^2. \end{aligned} \quad (6.11)$$



Again, using the fact that  $H_{(\gamma)}^\theta(\mathbf{J})$  and  $L_{(\theta)}^2(\mathbf{J})$  are interpolation spaces, it now follows that for  $0 \leq \theta \leq 2$  if  $u \in H_{(\gamma)}^\theta(\mathbf{J})$  then  $u \in L_{(\gamma-\theta)}^2(\mathbf{J})$  with  $\|u\|_{L_{(\gamma-\theta)}^2} \leq C \|u\|_{H_{(\gamma)}^\theta}$ .

The argument extends in an obvious manner to arbitrary  $s > 2$ . ■

**Theorem 6.2** *Let  $0 \leq s < 1$ ,  $\mu > -1$ , and  $\psi \in H_{(\mu)}^s(\mathbf{J})$ . Then  $x^p \psi \in H_{(\sigma)}^t(\mathbf{J})$  provided*

$$0 \leq t \leq s, \quad \sigma + 2p \geq \mu, \quad \sigma + 2p - t > -1, \quad \text{and} \quad \sigma + 2p + t \geq \mu + s. \quad (6.12)$$

*Additionally, when (6.12) is satisfied, there exists  $C > 0$  (independent of  $\psi$ ) such that  $\|x^p \psi\|_{H_{(\sigma)}^t(\mathbf{J})} \leq C \|\psi\|_{H_{(\mu)}^s(\mathbf{J})}$ .*

**Proof:** Firstly, for  $s = t = 0$ ,

$$\begin{aligned} \|x^p \psi\|_{H_{(\sigma)}^0}^2 &= \|x^p \psi\|_{H_{(\sigma)}^0}^2 = \int_{\mathbf{J}} x^\sigma (x^p \psi(x))^2 dx = \int_{\mathbf{J}} x^{\sigma+2p} (\psi(x))^2 dx \\ &\leq \int_{\mathbf{J}} x^\mu (\psi(x))^2 dx, \quad \text{provided } \sigma + 2p \geq \mu, \\ &= \|\psi\|_{H_{(\mu)}^0}^2 = \|\psi\|_{H_{(\mu)}^s}^2. \end{aligned} \quad (6.13)$$

For  $0 < s < 1$ , in addition to (6.13) we must also consider the semi-norm  $|x^p \psi|_{H_{(\sigma)}^t(\Lambda)}$ .

$$\begin{aligned} |x^p \psi|_{H_{(\sigma)}^t(\Lambda)}^2 &= \iint_{\Lambda} x^{\sigma+t} \frac{|x^p \psi(x) - y^p \psi(y)|^2}{|x - y|^{1+2t}} dy dx \\ &\lesssim \iint_{\Lambda} x^{\sigma+t} y^{2p} \frac{|\psi(x) - \psi(y)|^2}{|x - y|^{1+2t}} dy dx + \iint_{\Lambda} x^{\sigma+t} \psi^2(x) \frac{|x^p - y^p|^2}{|x - y|^{1+2t}} dy dx \\ &:= I_1 + I_2. \end{aligned} \quad (6.14)$$

Noting that in  $\Lambda$ ,  $y < \frac{3}{2}x$ , for  $I_1$  we have

$$\begin{aligned} I_1 &\lesssim \iint_{\Lambda} x^{\sigma+t} x^{2p} \frac{|\psi(x) - \psi(y)|^2}{|x - y|^{1+2t}} dy dx \\ &\leq \iint_{\Lambda} x^{\mu+s} \frac{|\psi(x) - \psi(y)|^2}{|x - y|^{1+2s}} dy dx, \quad \text{provided } \sigma + t + 2p \geq \mu + s, \text{ and } t \leq s, \\ &= |\psi|_{H_{(\mu)}^s(\Lambda)}^2. \end{aligned} \quad (6.15)$$

If  $p = 0$  then  $I_2 = 0$ . To bound  $I_2$  for  $p \neq 0$  we introduce the change of variable:  $y = \frac{1}{z}x$ , where  $\frac{2}{3} < z < \frac{3}{2}$ . With this change of variable, we have

$$\begin{aligned} dy &= \frac{-1}{z^2} x dz, \quad |x - y| = x \frac{1}{z} |z - 1|, \quad \text{and for } I_2 \\ I_2 &= \int_0^{1/2} x^{\sigma+2p-t} \psi^2(x) dx \int_{2/3}^{3/2} z^{-1+2t-2p} |1 - z|^{-1-2t} |z^p - 1|^2 dz \end{aligned} \quad (6.16)$$

$$\lesssim \|\psi\|_{L_{(\sigma+2p-t)}^2}^2 \cdot \int_{2/3}^{3/2} |1 - z|^{-1-2t} |z^p - 1|^2 dz. \quad (6.17)$$

Using Hardy's inequality [8, Lemma 3.2], we bound the integral in (6.17) as follows.

$$\begin{aligned}
\int_{2/3}^{3/2} |1-z|^{-1-2t} |z^p-1|^2 dz &= \int_{2/3}^1 (1-z)^{-1-2t} (z^p-1)^2 dz + \int_1^{3/2} (z-1)^{-1-2t} (z^p-1)^2 dz \\
&\lesssim \int_{2/3}^1 (1-z)^{-1-2t+2} (z^{p-1})^2 dz + \int_1^{3/2} (z-1)^{-1-2t+2} (z^{p-1})^2 dz \\
&\lesssim 1, \quad \text{provided } t < 1.
\end{aligned} \tag{6.18}$$

From Theorem 6.1, we have

$$\begin{aligned}
\|\psi\|_{L^2_{(\sigma+2p-t)}} &\lesssim \|\psi\|_{H^t_{(\sigma+2p)}}, \quad \text{provided } \sigma + 2p - t > -1, \\
&\leq \|\psi\|_{H^s_{(\mu)}}, \quad \text{provided } \sigma + 2p \geq \mu, \text{ and } t \leq s.
\end{aligned} \tag{6.19}$$

Finally, combining (6.13)-(6.19) we obtain the stated results. ■

**Theorem 6.3** *Let  $1 \leq s < 2$ ,  $\mu > -1$ , and  $\psi \in H^s_{(\mu)}(\mathbf{J})$ . Then  $x^p \psi \in H^t_{(\sigma)}(\mathbf{J})$  provided*

$$0 \leq t \leq s, \quad \sigma + 2p \geq \mu, \quad \sigma + 2p - t > -1, \quad \text{and} \quad \sigma + 2p + t \geq \mu + s. \tag{6.20}$$

*Additionally, when (6.20) is satisfied, there exists  $C > 0$  (independent of  $\psi$ ) such that  $\|x^p \psi\|_{H^t_{(\sigma)}(\mathbf{J})} \leq C \|\psi\|_{H^s_{(\mu)}(\mathbf{J})}$ .*

**Proof:** For  $0 \leq t < 1$  Theorem 6.2 applies. We assume that  $t \geq 1$ . Hence, of interest is

$$\|x^p \psi\|_{L^2_{(\sigma)}}, \quad \|D(x^p \psi)\|_{L^2_{(\sigma+1)}}, \quad |D(x^p \psi)|_{H^{t-1}_{(\sigma+1)}(\Lambda)}.$$

From Theorem 6.2,

$$\|x^p \psi\|_{L^2_{(\sigma)}} \lesssim \|\psi\|_{H^0_{(\mu)}}, \quad \text{provided } \sigma + 2p \geq \mu. \tag{6.21}$$

For  $D(x^p \psi)$  we have

$$\begin{aligned}
D(x^p \psi) &= x^p \psi' + p x^{p-1} \psi \quad \text{and} \\
(D(x^p \psi))^2 &\lesssim (x^p \psi')^2 + (x^{p-1} \psi)^2.
\end{aligned}$$

Then, using Hardy's inequality [8, Lemma 3.2] (using  $\sigma + 2p - t > -1$ , i.e.,  $\sigma + 2p > 0$ ),

$$\begin{aligned}
\|D(x^p \psi)\|_{L^2_{(\sigma+1)}}^2 &\lesssim \int_0^{3/4} x^{\sigma+1} (x^p \psi'(x))^2 dx + \int_0^{3/4} x^{\sigma+1} (x^{p-1} \psi(x))^2 dx \\
&= \int_0^{3/4} x^{\sigma+2p+1} (\psi'(x))^2 dx + \int_0^{3/4} x^{\sigma+2p-1} (\psi(x))^2 dx \\
&\lesssim \int_0^{3/4} x^{\sigma+2p+1} (\psi'(x))^2 dx \\
&\quad + \int_0^{3/4} x^{\sigma+2p+1} (\psi'(x))^2 dx + \int_0^{3/4} x^{\sigma+2p+1} (\psi(x))^2 dx \\
&\lesssim \|\psi\|_{H^1_{(\mu)}}^2, \quad \text{provided } \sigma + 2p \geq \mu.
\end{aligned} \tag{6.22}$$

Equation (6.22), together with (6.21) establishes the stated result for  $t = s = 1$ .

Next, for  $|D(x^p \psi)|_{H_{(\sigma+1)}^{t-1}(\Lambda)}$  we have

$$\begin{aligned}
|D(x^p \psi)|_{H_{(\sigma+1)}^{t-1}(\Lambda)}^2 &= \iint_{\Lambda} x^{\sigma+t} \frac{|D(x^p \psi(x)) - D(y^p \psi(y))|^2}{|x - y|^{1+2(t-1)}} dy dx \\
&\lesssim \iint_{\Lambda} x^{\sigma+t} \frac{|x^{p-1} \psi(x) - y^{p-1} \psi(y)|^2}{|x - y|^{1+2(t-1)}} dy dx + \iint_{\Lambda} x^{\sigma+t} \frac{|x^p \psi'(x) - y^p \psi'(y)|^2}{|x - y|^{1+2(t-1)}} dy dx \\
&:= I_3 + I_4.
\end{aligned} \tag{6.23}$$

For  $I_3$ , proceeding as in (6.14),

$$\begin{aligned}
I_3 &\lesssim \iint_{\Lambda} x^{\sigma+t} y^{2(p-1)} \frac{|\psi(x) - \psi(y)|^2}{|x - y|^{1+2(t-1)}} dy dx + \iint_{\Lambda} x^{\sigma+t} \psi^2(x) \frac{|x^{p-1} - y^{p-1}|^2}{|x - y|^{1+2(t-1)}} dy dx \\
&:= I_{3,1} + I_{3,2}.
\end{aligned}$$

For  $I_{3,1}$ , using  $y < \frac{3}{2}x$ , and the introducing the change of variable  $y = \frac{1}{z}x$ , where  $\frac{2}{3} < z < \frac{3}{2}$  we obtain

$$\begin{aligned}
I_{3,1} &\lesssim \int_{x=0}^{1/2} x^{\sigma+t+2(p-1)} \int_{z=2/3}^{3/2} \frac{z^{1+2(t-1)-2(p-1)}}{(x|z-1|)^{1+2(t-1)}} |\psi(x) - \psi(\frac{1}{z}x)|^2 z^{-2} x dz dx \\
&\lesssim \int_{x=0}^{1/2} x^{\sigma+2p-t} \left( \int_{z=2/3}^1 (1-z)^{1-2t} (\psi(x) - \psi(\frac{1}{z}x))^2 dz \right. \\
&\quad \left. + \int_{z=1}^{3/2} (z-1)^{1-2t} (\psi(x) - \psi(\frac{1}{z}x))^2 dz \right) dx \\
&\lesssim \int_{x=0}^{1/2} x^{\sigma+2p-t} \left( \int_{z=2/3}^1 (1-z)^{3-2t} z^{-4} x^2 (\psi'(\frac{1}{z}x))^2 dz \right. \\
&\quad \left. + \int_{z=1}^{3/2} (z-1)^{3-2t} z^{-4} x^2 (\psi'(\frac{1}{z}x))^2 dz \right) dx \\
&\quad \text{(using Hardy's inequality)} \\
&\lesssim \int_{x=0}^{1/2} x^{\sigma+2p-t+2} \int_{z=2/3}^{3/2} |1-z|^{3-2t} (\psi'(\frac{1}{z}x))^2 dz dx.
\end{aligned} \tag{6.24}$$

Next, letting  $w = \frac{1}{z}x$ , then  $dw = \frac{1}{z}dx$  and

$$\begin{aligned}
I_{3,1} &\lesssim \int_{z=2/3}^{3/2} |1-z|^{3-2t} \int_{w=0}^{1/2z} (zw)^{\sigma+2p-t+2} (\psi'(w))^2 z dw dz \\
&\lesssim \int_{2/3}^{3/2} |1-z|^{3-2t} dz \int_0^{3/4} w^{\sigma+2p-t+2} (\psi'(w))^2 dw \\
&\lesssim \|\psi'\|_{L^2_{(\sigma+2p-t+2)}}^2, \text{ as } t < 2, \\
&\lesssim \|\psi'\|_{H^{t-1}_{(\sigma+2p+1)}}^2, \text{ (using Theorem 6.1) provided } \sigma + 2p - t + 2 > -1, \\
&\lesssim \|\psi'\|_{H^{s-1}_{(\mu+1)}}^2, \text{ provided } \sigma + 2p \geq \mu, \text{ and } t \leq s, \\
&\lesssim \|\psi\|_{H^s_{(\mu)}}^2.
\end{aligned} \tag{6.25}$$

If  $p = 1$  then  $I_{3,2} = 0$ . For  $I_{3,2}$  with  $p \neq 1$ , proceeding as in the approach used to obtain the bound for  $I_2$ , (6.16) - (6.18),

$$\begin{aligned}
I_{3,2} &\lesssim \int_0^{1/2} x^{(\sigma+1)-(t-1)+2(p-1)} (\psi(x))^2 dx \int_{2/3}^{3/2} z^{-1+2(t-1)-2(p-1)} |1-z|^{-1-2(t-1)} |z^{p-1} - 1|^2 dz \\
&\lesssim \int_{x=0}^{1/2} x^{\sigma+2p-t} (\psi(x))^2 dx, \text{ provided } (t-1) < 1, \text{ i.e., } t < 2, \\
&\lesssim \int_0^{1/2} x^{\sigma+2p-t+2} (\psi'(x))^2 dx + \int_0^{1/2} x^{\sigma+2p-t+2} (\psi(x))^2 dx, \\
&\quad \text{(using Hardy's inequality) provided } \sigma + 2p - t > -1, \\
&\lesssim \|\psi'\|_{L^2_{(\sigma+2p-t+2)}}^2 + \|\psi\|_{L^2_{(\sigma+2p-t+2)}}^2 \\
&\lesssim \|\psi'\|_{H^{t-1}_{(\sigma+2p+1)}}^2 + \|\psi\|_{H^t_{(\sigma+2p+2)}}^2, \text{ (using Theorem 6.1) provided } \sigma + 2p - t + 2 > -1, \\
&\lesssim \|\psi'\|_{H^{s-1}_{(\mu+1)}}^2 + \|\psi\|_{H^s_{(\mu+2)}}^2, \text{ provided } \sigma + 2p \geq \mu \text{ and } t \leq s, \\
&\lesssim \|\psi\|_{H^s_{(\mu)}}^2.
\end{aligned} \tag{6.26}$$

With  $I_4$ , proceeding as in (6.14),

$$\begin{aligned}
I_4 &\lesssim \iint_{\Lambda} x^{\sigma+t} y^{2p} \frac{|\psi'(x) - \psi'(y)|^2}{|x - y|^{1+2(t-1)}} dy dx + \iint_{\Lambda} x^{\sigma+t} (\psi'(x))^2 \frac{|x^p - y^p|^2}{|x - y|^{1+2(t-1)}} dy dx \\
&\lesssim |\psi'|_{H^{s-1}_{(\mu+1)}(\Lambda)}^2 \text{ (provided } \sigma + 2p + t \geq \mu + s, \text{ and } t \leq s) \\
&\quad + \|\psi'\|_{H^{s-1}_{(\mu+1)}}^2 \text{ ( provided } \sigma + 2p - t + 2 > -1, \sigma + 2p \geq \mu, \text{ and } 1 < t \leq s) \\
&\lesssim \|\psi\|_{H^s_{(\mu)}}^2.
\end{aligned} \tag{6.27}$$

Combining (6.21) - (6.27) we obtain the stated result. ■

We are now in a position to state the general result.

**Theorem 6.4** *Let  $s \geq 0$ ,  $\mu > -1$ , and  $\psi \in H_{(\mu)}^s(\mathbb{J})$ . Then  $x^p \psi \in H_{(\sigma)}^t(\mathbb{J})$  provided*

$$0 \leq t \leq s, \quad \sigma + 2p \geq \mu, \quad \sigma + 2p - t > -1, \quad \text{and} \quad \sigma + 2p + t \geq \mu + s. \quad (6.28)$$

*Additionally, when (6.28) is satisfied, there exists  $C > 0$  (independent of  $\psi$ ) such that  $\|x^p \psi\|_{H_{(\sigma)}^t(\mathbb{J})} \leq C \|\psi\|_{H_{(\mu)}^s(\mathbb{J})}$ .*

**Proof:** The proof is an induction argument, using Theorem 6.3 as the initial step.

For  $t = s = n$  we have,

$$\begin{aligned} \|D^t(x^p \psi(x))\|_{L_{(\sigma+t)}^2}^2 &\lesssim \sum_{j=0}^n \|x^{p-j} D^{n-j} \psi(x)\|_{L_{(\sigma+n)}^2}^2 \\ &= \sum_{j=0}^n \int_0^{3/4} x^{\sigma+n} (x^{p-j} D^{n-j} \psi(x))^2 dx \\ &= \sum_{j=0}^n \int_0^{3/4} x^{\sigma+n+2(p-j)} (D^{n-j} \psi(x))^2 dx \\ &\quad (\text{applying Hardy's inequality } j \text{ times, using } \sigma + 2p - n > -1) \\ &\lesssim \sum_{j=0}^n \sum_{k=0}^j \int_0^{3/4} x^{\sigma+2p+n} (D^{n-j+k} \psi(x))^2 dx \\ &\lesssim \sum_{j=0}^n \|D^j \psi\|_{L_{(\sigma+2p+n)}^2}^2 \\ &\lesssim \sum_{j=0}^n \|D^j \psi\|_{L_{(\mu+n)}^2}^2 \quad (\text{using } \sigma + 2p \geq \mu) \\ &\lesssim \sum_{j=0}^n \|D^j \psi\|_{L_{(\mu+j)}^2}^2 \lesssim \|\psi\|_{H_{(\mu)}^n}^2 = \|\psi\|_{H_{(\mu)}^s}^2. \end{aligned} \quad (6.29)$$

Equation (6.29), together with the induction assumption establishes the result for  $t = s = n$ .

For  $n < t, s < n + 1$  we also need to consider  $|D^n(x^p \psi(x))|_{H_{(\sigma+n)}^{t-n}(\Lambda)}$ .

$$\begin{aligned}
|D^n(x^p \psi(x))|_{H_{(\sigma+n)}^{t-n}(\Lambda)}^2 &= \sum_{j=0}^n \iint_{\Lambda} x^{\sigma+t} \frac{|x^{p-j} D^{n-j} \psi(x) - y^{p-j} D^{n-j} \psi(y)|^2}{|x - y|^{1+2(t-n)}} dy dx \\
&\lesssim \sum_{j=0}^n \left( \iint_{\Lambda} x^{\sigma+t} y^{2(p-j)} \frac{|D^{n-j} \psi(x) - D^{n-j} \psi(y)|^2}{|x - y|^{1+2(t-n)}} dy dx \right. \\
&\quad \left. + \iint_{\Lambda} x^{\sigma+t} (D^{n-j} \psi(x))^2 \frac{|x^{p-j} - y^{p-j}|^2}{|x - y|^{1+2(t-n)}} dy dx \right) \\
&= \iint_{\Lambda} x^{\sigma+t} y^{2p} \frac{|D^n \psi(x) - D^n \psi(y)|^2}{|x - y|^{1+2(t-n)}} dy dx \\
&\quad + \sum_{j=1}^n \iint_{\Lambda} x^{\sigma+t} y^{2(p-j)} \frac{|D^{n-j} \psi(x) - D^{n-j} \psi(y)|^2}{|x - y|^{1+2(t-n)}} dy dx \\
&\quad + \sum_{j=0}^n \iint_{\Lambda} x^{\sigma+t} (D^{n-j} \psi(x))^2 \frac{|x^{p-j} - y^{p-j}|^2}{|x - y|^{1+2(t-n)}} dy dx. \tag{6.30}
\end{aligned}$$

The first term in (6.30) is bounded in a similar manner to  $I_1$  in Theorem 6.2, with  $\sigma \rightarrow \sigma + n$ ,  $t \rightarrow t - n$ , to obtain

$$\iint_{\Lambda} x^{\sigma+t} y^{2p} \frac{|D^n \psi(x) - D^n \psi(y)|^2}{|x - y|^{1+2(t-n)}} dy dx \lesssim |\psi|_{H_{(\mu)}^s(\Lambda)}^2, \text{ provided } \sigma + 2p + t \geq \mu + s, \text{ and } t \leq s. \tag{6.31}$$

For the second term in (6.30) the terms in the summation are bounded in a similar manner to  $I_{3,1}$  in Theorem 6.3,  $\sigma \rightarrow \sigma + n - 1$ ,  $t \rightarrow t - n + 1$ ,  $p \rightarrow p + 1 - j$ , to obtain

$$\begin{aligned}
&\iint_{\Lambda} x^{\sigma+t} y^{2(p-j)} \frac{|D^{n-j} \psi(x) - D^{n-j} \psi(y)|^2}{|x - y|^{1+2(t-n)}} dy dx \\
&\lesssim \|D^{n-j+1} \psi\|_{L_{(\sigma+2p+n-j+1-(t-n+j-1))}^2}^2 \\
&\lesssim \|D^{n-j+1} \psi\|_{H_{(\sigma+2p+n-j+1)}^{t-(n-j+1)}}^2, \text{ provided } \sigma + 2p + n - j + 1 - (t - n + j - 1) > -1 \\
&\lesssim \|D^{n-j+1} \psi\|_{H_{(\mu+n-j+1)}^{s-(n-j+1)}}^2, \text{ provided } \sigma + 2p \geq \mu \text{ and } t \leq s, \\
&\lesssim \|\psi\|_{H_{(\mu)}^s}^2. \tag{6.32}
\end{aligned}$$

For the third term in (6.30) the terms in the summation are bounded in a similar manner to  $I_2$  in

Theorem 6.2,  $\sigma \rightarrow \sigma + n$ ,  $t \rightarrow t - n$ ,  $p \rightarrow p - j$ , to obtain

$$\begin{aligned}
& \iint_{\Lambda} x^{\sigma+t} (D^{n-j}\psi(x))^2 \frac{|x^{p-j} - y^{p-j}|^2}{|x - y|^{1+2(t-n)}} dy dx \\
& \lesssim \|D^{n-j}\psi\|_{L^2_{(\sigma+2(p-j)+n-(t-n))}}^2 = \|D^{n-j}\psi\|_{L^2_{(\sigma+2p+n-j-(t-n+j))}}^2, \quad \text{provided } t < n+1, \\
& \lesssim \|D^{n-j}\psi\|_{H^{t-(n-j)}_{(\sigma+2p+n-j)}}^2, \quad \text{provided } \sigma + 2p + n - j - (t - n + j) > -1 \\
& \lesssim \|D^{n-j}\psi\|_{H^{s-(n-j)}_{(\mu+n-j)}}^2, \quad \text{provided } \sigma + 2p \geq \mu \text{ and } t \leq s, \\
& \lesssim \|\psi\|_{H^s_{(\mu)}}^2.
\end{aligned} \tag{6.33}$$

Combining (6.29) - (6.33) the stated result follows. ■

## 7 Regularity of the solution to the fractional diffusion, advection, reaction equation (1.3)

In this section we present the analysis for the regularity of the solution to (1.3). The solution's regularity can be influenced by the regularity of the coefficients  $b(x)$  and  $c(x)$ . We begin with two lemmas which enables us to insulate the influence of these terms.

Introduce the space  $W_w^{k,\infty}(\mathbf{I})$  and its associated norm, defined for  $k \in \mathbb{N}_0$ , as

$$W_w^{k,\infty}(\mathbf{I}) := \left\{ f : (1-x)^{j/2} x^{j/2} D^j f(x) \in L^\infty(\mathbf{I}), j = 0, 1, \dots, k \right\}, \tag{7.1}$$

$$\|f\|_{W_w^{k,\infty}} := \max_{0 \leq j \leq k} \|(1-x)^{j/2} x^{j/2} D^j f(x)\|_{L^\infty(\mathbf{I})}. \tag{7.2}$$

The subscript  $w$  denotes the fact that  $W_w^{k,\infty}(\mathbf{I})$  is a weaker space than  $W^{k,\infty}(\mathbf{I})$  in that the derivative of functions in  $W_w^{k,\infty}(\mathbf{I})$  may be unbounded at the endpoints of the interval.

**Lemma 7.1** *Let  $\alpha, \beta > -1$ ,  $0 \leq s \leq k \in \mathbb{N}_0$ , and  $f \in W_w^{k,\infty}(\mathbf{I})$ . Then, for*

$$g \in H^s_{(\alpha,\beta)}(\mathbf{I}) \text{ we have that } fg \in H^s_{(\alpha,\beta)}(\mathbf{I}). \tag{7.3}$$

**Proof:** We establish Lemma 7.1 for  $s = 0$ ,  $s = 1$ , and  $0 < s < 1$ . The proof extends in an obvious manner for  $s > 1$ .

If  $s = 0$  then  $k \geq 0$ , and

$$\begin{aligned}
\|fg\|_{H^s_{(\alpha,\beta)}(\mathbf{I})}^2 &= \|fg\|_{H^0_{(\alpha,\beta)}(\mathbf{I})}^2 = \int_0^1 (1-x)^\alpha x^\beta (f(x)g(x))^2 dx \\
&\lesssim \|f\|_{L^\infty(\mathbf{I})}^2 \|g\|_{H^0_{(\alpha,\beta)}(\mathbf{I})}^2 = \|f\|_{W_w^{0,\infty}(\mathbf{I})}^2 \|g\|_{H^0_{(\alpha,\beta)}(\mathbf{I})}^2.
\end{aligned} \tag{7.4}$$

Hence (7.3) is established for  $s = 0$ .

For  $s = 1$  then  $k \geq 1$ , and

$$\begin{aligned}
\|fg\|_{H_{(\alpha,\beta)}^1(\mathbf{I})}^2 &= \|fg\|_{L_{(\alpha,\beta)}^2(\mathbf{I})}^2 + \|D(fg)\|_{L_{(\alpha+1,\beta+1)}^2(\mathbf{I})}^2 \\
&= \int_0^1 (1-x)^\alpha x^\beta (f(x)g(x))^2 dx + \int_0^1 (1-x)^{\alpha+1} x^{\beta+1} (g(x)Df(x) + f(x)Dg(x))^2 dx \\
&\leq \int_0^1 f(x)^2 (1-x)^\alpha x^\beta (g(x))^2 dx + 2 \int_0^1 (1-x)^1 x^1 (Df(x))^2 (1-x)^\alpha x^\beta (g(x))^2 dx \\
&\quad + 2 \int_0^1 f(x)^2 (1-x)^{\alpha+1} x^{\beta+1} (Dg(x))^2 dx \\
&\lesssim \left( \|f\|_{L^\infty(\mathbf{I})}^2 + \|(1-x)^{1/2} x^{1/2} Df(x)\|_{L^\infty(\mathbf{I})}^2 \right) \left( \|g\|_{L_{(\alpha,\beta)}^2(\mathbf{I})}^2 + \|Dg\|_{L_{(\alpha+1,\beta+1)}^2(\mathbf{I})}^2 \right) \\
&= \|f\|_{W_w^{1,\infty}(\mathbf{I})}^2 \|g\|_{H_{(\alpha,\beta)}^1(\mathbf{I})}^2. \tag{7.5}
\end{aligned}$$

Hence (7.3) is established for  $s = 1$ .

Note that for  $f \in W_w^{1,\infty}(\mathbf{I})$ , from (7.4),  $\|fg\|_{H_{(\alpha,\beta)}^0(\mathbf{I})} \leq \|f\|_{W_w^{1,\infty}(\mathbf{I})} \|g\|_{H_{(\alpha,\beta)}^0(\mathbf{I})}$ . Combining this with (7.5) and the fact that  $H_{(\alpha,\beta)}^s(\mathbf{I})$  are interpolation spaces, it follows that for  $0 < s < 1$ ,  $\|fg\|_{H_{(\alpha,\beta)}^s(\mathbf{I})} \leq \|f\|_{W_w^{1,\infty}(\mathbf{I})} \|g\|_{H_{(\alpha,\beta)}^s(\mathbf{I})}$ . ■

Arising in the analysis is the product of a function and a functional (e.g.,  $b(x)Du(x)$ ). To define such a product it is convenient to consider  $L_{(\alpha,\beta)}^2(\mathbf{I}) = H_{(\alpha,\beta)}^0(\mathbf{I})$  as the pivot space for  $H_{(\alpha,\beta)}^s(\mathbf{I})$  and  $H_{(\alpha,\beta)}^{-s}(\mathbf{I})$ , with  $H_{(\alpha,\beta)}^{-s}(\mathbf{I})$  characterized as the closure of  $L_{(\alpha,\beta)}^2(\mathbf{I})$  with respect to the operator norm

$$|||v||| := \sup_{h \in H_{(\alpha,\beta)}^s(\mathbf{I})} \frac{\langle v, h \rangle_{\rho^{(\alpha,\beta)}}}{\|h\|_{H_{(\alpha,\beta)}^s(\mathbf{I})}}.$$

**Definition:** Product of a function and a functional.

Let  $g \in H_{(\alpha,\beta)}^{-s}(\mathbf{I})$ . Then there exists  $\{g_i\}_{i=1}^\infty \subset L_{(\alpha,\beta)}^2(\mathbf{I})$ , such that  $\lim_{i \rightarrow \infty} |||g - g_i||| = 0$ . Thus, for any  $h \in H_{(\alpha,\beta)}^s(\mathbf{I})$ ,  $g(h) = \lim_{i \rightarrow \infty} \langle g_i, h \rangle_{\rho^{(\alpha,\beta)}}$ .

For  $f \in W_w^{k,\infty}(\mathbf{I})$ , let  $fg$  be defined as

$$fg(h) := \lim_{i \rightarrow \infty} \langle fg_i, h \rangle_{\rho^{(\alpha,\beta)}}. \tag{7.6}$$

**Lemma 7.2** *Let  $\alpha, \beta > -1$ ,  $0 \leq s \leq k \in \mathbb{N}_0$ , and  $f \in W_w^{k,\infty}(\mathbf{I})$ . Then, for*

$$g \in H_{(\alpha,\beta)}^{-s}(\mathbf{I}) \text{ we have that } fg \in H_{(\alpha,\beta)}^{-s}(\mathbf{I}). \tag{7.7}$$

**Proof:** To establish that  $fg$  is well defined and contained in  $H_{(\alpha,\beta)}^{-s}(\mathbf{I})$ , we show that  $\{fg_i\}_{i=1}^\infty$  is a Cauchy sequence with respect to the norm  $||| \cdot |||$ .



As (7.7) trivially holds for  $f = 0$ , assume  $0 \neq f \in W_w^{k,\infty}(\mathbf{I})$ . Then,

$$\begin{aligned}
\|f g_i - f g_j\| &= \sup_{h \in H_{(\alpha,\beta)}^s(\mathbf{I})} \frac{\langle f g_i - f g_j, h \rangle_{\rho^{(\alpha,\beta)}}}{\|h\|_{H_{(\alpha,\beta)}^s(\mathbf{I})}} = \|f\|_{W_w^{k,\infty}(\mathbf{I})} \sup_{h \in H_{(\alpha,\beta)}^s(\mathbf{I})} \frac{\langle g_i - g_j, f h \rangle_{\rho^{(\alpha,\beta)}}}{\|f\|_{W_w^{k,\infty}(\mathbf{I})} \|h\|_{H_{(\alpha,\beta)}^s(\mathbf{I})}} \\
&\leq \|f\|_{W_w^{k,\infty}(\mathbf{I})} \sup_{h \in H_{(\alpha,\beta)}^s(\mathbf{I})} \frac{\langle g_i - g_j, f h \rangle_{\rho^{(\alpha,\beta)}}}{\|f h\|_{H_{(\alpha,\beta)}^s(\mathbf{I})}} \quad (\text{using Lemma 7.1}), \\
&\leq \|f\|_{W_w^{k,\infty}(\mathbf{I})} \sup_{\tilde{h} \in H_{(\alpha,\beta)}^s(\mathbf{I})} \frac{\langle g_i - g_j, \tilde{h} \rangle_{\rho^{(\alpha,\beta)}}}{\|\tilde{h}\|_{H_{(\alpha,\beta)}^s(\mathbf{I})}} \\
&\leq \|f\|_{W_w^{k,\infty}(\mathbf{I})} \|g_i - g_j\|.
\end{aligned}$$

As  $\{g_i\}_{i=1}^\infty$  is a Cauchy sequence in  $H_{(\alpha,\beta)}^{-s}(\mathbf{I})$ , then it follows that  $\{f g_i\}_{i=1}^\infty$  is also a Cauchy sequence with limit in  $H_{(\alpha,\beta)}^{-s}(\mathbf{I})$ . Hence, (7.6) defines a linear functional in  $H_{(\alpha,\beta)}^{-s}(\mathbf{I})$ . ■

**Theorem 7.1** *Let  $s > -1$ ,  $\beta$  be determined by **Condition A**,  $c \in W_w^{[\min\{s, \alpha + (\alpha - \beta) + 1, \alpha + \beta + 1\}], \infty}(\mathbf{I})$  satisfying  $c(x) \geq 0$  and*

$$f \in H^{-\alpha/2}(\mathbf{I}) \cap H_{(\beta, \alpha - \beta)}^s(\mathbf{I}). \quad (7.8)$$

*Then there exists a unique solution  $u(x) = (1 - x)^{\alpha - \beta} x^\beta \phi(x)$ , with  $\phi(x) \in H_{(\alpha - \beta, \beta)}^{\alpha + \min\{s, \alpha + (\alpha - \beta) + 1 - \epsilon, \alpha + \beta + 1 - \epsilon\}}(\mathbf{I})$  for arbitrary  $\epsilon > 0$ , to*

$$\mathcal{L}_r^\alpha u(x) + c(x) u(x) = f(x), \quad x \in \mathbf{I}, \quad \text{subject to } u(0) = u(1) = 0. \quad (7.9)$$

**Proof:** The stated result is established using two steps. In Step 1 existence of a solution  $u \in H_0^{\alpha/2}(\mathbf{I})$  to (7.9) is shown. Then, in Step 2 a boot strapping argument is applied to improve the regularity of  $u$ .

Step 1: For  $f$  satisfying (7.8), from [14], there exists a unique solution  $u \in H_0^{\alpha/2}(\mathbf{I})$  to (7.9).

Step 2: For  $u \in H_0^{\alpha/2}(\mathbf{I})$ , from [18, Theorem 1.2.16],  $u(x) = (1 - x)^{\alpha/2} x^{\alpha/2} g(x)$ , where  $g \in L^2(\mathbf{I})$ . Then, as for  $0 < \epsilon \leq \min\{\alpha - \beta, \beta\}$

$$\int_0^1 (1 - x)^{-1 + \epsilon} x^{-1 + \epsilon} (u(x))^2 dx = \int_0^1 (1 - x)^{-1 + \epsilon + \alpha} x^{-1 + \epsilon + \alpha} (g(x))^2 dx < \int_0^1 (g(x))^2 dx < \infty,$$

it follows that  $u \in H_{(-1 + \epsilon, -1 + \epsilon)}^0(\mathbf{I}) \subset H_{(\beta - 1, \alpha - \beta - 1)}^0(\mathbf{I}) \subset H_{(\beta, \alpha - \beta)}^0(\mathbf{I})$ , and using Lemma 7.1 (with the association  $g(x) = u(x) \in H_{(\beta, \alpha - \beta)}^0(\mathbf{I})$ ,  $f(x) = c(x) \in W_w^{0, \infty}(\mathbf{I})$ )

$$c(x) u(x) \in H_{(\beta, \alpha - \beta)}^0(\mathbf{I}). \quad (7.10)$$

Using (7.10), the solution  $u$  of (7.9) satisfies

$$\mathcal{L}_r^\alpha u(x) = f(x) - c(x) u(x) := f_1(x) \in H_{(\beta, \alpha - \beta)}^{\min\{s, 0\}}(\mathbf{I}).$$

From Corollary 5.1 it follows that

$$u(x) = (1-x)^{\alpha-\beta} x^\beta \phi_1(x), \text{ where } \phi_1 \in H_{(\alpha-\beta, \beta)}^{\alpha + \min\{s, 0\}}(\mathbf{I}).$$

Using Theorem 6.4 (with its parameters  $s, \mu, p, \sigma$  replaced by  $\min\{s+\alpha, \alpha\}, \alpha-\beta, \alpha-\beta, \beta$ , respectively; and in the second instance, with its parameters  $s, \mu, p, \sigma$  replaced by  $\min\{s+\alpha, \alpha\}, \beta, \beta, \alpha-\beta$ , respectively) we have that, for arbitrary  $\epsilon > 0$ ,  $u \in H_{(\beta, \alpha-\beta)}^{\min\{s+\alpha, \alpha, \alpha+(\alpha-\beta)+1-\epsilon, \alpha+\beta+1-\epsilon\}}(\mathbf{I})$  and using Lemma 7.1,

$$c(x)u(x) \in H_{(\beta, \alpha-\beta)}^{\min\{s, \alpha, \alpha+(\alpha-\beta)+1-\epsilon, \alpha+\beta+1-\epsilon\}}(\mathbf{I}) \quad (7.11)$$

Again, using that the solution  $u$  of (7.9) satisfies

$$\mathcal{L}_r^\alpha u(x) = f(x) - c(x)u(x) := f_2(x) \in H_{(\beta, \alpha-\beta)}^{\min\{s, \alpha, \alpha+(\alpha-\beta)+1-\epsilon, \alpha+\beta+1-\epsilon\}}(\mathbf{I}),$$

and from Corollary 5.1,

$$u(x) = (1-x)^{\alpha-\beta} x^\beta \phi_2(x), \text{ where } \phi_2 \in H_{(\alpha-\beta, \beta)}^{\alpha + \min\{s, \alpha, \alpha+(\alpha-\beta)+1-\epsilon, \alpha+\beta+1-\epsilon\}}(\mathbf{I}).$$

Using Theorem 6.4 (with its parameters  $s, \mu, p, \sigma$  replaced by  $\min\{s+\alpha, 2\alpha\}, \alpha-\beta, \alpha-\beta, \beta$ , respectively; and in the second instance, with its parameters  $s, \mu, p, \sigma$  replaced by  $\min\{s+\alpha, 2\alpha\}, \beta, \beta, \alpha-\beta$ , respectively), and Lemma 7.1,

$$c(x)u(x) \in H_{(\beta, \alpha-\beta)}^{\min\{s, 2\alpha, \alpha+(\alpha-\beta)+1-\epsilon, \alpha+\beta+1-\epsilon\}}(\mathbf{I}),$$

from which it then follows that

$$u(x) = (1-x)^{\alpha-\beta} x^\beta \phi_2(x), \text{ where } \phi_2 \in H_{(\alpha-\beta, \beta)}^{\alpha + \min\{s, 2\alpha, \alpha+(\alpha-\beta)+1-\epsilon, \alpha+\beta+1-\epsilon\}}(\mathbf{I}).$$

Noting that  $4\alpha \geq \min\{\alpha + (\alpha - \beta) + 1, \alpha + \beta + 1\}$  for  $1 < \alpha < 2$ , repeating the boot strapping argument two more times establishes the stated result. ■

The inclusion of an advection term can significantly reduced the regularity of the solution.

**Theorem 7.2** *Let  $s > -1$ ,  $\beta$  be determined by **Condition A**,*

*$b \in W_w^{\max\{1, \lceil \min\{s, \alpha + (\alpha - \beta) - 1, \alpha + \beta - 1\} \rceil, \infty\}}(\mathbf{I})$ ,  $c \in W_w^{\lceil \min\{s, \alpha + (\alpha - \beta) - 1, \alpha + \beta - 1\} \rceil, \infty}(\mathbf{I})$  satisfying  $c(x) - 1/2Db(x) \geq 0$ , and*

$$f \in H^{-\alpha/2}(\mathbf{I}) \cap H_{(\beta, \alpha-\beta)}^s(\mathbf{I}). \quad (7.12)$$

*Then there exists a unique solution  $u(x) = (1-x)^{\alpha-\beta} x^\beta \phi(x)$ , with  $\phi(x) \in H_{(\alpha-\beta, \beta)}^{\alpha + \min\{s, \alpha + (\alpha - \beta) - 1 - \epsilon, \alpha + \beta - 1 - \epsilon\}}(\mathbf{I})$  for arbitrary  $\epsilon > 0$ , to*

$$\mathcal{L}_r^\alpha u(x) + b(x)Du(x) + c(x)u(x) = f(x), \quad x \in \mathbf{I}, \quad \text{subject to } u(0) = u(1) = 0. \quad (7.13)$$

**Proof:** The proof follows the same two steps as in Theorem . Step 1, establishing the existence of a solution is exactly the same. In Step 2 the boot strapping argument is applied  $m$  times, where  $m$  is the least integer such that  $m(\alpha - 1) \geq \min\{\alpha + (\alpha - \beta), \alpha + \beta\}$ , to obtain the stated result.

Step 2: For  $u \in H_0^{\alpha/2}(\mathbf{I})$  and  $0 < \epsilon \leq \min\{\alpha - \beta, \beta\}$ ,  $u \in H_{(-1+\epsilon, -1+\epsilon)}^0(\mathbf{I}) \subset H_{(\beta-1, \alpha-\beta-1)}^0(\mathbf{I})$ . Then using Lemma 4.5,  $Du \in H_{(\beta, \alpha-\beta)}^{-1}(\mathbf{I})$ . Hence we have using (7.10) and (7.7),

$$c(x)u(x) \in H_{(\beta, \alpha-\beta)}^0(\mathbf{I}) \quad \text{and} \quad b(x)Du(x) \in H_{(\beta, \alpha-\beta)}^{-1}(\mathbf{I}).$$

This leads to the solution of (7.13) satisfying

$$\mathcal{L}_r^\alpha u(x) = f(x) - b(x)Du(x) - c(x)u(x) := f_1(x) \in H_{(\beta, \alpha-\beta)}^{\min\{s, -1\}}(\mathbf{I}).$$

From Corollary 5.1 it follows that

$$u(x) = (1-x)^{\alpha-\beta} x^\beta \phi_1(x), \quad \text{where } \phi_1 \in H_{(\alpha-\beta, \beta)}^{\alpha + \min\{s, -1\}}(\mathbf{I}).$$

Using Theorem 6.4 (with its parameters  $s, \mu, p, \sigma$  replaced by  $\min\{s + \alpha, \alpha - 1\}$ ,  $\alpha - \beta$ ,  $\alpha - \beta$ ,  $\beta - 1$ , respectively; and in the second instance, with its parameters  $s, \mu, p, \sigma$  replaced by  $\min\{s + \alpha, \alpha - 1\}$ ,  $\beta$ ,  $\beta$ ,  $\alpha - \beta - 1$ , respectively) we have that, for arbitrary  $\epsilon > 0$ ,  $u \in H_{(\beta-1, \alpha-\beta-1)}^{\min\{s+\alpha, \alpha-1, \alpha+(\alpha-\beta)-\epsilon, \alpha+\beta-\epsilon\}}(\mathbf{I}) = H_{(\beta-1, \alpha-\beta-1)}^{\min\{s+\alpha, \alpha-1\}}(\mathbf{I})$  and using Lemmas 7.1 and 4.5

$$Du(x) \in H_{(\beta, \alpha-\beta)}^{-1+\min\{s+\alpha, \alpha-1\}}(\mathbf{I}), \quad c(x)u(x) \in H_{(\beta-1, \alpha-\beta-1)}^{\min\{s, \alpha-1\}}(\mathbf{I}) \quad \text{and} \quad b(x)Du(x) \in H_{(\beta, \alpha-\beta)}^{\min\{s, \alpha-2\}}(\mathbf{I}). \quad (7.14)$$

The solution  $u$  of (7.9) then must satisfies

$$\mathcal{L}_r^\alpha u(x) = f(x) - b(x)Du(x) - c(x)u(x) := f_2(x) \in H_{(\beta, \alpha-\beta)}^{\min\{s, \alpha-2\}}(\mathbf{I}),$$

and from Corollary 5.1,

$$u(x) = (1-x)^{\alpha-\beta} x^\beta \phi_2(x), \quad \text{where } \phi_2 \in H_{(\alpha-\beta, \beta)}^{\alpha + \min\{s, \alpha-2\}}(\mathbf{I}).$$

Using Theorem 6.4 (with its parameters  $s, \mu, p, \sigma$  replaced by  $\min\{s + \alpha, 2\alpha - 2\}$ ,  $\alpha - \beta$ ,  $\alpha - \beta$ ,  $\beta - 1$ , respectively; and in the second instance, with its parameters  $s, \mu, p, \sigma$  replaced by  $\min\{s + \alpha, 2\alpha - 2\}$ ,  $\beta$ ,  $\beta$ ,  $\alpha - \beta - 1$ , respectively), we have that  $u \in H_{(\beta-1, \alpha-\beta-1)}^{\min\{s+\alpha, 2\alpha-2, \alpha+(\alpha-\beta)-\epsilon, \alpha+\beta-\epsilon\}}(\mathbf{I})$ , and  $Du(x) \in H_{(\beta, \alpha-\beta)}^{-1+\min\{s+\alpha, 2\alpha-2, \alpha+(\alpha-\beta)-\epsilon, \alpha+\beta-\epsilon\}}(\mathbf{I})$ ,

$$c(x)u(x) \in H_{(\beta-1, \alpha-\beta-1)}^{\min\{s, 2\alpha-2, \alpha+(\alpha-\beta)-\epsilon, \alpha+\beta-\epsilon\}}(\mathbf{I}) \quad \text{and} \quad b(x)Du(x) \in H_{(\beta, \alpha-\beta)-1}^{\min\{s, 2\alpha-3, \alpha+(\alpha-\beta)-1-\epsilon, \alpha+\beta-1-\epsilon\}}(\mathbf{I}). \quad (7.15)$$

The solution  $u$  of (7.9) then must satisfies

$$\mathcal{L}_r^\alpha u(x) = f(x) - b(x)Du(x) - c(x)u(x) := f_2(x) \in H_{(\beta, \alpha-\beta)}^{\min\{s, 2\alpha-3, \alpha+(\alpha-\beta)-1-\epsilon, \alpha+\beta-1-\epsilon\}}(\mathbf{I}),$$

and from Corollary 5.1,

$$u(x) = (1-x)^{\alpha-\beta} x^\beta \phi_3(x), \quad \text{where } \phi_3 \in H_{(\alpha-\beta, \beta)}^{\alpha + \min\{s, 2\alpha-3, \alpha+(\alpha-\beta)-1-\epsilon, \alpha+\beta-1-\epsilon\}}(\mathbf{I}).$$

Repeatedly applying this boot stepping procedure we obtain after  $(m-2)$  additional steps  $u(x) = (1-x)^{\alpha-\beta} x^\beta \phi_{m+1}(x)$ , where

$$\phi_{m+1} \in H_{(\alpha-\beta, \beta)}^{\alpha + \min\{s, m\alpha-(m+1), \alpha+(\alpha-\beta)-1-\epsilon, \alpha+\beta-1-\epsilon\}}(\mathbf{I}) = H_{(\alpha-\beta, \beta)}^{\alpha + \min\{s, \alpha+(\alpha-\beta)-1-\epsilon, \alpha+\beta-1-\epsilon\}}(\mathbf{I}).$$

■

## 8 Regularity of the solution to the fractional diffusion, advection, reaction equation in unweighted Hilbert spaces

To connect the regularity results for the solution of (1.3), given in Theorems 7.1 and 7.2 to the usual (unweighted) Hilbert spaces we use four steps. Step 1 uses Theorem 6.4 to determine  $H_{(\sigma)}^t(\mathbf{J})$  such that  $u_0(x) = x^\beta \psi_0(x) \in H_{(\sigma)}^t(\mathbf{J})$ , for  $\psi_0(x) \in H_{(\beta)}^{s*}(\mathbf{J})$ . Step 2 applies an embedding theorem (Corollary 8.1) to then obtain  $u_0 \in H^{v_0}(\mathbf{J})$ . Step 3 repeats Steps 1 and 2 for  $u_1(x) = x^{\alpha-\beta} \psi_1(x)$ , for  $\psi_1(x) \in H_{(\alpha-\beta)}^{s*}(\mathbf{J})$  to obtain  $u_1 \in H^{v_1}(\mathbf{J})$ . The final step combines Steps 2 and 3 to conclude that  $u \in H^{\min\{v_0, v_1\}}(\mathbf{I})$ .

To begin we introduce the space  $W_{(c,d)}^{s,2}(\mathbf{I}) := \{f : f \text{ is measurable and } \|f\|_{W_{(c,d)}^{s,2}(\mathbf{I})} < \infty\}$ , where

$$\|f\|_{W_{(c,d)}^{s,2}(\mathbf{I})}^2 := \begin{cases} \sum_{j=0}^s \|D^j f\|_{L_{(c,d)}^2(\mathbf{I})}^2, & \text{for } s \in \mathbb{N}_0 \\ \sum_{j=0}^{\lfloor s \rfloor} \|D^j f\|_{L_{(c,d)}^2(\mathbf{I})}^2 + |f|_{W_{(c,d)}^{s,2}(\mathbf{I})}^2, & \text{for } s \in \mathbb{R}^+ \setminus \mathbb{N}_0 \end{cases},$$

$$\text{for } |f|_{W_{(c,d)}^{s,2}(\mathbf{I})}^2 := \iint_{\tilde{\Lambda}} (1-x)^c x^d \frac{|D^{\lfloor s \rfloor} f(x) - D^{\lfloor s \rfloor} f(y)|^2}{|x-y|^{1+2(s-\lfloor s \rfloor)}} dy dx,$$

and  $\tilde{\Lambda}$  as defined in (6.1).

Following (6.5), also introduce  $W_{(\delta)}^{s,2}(\mathbf{J}) := \{f : f \text{ is measurable and } \|f\|_{W_{(\delta)}^{s,2}(\mathbf{J})} < \infty\}$ , where

$$\|f\|_{W_{(\delta)}^{s,2}(\mathbf{J})}^2 := \begin{cases} \sum_{j=0}^s \|D^j f\|_{L_{(\delta)}^2(\mathbf{J})}^2, & \text{for } s \in \mathbb{N}_0 \\ \sum_{j=0}^{\lfloor s \rfloor} \|D^j f\|_{L_{(\delta)}^2(\mathbf{J})}^2 + |f|_{W_{(\delta)}^{s,2}(\mathbf{J})}^2, & \text{for } s \in \mathbb{R}^+ \setminus \mathbb{N}_0 \end{cases},$$

$$\text{and } |f|_{W_{(\delta)}^{s,2}(\mathbf{J})}^2 := \iint_{\Lambda} x^\delta \frac{|D^{\lfloor s \rfloor} f(x) - D^{\lfloor s \rfloor} f(y)|^2}{|x-y|^{1+2(s-\lfloor s \rfloor)}} dy dx + \iint_{\Lambda_1} x^\delta \frac{|D^{\lfloor s \rfloor} f(x) - D^{\lfloor s \rfloor} f(y)|^2}{|x-y|^{1+2(s-\lfloor s \rfloor)}} dy dx.$$

From [9] we have the following embedding result.

**Theorem 8.1** [9, See Theorem 1.d.2] *Let  $\mu, \sigma > -1$ , and  $v, w$  be two real numbers such that  $0 \leq v \leq w$ . Then, if*

$$\begin{cases} v - \frac{\sigma}{2} < w - \frac{\mu}{2} \\ \text{or} \\ v - \frac{\sigma}{2} = w - \frac{\mu}{2} \text{ with } w - \frac{\mu}{2} - \frac{1}{2} \notin \mathbb{N} \end{cases}, \quad (8.1)$$

*we have  $W_{(\mu)}^{w,2}(\mathbf{J}) \subset W_{(\sigma)}^{v,2}(\mathbf{J})$ .*

**Corollary 8.1** *Let  $\gamma > -1$ , and  $v, w$  be two real numbers such that  $0 \leq v \leq w$ . Then, if*

$$\begin{cases} v < \frac{w-\gamma}{2} \\ \text{or} \\ v = \frac{w-\gamma}{2} \text{ with } \frac{w-\gamma}{2} - \frac{1}{2} \notin \mathbb{N} \end{cases}, \quad (8.2)$$

*we have  $H_{(\gamma)}^w(\mathbf{J}) \subset H^v(\mathbf{J})$ .*

**Proof:** From [29, Theorem 3.3], it follows that  $H_{(a,b)}^s(\mathbf{I})$  and  $W_{(a+s,b+s)}^{s,2}(\mathbf{I})$  are equivalent spaces, as are  $H_{(\gamma)}^s(\mathbf{J})$  and  $W_{(\gamma+s)}^{s,2}(\mathbf{J})$ . Using that  $H^r(\mathbf{I})$  and  $W_{(0,0)}^{r,2}(\mathbf{I})$  are equivalent spaces, the stated result follows from Theorem 8.1 for  $\sigma = 0$  and  $\mu = w + \gamma$ . ■

**Corollary 8.2 (See Corollary 5.1.)** *Let  $s \geq -\alpha$ ,  $f \in H_{(\beta, \alpha-\beta)}^s(\mathbf{I})$ , and  $s^* := s + \alpha$ . Then the unique solution to (1.1), (1.2), satisfies for any  $\epsilon > 0$*

$$u \in H^{\min\{\frac{s^*+(\alpha-\beta)}{2}, \frac{s^*+\beta}{2}, (\alpha-\beta)+\frac{1}{2}-\epsilon, \beta+\frac{1}{2}-\epsilon\}}(\mathbf{I}). \quad (8.3)$$

In particular, for  $s > -\alpha + 1 + \min\{(\alpha - \beta), \beta\}$ ,

$$u \in H^{\min\{(\alpha-\beta), \beta\}+\frac{1}{2}-\epsilon}(\mathbf{I}). \quad (8.4)$$

**Proof:** Proceeding as described at the beginning of this section, consider  $u_0(x) = x^\beta \psi_0(x)$ , for  $\psi_0(x) \in H_{(\beta)}^{s^*}(\mathbf{J})$ . Using Theorem 6.4, the most regular (i.e., “nicest”) weighted Sobolev space that  $u_0$  lies in is given by the largest value for  $t$  and the smallest value for  $\sigma$  such that the conditions stated in (6.28) are satisfied. To apply Theorem 6.4 in this case we have:  $s \rightarrow s^*$ ,  $\mu \rightarrow \beta$ ,  $p \rightarrow \beta$ . Equation (6.28) then require that  $\sigma$  and  $t$  satisfy

$$0 \leq t \leq s^*, \quad \sigma \geq -\beta, \quad \sigma > t - 2\beta - 1, \quad \sigma \geq s^* - t - \beta. \quad (8.5)$$

Two cases arise for consideration.

Case 1. If  $s^* < \beta + 1$  then  $t$  and  $\sigma$  satisfying (8.5) are determined by:  $0 \leq t \leq s^*$ , and  $\sigma \geq -\beta$  (see Figure 8.1).

With the choices  $t = s^*$ ,  $\sigma = -\beta$ , using Corollary 8.1 we obtain

$$u_0 \in H_{(-\beta)}^{s^*}(\mathbf{J}) \subset H_{(-\beta)}^{\frac{s^*+\beta}{2}}(\mathbf{J}). \quad (8.6)$$

Case 2. If  $s^* \geq \beta + 1$  then  $t$  and  $\sigma$  satisfying (8.5) are determined by:  $0 \leq t \leq s^*$ ,  $\sigma > t - 2\beta - 1$ , and  $\sigma \geq s^* - t - \beta$  (see Figure 8.2). With  $t \leq s^*$  and  $t - \sigma < 2\beta + 1$ , using Corollary 8.1 we obtain, for  $\epsilon > 0$ ,

$$u_0 \in H_{(\sigma)}^t(\mathbf{J}) \subset H^{\beta+\frac{1}{2}-\epsilon}(\mathbf{J}). \quad (8.7)$$

Combining (8.6) and (8.7) yields

$$u_0 \in H^{\min\{\frac{s^*+\beta}{2}, \beta+\frac{1}{2}-\epsilon\}}(\mathbf{J}). \quad (8.8)$$

For  $u_1(x) = x^{\alpha-\beta} \psi_1(x)$ , , for  $\psi_1(x) \in H_{(\alpha-\beta)}^{s^*}(\mathbf{J})$ , a similar analysis leads to

$$u_1 \in H^{\min\{\frac{s^*+(\alpha-\beta)}{2}, (\alpha-\beta)+\frac{1}{2}-\epsilon\}}(\mathbf{J}). \quad (8.9)$$

Combining (8.8) and (8.9) we obtain

$$u \in H^{\min\{\frac{s^*+(\alpha-\beta)}{2}, \frac{s^*+\beta}{2}, (\alpha-\beta)+\frac{1}{2}-\epsilon, \beta+\frac{1}{2}-\epsilon\}}(\mathbf{I}). \quad (8.10)$$




Figure 8.1: Illustration of Case 1 for constraints (8.5).




Figure 8.2: Illustration of Case 2 for constraints (8.5).

Noting that for  $s > -\alpha + 1 + \min\{(\alpha - \beta), \beta\}$  that  $s^* > \min\{(\alpha - \beta), \beta\} + 1$ , from (8.10),

$$u \in H^{\min\{(\alpha - \beta), \beta\} + \frac{1}{2} - \epsilon}(\mathbf{I}).$$

■

Corresponding to Theorem 7.1 we have the following.

**Corollary 8.3 (See Theorem 7.1.)** *Assuming the hypothesis of Theorem 7.1 are satisfied, and let  $s^* := \alpha + \min\{s, \alpha + (\alpha - \beta) + 1, \alpha + \beta + 1\}$ . Then the unique solution of (7.9) satisfies for any  $\epsilon > 0$*

$$u \in H^{\min\{\frac{s^* + (\alpha - \beta)}{2}, \frac{s^* + \beta}{2}, (\alpha - \beta) + \frac{1}{2} - \epsilon, \beta + \frac{1}{2} - \epsilon\}}(\mathbf{I}). \quad (8.11)$$

In particular, for  $s > -\alpha + 1 + \min\{(\alpha - \beta), \beta\}$ ,

$$u \in H^{\min\{(\alpha - \beta), \beta\} + \frac{1}{2} - \epsilon}(\mathbf{I}). \quad (8.12)$$

**Proof:** Proof follows exactly as that for Corollary 8.2.

■

Corresponding to Theorem 7.2 we have the following.

**Corollary 8.4 (See Theorem 7.2.)** *Assuming the hypothesis of Theorem 7.2 are satisfied, and let  $s^* := \alpha + \min\{s, \alpha + (\alpha - \beta) - 1, \alpha + \beta - 1\}$ . Then the unique solution of (7.13) satisfies for any  $\epsilon > 0$*

$$u \in H^{\min\{\frac{s^* + (\alpha - \beta)}{2}, \frac{s^* + \beta}{2}, (\alpha - \beta) + \frac{1}{2} - \epsilon, \beta + \frac{1}{2} - \epsilon\}}(\mathbf{I}). \quad (8.13)$$

In particular, for  $s > -\alpha + 1 + \min\{(\alpha - \beta), \beta\}$ ,

$$u \in H^{\min\{(\alpha - \beta), \beta\} + \frac{1}{2} - \epsilon}(\mathbf{I}). \quad (8.14)$$

**Proof:** Proof follows exactly as that for Corollary 8.2. ■

**Remark:** If the regularity of the right hand side function  $f$  is further restricted then the regularity of the solution may be improved. For example, if  $r = 1/2$  the operator  $\mathcal{L}_{1/2}^\alpha(\cdot)$  corresponds to the integral fractional Laplacian operator. For this operator Hao and Zhang in [21] showed that for  $f \in H^s(\mathbb{I})$  the solution of (7.13) satisfied  $u \in H^{\min\{s+\alpha, \alpha/2 + 1/2 - \epsilon\}}(\mathbb{I})$  for any  $\epsilon > 0$ .

## References

- [1] M. Abramowitz and I.A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [2] G. Acosta and J.P. Borthagaray. A fractional Laplace equation: Regularity of solutions and finite element approximations. *SIAM J. Numer. Anal.*, 55(2):472–495, 2017.
- [3] G. Acosta, J.P. Borthagaray, O. Bruno, and M. Maas. Regularity theory and high order numerical methods for the (1-d)-fractional Laplacian. *Math. Comp.*, 87:1821–1857, 2018.
- [4] G. Albanese, A. Fiscella, and E. Valdinoci. Gevrey regularity for integro-differential operators. *J. Math. Anal. Appl.*, 428(2):1225–1238, 2015.
- [5] H. Antil and C.N. Rautenberg. Sobolev spaces with non-Muckenhoupt weights, fractional elliptic operators, and applications. *SIAM J. Math. Anal.*, 51(3):2479–2503, 2019.
- [6] I. Babuška and B. Guo. Direct and inverse approximation theorems for the  $p$ -version of the finite element method in the framework of weighted Besov spaces. I. Approximability of functions in the weighted Besov spaces. *SIAM J. Numer. Anal.*, 39(5):1512–1538, 2001/02.
- [7] D. A. Benson, S.W. Wheatcraft, and M.M. Meerschaert. The fractional-order governing equation of Lévy motion. *Water Resour. Res.*, 36(6):1413–1424, 2000.
- [8] C. Bernardi, M. Dauge, and Y. Maday. Polynomials in the Sobolev world. Preprint IRMAR 07-14, Université de Rennes 1, (2007).
- [9] C. Bernardi, M. Dauge, and Y. Maday. Polynomials in weighted Sobolev spaces: basics and trace liftings. Internal Report 92039, Laboratoire Jacques-Louis Lions, Université Pierre et Marie Curie, Paris (1992).
- [10] A. Buades, B. Coll, and J.M. Morel. Image denoising methods. A new nonlocal principle. *SIAM Rev.*, 52(1):113–147, 2010. Reprint of “A review of image denoising algorithms, with a new one” [MR2162865].
- [11] M. Cozzi. Interior regularity of solutions of non-local equations in Sobolev and Nikol’skii spaces. *Ann. Mat. Pura Appl. (4)*, 196(2):555–578, 2017.
- [12] J.H. Cushman and T.R. Ginn. Nonlocal dispersion in media with continuously evolving scales of heterogeneity. *Transport Porous Media*, 13:123–138, 1993.

- [13] V.J. Ervin, N. Heuer, and J.P. Roop. Regularity of the solution to 1-D fractional order diffusion equations. *Math. Comp.*, 87:2273–2294, 2018.
- [14] V.J. Ervin and J.P. Roop. Variational formulation for the stationary fractional advection dispersion equation. *Numer. Methods Partial Differential Equations*, 22(3):558–576, 2006.
- [15] G. Fdez-Manin and R. Munoz-Sola. Polynomial approximation of some singular solutions in weighted Sobolev space. In A.V. Ilin and L.R. Scott, editors, *Proceedings of the Third International Conference on Spectral and Higher Order Methods*. Houston Journal of Mathematics, Houston Texas, 1995.
- [16] P. Gatto and J.S. Hesthaven. Numerical approximation of the fractional Laplacian via *hp*-finite elements, with an application to image denoising. *J. Sci. Comput.*, 65(1):249–270, 2015.
- [17] G. Gilboa and S. Osher. Nonlocal operators with applications to image processing. *Multiscale Model. Simul.*, 7(3):1005–1028, 2008.
- [18] P. Grisvard. *Singularities in boundary value problems*, volume 22 of *Recherches en Mathématiques Appliquées [Research in Applied Mathematics]*. Masson, Paris; Springer-Verlag, Berlin, 1992.
- [19] G. Grubb. Fractional Laplacians on domains, a development of Hörmander’s theory of  $\mu$ -transmission pseudodifferential operators. *Adv. Math.*, 268:478–528, 2015.
- [20] B.-y. Guo and L.-l. Wang. Jacobi approximations in non-uniformly Jacobi-weighted Sobolev spaces. *J. Approx. Theory*, 128(1):1–41, 2004.
- [21] Z. Hao, Guang L., and Z. Zhang. Error estimates of a spectral Petrov-Galerkin method for two-sided fractional reaction-diffusion equations. *Appl. Math. Comput.*, 374:125045, 13, 2020.
- [22] Z. Hao and Z. Zhang. Optimal regularity and error estimates of a spectral Galerkin method for fractional advection-diffusion-reaction equations. *SIAM J. Numer. Anal.*, 58(1):211–233, 2020.
- [23] L. Jia, H. Chen, and V.J. Ervin. Existence and regularity of solutions to 1-D fractional order diffusion equations. *Electron. J. Differential Equations*, 93:1–21, 2019.
- [24] R. Kress. *Linear integral equations*, volume 82 of *Applied Mathematical Sciences*. Springer-Verlag, Berlin, 1989.
- [25] F. Mainardi. Fractional calculus: Some basic problems in continuum and statistical mechanics. In *Fractals and fractional calculus in continuum mechanics (Udine, 1996)*, volume 378 of *CISM Courses and Lectures*, pages 291–348. Springer, Vienna, 1997.
- [26] F. Mainardi. *Fractional calculus and waves in linear viscoelasticity*. Imperial College Press, London, 2010. An introduction to mathematical models.
- [27] Z. Mao, S. Chen, and J. Shen. Efficient and accurate spectral method using generalized Jacobi functions for solving Riesz fractional differential equations. *Appl. Numer. Math.*, 106:165–181, 2016.
- [28] Z. Mao and G. Em Karniadakis. A spectral method (of exponential convergence) for singular solutions of the diffusion equation with general two-sided fractional derivative. *SIAM J. Numer. Anal.*, 56(1):24–49, 2018.



- [29] S. Nicaise. Jacobi polynomials, weighted Sobolev spaces and approximation results of some singularities. *Math. Nachr.*, 213:117–140, 2000.
- [30] J.P. Roop. *Variational solution of the fractional advection dispersion equation*. ProQuest LLC, Ann Arbor, MI, 2004. Thesis (Ph.D.)–Clemson University.
- [31] X. Ros-Oton and J. Serra. The Dirichlet problem for the fractional Laplacian: regularity up to the boundary. *J. Math. Pures Appl. (9)*, 101(3):275–302, 2014.
- [32] W. Rudin. *Principles of mathematical analysis*. McGraw-Hill Book Co., New York-Auckland-Düsseldorf, third edition, 1976. International Series in Pure and Applied Mathematics.
- [33] J. Schöberl. A short introduction to interpolation spaces. Preprint: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.725.2183&rep=rep1&type=pdf>, 2012.
- [34] M. F. Shlesinger, B. J. West, and J. Klafter. Lévy dynamics of enhanced diffusion: Application to turbulence. *Phys. Rev. Lett.*, 58(11):1100–1103, 1987.
- [35] G. Szegő. *Orthogonal polynomials*. American Mathematical Society, Providence, R.I., fourth edition, 1975. American Mathematical Society, Colloquium Publications, Vol. XXIII.
- [36] L. Tartar. *An introduction to Sobolev spaces and interpolation spaces*, volume 3 of *Lecture Notes of the Unione Matematica Italiana*. Springer, Berlin; UMI, Bologna, 2007.
- [37] G. M. Zaslavsky, D. Stevens, and H. Weitzner. Self-similar transport in incomplete chaos. *Phys. Rev. E (3)*, 48(3):1683–1694, 1993.
- [38] Z. Zhang. Error estimates of spectral Galerkin methods for a linear fractional reaction-diffusion equation. *J. Sci. Comput.*, 78(2):1087–1110, 2019.