

# Linkage of sets of Quaternion Algebras in characteristic 2

Adam Chapman

*School of Computer Science, Academic College of Tel-Aviv-Yaffo, Rabenu Yeruham St., P.O.B 8401 Yaffo, 6818211, Israel*

---

## Abstract

This note contains two new observations on the linkage properties of quaternion algebras over fields of characteristic 2: first, that a 3-linked field need not be 4-linked (a case which was left open in previous papers) and that three inseparably linked quaternion algebras are also cyclically linked when the base-field is odd-closed.

*Keywords:* Quaternion Algebras; Linkage; Symbol Algebras; Brauer Groups  
*2010 MSC:* 16K20 (primary); 11E81, 11E04 (secondary)

---

## 1. Introduction

A symbol  $p$ -algebra of prime degree  $p$  over a field  $F$  of  $\text{char}(F) = p$  is an algebra of the form  $[\alpha, \beta]_{p,F} = F\langle x, y : x^p - x = \alpha, y^p = \beta, yxy^{-1} = x + 1 \rangle$  for some  $\alpha \in F$  and  $\beta \in F^\times$ . The significance of these algebras derives from the fact (proven by Teichmüller, see [11, Section 9]) that  ${}_p\text{Br}(F)$  is generated by the Brauer classes of these algebras. Symbol  $p$ -algebras of degree 2 over a field  $F$  of  $\text{char}(F) = 2$  are the quaternion algebras over that field.

We say that symbol  $p$ -algebras  $A_1, \dots, A_m$  of degree  $p$  over  $F$  are linked if there exists a degree  $p$  field extension  $K/F$  that splits them all. We say they are inseparably linked if there exists a purely inseparable degree  $p$  field extension  $K = F[\sqrt[p]{\gamma}]$  that splits them all, and we say these algebras are cyclically linked if there exists a degree  $p$  cyclic extension  $K = F[\lambda : \lambda^p - \lambda = \delta]$  which splits them all. It was proven in [3] that if  $A_1$  and  $A_2$  are inseparably linked then they are also cyclically linked and counterexamples were provided for the converse statement. The special case of quaternion algebras had been proven earlier in [8], [12] and [9]. Note that quadratic field extensions are either cyclic or purely inseparable, and therefore if quaternion algebras  $A_1, \dots, A_m$  are linked then they are either cyclically or inseparably linked.

We say a field  $F$  is  $m$ -linked if any  $m$  quaternion algebras over  $F$  are linked. Examples of 2-linked fields that are not 3-linked exist:  $\mathbb{F}_2((\alpha))((\beta))$  is 2-linked by [1] but not 3-linked by [5]. The question of whether 3-linked fields which are not 4-linked exist was raised in [2]. Examples were provided in [7] for  $\text{char}(F) \neq 2$ . Similar questions in characteristic 2 were studied in [4], but this specific problem remained open because

---

*Email address:* adam1chapman@yahoo.com (Adam Chapman)

the tools used in those previous works proved inadequate. In this note we complete the picture by proving the existence of 3-linked fields of characteristic 2 which are not 4-linked. The tool is the  $w$ -invariant for valued division algebras defined and studied by Tignol in [14] and other works. We also extend the result from [3] from pairs of symbol  $p$ -algebras to triples, showing that if three symbol  $p$ -algebras of prime degree  $p$  over a  $p$ -special field  $F$  of characteristic  $p$  are inseparably linked then they are also cyclically linked.

## 2. 3-Linked Fields which are not 4-Linked

Let  $F = k((\alpha))((\beta))$  be the field of iterated Laurent series in two variables over a field  $k$  of  $\text{char}(k) = 2$ . Write  $v$  for the right-to-left  $(\alpha, \beta)$ -adic valuation on  $F$  and write  $\Gamma_F = \mathbb{Z} \times \mathbb{Z}$  for the underlying value group. This is a henselian valuation, and therefore it extends uniquely to any field extension  $K/F$  by the formula  $v(t) = \frac{1}{d}v(N(t))$  where  $N$  is the norm form  $K \rightarrow F$  and  $d = [K : F]$ . Consequently, the valuation extends to any central division algebra  $D/F$  with the analogous formula  $v(t) = \frac{1}{d}v(N(t))$  where  $N$  here is the reduced norm  $D \rightarrow F$  and  $d$  is the degree of  $D$  (see [15] for background on valuation theory for division algebras). Following [14], for any cyclic quadratic field extension  $K$  of  $F$ , we define  $w(K)$  to be  $\min\{v(\text{Tr}(t)) - v(t) : t \in K\}$  where  $\text{Tr} : K \rightarrow F$  is the trace map, given by  $\text{Tr}(ax + b) = a$  where  $K = F[\varphi^{-1}(\gamma)] = F[x : x^2 + x = \gamma]$ . Similarly, for any quaternion division algebra  $D$  over  $F$ , we define  $w(D)$  to be  $\min\{v(\text{Tr}(t)) - v(t) : t \in D\}$ , where  $\text{Tr} : D \rightarrow F$  is the reduced trace map given by  $\text{Tr}(a + bx + cy + xy) = b$  where  $D = [\gamma, \delta]_{2,F} = F\langle x, y : x^2 + x = \gamma, y^2 = \delta, yxy^{-1} = x + 1 \rangle$ . Note that for any quaternion division algebra  $D$  over  $F$  containing a quadratic cyclic extension  $K$  of  $F$ , the restriction of the reduced trace map on  $D$  to  $K$  coincides with the trace map on  $K$ , and therefore  $w(K) \geq w(D)$ .

**Lemma 2.1** ([14, Theorem 1.8, with  $p = 2$ ]). *If  $\gamma$  is an element in  $F$  with  $v(\gamma) < \vec{0}$  and  $v(\gamma) \notin 2\Gamma_F$ , then  $K = F[t : t^2 + t = \gamma]$  is a totally ramified quadratic cyclic field extension of  $F$  with  $w(K) = -v(t) = -\frac{1}{2}v(\gamma)$ .*

**Lemma 2.2.** *Consider the quaternion algebras  $[\alpha^{-1}, \beta]_{2,F}$ ,  $[\beta^{-1}, \alpha]_{2,F}$  and  $[\alpha^{-1}\beta^{-1}, \beta]_{2,F}$  over  $F = k((\alpha))((\beta))$  where  $k$  is a field of  $\text{char}(k) = 2$ . If a field  $K$  is a quadratic extension of  $F$  that splits these algebras, then  $w(K) \leq (\frac{1}{2}, \frac{1}{2})$ .*

*Proof.* These algebras do not share any quadratic inseparable splitting field by [4]. Note that they satisfy the conditions of [14, Proposition 2.1], and therefore they are division algebras. Suppose  $K$  is a common quadratic splitting field of theirs, and therefore it must be a cyclic extension of  $F$ . Since  $K$  is a quadratic splitting field of  $[\alpha^{-1}, \beta]_{2,F} = F\langle x, y : x^2 + x = \alpha^{-1}, y^2 = \beta, yxy^{-1} = x + 1 \rangle$ , it is also a subfield of that algebra, and since it is a cyclic extension of  $F$ , it must be generated over  $F$  by an element of trace 1, i.e. an element  $t$  of the form  $t = \lambda + x + ay + bxy$  for some  $\lambda, a, b \in F$ . Since  $x + ay + bxy$  generates  $K$  too and it is of trace 1, we can suppose  $\lambda = 0$  and take  $t = x + ay + bxy$  to be the generator of  $K$ . Now,  $v(x) = (-\frac{1}{2}, 0)$ ,  $v(y) = (0, \frac{1}{2})$  and  $v(xy) = (-\frac{1}{2}, \frac{1}{2})$ . Since they belong to three distinct classes modulo  $\Gamma_F$ , we have  $v(t) = \min\{v(x), v(ay), v(bxy)\}$ . Hence, if  $v(t) < (-\frac{1}{2}, -\frac{1}{2})$  then  $v(t) < (-\frac{1}{2}, 0)$  and so  $v(t) \neq v(x)$ , which means  $v(t) = \min\{v(ay), v(bxy)\}$ , and therefore the class

of  $v(t)$  modulo  $\Gamma_F$  is either  $(\frac{1}{2}, \frac{1}{2})$  or  $(0, \frac{1}{2})$ . Since  $w(K) = -v(t)$  by Lemma 2.1, if  $w(K) > (\frac{1}{2}, \frac{1}{2})$  then the class of  $w(K)$  modulo  $\Gamma_F$  is either  $(\frac{1}{2}, \frac{1}{2})$  or  $(0, \frac{1}{2})$ . Recalling that  $K$  is also a quadratic splitting field of  $[\beta^{-1}, \alpha]_{2,F}$ , the same argument shows that if  $w(K) > (\frac{1}{2}, \frac{1}{2})$  then the class of  $w(K)$  modulo  $\Gamma_F$  is either  $(\frac{1}{2}, \frac{1}{2})$  or  $(\frac{1}{2}, 0)$ . Similarly, since  $K$  is a quadratic splitting field of  $[\alpha^{-1}\beta^{-1}, \beta]_{2,F}$ , if  $w(K) > (\frac{1}{2}, \frac{1}{2})$  then the class of  $w(K)$  modulo  $\Gamma_F$  is either  $(0, \frac{1}{2})$  or  $(\frac{1}{2}, 0)$ . However, the intersection of  $\{(\frac{1}{2}, \frac{1}{2}), (0, \frac{1}{2})\}$ ,  $\{(\frac{1}{2}, \frac{1}{2}), (\frac{1}{2}, 0)\}$  and  $\{(\frac{1}{2}, 0), (0, \frac{1}{2})\}$  is trivial, and so it is impossible that  $w(K) > (\frac{1}{2}, \frac{1}{2})$ , which means that  $w(K) \leq (\frac{1}{2}, \frac{1}{2})$ .  $\square$

**Proposition 2.3.** *The algebras  $[\alpha^{-1}, \beta]_{2,F}$ ,  $[\beta^{-1}, \alpha]_{2,F}$ ,  $[\alpha^{-1}\beta^{-1}, \beta]_{2,F}$  and  $[\alpha^{-2}\beta^{-1}, \alpha]_{2,F}$  are not linked.*

*Proof.* All the quadratic splitting fields  $K$  the first three share satisfy  $w(K) \leq (\frac{1}{2}, \frac{1}{2})$ . On the other hand, by [14, Proposition 2.1]  $[\alpha^{-2}\beta^{-1}, \alpha]_{2,F}$  is a division algebra with  $w([\alpha^{-2}\beta^{-1}, \alpha]_{2,F}) = (1, \frac{1}{2})$ , which is greater than  $(\frac{1}{2}, \frac{1}{2})$ , so every quadratic splitting field  $K$  of  $[\alpha^{-2}\beta^{-1}, \alpha]_{2,F}$  has  $w(K) \geq w([\alpha^{-2}\beta^{-1}, \alpha]_{2,F}) > (\frac{1}{2}, \frac{1}{2})$ . Therefore there is no common quadratic splitting field of the first three algebras which is also a splitting field of the fourth.  $\square$

**Example 2.4.** The field  $F = k((\alpha))((\beta))$  is 3-linked when  $k$  is algebraically closed (e.g.,  $k = \mathbb{F}_2^{sep}$ ). The reason is that given any three quaternion algebras  $[\gamma_1, \delta_1]_{2,F}$ ,  $[\gamma_2, \delta_2]_{2,F}$  and  $[\gamma_3, \delta_3]_{2,F}$ , the system

$$\begin{aligned} b^2\gamma_1 + (c_1^2 + c_1d_1 + d_1^2\gamma_1)\delta_1 &= a_2^2 + a_2b + b^2\gamma_2 + (c_2^2 + c_2d_2 + d_2^2\gamma_2)\delta_2 \\ b^2\gamma_1 + (c_1^2 + c_1d_1 + d_1^2\gamma_1)\delta_1 &= a_3^2 + a_3b + b^2\gamma_3 + (c_3^2 + c_3d_3 + d_3^2\gamma_3)\delta_3 \end{aligned}$$

is a system of two homogeneous quadratic equations in 9 variables

$$a_2, a_3, b, c_1, c_2, c_3, d_1, d_2, d_3,$$

and since  $F$  is a  $C_2$ -field (see [13] and [10, Chapter 6]) and  $9 > 2 \cdot 2^2$ , the system has a nontrivial solution. If this nontrivial solution has  $b = 0$ , then  $F[\sqrt{(c_1^2 + c_1d_1 + d_1^2\gamma_1)\delta_1}]$  is a common inseparable splitting field of the three quaternion algebras, and if  $b \neq 0$  then  $F[t : t^2 + t = \gamma_1 + (\frac{c_1^2}{b^2} + \frac{c_1d_1}{b^2} + \frac{d_1^2}{b^2}\gamma_1)\delta_1]$  is a common cyclic splitting field of the three quaternion algebras. However,  $F$  is not 4-linked by Proposition 2.3. This provides a characteristic 2 analogue of [7].

### 3. Cyclic and Inseparable 3-Linkage of Symbol $p$ -Algebras

In [3] it was proven that if two symbol  $p$ -algebras of degree  $p$  over a field  $F$  of  $\text{char}(F) = p$  are inseparably linked then they are also cyclically linked. Here we prove that the same holds for three algebras, under the assumption that  $F$  is  $p$ -special, i.e. has no finite field extensions of degree prime to  $p$ .

**Theorem 3.1.** *Given a prime integer  $p$  and a field  $F$  of  $\text{char}(F) = p$ , if three symbol  $p$ -algebras of degree  $p$  over  $F$  are inseparably linked then they become cyclically linked under a prime-to- $p$  extension of  $F$  of degree at most  $2p - 1$ .*

*Proof.* The algebras are inseparably linked, so they can be written as  $A_1 = [\alpha_1, \beta]_{p,F}$ ,  $A_2 = [\alpha_2, \beta]_{p,F}$  and  $A_3 = [\alpha_3, \beta]_{p,F}$ . We can suppose that  $\langle [A_1], [A_2], [A_3] \rangle$  is a subgroup of  ${}_p\text{Br}(F)$  of order  $p^3$ , because otherwise it is enough to stress that two of them are cyclically linked (which is true by [3]) and the third shares the same cyclic field extension of  $F$  shared by the first two. This means that  $\alpha_1, \alpha_2, \alpha_3 \notin \wp(F) = \{\lambda^p - \lambda : \lambda \in F\}$  and in particular they are nonzero. Moreover,  $\alpha_2 \notin \mathbb{F}_p \alpha_1$ , and so  $(\frac{\alpha_1}{\alpha_2})^{p-1} \neq 1$ .

Now,  $A_i$  can also be written as  $[\alpha_i^p, \beta]_{p,F}$ , and a solution to the following system will show the algebras are cyclically linked:

$$\begin{aligned}\alpha_1^p + \beta(u_1^p - u_1 v_1^{p-1} + \alpha_1^p v_1^p) &= \alpha_2^p + \beta(u_2^p - u_2 v_2^{p-1} + \alpha_2^p v_2^p) \\ \alpha_1^p + \beta(u_1^p - u_1 v_1^{p-1} + \alpha_1^p v_1^p) &= \alpha_3^p + \beta(u_3^p - u_3 v_3^{p-1} + \alpha_3^p v_3^p).\end{aligned}$$

Take  $u_1 = u_2 = u_3$ , and then the equations reduce to

$$\begin{aligned}\alpha_1^p + \beta(-u_1 v_1^{p-1} + \alpha_1^p v_1^p) &= \alpha_2^p + \beta(-u_2 v_2^{p-1} + \alpha_2^p v_2^p) \\ \alpha_1^p + \beta(-u_1 v_1^{p-1} + \alpha_1^p v_1^p) &= \alpha_3^p + \beta(-u_3 v_3^{p-1} + \alpha_3^p v_3^p).\end{aligned}$$

and then isolate  $u_1$  in both equations

$$\begin{aligned}\beta(u_1 v_2^{p-1} - u_1 v_1^{p-1}) &= \alpha_2^p - \alpha_1^p + \beta(\alpha_2^p v_2^p - \alpha_1^p v_1^p) \\ \beta(u_1 v_3^{p-1} - u_1 v_1^{p-1}) &= \alpha_3^p - \alpha_1^p + \beta(\alpha_3^p v_3^p - \alpha_1^p v_1^p).\end{aligned}$$

Take  $v_2 = \frac{\alpha_1}{\alpha_2} v_1$  and  $v_3 = 1$ , and then

$$\begin{aligned}\beta\left(\left(\frac{\alpha_1}{\alpha_2}\right)^{p-1} - 1\right)u_1 v_1^{p-1} &= \alpha_2^p - \alpha_1^p \\ \beta(u_1 - u_1 v_1^{p-1}) &= \alpha_3^p - \alpha_1^p + \beta(\alpha_3^p - \alpha_1^p v_1^p).\end{aligned}$$

Plugging in the first equation in the second gives

$$\beta u_1 - \frac{\alpha_2^p - \alpha_1^p}{(\frac{\alpha_1}{\alpha_2})^{p-1} - 1} = \alpha_3^p - \alpha_1^p + \beta(\alpha_3^p - \alpha_1^p v_1^p), \quad \text{and so}$$

$$\beta u_1 = \alpha_3^p - \alpha_1^p + \frac{\alpha_2^p - \alpha_1^p}{(\frac{\alpha_1}{\alpha_2})^{p-1} - 1} + \beta(\alpha_3^p - \alpha_1^p v_1^p).$$

Plugging this into the first equation gives

$$(\alpha_3^p - \alpha_1^p + \frac{\alpha_2^p - \alpha_1^p}{(\frac{\alpha_1}{\alpha_2})^{p-1} - 1} + \beta(\alpha_3^p - \alpha_1^p v_1^p))\left(\left(\frac{\alpha_1}{\alpha_2}\right)^{p-1} - 1\right)v_1^{p-1} = \alpha_2^p - \alpha_1^p,$$

a degree  $2p - 1$  equation in one variable  $v_1$ , that must have a solution under some prime-to- $p$  extension of  $F$  of degree up to  $2p - 1$ .  $\square$

**Corollary 3.2.**

1. For  $p$ -special fields, inseparable linkage of three symbol  $p$ -algebras of degree  $p$  implies cyclic linkage.
2. When  $p = 2$ , every three inseparably linked quadratic 2-fold Pfister forms over  $F$  are either cyclically linked or become cyclically linked over a degree 3 extension of  $F$ .

Note that three cyclically linked symbol  $p$ -algebras need not be inseparably linked. For example, the algebras  $[\alpha^{-1}, \beta_1]_{p,F}$ ,  $[\alpha^{-1}, \beta_2]_{p,F}$  and  $[\alpha^{-1}, \beta_3]_{p,F}$  over  $F = k(\langle\langle\alpha\rangle\rangle\langle\langle\beta_1\rangle\rangle\langle\langle\beta_2\rangle\rangle\langle\langle\beta_3\rangle\rangle)$  (or over  $F_0 = k(\alpha, \beta_1, \beta_2, \beta_3)$ ) are pair-wise not inseparably linked by the same argument as in [3, Example 4.2].

When  $p = 2$ , these algebras remain pair-wise not inseparably linked under any odd field extension of  $F$ . The reason is that  $[\alpha^{-1}, \beta_i]_{2,K}$  and  $[\alpha^{-1}, \beta_j]_{2,K}$  are inseparably linked if and only if the 3-fold Pfister form  $\langle\langle\beta_i, \beta_j, \alpha^{-1}\rangle\rangle_K$  is hyperbolic by [6], and since  $\langle\langle\beta_i, \beta_j, \alpha^{-1}\rangle\rangle_F$  is anisotropic, it remains anisotropic under scalar extension to any odd degree extension  $K/F$ . Thus, if  $K$  is the odd closure of  $F$ , then the quaternion algebras  $[\alpha^{-1}, \beta_1]_{2,K}$ ,  $[\alpha^{-1}, \beta_2]_{2,K}$  and  $[\alpha^{-1}, \beta_3]_{2,K}$  are not inseparably linked.

## Acknowledgements

The author thanks Jean-Pierre Tignol for his comments on the manuscript.

## References

- [1] R. Aravire and B. Jacob.  $p$ -algebras over maximally complete fields. In *K-theory and algebraic geometry: connections with quadratic forms and division algebras* (Santa Barbara, CA, 1992), volume 58 of *Proc. Sympos. Pure Math.*, pages 27–49. Amer. Math. Soc., Providence, RI, 1995. With an appendix by Jean-Pierre Tignol.
- [2] K. J. Becher. Triple linkage. *Ann. K-Theory*, 3(3):369–378, 2018.
- [3] A. Chapman. Common subfields of  $p$ -algebras of prime degree. *Bull. Belg. Math. Soc. Simon Stevin*, 22(4):683–686, 2015.
- [4] A. Chapman. Common slots of bilinear and quadratic Pfister forms. *Bull. Aust. Math. Soc.*, 98(1):38–47, 2018.
- [5] A. Chapman, A. Dolphin, and D. B. Leep. Triple linkage of quadratic Pfister forms. *Manuscripta Math.*, 157(3-4):435–443, 2018.
- [6] A. Chapman, S. Gilat, and U. Vishne. Linkage of quadratic Pfister forms. *Comm. Algebra*, 45(12):5212–5226, 2017.
- [7] A. Chapman and J.-P. Tignol. Linkage of Pfister forms over  $\mathbb{C}(x_1, \dots, x_n)$ . *Ann. K-Theory*, 4(3):521–524, 2019.
- [8] P. Draxl. *Skew Fields*, volume 81 of *London Math. Soc. Lect. Notes*. Cambridge University Press, 1983.

- [9] A. Elduque and O. Villa. A note on the linkage of Hurwitz algebras. *Manuscripta Math.*, 117(1):105–110, 2005.
- [10] A. J. Engler and A. Prestel. *Valued fields*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2005.
- [11] P. Gille and T. Szamuely. *Central simple algebras and Galois cohomology*, volume 101 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2006.
- [12] T. Y. Lam. On the linkage of quaternion algebras. *Bull. Belg. Math. Soc. Simon Stevin*, 9(3):415–418, 2002.
- [13] S. Lang. On quasi algebraic closure. *Ann. of Math. (2)*, 55:373–390, 1952.
- [14] J.-P. Tignol. Classification of wild cyclic field extensions and division algebras of prime degree over a Henselian field. In *Proceedings of the International Conference on Algebra, Part 2 (Novosibirsk, 1989)*, volume 131 of *Contemp. Math.*, pages 491–508. Amer. Math. Soc., Providence, RI, 1992.
- [15] J.-P. Tignol and A. R. Wadsworth. *Value Functions on Simple Algebras, and Associated Graded Rings*. Springer Monographs in Mathematics. Springer, 2015.