

ON THE SHAFAREVICH CONJECTURE FOR ENRIQUES SURFACES

TEPPEI TAKAMATSU

ABSTRACT. Enriques surfaces are minimal surfaces of Kodaira dimension 0 with $b_2 = 10$. If we work with a field of characteristic away from 2, Enriques surfaces admit double covers which are K3 surfaces. In this paper, we prove the Shafarevich conjecture for Enriques surfaces by reducing the problem to the case of K3 surfaces. In our formulation of the Shafarevich conjecture, we use the notion “admitting a cohomological good K3 cover”, which includes not only good reduction but also flower pot reduction.

1. INTRODUCTION

The Shafarevich conjecture for abelian varieties states the finiteness of isomorphism classes of abelian varieties of a fixed dimension over a fixed number field admitting good reduction away from a fixed finite set of finite places. The purpose of this paper is to formulate and prove this conjecture for Enriques surfaces. Our main theorem is the following.

Theorem 1.1 (Theorem 3.7). *Let F be a finitely generated field over \mathbb{Q} , and R be a finite type algebra over \mathbb{Z} which is a normal domain with the fraction field F . Then, the set*

$$\left\{ X \left| \begin{array}{l} X : \text{Enriques surface over } F, \\ \text{for any height 1 prime ideal } \mathfrak{p} \in \text{Spec } R, \\ X \text{ admits a cohomological good K3 cover at } \mathfrak{p} \end{array} \right. \right\} / F\text{-isom}$$

is finite.

Here, we say that X admits a cohomological good K3 cover at \mathfrak{p} if there exists a K3 double cover of X whose ℓ -adic cohomologies are unramified at \mathfrak{p} (see Definition 3.2). In particular, we prove the original Shafarevich conjecture for Enriques surfaces over finitely generated fields over \mathbb{Q} .

Corollary 1.2 (Corollary 3.8). *Let F be a finitely generated field over \mathbb{Q} , and R be a finite type algebra over \mathbb{Z} which is a normal domain with the fraction field F . Then, the set*

$$\left\{ X \left| \begin{array}{l} X : \text{Enriques surface over } F, \\ X \text{ admits good reduction at any height 1 prime ideal } \mathfrak{p} \in \text{Spec } R \end{array} \right. \right\} / F\text{-isom}$$

is finite.

Date: April 8, 2024.

2010 Mathematics Subject Classification. Primary 14J28; Secondary 11G35.

Key words and phrases. Enriques surfaces, Shafarevich conjecture.

Theorem 1.1 is, in fact, a direct corollary of the Shafarevich conjecture for K3 surfaces ([She17], [Tak18]). To conclude Corollary 1.2 from Theorem 1.1, we recall the properties of good reduction of Enriques surfaces (see Proposition 2.4 and Lemma 3.4). We shall make some remarks on the statement of Theorem 1.1. It is known that there exists an Enriques surface over a local field which does not admit good reduction, but whose K3 double cover admits good reduction (see [LM18, Theorem 3.6]). Our Theorem 1.1 generalizes the original Shafarevich conjecture (Corollary 1.2) to deal with such phenomena.

Acknowledgments. The author is deeply grateful to his advisor Naoki Imai for his deep encouragement and helpful advice. The author also thanks Tetsushi Ito for the proof of Proposition 2.4, Yohsuke Matsuzawa for helpful suggestions, and Yuya Matsumoto for pointing out my misunderstandings about cohomologies of K3 double covers (Remark 3.3).

2. ENRIQUES SURFACES AND GOOD REDUCTION

First, we recall the definition of Enriques surfaces.

Definition 2.1. For any field k , an *Enriques surface over k* is a smooth projective surface over k satisfying $\omega_{X_{\bar{k}}/\bar{k}} \equiv \mathcal{O}_{X_{\bar{k}}}$ and $b_2(X_{\bar{k}}) = 10$, where \equiv denotes numerical equivalence.

If k is a field of characteristic away from 2, it is well-known that an Enriques surface over k has a K3 double cover (see [Băd01, Proposition 10.14]).

The main purpose of this paper is to study the Shafarevich conjecture for Enriques surfaces by reducing the problem to the case of K3 surfaces. In the rest of this section, we will define the good reduction of Enriques surfaces, and study their properties.

Definition 2.2. Let K be a discrete valuation field, and X be an Enriques surface over K . We say X admits good reduction if there exists a smooth proper algebraic space \mathcal{X} over \mathcal{O}_K such that $\mathcal{X}_K \simeq X$.

First, we recall the lifting property of line bundles on Enriques surfaces, which was proved by Liedtke.

Lemma 2.3. *Let R be a complete discrete valuation ring with the residue field k . Let $\mathcal{X} \rightarrow \mathrm{Spf} R$ be a formal deformation of Enriques surfaces, i.e. proper smooth formal scheme with the special fiber \mathcal{X}_k which is an Enriques surface. If $\mathcal{L} \in \mathrm{Pic}(\mathcal{X}_k)$, then $\mathcal{L}^{\otimes 2}$ extends to $\mathrm{Pic}(\mathcal{X})$. Moreover, if $H^2(\mathcal{X}_k, \mathcal{O}) = 0$ (e.g. the case of $\mathrm{char} k \neq 2$), then \mathcal{L} extends to $\mathrm{Pic}(\mathcal{X})$.*

Proof. This is a special case of [Lie15, Proposition 4.4 (5)]. We note that Liedtke's argument also works for the case where k is not algebraically closed. For the sake of completeness, we recall the proof. We put $\mathcal{X} = \mathrm{colim}_n \mathcal{X}_n$, where \mathcal{X}_n is a smooth proper scheme over $S_n := \mathrm{Spec}(R/\mathfrak{m}^n)$. Here, \mathfrak{m} is the maximal ideal of R . Let \tilde{R} be a strict Henselization of R , and $\tilde{\mathfrak{m}}$ be the maximal ideal of \tilde{R} . Let $\tilde{\mathcal{X}}_n := (\mathcal{X}_n)_{\tilde{S}_n}$ be the base change. By the Hochschild-Serre spectral sequence with respect to the Galois covering $\tilde{\mathcal{X}}_n \rightarrow \mathcal{X}_n$, we have the exact sequence

$$0 \rightarrow H^1(G_k, H^0(\tilde{\mathcal{X}}_n, \mathcal{O}^\times)) \rightarrow \mathrm{Pic}(\mathcal{X}_n) \rightarrow \mathrm{Pic}(\tilde{\mathcal{X}}_n)^{G_k} \rightarrow H^2(G_k, H^0(\tilde{\mathcal{X}}_n, \mathcal{O}^\times)),$$

where G_k is an absolute Galois group of k . Note that the left end is 0, and the right end is no other than the Brauer group $\mathrm{Br}(R/\mathfrak{m}^n) := H_{\mathrm{et}}^2(\mathrm{Spec} R/\mathfrak{m}^n, \mathbb{G}_m)$. We recall that there exists the natural isomorphisms $\mathrm{Br}(R/\mathfrak{m}^n) \simeq \mathrm{Br}(k)$ (see [Gro95a, COROLLAIRE 6.2], [Gro95b, COROLLAIRE 2.5]). By [Lie15, Proposition 4.4 (1)-(3)], for any positive integer n , $\mathcal{L}^{\otimes 2}$ lifts to a rational point of the Picard scheme $\mathcal{M}_n \in \mathrm{Pic}_{\mathcal{X}_n/S_n}(S_n)$, which gives an element of $\mathrm{Pic}(\tilde{\mathcal{X}}_n)^{G_k}$. By the above exact sequence, the line bundle \mathcal{M}_n descends to the line bundle on \mathcal{X}_n if and only if its image in $\mathrm{Br}(R/\mathfrak{m}^n)$ is zero. If $n = 1$, this image is zero, thus for any n , it is zero. Therefore, \mathcal{M}_n is an extension of $\mathcal{L}^{\otimes 2}$ to $\mathrm{Pic}(\mathcal{X})$. The final statement follows from [FGI⁺05, Corollary 8.5.5]. \square

Tetsushi Ito taught me the following proposition.

Proposition 2.4. *Let K be a complete discrete valuation field, X be an Enriques surface over K . The following are equivalent.*

- (1) X admits good reduction.
- (2) There exists a smooth projective scheme \mathcal{X} over \mathcal{O}_K such that $\mathcal{X}_k \simeq X$.
- (3) There exists a smooth proper scheme \mathcal{X} over \mathcal{O}_K such that $\mathcal{X}_k \simeq X$.

Proof. Clearly (2) \Rightarrow (3) \Rightarrow (1). So we shall prove (1) \Rightarrow (2). Let k be a residue field of K . Take a smooth proper algebraic space model \mathcal{X} over \mathcal{O}_K . Let \mathcal{X}_k be the special fiber of \mathcal{X} , and $\hat{\mathcal{X}}$ be the completion of \mathcal{X} along the special fiber \mathcal{X}_k . By Lemma 2.3, there exists an ample line bundle of \mathcal{X}_k which can be lifted to a line bundle on \mathcal{X} . Therefore by [FGI⁺05, Corollary 8.5.6], we get the desired smooth projective model. \square

Remark 2.5. As in [Mat15, Example 5.2], the above proposition does not hold in the case of K3 surfaces.

3. THE SHAFAREVICH CONJECTURE FOR ENRIQUES SURFACES

First, we recall the definition of a K3 double cover to introduce a notion which generalizes good reduction of Enriques surfaces (see also [CL16, Section 7]).

Definition 3.1. Let k be a field with characteristic away from 2, and X a K3 surface over k . By fixing an isomorphism $\omega_{X/k}^{\otimes 2} \simeq \mathcal{O}_X$, we can equip the \mathcal{O}_X -module $\mathcal{O}_X \oplus \omega_{X/k}$ with an \mathcal{O}_X -algebra structure. We refer to a relative spectrum $\tilde{X} := \mathrm{Spec}_X(\mathcal{O}_X \oplus \omega_{X/k})$ as a K3 double cover of X .

Definition 3.2. Let K be a discrete valuation field with the residual characteristic $p \neq 2$, and let X be an Enriques surface over K . We say X admits a cohomological good K3 cover if there exists a K3 double cover \tilde{X} of X such that $H_{\mathrm{et}}^2(\tilde{X}_{\bar{K}}, \mathbb{Q}_{\ell})$ is an unramified representation, for $\ell \neq p$ (which is independent of ℓ as in [Tak18, Lemma 5.0.1]). Yuya Matsumoto taught me that this definition depends on the choice of \tilde{X} (see Remark 3.3).

Remark 3.3. Let $\iota_1, \iota_2: \omega_{X/K}^{\otimes 2} \simeq \mathcal{O}_X$ be isomorphisms satisfying that $\iota_1 \circ \iota_2^{-1}: \mathcal{O}_X \rightarrow \mathcal{O}_X$ sends the global section 1 to $\alpha \in K$. Let $\tilde{X}_i \rightarrow X$ be K3 double covers defined by ι_1 and ι_2 . For $\sigma \in \mathrm{Gal}(\bar{K}/K)$, let $\sigma_i: \tilde{X}_{i,\bar{K}} \rightarrow \tilde{X}_{i,\bar{K}}$ be the Galois action defined by the K -structure \tilde{X}_i . By the definition of a K3 double cover, there exists an isomorphism $\tilde{X}_{1,K[\sqrt{\alpha}]} \simeq \tilde{X}_{2,K[\sqrt{\alpha}]}$ over $X_{K[\sqrt{\alpha}]}$. Via this isomorphism, for any σ with $\sigma(\sqrt{\alpha}) = -\sqrt{\alpha}$,

σ_1 is equal to $\tau \circ \sigma_2$, where τ is the Enriques involution. So if the order of α is odd, then at most one of $H_{\text{ét}}^2(\tilde{X}_i, \mathbb{Q}_\ell)$ ($i = 1, 2$) is unramified. We also note that if the order of α is even, then the unramifiedness is preserved.

Lemma 3.4. *Let K be a discrete valuation field with the residual characteristic $p \neq 2$, and let X be an Enriques surface over K . If X admits good reduction, then X admits a cohomological good K3 cover.*

Proof. Taking a completion, we may assume K is complete (see Remark 3.3). By Proposition 2.4, one can take a smooth projective scheme model \mathcal{X} over \mathcal{O}_K . By the separatedness of the Picard functor [BLR90, Theorem 8.4.3], we have an isomorphism $\omega_{\mathcal{X}/\mathcal{O}_K}^{\otimes 2} \simeq \mathcal{O}_{\mathcal{X}}$. Then we can attach an $\mathcal{O}_{\mathcal{X}}$ -algebra structure on the $\mathcal{O}_{\mathcal{X}}$ -module $\mathcal{O}_{\mathcal{X}} \oplus \omega_{\mathcal{X}/\mathcal{O}_K}$. Then a double covering $\tilde{\mathcal{X}} := \text{Spec}_{\mathcal{X}}(\mathcal{O}_{\mathcal{X}} \oplus \omega_{\mathcal{X}/\mathcal{O}_K})$ is finite étale over \mathcal{X} by the assumption on the residual characteristic. So $\tilde{\mathcal{X}}$ gives a smooth projective scheme model of \tilde{X}_K , which is a K3 double cover of X . \square

Remark 3.5. There exists an Enriques surface which does not admit good reduction, but whose K3 double cover admits good reduction. See [LM18, Theorem 3.6] for such an example having flower pot reduction.

Lemma 3.6. *Let k be a field of characteristic away from 2. Let Y be a K3 surface over k . Then there exists only finitely many k -isomorphism classes of Enriques surfaces which is a $\mathbb{Z}/2\mathbb{Z}$ -quotient of Y .*

Proof. It suffices to show the finiteness of conjugacy classes of finite order elements in $\text{Aut}_F(Y)$. By [BLvL19, Proposition 3.10], there exists a group homomorphism $\text{Aut}_F(Y) \rtimes R_Y \rightarrow \text{O}(\text{Pic} Y)$ which has finite kernel and image of finite index, where R_Y is the Galois fixed part of the Weyl group of $Y_{\bar{F}}$. Here, by [Bor63, Section 5, (a)], there exist only finitely many conjugacy classes of finite order elements in $\text{O}(\text{Pic} Y)$. So the desired property of $\text{Aut}_F(Y)$ follows from [Oha07, Lemma 1.4]. \square

Theorem 3.7. *Let F be a finitely generated field over \mathbb{Q} , and R be a finite type algebra over \mathbb{Z} which is a normal domain with the fraction field F . Then, the set*

$$\text{Shaf} := \left\{ X \left| \begin{array}{l} X: \text{Enriques surface over } F, \\ \text{for any height 1 prime ideal } \mathfrak{p} \in \text{Spec } R, \\ X \text{ admits a cohomological good K3 cover at } \mathfrak{p} \end{array} \right. \right\} / F\text{-isom}$$

is finite.

Proof. First, we note that $\text{Pic}(\text{Spec } R)$ is finitely generated ([Lan83, Chapter 2, Theorem 7.6]). Let $\mathcal{O}(D_1), \dots, \mathcal{O}(D_r)$ be its generators, where D_i are Cartier divisors. Then by removing $\text{Supp}(D_i)$, we may assume that $\text{Pic}(\text{Spec } R)$ is trivial. Moreover, by shrinking $\text{Spec } R$ again, we may assume $1/2 \in R$.

Next, we will show that for any $X \in \text{Shaf}$, there exists a K3 double cover \tilde{X} such that for any height 1 prime $\mathfrak{p} \in \text{Spec } R$, $H_{\text{ét}}^2(X_{\bar{F}}, \mathbb{Q}_\ell)$ is unramified at \mathfrak{p} for $\ell \notin \mathfrak{p}$. Take an element $X \in \text{Shaf}$. Fix an isomorphism $\iota: \omega_{X/F}^{\otimes 2} \simeq \mathcal{O}_X$, and let \tilde{X}' be a corresponding K3 double cover. One can take a finite set S of height 1 prime ideals of $\text{Spec } R$ such that for any $\mathfrak{p} \notin S$, $H_{\text{ét}}^2(\tilde{X}', \mathbb{Q}_\ell)$ is unramified for $\ell \notin \mathfrak{p}$. Moreover, for any height 1 prime $\mathfrak{p} \in \text{Spec } R$, there exists an isomorphism $\iota_{\mathfrak{p}}: \omega_{X/F}^{\otimes 2} \simeq \mathcal{O}_X$ such that the corresponding K3 double cover has ℓ -adic cohomologies which are unramified at \mathfrak{p} . Suppose that $\iota_{\mathfrak{p}} \circ \iota^{-1}: \mathcal{O}_X \rightarrow \mathcal{O}_X$ sends the global section 1 to $\alpha_{\mathfrak{p}} \in F$. We may

assume that $\alpha_{\mathfrak{p}} = 1$ for $\mathfrak{p} \notin S$. Since $\text{Pic}(\text{Spec } R)$ is trivial, there exists $\alpha \in F$ satisfying that $\text{ord}_{\mathfrak{p}}(\alpha) = \text{ord}_{\mathfrak{p}}(\alpha_{\mathfrak{p}})$ for any height 1 prime $\mathfrak{p} \in \text{Spec } R$. Then the composition $\alpha \circ \iota: \omega_{X/F}^{\otimes 2} \simeq \mathcal{O}_X$ induces a desired K3 double cover \tilde{X} (see Remark 3.3).

Therefore, by Lemma 3.6, we can reduce the problem to the finiteness of K3 double covers. So this theorem follows from [Tak18, Theorem 1.0.1]. \square

Corollary 3.8. *Let F be a finitely generated field over \mathbb{Q} , and R be a finite type algebra over \mathbb{Z} which is a normal domain with the fraction field F . Then, the set*

$$\left\{ X \left| \begin{array}{l} X: \text{Enriques surface over } F, \\ X \text{ admits good reduction at any height 1 prime ideal } \mathfrak{p} \in \text{Spec } R \end{array} \right. \right\} / F\text{-isom}$$

is finite.

Proof. Remark that we may assume $1/2 \in R$. Therefore, the desired finiteness follows from Lemma 3.4 and Theorem 3.7. \square

REFERENCES

- [Băd01] Lucian Bădescu, *Algebraic surfaces*, Universitext, Springer-Verlag, New York, 2001, Translated from the 1981 Romanian original by Vladimir Măşek and revised by the author.
- [BLR90] Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 21, Springer-Verlag, Berlin, 1990.
- [BLvL19] Martin Bright, Adam Logan, and Ronald van Luijk, *Finiteness results for K3 surfaces over arbitrary fields*, European Journal of Mathematics (2019).
- [Bor63] Armand Borel, *Arithmetic properties of linear algebraic groups*, Proc. Internat. Congr. Mathematicians (Stockholm, 1962), Inst. Mittag-Leffler, Djursholm, 1963, pp. 10–22.
- [CL16] Bruno Chiarellotto and Christopher Lazda, *Combinatorial degenerations of surfaces and Calabi-Yau threefolds*, Algebra Number Theory **10** (2016), no. 10, 2235–2266.
- [FGI⁺05] Barbara Fantechi, Lothar Göttsche, Luc Illusie, Steven L. Kleiman, Nitin Nitsure, and Angelo Vistoli, *Fundamental algebraic geometry*, Mathematical Surveys and Monographs, vol. 123, American Mathematical Society, Providence, RI, 2005, Grothendieck’s FGA explained.
- [Gro95a] Alexander Grothendieck, *Le groupe de Brauer. I. Algèbres d’Azumaya et interprétations diverses*, Séminaire Bourbaki, Vol. 9, Soc. Math. France, Paris, 1995, pp. Exp. No. 290, 199–219.
- [Gro95b] ———, *Le groupe de Brauer. II. Théorie cohomologique*, Séminaire Bourbaki, Vol. 9, Soc. Math. France, Paris, 1995, pp. Exp. No. 297, 287–307.
- [Lan83] Serge Lang, *Fundamentals of Diophantine geometry*, Springer-Verlag, New York, 1983.
- [Lie15] Christian Liedtke, *Arithmetic moduli and lifting of Enriques surfaces*, J. Reine Angew. Math. **706** (2015), 35–65.
- [LM18] Christian Liedtke and Yuya Matsumoto, *Good reduction of K3 surfaces*, Compos. Math. **154** (2018), no. 1, 1–35.
- [Mat15] Yuya Matsumoto, *Good reduction criterion for K3 surfaces*, Math. Z. **279** (2015), no. 1-2, 241–266.
- [Oha07] Hisanori Ohashi, *On the number of Enriques quotients of a K3 surface*, Publ. Res. Inst. Math. Sci. **43** (2007), no. 1, 181–200.
- [She17] Yiwei She, *The unpolarized Shafarevich Conjecture for K3 Surfaces*, preprint (2017), arXiv:1705.09038.
- [Tak18] Teppei Takamatsu, *On a cohomological generalization of the Shafarevich conjecture for K3 surfaces*, preprint (2018), arXiv:1810.12279.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, KOMABA, TOKYO, 153-8914, JAPAN

E-mail address: teppei@ms.u-tokyo.ac.jp