

On self-similar solutions to degenerate compressible Navier-Stokes equations

Pierre GERMAIN, Tsukasa IWABUCHI, Tristan LÉGER

ABSTRACT. We study cavitating self-similar solutions to compressible Navier-Stokes equations with degenerate density-dependent viscosity. We prove both existence of expanders and non-existence of small shrinkers.

CONTENTS

1. Introduction	1
1.1. The equations	1
1.2. Background	3
1.3. Results	3
2. Existence of expanders	4
2.1. Existence when $0 < \alpha < 1$	5
2.2. Existence when $\alpha = 1$	14
3. Non-existence of shrinkers	17
3.1. Set-up	19
3.2. Basic properties of the matrix A	20
3.3. Weighted energy estimate	22
References	27

1. INTRODUCTION

1.1. The equations. In this article we consider the compressible Navier-Stokes-Fourier system, which describes the motion of a heat conducting gas:

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla \pi = \operatorname{div}(\tau), \\ \partial_t \left[\rho \left(\frac{|u|^2}{2} + e \right) \right] + \operatorname{div} \left[u \left(\rho \left(\frac{|u|^2}{2} + e \right) + \pi \right) \right] - \operatorname{div} q \\ \quad \quad \quad = \operatorname{div}(\tau \cdot u), \end{cases}$$

where $\rho : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ denotes the density of the fluid, $u : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ its velocity field, $\pi : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ its pressure, $\tau : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^{d^2}$ its stress tensor, $e : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ its

Mathematics Subject Classification (2010): 35C06, 76N10

Keywords: Compressible Navier-Stokes equations; self-similar solutions.

Acknowledgements: P. Germain was partially supported by the NSF grant DMS-1501019. T. Iwabuchi was supported by JSPS KAKENHI Grant 17H04824.

Addresses: Pierre Germain, Courant Institute of Mathematical Sciences, New York University, 251 Mercer Street, New York, N.Y. 10012-1185, USA, pgermain@cims.nyu.edu

Tsukasa Iwabuchi, Mathematical Institute, Tohoku University, Sendai City, 980-8578, JAPAN, t-iwabuchi@tohoku.ac.jp

Tristan Léger, Courant Institute of Mathematical Sciences, New York University, 251 Mercer Street, New York, N.Y. 10012-1185, USA, tleger@cims.nyu.edu

internal energy and $q : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ its internal energy flux.

We will assume that e is given by Joule's law, that is $e = C_V \theta$ (C_V is the heat constant, and $\theta(t, x)$ the temperature of the gas). We also take q to be proportional to the gradient of θ , in accordance with Fourier's law: $q = -\kappa \nabla \theta$, where $\kappa > 0$ denotes the thermal conductivity. The pressure will be given by the ideal gas law: $\pi = \rho R \theta$. We will also restrict our attention to newtonian gases, for which τ is given by

$$\tau = \lambda \operatorname{div} u \operatorname{Id} + 2\mu D(u), \quad D(u) := \left(\frac{\partial_i u_j + \partial_j u_i}{2} \right)_{1 \leq i \leq d, 1 \leq j \leq d},$$

where the Lamé coefficients λ, μ are such that

$$\mu > 0, \quad 2\mu + d\lambda \geq 0.$$

Moreover it is physically relevant (from the point of view of kinetic theory for example) to consider the coefficients λ and μ that depend on the density ρ . More precisely, we postulate the following laws:

$$\lambda(\rho) = \lambda_0 \rho^\alpha, \quad \mu(\rho) = \mu_0 \rho^\alpha, \quad (1.1)$$

where $0 < \alpha \leq 1$.

The equations read

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + \nabla(\rho R \theta) = \operatorname{div}(\lambda_0 \rho^\alpha \operatorname{div} u \operatorname{Id} + \mu_0 \rho^\alpha (\nabla u + \nabla u^T)), \\ \partial_t \left[\rho \left(\frac{|u|^2}{2} + C_V \theta \right) \right] + \operatorname{div} \left[u \left(\rho \left(\frac{|u|^2}{2} + C_V \theta \right) + \rho R \theta \right) \right] - \kappa \Delta \theta \\ \quad = \operatorname{div}(\lambda_0 \rho^\alpha (\operatorname{div} u) u + \mu_0 \rho^\alpha (\nabla u + \nabla u^T) \cdot u), \end{cases} \quad (\text{cNS})$$

Note that the system is scaling invariant: if (ρ, u, θ) denotes a solution, then $(\rho_\lambda, u_\lambda, \theta_\lambda)$, where

$$\rho_\lambda(t, x) := \rho(\lambda^2 t, \lambda x), \quad u_\lambda(t, x) := \lambda u(\lambda^2 t, \lambda x), \quad \theta_\lambda(t, x) := \lambda^2 \theta(\lambda^2 t, \lambda x)$$

is also a solution.

This leads us to considering self-similar solutions to (cNS). In this paper, we will therefore restrict our attention to self-similar, radially symmetric solutions. Moreover since a well-known difficulty in the case of degenerate viscosity is the presence of vacuum, we will also assume that our solutions exhibit cavitation at the origin. More precisely, we study both expanders of the form

$$\begin{aligned} \rho(t, x) &= P \left(\frac{|x|}{\sqrt{t}} \right), \quad u(t, x) = \frac{1}{\sqrt{t}} U \left(\frac{|x|}{\sqrt{t}} \right) \frac{x}{|x|}, \\ \theta(t, x) &= \frac{1}{t} \Theta \left(\frac{|x|}{\sqrt{t}} \right), \end{aligned} \quad (1.2)$$

and shrinkers:

$$\begin{aligned} \rho(t, x) &= P \left(\frac{|x|}{\sqrt{T-t}} \right), \quad u(t, x) = \frac{1}{\sqrt{T-t}} U \left(\frac{|x|}{\sqrt{T-t}} \right) \frac{x}{|x|}, \\ \theta(t, x) &= \frac{1}{T-t} \Theta \left(\frac{|x|}{\sqrt{T-t}} \right), \end{aligned} \quad (1.3)$$

where $T > 0$. We will require that $P(0) = 0$ to signify that there is cavitation at the origin.

1.2. Background.

1.2.1. *Weak solutions.* The system (cNS) and related models have been extensively studied in the literature. In the constant coefficient case, A. Matsumura and T. Nishida proved global existence for small data in [22]. Note that in this work the density is bounded away from 0, therefore no vacuum is present. P.-L. Lions constructed weak solutions for barotropic compressible Navier-Stokes with large initial data and possibly vacuum states in [20]. His result was subsequently refined and extended to the full system by E. Feireisl, A. Novotný and H. Petzeltová in [7] and E. Feireisl in [6].

In the case of degenerate density-dependent viscosity (for example of the type (1.1)), the situation becomes more involved. A work of major importance in this direction is the paper by D. Bresch, B. Desjardins and C.-K. Lin ([1]), where this problem is treated for various compressible models, including the shallow water model away from vacuum. Recently A. Vasseur and C. Yu were able to construct weak solutions to the barotropic compressible Navier-Stokes with vacuum in [25].

1.2.2. *Self-similar solutions.* Self-similar solutions have been the subject of investigation since the seminal work of J. Leray [18] on the incompressible Navier-Stokes equation. Indeed he noticed that the existence of a backward solution of this type would imply singularity formation. Forward self-similar solutions are also of interest since they are expected to describe the continuation of the backward solution after the singular time. Such small expander type solutions were constructed by Cannone and Planchon in [2]. Recently V. Šverák and H. Jia were able to construct solutions to the incompressible Navier-Stokes equation with large self-similar initial data in [15]. Regarding backward self-similar solutions, J. Nečas, M. Růžička, V. Šverák proved their non existence in the natural energy class of the Navier-Stokes equation. This result was improved by Tsai who showed that in fact this result still holds if only local energy inequalities are assumed.

For compressible Navier Stokes equations, Z. Guo and S. Jiang showed in [10] that in the 1D isothermal barotropic case, there exist neither forward nor backward self-similar solutions. In the case of (cNS) with constant viscosity, the first two authors constructed expanders both with and without cavitation in [8]. Finally in the companion paper [9], we prove non existence of small shrinkers, still in the constant coefficient case.

1.3. **Results.** We prove two types of results in this paper: existence of small expanders in Section 2, and non-existence of small shrinkers in Section 3. The construction of forward solutions follows the same pattern as in the paper of the first two authors [8], although the proof is more involved due to the degeneracy of the coefficients at vacuum. It requires a more precise control of the velocity profile near the origin. For technical reasons, we distinguish between the cases $0 < \alpha < 1$ and $\alpha = 1$. In the first situation ($0 < \alpha < 1$), here is simplified statement of our result:

Theorem 1.1. *Let $d \geq 3$. Assume $2\mu_0 + d\lambda_0 = 0$. Then if*

$$A + P_\delta + \delta + P_\delta^{1-\alpha}\delta^2 + \frac{P_\delta^{1-\alpha}\Theta_0}{A} + \frac{A\delta}{\Theta_0} + \frac{A^2}{\Theta_0} + P_\delta A$$

3

is small enough, there exists a solution to (cNS) of the form (1.2) such that

$$P(\delta) = P_\delta, U(0) = 0, U'(0) = A, \Theta(0) = \Theta_0 > 0, \Theta'(0) = 0.$$

In the second case ($\alpha = 1$), we prove

Theorem 1.2. *Let $d \geq 3$. Assume $2\mu_0 + d\lambda_0 > 0$. Then if $A \log \delta^{-1} + P_\delta + \delta$ is small enough, there exists a solution to (cNS) of the form (1.2) such that*

$$P(\delta) = P_\delta, U(0) = 0, U'(0) = A, \Theta(0) = \frac{2\mu_0 + \lambda_0}{R} A > 0, \Theta'(0) = 0.$$

Note that for $0 < \alpha < 1$, we construct a three parameter family of solutions, and for $\alpha = 1$ we obtain a two dimensional family (the choice of Θ_0 is not free). In both cases we also obtain a precise description of the shape of the profiles. Details can be found in the full statements of these theorems in Section 2.

In the second part of the paper, we show that (cNS) does not have small solutions of shrinker type (1.3). We prove the following result (in simplified form here):

Theorem 1.3. *Consider a cavitating solution to (cNS) of the form (1.3). Let $\varepsilon > 0$. Assume that*

$$\sup_{r>0} \left(\langle r \rangle^2 \Theta + P^{1-\alpha} + \left| \frac{U}{r\Theta} \right| \right) + \sup_{r>\varepsilon} \left| \frac{U'}{r\Theta'} \right|$$

is sufficiently small.

Then $U \equiv \Theta \equiv 0, P = \text{Constant}$.

We show a similar result in the companion paper [9], when the Lamé coefficients are constant. Note however that the two approaches are different. Indeed some key cancellations are no longer available in the density-dependent viscosity case. Therefore we develop a different method tailored to our setting of solutions exhibiting cavitation in the present work.

2. EXISTENCE OF EXPANDERS

In this section we construct expander solutions to (cNS), that is solutions of the type (1.2). Plugging this ansatz into (cNS), we obtain the following system of ODEs for the

profiles P, U and Θ :

$$\left\{ \begin{array}{l} -\frac{1}{2}rP' + P'U + P\left(U' + \frac{d-1}{r}U\right) = 0, \\ -\frac{1}{2}PU - \frac{1}{2}r(PU)' + (PU^2)' + \frac{d-1}{r}PU^2 + (PR\Theta)' \\ \quad = (2\mu_0 + \lambda_0)P^\alpha\left(U'' + \frac{d-1}{r}U' - \frac{d-1}{r^2}U\right) \\ \quad \quad + (2\mu_0 + \lambda_0)(P^\alpha)'U' + \lambda_0(P^\alpha)'\frac{d-1}{r}U, \\ -P\left(\frac{U^2}{2} + C_V\Theta\right) - \frac{1}{2}r\left(P\left(\frac{U^2}{2} + C_V\Theta\right)\right)' + \left(UP\left(\frac{U^2}{2} + C_V\Theta\right) + UPR\Theta\right)' \\ \quad + \frac{d-1}{r}\left(UP\left(\frac{U^2}{2} + C_V\Theta\right) + UPR\Theta\right) - \kappa\left(\Theta'' + \frac{d-1}{r}\Theta'\right) \\ \quad = 2\mu_0P^\alpha\left((U')^2 + \frac{d-1}{r^2}U^2\right) + \lambda_0P^\alpha\left(U' + \frac{d-1}{r}U\right)^2 \\ \quad \quad + (2\mu_0 + \lambda_0)P^\alpha\left(U'' + \frac{d-1}{r}U' - \frac{d-1}{r^2}U\right)U \\ \quad \quad + (2\mu_0 + \lambda_0)(P^\alpha)'U'U + \lambda_0(P^\alpha)'\frac{d-1}{r}U^2. \end{array} \right. \quad (2.1)$$

As announced in the introduction, we distinguish two cases for the existence of expanders: when $0 < \alpha < 1$ and when $\alpha = 1$. There are substantial differences between the two: we do not use the same integro-differential formulation for both, and require different conditions on the parameters. Most strikingly, Θ_0 is free for $0 < \alpha < 1$ but not for $\alpha = 1$.

2.1. Existence when $0 < \alpha < 1$. In this section, we show the following:

Theorem 2.1. *Let $d \geq 3$ and $0 < \alpha < 1$. Fix $(C_V, \kappa, R, \mu_0, \lambda_0) \in (0, \infty)^5$ such that $2\mu_0 + d\lambda_0 = 0$.*

Then there exists a constant $C(C_V, \kappa, R, \mu_0, \lambda_0) := C$ such that if

$$A + P_\delta + \delta + P_\delta^{1-\alpha}\delta^2 + \frac{P_\delta^{1-\alpha}\Theta_0}{A} + \frac{A\delta}{\Theta_0} + \frac{A^2}{\Theta_0} + P_\delta A < C,$$

there exists a solution $(P, U, \Theta) \in \mathcal{C}^{\frac{dA}{1/2-A}} \times \mathcal{C}^1([0, \infty)) \times \mathcal{C}^1([0, \infty))$ to (2.1) such that

$$P(0) = 0, P(\delta) = P_\delta, U(0) = 0, U'(0) = A, \Theta(0) = \Theta_0 > 0, \Theta'(0) = 0.$$

Near 0, the behavior of the solution is given by (here $0 < \varepsilon < 1 - \alpha$):

$$\begin{aligned} P(r) &= P_\delta\left(\frac{r}{\delta}\right)^{\frac{dA}{1/2-A}} + O\left(\left(\frac{r}{\delta}\right)^{1+\frac{dA}{1/2-A}+(1-\alpha-\varepsilon)dA}\right), \\ U(r) &= Ar + O(r^{1+(1-\alpha-\varepsilon)dA}), \\ \Theta(r) &= \Theta_0 + O(r^2). \end{aligned}$$

Moreover the profiles satisfy the following global bounds:

$$P(r) \simeq P_\delta \min \left[1, \left(\frac{r}{\delta} \right)^{\frac{2dA}{1-2A}} \right],$$

$$|U(r)| \lesssim \frac{Ar}{(1 + P_\delta^{1/2-\alpha/2} r)^2}, \quad |U'(r)| \lesssim \frac{A}{(1 + P_\delta^{1/2-\alpha/2} r)^2},$$

$$0 \leq \Theta(r) \lesssim \frac{1}{(1 + \sqrt{P_\delta} r)^2}, \quad |\Theta'(r)| \lesssim \frac{\sqrt{P_\delta} r}{(1 + \sqrt{P_\delta} r)^2}.$$

Finally as $r \rightarrow +\infty$, there exist $P_\infty > 0, U_\infty > 0, \Theta_\infty > 0$ such that

$$P(r) = P_\infty + O\left(\frac{1}{r^2}\right), \quad U(r) = \frac{U_\infty}{r} + O\left(\frac{1}{r^3}\right), \quad \Theta(r) = \frac{\Theta_\infty}{r^2} + O\left(\frac{1}{r^4}\right).$$

The proof is modelled after that of a similar result in [8] (Section 4): first we find an integro-differential formulation of the problem. Then we construct the solution locally near 0 using a fixed-point argument. After that, we prove global existence, and finally we study the asymptotic behavior of the solutions.

Note that the main difference compared to [8] is in the derivation of the integro-differential equation and the local existence part of the proof. These steps are more involved in the present paper due to the degeneracy of the Lamé coefficients at 0, and the presence of vacuum.

2.1.1. Integro-differential formulation. We are aiming at constructing solutions such that for $0 < A < 1/2$ and $\Theta_0 > 0$,

$$P(r) = O(r^{\frac{2dA}{1-2A}}), \quad U(r) = Ar + O(r^{1+(1-\alpha-\varepsilon)\frac{2dA}{1-2A}}), \quad \Theta(r) = \Theta_0 + O(r^2), \quad r \rightarrow 0.$$

First starting with the equation on P , we directly obtain after integration that

$$P(r) = P_\delta e^{V(r)-V(\delta)},$$

where

$$V(r) = \int_0^r \frac{U' + \frac{d-1}{r_1} U}{\frac{1}{2}r_1 - U} dr_1. \quad (2.2)$$

Next we move to the equation satisfied by U :

$$\begin{aligned} & -\frac{1}{2}PU - \frac{1}{2}r(PU)' + (PU^2)' + \frac{d-1}{r}PU^2 + (PR\Theta)' \\ & = (2\mu_0 + \lambda_0) \left\{ P^\alpha \left(U' + \frac{d-1}{r}U \right) \right\}' - 2\mu_0(P^\alpha)' \cdot \frac{d-1}{r}U. \end{aligned}$$

Using the assumption $2\mu_0 + d\lambda_0 = 0$, we can write the right-hand side

$$(2\mu_0 + \lambda_0)P^\alpha \left\{ \left(U' + \frac{d-1}{r}U \right)' + \alpha \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r - U} \left(U' - \frac{1}{r}U \right) \right\}.$$

After dividing by P^α , we have

$$\begin{aligned} & -\frac{1}{2}(rP^{1-\alpha}U)' - \frac{\alpha}{2}rP^{1-\alpha} \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r - U}U + \frac{1}{P^\alpha}(PU^2)' + \frac{d-1}{r}P^{1-\alpha}U^2 + \frac{(PR\Theta)'}{P^\alpha} \\ & = (2\mu_0 + \lambda_0) \left\{ \left(U' + \frac{d-1}{r}U \right)' + \alpha \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r - U} \left(U' - \frac{1}{r}U \right) \right\}. \end{aligned} \quad (2.3)$$

The equation (2.3) can be written

$$\begin{aligned} & -\frac{1}{2}(rP^{1-\alpha}U)' - \frac{\alpha}{2}rP^{1-\alpha}\frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r - U} + \frac{1}{P^\alpha}(PU^2)' + \frac{d-1}{r}P^{1-\alpha}U^2 + \frac{(PR\Theta)'}{P^\alpha} \\ & = (2\mu_0 + \lambda_0) \left\{ \left(U' + \frac{d-1}{r}U \right)' + \alpha \frac{U' - \frac{1}{r}U}{\frac{1}{2}r - U} \left(U' + \frac{d-1}{r}U \right) \right\}. \end{aligned}$$

Let

$$\tilde{V}(r) := \int_0^r \frac{U' - \frac{1}{\tilde{r}}U}{\frac{1}{2}\tilde{r} - U} d\tilde{r}. \quad (2.4)$$

Multiplying the above by $e^{\alpha\tilde{V}(r)}$ we get

$$\begin{aligned} & -\frac{1}{2}(e^{\alpha\tilde{V}(r)}rP^{1-\alpha}U)' + e^{\alpha\tilde{V}(r)} \left\{ -\frac{\alpha}{2}rP^{1-\alpha}\frac{\frac{d}{r}U}{\frac{1}{2}r - U} + \frac{1}{P^\alpha}(PU^2)' + \frac{d-1}{r}P^{1-\alpha}U^2 + \frac{(PR\Theta)'}{P^\alpha} \right\} \\ & = (2\mu_0 + \lambda_0) \left\{ e^{\alpha\tilde{V}(r)} \left(U' + \frac{d-1}{r}U \right) \right\}'. \end{aligned}$$

Now we integrate over $[0, r]$ and divide by $e^{\alpha\tilde{V}(r)}$:

$$\begin{aligned} & -\frac{1}{2}rP^{1-\alpha}U \\ & + e^{-\alpha\tilde{V}(r)} \int_0^r e^{\alpha\tilde{V}(r_2)} \left\{ -\frac{\alpha}{2}r_2P^{1-\alpha}\frac{\frac{d}{r_2}U}{\frac{1}{2}r_2 - U} + \frac{1}{P^\alpha}(PU^2)' + \frac{d-1}{r_2}P^{1-\alpha}U^2 + \frac{(PR\Theta)'}{P^\alpha} \right\} dr_2 \\ & = (2\mu_0 + \lambda_0) \left(U' + \frac{d-1}{r}U \right) - e^{-\alpha\tilde{V}(r)}(2\mu_0 + \lambda_0)dA. \end{aligned}$$

Let

$$W(r) := \frac{1}{2\mu_0 + \lambda_0} \int_0^r \frac{P(\tilde{r})^{1-\alpha}\tilde{r}}{2} d\tilde{r}. \quad (2.5)$$

Multiplication by $r^{d-1}e^{W(r)}$ then yields

$$(2\mu_0 + \lambda_0) \left(r^{d-1}e^{W(r)}U \right)' = r^{d-1}e^{W(r)}F_U(r),$$

where

$$\begin{aligned} F_U(r) & := e^{-\alpha\tilde{V}(r)}(2\mu_0 + \lambda_0)A \\ & + e^{-\alpha\tilde{V}(r)} \int_0^r e^{\alpha\tilde{V}(r_2)} \left\{ -\frac{\alpha}{2}r_2P^{1-\alpha}\frac{\frac{d}{r_2}U}{\frac{1}{2}r_2 - U} + \frac{1}{P^\alpha}(PU^2)' + \frac{d-1}{r_2}P^{1-\alpha}U^2 + \frac{(PR\Theta)'}{P^\alpha} \right\} dr_2. \end{aligned} \quad (2.6)$$

We obtain the desired integro-differential equation after integrating the above.

For the last equation on Θ , we proceed as in [8]. Therefore we omit the details.

We obtain the following integro-differential formulation:

$$\begin{cases} P(r) = e^{V(r)-V(\delta)} & \text{for given } \delta > 0, \\ U(r) = \frac{r^{-d+1}}{2\mu_0 + \lambda_0} \int_0^r r_1^{d-1} e^{-W(r)+W(r_1)} F_U(r_1) dr_1, \\ \Theta(r) = (d-2)r^{-d+2} \int_0^r r_1^{d-3} e^{-Z(r)+Z(r_1)} dr_1 \Theta_0 - \frac{U^2}{2C_V} + \frac{r^{-d+2}}{\kappa} \int_0^r r_1^{d-2} e^{-Z(r)+Z(r_1)} F_\Theta(r_1) dr_1, \end{cases} \quad (2.7)$$

where V, \tilde{V}, W and F_U are defined as in (2.2), (2.4), (2.5), (2.6), and

$$\begin{aligned} F_\Theta(r_1) := & UP \left(\frac{U^2}{2} + C_V \Theta \right) + UPR\Theta + \frac{d-2}{r_1} \int_0^{r_1} \left(UP \left(\frac{U^2}{2} + C_V \Theta \right) + UPR\Theta \right) dr_2 \\ & - (2\mu_0 + \lambda_0) \left(P^\alpha U U' + \frac{d-2}{r_1} \int_0^{r_1} P^\alpha U U' dr_2 \right) \\ & - \lambda_0(d-1) \left(\frac{P^\alpha U^2}{r_1} + \frac{d-2}{r_1} \int_0^{r_1} P^\alpha \frac{U^2}{r_2} dr_2 \right) \\ & + \frac{\kappa}{C_V} \left(U U' + \frac{d-2}{2r_1} U^2 \right), \end{aligned}$$

and

$$Z(r) := \frac{C_V}{\kappa} \int_0^r \frac{P(\tilde{r})\tilde{r}}{2} d\tilde{r}.$$

2.1.2. Local existence. In this section we construct a local solution to the above integro-differential formulation by a fixed point argument.

First we define the functional space in which we solve the equation:

Let $(1-\alpha)/2 < \varepsilon < 1-\alpha$. Define

$$\|(U, \Theta)\|^\delta := \sup_{0 < r < \delta} \left[r^{-1}|U(r)| + |U'(r)| + \frac{Ar(\frac{r}{\delta})^{-(1-\alpha-\varepsilon)dA}}{P_\delta^{1-\alpha}\Theta_0} \left| \left(\frac{U(r)}{r} \right)' \right| + \frac{A}{\Theta_0} |\Theta(r)| + \frac{A}{\Theta_0} r^{-1} |\Theta'(r)| \right],$$

$$E^\delta = \{(U, \Theta) \in \mathcal{C}^1(0, \delta) \text{ such that } \Theta(0) = \Theta_0, \|(U, \Theta)\|^\delta < \infty\}.$$

Note that E^δ differs from the definition in [8]. Here we need better control of the profile U near 0 to close the estimates.

We also define the map Φ on $B_{E^\delta}((Ar, \Theta_0), A/2)$ by the formula

$$\Phi : (U, \Theta) \mapsto$$

$$\left(\frac{r^{1-d}}{2\mu_0 + \lambda_0} \int_0^r e^{W(r_1)-W(r)} r_1^{d-1} F_U(r_1) dr_1, \Theta_0 + \frac{r^{2-d}}{\kappa} \int_0^r r_1^{d-2} e^{Z(r_1)-Z(r)} F_\Theta(r_1) dr_1 - \frac{U^2}{2C_V} \right).$$

The remainder of the subsection is dedicated to the proof of the following lemma, which shows local existence of solutions to (2.7).

Lemma 2.2. *Assume that*

$$A \ll \frac{1}{2}, \quad P_\delta \leq 1, \quad \delta + P_\delta^{1-\alpha} \delta^2 + \frac{P_\delta^{1-\alpha} \Theta_0}{A} + \frac{A\delta}{\Theta_0} + \frac{A^2}{\Theta_0} + P_\delta A \ll 1.$$

Then Φ is a contraction on $B_{E^\delta}((Ar, \Theta_0), A/2)$.

Proof. Stabilization: First we note that

$$|\tilde{V}(r)| = \left| \int_0^r \frac{\tilde{r}(\frac{U}{\tilde{r}})' }{\frac{1}{2}\tilde{r} - U} \right| \lesssim P_\delta^{1-\alpha} \Theta_0 \int_0^r \tilde{r}^{-1} \left(\frac{r}{\tilde{\delta}} \right)^{(1-\alpha-\varepsilon)dA} d\tilde{r} \lesssim \frac{P_\delta^{1-\alpha} \Theta_0}{A} \left(\frac{r}{\tilde{\delta}} \right)^{(1-\alpha-\varepsilon)dA},$$

which implies

$$|e^{\alpha\tilde{V}(r)} - 1| \lesssim \frac{P_\delta^{1-\alpha} \Theta_0}{A} \ll 1.$$

We have the estimates

$$\begin{aligned} P(r) &\lesssim P_\delta \left(\frac{r}{\delta} \right)^{dA}, \\ |F_U(r) - d(2\mu_0 + \lambda_0)A| &\lesssim \left(\frac{P_\delta^{1-\alpha} \Theta_0}{A} + P_\delta^{1-\alpha} A r^2 + \frac{P_\delta^{1-\alpha} \Theta_0}{A} + \frac{P_\delta^{1-\alpha} \Theta_0}{A} r^2 \right) A, \\ |F_\Theta(U)| &\lesssim P_\delta A^3 r^3 + P_\delta A \Theta_0 r + P_\delta A^2 r + A^2 r. \end{aligned}$$

Define $(\hat{U}, \hat{\Theta}) := \Phi(Ar, \Theta_0)$. We deduce that

$$\begin{aligned} |\hat{U} - Ar| &\lesssim \left(P_\delta^{1-\alpha} r^2 + \frac{P_\delta^{1-\alpha} \Theta_0}{A} + P_\delta^{1-\alpha} A r^2 + \frac{P_\delta^{1-\alpha} \Theta_0}{A} + \frac{P_\delta^{1-\alpha} \Theta_0}{A} r^2 \right) A r, \\ |\hat{U}' - A| &\lesssim \left(P_\delta^{1-\alpha} r^2 + \frac{P_\delta^{1-\alpha} \Theta_0}{A} + P_\delta^{1-\alpha} A r^2 + \frac{P_\delta^{1-\alpha} \Theta_0}{A} + \frac{P_\delta^{1-\alpha} \Theta_0}{A} r^2 \right) A, \\ |\hat{\Theta} - \Theta_0| &\lesssim (P_\delta \Theta_0 + A^2) r^2, \\ |\hat{\Theta}'| &\lesssim (P_\delta \Theta_0 + A^2) r. \end{aligned}$$

As for $(\hat{U}(r)/r)'$, we write

$$\left(\frac{\hat{U}(r)}{r} \right)' = -d \frac{U(r)}{r^2} + \frac{r^{-1}}{2\mu_0 + \lambda_0} F_U(r) - \frac{W'(r)}{r} U(r).$$

The third term satisfies

$$\left| \frac{W'(r)}{r} U(r) \right| \lesssim P_\delta^{1-\alpha} \left(\frac{r}{\delta} \right)^{(1-\alpha)dA} A r \lesssim \left(\frac{A r^2}{\Theta_0} \right) P_\delta^{1-\alpha} \Theta_0 r^{-1} \left(\frac{r}{\delta} \right)^{(1-\alpha)dA}.$$

By integration by parts, the sum of the first and the second terms is

$$\begin{aligned} &-d \frac{U(r)}{r^2} + \frac{r^{-1}}{2\mu_0 + \lambda_0} F_U(r) \\ &= \frac{r^{-d-1}}{2\mu_0 + \lambda_0} \int_0^r r_1^d e^{-W(r)+W(r_1)} \left(W'(r_1) F_U(r_1) + F'_U(r_1) \right) dr_1 \\ &= O \left(\left(\frac{A r^2}{\Theta_0} \right) P_\delta^{1-\alpha} \Theta_0 r^{-1} \left(\frac{r}{\delta} \right)^{(1-\alpha)dA} \right) + \frac{r^{-d-1}}{2\mu_0 + \lambda_0} \int_0^r r_1^d e^{-W(r)+W(r_1)} F'_U(r_1) dr_1. \end{aligned}$$

To handle the last term, we write

$$\begin{aligned} |F'_U(r)| &\lesssim |\tilde{V}'| e^{-\alpha\tilde{V}} A + P_\delta^{1-\alpha} A^2 r \left(\frac{r}{\delta} \right)^{(1-\alpha)dA} + P_\delta^{1-\alpha} A \Theta_0 r^{-1} \left(\frac{r}{\delta} \right)^{(1-\alpha)dA} + P_\delta^{1-\alpha} \Theta_0 r \left(\frac{r}{\delta} \right)^{(1-\alpha)dA} \\ &\lesssim \left(A + \frac{A^2}{\Theta_0} r^2 + r^2 \right) P_\delta^{1-\alpha} \Theta_0 r^{-1} \left(\frac{r}{\delta} \right)^{(1-\alpha)dA}, \end{aligned}$$

and hence,

$$\frac{r^{-d-1}}{2\mu_0 + \lambda_0} \int_0^r r_1^d e^{-W(r)+W(r_1)} F'_U(r_1) dr_1 \lesssim \left(A + \frac{A^2}{\Theta_0} r^2 + r^2\right) P_\delta^{1-\alpha} \Theta_0 r^{-1} \left(\frac{r}{\delta}\right)^{(1-\alpha)dA}.$$

We can then conclude that

$$\left| \left(\frac{\widehat{U}(r)}{r} \right)' \right| \lesssim \left(\frac{Ar^2}{\Theta_0} + A + \frac{A^2}{\Theta_0} r^2 + r^2 \right) P_\delta^{1-\alpha} \Theta_0 r^{-1} \left(\frac{r}{\delta}\right)^{(1-\alpha)dA}.$$

Contraction: Let $(U_i, \Theta_i) \in B_{E^\delta}((Ar, \Theta_0), A/2)$, $i = 1, 2$.

Let $D := \|(U_1, \Theta_1) - (U_2, \Theta_2)\|^\delta$.

Denote $(\widetilde{U}_i, \widetilde{\Theta}_i) = \Phi((U_i, \Theta_i))$, $i = 1, 2$.

Using that

$$\begin{aligned} |P_1^{1-\alpha} - P_2^{1-\alpha}| &\leq P_\delta^{1-\alpha} D \ln \left(\frac{\delta}{r} \right) \left(\frac{r}{\delta} \right)^{(1-\alpha)dA} \\ &\lesssim P_\delta^{1-\alpha} \frac{D}{A} \left(\frac{r}{\delta} \right)^{(1-\alpha-\varepsilon)dA}, \end{aligned}$$

we deduce the bounds

$$\begin{aligned} |W_1(r) - W_2(r)| &\lesssim P_\delta^{1-\alpha} \frac{D}{A} \left(\frac{r}{\delta} \right)^{(1-\alpha-\varepsilon)dA} r^2, \quad |Z_1(r) - Z_2(r)| \lesssim P_\delta \frac{D}{A} \left(\frac{r}{\delta} \right)^{(1-\varepsilon)dA} r^2, \\ |W'_1(r) - W'_2(r)| &\lesssim P_\delta^{1-\alpha} \frac{D}{A} \left(\frac{r}{\delta} \right)^{(1-\alpha-\varepsilon)dA} r, \quad |Z'_1(r) - Z'_2(r)| \lesssim P_\delta \frac{D}{A} \left(\frac{r}{\delta} \right)^{(1-\varepsilon)dA} r, \\ |F_{U_1}(r) - F_{U_2}(r)| &\lesssim \left(\frac{P_\delta^{1-\alpha} \Theta_0}{A} + P_\delta^{1-\alpha} A \delta^2 \right) D, \\ |F'_{U_1}(r) - F'_{U_2}(r)| &\lesssim \left(D + \frac{AD}{\Theta_0} \delta^2 + \frac{D}{A} \delta^2 \right) P_\delta^{1-\alpha} \Theta_0 r^{-1} \left(\frac{r}{\delta} \right)^{(1-\alpha-\varepsilon)dA}, \\ |F_{\Theta_1}(r) - F_{\Theta_2}(r)| &\lesssim \left(P_\delta + \frac{P_\delta A^2 \delta^2}{\Theta_0} + \frac{P_\delta^\alpha A}{\Theta_0} + \frac{A}{\Theta_0} \right) Dr \Theta_0. \end{aligned}$$

Then we can estimate

$$\begin{aligned} |\widetilde{U}_1(r) - \widetilde{U}_2(r)| &\lesssim r^{1-d} \int_0^r r_1^{d-1} |e^{-W_1(r)+W_1(r_1)} - e^{-W_2(r)+W_2(r_1)}| |F_{U_1}(r_1)| dr_1 \\ &\quad + r^{1-d} \int_0^r r_1^{d-1} e^{-W_2(r)+W_2(r_1)} |F_{U_1}(r_1) - F_{U_2}(r_1)| dr_1 \\ &\lesssim \left(\delta^2 P_\delta^{1-\alpha} + \frac{P_\delta^{1-\alpha} \Theta_0}{A} \right) Dr. \end{aligned} \tag{2.8}$$

In similar fashion, we have

$$|\widetilde{U}'_1(r) - \widetilde{U}'_2(r)| \lesssim \left(\delta^2 P_\delta^{1-\alpha} + \frac{P_\delta^{1-\alpha} \Theta_0}{A} \right) D. \tag{2.9}$$

Now we write

$$\begin{aligned}
\left| \left(\frac{\widetilde{U}_1}{r} \right)' - \left(\frac{\widetilde{U}_2}{r} \right)' \right| &\lesssim \frac{1}{r} |W_1'(r)U_1(r) - W_2'(r)U_2(r)| \\
&\quad + r^{-d-1} \int_0^r r_1^d \left| e^{-W_1(r)+W_1(r_1)} W_1'(r_1) F_{U_1}(r_1) - e^{-W_2(r)+W_2(r_1)} W_2'(r_1) F_{U_2}(r_1) \right| dr_1 \\
&\quad + r^{-d-1} \int_0^r r_1^d \left| e^{-W_1(r)+W_1(r_1)} F_{U_1}'(r_1) - e^{-W_2(r)+W_2(r_1)} F_{U_2}'(r_1) \right| dr_1 \\
&\lesssim \frac{P_\delta^{1-\alpha} \Theta_0}{A} r^{-1} \left(\frac{r}{\delta} \right)^{(1-\alpha-\varepsilon)dA} D \left(\frac{A}{\Theta_0} \delta^2 + A + \delta^2 \right). \tag{2.10}
\end{aligned}$$

Similarly, we obtain the following estimates on Θ :

$$|\widetilde{\Theta}_1(r) - \widetilde{\Theta}_2(r)| \lesssim \left(P_\delta + \frac{P_\delta A^2 \delta^2}{\Theta_0} + \frac{P_\delta^\alpha A}{\Theta_0} + \frac{A}{\Theta_0} \right) D r^2 \Theta_0 + D A r^2, \tag{2.11}$$

$$|\widetilde{\Theta}_1'(r) - \widetilde{\Theta}_2'(r)| \lesssim \left(P_\delta + \frac{P_\delta A^2 \delta^2}{\Theta_0} + \frac{P_\delta^\alpha A}{\Theta_0} + \frac{A}{\Theta_0} \right) D r \Theta_0 + D A r. \tag{2.12}$$

Putting all the estimates (2.8), (2.9), (2.10), (2.11) and (2.12) together, the desired result follows given our smallness assumptions. \square

To prove Hölder continuity of P , we can repeat an argument from [8] (page 20). We omit the details here.

2.1.3. *Global existence.* We define:

$$\begin{aligned}
Z(r) &= \sup_{0 < s < r} \frac{(1 + P_\delta^{1/2-\alpha/2} s)^2}{M_1 s} |U(s)| + \frac{(1 + P_\delta^{1/2-\alpha/2} s)^2}{M_1'} \left| U'(s) + \frac{d-1}{s} U(s) \right| \\
&\quad + \frac{(1 + \sqrt{P_\delta} s)^2}{M_2} \Theta(s) + \frac{(1 + \sqrt{P_\delta} s)^2}{M_2 P_\delta s} |\Theta'(s)|,
\end{aligned}$$

where the constants M_1, M_1', M_2 are chosen so that

$$M_2 \ll M_1 \ll M_1' \ll 1, A \ll M_1, \Theta_0 \ll M_2, A^2 \ll P_\delta M_2, M_1' \log \frac{1}{\delta^2 P_\delta^{1-\alpha}} \leq 1,$$

$$\frac{P_\delta^{1-\alpha}}{A} M_1' \leq 1, M_2 \ll M_1 P_\delta^{1/2+\alpha}, M_1 M_1' \ll P_\delta M_2, M_1^3 \ll P_\delta^{1-2\alpha} \Theta_0, M_1 M_1' \ll \Theta_0 P_\delta^{1-\alpha}.$$

We make use of Z and a bootstrap argument to prove global existence. This is contained in the following lemma:

Lemma 2.3. *We have $Z(\delta) \leq \frac{1}{2}$.*

Moreover if $Z(r) \leq 1$ for some $r > \delta$, then the stronger bound $Z(r) \leq \frac{1}{2}$ holds.

Proof. Starting point:

To ensure $Z(\delta) \ll 1$, we require, as in [8]:

$$A \ll M_1, \Theta_0 \ll M_2, A^2 \ll P_\delta M_2.$$

Estimate on \tilde{V} : We note that for $r > \delta$:

$$\begin{aligned} |\tilde{V}(r)| &\lesssim \int_0^\delta P_\delta^{1-\alpha} \Theta_0 r_1^{-1} \left(\frac{r_1}{\delta}\right)^{(1-\alpha-\varepsilon)dA} dr_1 + \int_\delta^r \frac{M'_1 + M_1}{r_1 (1 + P_\delta^{1/2-\alpha/2} r_1)^2} ds \\ &\lesssim \frac{P_\delta^{1-\alpha} \Theta_0}{A} + (M_1 + M'_1) \left[\log \frac{1}{P_\delta^{1-\alpha} \delta^2} + 1 \right] \lesssim 1, \end{aligned}$$

provided $(M'_1 + M_1) \log \frac{1}{\delta^2 P_\delta^{1-\alpha}} = O(1)$.

By a similar reasoning, we can show that under that same assumption,

$$P_\delta \lesssim P(r) \lesssim P_\delta,$$

for $r > \delta$.

Estimates on U : We begin with bounds on F_U .

Using the fact that $P(r) \leq P_\delta \left(\frac{r}{A}\right)^{dA}$ for $r \in [0, \delta]$, we write

$$\int_0^r e^{\alpha \tilde{V}(r_1)} \frac{(PR\Theta)'}{P^\alpha} dr_1 \lesssim \frac{P_\delta^{1-\alpha}}{A} M'_1 M_2 + P_\delta^{1-\alpha} M_2 \delta^2 + \left| \int_\delta^r e^{\alpha \tilde{V}(r_1)} \frac{(PR\Theta)'}{P^\alpha} dr_1 \right|.$$

After integrating by parts, we obtain

$$\left| \int_\delta^r e^{\alpha \tilde{V}(r_1)} \frac{(PR\Theta)'}{P^\alpha} dr_1 \right| \lesssim P_\delta^{1-\alpha} \Theta_0 + P_\delta^{1-\alpha} M_2 + M_2 (M_1 + M'_1) \log \frac{1}{P_\delta \delta^2} \lesssim P_\delta^{1-\alpha} \Theta_0 + M_2.$$

We deduce from the above that

$$|F_U(r)| \lesssim A + M_1 M'_1 + P_\delta^{1-\alpha} \Theta_0 + M_2 \quad (2.13)$$

$$|F'_U(r)| \lesssim \frac{M'_1 A + M_1 M'_1 + P_\delta^{1-\alpha} \Theta_0 + M_2}{r} \quad (2.14)$$

Together with the fact that

$$r^{1-d} \int_0^r r_1^{d-1} e^{-C P_\delta^{1-\alpha} (r^2 - r_1^2)} dr_1 \lesssim \frac{r}{(1 + P_\delta^{1/2-\alpha/2} r)^2},$$

(2.13) implies

$$|U(r)| \lesssim \frac{r}{(1 + P_\delta^{1/2-\alpha/2} r)^2} \left(A + M_1 M'_1 + P_\delta^{1-\alpha} \Theta_0 + M_2 \right).$$

Moving on to the derivative part, we first deal with $r \leq \frac{1}{P_\delta^{1/2-\alpha/2}}$. We write that

$$\begin{aligned} \left| U' + \frac{d-1}{r} U \right| &= \left| \frac{1}{2\mu_0 + \lambda_0} F_U(r) - W'(r) U(r) \right| \\ &\lesssim A + M_1 M'_1 + P_\delta \Theta_0 + \frac{M_2}{P_\delta^{1/2+\alpha}} + M_1. \end{aligned}$$

When $r \geq \frac{1}{P_\delta^{1/2-\alpha/2}}$, we write, using an integration by parts:

$$\left| U' + \frac{d-1}{r} U \right| = \left| W'(r) \frac{r^{1-d}}{2\mu_0 + \lambda_0} \int_0^r e^{-W(r)+W(s)} \partial_s \left(\frac{s^{d-1} F_U(s)}{W'(s)} \right) ds \right|. \quad (2.15)$$

We have by (2.14)

$$\left| \partial_s \left(\frac{s^{d-1} F_U(s)}{W'(s)} \right) \right| \lesssim r^{d-3} \frac{A + M_1 M'_1 + P_\delta^{1-\alpha} \Theta_0 + M_2}{P_\delta^{1-\alpha}} \left(1 + \left(\frac{r}{\delta} \right)^{-4dA(1-\alpha)} \right).$$

Together with (2.15) and the fact that

$$r^{1-d} \int_0^r r_1^{d-3} e^{-CP_\delta^{1-\alpha}(r^2-r_1^2)} dr_1 \lesssim \frac{1}{P_\delta^{1-\alpha} r^3} \text{ when } r \geq \frac{1}{P_\delta^{1/2-\alpha/2}},$$

we conclude that

$$\begin{aligned} \left| U' + \frac{d-1}{r} U \right| &\lesssim P_\delta^{1-\alpha} r \frac{1}{P_\delta^{1-\alpha} r^3} \frac{A + M_1 M'_1 + P_\delta^{1-\alpha} \Theta_0 + M_2}{P_\delta^{1-\alpha}} \\ &\lesssim \frac{1}{P_\delta^{1-\alpha} r^2} \left(A + M_1 M'_1 + P_\delta^{1-\alpha} \Theta_0 + M_2 \right). \end{aligned}$$

Estimates on Θ :

We start with the following estimates

$$|F_\Theta(r)| \lesssim M_1 M'_1 r, \text{ when } r \leq \frac{1}{\sqrt{P_\delta}}, \quad (2.16)$$

$$|F_\Theta(r)| \lesssim \frac{1}{r} \left(\frac{M_1^3}{P_\delta^{1-2\alpha}} + \frac{M_1 M'_1}{P_\delta^{1-\alpha}} \right), \text{ when } r \geq \frac{1}{\sqrt{P_\delta}}. \quad (2.17)$$

We deduce from (2.16) that

$$|\Theta(r)| \lesssim \Theta_0 + \frac{M_1 M'_1}{P_\delta}.$$

Now for the case of r large, using that

$$r^{2-d} \int_0^r r_1^{d-3} e^{-CP_\delta(r^2-r_1^2)} dr_1 \lesssim \frac{1}{(1 + \sqrt{P_\delta} r)^2},$$

we deduce from (2.17) that

$$|\Theta(r)| \lesssim \left(\frac{1}{\sqrt{P_\delta} r} \right)^2 \left(\Theta_0 + \frac{M_1^3}{P_\delta^{1-2\alpha}} + \frac{M_1 M'_1}{P_\delta^{1-\alpha}} \right).$$

Similarly for the derivative, we use

$$\left| \partial_r \left[r^{2-d} \int_0^r r_1^{d-3} e^{-CP_\delta(r^2-r_1^2)} dr_1 \right] \right| \lesssim \frac{P_\delta r}{1 + P_\delta r^2},$$

and deduce

$$\begin{aligned} |\Theta'(r)| &\lesssim P_\delta r \left(\Theta_0 + \frac{M_1^3}{P_\delta^{1-2\alpha}} + \frac{M_1 M'_1}{P_\delta^{1-\alpha}} \right), \text{ when } r \leq \frac{1}{\sqrt{P_\delta}}, \\ |\Theta'(r)| &\lesssim \frac{1}{r} \left(\Theta_0 + \frac{M_1^3}{P_\delta^{1-2\alpha}} + \frac{M_1 M'_1}{P_\delta^{1-\alpha}} \right), \text{ when } r \geq \frac{1}{\sqrt{P_\delta}}. \end{aligned}$$

□

Global existence directly follows from the previous lemma by a standard continuation argument.

2.1.4. *Asymptotic behavior.* Regarding the asymptotic behavior of P, U and Θ , the arguments are almost identical to those of [8]. Therefore we only state the condition needed on M_1, M'_1, M_2 to ensure positivity of Θ :

$$\frac{M_1^3}{P_\delta^{1-2\alpha}} + \frac{M_1 M'_1}{P_\delta^{1-\alpha}} \ll \Theta_0.$$

2.2. **Existence when $\alpha = 1$.** In this section, we show an existence theorem in the case where $\alpha = 1$. Unlike in the previous section, we now assume $2\mu_0 + d\lambda_0 > 0$. We show the following existence result:

Theorem 2.4. *Let $d \geq 3$. Fix $(C_V, \kappa, R, \mu_0, \lambda_0) \in (0, \infty)^5$ such that $2\mu_0 + d\lambda_0 > 0$. Then there exists a constant $C(C_V, \kappa, R, \mu_0, \lambda_0) := C > 0$ such that if*

$$A \log \delta^{-1} + P_\delta + \delta < C,$$

there exists a solution $(P, U, \Theta) \in \mathcal{C}^{\frac{dA}{1/2-A}} \times \mathcal{C}^1([0, \infty)) \times \mathcal{C}^1([0, \infty))$ to (2.1) such that $P(0) = 0, P(\delta) = P_\delta, U(0) = 0, U'(0) = A, \Theta(0) = \Theta_0 := \frac{1}{R}(2\mu_0 + d\lambda_0)A, \Theta'(0) = 0$.

Moreover the profiles satisfy the following global bounds:

$$\begin{aligned} P(r) &\simeq P_\delta \min \left[1, \left(\frac{r}{\delta} \right)^{\frac{2dA}{1-2A}} \right], \\ |U(r)| &\lesssim \frac{Ar}{(1+r)^2}, \quad |U'(r)| \lesssim \frac{A}{(1+r)^2}, \\ 0 \leq \Theta(r) &\lesssim \frac{1}{(1 + \sqrt{P_\delta}r)^2}, \quad |\Theta'(r)| \lesssim \frac{\sqrt{P_\delta}r}{(1 + \sqrt{P_\delta}r)^2}. \end{aligned}$$

Finally, there exist $P_\infty > 0, U_\infty > 0, \Theta_\infty > 0$ such that as $r \rightarrow +\infty$:

$$\begin{aligned} P(r) &= P_\infty + O\left(\frac{1}{r^2}\right), \\ U(r) &= \frac{U_\infty}{r} + O\left(\frac{1}{r^3}\right), \\ \Theta(r) &= \frac{\Theta_\infty}{r^2} + O\left(\frac{1}{r^4}\right). \end{aligned}$$

Remark 2.5. *Note that compared to Theorem 2.1, the choice of Θ_0 is not free.*

Proof. We proof is similar to that of Theorem 2.1 above. Therefore we only outline the parts of the argument that differ, namely the derivation of the integro-differential equation, and the proof of local existence.

Integro-differential formulation:

We consider the following initial conditions:

$$P(0) = 0, \quad U(0) = 0, \quad U'(0) = A, \quad \Theta(0) = \Theta_0 := \frac{2\mu_0 + d\lambda_0}{R}A, \quad \Theta'(0) = 0.$$

First, we integrate the equation on P , and obtain for some $\delta > 0$:

$$P(r) = e^{V(r)-V(\delta)} P(\delta), \tag{2.18}$$

where

$$V(r) - V(\delta) = \int_{\delta}^r \frac{U' + \frac{d-1}{r_1}U}{\frac{1}{2}r_1 - U} dr_1.$$

The equation on U reads:

$$\begin{aligned} & -\frac{1}{2}(rPU)' + (PU^2)' + \frac{d-1}{r}PU^2 - (2\mu_0 + \lambda_0)P\left(U' + \frac{d-1}{r}U\right)' + (PR\Theta)' \\ & = (2\mu_0 + \lambda_0)P'U' + \lambda_0P'\frac{d-1}{r}U. \end{aligned}$$

We integrate the previous expression, perform an integration by parts and divide by P to obtain:

$$\begin{aligned} & -\frac{1}{2}rU + U^2 + P^{-1} \int_0^r \frac{d-1}{r_1}PU^2 dr_1 - (2\mu_0 + \lambda_0)\left(U' + \frac{d-1}{r}U\right) + R\Theta \\ & = -2\mu_0P^{-1} \int_0^r P'\frac{d-1}{r_1}U dr_1. \end{aligned}$$

The necessity of the relation between A and Θ_0 can be seen in this equation. Since we expect the behavior $U(r) \simeq Ar$ near 0, equating the higher order terms on each side yields

$$-(2\mu_0 + \lambda_0)\left(U' + \frac{d-1}{r}U\right) + R\Theta + 2\mu_0P^{-1} \int_0^r P'\frac{d-1}{r_1}U dr_1 \simeq -(2\mu_0 + d\lambda_0)A + R\Theta_0.$$

This directly gives $-(2\mu_0 + d\lambda_0)A + R\Theta_0 = 0$.

Now we multiply by $r^{d-1}e^{W(r)}$ (where $W(r) = \frac{r^2}{4(2\mu_0 + \lambda_0)}$) and obtain

$$\begin{aligned} & -(2\mu_0 + \lambda_0)\left(r^{d-1}e^{W(r)}U\right)' = -r^{d-1}e^{W(r)}\left(R\Theta + U^2 + P^{-1} \int_0^r \frac{d-1}{r_1}PU^2 dr_1 \right. \\ & \quad \left. + 2\mu_0P^{-1} \int_0^r P'\frac{d-1}{r_1}U dr_1\right). \end{aligned}$$

We conclude with the integro-differential satisfied by U :

$$U(r) = \frac{r^{1-d}}{2\mu_0 + \lambda_0} \int_0^r e^{W(r_1)-W(r)} r_1^{d-1} F_U(r_1) dr_1,$$

where

$$F_U(r) = R\Theta + U^2 + P^{-1} \int_0^r \frac{d-1}{r_1}PU^2 dr_1 + 2\mu_0P^{-1} \int_0^r P'\frac{d-1}{r_1}U dr_1.$$

For Θ , we do not detail the argument since it is similar to the above. We get

$$\Theta(r) = (d-2)r^{2-d} \int_0^r r_1^{d-3} e^{-Z(r)+Z(r_1)} dr_1 \Theta_0 + \frac{r^{2-d}}{\kappa} \int_0^r r_1^{d-2} e^{Z(r_1)-Z(r)} F_{\Theta}(r_1) dr_1 - \frac{U^2}{2C_V},$$

where

$$\begin{aligned}
Z(r) &:= \frac{C_V}{2\kappa} \int_0^r \tilde{r} P(\tilde{r}) d\tilde{r}, \\
F_\Theta(r) &:= UP \left(\frac{U^2}{2} + C_V \Theta \right) + UPR\Theta + \frac{d-2}{r} \int_0^r \left(UP \left(\frac{U^2}{2} + C_V \Theta \right) + UPR\Theta \right) dr_2 \\
&\quad + \left(\frac{\kappa}{C_V} - (2\mu_0 + \lambda_0) \right) \frac{(U^2)'}{2} + \frac{\kappa}{C_V} \frac{d-2}{2r} U^2 \\
&\quad + \frac{2\mu_0 + \lambda_0}{r} \int_0^r \left((2-d)r_1^{-d+1} P + Pr_1^{-d+2} \frac{U' + \frac{d-1}{r} U}{\frac{1}{2}r - U} \right) (r_1^{d-1} UU') dr_1 \\
&\quad - \lambda \frac{d-1}{r} U^2 + \lambda_0 \frac{d-1}{r} \int_0^r (r_1^{-d+2} P)' (r_1^{d-2} U^2) dr_1 \\
&\quad + \frac{(2\mu_0 + \lambda_0)}{r} \int_0^r r_1 P' UU' dr_1 + \frac{(d-1)\lambda_0}{r} \int_0^r P' U^2 dr_1
\end{aligned}$$

Local existence:

Fix $\varepsilon > 0$ such that $A \ll \varepsilon \ll 1$. Assume that δ satisfies $\delta^\varepsilon/\varepsilon \ll 1$. We define the following norm (slightly different from the above):

$$\|(U, \Theta)\|^\delta := \sup_{0 < r < \delta} \left[r^{-1} |U(r)| + r^{1-\varepsilon} \left| \left(\frac{U}{r} \right)' \right| + \Theta(r) + r^{-1} |\Theta'(r)| \right].$$

We apply the fixed point argument with the above norm in the space

$$B_{E^\delta}((Ar, \Theta_0), A/2) \cap \left\{ \lim_{r \rightarrow 0} \frac{U(r)}{r} = A, \lim_{r \rightarrow 0} \Theta(r) = \Theta_0 \right\}.$$

Stabilization: We have the following estimates:

$$\begin{aligned}
|F_U(r) - d(2\mu_0 + \lambda_0)A| &\lesssim A \left(\delta^2 + \frac{\delta^\varepsilon}{\varepsilon} \right), \\
|F_\Theta(r)| &\lesssim (A^2 + P_\delta A)r,
\end{aligned}$$

where we used that $\Theta_0 = \frac{1}{R}(2\mu_0 + d\lambda_0)A$.

Denote $(\widehat{U}, \widehat{\Theta}) = \Phi((Ar, \Theta_0))$.

We can deduce from the above that

$$\begin{aligned}
|\widehat{U} - Ar| &\lesssim Ar \left(\delta^2 + \frac{\delta^\varepsilon}{\varepsilon} \right), \\
|\widehat{\Theta} - \Theta_0| &\lesssim (A^2 + P_\delta A)r^2, \\
|\widehat{\Theta}'| &\lesssim (A^2 + P_\delta A)r.
\end{aligned}$$

For the second U -part of the norm we can write, using that $P' > 0$ on $(0, \delta)$:

$$\begin{aligned}
|F'_U(r)| &\lesssim |\Theta'| + |UU'| + \left| \left(P^{-1} \int_0^r P \frac{U^2}{r_1} dr_1 \right)' \right| + \frac{A}{r} P^{-1} \int_0^r P' \left| \frac{(d-1)U(r_1)}{r_1} - \frac{(d-1)U(r)}{r} \right| dr_1 \\
&\lesssim Ar + \frac{A^2}{\varepsilon} r^{\varepsilon-1}.
\end{aligned}$$

We deduce the following estimate as we did in Section 2.1:

$$|(\frac{\widehat{U}}{r})'| \lesssim Ar^{-1+\varepsilon}(\delta^{2-\varepsilon} + \frac{A}{\varepsilon}).$$

Contraction: Let $(U_i, \Theta_i) \in B_{E^\delta}((Ar, \Theta_0), A/2)$. Denote $D := \|(U_1, \Theta_1) - (U_2, \Theta_2)\|^\delta$.

Finally, let $(\widetilde{U}_i, \widetilde{\Theta}_i) := \Phi((U_i, \Theta_i))$.

We have, arguing as in the stabilization part of the proof:

$$\begin{aligned} |\widetilde{U}_1(r) - \widetilde{U}_2(r)| &\lesssim Dr \left(\delta^2 + \frac{\delta^\varepsilon}{\varepsilon} \right), \\ |(\frac{\widetilde{U}_1}{r})' - (\frac{\widetilde{U}_2}{r})'| &\lesssim Dr^{-1+\varepsilon}(\delta^{2-\varepsilon} + \frac{A^2}{\varepsilon}), \\ |\Theta_1(r) - \Theta_2(r)| &\lesssim ADr^2, \\ |\Theta'_1(r) - \Theta'_2(r)| &\lesssim ADr. \end{aligned}$$

The proofs of global existence and asymptotic behavior of the solutions are done similarly as in Section 2.1, therefore details are omitted. \square

3. NON-EXISTENCE OF SHRINKERS

We now investigate shrinker-type solutions to the system (cNS), that is solutions of the type (1.3).

Plugging the corresponding ansatz into the system, we obtain the following ODEs satisfied by the profiles P, U and Θ :

For $r = |x| > 0$

$$\left\{ \begin{aligned} &\frac{1}{2}rP' + P'U + P\left(U' + \frac{d-1}{r}U\right) = 0, \\ &\frac{1}{2}PU + \frac{1}{2}r(PU)' + (PU^2)' + \frac{d-1}{r}PU^2 + (PR\Theta)' \\ &\quad = (2\mu_0 + \lambda_0)P^\alpha \left(U'' + \frac{d-1}{r}U' - \frac{d-1}{r^2}U \right) \\ &\quad \quad - (2\mu_0 + \lambda_0)\alpha \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U} P^\alpha U' - \lambda_0\alpha \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U} P^\alpha \frac{d-1}{r}U, \\ &P\left(\frac{U^2}{2} + C_V\Theta\right) + \frac{1}{2}r\left(P\left(\frac{U^2}{2} + C_V\Theta\right)\right)' + \left(UP\left(\frac{U^2}{2} + C_V\Theta\right) + UPR\Theta\right)' \\ &\quad + \frac{d-1}{r}\left(UP\left(\frac{U^2}{2} + C_V\Theta\right) + UPR\Theta\right) - \kappa\left(\Theta'' + \frac{d-1}{r}\Theta'\right) \\ &\quad = 2\mu_0P^\alpha\left((U')^2 + \frac{d-1}{r^2}U^2\right) + \lambda_0P^\alpha\left(U' + \frac{d-1}{r}U\right)^2 \\ &\quad + (2\mu_0 + \lambda_0)P^\alpha\left(U'' + \frac{d-1}{r}U' - \frac{d-1}{r^2}U\right)U \\ &\quad - (2\mu_0 + \lambda_0)\alpha \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U} P^\alpha U'U - \lambda_0\alpha \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U} P^\alpha \frac{d-1}{r}U^2. \end{aligned} \right. \quad (3.1)$$

The main result of this section is that (3.1) does not have non trivial small solutions.

As mentioned above, given the dependence laws that we have adopted for the Lamé coefficients, it is natural to restrict to solutions that exhibit cavitation. More precisely,

we will make the following mild assumptions on the density profile P throughout this section:

$$\begin{aligned} & \bullet \exists \varepsilon, P_\varepsilon > 0, \forall r \geq \varepsilon, P(r) \geq P_\varepsilon, \\ & \bullet \exists \Lambda > 0, \forall r \in (0, \varepsilon), \frac{r \int_0^r P^{1-\alpha}(r_1) dr_1}{\int_0^r P^{1-\alpha}(r_1) r_1 dr_1}, \frac{\int_0^r P}{P^\alpha \int_0^r P^{1-\alpha}} \leq \Lambda. \end{aligned} \tag{3.2}$$

Note that the functions P constructed in Section 2 satisfy this condition. Since the forward and backward systems are essentially equivalent near the origin, a shrinker exhibiting cavitation would be expected to behave similarly near 0.

The main theorem of this section is:

Theorem 3.1. *Let $d \geq 3$ and $0 < \alpha < 1$. Fix $(C_V, \kappa, R, \mu_0, \lambda_0) \in (0, \infty)^5$ such that $C_V \leq \frac{\kappa}{2\mu_0 + \lambda_0}$.*

Assume that the density function P satisfies the above condition (3.2).

Then there exists a constant $C(C_V, \kappa, R, \mu_0, \lambda_0, \varepsilon, P_\varepsilon, \Lambda) := C > 0$ such that if

$$\sup_{r>0} \left(\langle r \rangle^2 \Theta + P^{1-\alpha} + \left| \frac{U}{r\Theta} \right| \right) + \sup_{r>\varepsilon} \left| \frac{U'}{r\Theta'} \right| < C,$$

then $U \equiv \Theta \equiv 0, P = \text{Constant}$.

Remark 3.2. *The assumption $d \geq 3$ comes from the fact that we use Hardy's inequality in the proof.*

Remark 3.3. *Note that by scaling we expect the behavior $U(r) \sim \frac{U_\infty}{r}$ and $\Theta \sim \frac{\Theta_\infty}{r^2}$ at infinity. This makes the assumptions in the theorem critical.*

Remark 3.4. *The proof can be adapted with minor changes when $\alpha = 1$, or when the Lamé coefficients are not density dependent. For example this method could be applied in the case where $C_V \geq \frac{\kappa}{2\mu + \lambda}$. In this case the smallness condition would be written in terms of $\sup_{r>\varepsilon} \left| \frac{\Theta}{rU} \right|$ and $\sup_{r>\varepsilon} \left| \frac{\Theta'}{rU'} \right|$.*

Remark 3.5. *Note that our smallness assumptions imply that there exists $0 < b \ll 1$ such that*

$$\sup_{r>0} \left| \frac{U}{r} \right| \leq b.$$

Remark 3.6. *We notice that for any $r > 0$, $C_V \leq \frac{1}{2} \frac{\kappa}{2\mu_0 + \lambda_0} \frac{1}{P^\alpha}$, given the smallness assumption on P .*

The remainder of this section is dedicated to the proof of Theorem 3.1.

3.1. **Set-up.** Let us recall $\mu = \mu_0 P^\alpha, \lambda = \lambda_0 P^\alpha$. We start by writing the equations on Θ and U from (3.1) in 2×2 matrix form:

$$\begin{aligned}
& \begin{pmatrix} C_V P(\frac{r}{2} + U) \frac{d}{dr} & PR\Theta \frac{d}{dr} \\ PR \frac{d}{dr} & P(\frac{r}{2} + U) \frac{d}{dr} \end{pmatrix} \begin{pmatrix} \Theta \\ U \end{pmatrix} \\
& - \begin{pmatrix} \kappa(\frac{d-1}{r} \frac{d}{dr} - \frac{d^2}{dr^2}) & 0 \\ 0 & (2\mu + \lambda)(\frac{d-1}{r} \frac{d}{dr} - \frac{d^2}{dr^2}) \end{pmatrix} \begin{pmatrix} \Theta \\ U \end{pmatrix} \\
& = \begin{pmatrix} -P(r)\Theta(r)(C_V + R\frac{d-1}{r}U) + 2\mu\left((U')^2 + (\frac{d-1}{r}U)^2\right) + \lambda\left(U' + \frac{d-1}{r}U\right)^2 \\ -\frac{1}{2}P(r)U(r) - (2\mu + \lambda)\frac{d-1}{r^2}U \end{pmatrix} \\
& + \begin{pmatrix} -(2\mu + \lambda)\alpha\frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U}U'U - \lambda\alpha\frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U}\frac{d-1}{r}U^2 \\ -(2\mu + \lambda)\alpha\frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U}U' - \lambda\alpha\frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U}\frac{d-1}{r}U \end{pmatrix}.
\end{aligned}$$

We denote

$$\begin{aligned}
B &= \begin{pmatrix} \kappa & 0 \\ 0 & 2\mu + \lambda \end{pmatrix}, \\
\tilde{A}(r) &= \begin{pmatrix} C_V P(\frac{r}{2} + U) & PR\Theta \\ PR & P(\frac{r}{2} + U) \end{pmatrix}.
\end{aligned}$$

The previous equation can then be written

$$\begin{aligned}
& \tilde{A}(r) \frac{d}{dr} \begin{pmatrix} \Theta \\ U \end{pmatrix} - B \left(\frac{d-1}{r} - \frac{d^2}{dr^2} \right) \begin{pmatrix} \Theta \\ U \end{pmatrix} \\
& = \begin{pmatrix} -P(r)\Theta(r)(C_V + R\frac{d-1}{r}U) + 2\mu\left((U')^2 + (\frac{d-1}{r}U)^2\right) + \lambda\left(U' + \frac{d-1}{r}U\right)^2 \\ -\frac{1}{2}P(r)U(r) - (2\mu + \lambda)\frac{d-1}{r^2}U \end{pmatrix} \\
& + \begin{pmatrix} -(2\mu + \lambda)\alpha\frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U}U'U - \lambda\alpha\frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U}\frac{d-1}{r}U^2 \\ -(2\mu + \lambda)\alpha\frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U}U' - \lambda\alpha\frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U}\frac{d-1}{r}U \end{pmatrix}.
\end{aligned}$$

Hence after multiplication by B^{-1} and $\exp(-\int_0^r B^{-1}\tilde{A}(r')dr)$, we obtain:

$$\begin{aligned}
& -\nabla \cdot \left(\exp\left(-\int_0^r B^{-1}\tilde{A}(r')dr\right) \nabla \begin{pmatrix} \Theta \\ U \end{pmatrix} \right) \\
&= \exp\left(-\int_0^r B^{-1}\tilde{A}(r')dr\right) \\
&\times \left(-\frac{P(r)\Theta(r)}{\kappa} \left(C_V + R\frac{d-1}{r}U \right) + 2\frac{\mu}{\kappa} \left((U')^2 + \left(\frac{d-1}{r}U\right)^2 \right) + \frac{\lambda}{\kappa} \left(U' + \frac{d-1}{r}U \right)^2 \right. \\
&\quad \left. - \frac{1}{2(2\mu+\lambda)} P(r)U(r) - \frac{d-1}{r^2}U \right) \\
&+ \exp\left(-\int_0^r B^{-1}\tilde{A}(r')dr\right) \\
&\times \left(-\frac{(2\mu+\lambda)\alpha}{\kappa} \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U} U'U - \frac{\lambda}{\kappa} \alpha \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U} \frac{d-1}{r}U^2 \right. \\
&\quad \left. - \alpha \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U} U' - \frac{\lambda}{2\mu+\lambda} \alpha \frac{U' + \frac{d-1}{r}U}{\frac{1}{2}r + U} \frac{d-1}{r}U \right).
\end{aligned} \tag{3.3}$$

Denote from now on

$$\begin{aligned}
\exp(A(r)) &:= \exp\left(-\int_0^r B^{-1}\tilde{A}(r')dr\right) \\
&:= \exp\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}.
\end{aligned}$$

3.2. Basic properties of the matrix A . In this section we collect some elementary facts about the matrix A .

First note that we have explicit expressions for its coefficients:

$$\begin{aligned}
\alpha &= -\frac{C_V}{\kappa} \int_0^r P(r') \left(\frac{r'}{2} + U \right) dr', \\
\beta &= -\frac{R}{\kappa} \int_0^r P\Theta dr', \\
\gamma &= -R \int_0^r \frac{P}{2\mu+\lambda} dr', \\
\delta &= -\int_0^r \frac{P(r')}{2\mu+\lambda} \left(\frac{r'}{2} + U \right) dr'.
\end{aligned}$$

Remark 3.7. Given the assumption $C_V \leq \frac{\kappa}{2\mu+\lambda}$, we have $\delta < \alpha$.

The next lemma shows that the matrix A can be diagonalized.

Lemma 3.8. We have the following decomposition for A :

There exists a real-valued 2×2 matrix Q , with diagonal elements equal to one, such that:

$$A(r) = Q^{-1}(r) \begin{pmatrix} -\lambda_{\min} & 0 \\ 0 & -\lambda_{\max} \end{pmatrix} Q(r),$$

where $-\lambda_{\max} < -\lambda_{\min}$ denote the two real valued eigenvalues of A .

Proof. The characteristic polynomial of that matrix A is $X^2 - (\delta + \alpha)X + \alpha\delta - \beta\gamma$. Its discriminant is

$$\Delta = (\delta + \alpha)^2 - 4(\alpha\delta - \beta\gamma) = (\alpha - \delta)^2 + 4\beta\gamma.$$

Given that both β and γ are negative, $\Delta > 0$.

From this we can deduce the two eigenvalues:

$$-\lambda_{max} := \frac{1}{2}(\alpha + \delta - \sqrt{\Delta}), \quad -\lambda_{min} := \frac{1}{2}(\alpha + \delta + \sqrt{\Delta}).$$

We find the corresponding eigenvectors, and deduce that the matrix $A(r)$ can then be diagonalized after introducing the matrix $Q(r)$ defined as

$$\begin{aligned} Q(r) &= \begin{pmatrix} 1 & \frac{-\lambda_{max}-\delta}{\gamma} \\ \frac{\gamma}{-\lambda_{min}-\delta} & 1 \end{pmatrix}, \\ Q^{-1}(r) &= \frac{1}{D} \begin{pmatrix} 1 & -\frac{-\lambda_{max}-\delta}{\gamma} \\ -\frac{\gamma}{-\lambda_{min}-\delta} & 1 \end{pmatrix}, \end{aligned}$$

where $D := 1 - \frac{-\lambda_{max}-\delta}{-\lambda_{min}-\delta}$. □

We write

$$Q(r) = \begin{pmatrix} 1 & q_{12} \\ q_{21} & 1 \end{pmatrix}, \quad Q^{-1}(r) = \frac{1}{D} \begin{pmatrix} 1 & -q_{12} \\ -q_{21} & 1 \end{pmatrix}, \quad D = 1 - q_{12}q_{21}.$$

The following lemma justifies the fact that the matrix Q is a perturbation of the identity for large values of r :

Lemma 3.9. *We have the following estimates, valid for any $r > 0$:*

$$\begin{aligned} |q_{12}| &\leq \frac{2(2\mu_0 + \lambda_0)R\Lambda \sup_{r>0} P^\alpha \Theta}{\kappa(1/2 - b)r}, \\ |q_{21}| &\leq \frac{2R\Lambda}{(1/2 - b)r}, \\ |D| &\geq 1. \end{aligned}$$

Proof. We have

$$|q_{12}| = \left| \frac{-\lambda_{max} - \delta}{\gamma} \right| = \frac{2\beta}{\alpha - \delta + \sqrt{\Delta}} \leq \frac{\beta}{\alpha - \delta}.$$

Using remark 3.6 and the condition (3.2), we obtain the desired bound. Similarly, we obtain the bound on q_{21} .

For the last estimate on D , we notice that

$$1 - \frac{\alpha - \delta - \sqrt{\Delta}}{\alpha - \delta + \sqrt{\Delta}} = 1 - \frac{-4\beta\gamma}{(\alpha - \delta + \sqrt{\Delta})^2} \geq 1,$$

since both β and γ are negative. □

In what follows, we will systematically decompose Q as the sum of the identity matrix, and the matrix of its off-diagonal terms that will be treated like an error term. The following basic computation will be used repeatedly in the sequel:

Lemma 3.10. *We have, for real numbers a, b, c, d :*

$$\begin{aligned}
& \left\langle Q^{-1} \begin{pmatrix} e^{-\lambda_{\min}} & 0 \\ 0 & e^{-\lambda_{\max}} \end{pmatrix} Q \begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} c \\ d \end{pmatrix} \right\rangle \\
&= \left\{ \left(1 + \frac{q_{12}q_{21}}{D} \right) e^{-\lambda_{\min}} - \frac{q_{12}q_{21}}{D} e^{-\lambda_{\max}} \right\} ac \\
&+ \left\{ \left(1 + \frac{q_{12}q_{21}}{D} \right) e^{-\lambda_{\max}} - \frac{q_{12}q_{21}}{D} e^{\lambda_{\min}} \right\} bd \\
&+ (e^{-\lambda_{\min}} - e^{-\lambda_{\max}}) \frac{q_{12}}{D} bc + (-e^{-\lambda_{\min}} + e^{-\lambda_{\max}}) \frac{q_{21}}{D} ad.
\end{aligned}$$

We also have

$$\frac{1}{2} \leq 1 + \frac{q_{12}q_{21}}{D} \leq 1, \quad 0 < -q_{12}q_{21} \leq 1.$$

We end this section with rough bounds that are relevant to treat the case where r is close to 0:

Lemma 3.11. *We have the following estimate for any $r > 0$:*

$$|q_{12}| \leq \sqrt{\frac{2\mu_0 + \lambda_0}{\kappa} \sup_{r>0} P^\alpha \Theta}.$$

Moreover, we have

$$\sup_{0 < r < \varepsilon} e^{\sqrt{\Delta}} \leq 2, \quad \sup_{0 < r < \varepsilon} e^{\lambda_{\min}} \leq 2.$$

Proof. We write that

$$|q_{12}| = \left| \frac{-\lambda_{\max} - \delta}{\gamma} \right| = \frac{2|\beta|}{\alpha - \delta + \sqrt{(\alpha - \delta)^2 + 4\beta\gamma}} \leq \sqrt{\frac{\beta}{\gamma}} \leq \sqrt{\frac{2\mu_0 + \lambda_0}{\kappa} \sup_{r>0} P^\alpha \Theta}.$$

For the last two estimates, we use the following crude bounds:

$$\begin{aligned}
\sqrt{\Delta} &\leq \sqrt{4 \frac{R^2}{\kappa} \frac{\varepsilon^2}{2\mu_0 + \lambda_0} \sup_{r>0} P \Theta \sup_{r>0} P^{1-\alpha} + \frac{\varepsilon^4}{(2\mu_0 + \lambda_0)^2} \left(\sup_{r>0} P^{1-\alpha} \right)^2}, \\
\lambda_{\min} &\leq \frac{\varepsilon^2}{2\mu_0 + \lambda_0} \sup_{r>0} P^{1-\alpha}.
\end{aligned}$$

The estimates then follow from the smallness assumptions. \square

3.3. Weighted energy estimate. Now we take the inner product the equation (3.3) with $(\Theta; U)^T$, and integrate by parts in the left-hand side.

Then we split the expression between small and large values of r . Overall we get

$$\begin{aligned}
\text{LHS} &= \int_0^\varepsilon \langle \exp(A(r)) \begin{pmatrix} \Theta' \\ U' \end{pmatrix}, \begin{pmatrix} \Theta' \\ U' \end{pmatrix} \rangle dx \\
&+ \int_\varepsilon^\infty \langle \exp(A(r)) \begin{pmatrix} \Theta' \\ U' \end{pmatrix}, \begin{pmatrix} \Theta' \\ U' \end{pmatrix} \rangle dx \\
&:= \text{LHS}_1 + \text{LHS}_2.
\end{aligned}$$

We prove the following estimate on the left-hand side:

Lemma 3.12. *We have*

$$LHS \geq \int_0^\varepsilon e^{-\lambda_{max}} (\Theta'^2 + U'^2) dx + \frac{1}{4} \left(\int_\varepsilon^{+\infty} e^{-\lambda_{min}} \Theta'^2 dx + \int_\varepsilon^{+\infty} e^{-\lambda_{max}} U'^2 dx \right).$$

Proof. Bound on LHS₁ :

In this case the bound is straightforward:

$$LHS_1 \geq \int_0^\varepsilon e^{-\lambda_{max}} (U'^2 + \Theta'^2) dx.$$

Bound on LHS₂ :

As announced above, in this range, we use the diagonalization of A and we write the matrices Q and Q^{-1} as the sum of the identity and an error term. This yields, using Lemma 3.10:

$$\begin{aligned} LHS_2 &= \int_\varepsilon^\infty \left\{ \left(1 + \frac{q_{12}q_{21}}{D} \right) e^{-\lambda_{min}} - \frac{q_{12}q_{21}}{D} e^{-\lambda_{max}} \right\} \Theta'^2 dx \\ &\quad + \int_\varepsilon^\infty \left\{ \left(1 + \frac{q_{12}q_{21}}{D} \right) e^{-\lambda_{max}} - \frac{q_{12}q_{21}}{D} e^{-\lambda_{min}} \right\} U'^2 dx \\ &\quad + \int_\varepsilon^\infty (e^{-\lambda_{min}} - e^{-\lambda_{max}}) \frac{q_{12} - q_{21}}{D} U' \Theta' dx \end{aligned}$$

We notice that

$$\frac{1}{2} \leq 1 + \frac{q_{12}q_{21}}{D} \leq 1, \quad -q_{12}q_{21} > 0,$$

which makes the first two terms the main contribution. The last term is treated as an error.

Using the bounds from Lemma 3.9 and the smallness assumptions, we write that

$$\begin{aligned} \left| \int_\varepsilon^\infty \frac{q_{12}}{D} |e^{-\lambda_{min}} - e^{-\lambda_{max}}| U' \Theta' dx \right| &\leq \sup_{r \geq \varepsilon} \left| \frac{U'}{r \Theta'} \right| \frac{2(2\mu_0 + \lambda_0) R \Lambda \sup_{r > 0} P^\alpha \Theta}{\kappa(1/2 - b)} \int_\varepsilon^{+\infty} e^{-\lambda_{min}} \Theta'^2 dx \\ &\leq \frac{1}{100} \int_\varepsilon^{+\infty} e^{-\lambda_{min}} \Theta'^2 dx, \\ \left| \int_\varepsilon^\infty \frac{q_{21}}{D} |e^{-\lambda_{min}} - e^{-\lambda_{max}}| \Theta' U' dx \right| &\leq \sup_{r \geq \varepsilon} \left| \frac{U'}{r \Theta'} \right| \frac{2R \Lambda}{1/2 - b} \int_\varepsilon^{+\infty} e^{-\lambda_{min}} \Theta'^2 dx \\ &\leq \frac{1}{100} \int_\varepsilon^{+\infty} e^{-\lambda_{min}} \Theta'^2 dx. \end{aligned}$$

The other terms are easier to bound, therefore we omit the details. □

Now we move on to the right-hand side. We show that

Lemma 3.13. *We have*

$$\begin{aligned} RHS &\leq \frac{1}{20} \left(\int_{\mathbb{R}^d} e^{-\lambda_{min}} \Theta'^2 dx + \int_{\mathbb{R}^d} e^{-\lambda_{max}} U'^2 dx - \int_{\mathbb{R}^d} e^{-\lambda_{max}} \frac{P}{2\mu + \lambda} U^2 dx \right) \\ &\quad - \frac{C_V}{4\kappa} \int_{\mathbb{R}^d} e^{-\lambda_{min}} P \Theta^2 dx. \end{aligned}$$

Proof. We start by decomposing the right-hand side into three parts:

$$\begin{aligned}
& \left\langle \text{RHS of (3.3)}, \begin{pmatrix} \Theta \\ U \end{pmatrix} \right\rangle \\
&= - \left\langle \exp(A(r)) \begin{pmatrix} 0 \\ \frac{d-1}{r^2} U \end{pmatrix}, \begin{pmatrix} \Theta \\ U \end{pmatrix} \right\rangle \\
&- \left\langle \exp(A(r)) \begin{pmatrix} \frac{C_V}{\kappa} P(r) \Theta(r) \\ \frac{1}{2(2\mu + \lambda)} P(r) U(r) \end{pmatrix}, \begin{pmatrix} \Theta \\ U \end{pmatrix} \right\rangle \\
&+ \left\langle \exp(A(r)) \begin{pmatrix} N_1(P, \Theta, U) \\ N_2(P, \Theta, U) \end{pmatrix}, \begin{pmatrix} \Theta \\ U \end{pmatrix} \right\rangle \\
&:= \text{RHS}_1 + \text{RHS}_2 + \text{RHS}_3,
\end{aligned}$$

where

$$\begin{aligned}
N_1(P, \Theta, U) &= -\frac{R}{\kappa} P \Theta \frac{d-1}{r} U + 2\frac{\mu}{\kappa} \left((U')^2 + \left(\frac{d-1}{r} U \right)^2 \right) + \frac{\lambda}{\kappa} \left(U' + \frac{d-1}{r} U \right)^2 \\
&- \frac{2\mu + \lambda}{\kappa} \alpha \frac{U' + \frac{d-1}{r} U}{\frac{1}{2}r + U} U' U - \frac{\lambda}{\kappa} \alpha \frac{U' + \frac{d-1}{r} U}{\frac{1}{2}r + U} \frac{d-1}{r} U^2, \\
N_2(P, \Theta, U) &= -\alpha \frac{U' + \frac{d-1}{r} U}{\frac{1}{2}r + U} U' - \frac{\lambda}{2\mu + \lambda} \alpha \frac{U' + \frac{d-1}{r} U}{\frac{1}{2}r + U} \frac{d-1}{r} U.
\end{aligned}$$

Next we estimate RHS_1 , RHS_2 and RHS_3 . The result from Lemma 3.13 will then follow.

Bound on RHS_1 :

As we did above, we use the elementary computation from Lemma 3.10 to obtain the expression:

$$\text{RHS}_1 = - \int_{\mathbb{R}^d} \left\{ \left(1 + \frac{q_{12}q_{21}}{D} \right) e^{-\lambda_{\max}} - \frac{q_{12}q_{21}}{D} e^{\lambda_{\min}} \right\} \frac{d-1}{r^2} U^2 \, dx \quad (3.4)$$

$$- \int_{\mathbb{R}^d} (e^{-\lambda_{\min}} - e^{-\lambda_{\max}}) \frac{q_{12}}{D} \frac{d-1}{r^2} U \Theta \, dx \quad (3.5)$$

We know that the first term is negative. For the second term, we will distinguish between r large and r small. We introduce a cut-off function χ such that:

- $\chi = 1$ on $[0, \varepsilon]$.
- χ is supported on $[0, 2\varepsilon]$.
- $|\chi'| \leq C \mathbf{1}_{\{\varepsilon \leq |x| \leq 2\varepsilon\}}$ for some numerical constant C . Here $\mathbf{1}_A$ denotes the characteristic function of the set A .

Using this cut-off, we write:

$$|(3.5)| \leq \int_{\mathbb{R}^d} \frac{|q_{12}|}{D} e^{-\lambda_{\min}} \frac{d-1}{r^2} \chi^2 |U| \Theta \, dx \quad (3.6)$$

$$+ \int_{\mathbb{R}^d} \frac{|q_{12}|}{D} e^{-\lambda_{\min}} \frac{d-1}{r^2} (1 - \chi^2) |U| \Theta \, dx. \quad (3.7)$$

Bound on (3.6):

We write, using Lemma 3.11, Young's inequality, as well as the mean value theorem, we

obtain the bound:

$$|(3.6)| \leq \frac{1}{2} \sqrt{\frac{2\mu_0 + \lambda_0}{\kappa}} \sup_{r>0} P^\alpha \Theta \sup_{0<r<\varepsilon} (e^{2\sqrt{\Delta}}) \int_0^{+\infty} e^{-\lambda_{max} x} \chi^2 \left[\frac{(d-1)U^2}{r^2} + \frac{(d-1)\Theta^2}{r^2} \right] dx.$$

The first part containing U can be absorbed into (3.4) with the smallness assumptions and Lemma 3.11.

For the part containing Θ , we use that $-\lambda_{max} \leq 0$ as well as Hardy's inequality, and the properties of the cut-off χ to write:

$$\begin{aligned} \int_{\mathbb{R}^d} e^{-\lambda_{max} x} \chi^2 \frac{(d-1)\Theta^2}{r^2} dx &\leq C' \int_{\mathbb{R}^d} ((\chi\Theta)')^2 dx \\ &\leq 2C' \int_{\mathbb{R}^d} \Theta'^2 dx + 2C' \int_{\mathbb{R}^d} \chi'^2 \Theta^2 dx \\ &\leq 2C' \sup_{0<r<\varepsilon} e^{\lambda_{max}} \int_{\mathbb{R}^d} e^{-\lambda_{max} x} \Theta'^2 dx \\ &\quad + 2C' C^2 \sup_{0<r<\varepsilon} e^{\lambda_{min}} \frac{1}{P_\varepsilon} \int_{\mathbb{R}^d} e^{-\lambda_{min} x} P \Theta^2 dx, \end{aligned}$$

where C' denotes the constant in Hardy's inequality.

Given the smallness assumptions, these contributions are acceptable.

Bound on (3.7):

Using Lemma 3.11, we obtain:

$$\begin{aligned} |(3.7)| &\leq \sup_{r>\varepsilon} \left| \frac{U}{r\Theta} \right| \frac{2R\Lambda(2\mu_0 + \lambda_0) \sup_{r>0} P^\alpha \Theta}{\kappa(1/2 - b)} \int_{\mathbb{R}^d} (1 - \chi^2) e^{-\lambda_{min} x} \frac{d-1}{r^2} \Theta^2 dx \\ &\leq \frac{2R\Lambda(2\mu_0 + \lambda_0) \sup_{r>0} P^\alpha \Theta}{\varepsilon^2 \kappa P_\varepsilon (1/2 - b)} \sup_{r>\varepsilon} \left| \frac{U}{r\Theta} \right| \int_{\mathbb{R}^d} e^{-\lambda_{min} x} P \Theta^2 dx, \end{aligned}$$

which will be controlled by a part of RHS_2 . We can conclude with the smallness assumptions.

Bound on RHS_2 :

We use a similar reasoning for this part. With Lemma 3.10, we obtain:

$$\text{RHS}_2 = -\frac{C_V}{\kappa} \int_{\mathbb{R}^d} \left\{ \left(1 + \frac{q_{12}q_{21}}{D} \right) e^{-\lambda_{min} x} - \frac{q_{12}q_{21}}{D} e^{-\lambda_{max} x} \right\} P \Theta^2 dx \quad (3.8)$$

$$- \int_{\mathbb{R}^d} \frac{P}{2(2\mu + \lambda)} \left\{ \left(1 + \frac{q_{12}q_{21}}{D} \right) e^{-\lambda_{max} x} - \frac{q_{12}q_{21}}{D} e^{\lambda_{min} x} \right\} U^2 dx \quad (3.9)$$

$$+ \int_{\mathbb{R}^d} \left\{ (e^{-\lambda_{min} x} - e^{-\lambda_{max} x}) \frac{1}{D} \left(\frac{q_{12}}{2(2\mu + \lambda)} - \frac{q_{21}C_V}{\kappa} \right) \right\} P U \Theta dx. \quad (3.10)$$

Now we estimate (3.10). We only estimate the terms with the slowest exponential, the other two terms can clearly be estimated in the same way.

As we did previously, we distinguish between r small and r large.

We start with the slowest exponential term in (3.10):

$$\int_{\mathbb{R}^d} e^{-\lambda_{\min}} \frac{1}{D} \frac{q_{12}}{2\mu + \lambda} P U \Theta \, dx = \int_0^\varepsilon e^{-\lambda_{\min}} \frac{q_{12}}{D} \frac{P}{2\mu + \lambda} U \Theta \, dx \quad (3.11)$$

$$+ \int_\varepsilon^{+\infty} e^{-\lambda_{\min}} \frac{q_{12}}{D} \frac{P}{2\mu + \lambda} U \Theta \, dx, \quad (3.12)$$

and we estimate both pieces separately.

In the case where r is small, we use Lemma 3.11 (more precisely the bound $|q_{12}| \leq \sqrt{\frac{\beta}{\gamma}}$ in the proof), and get:

$$\begin{aligned} |(3.11)| &\leq \int_0^\varepsilon e^{-\lambda_{\min}} \sqrt{\frac{\int_0^r P \Theta}{\kappa P^\alpha \int_0^r P^{1-\alpha}}} \frac{\sqrt{P}}{\sqrt{2\mu + \lambda}} |U| \sqrt{P} \Theta \, dx \\ &\leq \frac{1}{2} \int_0^\varepsilon e^{-\lambda_{\min}} \sqrt{\frac{\int_0^r P \Theta}{\kappa P^\alpha \int_0^r P^{1-\alpha}}} \left(\frac{P}{2\mu + \lambda} U^2 + P \Theta^2 \right) dx \\ &\leq \frac{1}{2} \int_0^\varepsilon e^{-\lambda_{\min}} \sqrt{\frac{\Lambda \sup_{r>0} \Theta}{\kappa}} P \Theta^2 \, dx \\ &\quad + \frac{1}{2} \sup_{0 < r < \varepsilon} e^{2\sqrt{\Delta}} \int_0^\varepsilon e^{-\lambda_{\max}} \sqrt{\frac{\Lambda \sup_{r>0} \Theta}{\kappa}} \frac{P}{2\mu + \lambda} U^2 \, dx, \end{aligned}$$

and we can conclude with the smallness conditions.

Now we move on to the second part, using Lemma 3.9:

$$|(3.12)| \leq \frac{2R\Lambda \sup_{r>0} \Theta}{\kappa(1/2 - b)} \sup_{r>\varepsilon} \left| \frac{U}{r\Theta} \right| \int_{\mathbb{R}^d} e^{-\lambda_{\min}} P \Theta^2 \, dx,$$

and we can conclude using the smallness assumptions.

For the remaining term, we write

$$\frac{C_V}{\kappa} \int_{\mathbb{R}^d} \frac{q_{21}}{D} e^{-\lambda_{\min}} P \Theta U \, dx \leq \frac{2R\Lambda C_V}{\kappa(1/2 - b)} \sup_{r>0} \left| \frac{U}{r\Theta} \right| \int_{\mathbb{R}^d} e^{-\lambda_{\min}} P \Theta^2 \, dx.$$

Bound on RHS₃:

All these terms are treated as error terms, and the proofs are easier or similar to the above. Therefore we only show how to estimate the main terms.

Main contribution from N_1 :

For the first term, we simply use Remark 3.5 and write

$$\int_{\mathbb{R}^d} \frac{R}{\kappa} e^{-\lambda_{\min}} P \Theta^2 \frac{d-1}{r} U \, dr \leq \frac{R(d-1)b}{\kappa} \int_{\mathbb{R}^d} e^{-\lambda_{\min}} P \Theta^2 \, dr.$$

Next, we write

$$\begin{aligned} \int_{\mathbb{R}^d} \frac{\mu}{\kappa} e^{-\lambda_{\min}} U'^2 \Theta \, dx &\leq \frac{\mu_0}{\kappa} \sup_{r>0} P^\alpha \Theta \sup_{0 < r < \varepsilon} e^{2\sqrt{\Delta}} \int_0^\varepsilon e^{-\lambda_{\max}} U'^2 \, dx \\ &\quad + \frac{\mu_0}{\kappa} \sup_{r>\varepsilon} \left| \frac{U'}{r\Theta'} \right| \sup_{r>0} P^\alpha r^2 \Theta \int_\varepsilon^{+\infty} e^{-\lambda_{\min}} \Theta'^2 \, dx. \end{aligned}$$

This is acceptable given our smallness assumptions.
Similarly

$$\begin{aligned} \int_{\mathbb{R}^d} \frac{\mu}{\kappa} e^{-\lambda_{\min}} \frac{(d-1)U^2}{r^2} \Theta \, dx &\leq \frac{\mu_0}{\kappa} \sup_{r>0} P^\alpha \Theta \sup_{0<r<\varepsilon} e^{2\sqrt{\Delta}} \int_0^\varepsilon e^{-\lambda_{\max}} \frac{(d-1)^2 U^2}{r^2} \, dx \\ &+ \frac{\mu_0}{\kappa} \sup_{r>\varepsilon} P^{\alpha-1} \Theta \frac{U^2}{r^2 \Theta^2} \int_\varepsilon^{+\infty} e^{-\lambda_{\min}} P \Theta^2 \, dx. \end{aligned}$$

The other main terms are bounded in the same way.

Main contribution from N_2 :

We write that

$$\int_{\mathbb{R}^d} e^{-\lambda_{\max}} \alpha \frac{U}{r/2 + U} U'^2 \, dx \leq \frac{\alpha b}{\frac{1}{2} - b} \int_{\mathbb{R}^d} e^{-\lambda_{\max}} U'^2 \, dx.$$

The other main terms are bounded in the same way. \square

Conclusion of the proof of Theorem 3.1. The result follows directly by putting together Lemmas 3.12 and 3.13. \square

REFERENCES

- [1] D. Bresch, B. Desjardins, C.-K. Lin, On compressible fluid models: Korteweg, lubrication and shallow water systems, *Comm. Partial Differential Equations* 28 (2003), no. 3-4, 843–868
- [2] M. Cannone, F. Planchon, Self-similar solutions for Navier-Stokes equations in \mathbf{R}^3 , *Comm. Partial Differential Equations* 21 (1996), no. 1-2, 179–193.
- [3] J.-Y. Chemin, Théorèmes d’unicité pour le système de Navier-Stokes tridimensionnel, *J. Anal. Math.* 77 (1999), 27–50.
- [4] N. Chikami, R. Danchin, On the well-posedness of the full compressible Navier-Stokes system in critical Besov spaces, *J. Differential Equations* 258 (2015), no. 10, 3435–3467.
- [5] R. Danchin, Global existence in critical spaces for flows of compressible viscous and heat-conductive gases, *Arch. Ration. Mech. Anal.* 160 (2001), no. 1, 1–39.
- [6] E. Feireisl, *Dynamics of viscous compressible fluids*. Oxford Lecture Series in Mathematics and its Applications, 26. Oxford University Press, Oxford, 2004.
- [7] E. Feireisl, A. Novotný, H. Petzeltová, On the existence of globally defined weak solutions to the Navier-Stokes equations, *J. Math. Fluid Mech.* 3 (2001), no. 4, 358–392.
- [8] P. Germain, T. Iwabuchi, Self-similar solutions for compressible Navier-Stokes equations, *arXiv:1903.09958*
- [9] P. Germain, T. Iwabuchi, T. Léger, Backward self-similar solutions for compressible Navier-Stokes equations, *arXiv:1911.00339*
- [10] Z. Guo, S. Jiang, Self-similar solutions to the isothermal compressible Navier-Stokes equations, *IMA J. Appl. Math.* 71 (2006), 658–669.
- [11] D. Hoff, J. Smoller, Non-formation of vacuum states for compressible Navier-Stokes equations, *Comm. Math. Phys.* 216 (2001), no. 2, 255–276.
- [12] X. Huang, J. Li, , Global classical and weak solutions to the three-dimensional full compressible Navier-Stokes system with vacuum and large oscillations, *Arch. Ration. Mech. Anal.* 227 (2018), no. 3, 995–1059.
- [13] J. Jang, N. Masmoudi, Well-posedness of compressible Euler equations in a physical vacuum, *Comm. Pure Appl. Math.* 68 (2015), no. 1, 61–111.
- [14] J. Jang, N. Masmoudi, *Vacuum in gas and fluid dynamics*. Nonlinear conservation laws and applications, 315–329, IMA Vol. Math. Appl., 153, Springer, New York, 2011.
- [15] H. Jia, V. Šverák, Local-in-space estimates near initial time for weak solutions of the Navier-Stokes equations and forward self-similar solutions, *Invent. Math.* 196 (2014), no. 1, 233–265.

- [16] H. Koch and D. Tataru, Well-posedness for the Navier-Stokes equations, *Adv. Math.* 157 (2001), no. 1, 22–35.
- [17] T. Li, P. Chen, J. Xie, Self-similar solutions of the compressible flow in one-space dimension, *J. Appl. Math.* (2013), Art. ID 194704, 5.
- [18] J. Leray, Sur le mouvement d'un liquide visqueux emplissant l'espace, *Acta Math.* 63 (1934), 193 - 248
- [19] P.-L. Lions, *Mathematical topics in fluid mechanics. Vol. 1. Incompressible models.* Oxford Lecture Series in Mathematics and its Applications, 3. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1996.
- [20] P.-L. Lions, *Mathematical topics in fluid mechanics. Vol. 2. Compressible models.* Oxford Lecture Series in Mathematics and its Applications, 10. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1998.
- [21] J. Nečas, M. Růžička, V. Šverák, On Leray's self-similar solutions of the Navier-Stokes equations, *Acta Math.* 176 (1996), 283 - 294
- [22] A. Matsumura, T. Nishida, The initial value problem for the equations of motion of viscous and heat-conductive gases, *J. Math. Kyoto Univ.* 20 (1980), no. 1, 67–104.
- [23] Y. Qin, X. Su, S. Deng, Remarks on self-similar solutions to the compressible Navier-Stokes equations of a 1D viscous polytropic ideal gas, *Appl. Math. Sci. (Ruse)* 2 (2008), 1493–1506.
- [24] T.-P. Tsai, On Leray's self-similar solutions of the Navier-Stokes equations satisfying local energy estimates, *Arch. Rational Mech. Anal.* 143 (1998), 29-51
- [25] A. Vasseur, C. Yu, Existence of global weak solutions for 3D degenerate compressible Navier-Stokes equations, *Invent. Math.* 206 (2016), no. 3, 935–974
- [26] Z. Xin, Blowup of smooth solutions to the compressible Navier-Stokes equation with compact density, *Comm. Pure Appl. Math.* 51 (1998), no. 3, 229–240.