

ON ORTHOGONAL PROJECTIONS OF SYMPLECTIC BALLS

NUNO C. DIAS, MAURICE A. DE GOSSON, AND JOÃO N. PRATA

ABSTRACT. We study the orthogonal projections of symplectic balls in \mathbb{R}^{2n} on complex subspaces. In particular we show that these projections are themselves symplectic balls under a certain complexity assumption. Our main result is a refinement of a recent very interesting result of Abbondandolo and Matveyev extending the linear version of Gromov's non-squeezing theorem. We use a conceptually simpler approach where the Schur complement of a matrix plays a central role.

CONTENTS

1. Introduction	1
1.1. What is known	1
1.2. What we will do	2
2. Preliminaries	3
2.1. Williamson's symplectic diagonalization	3
2.2. Block-matrix partitions and Schur complements	4
2.3. Orthogonal projections of ellipsoids in \mathbb{R}^{2n}	4
3. Orthogonal Projections of Symplectic Balls	5
3.1. The main result: statement and proof	5
3.2. Discussion and extension	7
4. Perspectives	9
References	9

1. INTRODUCTION

1.1. **What is known.** Let $\sigma = dp_1 \wedge dx_1 + \cdots + dp_n \wedge dx_n$ be the standard symplectic form on $\mathbb{R}^{2n} \cong \mathbb{R}^n \times \mathbb{R}^n$; we call *symplectic ball* the image of the ball

$$B^{2n}(z_0, R) = \{z \in \mathbb{R}^{2n} : |z - z_0| \leq R\}$$

by a symplectic automorphism $S \in \mathrm{Sp}(n)$ (the symplectic group of $(\mathbb{R}^{2n}, \sigma)$). As a consequence of Gromov's non-squeezing theorem [7, 9] the orthogonal projection of a symplectic ball $S(B^{2n}(z_0, R))$ on any two-dimensional symplectic subspace of $(\mathbb{R}^{2n}, \sigma)$ has area at least equal to πR^2 . Abbondandolo

Key words and phrases. Symplectic ball, orthogonal projection, Gromov's non-squeezing theorem,

and Matveyev asked in [1] the question whether this result could be generalized to subspaces with higher dimensions. They showed that the orthogonal projection $\Pi_{\mathbb{V}}S(B^{2n}(z_0, R))$ of $S(B^{2n}(z_0, R))$ onto an arbitrary complex symplectic subspace $(\mathbb{V}, \sigma_{\mathbb{V}})$ of $(\mathbb{R}^{2n}, \sigma)$ such that $\dim \mathbb{V} = 2k$ satisfies

$$(1.1) \quad \text{Vol}_{\mathbb{V}} \Pi_{\mathbb{V}}S(B^{2n}(z_0, R)) \geq \frac{(\pi R^2)^k}{k!}$$

where $\text{Vol}_{\mathbb{V}}$ is the volume element on \mathbb{V} . Notice that $(\pi R^2)^k/k!$ is the volume of the ball $B_{\mathbb{V}}(\Pi_{\mathbb{V}}z_0, R)$ in \mathbb{V} :

$$(1.2) \quad \text{Vol}_{\mathbb{V}}(B_{\mathbb{V}}(\Pi_{\mathbb{V}}z_0, R)) = \frac{(\pi R^2)^k}{k!}.$$

They moreover proved that equality holds in (1.1) if and only if $S^T\mathbb{V}$ is itself a complex subspace of \mathbb{R}^{2n} . The inequality (1.1) implies the linear version of Gromov's theorem when $\dim \mathbb{V} = 2$ and conservation of volume by linear symplectomorphisms when $\mathbb{V} = \mathbb{R}^{2n}$. Abbondandolo and Matveyev proved their results using an ingenious but complicated argument involving the Wirtinger inequality for 2-forms on Kähler manifolds [3]. Results of this type are more subtle and difficult than they might appear at first sight; for instance as Abbondandolo and Matveyev show the inequality (1.1) does not hold when one replaces S by a nonlinear symplectomorphism f . In fact, one can construct examples where $\text{Vol}_{\mathbb{V}} \Pi_{\mathbb{V}}f(B^{2n}(R))$ can become arbitrarily small. They however make an interesting conjecture, to which we will come back at the end of this paper.

1.2. What we will do. We will prove by elementary means a stronger version of (1.1) and of its extension. We will actually prove (Theorem 3.1) that the orthogonal projection of a symplectic ball on a symplectic subspace contains a symplectic ball with the same radius in this subspace, and is itself a symplectic ball when the subspace under consideration is complex. The proof will be done in the particular case where the symplectic space \mathbb{V} is of the type $\mathbb{R}^{2n_A} \oplus 0$ in which case the symplectic orthogonal \mathbb{V}^{σ} is $0 \oplus \mathbb{R}^{2n_B}$; our refinement of (1.1) says that for every $S \in \text{Sp}(n)$ there exists $S_A \in \text{Sp}(n_A)$ (the symplectic group of $\mathbb{R}^{2n_A} \oplus 0 \equiv \mathbb{R}^{2n_A^1}$) such that

$$(1.3) \quad \Pi_{\mathbb{V}}S(B^{2n}(z_0, R)) \supseteq S_A(B^{2n_A}(z_{0,A}, R))$$

and $z_{0,A} = \Pi_{\mathbb{V}}z_0$.

This will be done using the theory of Schur complements and the notion of symplectic spectrum of a positive definite matrix. Since symplectomorphisms are volume-preserving, (1.3) implies (1.1). It is however a much stronger statement than (1.1) because, given two measurable sets Ω and Ω' with the same volume, there does not in general exist a symplectomorphism (let alone a linear one) taking Ω to Ω' as soon as the dimension of the symplectic space exceeds two [9].

¹For the sake of simplicity, we make the identification $\mathbb{R}^{2n_A} \oplus 0 \equiv \mathbb{R}^{2n_A}$. In particular, we write by abuse of language $\Pi_{\mathbb{V}}(z_A, z_B) = z_A$ instead of $(z_A, 0)$.

Note that these results are invariant under phase space translations. We will therefore assume henceforth that $z_0 = 0$.

We finally discuss in section 4 some possible extensions to the non-linear case, pointing out the difficulties.

2. PRELIMINARIES

In what follows M will be a real $2n \times 2n$ positive definite matrix; we will write $M > 0$. We denote by J the standard symplectic matrix $\begin{pmatrix} 0_n & I_n \\ -I_n & 0_n \end{pmatrix}$. We have, $\sigma(z, z') = Jz \cdot z'$ when $z = (x, p)$, $z' = (x', p')$. In this notation the condition $S \in \text{Sp}(n)$ is equivalent to $S^T JS = J$ (or $SJS^T = J$) where S^T is the transpose of S .

2.1. Williamson's symplectic diagonalization. By definition the symplectic spectrum of M is the increasing sequence $\lambda_1^\sigma(M) \leq \lambda_2^\sigma(M) \leq \dots \leq \lambda_n^\sigma(M)$ of numbers $\lambda_j^\sigma(M) > 0$ where the $\pm i\lambda_j^\sigma(M)$ are the eigenvalues of JM (which are the same as those of the antisymmetric matrix $M^{1/2}JM^{1/2}$). We will use the following property, known in the literature as ‘‘Williamson's symplectic diagonalization theorem’’ [4, 9]: there exists $S \in \text{Sp}(n)$ such that

$$M = S^T DS, \quad D = \begin{pmatrix} \Lambda & 0 \\ 0 & \Lambda \end{pmatrix}$$

where Λ is the diagonal matrix whose eigenvalues are the numbers $\lambda_j^\sigma(M)$ (all matrices corresponding here to the standard splitting $z = (x, p)$). The symplectic spectra of M and M^{-1} are inverses of each other in the sense that:

$$(2.1) \quad \lambda_j^\sigma(M^{-1}) = \lambda_{n-j}^\sigma(M)^{-1} \quad \text{for } 1 \leq j \leq n.$$

We also have the less obvious property ([4], section 8.3.2)

$$(2.2) \quad M \leq N \implies \lambda_j^\sigma(M) \leq \lambda_j^\sigma(N) \quad \text{for } 1 \leq j \leq n$$

where $M \leq N$ means $N - M \geq 0$.

The following simple result characterizing positive semi-definiteness in terms of the symplectic spectrum will be very useful for proving Theorem 3.1:

Lemma 2.1. *The Hermitian matrix $M + iJ$ is positive semi-definite: $M + iJ \geq 0$ if and only if $\lambda_j^\sigma(M) \geq 1$ for $1 \leq j \leq n$.*

Proof. Let $M = S^T DS$ be a Williamson diagonalization of M ; since $S^T JS = J$ the condition $M + iJ \geq 0$ is equivalent to $D + iJ \geq 0$. The characteristic polynomial of $D + iJ$ is the product $P(\lambda) = P_1(\lambda) \cdots P_n(\lambda)$ of polynomials

$$P_j(\lambda) = \lambda^2 - 2\lambda_j^\sigma(M)\lambda + \lambda_j^\sigma(M)^2 - 1$$

and the eigenvalues of $M + iJ$ are thus the numbers $\lambda_j = \lambda_j^\sigma(M) \pm 1$. The condition $M + iJ \geq 0$ is equivalent to $\lambda_j \geq 0$, that is to $\lambda_j^\sigma(M) \geq 1$ for $j = 1, \dots, n$. \square

Notice that if $S \in \text{Sp}(n)$ we have

$$(2.3) \quad S^T S + iJ \geq 0 \quad \text{and} \quad S S^T + iJ \geq 0$$

since $\lambda_j^\sigma(S^T S) = \lambda_j^\sigma(S S^T) = 1$ for all j (because $D = I$ in view of Williamson's diagonalization result).

2.2. Block-matrix partitions and Schur complements. Let $\mathbb{R}^{2n_A} \equiv \mathbb{R}^{2n_A} \oplus 0$ and $\mathbb{R}^{2n_B} \equiv 0 \oplus \mathbb{R}^{2n_B}$ be two symplectic subspaces of \mathbb{R}^{2n} . We split $(\mathbb{R}^{2n}, \sigma)$ as a direct sum $(\mathbb{R}^{2n_A} \oplus \mathbb{R}^{2n_B}, \sigma_A \oplus \sigma_B)$ where σ_A and σ_B are, respectively, the restrictions of σ to \mathbb{R}^{2n_A} and \mathbb{R}^{2n_B} . We write $z \in \mathbb{R}^{2n}$ as $z = (z_A, z_B) = z_A \oplus z_B$ with $z_A = (x_A, p_A) \in \mathbb{R}^{2n_A}$ and $z_B = (x_B, p_B) \in \mathbb{R}^{2n_B}$.

We denote by Π_A (*resp.* Π_B) the orthogonal projection $\mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n_A}$ (*resp.* \mathbb{R}^{2n_B}). We choose symplectic bases $\mathcal{B}_A, \mathcal{B}_B$ in \mathbb{R}^{2n_A} and \mathbb{R}^{2n_B} and identify linear mappings $\mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ with their matrices in the symplectic basis $\mathcal{B} = \mathcal{B}_A \oplus \mathcal{B}_B$ of \mathbb{R}^{2n} . Such a matrix will be written as

$$(2.4) \quad M = \begin{pmatrix} M_{AA} & M_{AB} \\ M_{BA} & M_{BB} \end{pmatrix}$$

the blocks $M_{AA}, M_{AB}, M_{BA}, M_{BB}$ having dimensions $2n_A \times 2n_A, 2n_A \times 2n_B, 2n_B \times 2n_A, 2n_B \times 2n_B$, respectively. Similarly, the standard symplectic matrix J will be split as

$$J = J_A \oplus J_B \equiv \begin{pmatrix} J_A & 0 \\ 0 & J_B \end{pmatrix}$$

where J_A (*resp.* J_B) is the standard symplectic matrix in $(\mathbb{R}^{2n_A}, \sigma_A)$ (*resp.* $(\mathbb{R}^{2n_B}, \sigma_B)$).

Since M is positive definite and symmetric the upper-left and lower-right blocks in (2.4) are themselves positive-definite and symmetric: $M_{AA} > 0$ and $M_{BB} > 0$. In particular the Schur complements

$$(2.5) \quad M/M_{BB} = M_{AA} - M_{AB}M_{BB}^{-1}M_{BA}$$

$$(2.6) \quad M/M_{AA} = M_{BB} - M_{BA}M_{AA}^{-1}M_{AB}$$

are well defined and invertible [13], and the inverse of the matrix M is given by the formula

$$(2.7) \quad M^{-1} = \begin{pmatrix} (M/M_{BB})^{-1} & -(M/M_{BB})^{-1}M_{AB}M_{BB}^{-1} \\ -M_{BB}^{-1}M_{BA}(M/M_{BB})^{-1} & (M/M_{AA})^{-1} \end{pmatrix}.$$

2.3. Orthogonal projections of ellipsoids in \mathbb{R}^{2n} . We will also need the following general characterization of the orthogonal projection of an ellipsoid on a subspace:

Lemma 2.2. *Let Π_A be the orthogonal projection $\mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n_A}$ and*

$$\Omega = \{z \in \mathbb{R}^{2n} : Mz^2 \leq R^2\}.$$

We have

$$(2.8) \quad \Pi_A \Omega = \{z_A \in \mathbb{R}^{2n_A} : (M/M_{BB})z_A^2 \leq R^2\}.$$

Proof. Let us set $Q(z) = Mz^2 - R^2$; the boundary $\partial\Omega$ of the hypersurface $Q(z) = 0$ is defined by

$$(2.9) \quad M_{AA}z_A^2 + 2M_{BA}z_A \cdot z_B + M_{BB}z_B^2 = R^2.$$

A point z_A belongs to $\partial\Pi_A\Omega$ if and only if the normal vector to $\partial\Omega$ at the point $z = (z_A, z_B)$ is parallel to \mathbb{R}^{2n_A} hence the constraint

$$\partial_z Q(z) = 2Mz \in \mathbb{R}^{2n_A} \oplus 0.$$

This is equivalent to the condition $M_{BA}z_A + M_{BB}z_B = 0$, that is to $z_B = -M_{BB}^{-1}M_{BA}z_A$. Inserting z_B in (2.9) shows that the boundary $\partial\Pi_A\Omega$ is the set of all $z_A \in \mathbb{R}^{2n_A}$ such that $(M/M_{BB})z_A^2 = R^2$ hence formula (2.8). \square

Interchanging A and B the orthogonal projection of Ω on \mathbb{R}^{2n_B} is similarly given by

$$(2.10) \quad \Pi_B \Omega = \{z_B \in \mathbb{R}^{2n_B} : (M/M_{AA})z_B^2 \leq R^2\}.$$

3. ORTHOGONAL PROJECTIONS OF SYMPLECTIC BALLS

3.1. The main result: statement and proof. Let us now prove the main result. We assume again the matrix M is written in block-form (2.4). To simplify notation we also assume that all balls $B^{2n}(z_0, R)$ are centered at the origin and set $B^{2n}(0, R) = B^{2n}(R)$. The case of a general ball $B^{2n}(z_0, R)$ trivially follows using the translation $z \mapsto z + z_0$.

Theorem 3.1. *Let $S \in \text{Sp}(n)$. (i) There exists $S_A \in \text{Sp}(n_A)$ such that*

$$(3.1) \quad \Pi_A(S(B^{2n}(R))) \supseteq S_A(B^{2n_A}(R));$$

(ii) We have

$$(3.2) \quad \Pi_A(S(B^{2n}(R))) = S_A(B^{2n_A}(R))$$

if and only if $S = S_A \oplus S_B$ for some $S_B \in \text{Sp}(n_B)$, in which case we also have

$$(3.3) \quad \Pi_B(S(B^{2n}(R))) = S_B(B^{2n_B}(R))$$

Proof. (i) The symplectic ball $S(B^{2n}(R))$ consists of all $z \in \mathbb{R}^{2n}$ such that $Mz^2 \leq R^2$ where $M = (SS^T)^{-1}$. It follows from Lemma 2.2 that $\Pi_A S(B^{2n}(R))$ is determined by the inequality $(M/M_{BB})z_A^2 \leq R^2$. We are going to show that the symplectic eigenvalues $\lambda_j^{\sigma_A}(M/M_{BB})$ are all ≤ 1 . The inclusion (3.1) will then follow since we have, in view of Williamson's diagonalization result,

$$(3.4) \quad M/M_{BB} = (S_A^{-1})^T D_A S_A^{-1}$$

for some $S_A \in \text{Sp}(n_A)$ and

$$(3.5) \quad D_A = \begin{pmatrix} \Lambda_A & 0 \\ 0 & \Lambda_A \end{pmatrix}, \quad \Lambda_A = \text{diag}(\lambda_1^{\sigma_A}(M/M_{BB}), \dots, \lambda_{n_A}^{\sigma_A}(M/M_{BB})).$$

It follows that:

$$(3.6) \quad \begin{aligned} (M/M_{BB}) z_A^2 &= ((S_A^{-1})^T D_A S_A^{-1}) z_A^2 \\ &= D_A (S_A^{-1} z_A)^2 \leq |S_A^{-1} z_A|^2 \end{aligned}$$

The condition $z \in S_A(B^{2n_A}(R))$ being equivalent to $|S_A^{-1} z_A|^2 \leq R^2$ thus implies $(M/M_{BB}) z_A^2 \leq R^2$ and hence $S_A(B^{2n_A}(R)) \subseteq \Pi_A S(B^{2n}(R))$.

To prove that we indeed have

$$(3.7) \quad \lambda_j^{\sigma_A}((M/M_{BB})) \leq 1 \text{ for } 1 \leq j \leq n_A$$

we begin by noting that the symplectic eigenvalues $\lambda_j^\sigma(M)$ of $M = (SS^T)^{-1}$ are all trivially equal to one, and hence also those of its inverse M^{-1} : $\lambda_j^\sigma(M^{-1}) = 1$ for $1 \leq j \leq n$. In view of Lemma 2.1 the Hermitian matrix $M^{-1} + iJ$ is positive semidefinite:

$$(3.8) \quad M^{-1} + iJ \geq 0$$

which implies that (cf.(2.7))

$$(3.9) \quad (M/M_{BB})^{-1} + iJ_A \geq 0$$

(recall that $J = J_A \oplus J_B$). Applying now Lemma 2.1 to $(M/M_{BB})^{-1}$ this implies that the inequalities (3.7) must hold (cf.(2.1)).

(ii) In view of (2.8) in Lemma 2.2 the equality (3.2) will hold if and only if $M/M_{BB} = (S_A S_A^T)^{-1}$. If $S = S_A \oplus S_B$, then $M = (SS^T)^{-1}$ implies $M/M_{BB} = (S_A S_A^T)^{-1}$ and we have the equality (3.2).

Conversely, suppose the equality (3.2) holds. From the inversion formula (2.7) we have

$$(3.10) \quad M^{-1} = \begin{pmatrix} (M/M_{BB})^{-1} & X \\ X^T & (M/M_{AA})^{-1} \end{pmatrix}.$$

where $X = -(M/M_{BB})^{-1} M_{AB} M_{BB}^{-1}$; since $M^{-1} \in \text{Sp}(n)$ is symmetric it satisfies the symplecticity condition

$$M^{-1}(J_A \oplus J_B)M^{-1} = J_A \oplus J_B$$

which is in turn equivalent to the set of conditions

$$(3.11) \quad (M/M_{BB})^{-1} J_A (M/M_{BB})^{-1} + X J_B X^T = J_A$$

$$(3.12) \quad (M/M_{BB})^{-1} J_A X + X J_B (M/M_{AA})^{-1} = 0$$

$$(3.13) \quad X^T J_A X + (M/M_{AA})^{-1} J_B (M/M_{AA})^{-1} J_B.$$

Since

$$(3.14) \quad M/M_{BB} = (S_A S_A^T)^{-1} \in \text{Sp}(n_A),$$

it follows from (3.11) that

$$(3.15) \quad X J_B X^T = 0.$$

Multiplying the identity (3.12) on the right by $(M/M_{AA})X^T$ and using (3.15), we obtain

$$(M/M_{BB})^{-1}J_A X(M/M_{AA})X^T = 0$$

that is

$$(3.16) \quad X(M/M_{AA})X^T = 0.$$

Since $(M/M_{AA}) > 0$, this is possible if and only if $X = 0$. Finally, from (3.13) we conclude that $(M/M_{AA})^{-1} \in \text{Sp}(n_B)$. Moreover, since $(M/M_{AA})^{-1}$ is symmetric and positive definite, there exists $S_B \in \text{Sp}(n_B)$, such that $(M/M_{AA})^{-1} = S_B S_B^T$.

Altogether

$$M^{-1} = S S^T = \begin{pmatrix} S_A S_A^T & 0 \\ 0 & S_B S_B^T \end{pmatrix}$$

hence $S = S_A \oplus S_B$, which concludes the proof. \square

We remark that the proof above actually provides the means to calculate explicitly the symplectic automorphisms S_A in (3.1). Recapitulating, it is constructed as follows: given $S \in \text{Sp}(n)$ calculate

$$M = (S S^T)^{-1} = \begin{pmatrix} M_{AA} & M_{AB} \\ M_{BA} & M_{BB} \end{pmatrix}$$

and then obtain the Schur complement (2.5)

$$M/M_{BB} = M_{AA} - M_{AB} M_{BB}^{-1} M_{BA}.$$

The matrix S_A is then obtained from (3.14) (observe that S_A is only defined up to a symplectic rotation, but this ambiguity is irrelevant since $B^{2n_A}(R)$ is rotationally invariant).

3.2. Discussion and extension. Theorem 3.1 implies *de facto* the Abbondandolo and Matveyev result (1.1) since formula (3.1) has the immediate consequence that

$$(3.17) \quad \text{Vol}_{2n_A} \Pi_A(S(B^{2n}(R))) \geq \frac{(\pi R^2)^{n_A}}{n_A!}$$

noting that $S_A \in \text{Sp}(n_A)$ is volume-preserving. Similarly, the equality (3.2) implies

$$(3.18) \quad \text{Vol}_{2n_A} \Pi_A(S(B^{2n}(R))) = \frac{(\pi R^2)^{n_A}}{n_A!}.$$

Abbondandolo and Matveyev's [1] however prove these relations for projections on a general complex symplectic subspace \mathbb{V} of $(\mathbb{R}^{2n}, \sigma)$ (that is such that $J\mathbb{V} = \mathbb{V}$) and they show that the equality in (1.1) holds if and only if the subspace $S^T\mathbb{V}$ is complex, that is if $J S^T\mathbb{V} = S^T\mathbb{V}$. Let us check that these conditions are satisfied when $\mathbb{V} = \mathbb{R}^{2n_A} \oplus 0$. We first note that

$$J\mathbb{V} = (J_A \oplus J_B)(\mathbb{R}^{2n_A} \oplus 0) = \mathbb{R}^{2n_A} \oplus 0 = \mathbb{V}$$

hence $\mathbb{R}^{2n_A} \oplus 0$ is complex in the sense above. Similarly, the condition $JS^T\mathbb{V} = S^T\mathbb{V}$ is satisfied if $S = S_A \oplus S_B$ since

$$(J_A \oplus J_B)(S_A^T \oplus S_B^T)(\mathbb{R}^{2n_A} \oplus 0) = \mathbb{R}^{2n_A} \oplus 0;$$

conversely the condition $JS^T\mathbb{V} = S^T\mathbb{V}$ is equivalent to

$$(SS^T)^{-1}J\mathbb{V} = \mathbb{V}$$

and if $\mathbb{V} = \mathbb{R}^{2n_A} \oplus 0$ this is possible for $S \in \text{Sp}(n)$ if and only if $S = S_A \oplus S_B$.

To show that Theorem 3.1 implies Abbondandolo and Matveyev's result for arbitrary complex subspaces \mathbb{V} is actually straightforward. To see this, let \mathbb{V}^σ be the symplectic orthocomplement of \mathbb{V} in $(\mathbb{R}^{2n}, \sigma)$. Choose symplectic bases $\mathcal{B}_\mathbb{V}$ of \mathbb{V} and $\mathcal{B}_{\mathbb{V}^\sigma}$ of \mathbb{V}^σ such that their union \mathcal{B} is a symplectic orthonormal basis of \mathbb{R}^{2n} (this is easily done using the symplectic version of the Gram–Schmidt construction for orthogonal bases; see [4]). Set $\dim \mathbb{V} = 2n_A$ and $\dim \mathbb{V}^\sigma = 2n_B$ (hence $n = n_A + n_B$) and let \mathcal{B}_A and \mathcal{B}_B be symplectic bases of $(\mathbb{R}^{2n_A}, \sigma_A)$ and $(\mathbb{R}^{2n_B}, \sigma_B)$, respectively, such that $\mathcal{B}_A \cup \mathcal{B}_B$ is a symplectic orthonormal basis of $(\mathbb{R}^{2n}, \sigma)$. Let U be the linear mapping $\mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ defined by $\mathcal{B}_\mathbb{V} = U(\mathcal{B}_A)$ and $\mathcal{B}_{\mathbb{V}^\sigma} = U(\mathcal{B}_B)$; clearly U is symplectic and orthogonal. The mapping $\Pi_\mathbb{V} : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ defined by $\Pi_\mathbb{V} = U\Pi_A U^{-1}$ is then the projection onto \mathbb{V} along \mathbb{V}^σ and we thus have

$$\Pi_\mathbb{V}(S(B^{2n}(R))) = U\Pi_A U^{-1}(S(B^{2n}(R))).$$

Theorem 3.1 implies that there exists $S'_A \in \text{Sp}(n_A)$ such that

$$\Pi_A(U^{-1}(S(B^{2n}(R)))) \supseteq S'_A(B^{2n_A}(R))$$

and hence, since U and S'_A are volume preserving,

$$(3.19) \quad \text{Vol } \Pi_\mathbb{V}(S(B^{2n}(R))) \geq \frac{(\pi R^2)^{n_A}}{n_A!}$$

which is (1.1). The “exact” case in (3.19) is equivalent to

$$(3.20) \quad \text{Vol } \Pi_A(S'(B^{2n}(R))) = \frac{(\pi R^2)^{n_A}}{n_A!}$$

where $S' = U^{-1}S$ with $U \in U(n)$. Theorem 3.1 implies that the projection $\Pi_A(S'(B^{2n}(R)))$ is either identical to $S'_A(B^{2n_A}(R))$ (for some $S'_A \in \text{Sp}(n_A)$) or strictly larger than $S'_A(B^{2n_A}(R))$ (in which case its volume cannot be $(\pi R^2)^{n_A}/n_A!$); hence (3.20) holds if and only if

$$\Pi_A(S'(B^{2n_A}(R))) = S'_A(B^{2n_A}(R))$$

in which case $S' = S'_A \oplus S'_B$ for some $S'_B \in \text{Sp}(n_B)$ (cf. Theorem 3.1). Finally, in view of the discussion above we have the equivalences

$$S' = S'_A \oplus S'_B \Leftrightarrow JS'^T(\mathbb{R}^{2n_A} \oplus 0) = S'^T(\mathbb{R}^{2n_A} \oplus 0) \Leftrightarrow JS^T\mathbb{V} = S^T\mathbb{V}$$

where we used the identities $S'^T = S^T U$ and $U(\mathbb{R}^{2n_A} \oplus 0) = \mathbb{V}$.

4. PERSPECTIVES

A first natural question that arises is whether Theorem 3.1 can be extended in some way to non-linear symplectic mappings, that is to general symplectomorphisms of $(\mathbb{R}^{2n}, \sigma)$. The first answer is that there are formidable roadblocks to the passage from the linear to the nonlinear case, as shortly mentioned in the Introduction. For instance, Abbondandolo and Matveyev [1] show, elaborating on ideas of Guth [8], that for every $\varepsilon > 0$ one can find a symplectomorphism f of $(\mathbb{R}^{2n}, \sigma)$ defined near $B^{2n}(0, 1)$ such that

$$\text{Vol}(\Pi_{\mathbb{V}} f(B^{2n}(0, 1))) < \varepsilon.$$

They however speculate in [1] that their projection result might still hold true when the linear symplectic automorphism $S \in \text{Sp}(n)$ is replaced with a symplectomorphism f of $(\mathbb{R}^{2n}, \sigma)$ close to a linear one. It would be interesting to apply our methods to tackle this difficult problem.

Also, Theorem 3.1 could be used to shed some light on packing problems (see the review [12] by Schlenk) which form a notoriously difficult area of symplectic topology.

Given the partitioning of $\mathbb{R}^{2n} = \mathbb{R}^{2n_A} \oplus \mathbb{R}^{2n_B}$ it seems natural to expect some connection between orthogonal projections of symplectic balls and the separability/entanglement problem in quantum mechanics [6, 10, 11]. We intend to address this problem in a future work.

REFERENCES

- [1] A. Abbondandolo and R. Matveyev, How large is the shadow of a symplectic ball?, *J. Topol. Anal.* 5(01), (2013) 87–119
- [2] A. Abbondandolo and P. Majer, A non-squeezing theorem for convex symplectic images of the Hilbert ball, *Calc. Var.* 54, (2015) 1469–1506
- [3] H. Federer, *Geometric measure theory*, Springer, 1969
- [4] M. de Gosson, *Symplectic geometry and quantum mechanics*, Vol. 166. Springer Science & Business Media, 2006
- [5] M. de Gosson, *Symplectic methods in harmonic analysis and in mathematical physics*, Vol. 7. Springer Science & Business Media, 2011
- [6] M. de Gosson, On Density Operators with Gaussian Weyl Symbols, in *Proceedings to the Conference MLTFA18 – Microlocal and Time-Frequency Analysis*. Birkhäuser, series Applied Numerical and Harmonic Analysis (series editor: John J. Benedetto), 2018
- [7] M. Gromov, Pseudoholomorphic curves in symplectic manifolds. *Inv. Math.* 82(2), (1985) 307–347
- [8] L. Guth, Symplectic embeddings of polydisks, *Invent. Math.* 172 (2008), 477–489
- [9] H. Hofer and E. Zehnder, *Symplectic Invariants and Hamiltonian Dynamics*, Birkhäuser Advanced texts, (Basler Lehrbücher, Birkhäuser Verlag, (1994)
- [10] L. Lami, A. Serafini, and G. Adesso, Gaussian entanglement revisited, *New J. of Phys.* 20, (2018) 023030
- [11] R. F. Werner and M. M. Wolf, Bound Entangled Gaussian States, *Phys. Rev. Lett.* 86(16) (2001) 3658
- [12] F. Schlenk, Symplectic embedding problems, old and new, *Bull. Amer. Math. Soc.* 55, 139–182 (2018)
- [13] F. Zhang, *The Schur Complement and its Applications*, Springer, Berlin, 2005

Author's addresses:

- **Nuno Costa Dias** and **João Nuno Prata:** Escola Superior Náutica Infante D. Henrique. Av. Eng. Bonneville Franco, 2770-058 Paço d'Arcos, Portugal and Grupo de Física Matemática, Universidade de Lisboa, Av. Prof. Gama Pinto 2, 1649-003 Lisboa, Portugal
- **Maurice A. de Gosson:** University of Vienna, Faculty of Mathematics (NuHAG), Vienna, Austria.

E-mail address (N.C. Dias): ncdias@meo.pt

E-mail address (M. de Gosson): maurice.de.gosson@univie.ac.at

E-mail address (J.N. Prata): joao.prata@mail.telepac.pt
