arXiv:1911.03820v1 [math.NT] 10 Nov 2019

A CONTINUITY OF CYCLE INTEGRALS OF MODULAR FUNCTIONS

YUYA MURAKAMI

Abstract. In this paper we study a continuity of the "values" of modular functions at the real quadratic numbers which are defined in terms of their cycle integrals along the associated closed geodesics. Our main theorem reveals a more finer structure of the continuity of these values with respect to continued fraction expansions and it turns out that it is different from the continuity with respect to Euclidean topology.

1. Introduction

Let $\mathbb{H} := \{z = x + y\sqrt{-1} \in \mathbb{C} \mid x, y \in \mathbb{R}, y > 0\}$ be the upper half plane and $f \colon \mathbb{H} \to \mathbb{C}$ be a holomorphic modular function with respect to $\mathrm{SL}_2(\mathbb{Z})$. Kaneko [4] introduced the "value" of f at any real quadratic number w as follows. For such a w and its non-trivial Galois conjugate w' we define

$$\eta_w := \left(\frac{1}{z-w'} - \frac{1}{z-w}\right)dz.$$

Let $\mathrm{SL}_2(\mathbb{Z})_w$ be the stabilizer of w and $\gamma_w = \begin{pmatrix} * & * \\ c & d \end{pmatrix}$ be the unique element of $\mathrm{SL}_2(\mathbb{Z})_w$ such that $\mathrm{SL}_2(\mathbb{Z})_w = \{\pm\gamma_w^n \mid n \in \mathbb{Z}\}, \varepsilon_w := j(\gamma_w, w) := cw + d > 1$. For $z_0 \in \mathbb{H}$ we define

$$\widetilde{f}(w) := \int_{z_0}^{\gamma_w z_0} f\eta_w, \ \widetilde{1}(w) := \int_{z_0}^{\gamma_w z_0} \eta_w.$$

The integral is independent of z_0 since $f \circ \gamma_w = f, \gamma_w^* \eta_w = \eta_w$ and f is holomorphic. We set

$$f(w) := \frac{\widetilde{f}(w)}{\widetilde{1}(w)}.$$

This value has a representation as a cycle integral

$$f(w) = \frac{1}{\mathrm{length}(\mathrm{SL}_2(\mathbb{Z})_w \backslash S_w)} \int_{\mathrm{SL}_2(\mathbb{Z})_w \backslash S_w} f ds$$

where $ds := y^{-1}\sqrt{dx^2 + dy^2}$, S_w is the geodesic in $\mathbb{H}$ from w' to w and

$$\mathrm{length}(\mathrm{SL}_2(\mathbb{Z})_w \backslash S_w) := \int_{\mathrm{SL}_2(\mathbb{Z})_w \backslash S_w} ds = \widetilde{1}(w).$$

The values $f(w)$ and $\widetilde{f}(w)$ have been studied in the last decade. Duke, Imamoḡlu and Tóth [3] proved that a generating function of traces of these values is a mock modular form of weight $1/2$ for $\Gamma_0(4)$. This work is a real quadratic analog of Zagier's work on traces of singular moduli in [7]. Masri [5] studied equidistribution of traces of these values. Päpcke [6] showed the interesting formula

$$\lim_{n\to\infty} j_1\left(\frac{n + \sqrt{n^2-4}}{2}\right) = 0.$$

Date: March 13, 2024.

Key words and phrases. elliptic modular functions; cycle integrals; j-invariant; values at real quadratic numbers; continued fractions.

Mathematical Institute, Tohoku University, 6-3, Aoba, Aramaki, Aoba-Ku, Sendai 980-8578, JAPAN *E-mail address*: yuya.murakami.s8@dc.tohoku.ac.jp.

We can regard it as a certain continuity of $f(w)$. We remark that the real quadratic numbers in the left hand side have the continued fraction expansions

$$\frac{n+\sqrt{n^2-4}}{2} = n + \cfrac{1}{n + \cfrac{1}{n + \cfrac{1}{\ddots}}}.$$

Kaneko [4] conjectured another continuity of $f(w)$ when w is a Markov quadratic number and this conjecture is proved by Bengoechea and Imamoḡlu [2]. In this paper, we pursue more finer structure in the continuity with respect to the continued fraction expansions. Our main result is to show that the function $f(w)$ has a continuity with respect to how many cyclic parts appear in a continued fraction expansion of w.

Each real number can be represented by the unique continued fraction

$$[k_1, k_2, k_3, \dots] := k_1 + \cfrac{1}{k_2 + \cfrac{1}{k_3 + \cfrac{1}{\ddots}}}$$

where k_1 is an integer and $k_2, k_3, \dots$ are positive integers. Also we set a periodic continued fraction

$$[k_1, \dots, k_r, \overline{k_{r+1}, \dots, k_s}] := [k_1, \dots, k_r, k_{r+1}, \dots, k_s, k_{r+1}, \dots, k_s, k_{r+1}, \dots, k_s, \dots].$$

For a real number, being quadratic is equivalent to that it has a periodic continued fraction expansion.

For an even number $r \in 2\mathbb{Z}_{\geq 0}$ and positive integers $k_1, \dots, k_r$, we call $W = (k_1, \dots, k_r)$ an even word. For even words $W_1 = (k_1^{(1)}, \dots, k_{r_1}^{(1)}), \dots, W_n = (k_1^{(n)}, \dots, k_{r_n}^{(n)})$, define its product

$$W_1 \cdots W_n := (k_1^{(1)}, \dots, k_{r_1}^{(1)}, \dots, k_1^{(n)}, \dots, k_{r_n}^{(n)}).$$

For an even word $W = (k_1, \dots, k_r)$, let

$$[\overline{W}] := [\overline{k_1, \dots, k_r}],$$

$$N(W) := \max\{n \in \mathbb{Z}_{>0} \mid \exists \text{ an even word } W_1 \text{ s.t. } W = W_1^n\}.$$

Here, if $W = W_1^{N(W)}$, then $N(W_1) = 1$. For example, $N((1,2,1,2)) = 2, N((1,2)) = 1$ and $N((1,1)) = 1$.

Our main result is the following:

Theorem 1.1. *For sequences $\{a_n^{(1)}\}_{n=1}^{\infty}, \dots, \{a_n^{(k)}\}_{n=1}^{\infty}$ in $\mathbb{Z}_{\geq 0}$ diverging to the positive infinity, let*

$$a^{(i)} := \lim_{n\to\infty} \frac{a_n^{(i)}}{a_n^{(1)} + \cdots + a_n^{(k)}},\ 1 \leq i \leq k.$$

For even words $V_0, \dots, V_k$ and non-empty even words $W_1, \dots, W_k$, let

$$w_i := [\overline{W_i}],\ w^{(n)} := \overline{[V_0 W_1^{a_n^{(1)}} V_1 W_2^{a_n^{(2)}} V_2 \dots W_k^{a_n^{(k)}} V_k]}.$$

Then we have

$$\lim_{n\to\infty} f(w^{(n)}) = \frac{a^{(1)}N(W_1)\widetilde{f}(w_1) + \cdots + a^{(k)}N(W_k)\widetilde{f}(w_k)}{a^{(1)}N(W_1)\widetilde{1}(w_1) + \cdots + a^{(k)}N(W_k)\widetilde{1}(w_k)}.$$

Remark 1.2. (i) *For the case when $k = 1, a_n^{(1)} = n$, our main theorem says that*

$$\lim_{n\to\infty} f(\overline{[V_0 W_1^n V_1]}) = f(w_1).$$

In particular, if V_0, V_1, W_1 is products of $(1,1), (2,2)$ and $V_0 = \emptyset$ or $V_1 = \emptyset$, then this result is conjectured in [4] *and proved in* [2]. *Here a sequence $[\overline{V_0 W_1^n V_1}]$ converges to $\gamma_{V_0} w_1 = [V_0 \overline{W_1}]$ and its non-trivial Galois conjugate converges to w_1' in the sense of Euclidean metric by Lemma* 1.28 *in* [1].

(ii) *For the case when* $k = 2, a_n^{(1)} = a_n^{(2)} = n$ *and* $V_0 = V_1 = V_2 = \emptyset$*, our main theorem says that*

$$\lim_{n\to\infty} f([\overline{W_1^n W_2^n}]) = \frac{\widetilde{f}(w_1) + \widetilde{f}(w_2)}{\widetilde{1}(w_1) + \widetilde{1}(w_2)}.$$

Here a sequence $[\overline{W_1^n W_2^n}]$ *converges to* w_1 *and its non-trivial Galois conjugate converges to* w_2' *in the sense of Euclidean metric by Lemma* 1.28 *in* [1]*. Since* $\widetilde{f}(w_2) = \widetilde{f}(w_2')$ *and* $\widetilde{1}(w_2) = \widetilde{1}(w_2')$ *by a proof of Proposition* (ii) *in* [4]*, we can reconstruct the limit of* $f([\overline{W_1^n W_2^n}])$ *from the limits of* $[\overline{W_1^n W_2^n}]$ *and* $[\overline{W_1^n W_2^n}]'$ *in the sense of Euclidean metric.*

(iii) *For the case when* $k = 3, a_n^{(1)} = a_n^{(2)} = a_n^{(3)} = n$ *and* $V_0 = V_1 = V_2 = V_3 = \emptyset$*, our main theorem says that*

$$\lim_{n\to\infty} f([\overline{W_1^n W_2^n W_3^n}]) = \frac{\widetilde{f}(w_1) + \widetilde{f}(w_2) + \widetilde{f}(w_3)}{\widetilde{1}(w_1) + \widetilde{1}(w_2) + \widetilde{1}(w_3)}.$$

Here a sequence $[\overline{W_1^n W_2^n W_3^n}]$ *converges to* w_1 *and its non-trivial Galois conjugate converges to* w_3' *in the sense of Euclidean metric by Lemma* 1.28 *in* [1]*. Thus we cannot reconstruct* w_2 *from* $[\overline{W_1^n W_2^n W_3^n}]$ *by Euclidean metric and the function* f *on the set of real quadratic numbers does not satisfy the continuity with respect to Euclidean metric.*

Remark 1.3. *We set the sign of* η_w *in the definition because of the fact* $\widetilde{1}(w) = 2\log\varepsilon_w$*. We prove it in section* 2*, although we do not need it to prove our main theorem.*

2. Preliminaries

In this section, we prepare several notation.

Definition 2.1. *For non-empty even words* $V = (k_1, \dots, k_r), W = (\ell_1, \dots, \ell_s)$*, define a real quadratic number*

$$[V\overline{W}] := [k_1, \dots, k_r, \overline{\ell_1, \dots, \ell_s}].$$

Definition 2.2. *For a non-empty even word* $W = (k_1, \dots, k_r)$*, define*

$$\gamma_W := \begin{pmatrix} k_1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} k_2 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} k_r & 1 \\ 1 & 0 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}).$$

For the empty word $\emptyset$*, define*

$$\gamma_\emptyset := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Lemma 2.3. *We have the following properties.*

(i) *For even words* $W_1, \dots, W_n$*, we have* $\gamma_{W_1 \cdots W_n} = \gamma_{W_1} \cdots \gamma_{W_n}$*.*

(ii) *For even words* $V = (k_1, \dots, k_r), W = (\ell_1, \dots, \ell_s)$ *and a real quadratic number* $w = [\overline{W}]$*, we have* $\gamma_V w = [V\overline{W}]$*.*

In particular, $\gamma_W w = w$*.*

(iii) *For a reduced real quadratic number* w*, there exists the minimal even word* W *such that* $w = [\overline{W}]$*. Then we have* $N(W) = 1$*. In this case,* $\gamma_W = \gamma_w$*.*

Generally, for a non-empty even word W *and a real quadratic number* $w = [\overline{W}]$*, we have* $\gamma_W = \gamma_w^{N(W)}$*.*

(iv) *For a real quadratic number* w*, there exist even words* V, W *such that* $w = [V\overline{W}]$*. Then we have* $\gamma_V \gamma_W \gamma_V^{-1} = \gamma_w^{N(W)}$*.*

Proof. (i) and (ii) follows by definition. For (iii), it follows that $N(W) = 1$ by the minimality of W. We have $\gamma_W \in \mathrm{SL}_2(\mathbb{Z})_w$ by (ii) and $j(\gamma_W, w) > 1$ since $w > 1$ and all entries of γ_W are positive. Thus $\gamma_W = \gamma_w$ by the minimality of W.

For (iv), since any real quadratic number has a periodic continued fraction expansion, one can find such V, W in the claim. Let $w_0 := [\overline{W}]$. We have $\gamma_V^{-1} w = w_0$ by (i) and thus $\mathrm{SL}_2(\mathbb{Z})_w = \gamma_V^{-1} \mathrm{SL}_2(\mathbb{Z})_{w_0} \gamma_V$. Here

γ_w, γ_{w_0} are the unique generators of $\mathrm{SL}_2(\mathbb{Z})_w, \mathrm{SL}_2(\mathbb{Z})_{w_0}$ such that $j(\gamma_w, w) > 1, j(\gamma_{w_0}, w_0) > 1$ respectively. Since

$$\begin{aligned} &j(\gamma_V^{-1}\gamma_{w_0}\gamma_V, w) = j(\gamma_V^{-1}\gamma_{w_0}\gamma_V, \gamma_V^{-1}w_0) = j(\gamma_V^{-1}, \gamma_{w_0}\gamma_V\gamma_V^{-1}w_0)j(\gamma_{w_0}, \gamma_V\gamma_V^{-1}w_0)j(\gamma_V, \gamma_V^{-1}w_0) \\ &= j(\gamma_V^{-1}, w_0)j(\gamma_{w_0}, w_0)j(\gamma_V^{-1}, w_0)^{-1} = j(\gamma_{w_0}, w_0) > 1, \end{aligned}$$

we have $\gamma_V\gamma_{w_0}\gamma_V^{-1} = \gamma_w$. By (iii), we obtain $\gamma_V\gamma_W\gamma_V^{-1} = \gamma_w^{N(W)}$. □

Definition 2.4. *Set*

$$\infty := \lim_{t\to+\infty} t\sqrt{-1},\ \mathbb{P}^1(\mathbb{R}) := \mathbb{R} \cup \{\infty\}.$$

For $x', x \in \mathbb{P}^1(\mathbb{R})$, define

$$\eta_{x',x}(z) := \left(\frac{1}{z-x'} - \frac{1}{z-x}\right)dz.$$

Here, let $1/(z-\infty) := 0$ if $x = \infty$ or $x' = \infty$.

The following lemma follows from a direct calculation and there details are omitted.

Lemma 2.5. *For any $x, x' \in \mathbb{P}^1(\mathbb{R})$ and $\gamma \in \mathrm{SL}_2(\mathbb{R})$, we have $\gamma^*\eta_{x',x} = \eta_{\gamma^{-1}x',\gamma^{-1}x}$. In particular, we have $\gamma^*\eta_w = \eta_{\gamma^{-1}w}$ for a real quadratic number w and $\gamma \in \mathrm{SL}_2(\mathbb{Z})$.*

Definition 2.6. *For a non-empty even word W with $w = [\overline{W}]$ and $z_0 \in \mathbb{H}$, define*

$$\eta_W := \eta_w,\ \widetilde{f}(W) := \int_{z_0}^{\gamma_W z_0} f\eta_W.$$

Lemma 2.7. *For a non-empty even word W with $w = [\overline{W}]$, we have*

$$\widetilde{f}(W) = N(W)\widetilde{f}(w).$$

Thus $\widetilde{f}(W)$ is independent of z_0 and we have

$$f(w) = \frac{\widetilde{f}(W)}{\widetilde{1}(W)}.$$

Proof. it follows that $\gamma_W = \gamma_w^{N(W)}$ by Lemma 2.3 (iii), $\gamma_w^*\eta_w = \eta_w$ by Lemma 2.5 and $f \circ \gamma_w = f$. Thus we have $\widetilde{f}(W) = N(W)\widetilde{f}(w)$. □

For a real quadratic number w, there exist integers a, b, c such that

$$aw^2 + bw + c = 0,\ \gcd(a,b,c) = 1,\ w = \frac{-b + \sqrt{b^2-4ac}}{2a}.$$

Let $D := b^2 - 4ac > 0$ and $K := \mathbb{Q}(\sqrt{D})$. We have a group isomorphism

$$\begin{array}{ccc} \mathrm{GL}_2(\mathbb{Z})_w & \xrightarrow{\sim} & \mathbb{Z}[aw]^\times \\ \begin{pmatrix} p & q \\ r & s \end{pmatrix} & \longmapsto & rw + s \\ \begin{pmatrix} \frac{x-by}{2} & -cy \\ ay & \frac{x+by}{2} \end{pmatrix} & \longleftarrow\!\shortmid & \frac{x+y\sqrt{D}}{2} \end{array}$$

such that the diagram

$$\begin{array}{ccc} \mathrm{GL}_2(\mathbb{Z})_w & \xrightarrow{\quad\sim\quad} & \mathbb{Z}[aw]^\times \\ & {\scriptstyle\det}\searrow \quad \circlearrowright \quad \swarrow{\scriptstyle N_{K/\mathbb{Q}}} & \\ & \{\pm 1\} & \end{array}$$

commutes. Here the symbol $N_{K/\mathbb{Q}}$ is the norm map $N_{K/\mathbb{Q}}\colon K^\times \to \mathbb{Q}^\times$. Thus we have a group isomorphism

$$\mathrm{SL}_2(\mathbb{Z})_w \cong \{\varepsilon \in \mathbb{Z}[aw]^\times \mid N_{K/\mathbb{Q}}(\varepsilon) = 1\}.$$

The following is a fundamental property of S_w.

Proposition 2.8. *For a real quadratic number w, we have $\widetilde{1}(w) = 2\log\varepsilon_w$.*

Proof. Let

$$\varepsilon_w = (x_0+y_0)/2,\ \sigma := \begin{pmatrix} w' & w/(w'-w) \\ 1 & 1/(w'-w) \end{pmatrix} \in \mathrm{SL}_2(\mathbb{R}).$$

Then

$$\gamma_w = \begin{pmatrix} \frac{x_0-by_0}{2} & -cy_0 \\ ay_0 & \frac{x_0+by_0}{2} \end{pmatrix},\ \sigma^{-1}\gamma_w\sigma = \begin{pmatrix} \varepsilon_w^{-1} & 0 \\ 0 & \varepsilon_w \end{pmatrix}.$$

Fix $z_0 \in \mathbb{H}$ and let $z_1 := \sigma^{-1}z_0$. Since $\sigma^*\eta_w = \eta_{\infty,0} = -z^{-1}dz$ by Lemma 2.5, we have

$$\widetilde{1}(w) = \int_{z_0}^{\gamma_w z_0} \eta_w = \int_{\sigma z_1}^{\sigma\sigma^{-1}\gamma_w\sigma z_1} \eta_w = \int_{z_1}^{\sigma^{-1}\gamma_w\sigma z_1} \sigma^*\eta_w = -\int_{z_1}^{\varepsilon_w^{-2}z_1} \frac{dz}{z} = 2\log\varepsilon_w.$$

□

3. A proof of our main theorem

Throughout this section, we fix a point $z_0 \in \mathbb{H}$. We start with some lemmas.

Lemma 3.1. *For non-empty even words $W_1, \dots, W_k$, we have*

$$\widetilde{f}(W_1\cdots W_k) = \sum_{1\le i\le k} \int_{z_0}^{\gamma_{W_i}z_0} f\eta_{W_i\cdots W_kW_1\cdots W_{i-1}}.$$

Proof. We have

$$\widetilde{f}(W_1\cdots W_k) = \int_{z_0}^{\gamma_{W_1}\cdots\gamma_{W_k}z_0} f\eta_{W_1\cdots W_k}$$

by Lemma 2.3 (i). Thus we obtain

$$\begin{aligned}\widetilde{f}(W_1\cdots W_k) &= \sum_{1\le i\le k} \int_{\gamma_{W_1}\cdots\gamma_{W_{i-1}}z_0}^{\gamma_{W_1}\cdots\gamma_{W_i}z_0} f\eta_{W_1\cdots W_k} \\ &= \sum_{1\le i\le k} \int_{z_0}^{\gamma_{W_i}z_0} f(\gamma_{W_1}\cdots\gamma_{W_{i-1}})^*\eta_{W_1\cdots W_k}.\end{aligned}$$

We have

$$\widetilde{f}(W_1\cdots W_k) = \sum_{1\le i\le k} \int_{z_0}^{\gamma_{W_i}z_0} f\eta_{W_i\cdots W_kW_1\cdots W_{i-1}}.$$

by Lemma 2.5 and the fact that

$$(\gamma_{W_1}\cdots\gamma_{W_{i-1}})^{-1}[\overline{W_1\cdots W_k}] = [\overline{W_i\cdots W_kW_1\cdots W_{i-1}}],$$

□

Lemma 3.2. *Let K be a compact subset in $\mathbb{H}$ and $y_0 := \min\{\mathrm{Im}(z) \mid z \in K\}$. For real quadratic numbers v and w, suppose the continued fraction expansions of v and w share first n terms. Then we have*

$$\left|\frac{1}{z-v} - \frac{1}{z-w}\right| \le \frac{2^{2-n}}{y_0^2}$$

for any $z \in K$.

Proof. Fix $z \in K$. Since $|z-a| \ge \mathrm{Im}(z) \ge y_0$ for any $a \in \mathbb{R}$, we have

$$\left|\frac{1}{z-v} - \frac{1}{z-w}\right| \le \frac{|v-w|}{|z-v||z-w|} \le \frac{|v-w|}{y_0^2}.$$

Also we get $|v-w| \le 2^{2-n}$ by Lemma 1.24 in [1]. □

Lemma 3.3. *Let W and $W_1, \dots, W_n, \dots$ be non-empty even words such that W_n and W_{n+1} coincide in the first n terms and the last n terms. Let $\{a_n\}_{n=1}^{\infty}$ be a sequence in $\mathbb{Z}_{\geq 0}$ diverging to positive infinity. Set $w := [\overline{W}]$. Then*

$$\left\{ \int_{z_0}^{\gamma_W z_0} f\eta_{W_n} \right\}_{n\geq 1}, \ \left\{ \int_{z_0}^{\gamma_W^{a_n} z_0} f\eta_{W^{a_n}W_n} - a_n N(W)\widetilde{f}(w) \right\}_{n\geq 1}$$

are Cauchy sequences. In particular, these sequences are bounded.

Proof. Let

$$A_n := \int_{z_0}^{\gamma_W z_0} f\eta_{W_n}, \ B_n := \int_{z_0}^{\gamma_W^{a_n} z_0} f\eta_{W^{a_n}W_n} - a_n N(W)\widetilde{f}(w).$$

Fix positive integers $m \leq n$ such that $a_m \leq a_n$. By Lemma 2.7 and Lemma 3.1, we have

$$B_n - B_m = \int_{z_0}^{\gamma_W z_0} f \left(\sum_{1\leq i\leq a_n} \eta_{W^i W_n W^{a_n - i}} - \sum_{1\leq i\leq a_m} \eta_{W^i W_m W^{a_m - i}} - (a_n - a_m)\eta_W \right).$$

This is equal to

$$-\int_{z_0}^{\gamma_W z_0} f\left(S_1(m,n,z) + S_2(m,n,z) - S_1'(m,n,z) - S_2'(m,n,z)\right) dz$$

where

$$\begin{aligned}
S_1(m,n,z) &:= \sum_{1\leq i\leq a_m} \left(\frac{1}{z - [\overline{W^i W_n W^{a_n - i}}]} - \frac{1}{z - [\overline{W^i W_m W^{a_m - i}}]} \right), \\
S_1'(m,n,z) &:= \sum_{1\leq i\leq a_m} \left(\frac{1}{z - [\overline{W^{a_n - i} W_n W^i}]'} - \frac{1}{z - [\overline{W^{a_m - i} W_m W^i}]'} \right), \\
S_2(m,n,z) &:= \sum_{a_m < i\leq a_n} \left(\frac{1}{z - [\overline{W^i W_n W^{a_n - i}}]} - \frac{1}{z - w} \right), \\
S_2'(m,n,z) &:= \sum_{a_m < i\leq a_n} \left(\frac{1}{z - [\overline{W^{a_n - i} W_n W^i}]'} - \frac{1}{z - w'} \right).
\end{aligned}$$

We evaluate these four sums. Let $y_0 := \min\{\mathrm{Im}(z_0), \mathrm{Im}(\gamma_W z_0)\}$.

The continued fraction expansions of $[\overline{W^i W_n W^{a_n - i}}]$ and $[\overline{W^i W_m W^{a_m - i}}]$ share at least first $i + m$ terms since W_m and W_n share first m terms by assumption. Thus we have

$$|S_1(m,n,z)| \leq \sum_{1\leq i\leq a_m} \frac{2^{2-i-m}}{y_0^2} = \frac{2^{2-m} - 2^{2-m-a_m}}{y_0^2}$$

by Lemma 3.2.

By Lemma 1.28 in [1], the continued fraction expansions of $-[\overline{W^{a_n - i} W_n W^i}]'$ and $-[\overline{W^{a_m - i} W_m W^i}]'$ share at least first $i + m$ terms since W_m and W_n share last m terms by the assumption. Thus we obtain

$$|S_1'(m,n,z)| \leq \frac{2^{2-m} - 2^{2-m-a_m}}{y_0^2}$$

by the same argument in Lemma 3.2.

Also we have

$$|S_2(m,n,z)| \leq \frac{2^{2-a_n} - 2^{2-a_m}}{y_0^2}, \ |S_2'(m,n,z)| \leq \frac{2^{2-a_n} - 2^{2-a_m}}{y_0^2}.$$

Let C be the maximal value of f in the contour for the integration. As seen in the discussion above, we obtain

$$|B_n - B_m| \leq 2C|\gamma_W z_0 - z_0| \frac{2^{2-m} - 2^{2-m-a_m} + 2^{2-a_n} - 2^{2-a_m}}{y_0^2}.$$

Since the right hand side converges to 0 as $m, n \to \infty$, $\{B_n\}_{n\geq 1}$ is a Cauchy sequence.

We can prove similarly that $\{A_n\}_{n\geq 1}$ is a Cauchy sequence. □

Proof of Theorem 1.1. Keep the notation in Theorem 1.1 and let

$$W^{(n)} := V_0 W_1^{a_n^{(1)}} V_1 W_2^{a_n^{(2)}} V_2 \cdots W_k^{a_n^{(k)}} V_k.$$

Then we have $w^{(n)} = [\overline{W^{(n)}}]$. It follows from Lemma 3.1 that

$$\begin{aligned} &N(W^{(n)})\widetilde{f}(w^{(n)}) \\ &= \sum_{0\leq i\leq k} \int_{z_0}^{\gamma_{V_i} z_0} f\eta_{V_i W_{i+1}^{a_n^{(i+1)}} \cdots V_k V_0 W_1^{a_n^{(1)}} V_1 \cdots W_i^{a_n^{(i)}}} + \sum_{1\leq i\leq k} \int_{z_0}^{\gamma_{W_i}^{a_n^{(i)}} z_0} f\eta_{W_i^{a_n^{(i)}} V_i \cdots V_k V_0 W_1^{a_n^{(1)}} V_1 \cdots V_{i-1}}. \end{aligned}$$

By Lemma 3.3, this can be written as

$$\sum_{1\leq i\leq k} a_n^{(i)} N(W_i)\widetilde{f}(w_i) + O(1)$$

as $n \to \infty$. Similarly we have

$$N(W^{(n)})\widetilde{1}(w^{(n)}) = \sum_{1\leq i\leq k} a_n^{(i)} N(W_i)\widetilde{1}(w_i) + O(1)$$

by substituting $f = 1$.

Putting everything together we have

$$\begin{aligned} \lim_{n\to\infty} f(w^{(n)}) &= \lim_{n\to\infty} \frac{N(W^{(n)})\widetilde{f}(w^{(n)})}{N(W^{(n)})\widetilde{1}(w^{(n)})} \\ &= \lim_{n\to\infty} \frac{a_n^{(1)} N(W_1)\widetilde{f}(w_1) + \cdots + a_n^{(k)} N(W_k)\widetilde{f}(w_k) + O(1)}{a_n^{(1)} N(W_1)\widetilde{1}(w_1) + \cdots + a_n^{(k)} N(W_k)\widetilde{1}(w_k) + O(1)} \\ &= \frac{a^{(1)} N(W_1)\widetilde{f}(w_1) + \cdots + a^{(k)} N(W_k)\widetilde{f}(w_k)}{a^{(1)} N(W_1)\widetilde{1}(w_1) + \cdots + a^{(k)} N(W_k)\widetilde{1}(w_k)}. \end{aligned}$$

□

Acknowledgements

I would like to show my greatest appreciation to Professor Takuya Yamauchi for introducing me to this topic and giving many advices. I am deeply grateful to Dr. Toshiki Matsusaka for many discussion and valuable comments.

References

[1] M. Aigner. *Aigner, Markov's theorem and 100 years of the uniqueness conjecture*. Springer, 1997.
[2] P. Bengoechea and Ö. Imamoglu. Cycle integrals of modular functions, Markov geodesics and a conjecture of Kaneko. *Algebra Number Theory*, 13(4):943–962, 2019.
[3] W. Duke, Ö. Imamoḡlu, and Á. Tóth. Cycle integrals of the j-function and mock modular forms. *Ann. of Math. (2)*, 173(2):947–981, 2011.
[4] M. Kaneko. Observations on the 'values' of the elliptic modular function $j(\tau)$ at real quadratics. *Kyushu Journal of Mathematics*, 63(2):353–364, 2009.
[5] R. Masri. The asymptotic distribution of traces of cycle integrals of the j-function. *Duke Math. J.*, 161(10):1971–2000, 2012.
[6] S. Päpcke. Values of the j-function and its relation to Markov numbers. Master's thesis, The Department of Mathematics at Eidgenössische Technische Hochschule Zürich, 2016.
[7] D. Zagier. Traces of singular moduli. In *Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998)*, volume 3 of *Int. Press Lect. Ser.*, pages 211–244. Int. Press, Somerville, MA, 2002.