

A COTLAR TYPE MAXIMAL FUNCTION ASSOCIATED WITH FOURIER MULTIPLIERS

RAJULA SRIVASTAVA

ABSTRACT. We prove the L^p boundedness of a maximal operator associated with a dyadic frequency decomposition of a Fourier multiplier, under a weak regularity assumption.

1. INTRODUCTION

Consider a Mikhlin-Hörmander multiplier m on \mathbb{R}^d satisfying the assumption

$$(1) \quad |\partial^\gamma m(\xi)| \leq A|\xi|^{-|\gamma|}$$

for all multi-indices γ with $|\gamma| \leq L$ for some integer $L > d$.

Let $\chi \in C_c^\infty(\mathbb{R})$ be supported in $(1/2, 2)$ such that $\sum_{j=-\infty}^{\infty} \chi(2^j t) = 1$ and let $\phi = \chi(|\cdot|)$. For a given Schwartz function f , let $Sf := \mathcal{F}^{-1}[m\hat{f}]$, and for $n \in \mathbb{Z}$ let S_n be defined by

$$(2) \quad \widehat{S_n f}(\xi) := \sum_{j \leq n} \phi(2^{-j}\xi) m(\xi) \hat{f}(\xi).$$

We are interested in bounds for the maximal function

$$(3) \quad S_* f(x) := \sup_{n \in \mathbb{Z}} |S_n f(x)|.$$

The above operator was studied by Guo, Roos, Seeger and Yung in [4], in connection with proving L^p bounds for a maximal operator associated with families of Hilbert transforms along parabolas. The multiplier m in [4] was assumed to satisfy the condition

$$(4) \quad \sup_{t>0} \|\phi m(t \cdot)\|_{\mathcal{L}_\beta^1} = B(m) < \infty$$

with $\beta > d$. Here \mathcal{L}_β^1 is the potential space of functions g with $(I - \Delta)^{\beta/2} g \in L^1$ (we note the analogy with condition (1) here). With the above hypothesis, the authors were able to prove a pointwise Cotlar-type inequality

$$(5) \quad S_* f(x) \leq \frac{1}{(1 - \delta)^{1/r}} (M(|Sf|^r)(x))^{1/r} + C_{d,\beta} \delta^{-1} B(m) Mf(x)$$

for $f \in L^p(\mathbb{R}^d)$ and for almost every x (with $r > 0$ and $0 < \delta \leq 1/2$). Here $M[f]$ denotes the standard Hardy-Littlewood Maximal function. (5) easily implies L^p and weak (1,1) bounds for the maximal operator S_* .

It is natural to ask if one could weaken the assumption (4) and still establish L^p bounds on the operator S_* , possibly without the intermediate step of proving a pointwise inequality of the form (5) for all exponents r . In this paper, we answer this question in the affirmative.

Date: 2022/01/21.

2010 *Mathematics Subject Classification.* 42B15, 42B20, 42B25.

In particular, we show that for the L^p bounds on S_* to hold, it is enough for the multiplier m to satisfy a much weaker condition

$$(6) \quad \sup_{t>0} \int |\mathcal{F}^{-1}[\phi m(t \cdot)](x)| (\log(2 + |x|))^\alpha dx := B(m) < \infty, \quad \alpha > 3.$$

Our main result is the following:

Theorem 1.1. *S_* , as defined in (3) for a multiplier m satisfying (6), is of weak-type $(1,1)$ and bounded on L^p for $p \in (1, \infty)$, with the respective operator norm $\lesssim_p B(m)$.*

We remark here that it is not possible to do away with the smoothness assumption entirely. In other words, the condition

$$\sup_{t>0} \int |\mathcal{F}^{-1}[\phi m(t \cdot)](x)| dx < \infty$$

alone is not enough to guarantee L^p bounds on the maximal operator S_* , or even on the singular operator S . For counterexamples, we refer to [10], Section 5 and [9], Section 3.

We shall denote the Littlewood-Paley pieces of m by m_j . More precisely, for $j \in \mathbb{Z}$, we define $m_j(\xi) := \phi(2^{-j}\xi)m(\xi)$. Furthermore let $a_j(\xi) = m_j(2^j\xi) = \eta(\xi)m(2^j\xi)$. Observe that $\text{supp}(m_j) \subset (2^{j-1}, 2^j)$ and $\text{supp}(a_j) \subset (1/2, 2)$. Let $K_j = \mathcal{F}^{-1}[m_j]$. Then (2) can be re-written as

$$(7) \quad S_n f(x) = \sum_{j \leq n} K_j * f(x).$$

However, in order to quantify the smoothness condition in (6), it is useful to partition K_j (for each $j \in \mathbb{Z}$) on the space side as well (see [2]). To this effect, let $\eta_0 \in C_c^\infty(\mathbb{R}^d)$ be such that η_0 is even, $\eta_0(x) = 1$ for $|x| \leq 1/2$ and η_0 is supported where $|x| \leq 1$. For $l \in \mathbb{N}$, let $\eta_l(x) = \eta_0(2^{-l}x) - \eta_0(2^{-l+1}x)$. For $l \in \mathbb{N} \cup \{0\}$ we define

$$(8) \quad K^{l,j}(x) = \eta_l(2^j x) \mathcal{F}^{-1}[m_j](x).$$

By the assumption (6), we then have

$$(9) \quad \|K^{l,j}\|_{L^1} \lesssim B(\log(2 + 2^l))^{-\alpha}.$$

Now the multiplier corresponding to $K^{l,j}$ is given by $2^{-jd} \hat{\eta}_l(2^{-j} \cdot) * m_j$, which, unlike m_j , is not compactly supported. However the rapid decay of $\hat{\eta}_l$ still leads to the multiplier "essentially" being supported in a slightly thicker (but still compact) dyadic annulus, with the other frequency regions contributing negligible error terms. We make these ideas rigorous in Section 2. The arguments used are similar in spirit to those in [1], Section 5. Another source of reference is [6]. As a corollary, we prove that the singular operator

$$(10) \quad S^l f(x) = \sum_{j \in \mathbb{Z}} K^{l,j} * f(x)$$

is bounded on L^2 , with operator norm $\lesssim B l^{-\alpha}$.

In Section 3, we use Bernstein's inequality (see [11]) to establish L^p bounds for the aforementioned portion with the major contribution and with compact frequency support. We also establish a pointwise estimate on its gradient.

In Section 4, using the estimates from Section 3, we prove that the Calderón-Zygmund operator S^l associated to the multiplier m is of weak type $(1,1)$, with the operator norm $\lesssim B l^{-\alpha+1}$. We do so by establishing the result for the operator $T^l f := \sum_{j \in \mathbb{Z}} H^{l,j} * f$, where $H^{l,j}$ is the portion of the kernel $K^{l,j}$ with the major contribution. The arguments flow

in the same vein as those in the proof of the Mikhlin-Hörmander Multiplier Theorem (see [3],[5]). We also establish L^p bounds on T^l using interpolation.

In Section 5, we investigate the properties of the truncated operator $T_n^l f := \sum_{j \leq n} H^{l,j} * f$ (for $n \in \mathbb{Z}$ and $l \geq 0$), with an aim of establishing L^p bounds for the associated maximal operator $T_*^l f = \sup |T_n^l f|$.

We wish to show that for $l \geq 0$, the operator T_*^l is bounded on L^p for $1 < p < \infty$ with the corresponding operator norm $\lesssim_p B l^{-\alpha+1}$ ($\lesssim B$ for $l = 0$). Since our assumption (6) on m is much weaker than that in [4], a pointwise inequality like (5) for all $r > 0$ seems out of reach. However, by using similar ideas, we are able to establish a Cotlar type inequality

$$T_*^l f(x) \lesssim_d \frac{1}{(1-\delta)^{1/p}} (M(|T^l f|^p)(x))^{1/p} + 2^{\frac{ld}{p}} l^{-\alpha+1} (1+\delta^{-1/p}) B(m) (M(|f|^p)(x))^{1/p}.$$

for each T_*^l and for a large enough exponent p (here $0 < \delta \leq 1/2$). Roughly speaking, a choice of $p = p_l \sim l$ will work. In particular, $p_l \rightarrow \infty$ as $l \rightarrow \infty$. Using this inequality, we can conclude that T_*^l is bounded (with norm $\lesssim_p B l^{-\alpha+1}$) for all $p \in [p_l, \infty)$. This idea of keeping track of the explicit dependence on the exponent l and using the decay in l to sum the pieces up has also been used in [2], albeit for a different maximal operator than the one considered here. For $p \in (1, p_l)$, however, we rely on a weak (1,1) estimate for T_*^l (which is not hard to obtain) and then an interpolation between 1 and p_l , which causes us to gain a power of l . In other words, we are only able to retain a decay of $l^{-\alpha+2}$ (hence the assumption $\alpha > 3$). This is a trade off of working with a weaker logarithmic regularity assumption.

Finally, in Section 6, we establish a weak (1,1) estimate for the maximal operator T_*^l and obtain L^p bounds for the sum $\sum_{l \geq 0} T_*^l$, and consequently for S_* (we use the decay in l and the condition $\alpha > 3$ here).

Acknowledgements. The author would like to thank her advisor Andreas Seeger for introducing this problem, for his guidance and several illuminating discussions. Research supported in part by NSF grant 1500162.

2. THE ERROR TERMS

Let $\Psi \in C^\infty$ with $\Psi = 1$ on $\{1/4 \leq |\xi| \leq 4\}$ and supported on $\{1/5 \leq |\xi| \leq 5\}$. For $j \in \mathbb{Z}$, define $\Psi_j(\xi) = \Psi(2^{-j}\xi)$. Then

$$(11) \quad \sum_{j \in \mathbb{Z}} K^{l,j} = \sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(\Psi_j) + \sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(1 - \Psi_j)$$

where $K^{l,j}$ is as defined in (8).

In this section, we will show that the contribution from the second sum above can be made as small as required using the rapid decay of $\hat{\eta}$. To control this sum, we study the corresponding multiplier given by

$$\sum_{j \in \mathbb{Z}} 2^{-jd} (1 - \Psi_j)(\xi) \hat{\eta}_l(2^{-j}\cdot) * m_j(\xi) = \sum_{j \in \mathbb{Z}} (1 - \Psi_j(\xi)) 2^{-jd} \int m_j(\omega) \hat{\eta}_l(2^{-j}(\xi - \omega)) d\omega.$$

Lemma 2.1. *Let $l \geq 0$. For any $N \in \mathbb{N}$, multi-index γ and $\xi \neq 0$, we have the estimate*

$$(12) \quad \sum_{j \in \mathbb{Z}} 2^{-jd} |(1 - \Psi_j) \partial_\xi^\gamma (\hat{\eta}_l(2^{-j}\cdot) * m_j)(\xi)| \lesssim_{\eta, N, \gamma, d} B 2^{l(d+|\gamma|-N)} |\xi|^{-|\gamma|}.$$

Proof. Fix $\xi \neq 0$. Let $k \in \mathbb{Z}$ be such that $2^{k-1} \leq |\xi| < 2^{k+1}$. As $\Psi_k = 1$ on $\{2^{k-2} \leq |\xi| \leq 2^{k+2}\}$, we can split the sum under consideration into two parts

$$\begin{aligned} & \sum_{j \in \mathbb{Z}} 2^{-jd} |(1 - \Psi_j) \partial_\xi^\gamma (\hat{\eta}_l(2^{-j} \cdot) * m_j)(\xi)| \leq \\ & \sum_{j > k+2} 2^{-jd} |\partial_\xi^\gamma (\hat{\eta}_l(2^{-j} \cdot) * m_j)(\xi)| + \sum_{j < k-2} 2^{-jd} |\partial_\xi^\gamma (\hat{\eta}_l(2^{-j} \cdot) * m_j)(\xi)|. \end{aligned}$$

First Term: We have

$$\sum_{j > k+2} 2^{-jd} |\partial_\xi^\gamma (\hat{\eta}_l(2^{-j} \cdot) * m_j)(\xi)| \leq \sum_{j > k+2} 2^{-jd} \int_{2^{j-1} \leq |\omega| \leq 2^{j+1}} |m_j(\omega)| |\partial_\xi^\gamma \hat{\eta}_l(2^{-j}(\xi - \omega))| d\omega.$$

Now we observe that for $2^{j-1} \leq |\omega| \leq 2^{j+1}$, we have $|\omega - \xi| \geq |\omega|/2$ and $2^{-j}|\omega|/2 \sim 1$. Hence, the Schwartz decay of $\hat{\eta}$ yields

$$\begin{aligned} & \sum_{j > k+2} 2^{-jd} |\partial_\xi^\gamma (\hat{\eta}_l(2^{-j} \cdot) * m_j)(\xi)| \\ & \lesssim_{\eta, N, \gamma} \sum_{j > k+2} 2^{-jd} \int_{2^{j-1} \leq |\omega| \leq 2^{j+1}} \|m\|_{L^\infty} 2^{l(d+|\gamma|)} 2^{-j|\gamma|} \left(2^{l-j}|\omega|/2\right)^{-N} d\omega \\ & \lesssim_{d, \eta, N, \gamma} \sum_{j > k+2} 2^{-jd} \|m\|_{L^\infty} 2^{l(d+|\gamma|-N)} 2^{-j|\gamma|} \int_{2^{j-1} \leq |\omega| \leq 2^{j+1}} d\omega \\ & \lesssim_{d, \eta, N, \gamma} \sum_{j > k+2} 2^d \|m\|_{L^\infty} 2^{l(d+|\gamma|-N)} 2^{-j|\gamma|} \lesssim_{d, \eta, N, \gamma} B 2^{l(d+|\gamma|-N)} 2^{-(k+2)|\gamma|} \lesssim_{d, \eta, N, \gamma} B 2^{l(d+|\gamma|-N)} |\xi|^{-|\gamma|}. \end{aligned}$$

Second Term: In this case for $2^{j-1} \leq |\omega| \leq 2^{j+1}$ we have that $|\omega - \xi| \geq |\xi|/2$ and $2^{-j}|\xi|/2 \geq 1$. Hence

$$\begin{aligned} & \sum_{j < k-2} 2^{-jd} |\partial_\xi^\gamma (\hat{\eta}_l(2^{-j} \cdot) * m_j)(\xi)| \\ & \lesssim_{\eta, N, \gamma} \sum_{j < k-2} 2^{-jd} \int_{2^{j-1} \leq |\omega| \leq 2^{j+1}} \|m\|_{L^\infty} 2^{l(d+|\gamma|)} 2^{-j|\gamma|} \left(2^{l-j}|\xi|/2\right)^{-N} d\omega \\ & \lesssim_{\eta, N, \gamma} \sum_{j < k-2} 2^{-jd} \|m\|_{L^\infty} 2^{l(d+|\gamma|-N)} \int_{2^{j-1} \leq |\omega| \leq 2^{j+1}} |\xi|^{-|\gamma|} \left(2^{-j}|\xi|/2\right)^{-N+|\gamma|} d\omega \\ & \lesssim_{\eta, N, \gamma, d} \|m\|_{L^\infty} 2^{l(d+|\gamma|-N)} |\xi|^{-|\gamma|} \sum_{j < k-2} 2^{-jd} 2^{(k-1)d} \lesssim_{\eta, N, \gamma, d} \|m\|_{L^\infty} 2^{l(d+|\gamma|-N)} |\xi|^{-|\gamma|}. \end{aligned}$$

□

Now a simple application of the Mikhlin-Hörmander Multiplier theorem gives us

Theorem 2.2. *Let $l \geq 0$. For any $N_0 \in \mathbb{N}$, $p \in (1, \infty)$ and Schwartz function f , we have*

$$\left\| \sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(1 - \Psi_j) * f \right\|_{L^p} \lesssim_d \max(p, (p-1)^{-1}) C_{\eta, N_0, p, \Psi} B 2^{-\log_0} \|f\|_{L^p}.$$

Furthermore, we also have

$$\left\| \sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(1 - \Psi_j) * f \right\|_{L^{1,\infty}} \lesssim_d C_{\eta, N_0, p, \Psi} B 2^{-\log_0} \|f\|_{L^1}.$$

Proof. We need to prove that

$$|\sum_{\beta \leq \gamma} c_{\beta, \gamma} \sum_{j \in \mathbb{Z}} \partial^\beta (1 - \Psi_j)(\xi) \partial^{\gamma - \beta} (2^{-jd} \hat{\eta}_l(2^{-j} \cdot) * m_j)(\xi)| \lesssim_d C_{\eta, N, p, \Psi} B 2^{-l \log_0} |\xi|^{-|\gamma|}$$

where γ is a multi-index with $|\gamma| \leq d/2 + 1$ and $\xi \neq 0$. We observe that for all values of j except for j_1, j_2 where $2^{j_1+2} \leq |\xi| < 5 \cdot 2^{j_1}$ or when $2^{j_2}/5 \leq |\xi| < 2^{j_2-2}$, we can use lemma 2.1 directly (as then $|(1 - \Psi_j)(\xi)|$ is a constant). Further, for the two remaining cases, we observe that $|\xi| \sim 2^{j_k}$ ($k = 1, 2$). Hence we can bound the term $|\partial^\beta (1 - \Psi_{j_k})|$ by $C_\Psi |\xi|^{-|\beta|}$ and apply the previous lemma to the term $|\partial^{\gamma - \beta} (2^{-jd} \hat{\eta}_l(2^{-j} \cdot) * m_j)(\xi)|$ to bound it above by $C_{\eta, N} B 2^{l(d+|\gamma|-|\beta|-N_0-|\gamma|)} |\xi|^{-(|\gamma|-|\beta|)}$. The result then follows by summing up. \square

As a consequence, we obtain the L^2 boundedness of S^l (as defined in (10)), with polynomial decay in l .

Theorem 2.3. *For $f \in L^2$ and $l > 0$, we have*

$$\|S^l(f)\|_{L^2} \lesssim B l^{-\alpha} \|f\|_{L^2}.$$

We also have

$$\|S^0(f)\|_{L^2} \lesssim B \|f\|_{L^2}.$$

Proof. It is enough to prove the above for a Schwartz function f . Now

$$S^l(f) = \sum_{j \in \mathbb{Z}} K^{l,j} * f = \sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(\Psi_j) * f + \sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(1 - \Psi_j) * f.$$

As we have already established the L_2 boundedness of the second term in Theorem 2.2 (with as good a decay in l as required), we only need to prove the theorem for the first term. To this effect, let

$$f = \sum_{k \in \mathbb{Z}} \Delta_k f$$

be a Littlewood-Paley decomposition of f . Then

$$\begin{aligned} \|\sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(\Psi_j) * f\|_{L^2} &\lesssim \|\sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(\Psi_j) * \Delta_k f\|_{L^2} \\ &\lesssim \|\sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(\Psi_j) * (\Delta_{j-1} + \Delta_j + \Delta_{j+1}) f\|_{L^2} \end{aligned}$$

where we have used the fact that the frequency support of $K^{l,j} * \mathcal{F}^{-1}(\Psi_j)$ is contained in $\{2^j/5 \leq |\xi| \leq 5 \cdot 2^j\}$. From (9), we also have that

$$\|K^{l,j} * \mathcal{F}^{-1}(\Psi_j)\|_{L^1} \leq \|K^{l,j}\|_{L^1} \|\mathcal{F}^{-1}(\Psi_j)\|_{L^1} \lesssim B (\log(2 + 2^l))^{-\alpha} \|\Psi\|_{L^\infty}.$$

Hence

$$\begin{aligned} \|\sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(\Psi_j) * f\|_{L^2} &\lesssim B (\log(2 + 2^l))^{-\alpha} \|\sum_{j \in \mathbb{Z}} (\Delta_{j-1} + \Delta_j + \Delta_{j+1}) f\|_{L^2} \\ &\lesssim B (\log(2 + 2^l))^{-\alpha} \|f\|_{L^2} \end{aligned}$$

which proves the theorem. \square

3. ESTIMATES FOR THE MAJORLY CONTRIBUTING PORTION OF THE KERNEL

We now turn our attention to the first term in (11), which is the one with the main contribution. The main advantage we have now is that the j th term in

$$\sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(\Psi_j)$$

has frequency supported in the annulus $\{2^j/5 \leq |\xi| \leq 5 \cdot 2^j\}$. Hence, we can use Bernstein's inequality to get bounds on the L_p norm (of each term, with $1 < p < \infty$) and L^∞ norm (of the derivative).

Proposition 3.1. *Let $q \in (1, \infty)$, and let q' denote the Hölder conjugate exponent of q . Then for all $l \in \mathbb{N} \cup \{0\}$ and $j \in \mathbb{Z}$, we have*

- (1) $\|K^{l,j} * \mathcal{F}^{-1}(\Psi_j)\|_{L^{q'}} \lesssim B 2^{\frac{jd}{q}} (\log(2 + 2^l))^{-\alpha}$.
- (2) For $x \in \mathbb{R}^d$, $|\nabla(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))(x)| \lesssim 2^j |K^{l,j} * \mathcal{F}^{-1}(\Psi_j)(x)|$.

Proof. For the first part, we have

$$\begin{aligned} & \|K^{l,j} * \mathcal{F}^{-1}(\Psi_j)\|_{L^{q'}} \\ & \leq \|K^{l,j}\|_{L^1} \|\mathcal{F}^{-1}(\Psi_j)\|_{L^{q'}} \lesssim B (\log(2 + 2^l))^{-\alpha} \|\Psi_j\|_{L^q} \lesssim_\Psi B (\log(2 + 2^l))^{-\alpha} 2^{jd/q} \end{aligned}$$

where in the second step we have used (9) and Hausdorff-Young's inequality.

For the second part, we recall that the Fourier transform of $\nabla(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))$ is supported on a dyadic annulus of radius 2^j . The assertion then follows from Bernstein's inequality (see [11], Proposition 5.3). \square

4. WEAK (1,1) BOUNDEDNESS OF S^l

Let $\mathbb{1}_l$ be the characteristic function of the set $\{x : 2^{l-1} \leq |x| \leq 2^{l+1}\}$ for $l > 0$ and of the set $\{x : |x| \leq 1\}$ for $l = 0$. We will denote $\mathbb{1}_l(2^j x)$ by $\mathbb{1}^{l,j}(x)$. Then (11) can be rewritten as

$$(13) \quad \sum_{j \in \mathbb{Z}} K^{l,j} = \sum_{j \in \mathbb{Z}} K^{l,j} \mathbb{1}^{l,j} = \sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(\Psi_j) \mathbb{1}^{l,j} + \sum_{j \in \mathbb{Z}} K^{l,j} * \mathcal{F}^{-1}(1 - \Psi_j) \mathbb{1}^{l,j}.$$

The advantage of (13) over (11) is that it preserves information about the compact support of the kernel $K^{l,j}$, a property we will exploit quite often in the forthcoming proofs. Now for $f \in L^1(\mathbb{R}^d)$, we can estimate

$$\left\| \sum_{j \in \mathbb{Z}} K^{l,j} * f \right\|_{L^{1,\infty}} \leq \left\| \sum_{j \in \mathbb{Z}} H^{l,j} * f \right\|_{L^{1,\infty}} + \left\| \sum_{j \in \mathbb{Z}} (K^{l,j} * \mathcal{F}^{-1}(1 - \Psi_j) \mathbb{1}^{l,j}) * f \right\|_{L^{1,\infty}}$$

where we define $H^{l,j} := (K^{l,j} * \mathcal{F}^{-1}(\Psi_j)) \mathbb{1}^{l,j}$. Also let the operator T^l be defined as $T^l f := \sum_{j \in \mathbb{Z}} H^{l,j} * f$.

By Theorem 2.2, we conclude that $\|\sum_{j \in \mathbb{Z}} (K^{l,j} * \mathcal{F}^{-1}(1 - \Psi_j) \mathbb{1}^{l,j}) * f\|_{L^{1,\infty}} \lesssim B l^{-\alpha+1} \|f\|_{L^1}$. Hence, in order to prove that S^l is of weak type $(1, 1)$ (with the respective norm $\lesssim_d B l^{-\alpha+1}$), it is enough to prove the same for T^l , which is the content of the next theorem. The proof we give here essentially uses the same ideas as Hörmander's original proof of the (Hörmander-)Mikhlin Multiplier Theorem (see [5], also [3]).

Theorem 4.1. *For all $f \in L^1(\mathbb{R}^d)$, we have*

$$\|T^l f\|_{L^{1,\infty}} \lesssim B l^{-\alpha+1} \|f\|_{L^1}$$

for $l \geq 1$ and

$$\|T^0 f\|_{L^{1,\infty}} \lesssim B \|f\|_{L^1}$$

Proof. We prove the result for $l > 0$. The result for $l = 0$ follows similarly. Also, for this proof, we can assume that $B = 1$. Let $f \in L^1(\mathbb{R}^d)$ and fix $\sigma > 0$. Let

$$f = f_0 + f_1$$

be the standard Calderón-Zygmund decomposition of f at the level $l^{\alpha-1}\sigma$. More precisely, let $\{I_k\}_{k \in \mathbb{N}}$ be axis-parallel cubes with centres $\{a_k\}_{k \in \mathbb{N}}$ respectively such that

$$l^{\alpha-1}\sigma < |I_k|^{-1} \int_{I_k} |f(y)| dy \leq 2^d l^{\alpha-1}\sigma,$$

$$|f(x)| \leq l^{\alpha-1}\sigma \text{ a.e. for } x \notin \bigcup_{k \in \mathbb{N}} I_k,$$

$$f_0(x) = \begin{cases} f(x) - |I_k|^{-1} \int_{I_k} f(y) dy & x \in I_k, \\ 0, & \text{otherwise.} \end{cases} \quad f_1(x) = \begin{cases} |I_k|^{-1} \int_{I_k} f(y) dy & x \in I_k, \\ f(x), & \text{otherwise.} \end{cases}$$

Now

$$\begin{aligned} & \text{meas}(\{x : |\sum_{j \in \mathbb{Z}} H^{l,j} * f(x)| > \sigma\}) \\ & \leq \text{meas}(\{x : |\sum_{j \in \mathbb{Z}} H^{l,j} * f_0(x)| > \sigma/2\}) + \text{meas}(\{x : |\sum_{j \in \mathbb{Z}} H^{l,j} * f_1(x)| > \sigma/2\}) \end{aligned}$$

We estimate $\text{meas}(\{x : |\sum_{j \in \mathbb{Z}} H^{l,j} * f_0(x)| > \sigma/2\})$

$$\begin{aligned} & \leq \text{meas}(\{x : |\sum_{j \in \mathbb{Z}} H^{l,j} * f_0(x)| > \sigma/2\} \cap (\bigcup_{k \in \mathbb{N}} 2I_k)^c) + \text{meas}(\bigcup_{k \in \mathbb{N}} 2I_k) \\ (a) \quad & \lesssim_d \text{meas}(\{x : |\sum_{j \in \mathbb{Z}} H^{l,j} * f_0(x)| > \sigma/2\} \cap (\bigcup_{k \in \mathbb{N}} 2I_k)^c) + \frac{l^{-\alpha+1} \|f\|_1}{\sigma}. \end{aligned}$$

Now since the mean value of f_0 over I_k vanishes, we have

$$\begin{aligned} & \int_{(\bigcup_{k \in \mathbb{N}} 2I_k)^c} |\sum_{j \in \mathbb{Z}} H^{l,j} * f_0(x)| dx \\ & \leq \sum_{k \in \mathbb{N}} \int_{I_k} \left(\int_{(\bigcup_{k \in \mathbb{N}} 2I_k)^c} |H^{l,j}(x-y) - H^{l,j}(x-a_k)| dx \right) |f_0(y)| dy \\ (b) \quad & \lesssim_d l^{-\alpha+1} \sum_{k \in \mathbb{N}} \int_{I_k} |f_0(y)| dy \lesssim_d l^{-\alpha+1} \|f\|_1, \end{aligned}$$

provided we prove that

$$(c) \quad \int_{(\bigcup_{k \in \mathbb{N}} 2I_k)^c} |\sum_{j \in \mathbb{Z}} H^{l,j}(x-y) - H^{l,j}(x-a_k)| dx \lesssim_d l^{-\alpha+1}, \quad y \in I_k.$$

We postpone the proof of (c) in order to conclude the estimates. By (a), (b) and (c), we obtain

(d)

$$\text{meas}(\{x: |\sum_{j \in \mathbb{Z}} H^{l,j} * f_0(x)| > \sigma/2\}) \lesssim_d \sigma^{-1} l^{-\alpha+1} \|f\|_1 + \sigma^{-1} l^{-\alpha+1} \|f\|_1 \lesssim_d \sigma^{-1} l^{-\alpha+1} \|f\|_1.$$

Set $p = 2l + 4$. Then by Theorem 2.3, we have $\|S^l f\|_2 \lesssim_d l^{\alpha-1} \|f\|_{L^2}$, and hence

$$\sigma^2 \text{meas}(\{x: |\sum_{j \in \mathbb{Z}} H^{l,j} * f_1(x)| > \sigma/2\}) \leq \|\sum_{j \in \mathbb{Z}} H^{l,j} * f_1\|_2^2 \lesssim l^{2(-\alpha+1)} \|f_1\|_2^2$$

$$\begin{aligned} &= l^{2(-\alpha+1)} \left(\sum_{k \in \mathbb{N}} |I_k|^{-1} \left| \int_{I_k} f(x) dx \right|^2 + \int_{(\bigcup_{k \in \mathbb{N}} I_k)^c} |f(x)|^2 dx \right) \\ &\lesssim_d l^{-\alpha+1} \sigma \left\{ \sum_{k \in \mathbb{N}} \left| \int_{I_k} f(x) dx \right| + \int_{(\bigcup_{k \in \mathbb{N}} I_k)^c} |f(x)| dx \right\} \end{aligned}$$

which enables us to conclude

$$(e) \quad \text{meas}(\{x: |\sum_{j \in \mathbb{Z}} H^{l,j} * f_1(x)| > \sigma/2\}) \lesssim \frac{l^{-\alpha+1} \|f\|_1}{\sigma}.$$

Combining the estimates (d) and (e) yields

$$\text{meas}(\{x: |\sum_{j \in \mathbb{Z}} H^{l,j} * f(x)| > \sigma\}) \lesssim_d \frac{l^{-\alpha+1} \|f\|_1}{\sigma}.$$

There remains the proof of (c). It is sufficient to prove that

$$\sum_{j \in \mathbb{Z}} \int_{|x| \geq 2t} |H^{l,j}(x-y) - H^{l,j}(x)| dx \lesssim l^{-\alpha+1} (|y| \leq t, t > 0).$$

Fix $t > 0$. Let $m \in \mathbb{Z}$ such that $|t| \sim 2^m$. Now for $j > -m$,

$$\int_{|x| \geq 2t} |H^{l,j}(x-y) - H^{l,j}(x)| dx \leq 2 \int_{|x| \geq 2t} |H^{l,j}(x)| dx \leq \int_{|x| \geq 2^m} |K^{l,j} * \mathcal{F}^{-1}(\Psi_j)(x)| \mathbb{1}^{l,j}(x) dx.$$

Now we observe that for the last term to be non-zero, $2^{l-j} \geq 2^m$. Hence we have

$$\begin{aligned} &\sum_{j > -m} \int_{|x| \geq 2t} |H^{l,j}(x-y) - H^{l,j}(x)| dx \\ &\leq \sum_{-m < j \leq l-m} \int_{|x| \geq 2^m} |K^{l,j} * \mathcal{F}^{-1}(\Psi_j)(x)| \mathbb{1}^{l,j}(x) dx \leq l \|K^{l,j}\|_{L^1} \|\mathcal{F}^{-1}(\Psi_j)\|_{L^1} \lesssim_{\Psi} l (\log(2 + 2^l))^{-\alpha} \\ &\lesssim l^{-\alpha+1} \end{aligned}$$

where we have used our assumption (6) to conclude the second to last inequality. Moreover, by Proposition 3.1, we have $|\nabla(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))(x)| \lesssim 2^j |(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))(x)|$ which yields

$$\begin{aligned}
\sum_{j \leq -m} \int_{|x| \geq t} |H^{l,j}(x-y) - H^{l,j}(x)| dx &\leq \sum_{j \leq -m} \int_0^1 \int_{\mathbb{R}^n} |\langle y, \nabla(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))(x - \tau y) \rangle| dx d\tau \\
&\lesssim \sum_{j \leq -m} t \|\nabla(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))\|_1 \\
&\lesssim \sum_{j \leq -m} t 2^j \|K^{l,j} * \mathcal{F}^{-1}(\Psi_j)\|_1 \lesssim_{\Psi} \sum_{j \leq -m} \|K^{l,j}\| t 2^j \leq l^{-\alpha}.
\end{aligned}$$

The last two estimates give (c), and the proof is complete. \square

The following theorem now follows almost immediately using standard L^p -interpolation theory.

Theorem 4.2. T^l defines a bounded operator on L^p for $p \in (1, \infty)$, with

$$\|T^l\|_{L^p} \lesssim_d \max(p, (p-1)^{-1} B l^{-\alpha+1})$$

for $l > 0$. We also have

$$\|T^0\|_{L^p} \lesssim_d \max(p, (p-1)^{-1} B).$$

Proof. The operator T^l is bounded on L^2 (by Theorem 2.3) and maps L^1 to $L^{(1,\infty)}$ (by Theorem 4.1), with norm $B l^{-\alpha+1}$ (norm B for $l = 0$). Interpolating between the L^1 and L^2 spaces then yields the required norm for the operator acting on L^p with $p \in (1, 2)$. Further, a duality argument yields the desired result for $p \in (2, \infty)$ as well. This concludes the proof of the theorem. \square

5. BOUNDEDNESS ON L^p FOR LARGE p

In this section, we investigate the properties of the truncated operator $T_n^l f := \sum_{j \leq n} H^{l,j} * f$ (for $n \in \mathbb{Z}$ and $l \geq 0$), with an aim of establishing $L^p \rightarrow L^p$ bounds for the associated maximal operator $T_*^l f = \sup |T_n^l f|$. Let $M[f]$ denote the standard Hardy-Littlewood Maximal function.

We wish to show that for $l \geq 0$, the operator T_*^l is bounded on L^p for $1 < p < \infty$ with the corresponding operator norm $\lesssim_p B l^{-\alpha+1}$ ($\lesssim_p B$ for $l = 0$). We will achieve this by proving a Cotlar type inequality

$$T_*^l f(x) \lesssim_d \frac{1}{(1-\delta)^{1/q}} (M(|T^l f|^q)(x))^{1/q} + 2^{\frac{ld}{q}} l^{-\alpha+1} (1 + \delta^{-1/q}) B(m) (M(|f|^q)(x))^{1/q}.$$

for each T_*^l and for a large enough exponent q (here $0 < \delta \leq 1/2$). Roughly speaking, a choice of $q = q_l \sim l$ will work. In particular, $q_l \rightarrow \infty$ as $l \rightarrow \infty$. Using this inequality, we can conclude that T_*^l is bounded (with norm $\lesssim_p B l^{-\alpha+1}$) for all $p \in (p_l, \infty)$.

The following lemma is the main step in establishing the Cotlar type inequality. The ideas used are similar to Lemma A.1 in [4].

Lemma 5.1. Fix $\tilde{x} \in \mathbb{R}^d$, $n \in \mathbb{Z}$ and $q > 1$. Let $g(y) = f(y) \mathbf{1}_{B(\tilde{x}, 2^{-n})}(y)$ and $h = f - g$. Then we have

$$\begin{aligned}
(i) \quad |T_n^l g(\tilde{x})| &\lesssim \begin{cases} B(M[f^q](\tilde{x}))^{1/q} & l = 0. \\ 0 & l > 0. \end{cases} \\
(ii) \quad |T_n^l h(\tilde{x}) - T^l h(\tilde{x})| &\lesssim \begin{cases} 0 & l = 0. \\ B 2^{ld/q} l^{-\alpha+1} (M[f^q](\tilde{x}))^{1/q} & l > 0. \end{cases}
\end{aligned}$$

(iii) For $|w - \tilde{x}| \leq 2^{-n-1}$, we have

$$|T^l h(\tilde{x}) - T^l h(w)| \lesssim \begin{cases} B(M[f^q](\tilde{x}))^{1/q} & l = 0. \\ B2^{ld/q} l^{-\alpha+1} (M[f^q](\tilde{x}))^{1/q} & l > 0. \end{cases}$$

Proof. We may assume without loss of generality that $B = 1$. To prove (i), we consider for $j \leq n$

$$\begin{aligned} |H^{l,j} * g(\tilde{x})| &\leq \int_{|\tilde{x}-y| \leq 2^{-n}} |H^{l,j}(\tilde{x} - y)g(y)| dy \\ &\leq \|K^{l,j} * \mathcal{F}^{-1}(\Psi_j)(\tilde{x} - y)\|_{L_{q'}} \left(\int_{|\tilde{x}-y| \leq 2^{-n}} |g(y)|^q |\mathbb{1}^{l,j}(\tilde{x} - y)|^q dy \right)^{1/q}. \end{aligned}$$

We note that $\mathbb{1}^{l,j}(\tilde{x} - y)$ is supported around \tilde{x} in either a dyadic annulus of radius $\sim 2^{-j+l}$ for $l > 0$ or a disc of radius 2^{-j} for $l = 0$. As $j \leq n$, the second term above is non-zero only when $l = 0$. In this case, we estimate

$$\begin{aligned} |T^{0,j} * g(\tilde{x})| &\lesssim 2^{jd/q} (\log 2)^{-\alpha} 2^{-nd/q} \left(2^{nd} \int_{|\tilde{x}-y| \leq 2^{-n}} |g(y)|^q dy \right)^{1/q} \quad (\text{using Proposition 3.1}) \\ &\lesssim 2^{(j-n)d/q} (M[g^q](\tilde{x}))^{1/q}. \end{aligned}$$

Summing up in $j < n$, the assertion now follows as $|g| \leq |f|$.

For (ii), we observe that $|T_n^l h(\tilde{x}) - T^l(\tilde{x})| \leq \sum_{j>n} |H^{l,j} * h(\tilde{x})|$. For $j > n$, we then have

$$\begin{aligned} |H^{l,j} * h(\tilde{x})| &\leq \int_{|\tilde{x}-y| \geq 2^{-n}} |H^{l,j}(\tilde{x} - y)h(y)| dy \\ &\leq \|K^{l,j} * \mathcal{F}^{-1}(\Psi_j)(\tilde{x} - y)\|_{L_{q'}} \left(\int_{|\tilde{x}-y| \geq 2^{-n}} |h(y)|^q |\mathbb{1}^{l,j}(\tilde{x} - y)|^q dy \right)^{1/q}. \end{aligned}$$

Again, using the support property of $\mathbb{1}^{l,j}$, we observe that the second term above is non-zero only when $l > 0$ and $j < l + n$. For each $j \in (n, l + n)$, we estimate

$$\begin{aligned} |H^{l,j} * h(\tilde{x})| &\leq 2^{jd/q} (\log(2 + 2^l))^{-\alpha} \left(\int_{|\tilde{x}-y| \geq 2^{-n}} |h(y)|^q |\mathbb{1}^{l,j}(2^j(\tilde{x} - y))|^q dy \right)^{1/q} \quad (\text{using Proposition 3.1}) \\ &\lesssim 2^{jd/q} l^{-\alpha} \left(\int_{|\tilde{x}-y| \sim 2^{-l+j}} |h(y)|^q dy \right)^{1/q} \leq 2^{jd/q} l^{-\alpha} 2^{(-j+l)d/q} \left(2^{(-j+l)d} \int_{|\tilde{x}-y| \leq 2^{-l+j}} |h(y)|^q dy \right)^{1/q} \\ &\leq l^{-\alpha} 2^{ld/q} (M[h^q](\tilde{x}))^{1/q}. \end{aligned}$$

Summing up in $n < j < n + l$ and noting that $|h| \leq |f|$, we get

$$\sum_{j>n} |H^{l,j} * h(\tilde{x})| \lesssim 2^{ld/q} l^{-\alpha+1} (M[f^q](\tilde{x}))^{1/q}.$$

Now for (iii), we consider the terms $H^{l,j} * h(\tilde{x}) - H^{l,j} * h(w)$ separately for $j \leq n$ and $j > n$. The sum $\sum_{j>n} |H^{l,j} * h(\tilde{x})|$ was already dealt with in (ii) and like before, only matters for

$l > 0$. Since $|w - \tilde{x}| \leq 2^{-n-1}$ we have $|w - y| \approx |\tilde{x} - y|$ for $|\tilde{x} - y| \geq 2^{-n}$ and for non-zero l , the previous calculation leads to

$$\sum_{j>n} |H^{l,j} * h(w)| \lesssim \begin{cases} 0 & l = 0. \\ B2^{ld/q} l^{-\alpha+1} (M[f^q](\tilde{x}))^{1/q} & l > 0. \end{cases}$$

It remains to consider the terms for $j \leq n$. We write

$$H^{l,j} * h(\tilde{x}) - H^{l,j} * h(w) = \int_0^1 \int_{|\tilde{x}-y| \geq 2^{-n}} \langle \tilde{x}-w, \nabla(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))(w+s(\tilde{x}-w)-y) \rangle h(y) dy ds.$$

Since $|w - \tilde{x}| \leq 2^{n-1}$ we can replace $|w + s(\tilde{x} - w) - y|$ in the integrand with $|\tilde{x} - y|$. Also Proposition 3.1 yields

$$|\nabla(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))(\tilde{x} - y)| \leq 2^j |(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))(\tilde{x} - y)|.$$

Thus we have

$$\begin{aligned} & |H^{l,j} * h(\tilde{x}) - H^{l,j} * h(w)| \\ & \lesssim 2^j |\tilde{x} - w| \int_{|\tilde{x}-y| \geq 2^{-n}} |(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))(\tilde{x} - y) h(y)| dy \\ & \leq 2^{j-n-1} \|(K^{l,j} * \mathcal{F}^{-1}(\Psi_j))(\tilde{x} - y)\|_{L_{q'}} \left(\int_{|\tilde{x}-y| \geq 2^{-n}} |h(y)|^q |\mathbb{1}^{l,j}((\tilde{x} - y))|^q dy \right)^{1/q}. \end{aligned}$$

Proposition 3.1 now gives us

$$\begin{aligned} |H^{l,j} * h(\tilde{x}) - H^{l,j} * h(w)| & \lesssim 2^{j-n-1} 2^{jd/q} (\log(2 + 2^l))^{-\alpha} \left(\int_{|\tilde{x}-y| \sim 2^{-j+l}} |h(y)|^q dy \right)^{1/q} \\ & \lesssim 2^{j-n-1} 2^{jd/q} (\log(2 + 2^l))^{-\alpha} 2^{(-j+l)d/q} (M[h^q](\tilde{x}))^{1/q} \\ & \lesssim \begin{cases} 2^{ld/q} 2^{j-n-1} (M[f^q](\tilde{x}))^{1/q} & l = 0. \\ 2^{ld/q} 2^{j-n-1} l^{-\alpha} (M[f^q](\tilde{x}))^{1/q} & l > 0. \end{cases} \end{aligned}$$

Summing in $j \leq n$ leads to

$$\sum_{j \leq n} |H^{l,j} * h(\tilde{x}) - H^{l,j} * h(w)| \lesssim \begin{cases} 2^{ld/q} (M[f^q](\tilde{x}))^{1/q} & l = 0. \\ 2^{ld/q} l^{-\alpha} (M[f^q](\tilde{x}))^{1/q} & l > 0. \end{cases}$$

□

We can now prove the main result of this section.

Proposition 5.2. *Let $\alpha > 3$, $q > 1$ and $B(m)$ be as in (6). Let f be a Schwartz function. Then for almost every x and for $0 < \delta \leq 1/2$, we have*

$$T_*^l f(x) \lesssim \frac{1}{(1 - \delta)^{1/q}} (M(|T^l f|^q)(x))^{1/q} + C_d A_l (1 + \delta^{-1/q}) (M(|f|^q)(x))^{1/q}$$

where

$$A_l = \begin{cases} B & l = 0 \\ B2^{ld/q} l^{-\alpha+1} & l > 0. \end{cases}$$

Proof. The proof is essentially the same as that of an analogous result in [4], which is in turn a modification of the argument for the standard Cotlar inequality regarding the truncation of singular integrals (see [8], sec 1.7).

Fix $\tilde{x} \in \mathbb{R}^d$ and $n \in \mathbb{Z}$ and define g , h and q as in the previous lemma. For w (to be chosen later) with $|w - \tilde{x}| \leq 2^{-n-1}$ we can write

$$(14) \quad \begin{aligned} T_n^l f(\tilde{x}) &= T_n^l g(\tilde{x}) + (T_n^l - T^l)h(\tilde{x}) + T^l h(\tilde{x}) \\ &= T_n^l g(\tilde{x}) + (T_n^l - T^l)h(\tilde{x}) + T^l h(\tilde{x}) - T^l h(w) + T^l f(w) - T^l g(w). \end{aligned}$$

By Lemma 5.1, we have

$$|T_n^l g(\tilde{x})| + |(T_n^l - T^l)h(\tilde{x})| + |T^l h(\tilde{x}) - T^l h(w)| \lesssim A_l (M[f^q](\tilde{x}))^{1/q}.$$

All that remains in this case is to consider the term $T^l f(w) - T^l g(w)$ for w in a substantial subset of $B(\tilde{x}, 2^{-n-1})$. By Theorem 4.1 we have that for all $f \in L^q(\mathbb{R}^d)$ and all $\lambda > 0$

$$\text{meas}\{x: |T^l f(x)| > \lambda\} \leq A_l^q \lambda^{-q} \|f\|_q^q.$$

Now let $\delta \in (0, 1/2)$ and consider the set

$$\Omega_n(\tilde{x}, \delta) = \{w: |w - \tilde{x}| < 2^{-n-1}, |T^l g(w)| > 2^{d/q} \delta^{-1/q} A_l (M[f^q](\tilde{x}))^{1/q}\}.$$

In (14) we can estimate the term $|T^l g(w)|$ by $A_l 2^{d/q} \delta^{-1/q} (M[f^q](\tilde{x}))^{1/q}$ whenever $w \in B(\tilde{x}, 2^{-n-1}) \setminus \Omega_n(\tilde{x}, \delta)$. Hence we obtain

$$(15) \quad |T_n^l f(\tilde{x})| \lesssim \inf_{w \in B(\tilde{x}, 2^{-n-1}) \setminus \Omega_n(\tilde{x}, \delta)} |T^l f(w)| + C_d A_l (1 + \delta^{-1/q}) (M[f^q](\tilde{x}))^{1/q}.$$

By the weak type inequality for T^l we have

$$\begin{aligned} \text{meas}(\Omega_n(\tilde{x}, \delta)) &\leq \frac{A_l^q \|g\|_q^q}{2^d \delta^{-1} A_l^q M[f^q](\tilde{x})} = \frac{\delta}{2^d M[f^q](\tilde{x})} \int_{|x-y| \leq 2^{-n}} |f(y)|^q dy \\ &\leq \delta 2^{-d} \text{meas}(B(\tilde{x}, 2^{-n})) = \delta \text{meas}(B(\tilde{x}, 2^{-n-1})). \end{aligned}$$

Hence $\text{meas}(B(\tilde{x}, 2^{-n-1}) \setminus \Omega_n(\tilde{x}, \delta)) \geq (1 - \delta) \text{meas}(B(\tilde{x}, 2^{-n-1}))$ and thus

$$\begin{aligned} \inf_{w \in B(\tilde{x}, 2^{-n-1}) \setminus \Omega_n(\tilde{x}, \delta)} |T^l f(w)| &\leq \left(\frac{1}{\text{meas}(B(\tilde{x}, 2^{-n-1}) \setminus \Omega_n(\tilde{x}, \delta))} \int_{B(\tilde{x}, 2^{-n-1})} |T^l f(w)|^q \right)^{1/q} \\ &\leq \left(\frac{1}{(1 - \delta) \text{meas}(B(\tilde{x}, 2^{-n-1}))} \int_{B(\tilde{x}, 2^{-n-1})} |T^l f(w)|^q \right)^{1/q}. \end{aligned}$$

We obtain

$$|T_n^l f(\tilde{x})| \lesssim \frac{1}{(1 - \delta)^{1/q}} (M(|T^l f|^q)(\tilde{x}))^{1/q} + A_l (1 + \delta^{-1/q}) (M(|f|^q)(\tilde{x}))^{1/q}$$

uniformly in n , which implies Proposition 5.2. \square

The above proposition, in conjunction with Theorem 4.2 immediately leads to the following:

Theorem 5.3. T_*^l is bounded on L^p for $p \in (l + 2, \infty)$.

Proof. Fix $\delta = 1/2$ and $q = l + 2$. We note that both $f \rightarrow T^l f$ and $f \rightarrow (M[f^q])^{1/q}$ are bounded operators on L^p for $p \in (l + 2, \infty)$, with operator norms bounded by $pBl^{-\alpha+1}$ (pB for $l = 0$) and $p/(p - l - 2)$ respectively (upto multiplication by a dimensional constant). Hence, by Proposition 5.2, we obtain

$$\|T_*^l\|_{L^p} \lesssim_d \begin{cases} B, & l = 0 \\ l^{-\alpha+1} \frac{p^2}{p-l-2} B, & l > 0. \end{cases}$$

□

6. WEAK (1,1) BOUNDEDNESS OF THE MAXIMAL OPERATOR

In this section, we will prove that each of the pieces T_*^l is of weak type $(1, 1)$, with the respective norm $\lesssim_d Bl^{-\alpha+1}$ ($\lesssim_d B$ for $l = 0$). Combining this result with Theorem 5.3 and interpolating, we will obtain bounds on the operator norm of T_*^l on L^p for all $p \in (1, \infty)$, with a decay of $l^{-\alpha+2}$. This will allow us to achieve our final goal of summing up the pieces together to get bounds on the operator S^* .

The proof we give here is essentially the same as the one for Theorem 4.1, except for one notable difference. In order to obtain a weak bound for the "good" function in the Calderón-Zygmund decomposition, we use the bounds on L^{p_l} (as given by Theorem 5.3, with $p_l = 4l$) in place of those on L^2 , noting that the operator norm in the former case is $\lesssim Bl^{-\alpha+2}$ (for $l > 0$). The upshot is that the power of l in the weak $(1, 1)$ norm of T_*^l goes up by 1. Instead of repeating the entire argument, we sketch an outline.

Theorem 6.1. T_*^l is weak $(1, 1)$ bounded with

$$\|T_*^l\|_{L^1 \rightarrow L^{1,\infty}} \lesssim \begin{cases} B, & l = 0, \\ Bl^{-\alpha+2}, & l > 0. \end{cases}$$

Proof. Let $n \in \mathbb{Z}$ and $f \in L^1(\mathbb{R}^d)$. Fix $\sigma > 0$. We sketch the proof for $l > 0$ (the one for $l = 0$ proceeds in almost the same way). Also, we might assume $B = 1$. As before, we make a Calderón-Zygmund decomposition of f at the level $l^{\alpha-1}\sigma$. For the "good" function f_1 , we use the bound $\|T_n^l\|_{L^{p_l}} \lesssim Bl^{-\alpha+2}$ on L^{p_l} with $p_l = 4l$ for $l > 0$ and $\|T_n^0\|_{L^3} \lesssim B$ on L^3 . We obtain for $l > 0$

$$\|T_n^l f_1\|_{L^{1,\infty}} \lesssim Bl^{-\alpha+2} \|f\|_{L^1}$$

(and the corresponding result for $l = 0$). The argument for the "bad" part f_0 proceeds the exact way as in proof of Theorem 5.3, with the index n playing no real role, and we get for $l > 0$

$$\|T_n^l f_0\|_{L^{1,\infty}} \lesssim Bl^{-\alpha+1} \|f\|_{L^1}$$

(and the corresponding result for $l = 0$). The result then follows by combining the two estimates followed by taking a supremum over n . □

Theorems 5.3 and 4.1 together via L^p interpolation lead to

Theorem 6.2. T_*^l is bounded on L^p for $p \in (1, \infty)$, with

$$\|T_*^l\|_{L^p} \lesssim_d \begin{cases} B \max(p, (p-1)^{-1}), & l = 0, p \in (1, \infty) \\ B \max(p, (p-1)^{-1}) l^{-\alpha+2}, & l > 0, p \in (1, 4l) \\ B p l^{-\alpha+1}, & l > 0, p \in [4l, \infty). \end{cases}$$

Proof. The result for $l = 0$ is a clear outcome of L^p interpolation and the estimates proved earlier. For $l > 0$ and $p \in [4l, \infty)$, it follows easily from Theorem 5.3 and the observation that $p^2/(p-l-2) \lesssim p$ for $p \geq 4l$. For $l > 0$ and $p \in (1, 4l)$, it is an outcome of interpolating between the weak (1,1) estimate in Theorem 6.1 and the one contained in Theorem 5.3 for $p = 4l$, again making the observation that $p^2/(p-l-2) \lesssim p = 4l$. \square

Finally, we give the proof of Theorem 1.1.

Proof. It is enough to prove the theorem for S_n for a fixed $n \in \mathbb{Z}$, as the result then follows by taking the supremum over $n \in \mathbb{Z}$. For $p \in (1, \infty)$ and a Schwartz function f , we have

$$\|S_n f\|_{L^p} \leq \sum_{l \geq 0} \|S_n^l f\|_{L^p} \leq \sum_{l \geq 0} \|T_n^l f\|_{L^p} + \sum_{l \geq 0} \left\| \sum_{j \leq n} K^{l,j} * \mathcal{F}^{-1}(1 - \Psi_j) * f \right\|_{L^p}.$$

Using Theorem 6.2 for the first sum and Theorem 2.2 for the second one (and noting that summing up for $j \leq n$ instead of $j \in \mathbb{Z}$ does not affect the proof), we get

$$\|S_n f\|_{L^p} \lesssim B \max(p, (p-1)^{-1}) \sum_{l \geq 1} (l^{-\alpha+2} + 2^{-l}) \lesssim_p B.$$

For weak (1,1) boundedness, we argue in a similar way, only using Theorem 6.1 this time in place of Theorem 6.2. \square

REFERENCES

- [1] Anthony Carbery, *Variants of the Calderón-Zygmund theory for L^p -spaces*, Revista Matemática Iberoamericana **2** (1986), no. 4, 381–396.
- [2] Michael Christ, Loukas Grafakos, Petr Honzík, and Andreas Seeger, *Maximal functions associated with Fourier multipliers of Mikhlin-Hörmander type*, Mathematische Zeitschrift **249** (2005), no. 1, 223–240.
- [3] Loukas Grafakos, *Classical Fourier analysis*, Vol. 2, Springer, 2008.
- [4] Shaoming Guo, Joris Roos, Andreas Seeger, and Po-Lam Yung, *A maximal function for families of Hilbert transforms along homogeneous curves*, arXiv preprint arXiv:1902.00096 (2019).
- [5] Lars Hörmander, *Estimates for translation invariant operators in L^p spaces*, Acta Mathematica **104** (1960Dec), no. 1, 93–140.
- [6] Andreas Seeger, *Some inequalities for singular convolution operators in L^p -spaces*, Transactions of the American Mathematical Society **308** (1988), no. 1, 259–272.
- [7] Elias M Stein, *Singular integrals and differentiability properties of functions (pms-30)*, Vol. 30, Princeton university press, 2016.
- [8] Elias M Stein and Timothy S Murphy, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Vol. 3, Princeton University Press, 1993.
- [9] Elias M Stein and Antoni Zygmund, *Boundedness of translation invariant operators on Hölder spaces and L^p -spaces*, Annals of Mathematics (1967), 337–349.
- [10] Charles McCarthy Walter Littman and Nestor Riviere, *L^p -multiplier theorems*, Studia Mathematica **30** (1968), 193–217.
- [11] Thomas H Wolff, *Lectures on harmonic analysis*, Vol. 29, American Mathematical Soc., 2003.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, 480 LINCOLN DR, MADISON, WI-53706, USA

E-mail address: rsrivastava9@wisc.edu