

A constrained minimum criterion for model selection

Min Tsao

Department of Mathematics & Statistics, University of Victoria, Canada

Abstract: We propose a hypothesis test based model selection criterion for the best subset selection of sparse linear models. We show it is consistent in that the probability of its choosing the true model approaches one and the parameter values of its chosen model converge in probability to that of the true model as the sample size goes to infinity. This criterion is capable of controlling the balance between the false active rate and false inactive rate of the selected model, and it can be applied with other methods of model selection such as the lasso. We also demonstrate its accuracy and advantages with a numerical comparison and an application.

Keywords: Sparse linear models; Model selection criterion; Hypothesis test; Best subset selection; Model selection consistency.

1 Introduction

Consider the linear regression model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}, \quad (1)$$

where $\mathbf{y} = (y_1, \dots, y_n)^T$ is the response vector, $\mathbf{X} = [\mathbf{1}, \mathbf{x}_1, \dots, \mathbf{x}_p]$ is the design matrix, $\boldsymbol{\beta} = (\beta_0, \beta_1, \dots, \beta_p)^T$ is the unknown vector of model parameters and $\boldsymbol{\varepsilon} \sim N(\mathbf{0}, \sigma^2 \mathbf{I})$ is the vector of random error. A variable \mathbf{x}_i is said to be active if its parameter $\beta_i \neq 0$ and inactive if $\beta_i = 0$. Suppose model (1) is sparse in that some of variables are inactive. The best subset selection is often used to help identify the true model that contains only and all active variables.

There are a number of model selection criteria for the best subset selection in both frequentist and Bayesian flavours and with a wide range of motivations. The

most commonly used ones include the adjusted R^2 , Akaike information criterion AIC (Akaike, 1974), Bayesian information criterion BIC (Schwarz, 1978), Mallows' C_p statistic (Mallows, 1973) and cross-validation. The AIC and BIC are penalized log-likelihood measures of a model with a penalty term proportional to the model size. The adjusted R^2 is a model size adjusted goodness-of-fit measure. Mallows' C_p statistic and cross-validation are based on measures of prediction error of a model. Under a given criterion, usually the model that minimizes or maximizes the corresponding measure is selected. The sparsity in the selected model is not explicitly pursued and from this standpoint it is only a by-product of the model selection process. There is a large body of literature on model selection criteria and methods. For a comprehensive review, see Kadane and Lazar (2004), Ding, Tarokh and Yang (2018) and Miller (1990).

In this paper, we propose a new model selection criterion based on a hypothesis test which pursues sparsity in the selected model explicitly. In frequentist inference, it is well known that a hypothesis test can be used to derive a point estimator and a confidence interval. The Hodges-Lehmann estimators (Hodges and Lehmann, 1983) derived from various tests are highly efficient. For estimating the unknown true parameter vector of model (1), which we denote by β^t , the least squares estimator may also be viewed as the Hodges-Lehmann estimator derived from the likelihood ratio test for $H_0 : \beta^t = \beta$ as it is of the highest p -value and in the centre of this test induced confidence region for β^t . However, this estimator cannot be used for model selection as it does not generate sparse solutions. In order to find a sparse solution, we derive from the likelihood ratio test a new sparse estimator for β^t by considering the collection of models with p -values above a given level. This collection is also the set of models not rejected by the test at the given level. Our proposed model selection criterion is to choose from this collection the most sparse member as the estimator for β^t . Let $\mathcal{M} = \{M_j\}_{j=1}^{2^p}$ be the collection of 2^p subsets of the p variables in the full model (1) where each M_j represents a subset or model. Let $\hat{\beta}_j$ be the least squares estimator for model M_j . Using the pseudo-norm $\|\beta\|_0$ that counts the number of non-zero elements in a vector β , the proposed criterion may be formulated

as the solution to the following constrained optimization problem,

$$\underset{\mathcal{M}}{\text{minimize}} \|\hat{\beta}_j\|_0 \quad \text{subject to} \quad \lambda(\hat{\beta}_j) \leq \kappa_n, \quad (2)$$

where $\lambda(\beta)$ is the likelihood ratio test statistic defined in (13) and κ_n is the critical value that depends on the significance level α_n of the test. Because of (2), we call the criterion the constrained minimum criterion or CMC for short. We call the solution vector to the optimization problem (2) the CMC solution and its corresponding model the CMC selection. There may be multiple solution vectors, and when this happens we choose the one with the highest likelihood as the CMC solution.

Unlike the existing criteria, the CMC explicitly seeks maximum sparsity from plausible models, i.e., models not rejected by the underlying test at level α_n . As such, it uses the sparsity condition more aggressively and directly than the existing criteria. We show that with a properly chosen α_n the CMC solution converges in probability to the true parameter vector β^t and the probability that the CMC selection is the true model approaches one when the sample size n goes to infinity. We compare the CMC with three popular model selection criteria through a simulation study to demonstrate its excellent accuracy and its ability to control the balance between the false active rate and false inactive rate through the choice of α_n . Furthermore, we demonstrate its consistency and observe that with a properly chosen α_n the CMC selection converges to the true model faster than the BIC selection. We also apply the CMC to perform model selection and estimation for a prostate cancer data set. The CMC is defined for the best subset selection as the optimization in (2) is over the set of all models \mathcal{M} , but it can be easily extended for other methods of model selection such as the lasso by replacing the set of models \mathcal{M} in (2) with the sequence of models found along the lasso path. The underlying idea of inverting a hypothesis test for model selection by choosing the most sparse model not rejected by the test generalizes the Hodges-Lehmann estimator in the context of model selection, and it can also be applied to model selection problems beyond linear models.

The rest of this paper is organized as follows. In Section 2, we present theoretical results about the CMC and the numerical comparison. We also give recommendations

on choosing the α_n level, including a default level. In Section 3, we apply the CMC to analyse the prostate cancer data. We conclude with a few remarks in Section 4. The proof of the theoretical results is relegated to the Appendix.

2 The constrained minimum criterion

We assume that $n > p$ and variables $\mathbf{x}_1, \dots, \mathbf{x}_p$ are linearly independent so that the least squares estimator for the unknown $\boldsymbol{\beta}^t$ of the full model (1),

$$\hat{\boldsymbol{\beta}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}, \quad (3)$$

is available. Although a model M_j may have fewer than p variables, we view its least squares estimator $\hat{\boldsymbol{\beta}}_j$ as a $q = p + 1$ dimensional vector having zeros in elements corresponding to variables not in M_j , e.g., suppose M_2 contains only \mathbf{x}_1 and \mathbf{x}_2 , then elements of $\hat{\boldsymbol{\beta}}_2$ representing the coefficients of $\mathbf{x}_3, \dots, \mathbf{x}_p$ are all set to zero. Similarly, we view the true parameter vector $\boldsymbol{\beta}^t$ as a q dimensional vector.

We now present the asymptotic properties of the CMC. In the following, for a constant $\alpha \in (0, 1)$ we use $F_{1-\alpha, q, n-q}$ to denote the $(1 - \alpha)$ th quantile of the F distribution with q and $(n - q)$ degrees of freedom. We need the following regularity conditions on the design matrix \mathbf{X} :

$$D_n = \frac{1}{n} \sum_{i=1}^n \mathbf{x}_{ri} \mathbf{x}_{ri}^T \rightarrow D, \quad (4)$$

where \mathbf{x}_{ri} is the i th row of \mathbf{X} and D is a $q \times q$ positive definite matrix, and

$$\frac{1}{n} \max_{1 \leq i \leq n} \mathbf{x}_{ri}^T \mathbf{x}_{ri} \rightarrow 0. \quad (5)$$

Conditions (4) and (5) are commonly used to ensure the least squares estimator of the full model $\hat{\boldsymbol{\beta}}$ is consistent and asymptotically normal; see, for example, Knight & Fu (2000). Under these conditions, we have

$$\sqrt{n}(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}^t) \rightarrow N(\mathbf{0}, \sigma^2 D^{-1}). \quad (6)$$

Denote by M_j^* the unknown true model containing only and all active variables and by $\hat{\boldsymbol{\beta}}_j^*$ the least squares estimator for M_j^* . The following theorem gives the consistency of the CMC.

Theorem 2.1 *Suppose conditions (4) and (5) hold. Let α_n be a sequence in $(0, 1)$ such that $\alpha_n \rightarrow 0$ and $F_{1-\alpha_n, q, n-q} = o(n)$, and let*

$$\kappa_n = qF_{1-\alpha_n, q, n-q}. \quad (7)$$

Denote by $\hat{\beta}_{\alpha_n}$ be the CMC solution of (2) with κ_n defined in (7). Then,

$$\hat{\beta}_{\alpha_n} \xrightarrow{p} \beta^t \quad (8)$$

as $n \rightarrow +\infty$, and the probability that the true model is chosen satisfies

$$\lim_{n \rightarrow +\infty} P(\hat{\beta}_{\alpha_n} = \hat{\beta}_j^*) = 1. \quad (9)$$

A proof of this theorem and a remark about the condition on α_n are given in the Appendix. Using the CMC for model selection amounts to using the CMC solution $\hat{\beta}_{\alpha_n} = (\hat{\beta}_{\alpha_n}^0, \hat{\beta}_{\alpha_n}^1, \dots, \hat{\beta}_{\alpha_n}^p)^T$ as a classifier for the p variables. We classify an \mathbf{x}_j as active if $\hat{\beta}_{\alpha_n}^j \neq 0$ and inactive if $\hat{\beta}_{\alpha_n}^j = 0$, so result (9) also has the interpretation that the probability of no misclassification goes to 1.

We refer to α_n as the α -level of the CMC. It controls the balance between the false active rate and false inactive rate in the variable classification process. To see this, the CMC solution $\hat{\beta}_{\alpha_n}$ is the minimizer of the model size under the constraint $\lambda(\hat{\beta}_j) \leq \kappa_n$ and κ_n depends on α_n . By (7), when α_n increases κ_n decreases, so the minimum is taken over an increasingly smaller set of $\hat{\beta}_j$ values and thus gets larger. The selected model will therefore have more and more variables. At the extreme value of $\alpha_n = 1$, $\kappa_n = 0$ and the CMC solution is the least squares estimator $\hat{\beta}$ for the full model which, with probability one, has no zeros in its elements. Thus, the CMC selection is the full model with a 100% false active rate and a 0% false inactive rate. At the other extreme value of $\alpha_n = 0$, $\kappa_n = \infty$ and the CMC selection is the empty model containing no variables, so it has a 0% false active rate and a 100% false inactive rate. By empirically exploring different α -levels between 0 and 1, we may be able to find one that gives the desired balance between the two rates. In real applications where the sample size n is fixed, we usually set the α -level to a convenient value such as 0.1, 0.5 or 0.9. See examples below.

We now present a numerical comparison on the accuracies of the CMC, the adjusted R^2 , Mallows' C_p which is equivalent to the AIC in the present context, and the BIC. We also provide recommendations on selecting the α -level for the CMC based on simulation results. The model used for this comparison is

$$\mathbf{y} = \beta_0 \mathbf{1} + \mathbf{X}' \boldsymbol{\beta}' + \boldsymbol{\varepsilon}, \quad (10)$$

where $\boldsymbol{\varepsilon} \sim N(\mathbf{0}, \sigma^2 \mathbf{I})$ with $\sigma^2 = 1$, $\mathbf{X}' = [\mathbf{x}_1, \dots, \mathbf{x}_p]$, and the parameter values are $\beta_0 = 1$ and $\boldsymbol{\beta}' = (\beta_1, \dots, \beta_{p^*}, 0, \dots, 0)^T$ with $\beta_1 = \dots = \beta_{p^*} = 1$, so only the first p^* variables are active. We first consider the case where elements in \mathbf{X}' are random numbers generated from a standard normal distribution. We call this the case of weakly correlated predictor variables as the observed correlations of the generated variables are small. Table 1 contains values of the (false inactive rate, false active rate) pairs for six model selection criteria at 12 different combinations of n , p and p^* . The false inactive rate is the observed proportion of active variables misclassified as inactive and the false active rate is that of inactive variables misclassified as active, so when both rates are zero the selected model is the true model. Table 1 is computed using R package “leaps” by Lumley (2020) and each entry pair is based on 1000 simulation runs. For each run, we first generate an \mathbf{X}' matrix and then through equation (10) a random \mathbf{y} vector. We then perform the best subset selection using \mathbf{X}' and \mathbf{y} with the six criteria to find their chosen models and compute their (false inactive rate, false active rate) pairs based on the chosen models. We repeat this process 1000 times and Table 1 records the average of the 1000 rate pairs. The subscript α in CMC_α is the α -level for the CMC. We make the following observations based on results in Table 1:

1. When the sample size n is not large relative to the number of variables p (the three cases in Table 1 where $n = 2p$), none of the six criteria performed very well. The adjusted R^2 , C_p and BIC have low false inactive rates but high false active rates. The $\text{CMC}_{0.5}$ and $\text{CMC}_{0.1}$ have high false inactive rates but low false active rates. If we treat false active and false inactive as equally serious errors and rank the six criteria by the overall error rate (defined as the sum of the

Table 1: Variable classification accuracy comparison for the case of weakly correlated predictor variables: the entries are simulated (false inactive rate, false active rate) of the adjusted R^2 , Mallows' C_p (AIC), BIC and three CMC_α criteria for 12 scenarios. The bold CMC results are those at recommended α -levels. Results in the table are the average of the two rates for 1000 simulation runs rounded to the second digit after the decimal point, and (0.00, 0.00)* means both rates are zero in all 1000 runs.

(n, p, p^*)	Adj R^2	C_p (AIC)	BIC	$\text{CMC}_{0.9}$	$\text{CMC}_{0.5}$	$\text{CMC}_{0.1}$
(20, 10, 5)	(0.03, 0.39)	(0.06, 0.21)	(0.05, 0.24)	(0.07, 0.16)	(0.18, 0.07)	(0.41, 0.03)
(30, 10, 5)	(0.00, 0.35)	(0.00, 0.18)	(0.01, 0.13)	(0.01, 0.11)	(0.04, 0.03)	(0.17, 0.01)
(40, 10, 5)	(0.00, 0.32)	(0.00, 0.17)	(0.00, 0.09)	(0.00, 0.10)	(0.01, 0.02)	(0.06, 0.00)
(50, 10, 5)	(0.00, 0.32)	(0.00, 0.18)	(0.00, 0.08)	(0.00, 0.09)	(0.00, 0.02)	(0.01, 0.00)
(40, 20, 10)	(0.00, 0.35)	(0.00, 0.18)	(0.00, 0.15)	(0.01, 0.06)	(0.06, 0.02)	(0.17, 0.02)
(60, 20, 10)	(0.00, 0.30)	(0.00, 0.17)	(0.00, 0.08)	(0.00, 0.04)	(0.01, 0.01)	(0.04, 0.00)
(80, 20, 10)	(0.00, 0.28)	(0.00, 0.16)	(0.00, 0.06)	(0.00, 0.03)	(0.00, 0.00)	(0.00, 0.00)
(100, 20, 10)	(0.00, 0.26)	(0.00, 0.16)	(0.00, 0.05)	(0.00, 0.03)	(0.00, 0.00)	(0.00, 0.00)*
(60, 30, 15)	(0.00, 0.31)	(0.00, 0.17)	(0.00, 0.11)	(0.00, 0.03)	(0.02, 0.01)	(0.08, 0.00)
(90, 30, 15)	(0.00, 0.27)	(0.00, 0.16)	(0.00, 0.07)	(0.00, 0.02)	(0.00, 0.01)	(0.01, 0.00)
(120, 30, 15)	(0.00, 0.24)	(0.00, 0.16)	(0.00, 0.04)	(0.00, 0.01)	(0.00, 0.00)	(0.00, 0.00)
(150, 30, 15)	(0.00, 0.22)	(0.00, 0.15)	(0.00, 0.04)	(0.00, 0.01)	(0.00, 0.00)	(0.00, 0.00)*

two error rates), then the $\text{CMC}_{0.9}$ is the best of the three CMC criteria. For such small sample cases, the α -level of the CMC needs to be set to a high level or the false inactive rate and the overall error rate could be high.

2. When n is large relative to p (cases where $n \geq 3p$), the CMC is very accurate. While all 6 criteria have zero or low false inactive rates, the three CMC criteria also have low false active rates, whereas the adjusted R^2 and C_p still have double-digit false active rates. For such large sample cases, the CMC outperforms the other three criteria in terms of the overall error rate. In particular, overall the $\text{CMC}_{0.5}$ is the most accurate among the six criteria.
3. Focusing now on the α -level of the CMC, we have noted that the false inactive rate decreases and the false active rate increases when the α -level increases. We see from Table 1 that for any given (n, p, p^*) combination, this is indeed the case. Based on our experience for the case of weakly correlated predictor variables, to achieve a small overall error rate we recommend setting the α -level to 0.9 if $n \leq 2p$, to 0.5 if $2p < n \leq 4p$, and to 0.1 only when $n > 4p$. The CMC results for such recommended levels are in bold font in Table 1. In real data analysis, it may be helpful to use two levels at the same time. See Section 3 for a real example and Section 4 for a discussion about this point.

Table 2 constructed using relevant results from Table 1 illustrates the consistency of the CMC selection in (9) that the probability of no classification errors approaches one. It shows that as n increases from 40 to 100 and α_n decreases from 0.90 to 0.10, the false inactive and false active rates both go down towards zero as predicted by (9). Noting that the rate values in Tables 1 and 2 are the averages of such rates in 1000 runs, the pair of zero rates for the case of $n = 100$ and $\alpha_n = 0.10$ implies none of the underlying 1000 runs had any classification error, i.e., the CMC selection is 100% accurate in these runs. In practice, α_n does not have to be very small in order for the CMC to be very accurate so long as the sample size is large; e.g., for the cases of $\alpha_n = 0.5$ in Table 1, we also observed zero or near zero error rates at large sample sizes of $n \geq 3p$. It is worth noting that the CMC is the only criterion that reached

Table 2: An example illustrating the consistency of the CMC (9): the probability of false inactive rate (FIR) and false active rate (FAR) both being zero approaches 1 as $n \rightarrow \infty$ and $\alpha_n \rightarrow 0$.

(n, p, p^*)	α_n	(FIR, FAR)
(40, 20, 10)	0.90	(0.01, 0.06)
(60, 20, 10)	0.50	(0.01, 0.01)
(100, 20, 10)	0.10	(0.00, 0.00)*

zero error rates at $n = 5p$ for $p = 20, 30$. While Adjusted R^2 and C_p criteria are not expected to reach zero rates, the BIC is consistent in the current setting and it should reach zero rates as n goes to infinity. The observation that it did not even at $n = 5p$ suggests that the error rates of the BIC converge to zero at a slower rate than that of the CMC.

We now consider a case where there are systematic correlations among the predictor variables and examine how such correlations affect the performance of the model selection criteria. For this case, we set $(p, p^*) = (20, 10)$ so that in model (10) $\mathbf{x}_1, \dots, \mathbf{x}_{10}$ are active variables and $\mathbf{x}_{11}, \dots, \mathbf{x}_{20}$ are inactive variables. We then introduce correlations among 5 active variables and do the same for 5 inactive variables as follows. Let $\mathbf{z}_1, \dots, \mathbf{z}_{22}$ be 22 independent n -variate standard normal random vectors. For some $w \in (0, 1)$, define

$$\mathbf{x}_i = (1 - w)\mathbf{z}_i + w\mathbf{z}_{21} \text{ for } i = 1, \dots, 5 \text{ and } \mathbf{x}_i = \mathbf{z}_i \text{ for } i = 6, \dots, 10; \text{ and}$$

$$\mathbf{x}_i = (1 - w)\mathbf{z}_i + w\mathbf{z}_{22} \text{ for } i = 11, \dots, 15, \text{ and } \mathbf{x}_i = \mathbf{z}_i \text{ for } i = 16, \dots, 20,$$

so the first 5 active variables are correlated with a theoretical pairwise correlation coefficient of $\rho = w^2/[(1 - w)^2 + w^2]$. The first 5 inactive variables also have the same pairwise correlation coefficient. When we generate the \mathbf{x}_i through the \mathbf{z}_i , the observed correlation coefficients between the correlated \mathbf{x}_i are approximately ρ which goes to 1 as w goes to 1. This allows us to simulate predictor variables with different degrees of correlation by changing the w value. Table 3 contains simulated (false inactive rate, false active rate) of the adjusted R^2 , Mallows' C_p (AIC), BIC and three CMC_α

criteria for 3 levels of correlations $\rho = 0.3, 0.5, 0.8$, representing moderate, medium and strong levels of correlation. With half of the variables correlated, the accuracies of all criteria are negatively affected when compared to the weakly correlated case in the middle section of Table 1 where (p, p^*) is also $(20, 10)$. For all six criteria, the biggest impact of the correlation in predictor variables is an increase in false inactive rates, especially at $\rho = 0.8$, when compared to the case of weak correlation in Table 1. This is not unexpected as it is known that when active variables are strongly correlated, some of them tend to be dropped in the variable selection process. For the CMC, this increase in the false inactive rate slows the speed at the which the two error rates converge to zero; e.g., in the middle section of Table 1, the rates reached zero when $n = 5p$, but in Table 3 when $\rho = 0.8$ the rates reached zero only when $n = 20p$. Other examples containing a large number of moderately to strongly correlated variables that we have studied also showed a similar slowing in convergence to zero. Further, for $n \leq 5p$ the most accurate CMC level is often the 0.9 level. Based on these findings, for situations with moderately to strongly correlated variables we recommend setting the α -level to 0.9 if $n \leq 5p$, and to 0.5 if $5p < n \leq 15p$. Level 0.1 may be used when $n > 15p$.

In practice, whether or not there are correlated predictor variables may be determined by examining the correlation matrix of the variables. If one is unsure after examining the correlation matrix or wishes to avoid looking at this issue when applying the CMC, we recommend the 0.9 level as the default α -level for such situations. Numerical results such as those in Tables 1 and 2 show that at this level, the CMC is either similar to or better than the BIC in terms of the overall error rate regardless the underlying correlations of the variables and sample size, and it is consistently better than the adjusted R^2 and Mallows' C_p (AIC). The availability of such a default level makes the CMC a practical alternative to other criteria as a user does not have to fine-tune its α -level in order to make CMC competitive; the CMC under the default level already has better performance than other criteria. Nevertheless, if we want to realize the full potential of the CMC, we need to optimize the choice of its α -level by examining the correlations of the predictor variables and following the detailed

Table 3: Variable classification accuracy comparison for a case of correlated predictor variables: the entries are simulated (false inactive rate, false active rate) of the adjusted R^2 , Mallows' C_p (AIC), BIC and three CMC_α criteria. The bold CMC results are those at recommended CMC levels. Results in the table are the average of the two rates for 1000 simulation runs for the case of $(p, p^*) = (20, 10)$.

(ρ, n)	Adj R^2	C_p (AIC)	BIC	$\text{CMC}_{0.9}$	$\text{CMC}_{0.5}$	$\text{CMC}_{0.1}$
(0.3, 40)	(0.02, 0.35)	(0.03, 0.18)	(0.03, 0.15)	(0.06, 0.07)	(0.14, 0.03)	(0.26, 0.01)
(0.3, 60)	(0.00, 0.30)	(0.00, 0.17)	(0.01, 0.09)	(0.01, 0.04)	(0.06, 0.01)	(0.13, 0.00)
(0.3, 100)	(0.00, 0.26)	(0.00, 0.16)	(0.00, 0.05)	(0.00, 0.03)	(0.01, 0.00)	(0.04, 0.00)
(0.3, 200)	(0.00, 0.20)	(0.00, 0.15)	(0.00, 0.03)	(0.00, 0.03)	(0.00, 0.00)	(0.00, 0.00)
(0.5, 40)	(0.04, 0.35)	(0.06, 0.18)	(0.07, 0.16)	(0.11, 0.08)	(0.18, 0.03)	(0.30, 0.02)
(0.5, 60)	(0.00, 0.29)	(0.00, 0.17)	(0.02, 0.09)	(0.04, 0.05)	(0.09, 0.02)	(0.17, 0.00)
(0.5, 100)	(0.00, 0.26)	(0.00, 0.17)	(0.00, 0.05)	(0.00, 0.03)	(0.02, 0.01)	(0.08, 0.00)
(0.5, 200)	(0.00, 0.20)	(0.00, 0.16)	(0.00, 0.03)	(0.00, 0.03)	(0.00, 0.00)	(0.00, 0.00)
(0.8, 40)	(0.11, 0.36)	(0.15, 0.20)	(0.16, 0.16)	(0.19, 0.09)	(0.26, 0.03)	(0.36, 0.01)
(0.8, 60)	(0.04, 0.31)	(0.07, 0.18)	(0.10, 0.09)	(0.11, 0.07)	(0.18, 0.02)	(0.25, 0.00)
(0.8, 100)	(0.01, 0.24)	(0.01, 0.16)	(0.04, 0.05)	(0.05, 0.04)	(0.10, 0.01)	(0.16, 0.00)
(0.8, 200)	(0.00, 0.18)	(0.00, 0.16)	(0.00, 0.03)	(0.00, 0.03)	(0.02, 0.00)	(0.08, 0.00)
(0.8, 400)	(0.00, 0.12)	(0.00, 0.16)	(0.00, 0.02)	(0.00, 0.03)	(0.00, 0.00)	(0.00, 0.00)

recommendations about the selection of the α -level given above.

Finally, for brevity of presentation we have only presented examples where parameters of active variables are all 1 and $p^* = p/2$. We have tried different p^* and used parameter values that differ in size and sign, and obtained similar observations concerning the relative performance of the six criteria.

3 Application to a prostate cancer data

We now apply the CMC to analyse a prostate cancer data set from Stamey et al. (1989). This data set had been analysed in the book by Hastie, Tibshirani and Friedman (2009). It is concerned with the correlation between the level of prostate specific antigen (PSA) and a number of clinical measures in 97 men who were about to receive a radical prostatectomy. There are 9 variables in the data set: the log of PSA (lpsa), log cancer volume (lcavol), log prostate weight (lweight), age, log of benign prostatic hyperplasia amount (lbph), seminal vesicle invasion (svi), log of capsular penetration (lcp), Gleason score (gleason), and percent of Gleason scores 4 or 5 (pgg45). The goal is to predict the lpsa by using the other 8 variables through a linear model. Scatter plots of the variables can be found in Figure 1.1 of the book which suggest a linear model is appropriate. Table 3.3 of the book contains the estimated model based on measurements from a subset of 67 men by six different methods: the ordinary least squares regression, the best subset selection, ridge regression, the lasso, principle component regression and partial least squares regression. The model selection and penalty parameter selection criterion used for the last five methods is cross-validation. To compute Table 3.3, the 8 predictor variables were all standardized.

Since we do not need to split the sample for cross-validation, we use all 97 observations in the best subset selection. Due to the large sample size relative to the number of variables and no strong correlations were found among the 8 variables, we choose to use α -levels 0.1 and 0.5. To facilitate comparison with Table 3.3 in Hastie, Tibshirani and Friedman (2009), we also standardized the predictor variables. Ta-

Table 4: Estimated models by the best subset selection method with five variable selection criteria. The $\text{CMC}_{0.1}$, $\text{CMC}_{0.5}$ and BIC selected the same model. Notation: “***”: significant at 0.001 level; “**”: significant at 0.01 level; “*”: significant at 0.05 level.

Variable	LS	Adj R^2	C_p	BIC	$\text{CMC}_{0.5}$	$\text{CMC}_{0.1}$
Intercept	2.478 ***	2.478	2.478	2.478	2.478	2.478
lcavol	0.665 ***	0.671	0.622	0.619	0.619	0.619
lweight	0.266 **	0.263	0.229	0.283	0.283	0.283
age	−0.158	−0.155				
lbph	0.140	0.141	0.114			
svi	0.315 **	0.311	0.292	0.275	0.275	0.275
lcp	−0.148	−0.146				
gleason	0.035					
pgg45	0.125	0.150				

Table 4 contains models chosen by five criteria. It shows that $\text{CMC}_{0.1}$, $\text{CMC}_{0.5}$ and the BIC selected the same model with 3 variables lcavol+lweight+svi. The C_p criterion selected one more variable, lbph. The adjusted R^2 criterion dropped only one variable, gleason, from the full model. Comparing Table 4 with Table 3.3 in Hastie, Tibshirani and Friedman (2009), we see that there are some differences in the least squares estimates which are due to our use of the full data set for estimation and model selection, but estimates for parameters that are significant are approximately equal. While the best subset selection based on the cross-validation in Table 3.3 selected only lcavol+lweight, that based on the five criteria in Table 4 all selected more variables. In particular, the three-variable model chosen by $\text{CMC}_{0.1}$, $\text{CMC}_{0.5}$ and BIC has one more variable than the best subset selection in Table 3.3 and one fewer than the lasso selection in Table 3.3 which is lcavol+lweight+svi+lbph. Since lbph is not significant in the full model, the estimated parameter value for lbph in the lasso selection is only 0.002 (which is substantially smaller than other values) and it

is known that the lasso with cross-validation tends to include false active variables, lbph is likely a false active variable. So our final selection is the three-variable model lcavol+lwight+svi chosen by the two CMC criteria and the BIC.

4 Concluding remarks

The CMC's ability to control the balance between the false active rate and false inactive rate through its α -level is rooted in the underlying hypothesis test's ability to control the Type-I and Type-II errors through its significance level. This feature of the CMC is particularly useful when one type of misclassification is considered more serious than another and we want to keep the rate of the more serious misclassification to a low level which can be achieved by adjusting the α -level. For real applications, we may consider one or two α -levels besides the recommended level and compare their CMC selections to make a final choice. Since a higher α -level gives the CMC selection a lower false inactive rate and a lower level gives it a lower false active rate, if and when both levels lead to the same selection as in the prostate cancer data example, we have added confidence in the accuracy of the CMC selection as both rates associated with this selection are expected to be low. This ability is an advantage to the CMC as other model selection criteria do not have control on the balance between the two rates.

The best subset selection used to be only viable for model selection problems with $p < 40$ due to its high computational cost. Recently, however, Bertsimas, King and Mazumder (2016) developed an efficient way to perform the best subset selection for p in the hundreds and even thousands. Thus, the CMC may be applied to perform the best subset selection for problems with a large p . When the p is too large and the best subset selection becomes impractical, we may apply the CMC with the lasso or variants of the lasso discussed in Hastie, Tibshirani and Wainwright (2015). When applied with the lasso, for example, the set of models under consideration is the sequence of models found along the lasso path which we denote with \mathcal{M}_L , so the minimization in (2) is done over \mathcal{M}_L , not over the set of all possible models. The

proof of Theorem 2.1 can be modified to show that in this case the CMC solution is still a consistent estimator for β^t and, under an extra condition that the true model is in \mathcal{M}_L with probability tending to one as the sample size increases, the CMC selection is also consistent.

The CMC requires the condition that $n > p$ to ensure the least squares estimators $\hat{\beta}_j$ and their likelihood ratios are all well-defined so that the likelihood ratio test is valid. For high dimensional situations where $p > n$, we may reduce the dimension using the lasso before applying the CMC. To do so, we first use the lasso with the cross-validation criterion to select a model and then use this model as the “full model” for the best subset selection with the CMC. This “full model” has fewer than n variables and it tends to have a high number of false active variables. Simulation results (not included here) show that the CMC is very effective at reducing its number of false active variables.

To conclude, the CMC may be extended to handle model selection problems for other types of models such as generalized linear models. To define a CMC for a model selection problem, the key ingredient we need is a powerful test for individual models with a corresponding Hodges-Lehmann estimator that is consistent and a corresponding confidence region that can be kept at $o_p(n)$ size when the confidence level approaches one. When there is such a powerful test, the resulting CMC would be a strong competitor to the existing criteria.

Appendix

Under model (1), the log-likelihood function for β is

$$l(\beta) = -\frac{n}{2} \ln(2\pi) - n \ln(\sigma) - \frac{1}{2\sigma^2} \|\mathbf{y} - \mathbf{X}\beta\|_2^2, \quad (11)$$

which is maximized at $\beta = \hat{\beta}$ for any fixed σ^2 . The likelihood ratio statistic for testing the null hypothesis $H_0 : \beta^t = \beta$ is

$$-2\{l(\beta) - l(\hat{\beta})\} = \frac{1}{\sigma^2} \left\{ \|\mathbf{y} - \mathbf{X}\beta\|_2^2 - \|\mathbf{y} - \mathbf{X}\hat{\beta}\|_2^2 \right\} \quad (12)$$

which contains the unknown parameter σ^2 . For model selection, σ^2 is a nuisance parameter and it does not affect the relative likelihood of different $\boldsymbol{\beta}$ values. Let $\hat{\sigma}^2$ be the residual mean square from the least squares regression for the full model (1). Since $\hat{\sigma}^2$ is a consistent estimator for σ^2 , we substitute it for the unknown σ^2 in (12) and the likelihood ratio test statistic becomes

$$\lambda(\boldsymbol{\beta}) = \frac{1}{\hat{\sigma}^2} \left\{ \|\mathbf{y} - \mathbf{X}\boldsymbol{\beta}\|_2^2 - \|\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}\|_2^2 \right\}. \quad (13)$$

Under H_0 , the asymptotic distribution of a likelihood ratio test statistic involving an estimated parameter is often a scaled or a weighted χ^2 distribution. That of $\lambda(\boldsymbol{\beta})$ in (13) is also a scaled χ^2 distribution, but since the exact finite sample distribution of $\lambda(\boldsymbol{\beta})$ is available, we will use the exact distribution.

Proof of Theorem 2.1. It may be verified that the likelihood ratio statistic $\lambda(\boldsymbol{\beta})$ defined in (13) can be expressed as

$$\lambda(\boldsymbol{\beta}) = \frac{1}{\hat{\sigma}^2} \left\{ (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T \mathbf{X}^T \mathbf{X} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}) \right\}, \quad (14)$$

and that under $H_0 : \boldsymbol{\beta}^t = \boldsymbol{\beta}$ the distribution of $q^{-1}\lambda(\boldsymbol{\beta})$ is an $F_{q,n-q}$ distribution. Thus, the collection of $\boldsymbol{\beta}$ values not rejected by the likelihood ratio test at the α_n level form the following $100(1 - \alpha_n)\%$ confidence region for $\boldsymbol{\beta}^t$,

$$\mathcal{C}_{\alpha_n} = \left\{ \boldsymbol{\beta} : \frac{(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T \mathbf{X}^T \mathbf{X} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})}{q\hat{\sigma}^2} \leq F_{1-\alpha_n, q, n-q} \right\}. \quad (15)$$

By (7) and (14), we may express \mathcal{C}_{α_n} in terms of $\lambda(\boldsymbol{\beta})$ and κ_n as

$$\mathcal{C}_{\alpha_n} = \{\boldsymbol{\beta} : \lambda(\boldsymbol{\beta}) \leq \kappa_n\}. \quad (16)$$

Since the CMC solution $\hat{\boldsymbol{\beta}}_{\alpha_n}$ for (2) satisfies $\lambda(\hat{\boldsymbol{\beta}}_{\alpha_n}) \leq \kappa_n$, (16) implies that $\hat{\boldsymbol{\beta}}_{\alpha_n} \in \mathcal{C}_{\alpha_n}$. This, (15) and $F_{1-\alpha_n, q, n-q} = o(n)$ imply

$$\frac{1}{n}(\hat{\boldsymbol{\beta}}_{\alpha_n} - \hat{\boldsymbol{\beta}})^T \mathbf{X}^T \mathbf{X} (\hat{\boldsymbol{\beta}}_{\alpha_n} - \hat{\boldsymbol{\beta}}) \leq \frac{1}{n}q\hat{\sigma}^2 F_{1-\alpha_n, q, n-q} = o_p(1), \quad (17)$$

where the right-hand side is $o_p(1)$ instead of $o(1)$ because $\hat{\sigma}^2$ is not a constant but it converges in probability to σ^2 . By (4) and (17), $\|\hat{\boldsymbol{\beta}}_{\alpha_n} - \hat{\boldsymbol{\beta}}\|_2^2 = o_p(1)$. Also, by (6) $\|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}^t\|_2^2 = O_p(n^{-1})$. It follows that

$$\|\hat{\boldsymbol{\beta}}_{\alpha_n} - \boldsymbol{\beta}^t\|_2^2 \leq \|\hat{\boldsymbol{\beta}}_{\alpha_n} - \hat{\boldsymbol{\beta}}\|_2^2 + \|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}^t\|_2^2 = o_p(1), \quad (18)$$

which proves the consistency of the CMC solution (8).

To prove the CMC selection consistency (9), we first examine which elements of vectors β in \mathcal{C}_{α_n} may be zero and which elements may not be zero when n is large. Define an event

$$A = \{\text{Elements of } \beta \text{ in } \mathcal{C}_{\alpha_n} \text{ corresponding to non-zero elements of } \beta^t \text{ are also non-zero}\}.$$

By the definition of \mathcal{C}_{α_n} in (15), vectors β in \mathcal{C}_{α_n} satisfy

$$\frac{1}{n}(\beta - \hat{\beta})^T \mathbf{X}^T \mathbf{X}(\beta - \hat{\beta}) \leq \frac{1}{n} q \hat{\sigma}^2 F_{1-\alpha_n, q, n-q} = o_p(1), \quad (19)$$

uniformly. Similar to (18), by the triangular inequality we have $\|\beta - \beta^t\|_2^2 = o_p(1)$ uniformly for all $\beta \in \mathcal{C}_{\alpha_n}$. This implies individual elements of β converge in probability to corresponding elements of β^t uniformly and thus $P(A) \rightarrow 1$ as the sample size n goes to infinity. When event A occurs, among the set of vectors $\{\hat{\beta}_j\}_{j=1}^{2^p}$ only those for models containing all active variables can be in \mathcal{C}_{α_n} , so $\hat{\beta}_j^*$ of the true model M_j^* would be the most sparse member of $\{\hat{\beta}_j\}_{j=1}^{2^p}$ that may possibly be in \mathcal{C}_{α_n} . It follows that $\{\hat{\beta}_j^* \in \mathcal{C}_{\alpha_n}\} \cap A$ implies $\{\hat{\beta}_{\alpha_n} = \hat{\beta}_j^*\}$, so

$$P(\hat{\beta}_{\alpha_n} = \hat{\beta}_j^*) \geq P(\{\hat{\beta}_j^* \in \mathcal{C}_{\alpha_n}\} \cap A) \rightarrow P(\hat{\beta}_j^* \in \mathcal{C}_{\alpha_n}) \quad (20)$$

as n goes to infinity. Also, $\hat{\beta}_j^* \in \mathcal{C}_{\alpha_n}$ if $\beta^t \in \mathcal{C}_{\alpha_n}$ since $\hat{\beta}_j^*$ has a higher likelihood and thus a smaller likelihood ratio than β^t . It follows that

$$P(\hat{\beta}_j^* \in \mathcal{C}_{\alpha_n}) \geq P(\beta^t \in \mathcal{C}_{\alpha_n}) = 1 - \alpha_n \rightarrow 1. \quad (21)$$

Equations (20) and (21) then imply (9). \square

We now remark on the condition of $\alpha_n \rightarrow 0$ and $F_{1-\alpha_n, q, n-q} = o(n)$ in Theorem 2.1. It requires that α_n approaches 0 slowly so that $F_{1-\alpha_n, q, n-q}/n$ approaches 0 when n goes to infinity. A sequence of α_n satisfying this condition can be constructed in many ways, e.g., by letting

$$1 - \alpha_n = P(F_{q, n-q} \leq n^\delta) \quad (22)$$

where δ is a small positive number so that $F_{1-\alpha_n, q, n-q} = n^\delta = o(n)$. Since $F_{q, n-q}$ converges to χ_q^2/q as n goes to infinity, (22) implies that when n is large α_n is essentially $1 - P(\chi_q^2 \leq qn^\delta)$ which satisfies $\alpha_n \rightarrow 0$. In real applications where n is fixed, we do not choose the α_n using (22). Instead, we set it to convenient values such as 0.1, 0.5 and 0.9 depending on the relative size of n and p and whether there are correlated variables, or we simply set it to the default value of 0.9, as we recommended in Section 2. Nevertheless, the condition is an important reminder that we should not let α_n to be too close to 0 when n is not large. Otherwise, the CMC selection may be inaccurate. For example, the $\text{CMC}_{0.1}$ for the small sample case of (20, 10, 5) in Table 1 has a high false inactive rate of 41%. On the other hand, when the sample size is really large, very small α -levels could be used. As an example of this, for all three combinations of p and p^* in Table 1, when $n = 300$ the simulated CMC error rates at the 0.1, 0.05 and 0.01 levels are all zero, whereas the false active rate of the BIC is still around 2% and that of the adjusted R^2 and C_p are still in double digits.

We have observed in Section 2 that the α -level does not have to be very small to obtain accurate results as long as the sample size is large. As a part of future research on the CMC, we will look for a theoretical explanation for this phenomenon which may lead to a more effective way of choosing the α -level.

References

- [1] Akaike, H. (1974). A new look at the statistical model identification *IEEE Transactions on Automatic Control*, 19, 716–723.
- [2] Bertsimas, D., King, A. and Mazumder, R. (2016). Best subset selection via a modern optimization lens. *Annal of Statistics*, 44, 813–852.
- [3] Ding, J., Tarokh, V. and Yang, Y. (2018). Model Selection Techniques: An Overview. *IEEE Signal Processing Magazine*, 35, 16–34.
- [4] Hastie, T., Tibshirani, R. and Friedman, J. (2009). *Elements of Statistical Learning: Data Mining, Inference and Predictions*. 2nd edition. Springer Verlag, New York.
- [5] Hastie, T., Tibshirani, R. and Wainwright, M. (2015). *Statistical Learning with Sparsity*. 2nd edition. CRC Press, Boca Raton, FL.

- [6] Hodges, J. L. Jr. and Lehmann, E. L. (1983). Hodges-Lehmann estimators. *Encyclopedia of Statistical Sciences* edited by S. Kotz, N. L. Johnson, and C. B. Read, **3**, 463–465.
- [7] Kadane, J. B. and Lazar, N. A. (2004). Methods and criteria for model selection, *Journal of the American Statistical Association*, 99, 279–290.
- [8] Knight, K. and Fu, W. (2000). Asymptotics for lasso-type estimators, *Annals of Statistics*, 28, 1356–1378.
- [9] Lumley, T. (2020). Package ‘leaps’, an R package available at <https://cran.r-project.org>.
- [10] Mallows, C. L. (1973). Some Comments on CP, *Technometrics*, 15, 661–675.
- [11] Miller, A. J. (1990). *Subset Selection in Regression*. Chapman and Hall, New York.
- [12] Schwarz, G. E. (1978). Estimating the dimension of a model, *Annals of Statistics*, 6, 461–464.
- [13] Stamey, T., Kabalin, J., McNeal, J., Johnstone, I., Freiha, F., Redwine, E. and Yang, N. (1989). Prostate specific antigen in the diagnosis and treatment of adenocarcinoma of the prostate II radical prostatectomy treated patients, *Journal of Urology*, 16, 1076–1083.