

DIXMIER TRACES FOR DISCRETE PSEUDO-DIFFERENTIAL OPERATORS

DUVÁN CARDONA, CÉSAR DEL CORRAL, AND VISHVESH KUMAR

Dedicated to Professor Lázaro Reicht on his 79 birthday

ABSTRACT. In this paper we provide sharp results for the Dixmier traceability of discrete pseudo-differential operators on $\ell^2(\mathbb{Z}^n)$. In this setting, we introduce a suitable notion of a class of classical symbols which provide a class of Dixmier traceable discrete pseudo-differential operators. We also present a formula for the Dixmier trace of a Dixmier traceable discrete pseudo-differential operator by using the Connes equivalence between the Wodzicki residue and the Dixmier trace. **MSC 2010.** Primary 47G30; Secondary 58B34, 47L20.

1. INTRODUCTION

In this paper we study the membership of global pseudo-differential operators associated with the Hörmander symbol classes on \mathbb{Z}^n , also known as discrete pseudo-differential operators, introduced by Botchway, Kibity and Ruzhansky [2] to the Dixmier ideal of operators on $\ell^2(\mathbb{Z}^n)$. For this purpose, we use a fundamental tool of non-commutative geometry, namely, the celebrated theorem of Connes on the equivalence of the Wodzicki residue and the Dixmier trace [9]. Recently, due to its applications in number theory, difference equations and many other problems involving the discretization of differential equations, the theory of pseudo-differential operators on \mathbb{Z}^n has gained an important role in harmonic analysis [31, 45, 33, 25, 30, 43, 44, 45]. Discrete pseudo-differential operators appear early in the classic Fourier analysis (see [17, 21]) as well as discrete counterparts of Calderón-Zygmund singular integral operators. Delgado and Wong [11] started the investigation of spectral properties and traces of discrete pseudo-differential operators. Further investigation has been carried out by several authors in different setting, for example see [12, 13, 14, 15] and references therein.

The question “whether the C^* -algebra of all bounded linear operators on a Hilbert space H had a unique non-trivial trace” led to the concept of the Dixmier trace Tr_ω . In 1966, Dixmier answered this question in negative in a note published in Comptes Rendus [16]. To obtain the trace he used an invariant mean ω on the $ax + b$ group. In contrast to the usual trace, his trace vanishes on the ideal of trace class operators. Applications of the Dixmier trace in differential geometry as well as in non-commutative geometry were facilitated by the celebrated theorem of Connes [9] which relates the trace to the Wodzicki residue [46] for

2010 *Mathematics Subject Classification.* Primary .

Key words and phrases. Discrete pseudo-differential operators; Compact operators; Dixmier ideal; Dixmier Traces; Marcinkiewicz ideal.

pseudo-differential operators on a closed manifold. Both the Dixmier trace and the Wodzicki residue play important roles in noncommutative geometry and its applications to fractal theory, foliation theory, spaces of non-commuting coordinates, perturbation theory, and quantum field theory [9].

In a more recent context, the symbolic calculus of the pseudo-differential operators on \mathbb{Z}^n , introduced by Molahajloo in [29], has been further developed in the work of Botchway, Kibiti, and Ruzhansky [2] where they presented its applications to difference equations. In this setting the pseudo-differential operators on \mathbb{Z}^n are defined by

$$t_m f(n') := \int_{\mathbb{T}^n} e^{i2\pi n' \cdot \xi} m(n', \xi) (\mathcal{F}f)(\xi) d\xi, \quad f \in \mathcal{S}(\mathbb{Z}^n), \quad n' \in \mathbb{Z}^n. \quad (1.1)$$

Under suitable conditions on the symbol m , we have that t_m is a continuous linear operator from the discrete Schwartz space $\mathcal{S}(\mathbb{Z}^n)$ into itself. This is the case of symbols in the discrete Hörmander classes (see e.g. [2]). The references Rabinovich[34, 35], and Rabinovich and Roch[36, 37] are predecessor works on the subject. Analytic and spectral properties as the ellipticity and Fredholmness, nuclearity, the spectral traces, and the L^p and weak- L^p -boundedness for these operators have been considered in Botchway, Kibiti, and Ruzhansky [2], Catana [7], Wong[48], Rodriguez [38], Dasgupta and Kumar [10], and the works of the first and third author [5]. In particular, the Calderón-Vaillancourt theorem and the Gohberg Lemma for the discrete Hörmander classes in [2] have been proved in Cardona [6].

In this paper we are interested in providing sharp conditions on m , in order that t_m in (1.1), belongs to the Dixmier ideal $\mathcal{L}^{1,\infty}(\ell^2(\mathbb{Z}^n))$. The Dixmier traceability of pseudo-differential operators on smooth manifolds is a classical problem. Indeed, the problem was treated by using the notion of local symbols in the works of Connes, Dixmier, Fedosov, Grubb, Schrohe, [8, 19, 20, 22, 32, 46] and references therein. We refer the reader to [4] in which the Dixmier traceability of pseudo-differential operators on compact manifolds has been studied by using the matrix-valued quantization (see [39]).

Our main result is Theorem 3.3, where we prove that, under the condition $t_m \in \Psi_{cl}^{-n}(\mathbb{Z}^n)$ (see Definition 2.4) t_m being a positive pseudo-differential operator, its Dixmier trace is given, surprisingly, by the formula

$$\mathrm{Tr}_\omega(t_m) = \frac{1}{n(2\pi)^n} \int_{\mathbb{T}^n} \int_{\mathbb{S}^{n-1}} [i^*(\tau)(x, \xi)]_{(-n)} d\Sigma(\xi) d_{\mathrm{Vol}}x, \quad \tau(x, k) := \overline{m(-k, x)}, \quad (1.2)$$

where $d\Sigma(\xi)$ is the surface measure associated to the sphere $\mathbb{S}^{n-1} \subset \mathbb{R}^n$, and $d_{\mathrm{Vol}}x$ is the volume form on $M = [0, 1]^n$. The integral formula (1.2) shows that we can capture the spectral information in the Dixmier trace of an operator in terms of the geometry of the manifold $\mathbb{T}^n \times \mathbb{S}^{n-1}$, where the density $d\Sigma(\xi) d_{\mathrm{Vol}}x$ appears, together with the Fourier analysis encoded in the global symbol $i^*\tau = i^*(\overline{m(-\cdot, \cdot)})$.

In order to prove this formula, we will provide some preliminaries in Section 2 and later we will present the proof in Section 3.

2. PSEUDO-DIFFERENTIAL CALCULUS ON \mathbb{Z}^n AND THE n -TORUS \mathbb{T}^n

In this paper we will use the usual notation and terminology given in the standard monographs or papers (see e.g. [18, 23, 24, 39, 47, 48, 41]). The discrete Schwartz space $\mathcal{S}(\mathbb{Z}^n)$ is the the space of all functions $\phi : \mathbb{Z}^n \rightarrow \mathbb{C}$ such that

$$\forall M \in \mathbb{R}, \exists C_M > 0, \text{ such that } |\phi(\xi)| \leq C_M \langle \xi \rangle^M, \quad (2.1)$$

where $\langle \xi \rangle = (1 + |\xi|^2)^{\frac{1}{2}}$. The toroidal Fourier transform is defined as follows: for any $f \in C^\infty(\mathbb{T}^n)$,

$$\widehat{f}(\xi) = \int_{\mathbb{T}^n} e^{-i2\pi \langle x, \xi \rangle} f(x) dx, \quad \xi \in \mathbb{Z}^n.$$

The inverse toroidal Fourier transform is given by the following formula: for $f \in C^\infty(\mathbb{T}^n)$

$$f(x) = \sum_{\xi \in \mathbb{Z}^n} e^{i2\pi \langle x, \xi \rangle} \widehat{f}(\xi), \quad x \in \mathbb{T}^n.$$

The fundamental tool in the analysis of periodic operators, that is, operators defined on function spaces on \mathbb{T}^n , are the Hörmander class of periodic symbols. For every $m \in \mathbb{R}$, the periodic symbol of Hörmander class of order m and of (ρ, δ) -type, denoted by $S_{\rho, \delta}^m(\mathbb{T}^n \times \mathbb{R}^n)$, $0 \leq \rho, \delta \leq 1$, consists of those functions $a(x, \xi)$ which are smooth in $(x, \xi) \in \mathbb{T}^n \times \mathbb{R}^n$ and which satisfy the following toroidal symbols inequalities

$$|\partial_x^\beta \partial_\xi^\alpha a(x, \xi)| \leq C_{\alpha, \beta} \langle \xi \rangle^{m - \rho|\alpha| + \delta|\beta|}. \quad (2.2)$$

It is well-known fact that Symbols in $S_{\rho, \delta}^m(\mathbb{T}^n \times \mathbb{R}^n)$ can be considered as symbols in $S_{\rho, \delta}^m(\mathbb{R}^n \times \mathbb{R}^n)$ (see [23, 39]) of order m which are 1-periodic in x . If $a(x, \xi) \in S_{\rho, \delta}^m(\mathbb{T}^n \times \mathbb{R}^n)$, the corresponding pseudo-differential operator is defined by

$$a(X, D_x)u(x) = \int_{\mathbb{T}^n} \int_{\mathbb{R}^n} e^{i2\pi \langle x-y, \xi \rangle} a(x, \xi) u(y) d\xi dy, \quad u \in C^\infty(\mathbb{T}^n). \quad (2.3)$$

The symbol class $S_{\rho, \delta}^m(\mathbb{T}^n \times \mathbb{Z}^n)$, $0 \leq \rho, \delta \leq 1$, consists of those functions $a(x, \xi)$ which are smooth in x for all $\xi \in \mathbb{Z}^n$ and satisfy

$$\forall \alpha, \beta \in \mathbb{N}^n, \exists C_{\alpha, \beta} > 0, |\Delta_\xi^\alpha \partial_x^\beta a(x, \xi)| \leq C_{\alpha, \beta} \langle \xi \rangle^{m - \rho|\alpha| + \delta|\beta|}. \quad (2.4)$$

The operator Δ_ξ in (2.4) is the difference operator which is defined as follows: if $f : \mathbb{Z}^n \rightarrow \mathbb{C}$ is a discrete function and $(e_j)_{1 \leq j \leq n}$ is the canonical basis of \mathbb{R}^n , then Δ_ξ^α is defined by

$$(\Delta_{\xi_j} f)(\xi) = f(\xi + e_j) - f(\xi). \quad (2.5)$$

If $k \in \mathbb{N}$, denote by $\Delta_{\xi_j}^k$ the composition of Δ_{ξ_j} with itself k -times. Finally, if $\alpha \in \mathbb{N}^n$, $\Delta_\xi^\alpha = \Delta_{\xi_1}^{\alpha_1} \cdots \Delta_{\xi_n}^{\alpha_n}$.

The toroidal pseudo-differential operator (or periodic pseudo-differential operator) corresponding to the symbol $a(x, \xi) \in S_{\rho, \delta}^m(\mathbb{T}^n \times \mathbb{Z}^n)$ is defined by

$$a(x, D_x)u(x) = \sum_{\xi \in \mathbb{Z}^n} e^{i2\pi \langle x, \xi \rangle} a(x, \xi) \widehat{u}(\xi), \quad u \in C^\infty(\mathbb{T}^n). \quad (2.6)$$

Similarly, the Hörmander classes on \mathbb{Z}^n , (see Bothcway, Kibity and Ruzhansky [2]) $S_{\rho,\delta}^m(\mathbb{Z}^n \times \mathbb{T}^n)$, $0 \leq \delta, \rho \leq 1$, are defined by those functions $m \in C^\infty(\mathbb{Z}^n \times \mathbb{T}^n)$, such that,

$$|\partial_x^\beta \Delta_{n'}^\alpha \sigma(x, n')| \leq C_{\alpha,\beta} (1 + |n'|)^{m - \rho|\alpha| + \delta|\beta|}, \quad 0 \leq \rho < 1. \quad (2.7)$$

In this case we denote $\Psi_{\rho,\delta}^m(\mathbb{Z}^n \times \mathbb{T}^n) = \{t_\sigma : \sigma \in S_{\rho,\delta}^m(\mathbb{Z}^n \times \mathbb{T}^n)\}$.

The following proposition says that there exists a method to interpolate the second argument of symbols on $\mathbb{T}^n \times \mathbb{Z}^n$ in a smooth way to get a symbol defined on $\mathbb{T}^n \times \mathbb{R}^n$. The proof can be found in [26, 39].

Proposition 2.1. *Let $0 \leq \delta \leq 1$, $0 < \rho \leq 1$. The symbol $a \in S_{\rho,\delta}^m(\mathbb{T}^n \times \mathbb{Z}^n)$ if only if there exists a Euclidean symbol $a' \in S_{\rho,\delta}^m(\mathbb{T}^n \times \mathbb{R}^n)$ such that $a = a'|_{\mathbb{T}^n \times \mathbb{Z}^n}$.*

It is an interesting but non trivial fact that the definition of pseudo-differential operator on a torus (2.6) given by Agranovich and by Hörmander (2.3) are equivalent. McLean [26] prove this for all Hörmander classes $S_{\rho,\delta}^m(\mathbb{T}^n \times \mathbb{Z}^n)$. A different proof to this fact can be found in [39, Corollary 4.6.13].

Proposition 2.2. *(Equality of Operators Classes). For $0 \leq \delta \leq 1$, $0 < \rho \leq 1$, we have $\Psi_{\rho,\delta}^m(\mathbb{T}^n \times \mathbb{Z}^n) = \Psi_{\rho,\delta}^m(\mathbb{T}^n \times \mathbb{R}^n)$.*

In our further analysis it will be useful record the following fact: there exists a bijection

$$i^* : S_{\rho,\delta}^m(\mathbb{T}^n \times \mathbb{Z}^n) \rightarrow S_{\rho,\delta}^m(\mathbb{T}^n \times \mathbb{R}^n) \quad (2.8)$$

which associates every symbol $\sigma \in S_{\rho,\delta}^m(\mathbb{T}^n \times \mathbb{Z}^n)$ to an Euclidean symbol $i^*\sigma \in S_{\rho,\delta}^m(\mathbb{T}^n \times \mathbb{R}^n)$ such that

$$\sigma(x, D_x)f(x) = (i^*\sigma)(X, D_x)f(x), \quad f \in C^\infty(\mathbb{T}^n). \quad (2.9)$$

The existence of the mapping mapping i^* in (2.10) is a natural consequence of Proposition 2.2. This can be constructed explicitly in terms of the periodisation technique developed by Ruzhansky and Turunen in [39, Section 4.2].

Definition 2.3 (Classical symbols on \mathbb{T}^n). A classical pseudo-differential operator $A = a(X, D_x)$ on \mathbb{T}^n is an operator with symbol $a \in S_{1,0}^m(\mathbb{T}^n \times \mathbb{R}^n)$, admitting an asymptotic expansion

$$a(x, \xi) \sim \sum_{j=0}^{\infty} a_{m-j}(x, \xi) \quad (2.10)$$

where $a_{m-j}(x, \xi)$ are homogeneous functions with respect to ξ of degree $m - j$ for ξ far from to zero.

The set of classical symbols of order m on \mathbb{T}^n will be denoted by $S_{cl}^m(\mathbb{T}^n \times \mathbb{R}^n)$ and the corresponding class of periodic pseudo-differential operators will be denoted by $\Psi_{cl}^m(\mathbb{T}^n)$.

Now, with the help of the classical symbols on \mathbb{T}^n , we introduce classical pseudo-differential operators on \mathbb{Z}^n . We start with the following definition of the classical symbols of \mathbb{Z}^n .

Definition 2.4 (Classical symbols on \mathbb{Z}^n). A symbol $\sigma \in S_{1,0}^m(\mathbb{Z}^n \times \mathbb{T}^n)$ is called classical, if the symbol $\tau(x, k) := \overline{\sigma(-k, x)}$ satisfies that $i^* \tau \in S_{cl}^m(\mathbb{T}^n \times \mathbb{R}^n)$.

The set of classical symbols of order m on \mathbb{Z}^n will be denoted by $S_{cl}^m(\mathbb{Z}^n \times \mathbb{T}^n)$ and the corresponding class of periodic pseudo-differential operators will be denoted by $\Psi_{cl}^m(\mathbb{Z}^n)$.

Periodic and also discrete Hörmander classes of pseudo-differential operators are closed under compositions, adjoints and parametrices. We refer the reader to [2, 39, 40] for details.

3. DIXMIER TRACES FOR PSEUDO-DIFFERENTIAL OPERATORS ON \mathbb{Z}^n

In this section, we present our main result. We begin this section by recalling some basic facts on Dixmier trace and related results.

Let H be a Hilbert space and let $\mathcal{L}(H)$ be the algebra of all bounded linear operators on H . We say that a linear operator $A \in \mathcal{L}(H)$ is in the Dixmier class $\mathcal{L}^{(1,\infty)}(H)$ if

$$\sum_{1 \leq n \leq N} s_n(A) = O(\log(N)), \quad N \rightarrow \infty, \tag{3.1}$$

where $s(A) := \{s_n(A)\}_n$ denotes the sequence of singular values of A , i.e. the square roots of the eigenvalues of the non-negative self-adjoint operator A^*A , and $s_n(A)$ are arranged in increasing order. So, $\mathcal{L}^{(1,\infty)}(H)$ is endowed with the norm

$$\|A\|_{\mathcal{L}^{(1,\infty)}(H)} = \sup_{N \geq 2} \frac{1}{\log(N)} \sum_{1 \leq n \leq N} s_n(A). \tag{3.2}$$

The Dixmier trace Tr_ω of a positive operator A in $\mathcal{L}^{(1,\infty)}(H)$ can be formulated by using an average function ω of the following form

$$\text{Tr}_\omega(A) = \lim_\omega \frac{1}{\log(N)} \sum_{1 \leq n \leq N} s_n(A),$$

where \lim_ω is a positive linear functional on $l^\infty(\mathbb{N})$, the set of bounded sequences, satisfying:

- $\lim_\omega(\alpha_n) \geq 0$ for all $\alpha_n \geq 0$;
- $\lim_\omega(\alpha_n) = \lim(\alpha_n)$ whenever the ordinary limit exists;
- $\lim_\omega(\alpha_1, \alpha_1, \alpha_2, \alpha_2, \alpha_3, \alpha_3 \dots) = \lim_\omega(\alpha_n)$.

The functional Tr_ω can be defined for positive operators, and later extended to the whole ideal $\mathcal{L}^{(1,\infty)}(H)$ by linearity (in this case Tr_ω is not necessarily a positive functional).

Let us assume that M is a closed manifold. In the framework of noncommutative geometry, a remarkable result due to Connes shows that any classical pseudo-differential operator, of order of the minus of the dimension of the manifold underlying, acting on $L^2(M)$ lies to the Dixmier measurable class, and moreover the Dixmier trace coincides with the noncommutative residue, which is the unique trace (up to by factors) in the algebra of pseudo-differential operators with classical symbols on a closed manifold.

Now, we recall that for a compact manifold M of dimension \varkappa without boundary, a classical pseudo-differential operator A on M of order m , is defined by means of local symbols, this means that for any local chart U , the operator A reads

$$Au(x) = \int_{T_x^*U} e^{ix\xi} \sigma^A(x, \xi) \widehat{u}(\xi) d\xi$$

where $\sigma^A(x, \xi)$ is a smooth function on $T^*U \cong U \times \mathbb{R}^n$, $T_x^*U = \mathbb{R}^n$, and admits an asymptotic expansion

$$\sigma^A(x, \xi) \sim \sum_{j=0}^{\infty} \sigma_{m-j}^A(x, \xi) \quad (3.3)$$

where $\sigma_{m-j}^A(x, \xi)$ are homogeneous functions with respect to ξ of degree $m-j$ for ξ far away from the section zero. The set of classical pseudo-differential operators of order m is denoted by $\Psi_{cl}^m(M)$. For $A \in \Psi_{cl}(M)$, for $x \in M$, $\int_{|\xi|=1} \sigma_{-\varkappa}(x, \xi) d\Sigma(\xi)$,

defines a local density which can be glued over M . So, a linear functional called the noncommutative residue is defined by

$$\text{res}(A) = \frac{1}{\varkappa(2\pi)^\varkappa} \int_M \int_{|\xi|=1} \sigma_{-\varkappa}^A(x, \xi) d\Sigma(\xi) d_{\text{Vol}}x. \quad (3.4)$$

which vanishes on non-integer order classical operators.

In order to provide a trace formula for pseudo-differential operators on \mathbb{Z}^n we will use the following identity in [2]

$$t_\sigma = \mathcal{F}_{\mathbb{Z}^n}(\tau(x, D_x)^*) \mathcal{F}_{\mathbb{Z}^n}^{-1}, \quad (3.5)$$

and the following fundamental result due to Connes [8].

Theorem 3.1 (A. Connes). *Let M be a closed manifold of dimension \varkappa . Then every classical and positive pseudo-differential operator A of order $-\varkappa$ lies in $\mathcal{L}^{(1, \infty)}(L^2(M))$ and*

$$\text{Tr}_w(A) = \text{res}(A). \quad (3.6)$$

Corollary 3.2. *Let $\tau(x, D_x)$ be a classical positive pseudo-differential operator with symbol $\tau \in S_{1,0}^{-n}(\mathbb{T}^n \times \mathbb{R}^n)$. Then $\tau(x, D_x)$ belongs to $\mathcal{L}^{(1, \infty)}(L^2(\mathbb{T}^n))$ and*

$$\text{Tr}_w(\tau(x, D_x)) = \text{res}(\tau(x, D_x)). \quad (3.7)$$

The following theorem is our main result on the Dixmier traceability of pseudo-differential operators on \mathbb{Z}^n .

Theorem 3.3. *Let us assume that $t_\sigma \in \Psi_{cl}^{-n}(\mathbb{Z}^n)$ is a positive discrete pseudo-differential operator. Then the operator t_σ is Dixmier traceable on $\ell^2(\mathbb{Z}^n)$ and its Dixmier trace is given by*

$$\text{Tr}_w(t_\sigma) = \frac{1}{n(2\pi)^n} \int_{\mathbb{T}^n} \int_{\mathbb{S}^{n-1}} [i^*(\tau)(x, \xi)]_{(-n)} d\Sigma(\xi) d_{\text{Vol}}x, \quad \tau(x, k) := \overline{\sigma(-k, x)}, \quad (3.8)$$

where $[i^*(\tau)(x, \xi)]_{(-n)}$ in the homogeneous term of order $-n$ is the asymptotic expansion of the symbol $i^*(\tau)(x, \xi)$ in homogeneous components as in (2.10).

Proof. Let us consider the following identity due to Botchway, Kibiti and Ruzhansky [2]

$$t_\sigma = \mathcal{F}_{\mathbb{Z}^n}(\tau(x, D_x)^*)\mathcal{F}_{\mathbb{Z}^n}^{-1}. \quad (3.9)$$

First we will prove that the set of the singular values of t_σ agrees with the set of singular values of $\tau(x, D_x)^*$. We show this by using the fact that the spectrum is stable under unitary transformations. Since the Fourier transform on \mathbb{Z}^n is a unitary operator, we have that

$$\begin{aligned} s(\tau(x, D_x)^*) &= \{\sqrt{\lambda} : \lambda \in \text{Spectrum}(\tau(x, D_x)\tau(x, D_x)^*)\} \\ &= \{\sqrt{\lambda} : \lambda \in \text{Spectrum}(\mathcal{F}_{\mathbb{Z}^n}\tau(x, D_x)\tau(x, D_x)^*\mathcal{F}_{\mathbb{Z}^n}^{-1})\} \\ &= \{\sqrt{\lambda} : \lambda \in \text{Spectrum}(\mathcal{F}_{\mathbb{Z}^n}\tau(x, D_x)\mathcal{F}_{\mathbb{Z}^n}^{-1}\mathcal{F}_{\mathbb{Z}^n}\tau(x, D_x)^*\mathcal{F}_{\mathbb{Z}^n}^{-1})\} \\ &= \{\sqrt{\lambda} : \lambda \in \text{Spectrum}(\mathcal{F}_{\mathbb{Z}^n}\tau(x, D_x)\mathcal{F}_{\mathbb{Z}^n}^{-1} \circ t_\sigma)\} \\ &= \{\sqrt{\lambda} : \lambda \in \text{Spectrum}(t_\sigma^*t_\sigma)\} = s(t_\sigma), \end{aligned}$$

where in the last line we have used $t_\sigma^* = \mathcal{F}_{\mathbb{Z}^n}\tau(x, D_x)\mathcal{F}_{\mathbb{Z}^n}^{-1}$. Now, it is clear that t_σ is Dixmier traceable if and only if $\tau(x, D_x)^*$ is Dixmier traceable and in a such case $\text{Tr}_\omega(t_\sigma) = \text{Tr}_\omega(\tau(x, D_x)^*)$. The positivity of t_σ implies that $\tau(x, D_x)$ is also positive. Because $t_\sigma \in \Psi_{cl}^{-n}(\mathbb{Z}^n)$ we have that $\tau(x, D_x) \in \Psi_{cl}^{-n}(\mathbb{T}^n)$ and consequently $\tau(x, D_x)$ is a compact operator on $L^2(\mathbb{T}^n)$, (see e.g. [27]). In particular $\tau(x, D_x)$ is bounded and taking into account its positivity we conclude that $\tau(x, D_x)$ is self-adjoint. So, if we assume the Dixmier traceability of $\tau(x, D_x) = \tau(x, D_x)^*$, we deduce $\text{Tr}_\omega(\tau(x, D_x)^*) = \text{Tr}_\omega(\tau(x, D_x))$. Since $\tau(x, D_x) \in \Psi_{cl}^{-n}(\mathbb{T}^n)$ by the Connes Theorem 3.1 in the form of Corollary 3.2, we conclude that $\tau(x, D_x)$ is Dixmier traceable and its Dixmier trace can be computed from its noncommutative residue. So, we have

$$\begin{aligned} \text{Tr}_\omega(\tau(x, D_x)) &= \text{Tr}_\omega((i^*\tau)(X, D_x)) = \text{res}((i^*\tau)(X, D_x)) \\ &= \frac{1}{n(2\pi)^n} \int_{\mathbb{T}^n} \int_{\mathbb{S}^{n-1}} [i^*(\tau)(x, \xi)]_{(-n)} d\Sigma(\xi) d_{\text{Vol}}x. \end{aligned}$$

Consequently we obtain

$$\text{Tr}_\omega(t_\sigma) = \frac{1}{n(2\pi)^n} \int_{\mathbb{T}^n} \int_{\mathbb{S}^{n-1}} [i^*(\tau)(x, \xi)]_{(-n)} d\Sigma(\xi) d_{\text{Vol}}x. \quad (3.10)$$

Thus, we finish the proof of the theorem. □

Acknowledgements. This work was started when Duván Cardona was supported by Pontificia Universidad Javeriana. César del Corral was supported by Universidad de Los Andes and Vishvesh Kumar was supported by CSIR India.

Currently, Duván Cardona and Vishvesh Kumar are supported by Odysseus I Project (by FWO, Belgium) of Prof. Michael Ruzhansky. They thank Prof. Michael Ruzhansky for his support and encouragement.

Es con mucho aprecio que dedicamos éste trabajo al Profesor Lázaro Recht, uno de los mejores matemáticos argentinos. Su contribución a las generaciones

matemáticas jóvenes, en Argentina, Venezuela y Colombia ha sido determinante, aunque él, modestia aparte, opine lo contrario.

REFERENCES

1. Agranovich, M. S.: Spectral properties of elliptic pseudodifferential operators on a closed curve *Funct. Anal. Appl.* **13**, 279-281 (1971)
2. Botchway L., Kibiti G., Ruzhansky M., Difference equations and pseudo-differential operators on \mathbb{Z}^n , arXiv:1705.07564.
3. Calderón, A. P., Zygmund, A. On the existence of certain singular integrals, *Acta. Math.* **88** (1952), 85–139.
J. Pseudo-Differ. Oper. Appl., 5(4), (2014) 507-515.
4. Cardona, D., Del Corral, C. The Dixmier trace and the non-commutative residue for multipliers on compact manifolds, submitted. arXiv:1703.07453.
5. Cardona, D., Kumar, V., L^p -boundedness and L^p -nuclearity of multilinear pseudo-differential operators on \mathbb{Z}^n and the torus \mathbb{T}^n , *J. Fourier Anal. Appl.* DOI: 10.1007/s00041-019-09689-7, arXiv:1809.08380
6. Cardona, D. Pseudo-differential operators on \mathbb{Z}^n with applications to discrete fractional integral operators. *Bull. Iran. Math. Soc.* To appear. DOI: 10.1007/s41980-018-00195-y. arXiv:1803.00231.
7. Catana, V. L^p -boundedness of multilinear pseudo-differential operators on \mathbb{Z}^n and \mathbb{T}^n . *Math. Model. Nat. Phenom.*, 9(5), (20014), 17–38.
8. Connes A. Noncommutative geometry. Academic Press Inc., San Diego, CA (1994)
9. Connes, A. The action functional in noncommutative geometry, *Comm. Math. Phys.* **117**, no. 4, 673–683, 1988.
10. Dasgupta, A. and Kumar, V.: Ellipticity and Fredholmness of pseudo-differential operators on $\ell^2(\mathbb{Z}^n)$, (2019). arXiv:1910.05582v2
11. Delgado, J., Wong, M.W.: L^p -nuclear pseudo-differential operators on \mathbb{Z} and \mathbb{S}^1 , *Proc. Amer. Math. Soc.*, **141**(11), (2013), 3935–394.
12. Delgado, J. and Ruzhansky, M.: L^p -nuclearity, traces, and Grothendieck-Lidskii formula on compact Lie groups. *J. Math. Pures Appl.*, (**9**), 102(1), 153-172 (2014).
13. Delgado, J. and Ruzhansky, M.: Schatten classes on compact manifolds: Kernel conditions. *J. Funct. Anal.*, **267**(3), 772-798 (2014)
14. Delgado, J. and Ruzhansky, M.: Kernel and symbol criteria for Schatten classes and r -nuclearity on compact manifolds. *C. R. Acad. Sci. Paris. Ser. I.*, **352**, 779–784 (2014).
15. Delgado, J. and Ruzhansky, M.: Fourier multipliers, symbols and nuclearity on compact manifolds. *J. Anal. Math.*, **135**(2) 757–800 (2018).
16. Dixmier, J. Existence de traces non normales, (French) *C. R. Acad. Sci. Paris* **262** (1966) 1107–1108.
17. Duoandikoetxea, J.: *Fourier Analysis*, Amer. Math. Soc. (2001)
18. Folland, G.B.: *Harmonic Analysis in phase space*. Princeton University Press, (1989)
19. Fedosov, B. Golse, F. Leichtnam, E. Schrohe, E. The Noncommutative Residue for Manifolds with Boundary. *J. Funct. Anal.* Vol. **142** (1), 1996, 1–31.
20. Gayral, V., Gracia-Bondía, J.M., Iochum, B. Schücker, T., Vàrilly, J.C. Moyal planes are spectral triples. *Comm. Math. Phys.* **246** (2004), 569623.
21. Grafakos, L., *Classical Fourier Analysis* Grad. Texts in Math., vol. 249, Springer-Verlag, New York (2008)
22. Grubb, G. Schrohe, E. Trace expansions and the noncommutative residue for manifolds with boundary, Univ. of Copenhagen, Preprint Series no. 20., 1999
23. Hörmander, L.: *Pseudo-differential Operators and Hypo-elliptic equations* Proc. Symposium on Singular Integrals, Amer. Math. Soc. **10**, 138-183 (1967)
24. Hörmander, L.: *The Analysis of the linear partial differential operators* Vol. III. Springer-Verlag, (1985)

25. Ionescu, A.D., Wainger, S. L^p boundedness of discrete singular Radon transforms, J. Amer. Math. Soc. 19(2) (2005), 357–383.
26. Mclean, W.M.: Local and Global description of periodic pseudo-differential operators, Math. Nachr. **150**, 151-161 (1991)
27. Molahajloo, S.: A characterization of compact pseudo-differential operators on \mathbb{S}^1 Oper. Theory Adv. Appl. Birkhäuser/Springer Basel AG, Basel. **213**, 25-29 (2011)
28. Molahajloo, S., Wong, M.W.: Pseudo-differential Operators on \mathbb{S}^1 . New developments in pseudo-differential operators, Eds. L. Rodino and M.W. Wong. 297-306 (2008)
29. Molahajloo, S., Wong, M. W. : Ellipticity, Fredholmness and spectral invariance of pseudo-differential operators on \mathbb{S}^1 . J. Pseudo-Differ. Oper. Appl. **1** 183-205 (2010)
30. Pierce, L. Discrete Analogues in Harmonic Analysis. P.h.D Thesis, Princeton University (2009).
31. Pierce, L. On discrete fractional integral operators and mean values of Weyl sums. Bull. London Math. Soc., 43 (2011) 597–612.
32. Schrohe, E. Noncommutative Residues, Dixmier’s Trace, and Heat Trace Expansions on Manifolds with Boundary. In B. Booss-Bavnbek and K. Wojciechowski (eds), Geometric Aspects of Partial Differential Equations. Contemporary Mathematics, vol. 242, Amer. Math. Soc. Providence, R.I., 1999, p. 161–186
33. Hooley, C. On Hypothesis K^* in Warings problem, Sieve Methods, Exponential Sums, and their Applications in Number Theory, London Math. Soc. Lecture Notes No. 237, Cambridge University Press, (1997), 175–185.
34. Rabinovich, V. Exponential estimates of solutions of pseudodifferential equations on the lattice $(\mu\mathbb{Z})^n$: applications to the lattice Schrödinger and Dirac operators. J. Pseudo-Differ. Oper. Appl., 1(2), (2010), 233–253.
35. Rabinovich, V. . Wiener algebra of operators on the lattice $(\mu\mathbb{Z})^n$ depending on the small parameter $\mu > 0$. Complex Var. Elliptic Equ., 58(6), (2013), 751–766.
36. Rabinovich, V. S., Roch, S. The essential spectrum of Schrödinger operators on lattices. J. Phys. A, 39(26), (2006), 8377–8394.
37. Rabinovich, V. S., Roch, S. Essential spectra and exponential estimates of eigenfunctions of lattice operators of quantum mechanics. J. Phys. A, 42(38), (2009), 385–207.
38. Rodriguez, C. A., L^p -estimates for pseudo-differential operators on \mathbb{Z}^n , J. Pseudo-Differ. Oper. Appl. 1, (2011), 183–205.
39. Ruzhansky, M., Turunen, V.: Pseudo-differential Operators and Symmetries: Background Analysis and Advanced Topics Birkhäuser-Verlag, Basel, (2010)
40. Ruzhansky, M., Turunen, V.: Quantization of Pseudo-Differential Operators on the Torus J Fourier Annal Appl. Vol. 16. pp. 943-982. Birkhäuser Verlag, Basel, (2010)
41. Ruzhansky, ., Turunen, V., On the toroidal quantization of periodic pseudodifferential operators, Numerical Functional Analysis and Optimization, 30 (2009), 1098–1124.
42. Turunen, V., Vainikko, G.: On symbol analysis of periodic pseudodifferential operators Z. Anal. Anwendungen. **17**, 9-22 (1998)
43. Stein, E., Wainger, S., Discrete analogues of singular Radon transforms, Bull. Amer. Math. Soc. (N.S.) 23 (1990), no. 2, 537-544.
44. Stein, E., Wainger, S., Discrete analogues in harmonic analysis, I. l^2 estimates for singular Radon transforms,” Amer. J. Math. 121(6) (1999), 1291–1336.
45. Stein, E., Wainger, S. Discrete analogues in harmonic analysis, II. Fractional integration, J. Anal. Math. 80 (2000), 335–355.
46. Wodzicki, M. Noncommutative residue I, Fundamentals, in K-theory, arithmetic and geometry (Moscow, 1984–1986), 320–399, Lecture Notes in Math., 12-89, Springer, Berlin, 1987.
47. Wong, M. W.: An introduction to Pseudo-Differential Operators Second Edition, World Scientific, (1999)
48. Wong, M. W.: Discrete Fourier Analysis Birkhäuser, (2011)

DUVÁN CARDONA:

DEPARTMENT OF MATHEMATICS
PONTIFICIA UNIVERSIDAD JAVERIANA
BOGOTÁ-COLOMBIA

CURRENT ADDRESS

DEPARTMENT OF MATHEMATICS: ANALYSIS LOGIC AND DISCRETE MATHEMATICS
GHENT UNIVERSITY
GHENT-BELGIUM

E-mail address duvanc306@gmail.com

CÉSAR DEL CORRAL:

DEPARTMENT OF MATHEMATICS
UNIVERSIDAD DE LOS ANDES
BOGOTÁ

COLOMBIA

E-mail address ce-del@uniandes.edu.co; cesar.math@gmail.com

VISHVESH KUMAR:

DEPARTMENT OF MATHEMATICS
INDIAN INSTITUTE OF TECHNOLOGY DELHI
HUAZ KHAS, NEW DELHI- 110016
DELHI, INDIA

CURRENT ADDRESS

DEPARTMENT OF MATHEMATICS: ANALYSIS LOGIC AND DISCRETE MATHEMATICS
GHENT UNIVERSITY
GHENT-BELGIUM

E-mail address vishveshmishra@gmail.com